PlantRNA-FM: An Interpretable RNA Foundation Model for Exploration

Functional RNA Motifs in Plants

Haopeng Yu[1,3†], Heng Yang[2,†], Wenqing Sun[1,3†], Zongyun Yan[1†], Xiaofei Yang[4,5], Huakun

Zhang[3*], Yiliang Ding[1,*] & Ke Li[2,*]

1 Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park,

Norwich NR4 7UH, UK

2 Department of Computer Science, University of Exeter, EX4 4QF, Exeter, UK

3 Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal

University, Changchun 130024, China

4 National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in

Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of

Sciences, Shanghai 200032, China.

5 CAS-JIC Center of Excellence for Plant and Microbial Sciences, Institute of Plant Physiology

and Ecology, Chinese Academy of Sciences, Shanghai 200032, China.

* To whom correspondence should be addressed. Tel: +44 1392 724557; Email:

k.li@exeter.ac.uk.

Correspondence may also be addressed to yiliang.ding@jic.ac.uk or zhanghk045@nenu.edu.cn

† The authors wish it to be known that, in their opinion, the first four authors should be regarded

as Joint First Authors.

ABSTRACT

The complex 'language' of plant RNA encodes a vast array of biological regulatory elements that orchestrate crucial aspects of plant growth, development, and adaptation to environmental stresses. Recent advancements in foundation models (FMs) have demonstrated their unprecedented potential to decipher complex 'language' in biology. In this study, we introduced PlantRNA-FM, a novel high-performance and interpretable RNA FM specifically designed based on RNA features including both sequence and structure. PlantRNA-FM was pre-trained on an extensive dataset, integrating RNA sequences and RNA structure information from 1,124 distinct plant species. PlantRNA-FM exhibits superior performance in plant-specific downstream tasks, such as plant RNA annotation prediction and RNA translation efficiency (TE) prediction. Compared to the second-best FMs, PlantRNA-FM achieved an $F1$ score improvement of up to 52.45% in RNA genic region annotation prediction and up to 15.30% in translation efficiency prediction, respectively. Our PlantRNA-FM is empowered by our interpretable framework that facilitates the identification of biologically functional RNA sequence and structure motifs, including both RNA secondary and tertiary structure motifs across transcriptomes. Through experimental validations, we revealed novel translation-associated RNA motifs in plants. Our PlantRNA-FM also highlighted the importance of the position information of these functional RNA motifs in genic regions. Taken together, our PlantRNA-FM facilitates the exploration of functional RNA motifs across the complexity of transcriptomes, empowering plant scientists with novel capabilities for programming RNA codes in plants.

Introduction

The transcriptome contains a wide array of RNA motifs that impact diverse biological functions such as translation[1–5]. These RNA motifs encompass both RNA sequence and structure features. Previous individual studies have revealed the functional importance of RNA

sequence features such as the Kozak sequence motif[6]. Recently, our studies along with others

have suggested that both RNA secondary and tertiary structure motifs play important roles in

diverse biological processes[7–13]. Particularly in plants, the relatively low habitat temperatures

(~20 °C) favour the folding of RNA structure motifs, including RNA tertiary motifs such as

RNA G-quadruplex (rG4)[12]. However, systematically identifying functional RNA motifs

across transcriptomes remains a formidable challenge due to the high level of complexity

arising from astronomical combinations of the four nucleotide bases into tens of thousands of

transcripts[8,14]. For example, for a 50-nucleotide sequence, the number of artificially

synthesized sequences would be on the order of 4^{50} (approximately 1.27×10^{30}), which is

impossible to achieve experimentally. Additionally, the functional readouts using the reporter

gene assay for measuring biological functions such as translation may not be sensitive enough

to detect differences in individual single-nucleotide mutations[15].

The recent rapid advancements of foundation models (FMs) in artificial intelligence

(AI) are set to show exciting promise for supercharging scientific advances in life sciences[16].

FMs are distinguished by their massive scale, often encompassing millions to billions of

parameters. They are first pre-trained in a self-supervised manner on diverse forms of

unlabelled data. This makes them ideally suitable for bioscience, where acquiring abundant

labelled data is both prohibitively expensive and time-consuming. More importantly, FMs are

highly adaptable through fine-tuning and are poised to aid bioscientists in customising

generalist FMs in unravelling complex biological processes, paving the way for unprecedented

capabilities in modulating gene functions. For FMs on DNA sequences, DNABERT2 is one of

the FMs pre-trained on the genome sequences across 135 species, including mammals, fungi

and bacteria[17]. By pre-training on diverse human and non-human genomes, the Nucleotide

Transformers (NT) family learns transferable representations that enable accurate molecular

phenotype prediction with limited annotated data, while focusing on key genomic elements

without supervision[18]. FMs have also achieved success in protein sequences, also known as

protein language models. For example, ESM2 (Evolutionary Scale Modeling) has achieved

remarkable breakthroughs in atomic-level structure representations by pretraining on a vast

amount of protein sequences and structures[19].

For building RNA FM, several FM models were pre-trained using RNA sequence

information that has demonstrated great performance in RNA molecule design[20–22]. However,

RNA sequence information is not sufficient since RNA is capable of forming secondary or

tertiary structure motifs that are important for its functions[23,24]. Therefore, it is important to

generate an RNA FM including both RNA sequence and structure information to facilitate the

exploration of functional RNA motifs. Here, we developed PlantRNA-FM, a groundbreaking

RNA FM designed to globally identify functional RNA motifs including both RNA sequences

and structure motifs in plants (Fig. 1). By incorporating RNA sequences, annotations, and

structure information from 1,124 distinct plant species, PlantRNA-FM captures the extensive

diversity of plant transcriptomes (Fig. 1). We validate the superior performance of PlantRNA-

FM in downstream tasks compared to existing FMs. Furthermore, we also established an

interpretable framework based on our PlantRNA-FM to determine the critical regions across

the 5' untranslated regions (5' UTRs) that significantly impact translation. Remarkably,

PlantRNA-FM identifies RNA motifs at the transcriptome-wide scale that are functionally

important to translation including both RNA sequences, and secondary and tertiary structure

motifs. We further experimentally validated these identified RNA motifs in plants. The

development of our PlantRNA-FM represents a significant leap forward in our ability to

decipher hidden regulatory codes among the extensive complexity of nucleotides across the

transcriptome, opening new avenues for RNA-based gene regulation.

1. Results

1.1. Our PlantRNA-FM integrates both RNA sequence and structure information of the transcriptomes across 1,124 plant species.

The plant kingdom encompasses approximately 500,000 species, exhibiting remarkable diversity. The One Thousand Plant Transcriptomes Initiative (1KP) sequenced the transcriptomes of 1,124 species, capturing the extensive diversity of plant transcriptomes[14]. Here, we took advantage of this unique resource and generated the pre-training dataset for our PlantRNA-FM (Fig. 1). Different from existing FMs, our PlantRNA-FM was designed to capture and learn both RNA sequences and RNA structure motifs. We employed *RNAfold*[25] to predict RNA structures of individual RNA sequences across 1,124 transcriptomes and integrated them into the pre-training dataset. Our PlantRNA-FM has 35 million parameters, including 12 transformer network layers, 24 attention heads, and an embedding dimension of 480, optimised for RNA understanding rather than generation (**Methods**). Our tokenization approach surpasses the constraints of conventional k-mers and BPE methods, ensuring the preservation of RNA structure motifs as coherent units throughout the pre-training process (**Methods**). In addition, we incorporated RNA annotation information (CDS and UTRs) and employed advanced pre-training techniques, such as sequence truncation, filtering and masked nucleotide modeling (**Methods**).

To assess the effectiveness of our PlantRNA-FM in RNA structure prediction tasks, we evaluated its performance (Fig. S1, Table S1) using three benchmark datasets: bpRNA, ArchiveII, and RNAstralign[26–28]. The $F1$ score, which is the harmonic mean of precision and recall, was used to measure the model's predictive performance on these datasets. The $F1$ scores achieved by our PlantRNA-FM on these three datasets were 0.750, 0.924, and 0.981, respectively, while *RNAfold* alone only obtained $F1$ scores of 0.278, 0.759, and 0.748 (Fig. S1, Table S1). When compared to other state-of-the-art FMs, PlantRNA-FM outperformed the second-best model by 22.10%, 27.49%, and 17.38% on the respective datasets (Fig. S1,

120   Table S1). Therefore, the unique integration of RNA structure information equips our

121   PlantRNA-FM with the ability to predict RNA structure more accurately.

1.2.   PlantRNA-FM demonstrates superior performance on plant-specific downstream tasks

To evaluate the performance of PlantRNA-FM, we curated a benchmark set consisting

of four other state-of-the-art FMs: DNABERT-2, Nucleotide Transformer, ESM2, and

cdsBERT. We assessed their performance in two plant-specific downstream tasks: genic region

annotation and translation efficiency (TE) prediction (Fig. 2a).

In the RNA genic region annotation prediction task, we aimed to identify and classify

different genic regions of given RNA sequences, such as the 5' UTR, coding sequence (CDS),

and 3' UTR. We used the transcriptomes of two model plant species, *Arabidopsis thaliana* (a

dicot model plant) and *Oryza sativa L.* ssp. *Japonica* (rice, a moncot model plant). Both of

them were not included in our pre-training dataset. For the RNA genic region annotation

prediction in these two species, our PlantRNA-FM outperformed other FM models, achieving

average F1 scores of 0.974 and 0.958 for *Arabidopsis* and rice, respectively, surpassing the

second-best model by 52.45% and 43.90% (Fig. 2b, Table 1).

For translation, one of the key RNA biological processes, previous research has

highlighted the critical role of the 5' UTR in regulating translation efficiencies[17–19,21,29–31]. To

evaluate the TE prediction performance of our PlantRNA-FM, we used the 5' UTR sequences

of both *Arabidopsis* and rice transcriptomes along with the corresponding TE values measured

by polysome profiling[8]. We first classified the TE datasets into high and low TE groups, using

the mean plus or minus the standard deviation as the threshold. In the TE prediction task,

PlantRNA-FM achieved F1 scores of 0.735 and 0.737 for *Arabidopsis* and rice, respectively,

outperforming the second-best model by 15.30% and 13.83% (Fig. 2c). Taken together, our

PlantRNA-FM is better suited for plant-specific downstream tasks compared to other FMs pre-

trained on non-plant datasets.

1.3. Interpretable PlantRNA-FM revealed RNA features important to

translation

A general roadblock in applying AI models to biology is that, while these models

demonstrate strong predictive capabilities, the key to their successful application lies in

interpreting them to uncover the biological principles learned by the AI. In this paper, we

established an interpretable framework to derive an attention contrast matrix from our

PlantRNA-FM (**Methods**). In particular, we are interested in extracting the key RNA features

within the 5' UTR that significantly impact RNA translation, i.e., elucidating the RNA motifs

associated with translation (Fig. 3a). We developed two models in parallel: one is the true

model, denoted as PlantRNA-FM(+), trained using the real TE dataset, while the other one is

called the background model, PlantRNA-FM(-), altered using the same dataset but with

randomly assigned labels (Fig. 3a). The $F1$ score achieved by the background model is

approximately 50%, which is close to the random chance (mean $F1 = 0.522$), while the true

model attained a significantly higher mean $F1$ score of 0.737. This indicates that PlantRNA-

FM(+) has successfully learned the RNA features in the 5' UTR sequences associated with

translation.

By subtracting the attention matrices of the background model from those of the true

model, we obtained an attention contrast matrix that highlighted the significance of nucleotides

in the 5' UTR contributing to TE (Fig. 3a). Across the transcriptomes, we observed an increase

in attention contrast scores as the position approached the AUG start codon in both *Arabidopsis*

and rice (Fig. 3b). This result indicates that positions close to the start codon contribute the

most to the TE values. By underlining the RNA sequence contents with high contrast attention

score (identified by a z-score > 2.326), our PlantRNA-FM successfully identified the Kozak

sequence motifs in both *Arabidopsis* and rice transcriptomes that are associated with TE (Fig.

3c, 3d). This result demonstrates that our PlantRNA-FM successfully identifies evolutionarily

conserved RNA motifs that are important to translation (Fig. 3c, 3d).

1.4. PlantRNA-FM globally identifies the translation-associated RNA secondary structure motifs

Since RNA structure is the unique RNA feature incorporated in our PlantRNA-FM, we

further identified the RNA secondary structure motifs important to translation through the

model's attention contrast matrix and an unsupervised hierarchical clustering strategy (Fig. 4a,

Methods). Overall, we identified 112 RNA secondary structure motifs that are important to

translation, including 63 low translation-associated and 49 high translation-associated RNA

secondary structure motifs (Table S2). Notably, we identified low translation-associated RNA

secondary structure motifs with high GC base pairs such as the RNA secondary structure motif

with four GC base pairs in the stem (Fig. 4b). Interestingly, we also identified high translation-

associated RNA structure motifs with a balanced ratio of GC and AU base pairs such as the

RNA structure motif with four base pairs formed by two repeats of ACGU (Fig. 4c).

To validate our identified RNA secondary structure motifs important to translation, we

conducted experimental validation using the dual luciferase reporter assay in plants[12]. For the

high translation-associated RNA secondary structure motif with four base pairs formed by two

repeats of ACGU, we changed the two AU base pairs to the two GC base pairs, resulting in a

significant decrease in TE with a reduction up to 5.3 -fold (Fig. 4d). In contrast, when we

exchanged the low translation-associated RNA secondary structure motif with four GC base

pair in the stem for the high translation-associated RNA secondary structure motif with a

balanced mix of GC and AU base pairs, we found a significant increase in TE (Fig. 4e).

Notably, when we completely disrupted this low translation-associated RNA structure motif,

resulting in complete single-strandedness, we observed an even greater enhancement of TE up

to 2.1-fold (Fig. 4f). Our results demonstrate that PlantRNA-FM is capable of determining

functional RNA secondary structure motifs in plants.

1.5. PlantRNA-FM globally identifies the translation-associated RNA

tertiary structure motifs

RNA G-quadruplexes (rG4s) are one of the RNA tertiary structure motifs formed by

the stacking of two or more G-quartets, composed of four guanines held together by both

Watson-Crick and Hoogsteen hydrogen bonds[8,32,33]. Previous studies have demonstrated the

important role of individual rG4s in repressing translation[34]. However, it is impossible to

identify all the rG4 motifs important to translation from tens of thousands of rG4 motifs across

the transcriptome. Therefore, we took advantage of our PlantRNA-FM to identify the

translation-associated rG4s at the transcriptome-wide scale.

We first obtained all rG4 motifs in the 5' UTRs from our G4Atlas database[33].

Subsequently, we identified all rG4 motifs associated with translation using our model's

attention contrast matrix across the transcriptome (**Methods**). Notably, we only identified rG4

motifs associated with low TE, particularly with both GGA and GGU repeat (Table S3).

Therefore, our results indicate that rG4 serves as a translation repressor, which agrees with

previous studies on individual rG4s[35–37]. To validate our identified translation-associated rG4

motifs, we conducted the experimental validation using dual luciferase reporter assay in

plants[12]. We fused the 5'UTRs containing our identified rG4 motif and the corresponding

disrupted rG4 motif with the luciferase reporter genes[12]. We then measured the corresponding

TEs in plants and observed a significant increase of up to 5.8-fold in the disrupted rG4 motif

compared to the TE in the native rG4 motif (Fig. 4g). These results indicate that our PlantRNA-

FM is also capable of identifying functional RNA tertiary structure motifs such as translation-

associated rG4 motifs throughout the transcriptome.

2. Discussion

In this study, we developed PlantRNA-FM, a high-performance and interpretable plant-

specific RNA FM. PlantRNA-FM (Fig. 1) is designed for understanding RNA sequence and

structure information rather than generation. This state-of-the-art model was specifically

designed based on the extensive plant RNA information from 1,124 plant species, thereby

capturing the remarkable diversity of plant RNA features. From the perspective of the dataset,

we have incorporated RNA sequence information of all the RNAs from the transcriptomes

across 1,124 plant species. We also incorporated the corresponding RNA annotation

information. The integration of RNA structure information in our PlantRNA-FM achieves

superior performance in RNA structure prediction tasks compared to other FMs (Fig. S1).

Regarding the model architecture, we adopted a fine-grained tokenization method with single-

nucleotide resolution. This contrasts with commonly used tokenization methods, such as byte

pair encoding (BPE) and k-mers, which rely on frequency-based tokenization and may

inadvertently fragment RNA structure motifs into arbitrary pieces. This strategy ensures the

precise extraction and preservation of RNA structure motifs as coherent units throughout the

pre-training process, thereby maintaining the integrity of crucial structure information.

Additionally, PlantRNA-FM integrates rotational position embedding (RoPE), a technique that

has proven effective in enhancing the modeling capabilities for long tokens in large FMs[38]. The

implementation of RoPE leads to a approximately 30% reduction in the number of parameters

in the embedding layer, consequently improving the efficiency of RNA tokenisation and

modeling.

The superior performance of PlantRNA-FM can be further demonstrated in the plant-

specific downstream tasks (Fig. 2a). Our PlantRNA-FM achieved the best *F1* scores of 0.974

and 0.958 for the genic region annotation in *Arabidopsis* and rice, while our PlantRNA-FM

also achieved much better performance in predicting TE compared to other FMs (Fig. 2b, 2c).

The outperformance of our PlantRNA-FM is likely due to the combination of both RNA

sequence and structure information in our pre-training dataset, highlighting the importance of

RNA structure, a key RNA feature, in regulating RNA biological processes.

Notably, we developed an interpretable framework for our PlantRNA-FM to explore

the RNA features within the 5' UTR that influence translation (Fig. 3a). Using the attention

contrast matrices, we found that the nucleotides in the regions close to the start codon affect

the translation the most, emphasizing the importance of positional information of functional

RNA motifs (Fig. 3b). In contrast to conventional meta-gene analysis, our PlantRNA-FM is

capable of providing positional information of RNA motifs across transcriptomes, which is

critical for biological regulatory functions. Furthermore, the Kozak sequence, an evolutionary

conserved translation-associated sequence motif across translation initiation sites was

successfully identified in both *Arabidopsis* and rice using our PlantRNA-FM (Fig. 3c, 3d). This

result successfully demonstrates the capability of our PlantRNA-FM in identifying the RNA

sequence motifs important to translation across the transcriptomes. By using an unsupervised

hierarchical clustering strategy to explore our attention contrast matrix, we further

systematically identified RNA secondary and tertiary structure motifs that are functionally

important to translation (Fig. 4a). Notably, we identified both high translation-associated and

low translation-associated RNA secondary structure motifs where their differences are mainly

in the strengths of the base pairs (Fig. 4b, 4c). This suggests that RNAs may adopt different

RNA structure motifs with diverse folding strengths in regulating biological processes such as

translation. In contrast to conventional meta-gene analysis, our PlantRNA-FM is capable of

delivering a comprehensive understanding of functional RNA motifs such as the type of RNA

motifs, the genic position of the RNA motifs, the positive or negative effects of the RNA motifs

on their functions, and the exact contributions of the RNA motifs to their functions. For

instance, high GC content in the 5' UTR has been shown to be anti-correlated with translation

efficiency[39–41]. However, these correlations are not able to facilitate understanding of which

type of regulatory motifs with high GC content repress translation. Here, our PlantRNA-FM

revealed diverse RNA structure motifs such as the RNA secondary structure motif with four

GC base pairs in the stem and rG4s, serving as low translation-associated RNA motifs. This

suggests the diversity of RNA regulatory motifs across the transcriptomes (Fig. 4b).

In summary, we have built the first interpretable RNA FM with both RNA sequence

and structure information. Our PlantRNA-FM was pre-trained using 1,124 plant transcriptomes.

We have demonstrated that our PlantRNA-FM is capable of identifying functional RNA motifs

such as translation-associated sequence and structure motifs across the transcriptomes.

Through our experimental validations, we have elucidated novel translation-associated RNA

motifs in plants. Our FM model can be extended to explore functional RNA motifs in other

kingdoms and investigate RNA motifs important for other biological functions such as RNA

decay and maturation. Our PlantRNA-FM is poised to transform the way we determine RNA

motifs for regulating gene expression, opening new horizons for programming RNA codes to

facilitate crop improvements and RNA-based applications.

3. Methods

3.1. Pre-training datasets curation

The plant transcriptome data used for pre-training PlantRNA-FM was obtained from

the one thousand plant transcriptomes project (1KP)[14]. Note that modeling genomic sequences

differs significantly from natural language modeling. For instance, while RNA sequences are

one-dimensional, they strictly follow biological genomic patterns and depend heavily on

certain structural characteristics. In contrast, natural language models are more resilient and

can tolerate linguistic errors such as typos and grammar mistakes. Thus, effective RNA

sequence curation is crucial to minimize the impact of noisy data and enhance modeling

performance. Specifically, our data curation protocol is as follows.

• **Sequence truncation and filtering**: We truncated RNA sequences exceeding 512

nucleotides to comply with the model's maximum length capacity and filtered out sequences

shorter than 20 nucleotides to eliminate noise, such as RNA fragment sequences.

• **RNA secondary structure annotation:** Given the significant impact of RNA

secondary structures on sequence function, we annotated the local RNA structures of all RNA

sequences using ViennaRNA (with parameters maxBPspan = 30)[25].

• **Annotation of CDS and UTR sequences:** After obtaining the assembled transcripts

and translated RNA regions from the dataset, we retrieve the CDS (translated RNA), 5' UTR,

and 3' UTR sequences (upstream and downstream of the translated RNA).

3.2. Model architecture

In this study, we developed PlantRNA-FM, a specialised language model based on the

transformer architecture (Fig. 1). PlantRNA-FM has 35 million parameters, including 12

transformer network layers, 24 attention heads, and an embedding dimension of 480. We

applied layer normalisation and residual connections both before and after the encoder block.

As our focus is on RNA understanding rather than generation, we only utilised the encoder

component of the transformer architecture. PlantRNA-FM is capable of processing sequences

up to 512 nucleotides in length, making it compatible with consumer-grade GPUs, such as the

Nvidia RTX 4090, with a batch size of 16. The model was trained on four A100 GPUs over a

period of three weeks, completing 3 epochs.

3.3. Pretraining strategies of PlantRNA-FM

313 To develop an RNA FM for exploiting all potential patterns within RNA sequences, we

314 investigated the biological domain knowledge of RNA sequences and propose three self-

315 supervised pre-training objectives to enhance the foundational model.

3.3.1. Pretraining with Masked nucleotides modeling

Inspired by the concept of masked language modelling (MLM) in NLP, we introduced

masked nucleotide modelling (MNM) for RNA sequences. This approach involves randomly

masking a portion of nucleotides and leveraging the model itself to reconstruct these masked

nucleotides. Note that the ability to accurately reconstruct masked nucleotides indicates that

the model is empowered with the capability of understanding RNA sequence. MNM

dynamically selects 20% of nucleotides for masking in each input sequence, as opposed to the

fixed 15% masking used in the classic MLM objective designed for shorter natural language

sentences. This increased masking ratio is chosen to enhance MNM's modeling capability,

considering that RNA sequences typically contain around one thousand bases. Specifically,

10% are replaced with a '<mask>' token, 5% with random nucleotides, and the remaining 5%

are left as is. This approach, which aims for token classification, employs cross-entropy as the

loss function to enhance the model's predictive accuracy for masked or replaced nucleotides.

The loss function $L_{MLM}(\theta)$ for MLM is defined as follows:

330
$$L_{MLM}(\theta) = -\frac{1}{|m|} \sum_{i \in m} \log p_\theta(x_i \mid x_{\backslash i}),$$

where θ and m are the parameter set inside the FM and the number of masked nucleotides.

$p_\theta(x_i \mid x_{\backslash i})$ indicates the probability of predicting the masked nucleotide x_i based on its

context $(x_{\backslash i})$.

3.3.2. Pretraining with RNA Structure Prediction

We hypothesise that effectively aligning RNA sequences with their corresponding

secondary structures is important during the pre-training phase. In practice, we annotated the

secondary structures within the 1KP dataset, which comprises 50 billion nucleotides. This

establishes a robust foundation for our model to recognise the critical role of secondary

structures. Based on these annotated data, we utilized cross-entropy as the loss function to

predict the RNA secondary structure:

341
$$L_{SSP}(\theta) = -\sum_{i=1}^{N} \sum_{c=1}^{C} y_{i,c} \log p(y_{i,c} \mid x; \theta),$$

where N is the length of the RNA sequence, i.e., the total number of nucleotides in the

sequence; C denotes the number of prediction for each nucleotide (e.g., '(', ')', '.'); $y_{i,c}$ is the

prediction of the i-th nucleotide c, and $p(y_{i,c} \mid x; \theta)$ is the probability predicted by the model

parameterised by θ. $L_{SSP}(\theta)$ is the loss function that quantifies the discrepancy between the

model's predicted probabilities for each nucleotide's secondary structure and the actual

structure, with the aim of minimising this loss to improve the model's accuracy in secondary

structure prediction.

3.3.3. Pretraining with RNA annotation prediction

RNA sequences exhibit significant variation across different regions, each serving

distinct functions within an organism. Beyond the two aforementioned training objectives, the

third one focuses on classifying regions within RNA sequences. The loss function is as follows:

$$L_{CLS}(\theta) = -\sum_{i=1}^{N} \sum_{r=1}^{R} y_{i,r} \log p(y_{i,r} \mid x; \theta),$$

where N is the length of the RNA sequence, i.e., the total number of nucleotides or segments

considered for classification. R represents the number of region categories we are classifying,

including CDS, 3' UTR, and 5' UTR. $y_{i,r}$ is the prediction of the i-th nucleotide r.

$p(y_{i,r} \mid x; \theta)$ is the probability predicted by the model, with parameters θ, for the i-th

nucleotide given the RNA sequence x. $L_{CLS}(\theta)$ is the cross-entropy loss function aimed at

training the model to identify different regions.

3.4. Fine-tuning of downstream tasks

After the pre-training phase, our FM can be fine-tuned to adapt to various downstream

tasks. The fine-tuning phase consists of three steps. First, we gathered an annotated dataset

specific to each downstream task, which consists of sequences and their corresponding labels.

Note that we pre-sliced any sequences that exceed the model's maximum length, to ensure

compatibility. Next, using the pre-trained FM as a starting point, we adapted the output layer

to accommodate the requirements of RNA modelling tasks, which may include outputting

sequences, labels, or scalar values. Finally, the training and inference processes are tailored to

the demands of each downstream task by selecting task-specific optimisers, loss functions, and

tuning hyperparameters to achieve optimal performance. The source code for our training and

inference can be found in our repository.

3.5. Polysome profiling mapping and data processing

Raw polysome profiling sequencing data for *A. thaliana* were obtained from published

research[12]. For rice, we performed polysome-seq using the same protocol as *Arabidopsis*[8]. The

genomes and annotation files of *O. sativa* and *A. thaliana* were obtained from Phytozome v13

with version of *Oryza sativa* v7.0 and TAIR10[44]. After extracting the transcriptome sequence

through the reference genome and annotation files, clean polysome profiling and RNA-Seq

reads were mapped to the reference transcriptome using HISAT2 and followed by library

normalisation and quantification using DESeq2[45,46]. Next, genes with an RPKM of less than

1 were removed, and the TE of each gene was calculated by dividing the polysome-associated

RNA levels (polysome profiling RNA-seq) by the corresponding RNA levels (RNA-Seq)[12].

Subsequently, the dataset was classified as high or low TE, using the mean plus or minus the

standard deviation as a threshold, and were respectively assigned the labels 1 and 0 for high

and low TE.

3.6. RNA structure motif identification approach

3.6.1. Extraction of the attention contrast matrix

To facilitate better model interpretation, we created two additional models. One is the

true model, denoted as PlantRNA-FM(+), trained using the real TE labels, while the other one

is the background model, PlantRNA-FM(-), altered using the same dataset but with randomly

assigned labels. Specifically, we fine-tuned the pre-trained PlantRNA-FM (+) and (-) on each

dataset for 100 epochs, using regular hyperparameter settings. To avoid overfitting, we

employed an early stopping strategy to terminate the fine-tuning process when the best *F1* score

remained unchanged for 30 epochs. Once the fine-tuning was completed, we used the fine-

tuned models to predict each dataset and derive the raw attention score matrices corresponding

to each RNA sequence. Since the raw attention score matrices are five-dimensional, we

reshaped them through average-based downsampling to generate attention contrast matrices.

Finally, we subtracted the attention contrast matrices of PlantRNA-FM (+) from those of

PlantRNA-FM (-). Furthermore, we padded any negative values in the attention contrast

matrices with zeros for better visualisation.

3.6.2.   Generation of the RNA structure motif seed library

To identify RNA structure motifs, we first generate a library of that contains RNA

structure motif seeds derived from RNA sequences across the transcriptomes. In this work, we

apply the Zuker algorithm from the Vienna RNA package to obtain all suboptimal RNA

structure foldings for each RNA in our dataset[35,36]. We restrict the length of the RNA structure

motifs to a maximum of thirty[48]. The folded RNA structures are then annotated using

"bpRNA". Subsequently, all RNA structure motifs are extracted to generate a seed library of

RNA structure motifs for the plant transcriptomes[26]. In order to obtain reliable RNA structure

motifs, we set the range of RNA structure stems from 4 to 7, and the loop length from 4 to 9.

3.6.3.   Identification of translation-associated RNA secondary structure motifs

From the previous step, we obtained all potential foldings of the RNA structure motif

in the 5' UTR and aligned them with the attention contrast matrix. For each RNA structure

motif, we evaluated it using a paired t-test to obtain a p-value. Then, we corrected the obtained

p-value using the Benjamini-Hochberg (BH) method. RNA structure motifs with p-values less

than 0.01 were considered significant and extracted as the high-attention RNA structure

motifs. Then we extracted their corresponding RNA sequence and converted them into

numerical matrices using the one-hot encoding method. Subsequently, we applied an

unsupervised hierarchical clustering strategy to classify the nucleotides corresponding to the

positions of the RNA structure pairs into 2 to 100 clusters[49]. For each cluster containing a

minimum of 30 high-attention RNA structure motifs, the significance was assessed using

Fisher's exact test. RNA motifs with an odds ratio over 1 and a p-value below 0.05 were

identified as high translation-associated motifs. On the contrary, those with an odds ratio less

than 1 and a p-value below 0.05 were associated with low TE. Additionally, we calculated the

mean information content of all bases, defined as the "Average Positional Information Content"

(APIC). RNA motifs with an APIC below 1.5 were excluded from further analysis.

3.6.4. Identification of translation-associated rG4s

425 We obtained all potential rG4 in rice from our G4Atlas database[33]. Next, we aligned

426 the rG4 sequences with the corresponding attention contrast matrix and employed the paired t-

427 test to assess the statistical significance. For each length of rG4, we adjusted its p-value using

428 the Benjamini-Hochberg (BH) correction method and selected rG4s with a p-value less than

429 0.01 as the high attention rG4s.

4. Data availability

431 The polysome-seq sequence data of *A. thaliana* was obtained from the Sequence Read Archive

432 (SRA) (https://www.ncbi.nlm.nih.gov/sra) under BioProject ID number PRJNA762705[8]. The

433 raw sequence data of *O. sativa* has been deposited in the Sequence Read Archive (SRA)

434 (https://www.ncbi.nlm.nih.gov/sra) under BioProject ID number PRJNA1112739.

5. Code availability

436 The source code of this study is freely available at Huggingface

437 (https://huggingface.co/yangheng/PlantRNA-FM).

6. References

1. Piao, M., Sun, L. & Zhang, Q. C. RNA Regulations and Functions Decoded by

Transcriptome-wide RNA Structure Probing. *Genomics, Proteomics & Bioinformatics* **15**,

267–278 (2017).

2. Komatsu, K. R. *et al.* RNA structure-wide discovery of functional interactions with

multiplexed RNA motif library. *Nature Communications* **11**, 6275 (2020).

3. Espah Borujeni, A., Channarasappa, A. S. & Salis, H. M. Translation rate is controlled by

coupled trade-offs between site accessibility, selective RNA unfolding and sliding at

upstream standby sites. *Nucleic Acids Research* **42**, 2646–2659 (2014).

4. Gorochowski, T. E., Ignatova, Z., Bovenberg, R. A. L. & Roubos, J. A. Trade-offs between

tRNA abundance and mRNA secondary structure support smoothing of translation

elongation rate. *Nucleic Acids Research* **43**, 3022–3032 (2015).

5. Mortimer, S. A., Kidwell, M. A. & Doudna, J. A. Insights into RNA structure and function

from genome-wide studies. *Nat Rev Genet* **15**, 469–479 (2014).

6. Kozak, M. An analysis of vertebrate mRNA sequences: intimations of translational control.

Journal of Cell Biology **115**, 887–903 (1991).

7. Ding, Y. *et al.* In vivo genome-wide profiling of RNA secondary structure reveals novel

regulatory features. *Nature* **505**, 696–700 (2014).

8. Yang, X. *et al.* RNA G-quadruplex structure contributes to cold adaptation in plants. *Nat*

Commun **13**, 6224 (2022).

9. Xu, B. *et al.* Recent advances in RNA structurome. *Sci. China Life Sci.* **65**, 1285–1324

(2022).

10. Yang, M. *et al.* Intact RNA structurome reveals mRNA structure-mediated regulation

of miRNA cleavage in vivo. *bioRxiv* 2019.12.21.885699 (2020) doi:10/ghccqf.

11. Yang, M. *et al.* In vivo single-molecule analysis reveals COOLAIR RNA structural

diversity. *Nature* 1–6 (2022) doi:10.1038/s41586-022-05135-9.

12. Xiaofei Yang & Haopeng Yu. Wheat in vivo RNA structure landscape reveals a

prevalent role of RNA structure in modulating translational subgenome expression

asymmetry. 26 (2021).

13.   Deng, H. *et al.* Rice In Vivo RNA Structurome Reveals RNA Secondary Structure

Conservation and Divergence in Plants. *Molecular Plant* **11**, 607–622 (2018).

14.   One Thousand Plant Transcriptomes Initiative. One thousand plant transcriptomes and

the phylogenomics of green plants. *Nature* **574**, 679–685 (2019).

15.   Cao, J. *et al.* High-throughput 5′ UTR engineering for enhanced protein production in

non-viral gene therapies. *Nat Commun* **12**, 4138 (2021).

16.   Consens, M. E. *et al.* To Transformers and Beyond: Large Language Models for the

Genome. Preprint at https://doi.org/10.48550/arXiv.2311.07621 (2023).

17.   Zhou, Z. *et al.* DNABERT-2: Efficient Foundation Model and Benchmark For Multi-

Species Genome. Preprint at https://doi.org/10.48550/arXiv.2306.15006 (2023).

18.   Dalla-Torre, H. *et al.* The Nucleotide Transformer: Building and Evaluating Robust

Foundation Models for Human Genomics. 2023.01.11.523679 Preprint at

https://doi.org/10.1101/2023.01.11.523679 (2023).

19.   Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a

language model. *Science* **379**, 1123–1130 (2023).

20.   Chu, Y. *et al.* A 5' UTR Language Model for Decoding Untranslated Regions of

mRNA and Function Predictions. 2023.10.11.561938 Preprint at

https://doi.org/10.1101/2023.10.11.561938 (2023).

21.   Hallee, L., Rafailidis, N. & Gleghorn, J. P. cdsBERT - Extending Protein Language

Models with Codon Awareness. *bioRxiv* 2023.09.15.558027 (2023)

doi:10.1101/2023.09.15.558027.

22.   Chen, K. *et al.* Self-supervised learning on millions of primary RNA sequences from

72 vertebrates improves sequence-based RNA splicing prediction. *Briefings in*

Bioinformatics **25**, bbae163 (2024).

23.   Yang, X., Yang, M., Deng, H. & Ding, Y. New Era of Studying RNA Secondary

Structure and Its Influence on Gene Regulation in Plants. *Front. Plant Sci.* **9**, (2018).

24.   Zhang, H., Chung, B. Y.-W., Wang, Z. & Ding, Y. Editorial: Plant RNA structure.

Front. Plant Sci. **14**, (2023).

25.   Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).

26.   Danaee, P. *et al.* bpRNA: large-scale automated annotation and analysis of RNA

secondary structure. *Nucleic Acids Research* **46**, 5381–5394 (2018).

27.   Sloma, M. F. & Mathews, D. H. Exact calculation of loop formation probability

identifies folding motifs in RNA secondary structures. *RNA* **22**, 1808–1818 (2016).

28.   Tan, Z., Fu, Y., Sharma, G. & Mathews, D. H. TurboFold II: RNA structural

alignment and secondary structure prediction informed by multiple homologs. *Nucleic*

Acids Research **45**, 11570–11581 (2017).

29.   Hardy, E. C. & Balcerowicz, M. Untranslated yet indispensable—UTRs act as key

regulators in the environmental control of gene expression. *Journal of Experimental*

Botany erae073 (2024) doi:10.1093/jxb/erae073.

30.   Dever, T. E., Ivanov, I. P. & Hinnebusch, A. G. Translational regulation by uORFs and

start codon selection stringency. *Genes Dev.* **37**, 474–489 (2023).

31.   Evfratov, S. A. *et al.* Application of sorting and next generation sequencing to study

5´-UTR influence on translation efficiency in Escherichia coli. *Nucleic Acids Research* **45**,

3487–3502 (2017).

32.   Lyu, K., Chow, E. Y.-C., Mou, X., Chan, T.-F. & Kwok, C. K. RNA G-quadruplexes

(rG4s): genomics and biological functions. *Nucleic Acids Research* **49**, 5426–5450 (2021).

33.   Yu, H., Qi, Y., Yang, B., Yang, X. & Ding, Y. G4Atlas: a comprehensive

transcriptome-wide G-quadruplex database. *Nucleic Acids Research* **51**, D126–D134

(2023).

34.   Song, J., Perreault, J.-P., Topisirovic, I. & Richard, S. RNA G-quadruplexes and their

potential regulatory roles in translation. *Translation* **4**, e1244031 (2016).

35.   Kumari, S., Bugaut, A., Huppert, J. L. & Balasubramanian, S. An RNA G-quadruplex

in the 5′ UTR of the NRAS proto-oncogene modulates translation. *Nat Chem Biol* **3**, 218–

221 (2007).

36.   Beaudoin, J.-D. & Perreault, J.-P. 5′-UTR G-quadruplex structures acting as

translational repressors. *Nucleic Acids Research* **38**, 7022–7036 (2010).

37.   Jia, L. *et al.* Decoding mRNA translatability and stability from the 5′ UTR. *Nat Struct*

Mol Biol **27**, 814–821 (2020).

38.   Su, J. *et al.* RoFormer: Enhanced transformer with Rotary Position Embedding.

Neurocomputing **568**, 127063 (2024).

39.   Araujo, P. R. *et al.* Before It Gets Started: Regulating Translation at the 5′ UTR.

International Journal of Genomics **2012**, e475731 (2012).

40.   Leppek, K., Das, R. & Barna, M. Functional 5′ UTR mRNA structures in eukaryotic

translation regulation and how to find them. *Nat Rev Mol Cell Biol* **19**, 158–174 (2018).

41.   van der Velden, A. W. & Thomas, A. A. M. The role of the 5′ untranslated region of an

mRNA in translation regulation during development. *The International Journal of*

Biochemistry & Cell Biology **31**, 87–106 (1999).

42.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

596, 583–589 (2021).

43.   Verkuil, R. *et al.* Language models generalize beyond natural proteins.

2022.12.21.521521 Preprint at https://doi.org/10.1101/2022.12.21.521521 (2022).

44.   Goodstein, D. M. *et al.* Phytozome: a comparative platform for green plant genomics.

Nucleic Acids Research **40**, D1178–D1186 (2012).

45.   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low

memory requirements. *Nat Methods* **12**, 357–360 (2015).

46.   Love, M. I., Huber, W. & Anders, S. *Moderated Estimation of Fold Change and*

Dispersion for RNA-Seq Data with DESeq2. http://biorxiv.org/lookup/doi/10.1101/002832

(2014) doi:10.1101/002832.

47.   Zuker, M. On Finding All Suboptimal Foldings of an RNA Molecule. *Science* **244**,

48–52 (1989).

48.   Fish, L. *et al.* A prometastatic splicing program regulated by SNRPA1 interactions

with structured RNA elements. *Science* **372**, eabc7531 (2021).

49.   Steinbach, M., Karypis, G. & Kumar, V. A Comparison of Document Clustering

Techniques. (2000).

552
553

7. Funding

This work was supported by National Key Research and Development Program of

China [2023YFA0913500] (HZ); National Key Research and Development Program of China

[2021YFF1000900] (HZ); National Natural Science Foundation of China [32170229] (HZ);

Fundamental Research Funds for the Central Universities [2412023YQ005] (HZ); the China

Scholarship Council [No.202206620047] (WS); the United Kingdom Biotechnology and

Biological Sciences Research Council (BBSRC) [BB/X01102X/1] (HY, YD); European

Research Council (ERC) [selected by the ERC, funded by BBSRC Horizon Europe Guarantee

[EP/Y009886/1] (YD); Human Frontier Science Program Fellowship [LT001077/2021-L]

(HY); UKRI Future Leaders Fellowship [MR/S017062/1, MR/X011135/1] (KL); Kan Tong

Po International Fellowship [KTP\R1\231017] (KL); Amazon Research Award (KL) and

National Natural Science Foundation of China [62376056, 62076056] (KL).

566
567

8. TABLE AND FIGURES

570 **Fig. 1. Schematic overview of the Pre-training Phase of PlantRNA-FM.** The pre-training

571 dataset comprises transcriptomic sequences from 1,124 plant species, consisting of

572 approximately 25.0M RNA sequences and 54.2B RNA bases. The green dots on the global

573 mean temperature map represent the geographical distribution of these plant species across the

574 world.

Fig. 2. Fine-tuning PlantRNA-FM on plant-specific datasets. a, Overview of fine-tuning PlantRNA-FM for RNA genic region annotation prediction and RNA translation efficiency (TE) prediction tasks. *A. thaliana* and *O. sativa* were selected as representative plant species. For the RNA genic region annotation prediction task, RNA sequences from these two species were included, along with three labels: 5' UTR, CDS, and 3' UTR. For the RNA TE prediction task, 5'UTR sequences from these two species were included, along with TE labels (high TE and low TE). b, c, Comparison of the model performance of different pre-trained models on RNA genic region annotation prediction and RNA TE prediction tasks. The error bars represent the standard deviation of the *F1* scores obtained from three fine-tuning replicates.

Fig. 3. Our model interpretable framework reveals translation-associated RNA features.

a, Schematic of the model interpretability approach. b, Transcriptome-wide attention contrast scores. The -1 position represents the first site upstream of the AUG. Different species are distinguished by colours. c, d, The information content of the 10 high-attention bases closest to the AUG start codon.

Fig 4. RNA structure motif identification approach reveals translation-associated RNA structure motifs. a, Overview of the RNA structure motif identification approach. RNA structures are predicted using RNAfold with a maximum length of 30 nucleotides to obtain RNA structure seeds. Predicted RNA G-quadruplexes were obtained from the G4Atlas database. b, c, Schematic diagram of high translation-associated RNA structure motifs and low translation-associated RNA structure motifs. Sequence logos show the information content of each nucleotide, with semicircles connecting paired bases. APIC stands for average positional information content. The *p-value* is derived from Fisher's exact test. d,e,f,g, Experimental validation of high and low translation-associated RNA structure motifs and low translation-associated RNA G-quadruplex. The bar plot represents the translational efficiency of the original (WT) and RNA structure-mutated (Mut) constructs from the dual luciferase reporter assay in plants. It represents the change from high translation-associated RNA structure motifs

to low translation-associated RNA structure motifs (d), the change from low translation-

associated RNA structure motifs to high translation-associated RNA structure motifs (e), the

complete disruption of low translation-associated RNA structure motifs (f), and the complete

disruption of low translation-associated rG4 (g). * indicates $P < 0.05$, ** indicates $P < 0.01$, by

Student's t-test, n = 3, error bars indicate se.

Table 1. Comparison of *F1* scores achieved by different pre-trained models on benchmark

datasets.

Tasks	Species	PlantRNA-FM	cdsBERT	DNABERT-2	Nucleotide Transformer	ESM2
RNA genic region annotation prediction	*A. thaliana*	0.974±0.003	0.254±0.003	0.602±0.001	0.635±0.002	0.639±0.008
	O. sativa	0.958±0.006	0.252±0.017	0.580±0.015	0.635±0.013	0.665±0.010
RNA translation efficiency prediction	*A. thaliana*	0.735±0.003	0.359±0.010	0.346±0.001	0.637±0.010	0.617±0.008
	O. sativa	0.737±0.004	0.359±0.010	0.627±0.012	0.631±0.020	0.649±0.011

616