CAUSAL, PREDICTIVE OR OBSERVATIONAL? DIFFERENT

UNDERSTANDINGS OF KEY EVENT RELATIONSHIPS FOR

ADVERSE OUTCOME PATHWAYS AND THEIR IMPLICATIONS ON

PRACTICE

Zheng Zhou[1,✉], Jeroen L.A. Pennings[2], and Ullrika Sahlin[1]

[1] Center for Environmental and Climate Science, Lund University, Sweden

[2] National Institute for Public Health and the Environment, the Netherlands

✉ Correspondence: Zheng Zhou <zheng.zhou@cec.lu.se>

1   INTRODUCTION

1.1 Background

Regulatory toxicology and environmental risk assessment in the twenty-first century need

to handle the assessment of ever-increasing numbers of chemicals, and at the same time

adhere to the 3R principles to reduce, refine and replace animal experiments (Pallocca et al.

2022). The Adverse Outcome Pathways (AOPs) methodology facilitates the extrapolation of

evidence at lower biological levels from animal and non-animal methods for the inference

of adverse outcomes at higher biological levels (Villeneuve et al. 2014a). An AOP is defined

as an evolving conceptual framework to organize existing knowledge pertaining to a

molecular initiating event (MIE), an apical adverse outcome of interest (termed adverse

outcome; AO), key measurable/ observable events (termed key events; KEs), and the

linkages among MIE, AO and KEs (termed key event relationships; KERs) (Villeneuve et al.

2014a). A list of definitions and abbreviations is presented in the supplementary

information (SI). The robustness and reliability of the extrapolation across events at

different biological levels therefore depends on the characterization and inference of KERs

(Villeneuve et al. 2024), which could be causal, observational and predictive.

1.2 Causal Inference and Its Challenges in Key Event Relationship Evaluation

Causal inference pursues the unbiased understanding of how one condition, the cause,

affects another condition, the outcome, free from the threats to internal validity (Guzelian

et al. 2005). Randomized experiments are the gold standard for causal inference (Guzelian

et al. 2005; James et al. 2015) as they can approximate unbiased counterfactual estimates

of treatment effects (Höfler 2005; Pearl 2009). Toxicological studies typically implement

randomized experiments in controlled laboratory conditions, which reveal unbiased

estimates of the association between the randomized assignment (e.g. dose) to the

compound of interest and the experimental outcome (e.g. response), thereby establishing a

causal relationship(Höfler 2005). The mechanism of randomized controlled trials is

reviewed in the SI.

When randomized experiments are not feasible, inference could rely on observational data.

Quasi-experimental methods involve selecting subgroups from observational data to

approximate unbiased average treatment effects, assuming specific conditions are met

(Angrist and Pischke 2009). These methods have been employed in various fields such as

economy, social science, psychology and epidemiology (Angrist and Pischke 2009; Bagiella

et al. 2015; Fréchette et al. 2022). Observational data can also be evaluated using evidence-

based approaches. The Bradford Hill considerations (Hill 1965), an early evidence-based

framework for causal inference on observational, epidemiological data, has been modified

to infer causal estimates in various contexts (Guzelian et al. 2005). For example, the

evolved, rank-ordered Hill considerations for mode of action human relevance framework

can evaluate the weights of evidence supporting hypothesized KERs in modes of action

(MOAs) (Meek et al. 2014). However, neither quasi-experimental methods nor evidence-

based frameworks fully guarantee unbiased estimates of causality. Therefore, results based

on observational data have lower reliability compared to those from randomized

experiments, which are termed as "causal estimates" in fields such as econometrics and

social science. Within this article, we use the term "observational" to distinguish causal

estimates from counterfactual, causal findings, emphasizing the observational nature of the

data.

Despite the involvement of experimental toxicological data, there could be challenges to

obtain the empirical evidence needed for causal inference of KERs (Spînu et al. 2022). The

association between the dose and an outcome is causal, while the association between two

outcomes (KEs) is not necessarily causal, due to the inability to randomize upstream KEs.

For example, consider two KEs observed from the same dose-response experiment, with a

hypothetical KER between them (Figure 1). This experiment generates two pairs of dose-

response data, one for KE_{up} and one for KE_{down} respectively. While this setup provides

unbiased estimates of the dose effects on KE_{up} and KE_{down} (Figure 1 left panel), KE_{up} itself

is the consequence of the treatment and thus not randomized among subjects. The lack of

randomization in KE_{up} implies that the estimates of the KER may not be causal (Figure 1

right panel). Direct evaluation of observed changes in KE_{down} conditional on KE_{up} is

vulnerable to the threats to internal validity (Lipton and Ødegaard 2005), such as omitted

variable bias from confounding. An instance of confounding bias is that, even if KE_{up} does

not actually cause change in KE_{down}, simultaneous changes in both KEs might still occur

due to a third, unobserved confounding event. The relationship between KE_{up}, KE_{down} and

a potential confounder differs from the concept of non-adjacent KERs described in the

handbook (Villeneuve et al. 2024). This is because a confounder might affect both KE_{up} and

KE_{down} while a non-adjacent KER is when a third KE, which may not affect KE_{up}, is bridged

over to KE_{down}. The estimated effect of KE_{up} on KE_{down} would be biased without isolating

the confounding bias. For example, overdose of acetaminophen can cause hepatotoxicity

(Hinson et al. 2010) and nephrotoxicity (Mazer and Perrone 2008). Both liver and kidney

damage are suggested to be mediated by the depletion of glutathione (Jaeschke and McGill

2015; Mazer and Perrone 2008). The normal function of glutathione is involved in the

detoxification of N-acetyl-p-benzoquinone imine, a toxic metabolite of acetaminophen

(Casarett 2008). If the role of glutathione depletion is not properly accounted for, it could

introduce confounding bias to falsely suggest hepatotoxicity leads to nephrotoxicity upon

acetaminophen overdose.

Figure 1: Directed acyclic graph of key event relationship evaluation. RCT: randomized

controlled trials. Left shows that treatment effects are causal on both key events respectively;

right shows what is needed for a causal evaluation

1.3 Distinguishing Causal and Predictive Inference under the Context of Adverse Outcome

Pathways

Predictive models have been used in the application of traditional mechanistic study of

toxicology (Helma 2005), such as physiologically based pharmacokinetic models (Sager et

al. 2015). It also contributes to the development and application of alternatives to animal

methods (Khabib et al. 2022), such as quantitative structure activity relationships (Barber

et al. 2024; OECD 2014) and quantitative system toxicology (Bloomingdale et al. 2017).

The difference between causal and predictive inference has led to debate and confusion

(Young 2019). Causal inference falls under the framework of treatment effect evaluation

(Young 2019), whose goal is to approximate unbiasedness (Young 2019) using randomized

experiments and, when those are not possible, non-experimental methods (quasi-

experimental (Angrist and Pischke 2009), evidence-based frameworks). In contrast, the

objective of predictive inference is to make accurate predictions. The term "predictive"

reflects the scope of utility and application, and falls outside the treatment effect

framework (Young 2019). Predictions can be accurate even when biased by confounders

(Pearl 2009). For example, social-economic status is recognized as a major confounder to

the adverse neurological effects of lead exposure during early childhood (Bellinger et al.

1992). However, some early studies accurately predicted neurological deficits based on

dentine lead levels among children without adjusting for social-economic status (Herbert et

al. 1979).

The difference between "causal" and "predictive" terms can be further understood in

relation to mechanism and quantification. Mechanism implies that a causal relationship

defines a directed linkage from the cause to the outcome (Russo and Williamson 2007),

whereas the opposite is the error of reverse-causality. There is also a temporal

requirement in a causal relationship that the cause must occur prior to the outcome, as

emphasized through various iterations of Hill considerations (Becker et al. 2015; Cox 2018;

Hill 1965; Meek et al. 2014). In contrast, a predictive relationship can be non-directional,

i.e., knowing the temporal order between events is not necessary for prediction.

Quantification implies that a predictive relationship is validated on a strong statistical

strength of association. In contrast, a causal relationship does not necessarily include a

description of the magnitude of causality. Two events with a causal relationship could be

even statistically independent of each other, i.e., zero correlation. We present two examples

in the supplementary information for these two points respectively.

Furthermore, practical differences between predictive and the treatment effect

frameworks exist in many commonly used methods. Consider a linear regression with

multiple covariates, whose parameters are optimized using maximum likelihood

estimation. This regression can be used for both prediction and causal inference (Young

2019). When this regression is used as a predictive model, the goal is to improve the

model's predictive performance (Young 2019), through the selection and optimization of

the design matrix. When this regression is used as a causal model, bias adjustment is

performed on the data prior to model construction, such as control selection (Cinelli et al.

2022) and backdoor elimination (Pearl 2009). The goal is to use the adjusted data and

identify predictors that have significant coefficients (p values using maximum likelihood

estimation and null hypothesis testing) (Angrist and Pischke 2009). The same model works

differently for prediction and treatment effect evaluation. On the one hand, bias adjustment

on data may not improve predictive performance (Brookhart et al. 2010; Davies et al.

2013). Including predictors of local significance may not improve global performance

(Obermeyer and Emanuel 2016). On the other hand, optimizing the design matrix can not

reduce the bias without prior bias adjustment (Chen et al. 2021).

The distinction between causal, observational and predictive inference is particularly

relevant to context of toxicology and AOP development. This difference should be

considered in the assembly of KEs and KERs. Traditionally, KEs and KERs are characterized

among measured/ observed events from existing data, which is retrospective and

evidence-based. In contrast, recent *in silico* predictive models enable prospective forecast

of KEs and KERs for new compounds with minimal structural analysis (Hemmerich and

Ecker 2020). The distinction between causal, observational and predictive inference should

also be considered taking the difference between MOAs and AOPs. A major distinction

between MOAs and AOPs is that MOAs are chemical specific while AOPs are not (Villeneuve

et al. 2014a; Villeneuve et al. 2024). Evaluation of KERs under the MOA framework (Boobis

et al. 2006) involves the evolved MOA Hill considerations (Meek et al. 2014), where the

confidence in KER causality is crucial along the MOA of the compound of interest. In the

context of AOP development, it has been suggested that the causality is necessary between

the AO and compounds on all AOP models including predictive ones (Spînu et al. 2022).

However, it remains unclear whether it is necessary to require causality in all KERs in a

chemical agnostic AOP. Given the chemical agnostic principle of AOPs, the supporting data

for AOP development may include a wide range of compounds, making it common to

observe varying KE activities across compounds. Therefore, it is crucial to consider the

appropriate application of causal, observational and predictive inference in AOP

development.

1.4 Study Objectives

Given the difference between the terms "causal", "observational" and "predictive" and its

relevance to KER characterization, we hypothesize that there could be misuse of the terms

and it hinders the effective evaluation of KERs. To test this hypothesis, the objectives of this

study are to: 1) examine the use of terminology describing KERs across AOP

documentations and 2) investigate the adaptation of different terms in the development

and application of AOPs. Achieving these objectives may facilitate a continuous

development of AOP guidelines and a pragmatic implementation of AOP framework in

academic, regulatory and industrial contexts.

2   METHODS

2.1   Terms for Key Event Relationships

We performed a text analysis on two AOP documentations focused on setting the definitions, principles and guidelines for AOP development: AOP developer's handbook (Villeneuve et al. 2024) (abbreviated as the handbook) and OECD Guidance Document for the Use of Adverse Outcome Pathways in Developing Integrated Approaches to Testing and Assessment (OECD 2017) (abbreviated as OECD guidance thereafter). For a comprehensive analysis on the descriptions of KERs, manual searches were performed in the handbook and OECD guidance with the following keywords: "key event relationship" OR "KER" OR ((relationship OR link) AND "key event"). We also noted the sentences before and after the located keywords for context.

2.2   Adaptation of terminology

We searched journal articles published by May 2024 to identify how different terminology has been adapted in development and application of AOPs. Our search was performed using online scholarly databases such as Google Scholar and PubMed. We gave preferences to sources from the "Animal-free Safety assessment of chemicals: Project cluster for Implementation of novel Strategies" (ASPIS) research cluster, including: ONTOX (ONTOX 2024), RISK-HUNT3R (RISK-HUNT3R 2024) and PrecisionTox (PrecisionTox 2024). We prioritized information from ASPIS due to the alignment between this article's emphasis on AOPs and ASPIS's perspective on non-animal methodologies. These searches aimed to identify exemplary cases to illustrate typical scenarios where the terms describing KERs have been adapted, rather than to conduct a systematic review of all relevant literature.

The identified information was organized by how the method of adaptation fits into the

stages of AOP development and quantitative AOP (qAOP). AOP development consists of

three operationally defined stages (OECD 2017; Villeneuve et al. 2014a). In contrast, there

is no established guidance or consensus on the definition or stages of qAOP. Therefore,

acting phases for qAOP application were based on pragmatic objectives from published

literatures (Conolly et al. 2017; Perkins et al. 2019; Spinu et al. 2020), including prediction

of AO (Jeong et al. 2018), derivation of point of departure (POD) (Perkins et al. 2019), and

optimization of testing strategies (OECD 2017; Perkins et al. 2019).

3   RESULTS

3.1   Usage of terminology

KER descriptions in the OECD guidance are primarily based on the AOP developer's

handbook, with minor editorial changes for the use in integrated testing strategies.

Therefore, results of our text analysis focus on KER descriptions identified from the

handbook, highlighting the text examples located from the sections where the definitions

and principles of AOP development are set. Results show that with a clear distinction

between the terms "causal" and "predictive", the handbook's usage of them varies across

sections (Table 1). For instance, in the handbook Section 3 KER Description on page 25,

KERs are described in different ways: as predictive *"Within the AOP framework, the*

predictive relationships that facilitate extrapolation are represented by the KERs" or as

causal *"KERs are always described in the form of a directed relationship (one-way arrow)*

linking an upstream "causing" event to a downstream "responding" event". In some instances

KERs are described with both "causal" and "predictive", e.g., on page 6 Table 1 Definitions

of Key Terms that *"A scientifically-based relationship that connects one key event to another,*

defines a causal and predictive relationship between the upstream and downstream event".

Compared to instances where "causal" or "predictive" are used separately, the inclusion of

both terms suggests the consideration of dual requirements in KER description. The dual

requirements do not necessarily imply a conflict against separate use of terms. When both

causal and predictive inference are needed for KER description, one or both may be

emphasized at different stages of AOP development.

The dual requirements are also evident in the handbook's protocol for evaluating KER

supporting evidence. The handbook evaluates KER supporting evidence (Section 3G)

through 1) biological plausibility, 2) empirical evidence and 3) current quantitative

understandings. The handbook's guidance to assess biological plausibility (annex 1) is

adapted from the evolved MOA Hill considerations (Meek et al. 2014), which is

observational as mentioned in the Introduction of this paper. Empirical evidence is

assessed (handbook page 31 subsection ii) for predictive utility that *"assessment of changes*

in the upstream KE (KEupstream) leading to, or being associated with, a predictable

subsequent change in the downstream KE (KEdownstream)". Moreover, within the same

section, the uncertainty and inconsistency of empirical evidence is evaluated that *"while*

there are expected patterns of concordance that support a causal linkage between the KEs in

the pair, it is also helpful to identify experimental details that may explain apparent

deviations from the expected patterns of concordance". The handbook's evaluation of current

quantitative understandings of KERs, considering the weights of empirical evidence

supporting KER causality, is dedicated to prediction *"to define the precision with which the*

state of the downstream KE can be predicted from knowledge of the state of the upstream KE"

(page 34). Therefore, it seems that the handbook sets the definitions, descriptions and

protocols for evaluation of KERs based on a dual focus of causal and predictive inference.

The term "observational" or the more commonly recognized "causal estimates" does not

appear in the handbook, nor the OECD guidance.

Table 1: Occurrence of terminology describing KERs in two AOP documentations

		Terminology	
Location of KER description	Quote	Causal	Predictive
OECD guidance document	***OECD guidance document 2017***	**OECD guidance document**	
AOP definition	*a sequential chain of causally linked events at different levels of biological organisation*	√	
	AOP handbook 2024 version 2.7		
Definitions of key terms, KER , page 6 Table 1	*A scientifically-based relationship that connects one key event to another, defines a causal and predictive relationship between the upstream and downstream event*	√	√
Section 1, AOP development tip #3, page 13 line 448	*an AOP is acceptable when there are multiple KEs, causally linked to the MIE and AO*	√	
Section 3 KER Description, page 25 line 929	*Within the AOP framework, the predictive relationships that facilitate extrapolation are represented by the KERs.*		√
Section 3 KER Description, page 25 line 941	*KERs are always described in the form of a directed relationship (one-way arrow) linking an upstream "causing" event to a downstream "responding" event.*	√	
Section 3G Evidence Supporting this KER, Empirical	*biological plausibility, empirical support, and the current quantitative understanding of the KER are evaluated with regard to the predictive relationships/associations between defined pairs of KEs as a basis for considering WoE (Section 4)*		√

		Terminology	
Location of KER description	Quote	Causal	Predictive
Evidence, page 29 line 1126			
Section 3G Evidence Supporting this KER, Empirical Evidence, page 31 line 1167	*evidence relevant to assessment of changes in the upstream KE (KEupstream) leading to, or being associated with, a predictable subsequent change in the downstream KE (KEdownstream).*		√
Section 3G Evidence Supporting this KER, Uncertainties and Inconsistencies, page 32 line 1242	*while there are expected patterns of concordance that support a causal linkage between the KEs in the pair, it is also helpful to identify experimental details that may explain apparent deviations from the expected patterns of concordance.*	√	
Section 4B Assess the Essentiality of ALL KEs, Direct evidence, page 39, line 1470	*evidence that overexpression of the object of the KE prevents or impacts the downstream KEs in a manner consistent with its causal, and essential, role in the pathway.*	√	
Section 4C Evidence Assessment, Review the Empirical Support for Each KER, page 41, line 1565	*Empirical support entails consideration of experimental data in terms of the associations between KEs – namely dose-response concordance and temporal relationships between and across multiple KEs.*		√

236 **3.2   Adaptation of terminology in AOP development and qAOP application**

237 *Table 2: Examples of adaptation of KER terminology in AOPs and qAOPs from published*

238 *literature*

Stage	Characteristics	Causal	Observational	Predictive
Putative AOP development	Assembly of hypothesized KEs and KERs	Through targeted blocking experiment (Skarnes 2011)	Through WoE supporting KE essentiality and KER biological plausibility (Becker et al. 2015; Jensen et al. 2023)	Through statistical inference: network analysis (Villeneuve et al. 2018) and machine learning (Gadaleta et al. 2022)
	Assembly of partial AOP with incomplete linkage between MIE and AO		As a result of known gaps and uncertainties (Bal-Price et al. 2015; Becker et al. 2015)	
Qualitative AOP development	Assembly of KEs		Through describing possibility of measurement (Becker et al. 2015)	
	Assembly of KERs		Through describing WoE of empirical concordance supporting KERs (Becker et al. 2015); Through evaluation of KERs using quasi-experimental methods (Landfors et al. 2023)	
	Evaluation of AOP		Through overall WoE supporting the AOP: biological plausibility, essentiality and empirical evidence (Becker et al. 2015)	
Quantitative AOP development	Assembly of KEs		Through describing accuracy and precision of measurements (Becker et al. 2015)	
	Assembly of KERs		Through evaluating WoE for characterizing the	Same as in observational but

Stage	Characteristics	Causal	Observational	Predictive
			magnitude and/or duration of change in upstream KE needed to evoke change in downstream KE (Becker et al. 2015)	with a focus on predictive performance of response-response models (Becker et al. 2015); Through characterizing known gaps and uncertainties between MIE and AO: known modulating factors (Becker et al. 2015)
Quantitative AOP application (acting)	Prediction of AO			Based on KEs of good accuracy and precision of measurement (Jeong et al. 2018)
	Derivation of POD			For gene expression/ transcription (Song et al. 2023) and AO (Perkins et al. 2019)
	Optimization of testing strategy			Through integrated approaches to testing and assessment (OECD 2017)

KE: key event; KER: key event relationship; AO: adverse outcome; MIE: molecular initiating event; AOP: adverse outcome pathway; POD: point of departure; WoE: weight of evidence

Empty cell indicates no example identified for the corresponding adaptation

The handbook's dual requirement on both causal and predictive KERs provides a

comprehensive framework for AOP development. Examples identified from published

literature are categorized by which term among "causal", "observational" and "predictive"

was adapted, and displayed throughout stages (Villeneuve et al. 2014a) of putative,

qualitative and quantitative AOP development (Table 2).

Adapting the term "causal" in the strict sense would require experiments implementing

direct randomization of upstream KEs. However, no examples of practical implementation

could be identified in the literature. The alternative approach in published studies is

targeted blocking experiments to infer the impact of KE_{up} on KE_{down} that a reduction or

complete absence of a KE_{down} when KE_{up} is blocked (Meek et al. 2014). Empirical evidence

from targeted blocking experiments are direct support and contribute to a high level of

confidence for evaluating the essentiality of KEs (Meek et al. 2014; Villeneuve et al. 2014b;

Villeneuve et al. 2024). The development and application of qAOPs often assumes that KEs

are essential (Jensen et al. 2023) and KERs are causal (Burgoon et al. 2020; Gou et al.

2023).

Observational analyses of KEs, KERs and overall AOP could involve weight-of-evidence

(WoE) assessments of supporting evidence and quasi-experimental assessments. The

evolved MOA Hill considerations (Meek et al. 2014) are adapted in the handbook and the

OECD guidance for the WoE assessment of the essentiality of KEs , the biological

plausibility of KERs, the concordance of KER empirical evidence and the overall AOP level

of confidence (OECD 2017; Villeneuve et al. 2024). Best practice has been recommended

for the adapted WoE assessments (Villeneuve et al. 2014b) to categorize confidence levels

as high, moderate or low (Becker et al. 2015). Characterizing known gaps and uncertainties

that modulate incomplete linkages between MIE and AO could further enhance the weights

of evidence in the overall AOP (Bal-Price et al. 2015; Villeneuve et al. 2024). When high

WoE is achieved, qualitative and quantitative descriptions of KE measurement methods

and their accuracy and precision are crucial (Villeneuve et al. 2014b). These descriptions

support the characterization and calibration of predictive modeling of relationships among

the sequence of KEs in the overall AOP (Becker et al. 2015). In addition to WoE

assessments, the instrumental variable method, a type of quasi-experimental design, shows

promise in providing strong evidence supporting KER causality (Landfors et al. 2023).

Predictive models are primarily involved in the putative and quantitative stages (OECD

2017). Assembly of KEs and KERs in the putative stage can be derived using machine

learning (Gou et al. 2023; Hemmerich and Ecker 2020) or probabilistic graphical network

models (Villeneuve et al. 2018). KEs could be screened from a pool of candidate molecular

and cellular events using feature selection techniques (Perkins et al. 2019). Similar

methods have been applied to screen for genes and genomic modules associated with

significant differential gene expressions (Callegaro et al. 2021; Hemmerich and Ecker

2020). Topological metrics are evaluated to identify nodes and edges that best improve

overall predictive performance on the AO as KEs and KERs, respectively (Villeneuve et al.

2018). Graphical network models might consist of deterministic and probabilistic nodes,

and undirected, directed, acyclic edges, that have been involved in existing studies of AOP

development and application (OECD 2017; Perkins et al. 2019; Villeneuve et al. 2018).

Quantitative AOP development focuses on the derivation of response-response models

[Villeneuve2024]. Studies of qAOP typically assume the causality of KERs has been

established (Burgoon et al. 2020; Gou et al. 2023; Jensen et al. 2023), supported by the

essentiality of KEs, the biological plausibility of KERs and the concordance in KER empirical

evidence. Furthermore, to assure the reliability of qAOPs within relevant applicability

domains, the handbook includes a guidance (annex 2) to evaluate the WoE of qAOPs by not

only predictive performance, also considerations for known modulating factors, feedback

loops and generalizability across the applicability domains. Examples of possible

regulatory application of qAOPs include prediction of AO given MIE (Jeong et al. 2018),

derivation of omics (Song et al. 2023) and *in vitro* (Perkins et al. 2019) PODs, and

optimization of testing strategies (OECD 2017).

In summary, the handbook's nuanced usage of terms is reflected in AOP development that

focus of adaptation varies across stages of AOP development. Given the challenge of

obtaining experimental evidence for achieving causality, targeted block experiments offer a

practical approach to balance weights of evidence and feasibility. Best practice (Becker et

al. 2015; Villeneuve et al. 2014b) has been recommended to implement observational

inference using WoE assessments based on the evolved MOA Hill considerations (Meek et

al. 2014). The stage of putative AOP development has a dual focus on observational and

predictive evaluations. The qualitative stage focuses on the WoE of evidence supporting

KEs, KERs and the overall AOP. The stage of qAOP application and application involves

various predictive approaches, including machine learning, network and response-

response models.

4   DISCUSSIONS

The nuanced terminology usage in the handbook and approaches of adaptation reflect the

complex nature of AOP development. Our results suggest several issues that may warrant

further discussion, including: 1) a mismatch between the terminology in guideline and the

methodology in practice, and 2) considerations of terminology for interdisciplinary

communication.

4.1   Mismatch between methodology and terminology

Some putative AOP studies inferred causality of KERs based on predictive models (Gou et

al. 2023; Jeong et al. 2018). We argue the use of predictive models in these studies

supported the predictive but not the causal aspect of KERs, highlighting a mismatch

between the treatment effect framework and predictive models. This mismatch could be

demonstrated through possible misinterpretations of probabilistic network models as

causal. Probabilistic network models (Koller and Friedman 2009) are based on directed

acyclic graphs (DAGs). DAGs are commonly involved in causal reasoning and designing of

structural equation modeling (Greenland et al. 1999), e.g. to approximate unbiasedness by

quasi-experimental methods. But the use of a DAG is not a guarantee for having a causal

model. It depends on context, if a model involving DAGs can provide unbiased average

treatment effects for causal inference. In Bayesian modelling, Directed acyclic graphs are

used to decompose the joint probability distribution of variables and parameters into a

product, with no implication of causality. A Bayesian network is usually referred to a

probabilistic network model with binary or categorical nodes. Bayesian networks comprise

a wide range of applications, where the network can be inferred from multivariate data,

e.g. in machine learning, or given as representative of causal relationships where the

probabilities or conditional probability distribution are informed by expert judgment,

e.g. in decision analysis. Bayesian networks have been applied to parameterize models

where confounding biases are adjusted using quasi-experimental methods such as the

backdoor criterion (Pearl 2009) and instrumental variables (Shingaki et al. 2021). With the

popularity of Bayesian networks (Kaikkonen et al. 2021; Spinu et al. 2020) comes

misconceptions of to what extent a Bayesian network indicates causality, which might

come from the use of a DAG. There are studies where results of a Bayesian network were

interpreted as causal without employing bias reduction (Chen and Pollino 2012; Jeong et al.

2018; Zou et al. 2010). For evaluating the uncertainty in a Bayesian network, beyond the

inherent uncertainty embedded within the probability model, it is crucial to consider the

process and context in which the Bayesian network is applied (Sahlin et al. 2021).

This mismatch is particularly crucial in the stage of putative AOP development, where the

assembly of KEs and KERs could be achieved through either causal inference or statistical

modeling. The reason is, although predictive inference of KEs and KERs is optional in the

recommended best practice for the putative stage (Villeneuve et al. 2014b), an AOP would

be incomplete without considerations on the essentiality of KEs, biological plausibility and

empirical support of KERs. Therefore, misrepresenting results from predictive models as

causal may falsely imply that the dual requirements have been fulfilled in the putative

stage, which could risk overlooking causal and observational assessment in the qualitative

stage.

Furthermore, this mismatch also highlights the importance of integrating predictive

models with the evidence from causal or observational inference in the qAOP stage.

Current qAOP development often assumes that causality of KERs is well-supported in

established AOPs (Burgoon et al. 2020; Gou et al. 2023; Jensen et al. 2023). Yet there are

limited approaches to propagate the assumed causality in predictive models. One potential

approach is the use of Bayesian methods to integrate known causality of KERs and prior

knowledge on KEs in qAOP (Spînu et al. 2022). Another approach involves incorporating

causal KERs on the target/ utility/ loss function of predictive models through

regularization techniques such as Lasso (Tibshirani 1996). While regularization has been

applied in in silico model to mitigate overfitting (Hemmerich and Ecker 2020), there has

been limited exploration of its use in adjusting the target/ utility/ loss function on known

causality. This may be due to concerns that regularizing on causality may penalize

predictive performance. However, it is crucial to re-emphasize that AOPs are only complete

with both causal and predictive characteristics (Becker et al. 2015) and good predictive

performance alone is insufficient for robust qAOP development. According to the

handbook's guideline on the WoE assessment of qAOPs (annex 2), known modulating

factors and feedback loops need to be considered along with predictive performance for a

qAOP to achieve high level of confidence (Villeneuve et al. 2024).

4.2   Careful considerations of terminology in AOP guidance

We would like to initiate a discussion for careful considerations on the usage of terms

"causality" and "essentiality" to facilitate interdisciplinary communication and

collaboration.

The use of term "causal" needs to be considered because the frameworks for evaluating

causality in the handbook (Villeneuve et al. 2024) and for the use in integrated testing

strategies (OECD 2017) may not necessarily conclude causality in the same sense employed

in toxicology and generic scientific reasoning. Even the recommended best practice (Becker

et al. 2015; Villeneuve et al. 2014b) may be insufficient to achieve unbiased estimates of KE

essentiality and KER causality. This is not a critique of current established AOP guidelines

and practice because, as shown by our results, the term "causal" is pronounced and utilized

in a consistent manner by the handbook, the OECD guidance and studies within the niche of

AOP development. Distinctive usage across fields of the same terminology is common.

However, adding clarifying annotations on the term "causal" could greatly facilitate

interdisciplinary communication and collaboration that potential bias may exist and that

the results could be "causal estimates" rather than "causal".

To demonstrate, the handbook's framework highlights the inclusion of direct and indirect

evidence and a prioritization of direct evidence in the assessment of essentiality of KEs and

biological plausibility of KERs. Direct evidence of KE essentiality (handbook section 4B,

Table 5) includes evidence from targeted blocking experiments, such as know-out

experiments (Skarnes et al. 2011). Knockout experiments pragmatically approximates

counterfactual inference and are more robust and reliable than observational evaluation of

multiple KEs (see supplementary information). However, results from targeted blocking

experiments are not always causal because the caveat of off-target issues applies in design

(Höijer et al. 2022). The instrumentation of blocking could be attributable to confounding

and uncontrolled mediation on downstream KEs (Skarnes et al. 2011). Unbiased causality

could be achieved only if the blocking methods induce blocking in the target KE_{up} but not

anything else that causes changes in KE_{down} or their modulating factors. If the instrument

to block KE_{up} activates a third, unobserved event that, e.g., increases the activity of KE_{down},

it is possible to observe no change in KE_{down}, because the reduction by blocking KE_{up}

could be canceled out by the activation from the third event.

In targeted knockout experiments, there could be at least two technical reasons leading to

the off-target issue. Firstly, the genetic target in knockout procedure is identified by

genome-wide association studies, which is association by definition. Confounders may exist

since the gene-protein associations are not always one-to-one matched (Olariu et al. 2021).

It is well known that the same protein can be coded by different genes or multiple copies of

the same gene at different loci (Campenhout et al. 2019). It is also not uncommon to

observe the synthesis of multiple proteins from the transcription of one gene. The

variational splicing of introns of one gene can be transcribed to different mRNAs, which

result in the production of multiple, distinctive proteins (De Conti et al. 2013). Targeted

mutation techniques have the potential to become quasi-experimental by implementing

bias adjustment methods such as instrumental variables (Landfors et al. 2023). Secondly,

assuming the target identification is free from confounding bias, unintended genetic

changes could be induced during gene editing (Höijer et al. 2022). Therefore, direct

evidence from targeted blocking experiments, as prescribed in the best practice, could be

insufficient to guarantee causality.

Indirect, non-experimental evidence is evaluated using a WoE framework based on the

evolved MOA Hill considerations (Meek et al. 2014). The evolved MOA Hill considerations

classifies weights of confidence into high, moderate or low rather than a dichotomous

judgement of causality. This is evident that the terms "causal", "causality" or etc. were

included only once (not including quotes) in the source paper (Meek et al. 2014) to

describe the consideration for KE essentiality that *"if following cessation of repeated*

exposure for various periods, effects are reverse, this constitutes relatively strong evidence

that key events are causal". Furthermore, the adaptations made by the handbook implement

no further bias adjustment and focus on the classification of WoE of supporting evidence.

In conclusion, the frameworks for KER evaluation in the handbook and recommended

practice could be insufficient to reach the causal requirement prescribed in the handbook.

We also would like to bring up a discussion on the use of the term "essential", particularly

in the essentiality of KEs. We argue that adding enhancing annotations on the term

"essential" could benefit its practical utility in the assessment of KE essentiality. It is crucial

to note that this discussion does not revolve around the distinction between mathematical

and toxicological context of "essential". The term "essential" has specialized mathematical

context and usage to describe the conditional relationship between two statements. To

claim mathematical essentiality of event A for event B, that is to say, if A does not occur,

then B cannot occur at all. The validation of mathematical essentiality requires a falsity test.

Any case where the occurrence of B without the occurrence of A falsifies essentiality.

Obviously, such a falsity test is practically difficult to perform in science (Graesser and Hu

2011), especially in biology and toxicology. Most biological events are the consequence of

one or several alternative, sufficient causes. The use of "essential" in toxicology is distinct

from the mathematical context, including the evolved MOA Hill considerations and AOP

development. Under the toxicological context, evaluation of essentiality considers the

impact of blocking or reversing KE_{up} on KE_{down} (Meek et al. 2014). A mathematically

essential event would cause complete absence in the consequence, while a toxicologically

essential event could describe a KE_{up} which, when blocked, prevent or significantly modify

KE_{down} (Meek et al. 2014; Villeneuve et al. 2024). The validation of mathematical

essentiality of KE_{up} for KE_{down} could be practically unfeasible. The guidelines for AOP

development needs to be a pragmatic framework that balances scientific rigor and

practicability to serve regulatory and industrial application in the real world. However, the

level of significance of modification/ reduction is subjective. Neither the evolved MOA Hill

considerations nor the handbook defines the level of modification in KE_{down} to which the

changes associated with blocking KE_{up} should be considered as significant and sufficient.

Furthermore, the level of modification/ reduction is diverse across systems, organs and

molecules by nature. For example, body weight change depends on a myriad of factors,

including time period, age, sex and baseline weight. It would be helpful to annotate the

handbook's defining question of KE essentiality by adding a description of the level at

which the modification in KE_{down} is both statistically significant and biologically sufficient.

Further research using the benchmark dose modeling methodology might be useful to

characterize the level of modification for KE essentiality.

5   CONCLUSIONS

This study aimed to explore the usage and adaptation of the terms "causal", "observational"

and "predictive" for KER description in AOP guidance documents. The results indicate that

KERs are described to fulfill both causal and predictive requirements across three stages of

AOP development, and distinct approaches are implemented to address different focuses of

adaptation across these stages. These findings suggest that the nuanced approach towards

the distinction between causal and predictive inference of KERs reveal a mismatch

between predictive models and causality in published studies of AOP development,

particularly pertaining to the putative stage. Further exploration is needed to integrate

causality in qAOP development and application. Further discussions could be helpful to

consider the handbook's usage of the terms "causal" and "essential" to facilitate more

comprehensive, transparent interdisciplinary communication and practical application of

AOP under the academic, regulatory and industrial contexts.

REFERENCE

Angrist JD, Pischke J-S. 2009. *Mostly harmless econometrics: An empiricist's companion.*

Princeton university press.

Bagiella E, Karamlou T, Chang H, Spivack J. 2015. Instrumental variable methods in clinical

research. The Journal of Thoracic and Cardiovascular Surgery 150:779–782;

doi:10.1016/j.jtcvs.2015.07.056.

Bal-Price A, Crofton KM, Sachana M, Shafer TJ, Behl M, Forsby A, et al. 2015. Putative

adverse outcome pathways relevant to neurotoxicity. Critical reviews in toxicology 45:83–

91; doi:10.3109/10408444.2014.981331.

Barber C, Heghes C, Johnston L. 2024. A framework to support the application of the OECD

guidance documents on (q)SAR model validation and prediction assessment for regulatory

decisions. Computational Toxicology 100305; doi:10.1016/j.comtox.2024.100305.

Becker RA, Ankley GT, Edwards SW, Kennedy SW, Linkov I, Meek B, et al. 2015. Increasing

scientific confidence in adverse outcome pathways: Application of tailored bradford-hill

considerations for evaluating weight of evidence. Regulatory Toxicology and Pharmacology

72:514–537; doi:10.1016/j.yrtph.2015.04.004.

Bellinger DC, Stiles KM, Needleman HL. 1992. Low-level lead exposure, intelligence and

academic achievement: A long-term follow-up study. Pediatrics 90:855–861;

doi:10.1542/peds.90.6.855.

Bloomingdale P, Housand C, Apgar JF, Millard BL, Mager DE, Burke JM, et al. 2017.

Quantitative systems toxicology. Translational Toxicology: Biomarkers 4:79–87;

doi:10.1016/j.cotox.2017.07.003.

Boobis AR, Cohen SM, Dellarco V, McGregor D, Meek ME, Vickers C, et al. 2006. IPCS

framework for analyzing the relevance of a cancer mode of action for humans. Critical

Reviews in Toxicology 36:781–92; doi:10.1080/10408440600977677.

Brookhart MA, Rassen JA, Schneeweiss S. 2010. Instrumental variable methods in

comparative safety and effectiveness research. Pharmacoepidemiology and Drug Safety

19:537–554; doi:10.1002/pds.1908.

Burgoon LD, Angrish M, Garcia-Reyero N, Pollesch N, Zupanic A, Perkins E. 2020. Predicting

the probability that a chemical causes steatosis using adverse outcome pathway bayesian

networks (AOPBNs). Risk Analysis 40:512–523; doi:10.1111/risa.13423.

Callegaro G, Kunnen SJ, Trairatphisan P, Grosdidier S, Niemeijer M, Hollander W den, et al.

2021. The human hepatocyte TXG-MAPr: Gene co-expression network modules to support

mechanism-based risk assessment. Arch Toxicol 95:3745–3775; doi:10.1007/s00204-021-

03141-w.

Campenhout CV, Cabochette P, Veillard A-C, Laczik M, Zelisko-Schmidt A, Sabatel C, et al. 2019. Guidelines for optimized gene knockout using CRISPR/Cas9. BioTechniques 66:295–302; doi:10.2144/btn-2018-0187.

Casarett LJ. 2008. *Casarett and doull's toxicology: The basic science of poisons.* McGraw-Hill New York.

Chen IY, Pierson E, Rose S, Joshi S, Ferryman K, Ghassemi M. 2021. Ethical machine learning in healthcare. Annual Review of Biomedical Data Science 4:123–144; doi:https://doi.org/10.1146/annurev-biodatasci-092820-114757.

Chen SH, Pollino CA. 2012. Good practice in bayesian network modelling. Environmental Modelling & Software 37: 134–145.

Cinelli C, Forney A, Pearl J. 2022. A crash course in good and bad controls. Sociological Methods & Research 00491241221099552; doi:10.1177/00491241221099552.

Conolly RB, Ankley GT, Cheng W, Mayo ML, Miller DH, Perkins EJ, et al. 2017. Quantitative adverse outcome pathways and their application to predictive toxicology. Environmental Science & Technology 51:4661–4672; doi:10.1021/acs.est.6b06230.

Cox LAJr. 2018. Modernizing the bradford hill criteria for assessing causal relationships in observational data. Critical Reviews in Toxicology 48:682–712; doi:10.1080/10408444.2018.1518404.

Davies NM, Smith GD, Windmeijer F, Martin RM. 2013. Issues in the reporting and conduct of instrumental variable studies: A systematic review. Epidemiology 24.

De Conti L, Baralle M, Buratti E. 2013. Exon and intron definition in pre-mRNA splicing.

WIREs RNA 4:49–60; doi:10.1002/wrna.1140.

Fréchette GR, Sarnoff K, Yariv L. 2022. Experimental economics: Past and future. Annual

Review of Economics 14:777–794; doi:https://doi.org/10.1146/annurev-economics-

081621-124424.

Gadaleta D, Spînu N, Roncaglioni A, Cronin MTD, Benfenati E. 2022. Prediction of the

neurotoxic potential of chemicals based on modelling of molecular initiating events

upstream of the adverse outcome pathways of (developmental) neurotoxicity.

International Journal of Molecular Sciences 23; doi:10.3390/ijms23063053.

Gou X, Ma C, Ji H, Yan L, Wang P, Wang Z, et al. 2023. Prediction of zebrafish embryonic

developmental toxicity by integrating omics with adverse outcome pathway. Journal of

Hazardous Materials 448:130958; doi:10.1016/j.jhazmat.2023.130958.

Graesser AC, Hu X. 2011. Commentary on causal prescriptive statements. Educational

Psychology Review 23:279–285; doi:10.1007/s10648-011-9163-x.

Greenland S, Pearl J, Robins JM. 1999. Causal diagrams for epidemiologic research.

Epidemiology 37–48.

Guzelian PS, Victoroff MS, Halmes NC, James RC, Guzelian CP. 2005. Evidence-based

toxicology: A comprehensive framework for causation. Human & Experimental Toxicology

24:161–201; doi:10.1191/0960327105ht517oa.

Helma C. 2005. *Predictive toxicology*. CRC Press.

Hemmerich J, Ecker GF. 2020. In silico toxicology: From structure–activity relationships

towards deep learning and adverse outcome pathways. WIREs Computational Molecular

Science 10:e1475; doi:10.1002/wcms.1475.

Herbert NL, Charles G, Alan L, Robert R, Henry P, Cornelius M, et al. 1979. Deficits in

psychologic and classroom performance of children with elevated dentine lead levels. New

England Journal of Medicine 300:689–695; doi:10.1056/NEJM197903293001301.

Hill AB. 1965. The environment and disease: Association or causation? Proceedings of the

Royal Society of Medicine 58:295–300; doi:10.1177/003591576505800503.

Hinson JA, Roberts DW, James LP. 2010. Mechanisms of acetaminophen-induced liver

necrosis. In: *Adverse drug reactions* (J. Uetrecht, ed). Springer Berlin Heidelberg:Berlin,

Heidelberg. 369–405.

Höfler M. 2005. Causal inference based on counterfactuals. BMC Medical Research

Methodology 5:28; doi:10.1186/1471-2288-5-28.

Höijer I, Emmanouilidou A, Östlund R, Schendel R van, Bozorgpana S, Tijsterman M, et al.

2022. CRISPR-Cas9 induces large structural variants at on-target and off-target sites in vivo

that segregate across generations. Nature Communications 13:627; doi:10.1038/s41467-

022-28244-5.

Jaeschke H, McGill MR. 2015. Cytochrome P450-derived versus mitochondrial oxidant

stress in acetaminophen hepatotoxicity. Toxicology Letters 235:216–217;

doi:10.1016/j.toxlet.2015.04.002.

James R, Britt J, Halmes N, Guzelian P. 2015. Evidence-based causation in toxicology: A 10-

year retrospective. Human & Experimental Toxicology 34:1245–1252;

doi:10.1177/0960327115601767.

Jensen MA, Blatz DJ, LaLone CA. 2023. Defining the biologically plausible taxonomic domain

of applicability of an adverse outcome pathway: A case study linking nicotinic acetylcholine

receptor activation to colony death. Environmental Toxicology and Chemistry 42:71–87;

doi:10.1002/etc.5501.

Jeong J, Song T, Chatterjee N, Choi I, Cha YK, Choi J. 2018. Developing adverse outcome

pathways on silver nanoparticle-induced reproductive toxicity via oxidative stress in the

nematode caenorhabditis elegans using a bayesian network model. Nanotoxicology

12:1182–1197; doi:10.1080/17435390.2018.1529835.

Kaikkonen L, Parviainen T, Rahikainen M, Uusitalo L, Lehikoinen A. 2021. Bayesian

networks in environmental risk assessment: A review. Integrated Environmental

Assessment and Management 17:62–78; doi:10.1002/ieam.4332.

Khabib MNH, Sivasanku Y, Lee HB, Kumar S, Kue CS. 2022. Alternative animal models in

predictive toxicology. Toxicology 465:153053; doi:10.1016/j.tox.2021.153053.

Koller D, Friedman N. 2009. *Probabilistic graphical models: Principles and techniques.* MIT

press.

Landfors F, Chorell E, Kersten S. 2023. Genetic mimicry analysis reveals the specific lipases

targeted by the ANGPTL3-ANGPTL8 complex and ANGPTL4. Journal of Lipid Research 64;

doi:10.1016/j.jlr.2022.100313.

Lipton R, Ødegaard T. 2005. Causal thinking and causal language in epidemiology: It's in the

details. Epidemiological Perspectives and Innovations 2: 8–9.

Mazer M, Perrone J. 2008. Acetaminophen-induced nephrotoxicity: Pathophysiology,

clinical manifestations, and management. Journal of Medical Toxicology 4:2–6;

doi:10.1007/BF03160941.

Meek M, Palermo CM, Bachman AN, North CM, Jeffrey Lewis R. 2014. Mode of action human

relevance (species concordance) framework: Evolution of the bradford hill considerations

and comparative analysis of weight of evidence. Journal of Applied Toxicology 34:595–606;

doi:10.1002/jat.2984.

Obermeyer Z, Emanuel EJ. 2016. Predicting the future - big data, machine learning, and

clinical medicine. The New England journal of medicine 375:1216–1219;

doi:10.1056/NEJMp1606181.

OECD. 2017. Guidance document for the use of adverse outcome pathways in developing

integrated approaches to testing and assessment (IATA).

OECD. 2014. *Guidance document on the validation of (quantitative) structure-activity*

relationship [(q)SAR] models.

Olariu V, Yui MA, Krupinski P, Zhou W, Deichmann J, Andersson E, et al. 2021. Multi-scale

dynamical modeling of t cell development from an early thymic progenitor state to lineage

commitment. Cell Reports 34:108622; doi:10.1016/j.celrep.2020.108622.

ONTOX. 2024. Scientific papers-ONTOX. ONTOX-project.

Pallocca G, Moné MJ, Kamp H, Luijten M, Van de Water B, Leist M. 2022. Next-generation

risk assessment of chemicals - rolling out a human-centric testing strategy to drive 3R

implementation : The RISK-HUNT3R project perspective. Alternatives to Animal

Experimentation : ALTEX 39:419–426; doi:10.14573/altex.2204051.

Pearl J. 2009. Causal inference in statistics: An overview. Statistics Surveys 3:96–146;

doi:10.1214/09-SS057.

Perkins EJ, Ashauer R, Burgoon L, Conolly R, Landesmann B, Mackay C, et al. 2019. Building

and applying quantitative adverse outcome pathway models for chemical hazard and risk

assessment. Environmental Toxicology and Chemistry 38:1850–1865;

doi:10.1002/etc.4505.

PrecisionTox. 2024. Publications-PrecisionTox.

RISK-HUNT3R. 2024. Publications-RISK-HUNT3R. RISK-HUNT3R.

Russo F, Williamson J. 2007. Interpreting causality in the health sciences. International

Studies in the Philosophy of Science 21:157–170; doi:10.1080/02698590701498084.

Sager JE, Yu J, Ragueneau-Majlessi I, Isoherranen N. 2015. Physiologically based

pharmacokinetic (PBPK) modeling and simulation approaches: A systematic review of

published models, applications, and model verification. Drug Metabolism and Disposition

43:1823; doi:10.1124/dmd.115.065920.

Sahlin U, Helle I, Perepolkin D. 2021. "This is what we don't know": Treating epistemic

uncertainty in bayesian networks for risk assessment. Integrated Environmental

Assessment and Management 17:221–232; doi:10.1002/ieam.4367.

Shingaki R, Kanda H, Kuroki M. 2021. Selection and integration of generalized instrumental

variables for estimating total effects. Statistical Papers 62:2355–2381;

doi:10.1007/s00362-020-01190-4.

Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, et al. 2011. A conditional

knockout resource for the genome-wide study of mouse gene function. Nature 474:337–

342; doi:10.1038/nature10163.

Song Y, Zheng K, Brede DA, Gomes T, Xie L, Kassaye Y, et al. 2023. Multiomics point of

departure (moPOD) modeling supports an adverse outcome pathway network for ionizing

radiation. Environmental Science & Technology 57:3198–3205;

doi:10.1021/acs.est.2c04917.

Spinu N, Cronin MTD, Enoch SJ, Madden JC, Worth AP. 2020. Quantitative adverse outcome

pathway (qAOP) models for toxicity prediction. Archives of Toxicology 94:1497–1510;

doi:10.1007/s00204-020-02774-7.

Spînu N, Cronin MTD, Madden JC, Worth AP. 2022. A matter of trust: Learning lessons

about causality will make qAOPs credible. Computational toxicology (Amsterdam,

Netherlands) 21:100205; doi:10.1016/j.comtox.2021.100205.

Tibshirani R. 1996. Regression shrinkage and selection via the lasso. Journal of the Royal

Statistical Society: Series B (Methodological) 58:267–288; doi:10.1111/j.2517-

6161.1996.tb02080.x.

Tollefsen KE, Scholz S, Cronin MT, Edwards SW, Knecht J de, Crofton K, et al. 2014. Applying

adverse outcome pathways (AOPs) to support integrated approaches to testing and

644 assessment (IATA). Regulatory Toxicology and Pharmacology 70:629–640;

645 doi:10.1016/j.yrtph.2014.09.009.

646 Villeneuve DL, Angrish MM, Fortin MC, Katsiadaki I, Leonard M, Margiotta-Casaluci L, et al.

647 2018. Adverse outcome pathway networks II: Network analytics. Environmental

648 Toxicology and Chemistry 37:1734–1748; doi:10.1002/etc.4124.

649 Villeneuve DL, Crump D, Garcia-Reyero N, Hecker M, Hutchinson TH, LaLone CA, et al.

650 2014a. Adverse outcome pathway (AOP) development i: Strategies and principles.

651 Toxicological Sciences 142:312–320; doi:10.1093/toxsci/kfu199.

652 Villeneuve DL, Crump D, Garcia-Reyero N, Hecker M, Hutchinson TH, LaLone CA, et al.

653 2014b. Adverse outcome pathway development II: Best practices. Toxicological Sciences

654 142:321–330; doi:10.1093/toxsci/kfu200.

655 Villeneuve D, Meek B, Viviani B, Burgdorf T, LaLone C, O'Brien J, et al. 2024. AOP

656 developers' handbook version 2.7.

657 Young C. 2019. The difference between causal analysis and predictive models: Response to

658 "comment on young and holsteen (2017)." Sociological Methods & Research 48:431–447;

659 doi:10.1177/0049124118782542.

660 Zou T, Caton AJ, Koretzky GA, Kambayashi T. 2010. Dendritic cells induce regulatory t cell

661 proliferation through antigen-dependent and -independent interactions. Journal of

662 immunology 185:2790–2799; doi:10.4049/jimmunol.0903740.