Working memory affects motor, but not perceptual timing

MohammadAmin Farajzadeh[1,2], Mehdi Sanayei[1]

[1]School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

[2]Shahid Beheshti University of Medical Sciences, School of Medicine, Tehran, Iran

Corresponding author: Mehdi Sanayei (mehdi.sanayei@gmail.com)

Abstract

Whether different timing tasks utilize the same brain processes is still debated. To approach this question, we investigated how working memory affects two different timing tasks: time reproduction and time discrimination. We found that delay interval led to an overestimation in the reproduction task but did not lead to any bias in the perception of time in the time discrimination task. Delay intervals affected perception of time when subjects had to actively reproduce the perceived interval, but not when subjects were going to just recall the content of the working memory. In a subsequent Bayesian modeling, we showed that in the reproduction task, subject updated their measurement of the stimulus on the current trial(likelihood), based on the delay interval, rather than changes in motor system, or updating prior based on the delay interval. Our findings suggest that brain processes that are involved in time reproduction and discrimination are not completely overlapped, and delay interval in working memory task leads to changes in updating the likelihood.

Keywords: Time perception, motor timing, perceptual timing, working memory

Introduction

Perception of time and duration of events is one of the most essential parts of everyday life and have a pivotal role in speech processing, decision making and motor control [1–3]. Perception of sensory information (including time) and keeping this information in memory are considered different processes, as studies reveal distinct brain circuits for perception vs. memory [4]. However, the capacity to perceive time and different durations is intertwined with the working memory process [5–9]. The classic psychological model for time perception, which is still in use today, consists of a pace-maker and an accumulator, in which the pace-maker emits pulses and these pulses are accumulated to be compared with the reference duration in working memory [5,10]. Despite the importance of the role of working memory in time perception, this topic has remained relatively under-studied (but see references 11 and 12).

In time perception literature, there are two main tasks used to study time perception. Perceptual timing tasks (e.g., time discrimination), in which the subject is presented with two intervals and has to report which interval is shorter or longer [13]. Subject does not actively reproduce either the first or second duration. On the other hand, there are motor timing tasks (e.g., time reproduction), which demands active engagement and manipulation of the perceived duration. In reproduction tasks, the experimenter presents a time interval to the subject, and the subject is asked to reproduce the presented time interval (e.g., by pressing a button for the perceived duration).

Working memory involves memorizing some information for short time periods [14]. Working memory has been studied using tasks that require delayed responses to stimuli presented earlier [4,15]. Various studies have demonstrated that as this delay interval increases, the subject's performance decreases [16]. Additionally, it has been shown that the presentation of certain types of distractor stimuli during the delay period leads to a larger decrease in working memory performance [17].

We can use delayed-response tasks to study the effect of working memory on time perception, but the delay interval itself is an elapsed time. So, the delay interval has dual effects on the perception of time. First, delay affects the task as a component of any working memory task. Second, given the sample interval and delay interval are both time intervals, delay intervals can interfere with the perception of the sample interval, like a serial dependence effect [18,19]. Serial dependence effect is a phenomenon in which stimuli from previous trials could have an effect on the perception of the current trial (i.e., local context [20]). Recent studies showed that this local context effect in time perception could be due to the strong effects of previous trials (updating prior). On the other hand, local context could be due to effects of previous trials on the signal to noise ratio of the stimulus on the current trial (updating likelihood). Although these studies investigated the effect of previous trials, none of them studied the effect of the delay interval, which comes after the to be timed interval, on time perception, either with a reproduction or a discrimination task.

In this study, we examined the effect of delay interval on time perception using both reproduction and discrimination tasks in the same subjects. We found that longer delay intervals led to an overestimation of the sample interval in the reproduction but not in discrimination task. We showed that, in the reproduction task, time perception attracted towards the delay interval duration. Furthermore, we proposed a computational model to explain this overestimation of sample interval. We found stronger evidence that the delay interval updates the likelihood function rather than the prior or changing the motor noise.

Material and Methods

Subjects

Fourteen subjects (5 female, 25.5 ± 5.16 years old) participated in this study. All subjects have normal or corrected-to-normal vision, and were naïve about the goal of the experiment, except the two authors (subject number 1 & 2). All subjects were enrolled in both reproduction and discrimination tasks and the order of the tasks was counterbalanced between them. This study was conducted under the approval of the ethics committee at the School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM). A written informed consent was taken from subjects prior to the experiment.

Apparatus

Experiments were performed on a PC with Linux operating system. We used the Psychtoolbox 3 extension with MATLAB (2016b) for designing our tasks [21–23]. Stimuli were presented on a 17" monitor (refresh rate 60Hz) placed ~60cm from subjects. Subjects sat on a chair in a dim room, and their head was stabilized with a chin and head rest. The position of the left eye was monitored (1KHz, EyeLink 1000, SR Research, Mississauga, Ontario) during the experiment.

Time reproduction task

On each trial, we presented a fixation dot (0.2°) and two targets (left target: 0.5° diameters, right target: 2° diameter, 10° eccentricity) to the subject (Fig. 1A). After the subject fixed the central dot for 200ms (within 2° window), a circular wheel-shaped stimulus (diameter = 2.5°) was shown. The duration of the stimulus was 400, 600 or 1000ms, chosen pseudo-randomly on each trial. After that, there was a delay in which only the fixation point and targets were present. The delay was selected pseudo-randomly on each trial from 500, 900, or 1700ms. After this delay, we presented a second stimulus, which was the same as the first one but rotated by 180°. The subject had to make a saccadic eye movement to the right target when the second stimulus' duration reached the first stimulus' duration. The reproduced time was measured from the onset of the second stimulus on the screen until the subject made the saccade. After response, the difference between the duration of the presented stimulus and the reproduced time was presented to the subject with millisecond resolution at the location of the right target. Each participant completed 9 blocks, and each block consisted of 72 trials. Before the main reproduction task, each subject did 4 blocks (18 trials each, 2 repetitions per condition) of the reproduction task, to get them familiarized with the task. The trials were excluded from any analysis.

Time Discrimination task

In this task, we presented two stimuli sequentially with a delay between them and then asked subjects to indicate which one of the presented stimuli was longer (Fig. 1B). The design and timing of the discrimination task were similar to the reproduction task, with the exception of the duration of the second stimulus. After the delay interval, the second stimulus' duration was the first stimulus' duration ± 6, 12, 24, 48% of the first stimulus duration. The subject had a 4-

second window to respond. If the second stimulus lasted longer than the first stimulus, the subject had to make a saccade to the right target. If the second stimulus lasted shorter than the first stimulus, the subject had to make a saccade to the left target. After that, the feedback was provided for the subjects with red and green filled circles, for correct and incorrect responses, respectively. Each participant carried out 18 blocks, and each block consisted of 72 trials. Before the start of this task, subjects did blocks of 18 trials, with only the easiest conditions. After a subject performed 100% on two consecutive blocks, we moved on to the main task. On average, subjects performed 4.35 ± 2.6 (mean and standard deviation) blocks to get familiarized with the task. These trials were not included for further analysis.

Analysis of behavioral data

For the reproduction task, we excluded any reproduced time that was larger than 1.5 IQR for each subject (0.7% of trials from all subjects excluded) [24]. For the remaining trials, we calculated the mean and standard deviation of the reproduced time for each sample duration and each delay interval. We plotted the mean reproduction time (t_r) against sample interval duration (t_s) for each subject and applied a linear regression function to the data from all subjects. These lines crossed the diagonal line at the 'indifference point', which represent an interval where subjects had a veridical perception of the physical time [25]. We also calculated the indifference point for each subject.

For the discrimination task, for each combination of sample stimulus duration and delay interval (9 in total), we fitted a cumulative Gaussian function to the percentage of 'longer' response as a function of the comparison stimulus duration. To assess how well the cumulative Gaussian fitted the data, we measured the goodness of fit and excluded poorly fitted conditions ($r^2 < 0.5$)(none was excluded) [24]. We measured the point of subjective equality (PSE) as the mean of the fitted Gaussian and its standard deviation as an index of variability for the discrimination task. Further, we fitted linear regressions to the PSE values for each delay interval as a function of sample interval duration to obtain indifference points the same as in the reproduction task. To investigate the effect of delay in each task on the indifference point, we applied a One-way repeated measure analysis of variance (RM-ANOVA) on each task, separately.

We applied a three-way RM-ANOVA to the mean of the reproduced time and PSE of the discrimination time to measure the effects of task (2 levels), sample interval (3 levels), and delay interval (3 levels) and their interaction on the accuracy of timing. We applied the same RM-ANOVA to standard deviation of both tasks to investigate the effects of these factors on the precision of timing. To investigate the effect of delay duration on the two tasks separately, we applied a two-way RM-ANOVA on the mean of time perception to measure the effect of sample interval duration and delay interval duration on the mean of the reproduced time (reproduction task), or the PSE (discrimination task).

To analyze the delay dependency of the reproduced interval, we used a deviation index. In the reproduction task, deviation index on each trial was computed as the reproduced duration minus the mean of reproduced durations for all trials with that sample interval. In the discrimination task, we first transformed data from the cumulative Gaussian format to a Gaussian distribution

with the same mean and standard deviation, for each sample and delay interval, per subject. Then we draw 500 samples for each sample and delay interval combination (9 conditions in total).We then treated these distributions the same as the reproduced time, as previously has been done [24,26], and calculate the deviation index similar to how we calculated in the reproduction task. Positive values for the deviation index showed that the reproduced interval for the current trial was longer than the mean reproduced duration and negative values indicated that the reproduced duration was shorter than the mean reproduced duration. To measure the effect of the delay duration on the deviation index, we fitted a simplified Derivative of Gaussian (DoG) curve to describe the deviation index as a function of difference between the current sample interval and delay duration as:

$$y = \frac{-ax}{b^2} e^{-\frac{x^2}{2b^2}} \quad (Eq.\,1)$$

Where y is the deviation index, x is the difference between sample interval and delay interval, a and b are fitted parameters. We fitted DoG at both the group and individual levels. To measure the goodness-of-fit, we computed r^2 for DoG model compared to non-model and linear interaction between deviation index and delay interval.

Modeling

To model the effect of delay interval on the reproduced time, we used a combination of Bayesian Observer[27], distractor [28], and mixed-Gaussian update models[20] . The basic observer consists of three different components: likelihood function, prior distribution and motor noise. Subjects could update any of these components based on the delay interval on each trial. To investigate which of these three possibilities explained our data best, we used modified versions of the classic Bayesian model for time perception [27]. We first explain the model in general form, and then the modifications that we applied for our three models.

This model breaks the process of time perception to measurement, estimation, and production. In the measurement step, the sample interval presented to the subject (t_s), is mapped to measurement time, t_m, using a Gaussian distribution, where the mean of this Gaussian distribution is equal to t_s, and its standard deviation is equal to $t_s \times w_m$, where w_m is a free parameter in the model. We denote this step with likelihood function λ.

$$\lambda_{t_m} = p(t_s, w_m) = \frac{1}{\sqrt{2\pi(w_m t_s)^2}} e^{\frac{-(t_s - t_m)^2}{2(w_m t_s)^2}} \quad (Eq.\,2)$$

We defined the prior distribution of t_s as a uniform distribution of times between t_s^{max} and t_s^{min} as follows:

$$\pi(t_s) = \begin{cases} \dfrac{1}{t_s^{max} - t_s^{min}} & t_s^{min} < t_s < t_s^{max} \\ 0 & otherwise \end{cases} \quad (Eq.3)$$

Posterior is the product of prior multiplied by likelihood function as:

$$\pi(t_m) = \frac{\pi(t_s)p(t_s)}{\int \pi(t_s)p(t_s)dt_s} \quad (Eq.4)$$

In the next step, which was estimation, the model combines this likelihood with the prior distribution of time samples based on the mean of the posterior, generating the estimated time (t_e):

$$t_e = f_{BLS}(t_m) = \frac{\int_{t_s^{min}}^{t_s^{max}} t_s p(t_s)dt_s}{\int_{t_s^{min}}^{t_s^{max}} p(t_s)dt_s} \quad (Eq.5)$$

In the final step, the values of t_e are mapped to the produced time (t_p) using a Gaussian distribution, where the mean of the distribution is equal to t_e and the standard deviation of the distribution is equal to $t_e \times w_p$, where w_p is a free parameter in the model.

$$p(t_e) = \frac{1}{\sqrt{2\pi(w_p t_e)^2}} e^{\frac{-(t_p - t_e)^2}{2(w_p t_e)^2}} \quad (Eq.6)$$

Finally, the direct relationship between the reproduced time and the sample time that was presented to the subject is explained by the following equation:

$$p(t_s, w_m, w_p) = \int p(t_p | f_{BLS}(t_m), w_p)p(t_m | t_s, w_m)dt_m \quad (Eq.7)$$

Now we are going to add component of distraction to model the effect of the delay duration (t_d). The distraction effect on any duration, t, can be modeled by weighted average between t_d and t:

$$\hat{t} = \frac{t + t_d w}{1 + w} \quad (Eq.8)$$

The weight of the distractor, w, was between 0 and 1. The greater the weight of the distractor, the more effect it has on the perceived duration ($\hat{t}$). w is computed as:

$$w = k^{|t - t_d|}; 0 \leq k \leq 1 \quad (Eq.9)$$

Where k is the leak factor determining the distractor weight on the interval. When $k = 1$ the weight will always be one. When it is any value between 0 and 1, as the difference between t_d and t increases the weight and the effect of the distractor would decrease. We should remember that k itself is a free parameter that was fitted for each subject.

In the first model, on each trial, we assumed that subject updated the likelihood function with respect to t_d as:

$$\lambda'_{t_m} = p(\hat{t}_s, w_m) = \frac{1}{\sqrt{2\pi(w_m \hat{t}_s)^2}} e^{\frac{-(\hat{t}_s - t_m)^2}{2(w_m \hat{t}_s)^2}} (Eq.10)$$

The direct relationship between the reproduced time and the sample time that was presented to the subject is explained as:

$$p(t_s, t_d, w_m, w_p, k) = \int p(t_p | f_{BLS}(t_m), w_p) p(t_m | \hat{t}_s, w_m) dt_m \quad (Eq.11)$$

We named this model 'Likelihood Model'.

For the second model, we applied the same logic of distractor to the value of t_e. The t_e in the third step is replaced with $\hat{t}_e$ as:

$$\hat{t}_e = f_{dis}(t_e, t_d, w) = \frac{t_e + t_d w}{1 + w} \quad (Eq.12)$$

And the final equation of this model as:

$$p(t_s, t_d, w_m, w_p, k) = \int p(t_p | f_{dis}(f_{BLS}(t_m)), w_p) p(t_m | f_{dis}(t_s), w_m) dt_m \quad (Eq.13)$$

We named this model as 'Motor Model'.

Delay interval can also affect time perception via updating the prior, similar to what was previously applied to sequential effects on timing [20]. To address this issue, we used an

updating mechanism in which subject added a normal distribution (mean $= t_d$, sd $= t_d*w_p$) to the initial prior. So, the updated prior for each trial computed as:

$$\pi'(t_s, t_d) = \frac{\pi(t_s) + r*N(t_d, t_d*w_p)}{\int [\pi(t_s) + r*N(t_d, t_d*w_p)]dt_s} \quad (Eq.\,14)$$

$\pi'(t_s, t_d)$ is the updated prior and r is the learning rate, so BLS function now incorporated the new prior to estimate the t_e and the BLS formulation is:

$$t_e = f'_{BLS}(t_m, t_d) = \frac{\int_{t_s^{min}}^{t_s^{max}} t_s p(t_s)\pi'(t_s, t_d)dt_s}{\int_{t_s^{min}}^{t_s^{max}} p(t_s)\pi'(t_s, t_d)dt_s} \quad (Eq.\,15)$$

The final relationship between t_s and t_p is:

$$p(t_s, t_d, w_m, w_p, r) = \int p(t_p | f'_{BLS}(t_m, t_d), w_p)p(t_m | t_s, w_m)dt_m \quad (Eq.\,16)$$

We called this third modeling the 'Prior Model'.

We maximized the likelihood of model parameters w_m, w_p, and k for the 'Likelihood Model' and 'Motor Model', and w_m, w_p, and r for the 'Prior Model', across all combination of t_s, t_p, and t_d for reproduction.

To compare the three fitted models, we first calculated Bayesian Information Criterion (BIC) as:

$$BIC = k\, ln\, ln\,(n) - 2\, ln\, ln\,(\hat{L}) \quad (Eq.\,17)$$

Which the $\hat{L}$ is the maximized value of the likelihood function of the model, n is the number of observations (in our case the number of included trials for the reproduction task) and k is the number of free parameters that are used in each model. We then calculated ΔBIC between the three models. Model with the smallest BIC would be the model with the best explanatory power (among the models that we studied).

Results

In the discrimination task, the fraction of the trials that subjects reported that the second interval was longer than the first interval are plotted for 400 (Fig. 2A), 600 (Fig. 2B), and 1000ms (Fig. 2C) sample intervals, separately for 500 (green), 900 (blue), and 1700 ms (red) delay intervals. We found that as the sample interval increased, the sensitivity of time perception decreased, as expected according to Weber's law, as can be appreciated by decreasing the slope of the fitted psychometric function as the sample interval increased. We then transformed the data based on

the mean and standard deviation of the fitted cumulative Gaussian functions to the format in Fig. 2B, to compare data from discrimination and reproduction tasks directly. We also observed the classic Vierdordt's effect of overestimation of shorter, and underestimation of longer durations in the time discrimination (Fig. 2E). Interestingly, it seems that the delay duration had no effect on time perception in the discrimination task. In the reproduction task (Fig. 2D), we observed Vierordt's effect and Weber's law, as expected. But in contrast to the discrimination task, the perceived time in the reproduction task was affected by the delay interval: as the delay interval increased, participants overestimated the sample interval.

In order to test the effects of the sample and delay intervals on the accuracy (i.e., mean) of the perceived time, we applied a three-way repeated measure ANOVA (Factor 1 = task: two levels; Factor 2 = sample interval: three levels; Factor 3 = delay interval: three levels). We found that while 'task' type ($F = 0.12$, df = 1, 0 = 0.73) or delay interval ($F = 2.60$, df = 3, $p = 0.09$) did not have any significant effects on the accuracy of time perception, sample interval had a significant effect ($F = 186.89$, df = 2, $p < 0.0001$). The interaction between sample interval $\times$ delay interval did not reach significant level ($F = 0.71$, df = 4, $p = 0.58$), but the interaction between task type $\times$ sample interval ($F = 5.72$, df = 2, $p < 0.008$) and task type $\times$ delay interval ($F = 9.53$, df = 2, $p < 0.0008$) reached significant level. The interaction between task type $\times$ sample interval $\times$ delay interval was not significant ($F = 0.99$, df = 4, $p = 0.41$).

Given the significant interaction between task type and other factors, we run a two-way RM ANOVA on the accuracy of time perception for discrimination and reproduction tasks, separately. In the discrimination task, while sample interval had a significant effect ($F = 85.23$, df = 2, $p < 0.0001$), delay interval did not ($F = 0.03$, df = 2, $p = 0.96$) on the accuracy. The interaction between sample interval $\times$ delay interval ($F = 0.60$, df = 4, $p = 0.66$) did not reach a significant level either. On the other hand, in the reproduction task, the effect of both sample interval ($F = 144.29$, df = 2, $p < 0.0001$) and delay interval ($F = 20.57$, df = 2, $p < 0.0001$) was significant. Their interaction, sample interval $\times$ delay interval, was not significant ($F = 1.17$, df = 4, $p = 0.33$). This result showed that delay duration affected the accuracy of time perception in the reproduction task (i.e., motor timing) but not in the discrimination task (perceptual timing).

Furthermore, we showed that the indifference point, the time of veridical judgment, in the discrimination task, was not different between delay durations: 636, 633, and 643ms in 500, 900 and 1700 ms delay durations (Fig. 2F, One-way ANOVA: $F = 0.16$, df = 2, $p = 0.84$). In the reproduction task we found that the indifference point increased from 571ms in 500ms delay to 713ms in 900ms delay and to 766ms in 1700 ms delay (Fig. 2G, One-way ANOVA: $F = 10.02$, df = 2, $p < 0.006$). The pairwise comparison was significant between all pairs: 500 vs. 900 (Signed Rank test, $Z = 2.66$, $p < 0.003$), 900 vs. 1700 ($Z = 1.85$, $p < 0.03$), and 500 vs. 1700 ms ($Z = 2.79$, $p < 0.002$).

To investigate the effect of the delay duration on the variability of perceived time, we applied a three-way RM ANOVA on standard deviation for each sample interval (3 levels), delay duration (three levels) and task (two levels). We found while task ($F = 7.71$, df = 1, $p < 0.01$) and sample interval ($F = 13.18$, df = 2, $p < 0.0001$) had a significant effect on the variability of time perception, delay interval did not ($F = 1.88$, df = 2, $p = 0.17$). The interaction between delay

interval × task type (F = 1.46, df = 2, p = 0.24), between delay interval × sample interval (F = 1.09, df = 4, p = 0.36), between task × sample interval (F = 1.90, df = 2, p = 0.16), or between task × sample interval × sample interval (F = 0.17, df = 4, p = 0.94) did not reach significant level. Comparing these data with the results from effects of delay interval on the accuracy of time perception, we found that while delay interval led to overestimation, it did not have any effects on the variability of time perception.

Previous reports showed that in a reproduction task, preceding trials has a significant effect on the perception of time on the current trial [18,19]. We were interested to see whether the delay interval, which comes after the sample interval in our task, had a similar effect on the current trial. To examine the effect of delay interval on sample interval, we calculated a deviation index. Deviation index is the difference between the reproduced time on a given trial and the subject's mean reproduced time, regardless of delay duration, calculated separately for discrimination and reproduction tasks. The effect of delay interval on time perception in the reproduction task is shown by the change in the deviation index as a function of the difference between delay interval and sample interval (Fig. 3A). This deviation effect peaked when the delay interval was larger than the sample interval by approximately 900ms and decreased for larger differences. This interaction between the deviation index and the difference of the delay and sample interval was captured by derivative of Gaussian curve ($r^2 = 0.33$), and also a linear fit ($y = kx$, $r^2 = 0.28$) but not with the assumption that the deviation index was not dependent on the difference between sample and delay intervals ($y = 0$, $r^2 = -0.004$). We applied the deviation index analysis to the data from the discrimination task as well (Fig. 3B). In line with the data presented in Fig. 2, we found that delay interval did not have any effects on the deviation index in the discrimination task ($r^2 = 0.001$, 0.001, 0, for DoG, linear, and y = 0, respectively).

So far, we showed that in the reproduction task, subjects overestimated time as a function of the delay duration. This could be explained by the fact that subjects were incorporating the delay interval into their timing process. This incorporation can happen at different levels. To tease apart underlying mechanisms quantitatively, we implemented models that took the delay interval into account, at three different stages of a classic Observer model of time perception.

In the classic version of the model subjects combined the information from likelihood (Fig. 4A) with prior to come up with a posterior distribution (Fig. 4B) and then map the mean of the posterior to the reproduced interval with Gaussian distribution (Fig. 4C). To investigate the possible source for our findings we used three modified versions of the Bayesian observer model. In the first version of the model (Likelihood Model), the delay duration altered the mean and standard deviation of the *measurement* step with a weight of 'w', determined by the leak factor (k). This incorporation changes mean and standard deviation of likelihood function (from solid to dotted line in Fig. 4D). In the second implementation, Prior Model, the delay interval alters the prior which is determined by the distribution of the sample interval (t_s) on each trial (Fig. 4E, from green shade to gray). In this model the prior that is incorporated in the estimation step updates on each trial. Observer adds a normal distribution with mean of t_d and standard deviation of $t_d w_p$ to uniform prior distribution with the learning rate (r). In the third version of the model (Motor Model), we incorporated the delay interval to the production stage of the

Observer model (Fig. 4F, from solid blue to dotted). In this case, the mean and standard deviation of the distribution that maps estimated time to reproduced time were modified by the delay interval. We calculated the Bayesian information criterion (BIC) for each version of the model, fitted to each subject separately, to see which model explained our data best. We performed these stages for the motor timing task, since only in the motor timing task we observed that delay interval affected time perception.

We found that in the motor timing task, adding delay interval to the measurement stage (Likelihood Model) out-performed the model that added the delay interval to the production stage (Motor Model) in 13 out of 14 subjects (Fig. 4G). The median of ΔBIC was 77.14 smaller in the Likelihood Model than in the Motor Model (Fig. 4J). We found that the Likelihood Model also outperformed the Prior Model in 13 subjects (Fig. 4H). The median of ΔBIC was 68.95 in favor of the likelihood Model (Fig. 4K). On the other hand, the Motor Model was a better fit than the Prior Model in 9 subjects (Fig. 4I). The median of ΔBIC was 14.28 in favor of the Motor Model relative to the Prior Model (Fig. 4L).

Discussion

We demonstrated that in a working memory task, increasing the delay interval, after a sample interval, led to an overestimation of the sample interval in a time reproduction task but not in the time discrimination task. More specifically, the perception of the sample interval was attracted towards the delay interval We did not observe any changes in performance in the time discrimination task across different delay intervals between the first (sample) and the second (test) intervals. The effect of the delay interval on time perception in the reproduction task, can be interpreted not only as decreasing the fidelity of the representation of stimulus held in working memory, but also as biasing that representation towards the delay interval duration. We also found that this attraction towards the delay interval in the reproduction task can be better explained by updating the likelihood function rather than updating the prior or changes in motor processes.

Previous studies have shown that factors not relevant to the task, such as delay in our study, have an effect on the perception of time [29]. One such study observed that the impact of inter-trial intervals on duration judgment is similar to that of delays [29].In another study, Heron et al., demonstrated that the presentation of auditory white noise before the sample duration as an adaptor could distract the perception of the subsequent sample duration. They showed that presentation of adaptor stimulus before a time discrimination task could cause attraction of points of subjective equality toward adaptor duration nonlinearly[30].

There are different hypotheses about the underlying mechanism for this phenomenon, including the multi-channel process of time[31]. According to this theory, there is a specific population of neurons that each neuron encodes for a narrow range of duration, with overlapping tuning curves. The subsequent activation of different populations of neurons could cause distraction in the perceived duration[31].

The observed overestimation of time in our reproduction task as a function of the delay interval has been reported before[32]. Fortin et al., hypothesized that this overestimation of sample duration in working memory tasks is due to fewer pulses accumulated in the pacemaker-accumulator model of time perception[33,34]. Based on these findings, we expected to observe that delay duration affects both motor and perceptual timing tasks, which is not what we found. We hypothesize that the underlying mechanism for this difference can be found in the two similar but slightly different aspects of memory: working memory and short-term retention of information. Tasks in which the subject is forced to perform a yes/no recognition instead of actively reproducing the stimulus held in memory are referred to as short-term retention tasks (like discrimination task in our study)[35]. On the other hand, tasks requiring active manipulation of the content of memory engage working memory. Different studies have shown that the circuits involved in each of these aspects of memory are distinct[36]. Postle et al., have shown that delayed recognition of the ordinal position of five randomly ordered letters in humans remains unaffected by repetitive transcranial magnetic stimulation (rTMS) of the mid-dorsolateral prefrontal cortex (dlPFC) during the delay period. However, impairment is observed with rTMS to the parietal cortex. On the other hand, rTMS of the dlPFC does affect performance when subjects are required to reorder these letters into alphabetical order[37]. So, our findings also show that increasing the delay duration affects the task which needs working memory (i.e., time reproduction) but not the task which only relies on the short-term retention of information (i.e., discrimination).

A previous study that used a delayed-response task to investigate the reproduction of time has reported different results from ours. Ueda et al., found that an increase in the delay duration caused an increase in the central bias of the reproduced time[11]. In their study, subjects overestimated the short duration more and underestimated the long duration also more as a function of the delay interval. Their data are not compatible with our finding that an increase in delay duration causes an increase in overestimation for both short and long durations. This difference in our data from theirs may be due to the difference in the time range between studies (0.4 to 1s in our case vs. 1 to 8s in theirs). Another possibility might be the difference in the modality of response (eye vs. hand[24]) and the fixation requirement in our design. In our task, subjects had to fixate on the center of the screen during the delay interval. We think that by maintaining fixation, our subjects were more aware of the passage of time compared to when they were free to look around. So, in our tasks, subjects might incorporate the delay duration into the sample duration. This can explain the discrepancy between their results and our data.

There is also another study which tackled the effect of working memory on time reproduction with a different task design[12]. In a n-back design, Gumus and Balci used a series of sample intervals, followed by an instruction to the subject to reproduce the nth sample interval. They demonstrated as the duration between the sample interval and time reproduction increased, the subject overestimated the sample interval. We believe that this effect is another manifestation of the effect observed in our data because the increase in n could be equivalent to an increase in the delay duration in our task. The similarity between the results of these two task designs shows that the overestimation due to an increase in working memory load is robust across different tasks.

To have a better understanding of how subjects incorporated the delay interval into their perception of time in the reproduction task, we used modified versions of the Bayesian observer model, introduced before[27]. It has been shown that the central tendency toward the mean of the prior distribution and scalar variability of time perception could be explained by Bayesian least squares estimator. Wang et al., investigated the serial dependence in time modality and showed that the preceding trial could affect the perception of the current trial by updating the prior[20]. Contrary to their result, we found that the model which updates likelihood better explains our data. We assume that this is due to the temporal organization of the delay interval in our task which is presented *after* the sample interval. We found that the prior is quite robust to the changes and manipulation which occur after the presentation of the stimulus, but at the same time, the likelihood function is prone to changes in temporal context (in our case, is presented after the stimulus). We also found that the likelihood model outperformed the model that changes the motor noise as a function of the delay interval duration.

In this study, we implemented the updating rule for likelihood in a way that it updates the mean and standard deviation of the likelihood function. There are also other implementations of likelihood update that could explain the biases away from the prior distribution. In this line, Wei and Stocker showed that the asymmetric likelihood duration could explain the repulsion from the mean of prior distribution[38]. We assume that these two implementations could change the posterior in a way that leads to an increase in the density of posterior, in the areas away from the mean of the prior. The stability of the prior and the trial-by-trial updates in the likelihood function show that subjects have both stability in their perception and sensitivity to changes in temporal context, which lead to a better estimation of a sensory stimulus[39].

Data availability

Data from subjects and corresponding codes for data analysis and modeling will be made available online upon publication of the manuscript.

Acknowledgment

We would like to thank Ahmad Pourmohammadi and Sepehr Sima for their helps in model implementation and comments on the previous version of the manuscript.

Reference:

1. Kotz, S. A., Schwartze, M. & Schmidt-Kassow, M. Non-motor basal ganglia functions: A review and proposal for a model of sensory predictability in auditory language perception. *Cortex* **45**, 982–990 (2009).
2. Bechara, A., Tranel, D., Damasio, H. & Damasio, A. R. Failure to Respond Autonomically to Anticipated Future Outcomes Following Damage to Prefrontal Cortex. *Cereb Cortex* **6**, 215–225 (1996).
3. Gallistel, C. R. Animal Cognition: The Representation of Space, Time and Number.
4. Pasternak, T. & Greenlee, M. W. Working memory in primate sensory systems. *Nature Reviews Neuroscience* **6**, 97–107 (2005).
5. Treisman, M. Temporal discrimination and the indifference interval: Implications for a model of the 'internal clock'. *Psychological Monographs: General and Applied* **77**, 1–31 (1963).
6. Ma, A. C., Cameron, A. D. & Wiener, M. Memorability shapes perceived time (and vice versa). *Nat Hum Behav* (2024) doi:10.1038/s41562-024-01863-2.
7. Nobre, A. C. The attentive homunculus: Now you see it, now you don't. *Neuroscience & Biobehavioral Reviews* **25**, 477–496 (2001).
8. Olton, D. S. Frontal cortex, timing and memory. *Neuropsychologia* **27**, 121–130 (1989).
9. Baudouin, A., Vanneste, S., Isingrini, M. & Pouthas, V. Differential involvement of internal clock and working memory in the production and reproduction of duration: A study on older adults. *Acta Psychologica* **121**, 285–296 (2006).
10. Gibbon, J., Church, R. M. & Meck, W. H. Scalar Timing in Memory. *Ann NY Acad Sci* **423**, 52–77 (1984).
11. Ueda, N., Tanaka, K. & Watanabe, K. Memory decay enhances central bias in time perception. *i-Perception* **13**, 20416695221140411 (2022).
12. Gümüş, G. & Balcı, F. Working memory for time intervals: Another manifestation of the central tendency effect. *Psychon Bull Rev* **30**, 2289–2295 (2023).
13. Grondin, S. *The Perception of Time: Your Questions Answered*. (Routledge, 2019). doi:10.4324/9781003001638.
14. Hitch, G. J. & Baddeley, A. D. Verbal Reasoning and Working Memory. *Quarterly Journal of Experimental Psychology* **28**, 603–621 (1976).
15. Regan, D. Storage of spatial-frequency information and spatial-frequency discrimination. *J. Opt. Soc. Am. A* **2**, 619 (1985).
16. Bisley, J. W. The Multiple Roles of Visual Cortical Areas MT/MST in Remembering the Direction of Visual Motion. *Cerebral Cortex* **10**, 1053–1065 (2000).
17. Magnussen, S., Greenlee, M. W., Asplund, R. & Dyrnes, S. Stimulus-specific mechanisms of visual short-term memory. *Vision Research* **31**, 1213–1219 (1991).
18. Wiener, M., Thompson, J. C. & Coslett, H. B. Continuous Carryover of Temporal Context Dissociates Response Bias from Perceptual Influence for Duration. *PLoS ONE* **9**, e100803 (2014).
19. Wehrman, J. J., Wearden, J. & Sowman, P. Decisional carryover effects in interval timing: Evidence of a generalized response bias. *Atten Percept Psychophys* **82**, 2147–2164 (2020).
20. Wang, T. *et al.* A unitary mechanism underlies adaptation to both local and global environmental statistics in time perception. *PLoS Comput Biol* **19**, e1011116 (2023).
21. Brainard, D. H. The Psychophysics Toolbox. *Spatial Vision* (1997).
22. Pelli. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision* (1997).
23. Kleiner M, Brainard D, Pelli D. What's new in Psychtoolbox-3. *Perception 36 ECVP Abstract Supplement* (2007).
24. Pourmohammadi, A. & Sanayei, M. Context-specific and context-invariant computations of interval timing. *Front. Neurosci.* **17**, 1249502 (2023).

25. Grondin, S. Timing and time perception: A review of recent behavioral and neuroscience findings and theoretical directions. *Attention, Perception, & Psychophysics* **72**, 561–582 (2010).
26. Merchant, H., Zarco, W. & Prado, L. Do We Have a Common Mechanism for Measuring Time in the Hundreds of Millisecond Range? Evidence From Multiple-Interval Timing Tasks. *Journal of Neurophysiology* **99**, 939–949 (2008).
27. Jazayeri, M. & Shadlen, M. N. Temporal context calibrates interval timing. *Nat Neurosci* **13**, 1020–1026 (2010).
28. Boned, J. & López-Moliner, J. Duration judgments are mediated by the similarity with the temporal context. *Sci Rep* **12**, 22575 (2022).
29. Lieving, L. M., Lane, S. D., Cherek, D. R. & Tcheremissine, O. V. Effects of delays on human performance on a temporal discrimination procedure: Evidence of a choose-short effect. *Behavioural Processes* **71**, 135–143 (2006).
30. Heron, J. *et al.* Duration channels mediate human time perception. *Proc. R. Soc. B.* **279**, 690–698 (2012).
31. Becker, M. W. & Rasmussen, I. P. The rhythm aftereffect: Support for time sensitive neurons with broad overlapping tuning curves. *Brain and Cognition* **64**, 274–281 (2007).
32. Fortin, C. & Rousseau, R. Time estimation as an index of processing demand in memory search. *Perception & Psychophysics* **42**, 377–382 (1987).
33. Fortin, C. & Breton, R. Temporal interval production and processing in working memory. *Perception & Psychophysics* **57**, 203–215 (1995).
34. Fortin, C., Rousseau, R., Bourque, P. & Kirouac, E. Time estimation and concurrent nontemporal processing: Specific interference from short-term-memory demands. *Perception & Psychophysics* **53**, 536–548 (1993).
35. Jolicoeur, P., Lefebvre, C. & Martinez-Trujillo, J. *Mechanisms of Sensory Working Memory*. (Academic Press, London, UK San Diego, CA, 2015).
36. Lewis-Peacock, J. A., Drysdale, A. T., Oberauer, K. & Postle, B. R. Neural Evidence for a Distinction between Short-term Memory and the Focus of Attention. *Journal of Cognitive Neuroscience* **24**, 61–79 (2012).
37. Postle, B. R. Working memory as an emergent property of the mind and brain. *Neuroscience* **139**, 23–38 (2006).
38. Wei, X.-X. & Stocker, A. A. A Bayesian observer model constrained by efficient coding can explain 'anti-Bayesian' percepts. *Nat Neurosci* **18**, 1509–1517 (2015).
39. Fritsche, M., Spaak, E. & De Lange, F. P. A Bayesian and efficient observer model explains concurrent attractive and repulsive history biases in visual perception. *eLife* **9**, e55389 (2020).

Figure Legends

Figure 1. The sequence of events in a trial for reproduction (A) and discrimination (B) tasks. In the reproduction task (A), after the subject acquired fixation, a sample stimulus presented on the screen for the duration of t_s. Then, the subject had to wait for a delay interval (t_d). After that, they reproduced the t_s by making a saccade to the right target. At the end of each trial, we showed the response error (difference between the sample interval and reproduced duration) to the subject in millisecond. In the discrimination task (B), the subject was shown two intervals, with a delay (t_d) between them and they had to compare them. The subject had to report which interval was longer with a saccadic eye movement to one of two targets. At the end of each trial visual feedback (correct or error) was presented to the subject.

Figure 2. A, B and C, the proportion of long responses for different test interval duration and fitted psychometric function for 400, 600, and 1000ms delay interval, respectively. D, The mean of reproduced time as a function of sample interval in the reproduction task. E, The point of subjective equality (PSE) of perceived interval as function of sample interval in the discrimination task. F, The indifference points for the discrimination task for different delay intervals. Arrows in each subplot represent the indifference point. G, The indifference points for the reproduction task for different delay intervals. Arrows in each subplot represent the indifference point. In F and G error bars represent ± one standard error of the intercept.

Figure 3. Effect of the delay interval on the deviation index in the reproduction (A) and discrimination (B) tasks. A: The mean of the deviation index as a function of difference between the delay interval and sample interval and fitted Derivative of Gaussian in reproduction task. Pale dots represent each subject's data, and dark dots represent the average data. The red line represents the fitted derivative of the Gaussian (DoG) fitted to the population data. The black line represents the linear fit to the population data. B: The mean of deviation index as a function of difference between delay interval and sample interval and fitted Derivative of Gaussian in the reproduction task. Pale dots represent each subject's data, and dark dots represent the average data. The blue line represents DoG fitted to the population data. The black line represents the linear fit to the population data.

Figure 4. The three components of the classic Bayesian observer model: likelihood (A), incorporating prior (B) and motor component (C). The vertical line represents the sample duration on a given trial. The Gaussian distribution represents the likelihood with the mean of t_s and standard deviation (SD) of $t_s \times w_m$. B, the green line represents the prior that was used in our study. The gray shaded area represents the posterior. The vertical line represents t_e which is the average of the posterior distribution. C, the vertical line represents t_e which is the same as t_e in B. The Gaussian distribution represent the mapping function from t_e to t_p with the SD of $t_e \times w_p$.

Modification applied to the Bayesian observer model in the likelihood function (D), prior distribution (E) and motor noise (F), which was performed independently and separately. D: The delay interval (t_d) updates the likelihood of the sample interval (t_s), from the solid, to the dotted red line, with the mean of $\hat{t}_s$ and the SD of $\hat{t}_s \times w_m$. E: In the Prior model, delay interval updates the prior and consequently the posterior. The updated prior (green shade) is the sum of the old prior with normal distribution centered on the t_d and SD of $t_d \times w_p$. The gray shade represents the consequently updated posterior. F: in the Motor model, the delay interval shifted the mean of production distribution from t_e (with SD of $t_e \times w_p$, solid blue line) to $\hat{t}_e$ (with the SD of $\hat{t}_e \times w_p$, dotted blue line).

G: ΔBIC between likelihood model and Motor model for each subject. Number on the y axis is subjects' id number. ΔBIC between likelihood model and Prior model (H) per subject, and between the Motor model and Prior model per subject (I). Distribution of ΔBIC between likelihood model and Motor noise model (J), between likelihood model and Prior model (K) and between Motor model and Prior model (L). In J, K, and L, the dotted vertical line represents the value of 0, and the solid vertical line represents the median of the ΔBIC distribution.

Fig 1

A)Reproduction Task

200ms fixation

400 or 600 or 1000 ms ts

500 or 900 Or 1700 ms Delay

Reproduced time

B)Discrimination Task

200ms fixation

400 or 600 or 1000 ms ts

500 or 900 Or 1700 ms Delay

Comparison interval

Decision

Fig 2

A Sample interval = 400 ms

B Sample interval = 600 ms

C Sample interval = 1000 ms

D Mean Reproduced Time

E Discrimination transformed to Reproduction

F Discrimination task

Delay=500ms — 641 ms
Delay=900ms — 628 ms
Delay=1700ms — 629 ms

G Reproduction task

Delay=500ms — 545 ms
Delay=900ms — 641 ms
Delay=1700ms — 684 ms

Fig 3

Fig 4