Insect metamorphosis is regulated differently between sexes by members of a microRNA

cluster

Chade Li[1,^], Ki Kei Chan[1,^], Wenyan Nong[1,^], ShanShan Chen[2,^], Wai Lok So[1], Zhe Qu[1], Heidi

Y.C. Wu[1], Ho Yin Yip[1], Chi Bun Chan[3], Stephen S. Tobe[4,#], William G. Bendena[5], Zhen Peng

Kai[6,*], Jerome H.L. Hui[1,*]

1. School of Life Sciences, Simon F.S. Li Marine Science Laboratory, State Key Laboratory of

Agrobiotechnology, Institute of Environment, Energy and Sustainability, The Chinese

University of Hong Kong, Hong Kong SAR, China

2. Institute of Agro-food Standard and Testing Technology, Shanghai Academy of Agricultural

Sciences, China

3. School of Biological Sciences, The University of Hong Kong, Hong Kong SAR, China

4. Department of Cell and System Biology, University of Toronto, Canada

5. Department of Biology, Queen's University, Canada

6. Shanghai Institute of Technology, Shanghai, China

^ = contributed equally, # = deceased

* = correspondence = kaizp@sit.edu.cn; jeromehui@cuhk.edu.hk

<u>Abstract</u>

Insects comprise the majority of all described animal species and dominate the terrestrial

habitats. The evolution of insect metamorphosis played a profound role in their successful

adaptation and radiation. Insect metamorphosis is dependent on hormones ecdysteroids and

sesquiterpenoids such as juvenile hormone. Despite the fact there are genuine differences

between sexes during insect metamorphosis which facilitate their successful mating, how such

sexual dimorphism in metamorphosis is being controlled is poorly known. We first generated

transcriptomic profiles of male and female flies in late larvae and early pupae stages. Using a

combination of genome-wide prediction and *in vitro* dual-luciferase validations, members of a

microRNA cluster miR-277/34 were found to potentially regulate the neuropeptide receptor

(*AstC-R1*) that when activated inhibits the sesquiterpenoid pathway and a juvenile hormone-

dependent transcription factor (*Kr-h1*) in fly *Drosophila melanogaster.* Loss-of-function

mutants were created deleting either miR-277 or miR-34, and expression levels of both *AstC-*

R1 and *Kr-h1* as well as ecdysteroid and sesquiterpenoid hormone titres were altered. Further

comparison of transcriptomes of the late larvae and early pupae of both sexes revealed

differential gene pathways being regulated by members of miR-277/34 between sexes during

metamorphosis. This study highlights how members of a microRNA cluster control hormonal

and developmental gene pathways in different sexes of insects during metamorphosis.

Keywords: Sesquiterpenoid, ecdysteroid, microRNA, cluster, metamorphosis, sexes

Background

The Arthropoda is the largest animal phylum, containing most of the described living
species in the world, while the Insecta (hexapods) is the most abundant class of arthropods.
Majority of insects undergo metamorphosis, which results in an alteration to the whole body-
plan during development (Truman and Riddiford 1999; Belles 2011). Such strategy of changing
in body form undeniably played an important role during the evolution and successful radiation
of insects, such as reduce competition of food source and increase exploration and settlement
in new environment (Cheong et al 2015; Truman 2019).

Insect metamorphosis is strictly programmed and precisely controlled by the hormonal
systems. The three major groups of hormones govern metamorphosis include ecdysteroids,
neuropeptides, and sesquiterpenoids (Bendena et al 1999; Tobe and Bendena 1999; Truman
and Riddiford 2002; Riddiford 2012; Cao et al 2017; Qu et al 2018; Belles 2017; Tsang et al
2020). In general, these three hormones work in a cooperative manner, where ecdysteroid 20-
hydroxyecdysone control the timing of ecdysis or molting, neuropeptide ecdysis-triggering
hormone act as regulators of ecdysteroid synthesis or release, and sesquiterpenoids such as
juvenile hormone (JH) regulate metamorphosis based on their titre levels.

During larva to larva ecdysis, JH binds to intracellular receptor *Methoprene-tolerant*
(Met) and subsequently the transcription factor *Krüppel homolog 1* (Kr-h1) to suppress
metamorphosis to pupa (Konopova and Jindra 2007; Parthasarathy et al 2008; Minakuchi et al
2009); and during larva to adult metamorphosis, in the absence of JH, transcription factor *E93*
induces metamorphosis via the MEKRE93 (Met-Kr-h1-E93) pathway (Ureña et al 2016).
MicroRNAs are conserved noncoding RNAs that play crucial roles in many biological
processes in insects by post-transcriptional regulation. In the last decade, a number of
microRNAs have been identified that contribute to the regulation of insect metamorphosis (e.g.
Caygill and Johnston 2008; Sokol et al 2008; Lozano et al 2015).

In the fruit fly *Drosophila melanogaste*r, microRNAs have been found to regulate
hormonal and metabolic pathways (e.g. Chan et al 2023; Qu et al 2017). Similar to other
holometabolous insects, the life cycle of *Drosophila* exhibits embryo, larva, pupa, and adult
stages. Notably, *Drosophila* also shows sexual dimorphism in many aspects, including
reproductive behaviours (Han et al 2022), abdomen segments (Wang et al 2011), and
responsiveness to ethanol (Devineni and Heberlein 2012). Interestingly, feeding males with
juvenile hormone inhibitor could lead to feminization, and this effect is reversible by the

simultaneous application juvenile hormone analogue, suggesting hormonal control of sexual

dimorphism (Belgacem and Martin 2002).

Furthermore, previous study revealed sex-biased expression of microRNAs in various

tissues of males and females flies (Fagegaltier et al 2014). For instance, let-7 expressed in the

heads of both sexes, while its expression in the gonads could only be found in the males.

Interestingly, let-7 expression could be induced by ecdysone, while the maintenance of sexual

identity in male adulthood is regulated by let-7 and ecdysone signalling pathway genes

(Sempere et al 2002). However, whether microRNA could be involved in hormonal control and

sexual dimorphism during insect metamorphosis, remains unknown.

Here we identified a microRNA cluster involved in metamorphosis by comprehensive

mRNA and small RNA sequencing together with genome-wide *in silico* and *in vitro* analyses.

Subsequent knockout of members of this microRNA cluster and analyses demonstrated that

developmental and hormonal pathway genes were differentially regulated contributing to the

sexual dimorphism of insect metamorphosis.

Results

Differential protein-coding genes and microRNAs expression in male and female flies

during metamorphosis

To reveal the protein-coding genes and microRNAs expression in different sexes of

flies during metamorphosis, we have first sequenced the transcriptomes of L3 larvae instars

and white pre-pupae from both sexes. A total of 2,548 protein-coding genes were found to be

differentially regulated in females during metamorphosis (981 upregulated and 1,567

downregulated), while 2,650 protein-coding genes were differentially regulated in males (1,002

upregulated and 1,648 downregulated) (Figure 1A; Supplementary Table S1, S2;

Supplementary Figure S1; Supplementary File 1). We then tested if any functional gene

ontology terms were enriched in either sex or during metamorphosis for these differentially

expressed genes, and the results are summarised in Supplementary Table S3. In sum, most of

the enriched gene pathways are revealed for the downregulated genes during metamorphosis,

including the *insect hormone biosynthesis* pathway in both males and females, while some sex-

specific enriched gene pathways, such as amino acid metabolic pathways in males and
ascorbate and aldarate metabolism in females, could also be identified (Figure 1A).

On the other hand, we have also sequenced and uncovered a total of 52 microRNAs
was differentially regulated in females during metamorphosis (13 upregulated and 39
downregulated), and 40 microRNAs were differentially regulated in males (22 upregulated and
18 downregulated) (Figure 1B; Supplementary Table S4, S5; Supplementary Figure 2;
Supplementary File 1). There were 9 microRNAs that were found to be differentially expressed
similarly between sexes during metamorphosis, including miR-314-5p, miR-317-5p, miR-34-
3p and miR-980-5p to be downregulated, and miR-100-5p, miR-100-3p, miR-125-5p, miR-
125-3p and let-7-5p to be upregulated in both sexes during metamorphosis. In addition, there
were 12 microRNAs that showed opposite trends in expression between sexes during
metamorphosis (e.g. up-regulated in male but down-regulated in female), including miR-2b-1-
5p, miR-7-5p, miR-9b-5p, miR-13b-1-5p, miR-190-5p, miR-275-5p, miR-275-3p, miR-4969-
5p, miR-9372-5p, miR-963-5p, miR-970-5p and miR-987-5p (Figure 1B; Supplementary
Figure 3 and 4).

Screening for microRNAs targeting the sesquiterpenoid signalling pathway genes

In insects, sesquiterpenoids is one of the key hormones that regulate metamorphosis,
and there are less than a handful of published studies on microRNA regulation of
sesquiterpenoids in fly *D. melanogaster*, such as microRNA *bantam* regulates gene transcript
of the JHAMT gene and sesquiterpenoid biosynthesis (Qu et al 2017). Using the newly
generated transcriptomic data, we further tested if there were other potential microRNAs
involved in regulation of other parts of the sesquiterpenoid system. Genome-wide *in silico*
prediction of microRNAs that can potentially regulate different genes involved in the
sesquiterpenoid system of fly was carried out (Marco et al 2010; Griffiths-Jones et al 2011; Qu
et al 2020), and a list of 17 common microRNAs that can potentially regulate the
sesquiterpenoid system (other than the biosynthetic pathway) were predicted (Supplementary
Figure 5; Supplementary File 1).

Utilising an established cell culture dual-luciferase reporter assay (Hui et al 2013); we
have validated 7 microRNAs (miR-8, miR-277, miR-958, miR-1000, miR-1002, miR-2283,
miR-2280) that can downregulate the expression of *Drosophila* sesquiterpenoid pathway genes
to varying degrees, including *allatostatin C* (*AstC*), *allatostatin C receptor 1* (*AstC-R1*),
allatostatin C receptor 2 (*AstC-R2*), *Krüppel homolog 1* (*Kr-h1*), *Chd64*, *juvenile hormone*
binding protein (JHBP), *juvenile hormone diol kinase* (*JHDK*) and *juvenile hormone esterase*

(*JHE*) (Figure 1C). This *in vitro* screening confirms a list of potential microRNAs and
sesquiterpenoid gene interactions.

Among these genes, Kr-h1 or Kruppel homolog 1, is a juvenile hormone dependent
transcription factor, and have been found to be regulated by another microRNA in cockroaches
(Lozano et al 2015); while AstC-R1 or allatostatin C receptor 1, is a G-protein coupled receptor
for neuropeptide allatostatin C which regulates juvenile hormone (Stay and Tobe 2007).
Interestingly, it can be found that *Kr-h1* and two *AstC* receptors were targets of either or both
members of miR-277/34 cluster in our *in vitro* analyses in flies above. To delineate whether
miR-277 can selectively bind to AstC-R1 and Kr-h1 transcript in the presence of multiple
transcripts *in vivo*, a labelled microRNA pull-down (LAMP) assay system (Hsu et al 2009; Qu
et al 2017) was applied, and miR-277 has been shown to directly bind and regulate AstC-R1
and Kr-h1 transcripts *in vivo*, in agreement with the above *in silico* and *in vitro* results (Figure
1D). We also investigated the expression of miR-277 and miR-34 during *D. melanogaster*
development with small RNA sequencing and qPCR, which revealed a correlative expression
pattern with the expression level of Kr-h1 and AstC-R1 (Figure 1E-H).

Functional analyses reveal Kr-h1 and AstC-R are regulated by miR-277/34

The question then becomes, how significant is the finding of the revealed microRNA
interaction, and could it really alter any biological activities or functions? Using the available
homozygous loss of function of miR-277/34 cluster fly line (w[*]; Df(3R)mir-277-34-KO,
TI{w[+mW.hs]=TI}mir-277-34-KO/TM3, P{w[+mC]=GAL4-twi.G}2.3, P{UAS-
2xEGFP}AH2.3, Sb[1] Ser[1]; Chen et al 2014), we found that the heterozygous knockout
individuals displayed almost no lethality from the mutation, while the survivorship for the
double knockout flies fell significantly with 77% of the double knockout flies dying within five
days after eclosion (Figure 2A).

To fully understand how microRNA cluster miR-277/34 could be involved in the
regulation of sesquiterpenoid via non-biosynthetic pathway genes in the fly, we utilised
CRISPR/Cas9 to generate two individual knockout mutants for each of the microRNAs (i.e.
miR-277 and miR-34). The mutant lines, which we named miR-277-knockout (miR-277-KO)
and miR-34-knockout (miR-34-KO), were successfully created with confirmation by both
Sanger sequencing and next-generation genome re-sequencing (Supplementary Figure S3A-B).

We then compared the expression of *Kr-h1* and *AstC-R1* of miR-277-KO, miR-33-KO,
and between KO mutants and wild type flies using real-time PCR. The expression level of *Kr-
h1* was found to be significantly downregulated in both sexes of larva and pupa miR-277-KO

mutants; whereas a reduction was only noted in the male larvae for miR-34-KO mutants
(Figure 2B).

On the other hand, expression level of *AstC-R1* was found to be upregulated during
metamorphosis in wild-type (*w1118*) flies, and such change becomes insignificant in both miR-
277-KO and miR-34-KO mutants (Figure 2C; Supplementary Figure 6). In addition, we found
that the expression levels of *AstC-R1* in pupae of both miR-277-KO and mir-34-KO flies were
significantly different to that of the *w1118* flies (Figure 2C). All these data demonstrated that
Kr-h1 and *AstC-R* are stage-specifically regulated by miR-277/34 *in vivo*.

Sesquiterpenoid (JHB3, JHIII and FA) and ecdysteroid titres are altered differently
between sexes in miR-277 and miR-34 knockout mutants
During larva-pupal transitions in flies, either the absence or excess of sesquiterpenoids
results in abnormal phenotypes, including pupa lethality (Riddiford and Ashburner 1991;
Riddiford et al 2010; Wen et al 2015). To assess the roles of miR-277 and miR-34 in regulation
of sesquiterpenoid levels and other key metamorphosis hormone ecdysteroids in flies, we
measured the titres of sesquiterpenoid hormones (JHB3, JH III, and FA) and ecdysteroid 20-
hydroxylecdysone (20E) on the whole body of late L3 larvae and early pupae of both males
and females of miR-277-KO, miR-34-KO, and *w1118* (as control) (Supplementary Table S6.).

In both the sesquiterpenoid and ecdysteroid titres measurement, we observed
differences between males and females in the microRNA knockout mutants as well as the
controls (Figure 2D). For example, we could only observe significant differences of
sesquiterpenoid level changes during metamorphosis in females but not males between the
control, miR-277-KO, miR-34-KO (Figure 2D). These data showed that both sesquiterpenoids
and ecdysteroids were altered by miR-277 and miR-34.

Members of miR-277/34 regulate different sesquiterpenoid and related gene pathways
To characterize global gene expression changes triggered by losses of miR-277 and
miR-34, we compared the transcriptomes of RNA extracted from wandering 3rd instar larvae
(L3) and white prepupae from both sexes of miR-277-KO and miR-34-KO (Supplementary
Figure S8-10). In the gene pathway enrichment analyses, we revealed that the insect hormone
biosynthetic pathways were enriched in both control and miR-277-KO males (Figure 2F,
Supplementary Figure S8-9). In addition, in evaluation of gene expression of those involved in
sesquiterpenoid and ecdysteroid hormones biosynthetic pathways during larvae-pupae

transition, we also found that miR-277 and miR-34 affect their gene expression to different
degrees and trends in agreement to the hormonal measurement (Figure 2F and 2G).

The locus of microRNA cluster miR-277/34 has a common promoter and produces
three noncoding RNA transcripts including CR43459-RB and CR43459-RC. CR43459-RB
will encode miR-277, while CR43459-RC will encode both miR-277 and miR-34 (Figure 3A).
These two noncoding RNAs CR43459-RB and CR43459-RC have differ expression profile,
with CR43459-RC has a specific expression only at the adult male, and CR43459-RB has
higher expression in embryonic, early larval, pupae, and adult stages but drops to low levels
only during the late pupal stage before metamorphosis (Figure 3A). These expression profiles
suggesting that miR-277 and miR-34 maybe functioning in a cooperative manner.

To understand which target genes can be regulated by members of this miR-277/34
cluster, we first compared the enriched gene pathways in the miR-277-KO and miR-34-KO
mutants at different developmental stages and sexes when comparing to the wild-type *w1118*
(Supplementary Figure S8-9). We found that there are both similar and unique gene pathways
regulated by miR-277 and miR-34. For example, in the miR-277-KO mutants, alterations to
the *Hippo signaling*, *MAPK signaling*, *TGF-beta signaling*, *FoxO signaling*, *Wnt signaling*,
and *Dorso-ventral axis formation* gene pathways related to the sesquiterpenoid hormones are
revealed (Figure 3B-D; Supplementary Figure S11-13); while the *Broad complex* (*Br-C*) which
is a downstream transcription factor of the ecdysteroid hormonal pathway is significantly
downregulated in miR-277-KO flies but only in miR-34-KO pupae females (Figure 3C),
suggesting that the microRNA members in this miR-277/34 cluster regulate the sesquiterpenoid
and related gene pathways in a cooperative manner (summarized in Figure 3B and D).

Differential gene pathways regulated by miR-277/34 between sexes during
metamorphosis

Further dissection of the above altered gene pathways in miR-277-KO mutants revealed
that the *Hippo signaling* and *MAPK signaling pathways* are mutually downregulated regardless
of sexes and life stages, whereas the *TGF-beta signaling* and *FoxO signaling* are
downregulated exclusively in female pupae (Figure 3B, Supplementary Figure S8). In addition,
Wnt signaling pathway is often downregulated in pupae, while the *Dorso-ventral axis*
formation pathway is significantly downregulated in females but not in males. This data
prompted us to consider that miR-277 while having conserved regulatory roles of

sesquiterpenoid and related pathways, could also play divergent roles between sexes and
metamorphosis stages.

Based on the new findings, we have compared differentially expressed genes during
metamorphosis (i.e. larvae vs pupae) in different sexes between miR-277-KO, miR-34-KO,
and controls. Genes in the *Toll and Imd signaling pathway*s were mainly downregulated in
miR-277-KO females and miR-34-KO males, but slightly upregulated in miR-277-KO males
through larvae-pupae transition (Figure 4A, Supplementary Figure S1 and S14). Interestingly,
there are more male-specific downregulated gene pathways exclusively revealed in miR-277-
KO males during larvae-pupae transition, including the *arginine and proline metabolism, fatty*
acid degradation, beta-alanine metabolism, *cysteine and methionine metabolism*, *fatty acid*
degradation, pantothenate and CoA biosynthesis, and *valine leucine and isoleucine*
degradation (Figure 4A and B, Supplementary Figure 1). On the other hand, there are much
fewer enriched downregulated gene pathways in miR-34-KO mutants during metamorphosis,
such as the *other types of O-glycan biosynthesis* gene pathways which were specific to only
the male flies (Supplementary Figure 1, Supplementary Table S7). These data added up that
the members of microRNA cluster miR-277/34 also play differential gene regulatory roles
between sexes during metamorphosis (Figure 4B).

Discussion

Animals display sexual dimorphism, and in the fruit fly *Drosophila*, sex-specific
differences in body morphology, behaviour, and gene regulation have been previously
described (e.g. Han et al 2022; Wang et al 2011; Coyne and Oyama 1995). After fly eclosion,
different titres of ecdysteroids have been revealed in males and females potentially due to body
size differences (Bownes et al 1984; Testa et al 2013). There is a plethora of studies showing
sex-biased expression of micoRNAs across metazoans, however, there remains limited
examples demonstrating how microRNAs could contribute the differences in insect sexes
during metamorphosis and related hormonal control. We have first generated transcriptomes of
both mRNA and microRNAs from both sexes in late larvae and early pupae stages, carried out
genome-wide *in silico* prediction, *in vitro* luciferase assays, and *in vivo* pull-down assay, and
narrowed down a microRNA candidate miR-277, as part of the miR-277/34 cluster, which can
potentially involve in regulation of AstC-R1 and Kr-h1 transcripts. Previous studies have
demonstrated miR-277 play various functional roles in flies, including circadian rhythm (Anna
et al 2023), life span (Esslinger et al 2013), neurodegeneration and neuromuscular control (Tan
et al 2012; Deshpande et al 2024; Cerro-Herreros et al 2016), homeostasis (Jones et al 2013;

Li et al 2020), fatty acid metabolism (Zipper et al 2022), and amino acid metabolism (Esslinger

et al 2013; Stark et al 2003). On the other hand, miR-34 has also been shown to play other roles

in flies such as in autophagy, brain development, life span, survivorship, and regulation of

ecdysone (Liu et al 2012; Lai et al 2016; Kennerdell et al 2018, McNeill et al 2020; Srinivasan

et al 2022; Lai et al 2020; Xiong et al 2016). Together with the correlated expression data

(AstC-R1, Kr-h1, miR-277 and miR-34), *in vivo* mutants construction, hormone measurement

and transcriptomic analyses, this study strongly demonstrated that miR-277 and miR-34 can

play cooperative biological roles in sesquiterpenoid hormones in the fly sexes during

metamorphosis.

Unlike plants and fungi, metazoans including insects usually have significant

proportion of microRNAs that are clustered in their genomes (e.g. Bartel 2018; Tanzer and

Stadler 2004). Members of these microRNA clusters often transcribed as a single unit of

polycistronic transcript regulated by a single promoter (Kozomara and Griffiths-Jones 2014).

For instance, misregulation of clustered microRNAs could lead to disease and cancer formation

in humans (Dambal et al 2015; Ventura et al 2008; Kim et al 2009), and thus microRNAs in

the same cluster are proposed to possess similar targeting properties or regulate genes in the

same pathway (e.g. Hausser and Zavolan 2014). In *Drosophila*, microRNA cluster miR-

6/5/4/286/3/309 has been demonstrated that new members can reinforce the gene regulatory

network of existing microRNAs (Qu et al 2020), and let-7/miR-125 cluster are also well known

to be controlled by ecdysteroid hormones (e.g. Garbuzov and Tatar 2010). This study now

further provided evidence for functional constraints in microRNA clustering in insects, and in

this case of miR-277/34, via regulating different hormonal and related gene pathways in

different sexes during metamorphosis.

Materials and Methods

Animal husbandry, sex differentiation, and transcriptomes

D. melanogaster w1118 was reared at 25□C, 45-50% relative humidity and 14:10 hour

light-dark cycle. Sexes of wandering L3 larvae instar were differentiated under the microscope.

Total RNA was extracted from different sexes of L3 larvae instar and white pre-pupae stages

with TRIzol® Reagent (Thermo Fisher Scientific, US). The integrity and quantity of RNA was

verified through gel electrophoresis and NanoDrop One/OneC spectrophotometer (Thermo

Fisher Scientific, US). Total RNA was submitted to Novogene Hong Kong for library

construction and sequencing on the Illumina Novaseq6000 (PE150) platform. Raw reads were

trimmed and filtered using Trimmomatic (version 0.39, with parameters 'ILLUMINACLIP: TruSeq3-PE.fa:2:30:10 SLIDING WINDOW:4:5 LEADING:5 TRAILING:5 MINLEN:25') (Bolger et al 2014), while contamination were removed using Kraken2 (version 2.0.8) (Wood et al 2019). The processed reads were then aligned using HISAT2 (v2.0.5) (Kim et al 2019) and gene matrix count tables were generated using StringTie (v2.1.1) (Kovaka et al 2019) with references downloaded from Flybase (version FB2023_02). Details can be found in Supplementary Table S8.

In addition, small RNA was also extracted from different sexes of L3 larvae instar and white pre-pupae stages with mirVana™ miRNA Isolation Kit (Ambion). The integrity and quantity of RNA was checked using an Agilent 2100 Bioanalyser (Agilent RNA 6000 Nano Kit), and submitted to BGI Hong Kong for HiSeq small RNA library construction and 50-bp single end sequencing. Small RNA sequencing reads were analysed as previously described (Qu et al 2020). In brief, adapters were trimmed using cutadapt (version 1.11) (Martin 2011) and quality control was performed using fastqc (Andrews 2010). Qualified reads within 18 and 26 bp were used to generate a matrix count table using the 'quantifier.pl' module of the mirDeep2 package (Friedländer et al., 2012), with '-g 0' to allow for 0 mismatch when mapping reads to precursors downloaded from miRBase (version 22.1) (Kozomara et al 2019). The resulting count matrix was used for expression analyses in edgeR (Robinson et al 2010). Clean reads were also aligned to precursor sequences using Bowtie (Langmead et al 2009) with the parameters "-l 18 -v 0 -a --best --norc --strata" followed by manual inspection and checking. Details can be found in Supplementary Table S8.

MicroRNA target prediction and validation (dual-luciferase reporter, LAMP, and quantitative microRNA Taqman assays)

MicroRNA target prediction and in vitro validation were carried out as previously described (Qu et al 2017, 2020). In brief, mature microRNA sequences were retrieved from miRbase (version 22.1) (Kozomara et al 2019), and used to predict targets using the miRanda algorithm (Enright et al 2003). In addition, target prediction was also performed with TargetScanFly (release 7.2) (Agarwal et al 2018).

To validate the predicted microRNA targets, a dual-luciferase reporter assays were used (Qu et al 2017, 2020). MicroRNAs were amplified from D. melanogaster with amplicons cloned into pAC5.1 vector (Invitrogen), while gene targets were cloned into psi-check2 vector (Promega). All constructs were sequenced to confirm their identities. Drosophila S2 cells were kept at 23-25°C in Schneider Drosophila medium with 10% (v/v) heat-inactivated fetal bovine

serum and 1:100 penicillin–streptomycin (Gibco, Life Technologies). The pAC5.1-microRNA (200 ng) and psi-check2 vector (100ng) were transfected into Drosophila S2 cells using Effectene (Qiagen) per the manufactures' instructions. Luciferase activities were measured at 36-48 hours post-transfection using Dual-Luciferase Reporter Assay System (Promega) using Tecan Spark Multimode Microplate Reader (Eastwin International Trading Ltd).

Further validations were also carried out for miR-277 and miR-34 with labelled microRNA pull down assays (LAMP) as well as expression level analyses (TaqMan™ MicroRNA Reverse Transcription Kit and TaqMan™ MicroRNA Assay (Applied Biosystems), Assay IDs are 464452_mat, 463733_mat, 000298, and 463695_mat) as previously described (Qu et al 2017, 2020). Biotin-labelled oligos synthesized by IDT company while the Dynabeads M-280 streptavidin were obtained from the Invitrogen. The PCR was performed with the StepOnePlus™ Real-Time PCR System (Applied Biosystems). Sequences of oligos are provided in Supplementary Table S9.

miR-277 and miR-34 knockout mutants generation and confirmation

To prepare the CRISPR deletion constructs of miR-277 and miR-34, the corresponding target sequences and sequences for homologous recombination repairing were cloned into the pU6-3-Bsal-gRNA and pHD-DsRed-attP vectors (primer information is provided in Supplementary Table S9). Constructs were sequenced prior to injection into D. melanogaster w1118 embryos. D. melanogaster homozygous miR-277 and miR-34 CRISPR knockout mutants were screened and selected based on markers and phenotypes. Genomic DNA of each mutant line was extracted with PureLink DNA extraction kit (Invitrogen) following the manufacturer's protocol. PCR was performed with the primer sequences provided in the supplementary Table S9 to confirm that the respective microRNA has been taken out. In addition, whole genome resequencing was also performed on the these mutants on Illumina HiSeq platform at the Novogene Hong Kong. Sequenced raw reads were trimmed and cleaned using Trimmomatic (v0.39) (Bolger et al., 2014) and Kraken 2 (Wood et al., 2019) and aligned to the D. melanogaster reference genome downloaded from Flybase (r6.45) using BWA-MEM (Li, 2013) with parameters "-M -R", and the alignment was checked manually using samtools tview.

Ecdysteroid hormone measurement

Total body ecdysteroids were extracted from w1118, mir-277 knockout mutants (miR-277-KO) and mir-34 knockout mutants (miR-34-KO) L3 larvae and early pupae from both

sexes (n=3, 50 individuals per sample using methanol (Furuta et al., 2010), and the levels were

determined using the 20-hydroxyecdysone (20E) assay described previously (Kai et al., 2021).

Sesquiterpenoid hormone measurement

Total body sesquiterpenoid juvenile hormones were collected from w1118, miR-277-

KO and miR-34-KO L3 larvae and early pupae from both sexes (n=3, 50 individuals per sample)

and freeze-dried. The titre of FA, JH III and JHB3 were determined by adapting a protocol

from a previous study (Kai et al. 2018). In brief, the freeze-dried homogenates were transferred

to 10 ml glass centrifuge tubes containing 1 mL acetonitrile, 1 mL 0.9% (w/v) sodium chloride

solution and 10 ng of JH III-D3 as an internal standard, ultrasonicated for 1 min, and then

vortexed and extracted twice with 2 mL hexane. The hexane phase (upper layer) was removed

and transferred to a new glass vial, and then dried under nitrogen flow. The residue was

dissolved in 1 mL acetonitrile. The measurement of FA, JH III and JHB3 was determined based

on the previous method with little modifications reported by Ramirez et al. (2020). The assay

was performed on the Acquity UPLC (Waters) system connected to the AB SCIEX 5500 triple-

quadrupole mass spectrometer (Framingham, MA, USA). Isolated the target analytes on a BEH

C18 column (100 mm* ×2.1 mm, 1.7 µm particle size) maintained at 30°C. The mobile phases

were methanol (A) and 0.1% formic acid in ultrapure water (B), and the flow rate was 0.3 mL

min−1. The gradient program started at 10% A hold on for 1 min, changed to 90% A (in 1–7

min), and decreased to 10% A (7–8 min). The injection volume was 8 µL. The ion spray voltage

was set at 5500 V, 99.99% N2 was used as the desolvation/nebulizer gas, and 99.99% Ar as the

collision gas. The temperature of the block source was maintained at 500°C, while the pressure

of nebulizer gas and turbo gas was set at 50 psi. The curtain gas (CUR) pressure was 38 psi,

and the collision gas (CAD) value was set to 8. AB Sciex v1.6 analyst software was used for

controlling instruments, data acquisition and processing. The optimal MRM transitions and

other parameters of UPLC-MS/MS for the test JHs were given in Supplementary Table S6.

Transcriptomic analyses of mutants metamorphosis and between sexes

Total RNA was extracted from L3 larvae and pre-pupae of miR-277-KO and miR-34-

KO flies using TRIzol® Reagent (Thermo Fisher Scientific) and sequenced on Illumina HiSeq

platform. Raw reads were processed as above, and the processed reads were aligned using

HISAT2 (v2.0.5) (Kim et al 2019). Gene matrix count tables were generated using StringTie

(v2.1.1) (Kovaka et al 2019) with references downloaded from Flybase (version FB2023_02).

Differentially expressed genes (DEGs) were identified using iDEP 1.0 (Ge et al 2018) to

generate the hierarchical clustering heatmap, principal component analysis (PCA), enrichment
analysis and enriched pathway analysis with default parameters. The DEGs were also analysed
with ShinyGO (v0.77) (Ge et al 2020) for gene enrichment analyses. Details can be found in
Supplementary Table S7.
416

417 **Availability of data and materials**
418 The raw reads generated in this study have been deposited to the NCBI database under
419 the BioProject accessions: PRJNA974543.
420

421 **Acknowledgments**

422 This work was supported by the Hong Kong Research Grant Council General Research
423 Fund (14100420). KKC was supported by a PhD studentship supported by The Chinese
424 University of Hong Kong.

Figure 1. A) Enriched gene pathways and differentially expressed genes (DEGs) of *w*[1118] male and female flies during larvae-pupae transition; B) Differentially expressed microRNAs of *w*[1118] male and female flies in the larvae-pupae transition; C) Screening of potential microRNA candidates that can regulate sesquiterpenoid pathway gene candidates picked up by genome-wide *in silico* prediction; D) MicroRNA pull-down assay showing interactions of miR-277-3p and *Kr-h1* and *AstC-R1* transcripts in *Drosophila* S2 cells and 3[rd] instar larvae *in vivo*; E-F) Expression profiles of miR-277-3p, miR-277-5p, miR-34-3p and miR-34-5p read counts in fly embryos, larval and pupal stages; G-H) Expression level of *Kr-h1* and *AstC-R1* at larval and pupal stages and pupae stages.

Figure 2. A) Mortality of heterozygous miR-277 and miR-34 double knockout and homozygous miR-277 and miR-34 double knockout flies after eclosion within 5 days; B) *Kr-h1* and C) *AstC-R1* relative expression level in larvae male (LM), larvae female (LF), pupae male (PM) and pupae female (PF) of w^{1118}, *miR-277-KO*, and *miR-34-KO* flies; D) i. JHB3, ii. JH-III, iii. FA and iv. 20E hormone titre of w^{1118}, *miR-277-KO* and *miR-34-KO* for LM, LF, PM and PF; E) Differentially expressed genes showing up- (green) or down-regulation (red) in the sesquiterpenoid hormone biosynthesis pathway in w^{1118}, *miR-277-KO*, and *miR-34-KO* flies; G) Differentially expressed genes showing up- (green) or down-regulation (red) in the ecdysteroid hormone biosynthesis pathway in w^{1118}, *miR-277-KO*, and *miR-34-KO* flies.

Figure 3. A) Expression level of transcripts in the miR-277/miR-34 locus retrieved from Flybase (version FB2023_0); B) Enriched gene pathways in *miR-277-KO* and *miR-34-KO* flies when comparing to *w1118* at different stages in male and female; C) Differentially expressed genes showing up- (green) or down-regulation (red) in *dorsal-ventral axis formation pathway* in *miR-277-KO* and *miR-34-KO* in males and females at larvae and pupae stages. C) Genes showing downregulation in the *FoxO signalling*, *MAPK signalling*, *Hippo signalling*, and *dorsal-ventral axis formation pathways* in *miR-277-KO* and *miR-34-KO* in males and females at larvae and pupae stages.

Figure 4. A) Enriched gene pathways in males and females of *miR-277-KO* and *miR-34-KO* that are unique to one sex during larvae-pupae transition; B) Schematic summary of gene pathways enriched with differentially expressed genes during larvae-pupae transition.

Reference:

Agarwal, V., Subtelny, A. O., Thiru, P., Ulitsky, I., & Bartel, D. P. (2018). Predicting microRNA targeting efficacy in Drosophila. *Genome biology*, *19*(1), 152.

Andrews, S. (2010). FastQC: a quality control analysis tool for high throughput sequencing data. *Github*.

Anna, G., John, M., & Kannan, N. N. (2023). miR-277 regulates the phase of circadian activity-rest rhythm in Drosophila melanogaster. *Frontiers in physiology*, *14*, 1082866.

Bartel D. P. (2018). Metazoan MicroRNAs. *Cell*, *173*(1), 20–51.

Belgacem, Y. H., & Martin, J. R. (2002). Neuroendocrine control of a sexually dimorphic behavior by a few neurons of the pars intercerebralis in Drosophila. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(23), 15154–15158.

Belles, X. (2011). Origin and evolution of insect metamorphosis. *Encyclopedia of life sciences (ELS)*, 1-11.

Belles X. (2017). MicroRNAs and the Evolution of Insect Metamorphosis. *Annual review of entomology*, *62*, 111–125.

Bendena, W. G., Donly, B. C., & Tobe, S. S. (1999). Allatostatins: a growing family of neuropeptides with structural and functional diversity. *Annals of the New York Academy of Sciences*, *897*, 311–329.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, *30*(15), 2114–2120.

Bownes, M., Dübendorfer, A., & Smith, T. (1984). Ecdysteroids in adult males and females of Drosophila melanogaster. *Journal of insect physiology*, *30*(10), 823-830.

Cao, J. Q., Tong, W. S., Hiu, Y. Y., Tobe, S. S., Bendena, W. G., & Hui, J. H. (2017). The role of MicroRNAs in Drosophila regulation of insulin-like peptides and ecdysteroid signalling: where are we now?. In *Advances in Insect Physiology* (Vol. 53, pp. 55-85). Academic Press.

Caygill, E. E., & Johnston, L. A. (2008). Temporal regulation of metamorphic processes in Drosophila by the let-7 and miR-125 heterochronic microRNAs. *Current biology : CB*, *18*(13), 943–950.

Cerro-Herreros, E., Fernandez-Costa, J. M., Sabater-Arcis, M., Llamusi, B., & Artero, R.

(2016). Derepressing muscleblind expression by miRNA sponges ameliorates myotonic dystrophy-like phenotypes in Drosophila. *Scientific reports*, *6*, 36230.

Chan, K. K., Chan, T. F., Bendena, W., & Hui, J. H. L. (2023). Noncoding RNA Regulation of Hormonal and Metabolic Systems in the Fruit Fly Drosophila. *Metabolites*, *13*(2), 152.

Chen, Z. X., Sturgill, D., Qu, J., Jiang, H., Park, S., Boley, N., Suzuki, A. M., Fletcher, A. R., Plachetzki, D. C., FitzGerald, P. C., Artieri, C. G., Atallah, J., Barmina, O., Brown, J. B., Blankenburg, K. P., Clough, E., Dasgupta, A., Gubbala, S., Han, Y., Jayaseelan, J. C., … Richards, S. (2014). Comparative validation of the D. melanogaster modENCODE transcriptome annotation. *Genome research*, *24*(7), 1209–1223.

Cheong, S. P., Huang, J., Bendena, W. G., Tobe, S. S., & Hui, J. H. (2015). Evolution of Ecdysis and Metamorphosis in Arthropods: The Rise of Regulation of Juvenile Hormone. *Integrative and comparative biology*, *55*(5), 878–890.

Coyne, J. A., & Oyama, R. (1995). Localization of pheromonal sexual dimorphism in Drosophila melanogaster and its effect on sexual isolation. *Proceedings of the National Academy of Sciences of the United States of America*, *92*(21), 9505–9509.

Dambal, S., Shah, M., Mihelich, B., & Nonn, L. (2015). The microRNA-183 cluster: the family that plays together stays together. *Nucleic acids research*, *43*(15), 7173–7188.

Deshpande, P., Chen, C. Y., Chimata, A. V., Li, J. C., Sarkar, A., Yeates, C., Chen, C. H., Kango-Singh, M., & Singh, A. (2024). miR-277 targets the proapoptotic gene-hid to ameliorate Aβ42-mediated neurodegeneration in Alzheimer's model. *Cell death & disease*, *15*(1), 71.

Devineni, A. V., & Heberlein, U. (2012). Acute ethanol responses in Drosophila are sexually dimorphic. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(51), 21087–21092.

Enright, A. J., John, B., Gaul, U., Tuschl, T., Sander, C., & Marks, D. S. (2003). MicroRNA targets in Drosophila. *Genome biology*, *5*(1), R1.

Esslinger, S. M., Schwalb, B., Helfer, S., Michalik, K. M., Witte, H., Maier, K. C., Martin, D., Michalke, B., Tresch, A., Cramer, P., & Förstemann, K. (2013). Drosophila miR-277 controls branched-chain amino acid catabolism and affects lifespan. *RNA biology*, *10*(6), 1042–1056.

Fagegaltier, D., König, A., Gordon, A., Lai, E. C., Gingeras, T. R., Hannon, G. J., &

Shcherbata, H. R. (2014). A genome-wide survey of sexually dimorphic expression of
Drosophila miRNAs identifies the steroid hormone-induced miRNA let-7 as a regulator of
sexual identity. *Genetics*, *198*(2), 647–668.

Friedländer, M. R., Mackowiak, S. D., Li, N., Chen, W., & Rajewsky, N. (2012). miRDeep2
accurately identifies known and hundreds of novel microRNA genes in seven animal clades.
Nucleic acids research, *40*(1), 37–52.

Furuta, K., Fujita, N., Ibushi, T., Shiotsuki, T., Yamada, N., & Kuwano, E. (2010). Synthesis
and anti-juvenile hormone activity of ethyl 4-[(6-substituted 2, 2-dimethyl-2H-chromen-7-yl)
methoxy] benzoates. *Journal of Pesticide Science*, *35*(4), 405-411.

Garbuzov, A., & Tatar, M. (2010). Hormonal regulation of Drosophila microRNA let-7 and
miR-125 that target innate immunity. *Fly*, *4*(4), 306–311.

Ge, S. X., Jung, D., & Yao, R. (2020). ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinformatics (Oxford, England)*, *36*(8), 2628–2629.

Ge, S. X., Son, E. W., & Yao, R. (2018). iDEP: an integrated web application for differential
expression and pathway analysis of RNA-Seq data. *BMC bioinformatics*, *19*(1), 534.

Griffiths-Jones, S., Hui, J. H., Marco, A., & Ronshaugen, M. (2011). MicroRNA evolution by
arm switching. *EMBO reports*, *12*(2), 172–177.

Han, C., Peng, Q., Sun, M., Jiang, X., Su, X., Chen, J., Ma, M., Zhu, H., Ji, X., & Pan, Y.
(2022). The doublesex gene regulates dimorphic sexual and aggressive behaviors in
Drosophila. *Proceedings of the National Academy of Sciences of the United States of*
America, *119*(37), e2201513119.

Hausser, J., & Zavolan, M. (2014). Identification and consequences of miRNA-target
interactions--beyond repression of gene expression. Nature reviews. *Genetics*, *15*(9), 599–
612.

Hsu, R. J., Yang, H. J., & Tsai, H. J. (2009). Labeled microRNA pull-down assay system: an
experimental approach for high-throughput identification of microRNA-target mRNAs.
Nucleic acids research, *37*(10), e77.

Hui, J. H., Bendena, W. G., & Tobe, S. S. (2013). 2 Future Perspectives. *Juvenile Hormones*
and Juvenoids: Modeling Biological Effects and Environmental Fate, 15.

Jones, C. I., Grima, D. P., Waldron, J. A., Jones, S., Parker, H. N., & Newbury, S. F. (2013).
The 5'-3' exoribonuclease Pacman (Xrn1) regulates expression of the heat shock protein
Hsp67Bc and the microRNA miR-277-3p in Drosophila wing imaginal discs. *RNA biology*,
10(8), 1345–1355.

Kai, Z. P., Qiu, Y., Zhang, X. W., & Chen, S. S. (2021). Effects of fragrance compounds on
growth of the silkworm Bombyx mori. *PeerJ*, *9*, e11620.

Kai, Z. P., Yin, Y., Zhang, Z. R., Huang, J., Tobe, S. S., & Chen, S. S. (2018). A rapid
quantitative assay for juvenile hormones and intermediates in the biosynthetic pathway using
gas chromatography tandem mass spectrometry. *Journal of chromatography. A*, *1538*, 67–74.

Kennerdell, J. R., Liu, N., & Bonini, N. M. (2018). MiR-34 inhibits polycomb repressive
complex 2 to modulate chaperone expression and promote healthy brain aging. *Nature
communications*, *9*(1), 4188.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature biotechnology*, *37*(8),
907–915.

Kim, Y. K., Yu, J., Han, T. S., Park, S. Y., Namkoong, B., Kim, D. H., Hur, K., Yoo, M. W.,
Lee, H. J., Yang, H. K., & Kim, V. N. (2009). Functional links between clustered
microRNAs: suppression of cell-cycle inhibitors by microRNA clusters in gastric cancer.
Nucleic acids research, *37*(5), 1672–1681.

Konopova, B., & Jindra, M. (2007). Juvenile hormone resistance gene Methoprene-tolerant
controls entry into metamorphosis in the beetle Tribolium castaneum. *Proceedings of the
National Academy of Sciences of the United States of America*, *104*(25), 10488–10493.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., & Pertea, M. (2019).
Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome
biology*, *20*(1), 278.

Kozomara, A., Birgaoanu, M., & Griffiths-Jones, S. (2019). miRBase: from microRNA
sequences to function. *Nucleic acids research*, *47*(D1), D155–D162.

Kozomara, A., & Griffiths-Jones, S. (2014). miRBase: annotating high confidence
microRNAs using deep sequencing data. *Nucleic acids research*, *42*(Database issue), D68–
D73.

Lai, Y. W., Chu, S. Y., Li, J. C., Chen, P. L., Chen, C. H., & Yu, H. H. (2020). Visualization of

Endogenous Type I TGF-β Receptor Baboon in the Drosophila Brain. *Scientific reports*,

10(1), 5132.

Lai, Y. W., Chu, S. Y., Wei, J. Y., Cheng, C. Y., Li, J. C., Chen, P. L., Chen, C. H., & Yu, H. H.

(2016). Drosophila microRNA-34 Impairs Axon Pruning of Mushroom Body γ Neurons by

Downregulating the Expression of Ecdysone Receptor. *Scientific reports*, *6*, 39141.

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. L. (2009). Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. *Genome biology*, *10*(3),

R25.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *arXiv preprint arXiv:1303.3997*.

Li, R., Zhou, H., Jia, C., Jin, P., & Ma, F. (2020). Drosophila Myc restores immune

homeostasis of Imd pathway via activating miR-277 to inhibit imd/Tab2. *PLoS genetics*,

16(8), e1008989.

Liu, N., Landreh, M., Cao, K., Abe, M., Hendriks, G. J., Kennerdell, J. R., Zhu, Y., Wang, L.

S., & Bonini, N. M. (2012). The microRNA miR-34 modulates ageing and neurodegeneration

in Drosophila. *Nature*, *482*(7386), 519–523.

Lozano, J., Montañez, R., & Belles, X. (2015). MiR-2 family regulates insect metamorphosis

by controlling the juvenile hormone signaling pathway. *Proceedings of the National Academy*

of Sciences of the United States of America, *112*(12), 3740–3745.

Marco, A., Hui, J. H., Ronshaugen, M., & Griffiths-Jones, S. (2010). Functional shifts in

insect microRNA evolution. *Genome biology and evolution*, *2*, 686–696.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet. journal*, *17*(1), 10-12.

McNeill, E. M., Warinner, C., Alkins, S., Taylor, A., Heggeness, H., DeLuca, T. F., Fulga, T.

A., Wall, D. P., Griffith, L. C., & Van Vactor, D. (2020). The conserved microRNA miR-34

regulates synaptogenesis via coordination of distinct mechanisms in presynaptic and

postsynaptic cells. *Nature communications*, *11*(1), 1092.

Minakuchi, C., Namiki, T., & Shinoda, T. (2009). Krüppel homolog 1, an early juvenile

hormone-response gene downstream of Methoprene-tolerant, mediates its anti-metamorphic

action in the red flour beetle Tribolium castaneum. *Developmental biology*, *325*(2), 341–350.

Parthasarathy, R., Tan, A., & Palli, S. R. (2008). bHLH-PAS family transcription factor
methoprene-tolerant plays a key role in JH action in preventing the premature development of
adult structures during larval-pupal metamorphosis. *Mechanisms of development*, *125*(7),
601–616.

Qu, Z., Bendena, W. G., Nong, W., Siggens, K. W., Noriega, F. G., Kai, Z. P., Zang, Y. Y.,
Koon, A. C., Chan, H. Y. E., Chan, T. F., Chu, K. H., Lam, H. M., Akam, M., Tobe, S. S., &
Lam Hui, J. H. (2017). MicroRNAs regulate the sesquiterpenoid hormonal pathway in
Drosophila and other arthropods. *Proceedings. Biological sciences*, *284*(1869), 20171827.

Qu, Z., Bendena, W. G., Tobe, S. S., & Hui, J. H. L. (2018). Juvenile hormone and
sesquiterpenoids in arthropods: Biosynthesis, signaling, and role of MicroRNA. *The Journal*
of steroid biochemistry and molecular biology, *184*, 69–76.

Qu, Z., Yiu, W. C., Yip, H. Y., Nong, W., Yu, C. W. C., Lee, I. H. T., Wong, A. Y. P., Wong, N.
W. Y., Cheung, F. K. M., Chan, T. F., Lau, K. F., Zhong, S., Chu, K. H., Tobe, S. S., Ferrier,
D. E. K., Bendena, W. G., & Hui, J. H. L. (2020). Micro-RNA Clusters Integrate
Evolutionary Constraints on Expression and Target Affinities: The miR-6/5/4/286/3/309
Cluster in Drosophila. *Molecular biology and evolution*, *37*(10), 2955–2965.

Ramirez, C. E., Nouzova, M., Michalkova, V., Fernandez-Lima, F., & Noriega, F. G. (2020).
Common structural features facilitate the simultaneous identification and quantification of the
five most common juvenile hormones by liquid chromatography-tandem mass spectrometry.
Insect biochemistry and molecular biology, *116*, 103287.

Riddiford L. M. (2012). How does juvenile hormone control insect metamorphosis and
reproduction?. *General and comparative endocrinology*, *179*(3), 477–484.

Riddiford, L. M., & Ashburner, M. (1991). Effects of juvenile hormone mimics on larval
development and metamorphosis of Drosophila melanogaster. *General and comparative*
endocrinology, *82*(2), 172–183.

Riddiford, L. M., Truman, J. W., Mirth, C. K., & Shen, Y. C. (2010). A role for juvenile
hormone in the prepupal development of Drosophila melanogaster. *Development*
(Cambridge, England), *137*(7), 1117–1126.

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinformatics (Oxford, England)*, *26*(1), 139–140.

Sempere, L. F., Dubrovsky, E. B., Dubrovskaya, V. A., Berger, E. M., & Ambros, V. (2002). The expression of the let-7 small regulatory RNA is controlled by ecdysone during metamorphosis in Drosophila melanogaster. *Developmental biology*, *244*(1), 170–179.

Sokol, N. S., Xu, P., Jan, Y. N., & Ambros, V. (2008). Drosophila let-7 microRNA is required for remodeling of the neuromusculature during metamorphosis. *Genes & development*, *22*(12), 1591–1596.

Srinivasan, A. R., Tran, T. T., & Bonini, N. M. (2022). Loss of miR-34 in Drosophila dysregulates protein translation and protein turnover in the aging brain. *Aging cell*, *21*(3), e13559.

Stark, A., Brennecke, J., Russell, R. B., & Cohen, S. M. (2003). Identification of Drosophila MicroRNA targets. *PLoS biology*, *1*(3), E60.

Stay, B., & Tobe, S. S. (2007). The role of allatostatins in juvenile hormone synthesis in insects and crustaceans. *Annual review of entomology*, *52*, 277–299.

Tan, H., Poidevin, M., Li, H., Chen, D., & Jin, P. (2012). MicroRNA-277 modulates the neurodegeneration caused by Fragile X premutation rCGG repeats. *PLoS genetics*, *8*(5), e1002681.

Tanzer, A., & Stadler, P. F. (2004). Molecular evolution of a microRNA cluster. *Journal of molecular biology*, *339*(2), 327–335.

Testa, N. D., Ghosh, S. M., & Shingleton, A. W. (2013). Sex-specific weight loss mediates sexual size dimorphism in Drosophila melanogaster. *PloS one*, *8*(3), e58936.

Tobe, S. S., & Bendena, W. G. (1999). The regulation of juvenile hormone production in arthropods. Functional and evolutionary perspectives. *Annals of the New York Academy of Sciences*, *897*, 300–310.

Truman J. W. (2019). The Evolution of Insect Metamorphosis. *Current biology : CB*, *29*(23), R1252–R1268.

Truman, J. W., & Riddiford, L. M. (2002). Endocrine insights into the evolution of metamorphosis in insects. *Annual review of entomology*, *47*, 467–500.

Truman, J. W., & Riddiford, L. M. (1999). The origins of insect metamorphosis. *Nature*,

401(6752), 447–452.

Tsang, S. S. K., Law, S. T. S., Li, C., Qu, Z., Bendena, W. G., Tobe, S. S., & Hui, J. H. L.

(2020). Diversity of Insect Sesquiterpenoid Regulation. *Frontiers in genetics*, *11*, 1027.

Ureña, E., Chafino, S., Manjón, C., Franch-Marro, X., & Martín, D. (2016). The Occurrence

of the Holometabolous Pupal Stage Requires the Interaction between E93, Krüppel-Homolog

1 and Broad-Complex. *PLoS genetics*, *12*(5), e1006020.

Ventura, A., Young, A. G., Winslow, M. M., Lintault, L., Meissner, A., Erkeland, S. J.,

Newman, J., Bronson, R. T., Crowley, D., Stone, J. R., Jaenisch, R., Sharp, P. A., & Jacks, T.

(2008). Targeted deletion reveals essential and overlapping functions of the miR-17 through

92 family of miRNA clusters. *Cell*, *132*(5), 875–886.

Wang, W., Kidd, B. J., Carroll, S. B., & Yoder, J. H. (2011). Sexually dimorphic regulation of

the Wingless morphogen controls sex-specific segment number in Drosophila. *Proceedings of

the National Academy of Sciences of the United States of America*, *108*(27), 11139–11144.

Wen, D., Rivera-Perez, C., Abdou, M., Jia, Q., He, Q., Liu, X., Zyaan, O., Xu, J., Bendena,

W. G., Tobe, S. S., Noriega, F. G., Palli, S. R., Wang, J., & Li, S. (2015). Methyl farnesoate

plays a dual role in regulating Drosophila metamorphosis. *PLoS genetics*, *11*(3), e1005038.

Wood, D. E., Lu, J., & Langmead, B. (2019). Improved metagenomic analysis with Kraken 2.

Genome biology, *20*(1), 257.

Xiong, X. P., Kurthkoti, K., Chang, K. Y., Li, J. L., Ren, X., Ni, J. Q., Rana, T. M., & Zhou,

R. (2016). miR-34 Modulates Innate Immunity and Ecdysone Signaling in Drosophila. *PLoS

pathogens*, *12*(11), e1006034.

Zipper, L., Batchu, S., Kaya, N. H., Antonello, Z. A., & Reiff, T. (2022). The MicroRNA

miR-277 Controls Physiology and Pathology of the Adult Drosophila Midgut by Regulating

the Expression of Fatty Acid β-Oxidation-Related Genes in Intestinal Stem Cells. *Metabolites*,

12(4), 315.