Title

Firing of *Streptomyces coelicolor* contractile injection system requires the membrane protein CisA

Authors

Bastien Casu[1], Joseph W. Sallmen[2], Peter E. Haas[1], Pavel Afanasyev[1], Jingwei Xu[1], Susan Schlimpert[2, *], Martin Pilhofer[1, *]

Affiliations

[1]Department of Biology, Institute of Molecular Biology & Biophysics, Eidgenössische Technische Hochschule Zürich, Otto-Stern-Weg 5, 8093 Zürich, Switzerland

[2]John Innes Center, Department of Molecular Microbiology, Norwich Research Park, Norwich, NR4 7UH, United Kingdom

Contact Info

* Correspondence: Susan.Schlimpert@jic.ac.uk (S.S.) and pilhofer@biol.ethz.ch (M.P.)

Abstract

Bacterial contractile injection systems (CIS) are phage tail-like macromolecular complexes that mediate cell-cell interactions by injecting effector proteins into target cells. CIS^{Sc} from *Streptomyces coelicolor* are distinct because they induce regulated cell death under stress, impacting the organism's life cycle. It remains unknown whether CIS^{Sc} require accessory proteins to function.

Here we characterize the putative membrane adaptor CisA, a conserved factor in CIS^{Sc} gene clusters across *Streptomyces* species. We show by cryo-electron tomography imaging and *in vivo* assays that CIS^{Sc} contraction and function depend on CisA. Using single-particle cryo-electron microscopy, we provide an atomic model of the extended CIS^{Sc} apparatus; however, CisA is not part of the complex. Instead, our findings show that CisA is a membrane protein with a cytoplasmic N-terminus predicted to interact with CIS^{Sc} components, thereby providing a possible mechanism for mediating CIS^{Sc} recruitment to the membrane and subsequent firing.

Our work shows that CIS function in multicellular bacteria is distinct from Type 6 Secretion Systems and extracellular CIS, and possibly evolved due to the role of CIS^{Sc} in regulated cell death.

Keywords: bacterial development/cellular microbiology/contractile phage tail/ghost cells/multicellularity

Introduction

Bacteria employ different types of contractile injection systems (CIS) to mediate cell-cell interactions or to function in regulated cell death[1]. CIS are evolutionarily and structurally related to contractile tails of bacteriophages and are comprised of core modules including a baseplate, a contractile sheath, and an inner tube [2,3]. CIS firing is triggered by a conformational change within the baseplate[4,5] and results in the contraction of the sheath, which in turn propels the inner tube along with the spike tip. Tube expulsion facilitates the release of associated effector proteins into target cells or into the extracellular space[6,7].

Bioinformatic analyses have shown that CIS are conserved across diverse microbial phyla, including Gram-negative and Gram-positive bacteria, as well as archaea[8,9]. Based on their distinct modes of action, CIS can be categorized into two main groups: intracellular type VI secretion systems (T6SS) and extracellular CIS (eCIS). T6SS are abundant among Gram-negative bacteria and in some archaea, and they are anchored to the cytoplasmic membrane during assembly and firing[10–12]. A crucial component of the T6SS is the baseplate, which acts as a nucleus for T6SS assembly and mediates the binding of the T6SS particles to the cytoplasmic membrane until contraction. Upon contraction, the spike and inner tube are propelled out of the cell and into an adjacent cell or into the medium. eCIS, on the other hand, are assembled as free-floating particles in the cytoplasm and are subsequently released into the extracellular space upon lysis of the producer cell [13–16]. Following the release, eCIS attach to the surface of target cells via their tail fibers, followed by contraction and puncturing of the target cell envelope.

Besides T6SS and eCIS, there is accumulating evidence of additional CIS assemblies with distinct modes of action in bacteria. For example, in multicellular cyanobacteria, CIS are anchored to the intracellular thylakoid membrane stacks via an extension of the baseplate components and have been proposed to function in cell lysis and the formation of "ghost cells" in response to stress conditions[17]. Furthermore, several recent studies reported the production of free-floating cytoplasmic CIS particles in the multicellular Gram-positive *Streptomyces* bacteria, affecting cellular development through presumably slightly different mechanisms[18–22].

Streptomyces are best known for producing a plethora of medically and industrially important

secondary metabolites, including molecules with antimicrobial, antifungal, anticancer or

immunosuppressive properties. The production of these molecules is tightly coordinated with

the *Streptomyces* developmental life cycle that encompasses two filamentous cell types,

including vegetative hyphae that grow by tip extension and branching to scavenge for nutrients,

as well as reproductive (aerial) hyphae that eventually differentiate into chains of spores. These

spores can then be dispersed to restart the life cycle[23,24]. Using *Streptomyces coelicolor* as a

model, we and others previously demonstrated that *S. coelicolor* CIS (CIS^{Sc}) mediate a form

of regulated cell death in response to exogenous stress and are also activated by unknown

cellular signals, thereby impacting the timing of sporulation and secondary metabolite

production (Figure 1a)[18,19].

It remains unclear whether CIS^{Sc} contraction occurs while the CIS^{Sc} particles are free-floating

in the cytoplasm or whether this is dependent on an interaction with the membrane. Moreover,

CIS encoded in *Streptomyces* genomes lack clear homologs for tail fiber components or a T6SS

trans-envelope complex, raising the question of how CIS^{Sc} particles could potentially interact

with the cytoplasmic membrane. We recently noted an uncharacterized protein[18,19], SCO4242,

that is encoded just downstream of the baseplate components in the CIS^{Sc} gene cluster, a

location that usually encodes tail fibers from conventional eCIS[13,16,25]. Importantly, SCO4242

is widely conserved in *Streptomyces cis* gene clusters[18,19]. We will therefore refer to the gene

product of SCO4242 as CisA for "CIS^{Sc}-associated protein A." Here we set out to assess the

importance of CisA for CIS^{Sc} function and its putative role as a membrane anchor.

Results and Discussion

CisA is required for CIS[Sc] contraction *in situ*

To explore the role of CisA (SCO4242) in CIS[Sc] contraction and function, we first generated an *S. coelicolor* Δ*cisA* null mutant. Negative stain electron microscopy (EM) imaging of CIS[Sc] particles that were purified from Δ*cisA* mutant cells showed assemblies in the contracted conformation, similar to the CIS particles purified from the wildtype (WT) (Fig. 1b/c). We then imaged vegetative hyphae of the WT, the Δ*cisA* mutant and the complemented mutant (Δ*cisA/cisA*[+]) by cryo-electron tomography (cryoET) (270 tomograms in total, n=3 experiments). The observed CIS[Sc] particles in intact hyphae of these three imaged strains did not reveal discernable differences, indicating that CisA is not required for CIS[Sc] assembly. As observed previously, we detected three states of cellular integrity of *Streptomyces* hyphae, including 'intact hyphae', 'partially lysed hyphae' and 'ghost cells' (EV1a-c)[18]. Importantly, *cisA*-deficient hyphae only contained CIS[Sc] particles in the extended conformation irrespective of the cellular integrity (Fig. 1e, EV1a-d). This is in stark contrast to 'ghost cells' derived from hyphae of the WT and the Δ*cisA/cisA*[+] complementation strain, in which CIS[Sc] particles were predominantly found in the contracted conformation (Fig. 1d/f). These results indicate that CisA is required for the contraction of CIS[Sc] *in situ*.

CryoEM structure of the extended CIS[Sc] assembly

The CIS[Sc] gene cluster consists of 19 predicted open reading frames (accessions *SCO4242-4260*) (Fig. 2a), including genes with sequence similarities to potential CIS structural components (*cis1-cis16*), as well as additional genes with unknown functions (including *cisA*)[18,19]. To determine whether CisA is an integral component of CIS[Sc], we set out to elucidate the high-resolution structure of purified CIS[Sc] particles from a non-contractile CIS[Sc] mutant[18]. The purified assemblies were then subjected to single particle cryo-electron microscopy (cryoEM) for structure determination (EV2), yielding the first high-resolution model of the CIS[Sc] in its extended conformation, which also presents the first example of a CIS IId subtype[8].

Our model showed that extended CIS[Sc] particles share the conserved CIS architecture, including three modules: cap, sheath-tube and baseplate (Fig. 2b). We processed the dataset using an established workflow that was previously employed to produce the atomic model for

other CIS (EV2)[13,17]. The resulting maps of the baseplate and the cap reached resolutions of

3.5 and 3.4 Å, respectively. The quality of the final maps from the two modules enabled *de*

novo structural modelling (EV3, Table EV1). The model was further refined by integrating

symmetry-related protein subunits into a complete model of the entire CIS^{Sc} (Fig. 2b).

The final model of CIS^{Sc} comprised 783 polypeptide chains that assemble into a bullet-shaped

particle measuring 268 nm in length (Fig. 2b/c). The distal end of the CIS^{Sc} is capped by a tail

terminator complex (Fig. 2c/e). This cap comprises one protein, Cis16, forming a C6-

symmetrical complex that terminates the inner tube and sheath, respectively. This minimal

architecture resembles the cap complexes of PVC or Pyocin R2 CISs, but it is different from

the more complex AlgoCIS from *A. machipongonensis* and AFP from *S. entomophila*

(EV4)[13,16,25,26]. The structures of the CIS^{Sc} sheath-tube module have been discussed

previously[18]. Here we found that the module consists of 61 sheath layers (Cis2-L_1–L_{61}) and 60

tube layers (Cis1-L_1-L_{60}) (Fig. 2b/d).

The baseplate module consists of a heteromeric assembly of the proteins Cis1b/7/8/9/11/12

(Fig. 2b/c). The symmetry transition from the inner tube (Cis1a, C6) to the VgrG-like spike

(Cis8, C3) is adapted by two layers of the tube initiators Cis1b and Cis7. The first layer of the

sheath (Cis2) is bound to the conserved gp25-like protein Cis9, which connects to the baseplate

'wedges' composed of Cis11 and Cis12 (Fig. 2b/c). Importantly, the CIS^{Sc} 'wedges' consist of

two copies of Cis11 and one copy of Cis12 (Fig. 2f), resembling the baseplate of Pyocin R2[26].

This is in contrast to other previously reported eCIS, whose baseplate wedge comprises only

one Cis11/12 component (AlgoCIS, AFP, PVC) and a Cis11 extension forming a cage around

the spike (AlgoCIS) (EV4)[13,16,25].

Out of the 19 predicted open reading frames in the CIS^{Sc} gene cluster (Fig. 2a), nine encoded

proteins were present in the final reconstruction, all with atomic models built (Fig. 2b/c).

However, we did not detect a density for CisA in our model, suggesting that CisA is not part

of purified CIS^{Sc} particles but may interact via a different mechanism. This is consistent with

mass spectrometry results that indicate the absence of CisA peptides from purified CIS^{Sc}

particles (Fig. 2a)[18]. We speculate that a CIS^{Sc}-CisA interaction may not be preserved by the

purification protocol, and/or the interaction is too transient and only occurs under certain

growth conditions. To better understand how CisA may be involved in mediating CIS^{Sc}

function, we set out to further characterize CisA *in vivo*.

CisA is a bitopic protein

Previous bioinformatic analyses[18,19] and Alphafold2 modeling of CisA (Fig. 3a) suggest that CisA consists of a largely unstructured N-terminal region, a transmembrane segment, and a conserved immunoglobulin-like (IgG) domain at the C-terminus. To characterize the intracellular localization of CisA, we first generated a strain in which *cisA* was fused to the N-terminus of *mcherry* and expressed *in trans* from a constitutive promoter in the *S. coelicolor* Δ*cisA* mutant. Using fluorescence light microscopy, we consistently observed a CisA-mCherry fluorescence signal along the hyphal periphery, indicative of a membrane-associated protein (Fig. 3b). In contrast, no fluorescence was observed in WT hyphae carrying an empty plasmid (Fig. 3c). In furtherance to this, we wanted to provide more support that CisA localized to the membrane and so, we performed cellular fractionation experiments to separate soluble proteins from membrane proteins using whole cell lysates from the *S. coelicolor* WT or from the Δ*cisA* mutant that was complemented with a *cisA-3xFLAG* fusion expressed *in trans* from a constitutive promoter. Our analysis confirmed that CisA-3xFLAG co-sedimented with the cell membranes, while the cytoplasmic transcription factor WhiA could only be detected in the soluble protein fraction (Fig. 3d).

Next, we investigated the topology of CisA in the membrane by using a dual *phoA-lacZ*α reporter system in *E. coli*. The reporter activity can be directly visualized on indicator plates based on the complementary activities of the cytoplasmic reporter β-galactosidase LacZ (magenta coloration) and the periplasmic reporter alkaline phosphatase PhoA (blue coloration)[27]. We engineered *E. coli* strains that expressed a C-terminal fusion of CisA variants to the PhoA-LacZα reporter, including full-length CisA and CisA variants that carried either a deletion in the predicted transmembrane domain and/or the putative periplasmic domain (Fig. 3e). In addition, we included three controls, including cells 1) carrying the empty reporter plasmid, resulting only in cytoplasmic β-galactosidase activity, and expressing a PhoA/LacZ fusion to 2) the *Streptomyces* cytoplasmic cell division protein SepH and 3) the periplasmic domain of the *Streptomyces* membrane protein RsbN[28,29]. The results of this assay confirmed that CisA is a bitopic protein with an N-terminus (amino acids 1-285) located in the cytoplasm and a C-terminus (amino acids 310-468) present in the periplasm of *E. coli* (Fig. 3f).

176   Collectively, our results demonstrate that CisA is a membrane protein comprised of a flexible,

177   cytoplasmic N-terminal domain and a C-terminal globular IgG-like domain that is exposed to

178   the outside of the cytoplasmic membrane.

CisA is essential for the cellular function of CIS[Sc]

We previously established a fluorescence-based assay to demonstrate that the production of

functional CIS[Sc] particles results in a significantly reduced viability of *Streptomyces* hyphae

that were exposed to exogenous stress, including treatment with the membrane-pore forming

antibiotic nisin, UV radiation or the protonophore carbonyl cyanide 3-chlorophenylhydrazone

(CCCP)[18]. The assay is based on the measurement of the ratio of two fluorescent markers,

namely cytoplasmic sfGFP produced by viable intact hyphae and the membrane dye FM5-95

to detect intact as well as partially lysed hyphae. Using this assay, we tested whether CisA was

essential for CIS[Sc]-mediated cell death. We therefore determined the relative viability of the

WT *cisA*, the Δ*cisA* and the complemented *cisA* (Δ*cisA/cisA⁺*) strains expressing cytosolic

sfGFP. We compared cells that were exposed to a sublethal concentration of nisin (1 µg/ml for

90 min) to cells that were not exposed to antibiotic stress. All three strains were found to

express CIS[Sc] assemblies under these growth conditions (EV6).

In the absence of nisin stress, all three strains exhibited a similar sfGFP/FM5-95 ratio,

indicating no significant difference in viability (Fig. 4a/c). As previously observed[18], the

viability of the WT was dramatically reduced in response to nisin stress. In contrast, the

viability of Δ*cisA* hyphae was unaffected by nisin stress and comparable to the viability of

untreated cells. This phenotype could be completely reversed by complementing the Δ*cisA*

deletion *in trans* (Fig. 4b/d). Notably, the Δ*cisA* strain phenocopies CIS[Sc] knockout and non-

contractile mutants that we analyzed previously[18]. Thus, we conclude that CisA is required for

CIS[Sc]-mediated cell death upon stress.

To further test the dependence of CIS[Sc] function on CisA, we conducted additional *in vivo*

assays. Previous studies showed that the absence of functional CIS[Sc] impacted the timely

differentiation of *Streptomyces* hyphae into chains of spores, resulting in accelerated cellular

development and reduced secondary metabolite production[18,19]. Here we repeated these

analyses using the WT, the Δ*cisA* mutant and the complemented mutant (Δ*cisA/cisA⁺*). All

strains consistently completed their life cycle and synthesized spores after 96 h of growth on

solid medium (Fig. 4e). Importantly, unlike the WT and the Δ*cisA/cisA*⁺strain, Δ*cisA* mutant

colonies sporulated markedly earlier (72 h). These findings were further corroborated by

quantifying the number of spores produced by each strain under the same experimental

conditions (Fig. 4f). Finally, we also tested the production of the two characteristic pigmented

secondary metabolites of *S. coelicolor*, actinorhodin (blue)[30] and undecylprodigiosin (red)[31].

Compared to the WT and the Δ*cisA/cisA*⁺ strain, the Δ*cisA* mutant showed significantly

reduced levels of secondary metabolite production (EV7). Our data therefore support the

conclusion that the cellular function of CIS^{Sc} in *S. coelicolor* is dependent on CisA and its

absence impacts the timely cellular and chemical differentiation of *S. coelicolor* hyphae.

Conclusion

Here we identify CisA as an essential membrane-associated factor that mediates the cellular function of CIS[Sc] particles in *Streptomyces coelicolor*. In the absence of CisA, CIS[Sc] are not able to contract in the cellular context (Fig. 1). Similar to a CIS[Sc] deletion or non-contractile mutant[18], this decreases the induction of regulated cell death upon stress, which in turn affects cellular development and secondary metabolite production (Fig. 5). The mechanistic aspects of this distinct mode of action of a CIS are discussed further below.

A conserved feature of all previously studied CIS is the requirement of the contractile apparatus to bind to a membrane before firing. This can be achieved in different ways, leading to the classification of CIS into different modes of action. First, T6SS are anchored to the cytoplasmic membrane of the host cell by a trans-envelope complex spanning from the cytoplasm to the inner membrane, periplasm, and outer membrane[32–34]. Second, contractile phages and eCIS bind to the surface of their target cell via tail fibers, which are typically connected to the baseplate and transmit the signal for firing to the baseplate[13,16,35]. Third, tCIS are anchored to thylakoid membrane stacks in cyanobacteria by an extension of their baseplate[36]. Our data suggest that CisA may be required to mediate an interaction of CIS[Sc] with the cytoplasmic membrane prior to firing. This interaction may occur directly via the binding of the CIS[Sc] particle to the membrane protein CisA, or through the interaction with a yet unknown factor. We performed an *in silico* prediction of protein-protein interactions between CisA and CIS[Sc] components using Alphafold2-Multimer. Interestingly, this analysis identified Cis11 as a significant hit, suggesting the possibility of a direct interaction of CisA with Cis11 (EV8a). Such a protein-protein interaction would be consistent with our cryoEM structure, showing a peripheral surface-exposed position of Cis11 within the baseplate (EV8b). We were unable, however, to confirm a direct interaction between CisA and Cis11 *in vivo* by co-immunoprecipitation with and without nisin stress. We did also not detect CisA in our CIS[Sc] purifications that were used for structure determination (Fig. 2). Importantly, however, by proteomics we did detect CisA peptides in a crude purification of CIS[Sc] from nisin-stressed cells (Table EV2). This would be consistent with a transient and/or short-lived interaction of CIS[Sc] particles with CisA triggered by an exogenous or cellular signal. In agreement with this idea, we rarely observed extended CIS[Sc] assemblies that were associated with the membrane via their baseplate in WT cells (Fig. 1d).

We speculate that CisA may not only serve as a mediator for the interaction of CIS^{Sc} with the
cytoplasmic membrane but also as a sensor and checkpoint for inducing cell death only under
specific stress conditions or in response to cellular signals. The CisA C-terminal IgG-like
domain that is localized outside the cytoplasm and in proximity of the Gram-positive cell wall
may sense stress signals that could potentially trigger conformational changes in CisA which
in turn could mediate the recruitment of CIS^{Sc} assemblies to the cytoplasmic membrane,
followed by firing. The fatal outcome of firing for the producing cell may have driven the
evolution of a stepwise mechanism involving a checkpoint that prevents self-inflicted cell death
by accidental firing. Such a mechanism would keep the CIS^{Sc} in a free-floating, non-functional
and safe state in the cytoplasm.

METHODS

Bacterial strains, plasmids, and oligonucleotides

Bacterial strains, plasmids, and oligonucleotides used in this study are listed in Supplementary Tables 3-4. Plasmids were generated using either standard restriction-ligation or assembled using the Gibson Assembly Master Mix (NEB) or *In Vivo* Assembly. All plasmids were verified by DNA sequencing. *Escherichia coli* strains were cultured in LB, SOB, or DNA medium. *E. coli* strains TOP10, DH5α and NEB5α were used to propagate plasmids and cosmids, *E. coli* strain BW25113/pIJ790 for recombineering cosmids and *E. coli* ET12567/pUZ8002 for interspecies conjugation. When required, media was supplemented with antibiotics at the following concentrations: 100 µg/ml carbenicillin, 50 µg/ml apramycin, 50 µg/ml kanamycin, 50 µg/ml hygromycin.

Streptomyces coelicolor strains were cultivated in LB, TSB, TSB-YEME, or R2YE liquid medium at 30 °C in baffled flasks or flasks with springs, at 250 rpm or grown on LB, SFM, R2YE medium solidified with 1.5 % (w/v) Difco agar[37]. Antibiotics were added at the following concentrations: 25 µg/ml apramycin, 50 µg/ml kanamycin, 25 µg/ml hygromycin, 12.5 µg/ml nalidixic acid.

Construction of the *S. coelicolor* Δ*cisA* mutant

The λ RED homologous recombination system was used to isolate gene replacement mutations using PCR-directed mutagenesis (ReDirect) of the *S. coelicolor* cosmid StD8A containing the *cis* gene cluster[38,39]. The *cisA* coding sequence (*SCO4242*) was replaced with the *aac3(IV)-oriT* resistance cassette from pIJ773. The mutant cosmid (pSS684) was introduced into *E. coli* ET12567/pUZ8002, followed by conjugation into *S. coelicolor* M145. Exconjugants that had successfully undergone double-homologous recombination were identified by screening for apramycin resistance and kanamycin sensitivity. The deletion of *cisA* was subsequently verified by PCR.

Sheath preparation of CIS^Sc

S. coelicolor strains were grown in 30 ml TSB, TSB-YEME or R2YE liquid medium for 48 h. Cells were pelleted by centrifugation (7,000 x g, 10 min, 4 °C), resuspended in 5 ml lysis buffer

(150 mM NaCl, 50 mM Tris-HCl, 0.5× CellLytic B (Sigma-Aldrich), 1 % Triton X 100, 200

µg/ml lysozyme, 50 µg/ml DNAse I, pH 7.4), and incubated for 1 h at 37 °C. Cell debris was

removed by centrifugation (15,000 × g, 15 min, 4 °C) and cleared lysates were subjected to

ultra-centrifugation (150,000 × g, 1 h, 4 °C). Pellets were resuspended in 150 µl resuspension

buffer (150 mM NaCl, 50 mM Tris-HCl, supplemented with protease inhibitor cocktail

[Roche], pH 7.4). CIS^{Sc} sheath preparations were analyzed by negative stain EM imaging[40] and

mass spectrometry at the Functional Genomics Center Zürich.

To purify non-contractile CIS^{Sc} particles from *S. coelicolor (*SS393*)*, cleared cell lysates were

subjected to ultracentrifugation using a sucrose cushion (20 mM Tris pH 8.0, 150 mM NaCl,

50 mM EDTA, 1 % Triton X 100, 50 % [w/v] sucrose)[18]. The sucrose cushion alongside 1 mm

of liquid above and residual bacterial contaminants were removed by centrifugation at 15,000

x g. Samples were then subjected to a second round of ultracentrifugation without a sucrose

cushion (150,000 × g, 1 h, 4 °C) and resulting cell pellets were resuspended in buffer (50 mM

Tris, 150 mM NaCl). These crude samples were then purified further by gradient

ultracentrifugation (10 %-50 % continuous gradient made with BIOCOMP gradient master IP

model 107 gradient forming instrument) at 100,000 x g for 1 h using a SW55 Ti rotor. The

gradient was analyzed in 11 fractions of 500 µl and checked by negative stain EM. The

fractions, which contained CIS^{Sc} particles, were pooled and used for further experiments.

Negative stain electron microscopy

303 4 µl of purified CIS^{Sc} sheath particles were adsorbed to glow-discharged, carbon-coated copper

304 grids (Electron Microscopy Sciences) for 60 s, washed twice with milli-Q water and stained

305 with 2 % phosphotungstic acid for 45 s. The grids were imaged at room temperature using a

306 Thermo Fisher Scientific Morgagni transmission electron microscope (TEM) operated at

307 80 kV.

Mass spectrometry analysis

To confirm the presence of predicted CIS^{Sc} components, isolated sheath particles were

subjected to liquid chromatography–mass spectrometry analysis (LC–MS/MS). These

experiments were conducted at the Functional Genomics Center Zürich. First, the samples were

digested with 5 µl of trypsin (100 ng/µl in 10 mM HCl) and microwaved for 30 min at 60 °C.

The samples were then dried, dissolved in 20 µl ddH₂0 with 0.1 % formic acid, diluted in 1:10

and transferred to autosampler vials for liquid chromatography with tandem mass spectrometry

analysis. A total of 1 µl was injected on a nanoAcquity UPLC coupled to a Q-Exactive mass

spectrometer (ThermoFisher). Database searches were performed by using the Mascot

swissprot and tremble_streptomycetes search programs. For search results, stringent settings

have been applied in Scaffold (1 % protein false discovery rate, a minimum of two peptides

per protein, 0.1 % peptide false discovery rate). The results were visualized by Scaffold

software ([Proteome Software Inc.], Version 4.11.1).

Protein gel electrophoresis and western blotting

For general protein analysis, protein samples were boiled at 95 °C for 5 min and separated by

a 4–20% Mini-PROTEAN® TGX Stain-Free™ SDS PAGE (BioRad) or 12 % Tris-Glycine

SDS PAGA (Invitrogen) and visualized using InstantBlue Coomassie Protein Stain or

ReadyBlue Protein Gel Stain (Sigma Aldrich).

For western blotting, cells were grown in biological triplicate for 48 h in TSB, 2 ml aliquots of

each culture were pelleted and washed once with 1x PBS. Cell pellets were resuspended in 0.4

ml of sonication buffer (20 mM Tris pH 8.0, 5 mM EDTA, 1x EDTA-free protease inhibitors

[Sigma Aldrich]) and subjected to sonication at 4.5-micron amplitude for 7 cycles of 15 s on /

15 s off. Samples were centrifuged at 17,000 x *g* for 15 min at 4 °C. Protein concentration in

cleared cell lysate supernatants was determined by Bradford Assay (Biorad). Equivalent total

protein concentrations (1 mg/ml) were assayed using the semi-dry western blot transfer. The

gel was transferred to a PVDF membrane (Biorad) and probed with the appropriate antibody

diluted in TBS-T or using the iBIND buffer system (Thermo Fisher): Rabbit anti-FLAG

([Sigma Aldrich] F7425, 1:2500), Rabbit anti-WhiA ([41], 1:2500), and Goat Anti-Rabbit-HRP

(Abcam Ab6721, 1:5000). To visualize HRP-conjugated antibodies, membranes were

incubated with SuperSignal West Femto ECL solution (Thermo Fisher).

Cellular fractionation

To fractionate membrane and soluble proteins from *S. coelicolor* (WT, SS576), cells were
grown for 48 h in 50 ml TSB/YEME and harvested by centrifugation. Cell pellets were
resuspended in 1/10 volume of lysis buffer (0.2 M Tris-HCl, pH 8, 10 mg/mL lysozyme, and
1× EDTA-free protease inhibitors; [Roche]), incubated for 30 min at 37 °C, and then briefly
cooled on ice before lysed by sonication (11 x 15 sec on/off at 50 % power at 8 microns on
ice). Cell debris was removed by centrifugation at 16,000 × g for 20 min and supernatant was
subsequently subjected to two additional rounds of centrifugation. The cleared cell lysate was
subjected to ultracentrifugation for 1 h at 150,000 × g at 4 °C to separate the soluble proteins
from membrane proteins. Sedimented membrane proteins were resuspended and washed twice
in 1 volume of wash buffer (60 mM Tris-HCl, 200 mM NaCl, pH 8, 0.2 mM EDTA, and 0.2
M sucrose) at 150,000 × g at 4 °C for 1 h. The wash step was repeated one final time with the
wash buffer containing 8 M urea to remove traces of membrane-associated proteins[42]. The final
pellet was dissolved in 1/10 of the initial volume with wash buffer (no urea). Equi-volume
amounts of fractions were mixed with 2x SDS sample buffer and analyzed by immunoblotting.

Membrane protein topology analysis in *E. coli*

The coding sequence of CisA was inserted in the dual pho-lac reporter plasmid pKTop, which
consists of an *E. coli* alkaline phosphatase fragment $PhoA_{22-472}$ fused in frame after the α-
fragment of β-galactosidase $LacZ_{4-60}$ (Table EV3)[27]. For membrane protein topology, *E. coli*
TG1 was transformed with the pKTop and derivatives. Membrane protein topology was
assayed by plating the resulting reporter strains on dual-indicator plates containing LB agar
supplemented with 80 μg/ml 5-bromo-4-chloro-3-indolyl phosphate disodium salt (X-Phos)
([Sigma Aldrich], RES1364C-A101X) and 100 μg/mL 6-chloro-3-indolyl-β-d-galactoside
(Red-Gal) ([Sigma Aldrich], B6149) as indicators, 1 mM IPTG, and 50 μg/ml kanamycin. A
periplasmic or extracellular location of the reporter fusion results in higher alkaline
phosphatase activity (blue color), whereas a cytosolic location of the reporter leads to higher
β-galactosidase activity (magenta color). Plates were incubated for two days at 37 °C and
scanned.

Vitrified sample preparations

For single-particle cryoEM (SPA), the *S. coelicolor* non-contractile CIS particles were purified as described above and vitrified using a Vitrobot Mark IV (Thermo Fisher Scientific). 4 µl of samples were applied twice on glow-discharged 200 mesh Quantifoil copper grids (R 2/2) which were manually coated with a layer of 1 nm carbon. Grids were blotted for 5.5 s and plunge-frozen in liquid ethane-propane mix (37 %/63 %). Frozen grids were stored in liquid nitrogen until loading into the microscope.

For cryo-electron tomography (cryoET), *Streptomyces* cells were mixed with 10 nm Protein A conjugated colloidal gold particles (1:10 v/v, [Cytodiagnostics]) and 4 µl of the mixture was applied to a glow-discharged holey-carbon copper EM grid (R2/1 or R2/2, [Quantifoil]). The grid was automatically blotted from the backside for 4-6 s in a Mark IV Vitrobot by using a Teflon sheet on the front pad, and plunge-frozen in a liquid ethane-propane mixture (37 %/63 %) cooled by a liquid nitrogen bath.

SPA data collection and image processing

44,925 movies were collected on Titan Krios microscope (Thermo Fisher Scientific), operated at 300 keV equipped with K3 direct electron detector, operating in counting mode, and using a slit width of 20 eV on a GIF-Quantum energy filter (Gatan). The automated data collection was conducted with EPU software (Thermo Fisher Scientific) with a final pixel size of 1.065Å/pix over 40 frames with a total dose of 50 e⁻/Å². The targeted defocus was set between -1.6 and -2.8 µm with 0.2 µm increment.

Movie-alignments with dose-weighting were performed in MotionCor2 software[43] with a gain reference estimated a posteriori[44] followed by standard in-house pipeline of manual inspection and selection of 38,845 micrographs[45] for further processing in cryoSPARC package v4.0.3[46]. Due to the low concentration of particles and the presence of other contaminants in the sample, automatic particle picking appeared challenging, and we developed CIS-specific approach for efficient picking of the cap and baseplates. In this approach, CIS^{Sc} particles were manually selected as filaments (start-to-end) on 920 micrographs and used for training a crYOLO[47] picking model (EV2). The large size of the training set appeared essential for successful picking; we could not get rid of large number of false-positive picks, but there were not so many false-negative picks. Then, CIS^{Sc} particles were picked in crYOLO as filaments,

followed by the extraction of the ends of the "filaments" using star_modif.py script[45]. Particle

coordinates were imported in cryoSPARC, where 211,041 particles were extracted at a large

box size of 756 pixels binned to 64 pixels for initial 2D-classification, which allowed us to

separate initial particle sets corresponding to the CIS^{Sc} cap (36,569 particles) and CIS^{Sc}

baseplate (43,087 particles) (EV2).

Two subsets of this dataset were processed in cryoSPARC separately. For the cap subset, a

standard cryoSPARC single-particle processing approach was applied, including gradual

iterative 2D-classification rounds, followed by selection of good classes and particle re-

extraction at finer sampling. The main challenge was the proper centering of the particles

selected on the end of elongated filament-like objects with repeating patterns (sheath-tube

module of CIS), hampering the alignments. Ab-initio model generation with applied C6

symmetry was hampered by poor centering of 2D-classes due to elongated nature of the

particles (EV2). Homo-refinement with this reference was performed followed by particle re-

extraction and one more round of ab-initio model generation with applied C6 symmetry (EV2).

The new 3D-reconstruction appeared reliable and was used for further iterative 3D-refinements

(Homo-refinement and NU-refinements) accompanied by 2D-classification and CTF-

refinement rounds to exclude particles with wrong Euler angle assignments. The final 3.4 Å

resolution map with imposed C6 symmetry resulted from 19,218 particles.

The baseplate subset was processed using the same approach, resulting in a 3D-reconstruction

with imposed C6 symmetry with a resolution of 3.5Å from 22,980 particles (EV2). The tip of

the baseplate has a C3 symmetry, therefore to solve that part of the map correctly, we exported

that particle stack to Relion4[48] and took advantage of 2D-classification in Relion to select

18,124 particles, corresponding to the best 2D classes averages. 3D-refinement with relaxed

symmetry resulted in a 3.8 Å 3D-resolution reconstruction with imposed C3 symmetry.

Cryo-electron tomography

Intact *Streptomyces* cells were imaged by cryoET as described previously[49]. Images were

recorded using a Titan Krios 300 kV microscope (Thermo Fisher Scientific) equipped with a

Quantum LS imaging filter operated at a 20 eV slit width and with K3 Summit direct electron

detectors (Gatan). Tilt series were collected using a bidirectional tilt-scheme from -60 to +60°

in 2° increments. Total dose was 130-150 e$^-$/Å^2 and defocus was kept at -8 μm. Tilt series were

acquired using SerialEM[50], drift-corrected using alignframes, and reconstructed using IMOD

program suite[51]. To enhance contrast, tomograms were deconvolved with a Wiener-like filter

'tom_deconv'[52].

Structural modeling

The map qualities of different modules of CIS[Sc] (cap and baseplate) allowed for *de novo*

structural modelling (EV2-3). The atomic models of SCO4243-4248, SCO4252-4253 and

SCO4260 were built *de novo* using COOT[53]. The resulting models were iteratively refined

using RosettaCM[54] and real-space refinement implemented in PHENIX[55]. In cases where

protein domains could only be partially modeled, side chains were not assigned. Final model

validation was done using MolProbity[55] and correlation between models and the corresponding

maps were estimated using mtriage[55].

Fluorescence microscopy

Fluorescence-based cell viability assays were performed as described previously[18]. Briefly, to

produce cytoplasmic sfGFP in *Streptomyces*, the coding sequence for sfGFP was introduced

downstream of the constitutive promoter *ermE** on an integrating plasmid. The plasmid was

introduced by conjugation to *S. coelicolor* strains (SS430, SS575, SS576. An equal level of

cytoplasmic sfGFP in the different strains was confirmed by Western blotting (EV9).

Streptomyces strains were grown in 30 ml of TSB liquid culture at 30 ˚C with shaking at 250

rpm for 48 h. Where appropriate, nisin was added to a final concentration of 1 µg/ml 90 min

prior to imaging. One ml aliquots of each culture were centrifuged for 5 min at 15,000 x *g*,

washed twice with 1xPBS, resuspended in 1 ml of 1xPBS containing 5 µg/ml of the red-

fluorescent membrane dye FM5-95 (Invitrogen) and then incubated in the dark at room

temperature for 10 min followed by two wash steps with 1x PBS. Washed cell pellets were

resuspended in a total volume of 50 µl PBS and 10 µl of each sample was spotted onto 1 %

agar pads and subsequently imaged using the Thunder imager 3D cell culture microscope

(Leica). First, tile scan images were acquired using the Las X Navigator plug-in software (Leica

Application Suite X, Version 3.7.4.23463), and 100 regions of interest were picked manually.

Then Z-stack images were acquired using a HC PL APO 100x objective with the following

excitation wavelengths: GFP (475 nm) and TRX (555 nm). Images were processed using LasX

software to apply thunder processing and maximum projection and FIJI to create segmentation

and quantify the live (sfGFP)/total cells (FM5-95) area ratio[56]. Statistical analyses were

performed on data from biological triplicates (n=100 images for each experiment) using a one-
way ANOVA and Tukey's post-test in GraphPad Prism 9 (Version 9.3.1).

For imaging the subcellular localization of the CIS adaptor protein in *Streptomyces*, cells (strain
JS69 or *S. coelicolor* M145) were grown for 48 h in TSB/YEME liquid medium. Two µl of
each strain were spotted on 1 % agarose pads and subsequently imaged. Images were acquired
using a Zeiss Axio Observer Z.1 inverted epifluorescence microscope fitted with a sCMOS
camera (Hamamatsu Orca FLASH 4), a Zeiss Colibri 7 LED light source, and a Hamamatsu
Orca Flash 4.0v3 sCMOS camera. mCherry fluorescence was detected using an
excitation/emission bandwidth of 577–603 nm/614–659 nm. Images were collected using Zen
Blue (Zeiss) and analyzed using Fiji[56].

Cover glass impression of *Streptomyces* spore chains

Spore titers of relevant strains were determined by dilution plating[18]. 10^7 colony forming units
(CFU) of *S. coelicolor* strains (WT, SS539, SS557) were spread onto R2YE agar plates and
grown at 30 °C. Sterile glass coverslips were gently applied to the top surface of each bacterial
lawn after 48 h, 72 h and 96 h post inoculation. Coverslips were then mounted onto glass
microscope slides and imaged using a 40x objective on a Leica Thunder Imager 3D Cell
Culture. Images were processed using FIJI[56].

Actinorhodin production assay

S. coelicolor strains (WT, SS539, SS557) were inoculated into 30 ml R2YE liquid media at a
final concentration of 1.5 x 10^6 CFU/ml. Cultures were grown in baffled flasks at 30 °C
overnight. Cultures were standardized to an OD_{450} of 0.5 and inoculated in 30 ml of fresh R2YE
liquid medium. For visual comparison of pigment production, images of the growing culture
were taken between t = 0 and t = 72 h (as indicated in EV7). For quantification of total
actinorhodin production, 480 µl of samples were collected at the same time points where
images were taken. 120 µl of 5M KOH was added, samples were vortexed and centrifuged at
5,000 x g for 5 min. The weight of each tube was recorded. A Synergy 2 plate reader (Biotek)
was used then to measure the absorbance of the supernatant at 640 nm. The absorbance was
normalized by the weight of the wet pellet.

Structure prediction

CisA structure was predicted using AlphaFold2[57] and further processed using UCSF Chimera[58]

or ChimeraX-1.7.1[59].

A pairwise *in silico* screen for possible interactions between CisA and components encoded by

the *S. coelicolor cis* gene cluster (*SCO4242-SCO4260*) was conducted using the AlphaFold2

Multimer-based LazyAF pipeline[57,60,61]. The confidence metric (ipTM) for the top model from

each pairwise interaction was tabulated and visualised using Prism (Version 10.2.2) with ipTM

scores >0.7, indicating a possible protein-protein interaction[62].

Data availability

Representative reconstructed tomograms (EMD-XXXXX, EMD-XXXXX, EMD-XXXXX, EMD-XXXXX, EMD-XXXXX, EMD-XXXXX, EMD-XXXXX…) and SPA cryoEM maps (EMD-XXXXX, EMD-XXXXX and EMD-XXXXX) have been deposited in the Electron Microscopy Data Bank. Atomic models (PDB: XXXX, PDB: XXXX, PDB: XXXX) have been deposited in the Protein Data Bank.

Acknowledgements

We thank ScopeM for instrument access at ETH Zürich. We thank the Functional Genomics Center Zürich for mass spectrometry support. Pilhofer Lab members are acknowledged for discussions. M.P. was supported by the Swiss National Science Foundation (31003A_179255/310030_212592), the European Research Council (679209), and the NOMIS foundation. Work in the lab of S.S. was supported by a Royal Society University Research Fellowship (URF\R1\180075) and BBSRC grant BB/T015349/1 to S.S. and by the BBSRC Institute Strategic Program grant BB/X01097X/1 to the John Innes Centre.

Author contributions statement

B.C., S.S. and M.P. conceived the project. B.C. optimized the sample preparation for SPA; B.C and P.A collected and processed the cryoEM data, reconstructed the cryoEM map; B.C, P.E.H and J.X built and refined the structural models; B.C. and P.E.H conducted and processed cryoET; B.C. conducted automated fluorescent light microscopy, determined sporulation efficiency and actinorhodin production; J.W.S and S.S. generated plasmids and *Streptomyces* strains; J.W.S. designed constructs and performed membrane topology assay. J.W.S. prepared samples and performed cellular fractionation experiment. J.W.S. imaged strains expressing fluorescently tagged CisA. B.C., J.W.S, S.S. and M.P. wrote the manuscript.

521   **Competing interests statement**

522   The authors declare no competing interests.

Figures legends

Figure 1: Contraction of CIS^{Sc} *in situ* is dependent on CisA.

a. Schematic of the mode of action of CIS^{Sc} in *Streptomyces coelicolor*[18,19]. CIS^{Sc} are assembled as free-floating particles in the cytoplasm of vegetative hyphae. In response to cellular stress and/or an unknown cellular signal, CIS^{Sc} particles contract, which results in regulated cell death (mediated by released effectors) and impacts cellular development.

b/c. Negative-stain electron micrographs of purified CIS^{Sc} particles from *S. coelicolor* wildtype (WT) (b) and the Δ*cisA* mutant (c), show that all CIS^{Sc} particles are contracted upon purification. These experiments were performed three independent times. Bars, 100 nm.

d-f. Shown are representative images of cryo-electron tomogram slices (thickness 11 nm) of vegetative hyphae (top: intact cells; bottom: ghost cells) of *S. coelicolor* WT (d)*,* Δ*cisA* mutant (e), and the complemented Δ*cisA/cisA$^+$* mutant (f). CIS^{Sc} particles remain almost exclusively in an extended state (white arrowheads) in the Δ*cisA* mutant, whereas in ghost cells derived from the WT and the complemented mutant CIS^{Sc} particles (black arrowheads) are mostly contracted. See also EV1. PG, peptidoglycan; CM, cytoplasmic membrane; Bars, 50 nm.

Figure 2: CryoEM structure of the extended CIS^{Sc} assembly reveals its composition.

a. Schematic illustrating the CIS^{Sc} gene cluster (adapted from [18]). Asterisks indicate gene
products that were detected by mass spectrometry analyzing crude preparations of CIS^{Sc}
particles (red). See also Table EV1.

b/c. Single-particle cryoEM structure of an extended CIS^{Sc} particle obtained from purified
particles from a non-contractile CIS^{Sc} mutant. Shown is the composite atomic model in surface
(left) and ribbon (right) rendering (b). Subunits are color-coded according to (a) and their copy
numbers in the assembly are indicated in (c).

d. Perpendicular views of the ribbon representation of the CIS^{Sc} model of the cap and baseplate
modules.

e. Top view (left) and side view (right) of ribbon diagrams showing the cap module of CIS^{Sc},
revealing that the cap is composed of a Cis16 hexamer.

f. Ribbon diagrams showing a single baseplate wedge subunit, which is composed of two
copies of Cis11 and one copy of Cis12.

Figure 3: CisA is a single-pass membrane protein.

a. Schematic and Alphafold2 model of CisA. CisA is predicted to be comprised of a largely unstructured N-terminal domain (grey), a predicted transmembrane domain (TM, purple) and a C-terminal immunoglobulin-like domain (green)[18,19].

b/c. Representative micrographs (left panels: phase contrast, Ph3; right panel: mCherry) of strains of *S. coelicolor* vegetative hyphae either constitutively expressing CisA-mCherry (*ΔcisA/cisA-mcherry*) *in trans* (b) or WT cells carrying an empty vector (e.v.) to control for background fluorescence (c). Box shows a magnified region of hyphae with CisA-mCherry accumulation in the cellular periphery. Bars, 5 μm.

d. Western blot of the cellular fractionation of samples from the *S. coelicolor* WT or the *ΔcisA* mutant constitutively expressing a CisA-3xFLAG fusion *in trans*. Lysate and soluble and membrane fractions were probed for the presence of CisA-3xFLAG with an α-FLAG antibody

(top). Fractionation efficiency was assessed using an α-WhiA antibody to detect the soluble
transcription factor WhiA (bottom). CisA-3xFLAG was detected in the lysate and the
membrane fraction. The experiment was performed in biological duplicates and shown is a
representative image.

e/f. Experimental determination of the CisA membrane topology using the dual *phoA-lacZα*
reporter in *E. coli*. The schematic diagram (e) depicts the four CisA constructs used in the assay
and indicates the relevant amino acid (aa) deleted in the three CisA mutant derivates. The
analysis of colony coloration (f); pink coloration for cytosolic proteins and blue coloration for
periplasmic proteins indicates that the CisA N-terminus is localized in the cytoplasm and the
CisA C-terminus localized to the periplasm and this topology is dependent on the
transmembrane domain (TM). *E. coli* strains carrying an empty reporter plasmid or reporter
plasmids with the *Streptomyces* genes for SepH (cytoplasmic) and RsbN (periplasmic) were
used as controls. Note that expression of full-length CisA in *E. coli* is toxic, leading to reduced
growth and lighter blue coloration.

Figure 4: CisA is required for regulated cell death and cellular function of CIS^{Sc}.

a-d. Light microscopy-based viability assay showing that CIS^{Sc} do not mediate cell death in CisA-deficient hyphae in response to exogenous stress. Shown are representative micrographs of the WT/sfGFP, the Δ*cisA*/sfGFP and the complemented Δ*cisA*/*cisA*$^+$/sfGFP mutant expressing cytosolic sfGFP (indicator for live cells). Strains were grown for 48 h and then either incubated without (a) or with (b) the membrane-disrupting antibiotic nisin (1 µg/ml nisin for 90 min). Samples were subsequently stained with the fluorescent membrane dye FM5-95 to visualize the cytoplasmic membrane. Bars, 10 µm. Z-stacks were taken of the samples and the relative ratio of fluorescence from sfGFP to FM5-95 was calculated in c/d. WT and Δ*cisA*/*cisA*$^+$ complemented cells showed a significantly higher rate of cell death in response to nisin treatment compared to Δ*cisA* cells. There is no significant difference in the viability of the tested strains in the absence of nisin. Experiments were performed in three biological replicates (green, grey and pink data points). Black lines indicate the mean ratio derived from biological triplicate experiments (n=100 images per replicate). ns (not significant) and **** (p < 0.0001) were determined using a one-way ANOVA and Tukey's post-test.

e. Visualization of spore production in WT, Δ*cisA* mutant and Δ*cisA*/*cisA*$^+$ mutant reveal accelerated cellular development of the Δ*cisA* mutant. Shown are representative brightfield images of surface impressions of plate-grown colonies of each of the strains taken at the indicated time points. Only sporulating hyphae will attach to the hydrophobic cover glass surface. Insets show magnified regions of the colony surface containing spores and spore chains. Bars, 50 µm.

f. Quantification of sporulation shown in (e) via counts of colony-forming units (c.f.u.). The graph shows a 6-fold higher c.f.u. count at 72 h in the Δ*cisA* mutant. Strains were grown on R2YE agar and spores were harvested after 48 h, 72 h and 96 h of incubation. Data show mean ± s.d. obtained from biological triplicate experiments.

Figure 5: Cytoplasmic CIS^{Sc} require the membrane protein CisA for contraction and
function.

Proposed model illustrating a possible role of CisA. In response to exogenous stress or an
unknown cellular signal, membrane-bound CisA either directly or indirectly mediates the
association of free-floating CIS^{Sc} particles with the cytoplasmic membrane, leading to CIS^{Sc}
firing and regulated cell death, which impacts the *Streptomyces* developmental life cycle.

Extended Data Figure 1: Sheath contraction *in situ* is linked to the presence of CisA.

a-c. Shown are additional cryo-tomographic slices (thickness 11 nm) of *S. coelicolor* Δ*cisA* vegetative hyphae, which were observed as 'Intact hyphae' (a), 'Partially lysed hyphae' (b) and 'Ghost cells' (c). Note that all visible CIS^{Sc} particles (white arrowheads) are in the extended conformation. PG, peptidoglycan; CM, cytoplasmic membrane/membranes. Bars, 50 nm.

d. Shown is a quantification of CIS^{Sc} assemblies per tomogram and their conformations as observed in different classes of Δ*cisA* hyphae. Almost all CIS^{Sc} particles were seen in the extended conformation, indicating that sheath contraction *in situ* correlates with the presence of CisA. Data shows mean values and standard deviations obtained from biological triplicate experiments, with n=30 tomograms for each class of hyphae.

Total 38,845 micrographs collected
crYOLO training: only 920 micrographs used
and manually picked as filaments (start-to-end)

crYOLO picking as filaments
Extraction of the ends of the "filaments" using star_modif.py script
211,041 particles imported in cryoSPARC
Then extracted at a large box size of 756 pixels
Binned to 64 pixels for initial 2D-classification

CIS^{Sc} cap - 36,569 particles

CIS^{Sc} baseplate - 43,087 particles

The processing approach was similar to that of the cap

Standard cryoSPARC single-particle processing:
Gradual iterative 2D-classification rounds
Selection of good classes
Particle re-extraction at finer sampling
Proper centering of the particles selected
Ab-initio model generation (C6 symmetry)

Final 3.5 Å resolution map
(C6 symmetry)
from 22,980 particles

Homo-refinement with initial reference
Particle re-extraction
One more round of *ab-initio* (C6 symmetry)

The tip of the baseplate has a C3 symmetry:
Export particle stack to Relion4
2D-classification in Relion to select 18,124 particles
3D-refinement with relaxed symmetry at 3.8 Å (C3 symmetry)

New 3D-reconstruction reliable
Iterative 3D-refinements
2D-classification
CTF-refinement rounds
Final 3.4 Å resolution map (C6 symmetry)
from 19,218 particles

Iterative 3D-refinements

Extended Data Figure 2: Workflow for the cryoEM structural determination of the extended CIS^{Sc}.

Flowchart for cryoEM reconstruction of the extended CIS^{Sc} particle. See METHODS and Table EV1 for details.

631
632

Extended Data Figure 3: Map qualities of different CIS^{Sc} modules.

a-f. Local resolution maps of the CIS^{Sc} baseplate with C6 symmetry (a), CIS^{Sc} baseplate with
C3 symmetry (c), and the CIS^{Sc} cap with C6 symmetry (e). (d-f) shows regions of the cryoEM
density map (mesh) that were superimposed with the atomic models (ribbons and sticks),
demonstrating the agreement between the observed and modeled amino acid side chains for
one beta-sheet (left) and one alpha helix (right). Shown are examples of the baseplate with C6
symmetry (b), the baseplate with C3 symmetry (d), and the cap with C6 symmetry (f).

Extended Data Figure 4: Structures of the cap modules of other CISs.

a-d. Top views (left) and side views (right) of ribbon representations of the structures of CIS

cap modules of Pyocin R2 (a, PDB 6U5B), PVC (b, PDB 6J0N), AFP (**c**, PDB 6RAP) and

AlgoCIS (d, PDB 7ADZ).

Extended Data Figure 5: Structures of the baseplate modules of other CISs.

a-d. Ribbon representations of the full structures (left) and the individual wedge subunit (right)

of the CIS baseplate components (Cis11/12) of Pyocin R2 (a, PDB 6U5F), PVC (b, PDB 6J0F),

AFP (c, PDB 6RAO) and AlgoCIS (d, PDB 7AEB).

Extended Data Figure 6: Control experiment showing that CIS^{Sc} particles are comparable in the strains used in the viability assay.

662 Representative negative-stain electron micrographs of purified CIS^{Sc} particles from the
663 WT/sfGFP, the Δ*cisA*/sfGFP mutant and the Δ*cisA*/Δ*cisA*$^+$/sfGFP complemented mutant
664 exposed to no stress (upper row) or 1 μg/ml nisin (lower row). No differences were observed
665 between the overall structure of CIS^{Sc} particles. These experiments were performed three
666 independent times. Bars, 100 nm.

Extended Data Figure 7: CisA impacts secondary metabolite production.

a. Comparison of cultures of WT, Δ*cisA* mutant and Δ*c*isA/Δ*cisA*$^+$ mutant in R2YE liquid medium. The coloration of the culture is indicative of actinorhodin (purple) and undecylprodigiosin (red) production[31]. Images of each culture flask were taken at the indicated time points.

b. Quantification of total actinorhodin production by the strains shown in (a), revealing reduced production in the Δ*cisA* mutant. Samples were taken at the indicated time points. Actinorhodin was extracted and quantified by measuring the optical density OD_{640} of the culture supernatant[63] and normalized to the wet pellet weight. Analysis was performed in biological triplicate experiments. Shown are the mean values and standard error. ns (not significant) and **** ($p < 0.0001$) were calculated using one-way ANOVA and Tukey's post-test.

Extended Data Figure 8: *In silico* **protein-protein interaction analyses of CisA and proteins of the CIS^{Sc} gene cluster.**

a. An AlphaFold2-Multimer based protein-protein interaction screen between CisA and the 19 gene products of the CIS^{Sc} gene cluster suggests an interaction between CisA and the baseplate component Cis11. The likelihood of a protein-protein interaction was ranked based on the interface pTM (ipTM) confidence score, with an ipTM greater than 0.7 indicating a potential protein-protein interaction.

b. Structure of the CIS^{Sc} baseplate illustrating the peripheral and surface-exposed position of Cis11.

Extended Data Figure 9: Western blot confirming similar levels of cytosolic sfGFP in strains used for viability assays.

Control experiments to verify similar protein levels of sfGFP which was expressed either from the φC31 or the φBT1 phage attachment site in the *S. coelicolor* genome. Expression of *sfgfp* was driven from the constitutive *ermE** promoter (P_{ermE*}). Equal amounts of total protein were loaded and sfGFP levels were detected with an anti-GFP antibody. As a control, the same cell lysates were also probed with an anti-WhiA antibody. Shown are representative results of duplicate experiments.

References

1. Lin, L. The expanding universe of contractile injection systems in bacteria. *Curr. Opin. Microbiol.* **79**, 102465 (2024).

2. Brackmann, M., Nazarov, S., Wang, J. & Basler, M. Using Force to Punch Holes: Mechanics of Contractile Nanomachines. *Trends Cell Biol.* **27**, 623–632 (2017).

3. Taylor, N. M. I., van Raaij, M. J. & Leiman, P. G. Contractile injection systems of bacteriophages and related systems. *Mol. Microbiol.* **108**, 6–15 (2018).

4. Taylor, N. M. I. *et al.* Structure of the T4 baseplate and its function in triggering sheath contraction. *Nature* **533**, 346–352 (2016).

5. Böck, D. *et al.* In situ architecture, function, and evolution of a contractile injection system. *Science* **357**, 713–717 (2017).

6. Allsopp, L. P. & Bernal, P. Killing in the name of: T6SS structure and effector diversity. *Microbiology* **169**, 001367 (2023).

7. Coulthurst, S. The Type VI secretion system: a versatile bacterial weapon. *Microbiol. Read. Engl.* **165**, 503–515 (2019).

8. Chen, L. *et al.* Genome-wide Identification and Characterization of a Superfamily of Bacterial Extracellular Contractile Injection Systems. *Cell Rep.* **29**, 511-521.e2 (2019).

9. Geller, A. M. *et al.* The extracellular contractile injection system is enriched in environmental microbes and associates with numerous toxins. *Nat. Commun.* **12**, 3743 (2021).

10. Basler, M., Pilhofer, M., Henderson, G. P., Jensen, G. J. & Mekalanos, J. J. Type VI secretion requires a dynamic contractile phage tail-like structure. *Nature* **483**, 182–186 (2012).

11. Durand, E. *et al.* Biogenesis and structure of a type VI secretion membrane core complex. *Nature* **523**, 555–560 (2015).

12. Zachs, T. *et al.* Archaeal type six secretion system mediates contact-dependent antagonism.

2024.04.11.588991 Preprint at https://doi.org/10.1101/2024.04.11.588991 (2024).

13. Xu, J. *et al.* Identification and structure of an extracellular contractile injection system from

the marine bacterium Algoriphagus machipongonensis. *Nat. Microbiol.* **7**, 397–410 (2022).

14. Shikuma, N. J. *et al.* Marine tubeworm metamorphosis induced by arrays of bacterial phage

tail-like structures. *Science* **343**, 529–533 (2014).

15. Hurst, M. R. H., Beard, S. S., Jackson, T. A. & Jones, S. M. Isolation and characterization

of the Serratia entomophila antifeeding prophage. *FEMS Microbiol. Lett.* **270**, 42–48 (2007).

16. Jiang, F. *et al.* Cryo-EM Structure and Assembly of an Extracellular Contractile Injection

System. *Cell* **177**, 370-383.e15 (2019).

17. Weiss, G. L. *et al.* Structure of a thylakoid-anchored contractile injection system in

multicellular cyanobacteria. *Nat. Microbiol.* **7**, 386–396 (2022).

18. Casu, B., Sallmen, J. W., Schlimpert, S. & Pilhofer, M. Cytoplasmic contractile injection

systems mediate cell death in Streptomyces. *Nat. Microbiol.* **8**, 711–726 (2023).

19. Vladimirov, M., Zhang, R. X., Mak, S., Nodwell, J. R. & Davidson, A. R. A contractile

injection system is required for developmentally regulated cell death in Streptomyces

coelicolor. *Nat. Commun.* **14**, 1469 (2023).

20. Nagakubo, T. *et al.* Phage tail-like nanostructures affect microbial interactions between

Streptomyces and fungi. *Sci. Rep.* **11**, 20116 (2021).

21. Nagakubo, T. *et al.* Intracellular Phage Tail-Like Nanostructures Affect Susceptibility of

Streptomyces lividans to Osmotic Stress. *mSphere* **8**, e00114-23 (2023).

22. Nagakubo, T., Nishiyama, T., Yamamoto, T., Nomura, N. & Toyofuku, M. Contractile

injection systems facilitate sporogenic differentiation of Streptomyces davawensis through

the action of a phage tapemeasure protein-related effector. *Nat. Commun.* **15**, 4442 (2024).

23. Schlimpert, S. & Elliot, M. A. The Best of Both Worlds—Streptomyces coelicolor and

Streptomyces venezuelae as Model Species for Studying Antibiotic Production and

Bacterial Multicellular Development. *J. Bacteriol.* **205**, e00153-23 (2023).

24. Flärdh, K. & Buttner, M. J. Streptomyces morphogenetics: dissecting differentiation in a

filamentous bacterium. *Nat. Rev. Microbiol.* **7**, 36–49 (2009).

25. Desfosses, A. *et al.* Atomic structures of an entire contractile injection system in both the

extended and contracted states. *Nat. Microbiol.* **4**, 1885–1894 (2019).

26. Ge, P. *et al.* Action of a minimal contractile bactericidal nanomachine. *Nature* **580**, 658–

662 (2020).

27. Alexeyev, M. F. & Winkler, H. H. Membrane topology of the Rickettsia prowazekii

ATP/ADP translocase revealed by novel dual pho-lac reporters 1 1Edited by G. von Heijne.

J. Mol. Biol. **285**, 1503–1513 (1999).

28. Ramos-León, F. *et al.* A conserved cell division protein directly regulates FtsZ dynamics

in filamentous and unicellular actinobacteria. *eLife* **10**, e63387 (2021).

29. Bibb, M. J., Domonkos, A., Chandra, G. & Buttner, M. J. Expression of the chaplin and

rodlin hydrophobic sheath proteins in Streptomyces venezuelae is controlled by σ(BldN)

and a cognate anti-sigma factor, RsbN. *Mol. Microbiol.* **84**, 1033–1049 (2012).

30. Malpartida, F. & Hopwood, D. A. Molecular cloning of the whole biosynthetic pathway of

a Streptomyces antibiotic and its expression in a heterologous host. *Nature* **309**, 462–464

(1984).

31. Tsao, S. W., Rudd, B. A., He, X. G., Chang, C. J. & Floss, H. G. Identification of a red

pigment from Streptomyces coelicolor A3(2) as a mixture of prodigiosin derivatives. *J.*

Antibiot. (Tokyo) **38**, 128–131 (1985).

32. Rapisarda, C. *et al.* In situ and high-resolution cryo-EM structure of a bacterial type VI

secretion system membrane complex. *EMBO J.* **38**, e100886 (2019).

33. Lien, Y.-W. *et al.* Mechanism of bacterial predation via ixotrophy. 2024.01.29.577165

Preprint at https://doi.org/10.1101/2024.01.29.577165 (2024).

34. Bongiovanni, T. R. *et al.* Assembly of a unique membrane complex in type VI secretion

systems of Bacteroidota. *Nat. Commun.* **15**, 429 (2024).

35. Ouyang, R., Ongenae, V., Muok, A., Claessen, D. & Briegel, A. Phage fibers and spikes:

a nanoscale Swiss army knife for host infection. *Curr. Opin. Microbiol.* **77**, 102429 (2024).

36. Weiss, G. L. *et al.* Structure of a thylakoid-anchored contractile injection system in

multicellular cyanobacteria. *Nat. Microbiol.* **7**, 386–396 (2022).

37. T. Kieser M. J. Bibb M. J. Buttner K. F. Chater and D. A. Hopwood. *Practical Streptomyces*

Genetics. (John Innes Foundation, 2000).

38. Gust, B., Challis, G. L., Fowler, K., Kieser, T. & Chater, K. F. PCR-targeted Streptomyces

gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene

soil odor geosmin. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 1541–1546 (2003).

39. Gust, B. *et al.* Lambda red-mediated genetic manipulation of antibiotic-producing

Streptomyces. *Adv. Appl. Microbiol.* **54**, 107–128 (2004).

40. Ohi, M., Li, Y., Cheng, Y. & Walz, T. Negative Staining and Image Classification -

Powerful Tools in Modern Electron Microscopy. *Biol. Proced. Online* **6**, 23–34 (2004).

41. Bush, M. J., Bibb, M. J., Chandra, G., Findlay, K. C. & Buttner, M. J. Genes Required for

Aerial Growth, Cell Division, and Chromosome Segregation Are Targets of WhiA before

Sporulation in Streptomyces venezuelae. *mBio* **4**, e00684-13 (2013).

42. Gallego, J. J., Severi, E., Chandra, G. & Palmer, T. Identification of novel tail--anchored

membrane proteins integrated by the bacterial twin--arginine translocase.

43. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for

improved cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

44. Afanasyev, P. *et al.* A posteriori correction of camera characteristics from large image data

sets. *Sci. Rep.* **5**, 10317 (2015).

45. P. Afanasyev. CryoEM tools. https://github.com/afanasyevp/cryoem_tools (2023).

46. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for

rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

47. Wagner, T. *et al.* SPHIRE-crYOLO is a fast and accurate fully automated particle picker

for cryo-EM. *Commun. Biol.* **2**, 1–13 (2019).

48. Scheres, S. H. W. RELION: implementation of a Bayesian approach to cryo-EM structure

determination. *J. Struct. Biol.* **180**, 519–530 (2012).

49. Weiss, G. L., Medeiros, J. M. & Pilhofer, M. In Situ Imaging of Bacterial Secretion Systems

by Electron Cryotomography. *Methods Mol. Biol. Clifton NJ* **1615**, 353–375 (2017).

50. Mastronarde, D. N. Automated electron microscope tomography using robust prediction of

specimen movements. *J. Struct. Biol.* **152**, 36–51 (2005).

51. Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-

dimensional image data using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996).

52. Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with

Warp. *Nat. Methods* **16**, 1146–1152 (2019).

53. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.

Acta Crystallogr. D Biol. Crystallogr. **66**, 486–501 (2010).

54. Song, Y. *et al.* High-resolution comparative modeling with RosettaCM. *Struct. Lond. Engl.*

1993 **21**, 1735–1742 (2013).

55. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular

structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–221 (2010).

56. Schindelin, J. *et al.* Fiji - an Open Source platform for biological image analysis. *Nat.*

Methods **9**, 10.1038/nmeth.2019 (2012).

57. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

58. Ef, P. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis.

J. Comput. Chem. **25**, (2004).

59. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and

analysis. *Protein Sci. Publ. Protein Soc.* **27**, 14–25 (2018).

60. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods* **19**,

679–682 (2022).

61. McLean, T. C. LazyAF, a pipeline for accessible medium-scale *in silico* prediction of

protein-protein interactions. Preprint at https://doi.org/10.1101/2024.01.29.577767 (2024).

62. O'Reilly, F. J. *et al.* Protein complexes in cells by AI -assisted structural proteomics. *Mol.*

Syst. Biol. **19**, e11544 (2023).

63. Zhang, Y., Wang, L., Zhang, S., Yang, H. & Tan, H. hmgA, transcriptionally activated by

HpdA, influences the biosynthesis of actinorhodin in Streptomyces coelicolor. *FEMS*

Microbiol. Lett. **280**, 219–225 (2008).

Table EV1. Cryo-EM data statistical analysis

	Cap	Baseplate_C6 (overall)	Baseplate_C3
Data collection and processing			
Nominal magnification		81,000	
Voltage (kV)		300	
Electron exposure (e⁻/Å)		~60 (K3 camera)	
Defocus range (μm)		1.0 - 3.0	
Pixel size (Å/pixel)		1.065	
Symmetry imposed	C6	C6	C3
Initial particles (No.)	36,569	43,087	36,569
Final particles (No.)	19,218	22,920	18,124
Map resolution (Å)	3.4	3.5	3.8
FSC threshold		0.143	
Refinement			
Model composition			
Atoms	33006 (Hydrogens: 243)	141138 (Hydrogens: 0)	14343 (Hydrogens: 0)
Protein residues	4266	18408	1923
Chains	18	60	3
R.M.S deviations			
Bond length (Å)	0.003	0.007	0.005
Bond angles (˚)	0.558	0.839	0.768
Validation			
MolProbity score	1.51	2.22	1.91
Clashscore	5.77	14.01	7.86
Rotamer outlier (%)	0.03	0.04	0
Ramachandran plot			
Favored (%)	96.81	89.51	92.18
Allowed (%)	3.19	10.46	7.82
Outlier (%)	0	0.44	0
Masked CC	0.77	0.79	0.77

Table EV2. Proteins detected by mass spectrometry in samples of purified CIS^{Sc} after crude sheath preparation from a *S. coelicolor* non-contractile CIS^{Sc} mutant without exogeneous stress and following nisin stress.

Protein ID	CIS ID	*S. coelicolor* CIS-N5	
		Without exogeneous stress	Upon nisin stress
SCO4242	CisA	-	5% coverage / 2 total unique peptide
SCO4243	Cis12	6% coverage / 1 total unique peptide	5% coverage / 1 total unique peptide
SCO4244	Cis11	13% coverage / 6 total unique peptide	4% coverage / 2 total unique peptide
SCO4245	Cis9	21% coverage / 2 total unique peptide	20% coverage / 2 total unique peptide
SCO4246	Cis8	18% coverage / 10 total unique peptide	13% coverage / 7 total unique peptide
SCO4247	Cis7	5% coverage / 1 total unique peptide	4% coverage / 1 total unique peptide
SCO4248	Cis1b	22% coverage / 2 total unique peptide	20% coverage / 2 total unique peptide
SCO4249	-	5% coverage / 1 total unique peptide	4% coverage / 1 total unique peptide
SCO4252	Cis1a	26% coverage / 4 total unique peptide	19% coverage / 3 total unique peptide
SCO4253	Cis2	23% coverage / 9 total unique peptide	30% coverage / 14 total unique peptide
SCO4254	-	10% coverage / 6 total unique peptide	10% coverage / 6 total unique peptide
SCO4256	-	17% coverage / 7 total unique peptide	22% coverage / 9 total unique peptide
SCO4257	-	-	-
SCO4258	-	-	-
SCO4259	Cis15	-	-
SCO4260	Cis16	20% coverage / 4 total unique peptide	17% coverage / 3 total unique peptide

Table EV3: Strains and plasmids used in this study

Strain	Description	Construction	Source
Escherichia coli strains			
TOP10	*F⁻ mcrA Δ(mrr-hsdRMS-mcrBC) Φ80lacZΔM15 ΔlacX74 recA1 araD139 Δ(ara leu) 7697 galU galK rpsL (StrR) endA1 nupG*	Cloning	Lab strain
ET12567/pUZ8002	*F⁻ dam13::Tn9 dcm6 hsdM hsdR recF143:: Tn10 galK2 galT22 ara-14 lacY1 xyl-5 leuB6 thi-1 tonA31 rpsL hisG4 tsx-78 mtl-1 glnV44*	ET12567 with helper plasmid pUZ8002	[1]
BW25113/pIJ790	*Δ(araD-araB)567 ΔlacZ4787(::rrnB-4) lacIp-4000(lacIQ), λ−rpoS369(Am) rph-1 Δ(rhaD-rhaB)568 hsdR514*	BW25113 containing λ RED recombination plasmid pIJ790	[2]
NEB5-alpha	*fhuA2Δ(argF-lacZ)U169 phoA glnV44 Φ80 Δ(lacZ)M15 gyrA96 recA1 relA1 endA1 thi-1 hsdR17*	Cloning	NEB
Rosetta (DE3)	*F⁻ ompT hsdSB(rB⁻ mB-) gal dcm (DE3) pRARE (CamR)*	Host strain for protein overexpression	Merck
TG1	*supE thi-1 Δ(lac-proAB) Δ(mcrB-hsdSM)5 (rK-mK) / F´ traD36 proAB lacIqZΔM15*	Topology Experiment	Lab Strain
Streptomyces coelicolor strains			
S. coelicolor M145	Wild type (SCP1-, SCP2-)		[3]
SS387	*ΔSCO4253-51::apr, aprR*		[4]

SS393	∆*SCO4253-51::apr* *attB* φ*BT1* *(SCO4253-N5-SCO4252-51),* *hyg^R*		[4]
SS430	WT φ*BT1* *(PermE*-sfgfp)*		[4]
SS539	∆*SCO4242::apr,* *apr^R*	chromosomal *SCO4242* *(cisA)* locus deleted using pSS684	This study
SS549	∆*SCO4242::apr* φ*BT1 (PermE*-sfgfp)*	pSS150 integrated at ΦBT1 attachment site of SS539	This study
SS557	∆*SCO4242* φ*BT1(PermE*-SCO4242-3xFLAG)* *hyg^R, apr^R*	pSS735 integrated at φBT1 attachment site of SS539	This study
SS574	WT φ*C31* *(PermE*-sfgfp),* *apr^R, thio^R*	pSS758 integrated at φC31 attachment site of WT	This study
SS575	∆*SCO4242::apr* φ*C31 (PermE*-sfGFP)* *apr^R, thio^R*	pSS758 integrated at φC31 attachment site of SS539	This study
SS576	∆*SCO4242:apr* φ*BT1(PermE*-SCO4242-3xFLAG)* φ*C31 (PermE*-sfGFP)* *apr^R, thio^R, hyg^R*	pSS758 integrated at φC31 attachment site of SS557	This study
JS65	∆*SCO4242:apr* φ*BT1(PermE*-)* *hyg^R, apr^R*	pIJ10257 integrated at φBT1 attachment site of SS387	This Study
JS69	∆*SCO4242:apr* φ*BT1(PermE*-SCO4242-mcherry)* *hyg^R, apr^R*	pSS705 integrated at φBT1 attachment site of SS387	This study
Cosmid and Plasmids			
StD8A	Cosmid vector containing coding sequence for *SCO4242* km^R, carb^R		http://strepdb. streptomyces.org.uk
pIJ773	pBluescript KS (+) containing the apramycin resistance gene *apr* and *oriT* of plasmid RP4,		[5]

	flanked by FRT sites (Apr^R). Used as template for the amplification of the *apr-oriT* cassette for 'REDIRECT' PCR targeting, apr^R		
pIJ10257	Cloning vector for the conjugal transfer of DNA (under control of the *ermE** constitutive promoter). Integrates at the *ΦBT1* attachment site, hyg^R		[6]
pET21b	Plasmid for overexpression of proteins in *E. coli*, carb^R		Lab stock
pKF351	Derivative of pKF280, integrates at the *ΦC31* attachment site, apr^R		[7]
pSS150	pIJ10257 carrying *PermE*-sfgfp*, hyg^R		[4]
pSS684	Mutated cosmid StD8A for REDIRECT containing *ΔSCO4242::apr, km^R, carb^R, apr^R*	The *SCO4242* coding sequence on the cosmid vector StD8A was replaced by an *oriT*-containing apramycin resistance cassette, which amplified from pIJ773 using primer 1676/1677	This study
pSS88	pIJ10257 carrying *mcherry*, hyg^R	*Streptomyces* Codon-optimised *mcherry* was inserted between the NdeI-XhoI site of pIJ10257.	This study
pSS705	*PermE*-SCO4242-mcherry, hyg^R*	*SCO4242* coding sequence amplified with primer 1731/1732 and assembled with pSS88 cut with NdeI-XhoI.	This study
pSS734	*PermE*-SCO4242- TEV-sfgfp, hyg^R*	A "GGS-TEV" amino acid sequence was added upstream of *sfgfp* from pSS150 by PCR using	This study

		primer 1800/1801. The PCR product was cut with XhoI-AvrI and inserted into pSS705 cut with the same enzymes.	
pSS735	*PermE*-SCO4242-3xFLAG, hyg^R*	*PermE*-SCO4242* fragment was isolated from pSS705 by restriction digestion with KpnI-XhoI and ligated with pIJ10257 cut with KpnI-XhoI.	This study
pSS758	pKF351 carrying *PermE*-sfgfp, apr^R, thio^R*	*sfgfp* amplified from pSS150 with primer 1851/1852 and assembled into pKF351 cut with BamHI-KpnI.	This study
Plasmid used to test membrane topology			
pKTop	Plasmid encoding a dual *pho-lac* reporter, km^R		[8]
pFRL1	Plasmid encoding the known *Streptomyces* cytoplasmic protein, SepH, in the dual *pho-lac* reporter, *km^R*		Felix Ramos-Leon (unpublished)
pFRL5	Plasmid encoding the *Streptomyces* membrane protein, RsbN, in the dual *pho-lac* reporter, *km^R*		Felix Ramos-Leon (unpublished)
pJUK131	pKTop carrying *SCO4242, km^R*	Codon optimized *SCO4242* amplified with primers JS111/JS112 from pJUK123 digested and inserted into *HinDIII*/*Kpn*I digested pKTop	This study
pJUK132	pKTop carrying *SCO4242* (1-310aa), *km^R*	Codon optimized *SCO4242* containing only codons for aa 1-310 amplified with primers JS111/JS113 from pJUK123 digested and inserted into *HinDIII*/*Kpn*I digested pKTop	This study
pJUK145	pKTop carrying *SCO4242* (1-285aa), *km^R*	Codon optimized *SCO4242* containing only codons for aa 1-285 amplified with primers JS111/JS114 from pJUK123 digested and inserted into HinDIII/KpnI digested pKTop	This study

pJUK146	pKTop carrying *SCO4242* (Δ295-305aa), *km^R*	Codon optimized SCO4242 omitting codons for aa 295-310 amplified with primers JS111/JS112 from pJUK133 digested and inserted into HinDIII/KpnI digested pKTop	This study

Table EV4: Oligonucleotides used in this study.

Name	Sequence (5'→3')
1676	GGAGAGATGACCACGCAGAACTGCGCCGAGTGCGGAACCATTCCGGGGATCCGTCGACC
1677	CGGCCCGGGTCAGGAGGAGCGGTTGGCGCTGGACGGGCCTGTAGGCTGGAGCTGCTTC
1691	ATATTCTAGAGATGACCACGCAGAACTGCGCC
1692	ATATGGTACCTCAGGAGGAGCGGTTGGCGC
1693	ATATAAGCTTGATGCCCCTGCCCTCTCCCAAC
1694	ATATGGTACCCGCGCGCTGTCCCCGATCACG
1731	TGGTAGGATCGTCTAGAACAGGAGGCCCCATATGACCACGCAGAACTGCGCC
1732	ATGTTGTCCTCCTCGCCCTTGGAGACCATCTCGAGGGAGGAGCGGTTGGCGCT
1767	AGCTGTTTCCTGTGTGAAATTGTTATCC
1800	ATATACTCGAGGGTGGCTCCGAAAACCTGTACTTCCAATCCATGTCCAAGGGCGAGGAGC
1801	AATTAACCTAGGTCACTTGTACAGCTCGTCCATGCC
1851	AAGTCGTGCTGCTTCATGTGGTCGGGGTACCCGACCCGAGCACGCG
1852	ATTACTGGACCGGATGAATTCACTTGGATCCGTTAATTAATCACTCGAGCTCCGGGCCCG
1930	CCAAGCTTGCATGCCTGCAGG
1931	TCACACAGGAAACAGCTATGACCACGCAGAACTGCGCC
1932	GGCATGCAAGCTTGGGATGCGGTCGAAGACGCGGC
1933	CCAAGCTTGCATGCCTGCAGG
1934	GGCATGCAAGCTTGGGATGCGGTCGAAGACGCGGC
JS105	CCTCCGCGAATCGGTCGTCTCTCGAGGGTGGCTCCGAAAACCT
JS106	GCGCAATTTTGGGTCGTCATATGTATATCTCCTTCTTAAAGTTAAACAAAATTATTTCTAGAGGGGAA
JS107	TTTAAGAAGGAGATATACATATGACGACCCAAAATTGCGCCGAATGT
JS108	TTTTCGGAGCCACCCTCGAGAGACGACCGATT
JS111	CATTAGAAGCTTTATGACGACCCAAAATTGCGCCG
JS112	CATTAGGGTACCGACGACCGATTCGCGGAGG
JS113	CATTAGGGTACCTGAACCCCGTCGGGAATATA
JS114	CATTAGGGTACCCCGATTCGATCGAAAACACGACGC

References:

1. Paget, M. S. B., Chamberlin, L., Atrih, A., Foster, S. J. & Buttner, M. J. Evidence that the Extracytoplasmic Function Sigma Factor ςE Is Required for Normal Cell Wall Structure in Streptomyces coelicolor A3(2). *J. Bacteriol.* **181**, 204–211 (1999).

2. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci.* **97**, 6640–6645 (2000).

3. T. Kieser M. J. Bibb M. J. Buttner K. F. Chater and D. A. Hopwood. *Practical Streptomyces Genetics*. (John Innes Foundation, Norwich, 2000).

4. Casu, B., Sallmen, J. W., Schlimpert, S. & Pilhofer, M. Cytoplasmic contractile injection systems mediate cell death in Streptomyces. *Nat. Microbiol.* **8**, 711–726 (2023).

5. Gust, B., Challis, G. L., Fowler, K., Kieser, T. & Chater, K. F. PCR-targeted Streptomyces gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 1541–1546 (2003).

6. Hong, H.-J., Hutchings, M. I., Hill, L. M. & Buttner, M. J. The role of the novel Fem protein VanK in vancomycin resistance in Streptomyces coelicolor. *J. Biol. Chem.* **280**, 13055–13061 (2005).

7. Schlimpert, S. *et al.* Two dynamin-like proteins stabilize FtsZ rings during Streptomyces sporulation. *Proc. Natl. Acad. Sci.* **114**, E6176–E6183 (2017).

8. Karimova, G., Robichon, C. & Ladant, D. Characterization of YmgF, a 72-Residue Inner Membrane Protein That Associates with the Escherichia coli Cell Division Machinery. *J. Bacteriol.* **191**, 333–346 (2009).