Altered thymic niche synergistically drives the massive proliferation of malignant thymocytes.

Erika Tsingos [1,*,#], **Advaita M. Dick** [2,*], and **Baubak Bajoghli** [2,3,#]

[1] Computational Developmental Biology Group, Institute of Biodynamics and Biocomplexity, Utrecht University, Utrecht, The Netherlands

[2] Department of Hematology, Oncology, Immunology, and Rheumatology, University, Hospital of Tübingen, Otfried-Müller-Strasse 10, 72076 Tübingen, Germany.

[3] Present address: Austrian BioImaging/CMI, Dr. Bohr-Gasse 7, 1030 Vienna, Austria

[*] These authors contributed equally to this work.

[#] Corresponding authors: Baubak Bajoghli (baubak.bajoghli@vbcf.ac.at) and Erika Tsingos (e.tsingos@uu.nl)

ORCID identifiers:

Erika Tsingos: 0000-0002-7267-160X

Advaita M. Dick: 0009-0000-4888-1271

Baubak Bajoghli: 0000-0002-7368-7523

ABSTRACT

The discovery of genetic alterations in patient samples over the last 20 years has promoted a cell-autonomous view of proliferative expansion during T-cell acute lymphoblastic leukemia (T-ALL) development in the thymus. However, the potential contribution of non-cell-autonomous factors, particularly the impact of thymic epithelial cells (TECs) within the thymic niche during the initiation phase, remains unexplored. In this study, we employ a unique combination of a cell-based computational model of the thymus and *in vivo* experiments. We systematically analyze the impact of 12 cell-autonomous and non-autonomous factors, either alone or in combinations, on the proliferation of normal and malignant thymocytes with interleukin-7 receptor (IL7R) gain-of-function mutations or elevated IL7R levels, as observed in T-ALL patients. By simulating over 1500 scenarios, we show that while a dense TEC network favored the proliferation of normal thymocytes, it inhibited the proliferation of malignant lineages, which achieved their maximal proliferative capacity when TECs were sparsely distributed. Our *in silico* model predicts that certain mutations could accelerate proliferative expansion within a few days. This was experimentally validated, revealing rapid onset of thymus lymphoma and infiltration of malignant T-cells into other organs within 8 days of medaka (*Oryzias latipes*) embryonic development, thus revealing that modifications in the thymic niche and oncogenes in thymocytes together accelerate the disease development. Our results also suggest that negative feedback from the proliferative state inhibits differentiation of thymocytes, thereby prolonging the proliferative state and further fueling malignant expansion. Overall, this work reveals the critical impact of TEC-thymocyte interactions in both the initiation and progression of disease.

INTRODUCTION

T-cell acute lymphoblastic leukemia (T-ALL) is a malignancy characterized by the proliferation of immature T cells, known as thymocytes, and is subclassified according to the stage of thymic maturation (Terwilliger and Abdul-Hay 2017; De Keersmaecker et al. 2013). Representing approximately 10-15% of pediatric cases and 25% of adult cases of all acute lymphoblastic leukemia (Vadillo et al. 2018), T-ALL originates in the thymus. This organ is the primary site where early T cell progenitors (ETPs) originating from the hematopoietic tissue differentiate into mature, immunocompetent T cells. Thymocytes follow a well-orchestrated migratory path within distinct thymic niches, receiving essential extrinsic signals for their proliferation, differentiation, and selection (Takahama 2006). Thymic epithelial cells (TECs) play a central role in this process, acting as the main cellular component of these niches, and influencing thymocyte migration and differentiation by expressing crucial growth factors, ligands (such as DLL4), or releasing chemokines (e.g. CCL25) and cytokines such as interleukin-7 (IL-7) into the thymic niche (Yui and Rothenberg 2014; Koch et al. 2008; Zamisch et al. 2005).

The etiology of T-ALL is a topic of extensive research. Isolation and genome sequencing of malignant thymocytes from patients have uncovered mutations in over 100 putative driver genes (Liu et al. 2017; Girardi et al. 2017). Notably, over 50% of T-ALL patients harbor gain-of-function mutations in the NOTCH1 gene, which encodes the receptor for DLL4, while around 10% of patients exhibit similar mutations in the *IL-7 receptor* (*IL7R*) gene (Liu et al. 2017). The leukemogenic potential of constitutive activation of these receptors has been demonstrated across various vertebrate models (González-García et al. 2009; Silva et al. 2021; Shochat et al. 2011; Zenatti et al. 2011; Oliveira et al. 2022; Chen et al. 2007; Blackburn et al. 2012), emphasizing the evolutionary conservation of thymic developmental mechanisms that become misrouted in T-ALL (Bajoghli et al. 2019). These studies have contributed valuable insights into the impact of genetic alterations and oncogenic pathways on T-ALL development. However, the gene-centric view provided only limited understanding of the initiation and progression of the disease. In recent years, the growing evidence of the role of an abnormal tumor microenvironment in carcinogenesis strongly supports a more integrative view that considers the convergence of cellular genetics and the surrounding malignant niche as crucial elements for disease progression (Vadillo et al. 2018). In the context of T-ALL, however, the impact of the thymic niche or the role of TECs has not been studied.

Computational and mathematical models are powerful tools for integrating complex biological processes and can be used to test new hypotheses in health and disease (Ji et al. 2017; Metzcar et al. 2019; King et al. 2021). Several such models have been developed to study population dynamics during T cell development (Robert et al. 2021; Aghaallaei et al. 2021; Efroni, Harel, and Cohen 2007; Thomas-Vaslin et al. 2008; Vibert and Thomas-Vaslin 2017; Souza-e-Silva et al. 2009). Our recently developed cell-based computational model, for example, simulates individual thymocytes using agents within a spatially-resolved "virtual thymus" to explore cell-level behaviors and their effects on thymocyte population dynamics. This model was originally developed to investigate the impact of thymic niche signals and intrathymic cell localization on the αβ/γδ T cell sublineage outcomes (Aghaallaei et al. 2021).

In this study, we developed new computational tools to enhance the virtual thymus model, enabling us to identify both cell-autonomous and non-autonomous factors in thymocytes and TECs that could drive the proliferative expansion of a lineage derived from a single progenitor cell (hereafter referred to as a clone) within the thymus. We conducted an unbiased systematic analysis of various parameters, including TEC shape and architecture, IL-7 signaling, cell cycle, duration of the proliferative phase, and cell migration. This analysis facilitated a detailed comparison of proliferative expansion between wildtype (WT) and lesioned clones with alterations in the IL7R and NOTCH1 receptors. Through simulations of over 1500 scenarios, we identified factors that promote substantial expansion of malignant clones. In particular, modifications in the TEC network and niche emerged as a previously uncharacterized factor that could synergistically accelerate clonal expansion. Subsequent *in vivo* experiments using medaka fish (*Oryzias latipes*) confirmed the outcomes of our *in silico* analysis, solidifying the predictive accuracy of our virtual model, and revealing the impact of TECs on the initiation and progression of T-ALL.

RESULTS AND DISCUSSION

A virtual thymus to model the medaka embryonic T cell development

The details of the virtual thymus model implementation and parameter estimation have been extensively described previously (see Supplementary Material of Aghaallaei et al. 2021), and in the following we will only delineate its main features. The model considers both spatial and temporal aspects of early thymus development at multiple scales: Cells are physically

represented as one or multiple spheres that mechanically interact with adhesive and repulsive forces to capture cell crowding; each cell has an independent internal state that changes dynamically based on subcellular signaling modeled with ordinary differential equations or with phenomenological rules; and at the tissue level a partial differential equation is used to solve the diffusion of extracellular molecules such as cytokines. To reduce computational cost, the computational representation focuses on a 5 μm deep slice of the lower half of the radially symmetric organ, approximately 1/10 of the total volume of a medaka embryonic thymus (Figure 1 - Supplement 1A, B). Consistent with observations in vertebrates, including in medaka (Bajoghli et al. 2009; 2015; Aghaallaei et al. 2021; 2022), ETPs within our virtual thymus mechanically and biochemically interact with TECs, proliferate and differentiate in response to signals from the thymic niche, and subsequently diverge into two distinct T cell sublineages before selection and exit from the thymus (Figure 1A). In this work, we define cells of a lineage derived from a single founder ETP as a "clone", and we do not consider differences in immunological clonotypes that emerge after recombination of the T cell receptor.

In the virtual thymus model, factors governing thymocyte motility, including cell speed (Figure 1B) and directionality (Figure 1C) were fine-tuned based on quantitative noninvasive imaging of the thymus using multiple medaka transgenic reporter lines (Bajoghli et al. 2009; 2015; Aghaallaei et al. 2021). Based on these experimental observations, we modeled entry into the organ by creating new cells at the bottom of the simulated domain (ventral to the organ), these cells move into the thymus and, some time after differentiation, either leave the organ by exiting from the ventral side as a naïve T cell or undergo cell death by gradually shrinking and eventually being removed from the simulation (Figure 1 – Supplement Movie 1-2). The model accounts for the time taken for a single progenitor to enter and exit the thymus, which is estimated at approximately 3 days in zebrafish (Hess and Boehm 2012) and medaka (Bajoghli et al. 2015). For time calibration, each simulation step was set to represent 15 seconds. With a total of 48,000 steps per simulation, the entire simulation spans 720,000 seconds, equivalent to about 8.3 days (Figure 1D). This time interval was chosen to recapitulate thymic development in medaka from 2.5 to 11 days post-fertilization (dpf), because at 2.5 dpf the first ETP enters the embryonic thymus (Bajoghli et al. 2019). Correspondingly, each simulation starts with entering of the first ETP into the virtual thymus and ends with a homeostatic cell population of roughly 100 cells (mean ± standard deviation: 98±7; Figure 1 – Supplement Movie 1).

Figure 1 | The virtual thymus model enables spatiotemporally resolved clonal analysis. (A) Schematic overview highlighting 6 key steps during T-cell development, which are integrated into the virtual thymus model: (1) ETPs enter the thymus niche; (2) they receive Dll4 and IL-7 signals from the TECs. Triggered by these signals, (3) thymocytes undergo proliferation and (4) differentiation into two distinct T-cell sublineages (5). Finally, thymocytes undergo selection, either leaving the organ or undergoing cell death (6). (B-C) Experimentally measured ("Medaka") and computationally modeled ("Virtual") cell speed (B) and cell directionality (C) for ETPs and thymocytes. Bars are mean values, and error bars standard deviation. (D) Timeline of medaka embryonic development (top) compared to simulated time (bottom). As indicated by the cartoon at the bottom, the first ETP enters the thymus roughly at 2.5 dpf (t = 0 h in the simulation). At roughly 60 h (roughly 5 dpf), the thymus reaches its homeostatic cell population, with both cell entry and exit. (E) Representative snapshots of the simulation at homeostasis showing spatial expression patterns and signaling activity of components of IL7R and Notch1 signaling pathways in TECs (top two panels) and thymocytes (bottom 4 panels). Data are shown in arbitrary units normalized to a scale from 0 to 1. (F) Left: Cartoon explaining the asynchronous arrival (homing) of new cells into the organ, their clonal expansion, and asynchronous cell death or exit. Right: Visualization of clonal diversity in a typical simulation; each color shade uniquely indicates a clonal lineage. The height of each colored patch indicates the number of cells in that clone, the length indicates simulation time. For illustrative purposes, a black arrow indicates the homing of a sample of lineages. (G) Data from the right panel in (F) was normalized to time of homing. This visualization highlights developmental phases of thymocyte lineages. Gray arrowheads indicate rounds of cell division. Scale bar 50 cells. Abbreviations: ETP = early thymic progenitor, T = Thymocyte, *N* = number of replicates (biological or computational).

At 11 dpf, the medaka thymus contains 877±146 (N=7) thymocytes. Considering that the virtual thymus represents 1/10 of the medaka embryonic thymus, the observed cell population in the virtual thymus model reproduces the temporal dynamics in medaka within the margin of error.

Each cell's internal state and decision to proliferate or differentiate depends on intrinsic and extrinsic factors that are integrated by signaling pathways. The virtual thymus includes Delta-like 4 (Dll4), the ligand of Notch1 receptor, and interleukin-7 (IL-7) cytokine as two cell-extrinsic factors provided by the TECs (Figure 1E). Consistent with medaka WT embryonic thymus (Aghaallaei et al. 2021; 2022), all virtual TECs in this model express the Dll4 (Figure 1E, top-right panel), with a subset spatially releasing IL-7 into the environment (Figure 1E, top-left panel), creating a short-ranged IL-7 cytokine gradient in the thymic cortex niche (Aghaallaei et al. 2021). In terms of cell-intrinsic factors, all ETPs uniformly express the Notch1 receptor (Figure 1E, middle-right panel), and each cell lineage expresses the IL-7 receptor (IL7R) at a constant level chosen randomly between 0 and 1 (Figure 1E, middle-left panel). In each simulation, new ETPs continually enter the thymic niche at regular intervals. Following engagement with TECs and receiving Notch1 and IL-7 signals (Figure 1E, bottom panels), they proliferate and differentiate (step 3 in Figure 1A). The cell cycle was modeled with an average duration of 7 hours and was subdivided into G1, S, G2, and M phases. In line with *in vivo* observations (Ruijtenberg and van den Heuvel 2016), virtual cells require external pro-proliferative signals during the G1 phase to commit to S through M. Once differentiated, cells can no longer enter the cell cycle. The dynamics of pro-proliferative signals and the proliferation stop after differentiation allow clones *in silico* to undergo up to four rounds of cell division (Aghaallaei et al. 2021).

In this work, we implemented a code that enables us to meticulously trace the unique identity of each cell upon entering the virtual thymus and subsequently track the identity of its descendants (Figure 1F, left panel; Figure 1 – Supplement 1 C). This tool identified various waves of clones because of the constant influx of new ETPs and negative selection and efflux of fully differentiated T cells (Figure 1F, arrows in the right panel, Figure 1 – Supplement Movie 1). We note a slight variability in the clone sizes, which could arise from three processes. First, random variations in the expression level of IL7R impact IL-7 signal transduction and, consequently, proliferation (Aghaallaei et al. 2021). Second, due to random cell motility and cell crowding effects, certain cells could coincidentally remain in close contact with IL-7-secreting TECs, maximizing their exposure to pro-proliferative

signals such as IL-7 and Notch ligand, and thus increasing their chances of entering the cell cycle. Third, each cell determines the duration of its next cell cycle by drawing an Erlang-distributed random variable with a mean of 7 hours and a standard deviation of ≅0.99 hours.

After normalizing all clones based on their time of entry into the thymus, the pattern of developmental phases became clear across all clones (Figure 1G). In this visualization, the now near-synchronous rounds of cell division across clones can be seen as bumps in the graph indicating clonal expansion (Figure 1G, gray arrowheads). During the commitment phase, which has a minimum duration of 24h but can be extended due to insufficient Notch signaling, thymocytes were competent for proliferation, leading to an increase in the total cell number per clone. In the differentiation phase, which has a fixed duration of 24h, clones reached a maximum size. Note that differentiation prevents further entry into the cell cycle but permits cells that passed G1 and therefore committed to S through M phases to complete their division. Finally, clones underwent selection and exit from the virtual thymus, thus leaving the simulation. Together, the enhanced virtual thymus model facilitated the investigation of the heterogeneity and temporal dynamics of individual clones originating from a founder progenitor cell.

Modeling the impact of TEC architecture on thymocyte population size

In vivo, individual TECs exhibit a distinctive star-shaped morphology, and protrusions of neighboring TECs contact each other forming a three-dimensional network (Figure 2A, left panel). Using confocal imaging, we estimated the relative area occupied by thymocytes (53±3%; N=4) and TECs (47±3%; N=4). A similar morphology and tissue density was replicated in our *in silico* model (Figure 2A, right panel), with thymocytes taking up on average 56±2% and TECs 44±2% of the volume in homeostasis. We wondered to what extent TEC morphology and density could impact thymocyte population size. This aspect is difficult to study experimentally and is, therefore, an ideal use case for the virtual thymus model. In a series of simulations, we combinatorially varied (i) the size of TECs, (ii) the number of their protrusions, and (iii) the TEC cell density (Figure 2B). The average homeostatic thymocyte population size in each simulation was then used as a readout. The results from 26 different tested conditions predicted that a higher density of larger TECs with more protrusions led to an almost two-fold increase in the number of thymocytes (200±13, N=3; scenario 26 in Figure 2C, Figure 2 – Supplement 1A bottom) compared to the reference

condition (98±7, N=19; Reference in Figure 2C). Conversely, scenarios where these parameters were reduced, particularly a reduced TEC size, also diminished the homeostatic thymocyte population size (scenarios 1-14 in Figure 2C). The condition with the least amount of cells, scenario 1, had a reduced TEC size, but an unchanged number of protrusions and density. The condition diametrically opposite to scenario 26 was scenario 6, with a reduced TEC size, reduced number of protrusions, and reduced TEC density (Figure 2 – Supplement 1A top). In both scenario 1 (45±1, N=3) and scenario 6 (51±2, N=3), the thymic population size was only about half of that in the reference condition (98∓7, N=19). Statistical analysis confirmed that all three parameters and their interactions were significant predictors of total thymocyte population size (Figure 2 – Supplement 1B-C). The statistically significant interaction can be explained by the fact that higher TEC density increases the number of TECs, and thereby amplifies the effect of an increased TEC radius or number of protrusions. Thus, these three alterations of the thymic niche morphology act synergistically to modulate thymocyte population size.

There are two possibilities that could explain the variations in total thymocyte numbers in simulations. Either over- or under-proliferation of a few thymocyte lineages, or the cumulative impact of subtle changes across several thymocyte clones. To distinguish between these possibilities, we used our clonal analysis tool for scenarios 6 and 26 (Figure 2D). This analysis revealed that variations in total thymocyte numbers were a consequence of changes in proliferation across all clones to a similar degree. These results predict that the TEC architecture has a direct impact on thymocyte proliferation rate, which fits well with the fact that TECs act as the main source of ligands, growth factors, cytokines, and chemokines (Gameiro, Nagib, and Verinaud 2010). Any changes in the TEC architecture could therefore influence the amount of cytokine production. In our virtual thymus model, this affects the extracellular spatial distribution of IL-7 cytokine (Figure 2E). To further explore this, we tested two additional scenarios. In scenario 27, TECs maintained their reference architecture and density but all expressed IL-7, leading to a more uniform cytokine distribution in the thymic environment (Figure 2E). The extracellular level of IL-7 (Figure 2F) and thymocyte population size (155±11, N=5; Figure 2G) in scenario 27 was similarly high as in scenario 26, where the size, protrusion number, and density of TECs were increased. Conversely, when TECs did not express IL-7 (scenario 28), the thymocyte population size (Figure 2G) was low (42±1, N=5), akin to scenario 6, where the size, protrusion, and density of TECs were reduced.

Figure 2 | TEC density influences thymocyte proliferation via IL7R signaling. (A) Left: Detail of confocal section highlighting the tight spatial interaction between TECs and thymocytes (T). Scale bar 5 μm. Right: Detail from a three-dimensional render of a simulation snapshot; TECs cyan, thymocytes green. (B) Parameter permutations tested. (C) Impact of parameter permutations on average homeostatic cell population. Bars indicate means, error bars standard deviation. (D) Clonal lineages normalized to time of thymic entry for reference (left), a scenario with low average population (middle) and a scenario with high average population (right). Scale bar: 100 cells. (E) Extracellular IL-7 gradient in simulations; sum projection of all z-planes of a simulated confocal stack. Units are arbitrary. (F) Mean extracellular IL-7 concentration in the entire simulated volume. Note that these values are lower than in (E) because of averaging over the volume, including empty space (black areas in (E)). (G) Thymocyte population size over time averaged over several simulation runs. Shaded area indicates standard deviation. (H) Illustration of experimental setup for IL-7^{KO} and TEC:IL-7^{HI}. At least 3 technical replicates were done for the injection of each construct. (I) Representative images of embryos stained with GFP (green) and pH3 (red) of WT, IL-7^{KO} and TEC:IL-7^{HI}. Scale bar 20μm. (J) Numbers of pH3 positive cells per thymus normalized to mean of WT. *N* represents the number of individual fish. Data show means ± standard deviation.

Interestingly, although the average extracellular IL-7 level is at its highest in scenario 27 with ubiquitous expression, the spatially restricted elevated IL-7 concentration in scenario 26 is more effective at driving thymic population growth (Figure 2E-G). Thymocytes first enter the organ at its periphery from below (at the ventrolateral site), where IL-7 levels are highest, and tend to migrate to the IL-7-depleted center of the thymus as they become non-proliferative (Aghaallaei et al. 2021; see also Figure 1 – Supplement Movie 1). Thus, in scenario 26, thymocytes in their proliferative phase immediately encounter elevated IL-7 at the organ periphery, stimulating cell proliferation. In contrast, in scenario 27, thymocytes that enter the organ periphery encounter comparable levels of IL-7 to the reference scenario, and only after entering deeper into the organ can the ubiquitous expression of IL-7 unfold its effects. This delay contingent to random cell migration (which is further exacerbated by cell crowding posing an obstacle), reduces the impact of elevated IL-7 in scenario 27. These results highlight how the spatial structure of the thymic niche can impact on thymocyte population dynamics.

Our simulations predict that the availability of IL-7 provided by TECs has a direct impact on thymocyte population size. To validate this prediction *in vivo*, we conducted two functional analyses using the medaka model organism (Figure 2H). In one experiment, employing the CRISPR-Cas9 technique, we knocked out the *il7* gene (Figure 2- Supplement 2A, B) in a medaka transgenic line, where thymocytes express green fluorescent protein (GFP). Consistent with our *in silico* outcomes, *il7* crispant embryos displayed fewer thymocytes and a smaller thymus size (Figure 2I, Figure 2 - Supplement 2C, D). To estimate the extent of cell proliferation, we counted mitotic cells throughout the entire thymus using the M-phase marker phospho-histone 3 (pH3) and normalized values to the mean of the control. This analysis further supported that a reduction in IL-7 availability in the thymus reduces thymocyte proliferation (Figure 2I, J). In a second experimental setup (Figure 2H), we artificially increased *il7* levels in the thymic niche by injecting the ccl25a:*il7* construct into embryos (hereafter called TEC:IL-7^{HI}), resulting in a strong upregulation of the levels of *il7* produced by the TECs (Aghaallaei et al. 2021). Compared to WT, the thymus of TEC:IL-7^{HI} embryos appeared visibly larger, and the number of pH3+ cells was significantly increased (Figure 2I, J). Therefore, the *in vivo* results confirm the critical role of IL-7 for thymocyte proliferation in medaka, a mechanism that is evolutionarily conserved among vertebrates (Iwanami et al. 2011). Additionally, the alterations in thymic size suggest a regulatory crosstalk between thymocyte proliferation and TEC architecture.

Figure 3 | High IL7R signaling promotes proliferation. (A) Illustration of scenarios without extracellular IL-7 depletion (left) and with extracellular IL-7 depletion (right). (B) Ordinary differential equation used to model the IL7R signaling activity. Parameters that were modified are highlighted. (C) Effect of IL7R signaling-related parameter permutation. Bars indicate means and error bars standard deviation. (D) Left: Mean extracellular IL-7 concentration in the entire simulated volume. Right: Extracellular IL-7 gradient in simulations; sum projection of all z-planes of a simulated confocal stack. Units are arbitrary. The introduction of IL-7 depletion by thymocytes leads to a slight reduction in extracellular IL-7 availability.

Modeling the impact of IL-7 signaling activity on thymocyte population size

Given the important contribution of extracellular IL-7 on thymocyte population size, we next evaluated the impact of (i) extracellular IL-7 depletion (e.g., via ligand internalization), (ii) the rate of signal transduction activation upon IL7R and IL-7 binding, and (iii) the rate of decay of signaling activity (Figure 3A, B). In our virtual thymus model, the IL-7 signaling activity (σ_{IL7}) of a thymocyte over time (t) is modeled phenomenologically using a function involving the IL7R concentration ([IL7R]), the average extracellular IL-7 concentration ($\langle$[IL-7$_{ex}$]$\rangle$), a parameter (a_{IL-7}) that scales the strength of signal transduction activation, and another parameter (d_{IL-7}) that scales the rate at which the signaling activity diminishes over time (Figure 3B). As before, we combinatorically perturbed these three factors, simulating 17 different scenarios, to assay their impact on thymocyte population size (Figure 3C). As expected, we observed a positive correlation between the level of signal transduction activation a_{IL-7} and thymocyte numbers.

Conversely, variations in signaling decay rate $d_{\text{IL-7}}$ showed the opposite
effect. The rate of signal transduction deactivation models cellular short-
term memory, thus a lower signaling deactivation rate indicates that cells
retain their IL-7 stimulus for a longer duration. Because IL-7 signaling
activity σ_{IL7} promotes cell cycle entry, we expect that parameter changes
that increase stimulus duration should promote cell division. Indeed, the
scenario with the highest number of thymocytes occurred when signaling
deactivation was low and activation was high (274±11, N=4; scenario 45 in
Figure 3C). The diametrically opposite combination had the least amount of
cells (42±1, N=4; scenario 29 in Figure 3C). Statistical analysis confirms that
signal transduction activation $a_{\text{IL-7}}$, and the signal transduction deactivation
rate $d_{\text{IL-7}}$ were both statistically significant predictors of thymocyte
population size (Figure 3 - Supplement 1). There was no statistically
significant effect of parameter interactions, likely because these parameters
exert their effect independently. There was only a limited reduction in
population size with IL-7 depletion (91±11, N=4; scenario 36 in Figure 3C)
compared to no depletion in our reference setting (98±7, N=19). Overall, IL-7
depletion did not have a significant effect on thymocyte population (Figure
3 - Supplement 1). However, we noted that population size difference was
more pronounced at higher cell numbers, e.g., comparing between
scenario 44 (237±7, N=4) and scenario 45 (274±11, N=4). Similarly, the mean
extracellular IL-7 content in the virtual thymus barely changed when
comparing the reference to scenario 36 but was noticeably reduced in
scenario 44 compared to scenario 45 (Figure 3D). Therefore, thymocyte
proliferation could self-inhibit via depletion of extracellular IL-7, but we
expect this effect to be small unless cell numbers are massively increased.

Together, the outcomes of our simulations reveal that molecular and
cellular changes, specifically those capable of increasing the extrinsic factor
IL-7 in the niche — such as a higher density of TECs or an elevated
expression level of IL-7 by individual TECs — directly contribute to an
increase in thymocyte population size. Tuning thymocytes' sensitivity to IL-
7 by manipulating IL7R signaling can further amplify proliferation. Finally,
we expect that cytokine internalization by thymocytes has only a mild
inhibitory effect on excessive population growth by limiting the availability
of the pro-proliferative IL-7 in the extracellular space.

A systematic approach identified synergistic factors promoting clonal expansion of IL7R-lesioned clones

Our 45 tested scenarios thus far reached a maximum three-fold induction
of thymocyte population size when parameter changes promoted

increased IL-7 signaling. While the activation of IL-7 signaling is linked to T-
ALL development, this pathology is associated with proliferation of clones
carrying somatic mutations (Oliveira et al. 2019; 2022; Silva et al. 2021). We
therefore modified the code of our virtual thymus model to enable the
introduction of parameter changes exclusively in a single clone and all of its
progeny (hereafter referred to as the lesioned clone). Inspired by
observations that 10% of T-ALL patients exhibit dominant active mutations
in the *IL7R* gene (Liu et al. 2017; Shochat et al. 2011; Zenatti et al. 2011), and a
significant subset of T-ALL patients with active IL7R signaling display
elevated *IL7R* expression levels (Silva et al. 2021), two types of lesions were
introduced: (1) a dominant active IL7R lesion (hereafter called IL7R^{DA} lesion),
(2) an IL7R overexpression lesion (hereafter called IL7R^{HI} lesion). For the
IL7R^{DA} lesion, we set the IL7R level of the lesioned clone to the maximum
WT level of 1 and allowed it to activate IL-7 signaling regardless of
extracellular IL-7 cytokine, mimicking the dominant activation mutations in
the *IL7R* gene (Figure 4A, middle panel). For the IL7R^{HI} lesion, we mimicked
overexpression of the *IL7R* gene by assigning a value of 10 for the IL7R level
in the lesioned clone, i.e. 10-fold higher than the maximum attainable in the
WT population (Figure 4A, right panel). In both scenarios, non-lesioned
thymocytes retained reference IL7R levels randomly ranging between 0 to
1 (Aghaallaei et al. 2021). In each simulation, a single lesioned clone entered
the thymic niche after the 60th hour of a simulation, that is shortly after the
establishment of a homeostatic population size (Figure 4B, Figure 4 -
Supplement Movie 1).

First, we evaluated the impact of IL7R^{HI} and IL7R^{DA} lesions in the reference
condition, meaning no additional modifications were made to our model.
The clonal analysis of simulations revealed that introducing a lesioned clone
did not markedly alter the division behavior of non-lesioned clones in the
same simulation (scenario with no lesion: 1.04±1.26 rounds of cell division;
scenario with IL7R^{DA}: 0.98±1.22 rounds of cell division; scenario with IL7R^{HI}:
1.04±1.23 rounds of cell division; Figure 4B-C). The large variance in non-
lesioned clones is consistent with our previous work (Aghaallaei et al. 2021),
which showed that clones expressing endogenously high levels of IL7R
undergo more rounds of cell division in the virtual thymus, whereas clones
with lower levels of IL7R proliferate very little or not at all. In contrast,
lesioned clones averaged just above 4 rounds of cell division (4.20±0.34 for
IL7R^{DA} and 4.10±0.13 for IL7R^{HI}). Consequently, the size of IL7R^{DA} and IL7R^{HI}
clones was nearly 8-fold larger than an average non-lesioned clone (Figure
4C). This result suggests that lesioned clones acquired a distinct proliferative
advantage over non-lesioned clones in our virtual thymus model.
Nevertheless, since the expansion of lesioned clones amounted to, at best,

only one additional round of cell division compared to the upper range of the non-lesioned clone distribution (Figure 4C), our virtual thymus model predicts that a single lesion in the *IL7R* gene will not be clinically impactful.

Therefore, we next explored whether additional modifications in both cell-autonomous and non-autonomous factors might substantially enhance the clonal expansion. To identify these factors, we decided to undertake a systematic approach and tested combinations of scenarios affecting the TEC architecture and IL-7 signaling parameters, as tested in Figures 2 and 3, together with changing other parameters affecting proliferation and differentiation of all clones (Figure 4D). Furthermore, we considered scenarios in which only lesioned clones exhibited additional modifications, such as reduced cell motility, slower differentiation time, or autocrine IL-7 production. The inclusion of the latter was inspired by a study showing that certain malignant T cells derived from T-ALL patients possess the ability to ectopically express IL-7 cytokine (Buffière et al. 2019). In theory, there are 104976 possible permutations of these scenarios. To reduce the space of possibilities and thus the computational cost, we prioritized alterations expected to increase proliferative potential based on outcomes shown in Figures 2 and 3. We also decided to focus on simulating the extremes; e.g., low and high levels shown in Figure 4D. In addition, we simulated a small sample of scenarios expected to decrease proliferative potential, such as a reduction in TEC density. In the end, 1580 scenarios were simulated. The clonal analysis tool was used to compare clone size and the number of cell division rounds between lesioned and non-lesioned clones under the same conditions. Overall, we observed a similar proliferation pattern among IL7R^{HI} and IL7R^{DA} clones in most of the tested scenarios. The lesioned clones consistently showed a higher proliferation rate than their counterpart clones (Figure 4E; Figure 4 – Source Data 1). This trend persisted even when parameters known to generally increase cell division for all cells, such as mean cycle duration and proliferation duration, were modified (Figure 4 - Supplement 1, Figure 4 – Supplement Movie 2). In fact, lesioned clones disproportionally gained up to 6 rounds of cell division by modulating proliferative parameters, while normal clones only gained 2 rounds of cell division in the same scenarios. This difference results from cell crowding creating a disproportional disadvantage to non-lesioned clones: In the scenario with increased proliferation, the first clones to colonize the organ can massively proliferate and completely surround the TECs, preventing later arriving clones from accessing TEC-derived pro-proliferative signals (Figure 4 – Supplement Movie 2, right column). Thus, these few "lucky" early colonizers prevent late cells from attaining their full proliferative potential. Lesioned clones carry an intrinsic pro-proliferative advantage due to their

Figure 4 | Systematic screen shows that lower TEC density promotes malignant expansion. (A) Schematic representation of the major lesions to IL7R that were implemented in the model. (B) Representative clonal plot highlighting in black the lesioned clone. (C) We defined clone size as the maximum number of terminal leaves of a lineage, regardless of eventual fate of each cell. Clone size distribution in the reference simulation (REF), compared to IL7R^{DA} and IL7R^{HI}. REF: N=20 simulations, with 2088 clones total: IL7R^{DA}: N=13 simulations with 1330 non-lesioned clones and 13 lesioned clones. IL7R^{HI}: N=13 simulations with 1336 non-lesioned clones and 13 lesioned clones. (D) Schematic of tested scenarios shown in (E). (E) Simulated permutations ordered along increasing mean lesioned clone size for the IL7R^{HI} condition. Expected effect on proliferation is a qualitative estimate based on preliminary simulations. A total of 1580 permutations were tested with a varying amount of replicates per permutation. Error bars indicate standard deviation. Note that in (C) and (E) the axis is in base-2 logarithmic scale to better illustrate rounds of cell division.

lesions in the IL7R, enabling proliferation despite a lack of direct contact with TECs. In contrast, modifying parameters that affected IL-7 activity only enhanced the proliferation rate of non-lesioned clones but showed no

impact on either IL7R^{HI} or IL7R^{DA} clones (Figure 4 - Supplement 2). While this outcome is expected for IL7R^{DA} clones, which are insensitive to IL-7, for IL7R^{HI} clones it indicates that 10-fold receptor overexpression is enough to activate downstream pro-proliferative effects of the pathway regardless of the parameter permutations that we tested. In contrast, proliferation of normal clones is strongly affected in these conditions, gaining up to 3 rounds of cell division between least and most proliferative conditions.

In our systematic approach, we did not detect an added effect on clonal expansion from combining reduced cell motility or autocrine IL-7 production with IL7R^{HI} or IL7R^{DA} lesions (Figure 4 - Supplement 3). Added IL-7 secretion by lesioned clones with the autocrine lesion had only a small effect on proliferation of non-lesioned clones in the same simulation. The most critical lesion in conferring a proliferative advantage to lesioned clones was a delay in differentiation. This delay essentially doubled the duration of the proliferative phase, and indeed we observed up to 8 rounds of cell division, double the level of clones having only a single lesion in the IL7R. This result fits well with *in vivo* data suggesting that perturbations in differentiation have been implicated in contributing to the initiation and progression of cancer (Ruijtenberg and van den Heuvel 2016).

Surprisingly, the expansion of lesioned clones was amplified when the TEC niche was at its sparsest – a result opposite to the non-lesioned clones which instead profit from a dense TEC niche (Figure 4 - Supplement 4). While non-lesioned clones gained 2 rounds of division with an increase in TEC density, lesioned clones lost 2 rounds of division. Indeed, parameter combinations with reduced density and size of TECs were overrepresented in the scenarios that most increased proliferation (Figure 4 - Supplement 5A-C). For example, in the most extreme scenario observed in our systematic approach, IL7R^{HI} or IL7R^{DA} clones underwent up to 18 rounds of cell division – far above the 12 rounds predicted from modifying each group of parameters in isolation (+6 from proliferation, +4 from added lesions, +2 from reduced TEC density, assuming an additive effect; Figure 4 – Supplement 1-4). This result suggests that a defective niche amplifies the effect of other pro-proliferative modulations such as slower differentiation and shorter cell cycle, conferring a synergistic advantage to IL7R^{HI} and IL7R^{DA} clones compared to their normal counterparts. Note that at 18 rounds of division lesioned clones were so massive that the total lesioned cell volume was over a 1000-fold larger than the volume of the simulated organ slice (Figure 4 - Supplement 5D). In comparison, in the homeostatic condition, the total volume occupied by both thymocytes and TECs amounted to only 0.7-fold of the available tissue volume. Though our simulations by default include

physical volume exclusion effects that prevent dense packing, we did not implement density-dependent feedback on proliferation, enabling cells to duplicate their volume at cell division and proliferate uncontrolled despite the lack of available space. Presumably, similar conditions *in vivo* would lead to thymus hyperplasia. Indeed, our experimental manipulations of IL-7 levels in the thymus had a direct effect on organ size (Figure 2I, Figure 2 - Supplement 2C, D).

Identifying a sparse TEC network as a new factor that could influence the clonal expansion of lesioned clones was an unexpected outcome. This is because, in the WT situation, a sparser TEC network results in decreased extracellular IL-7, leading to reduced clonal size (Figure 2). While being disadvantageous for non-lesioned clones, this compromised thymic niche proved to be optimal for a massive expansion of lesioned clones. Conversely, we observed that a denser TEC network – characterized by a higher density of TECs with more protrusions and larger size – positively influenced the population size of non-lesioned cells but had a negative impact on the IL7R^{HI} and IL7R^{DA} lesioned clones (Figure 4 - Supplement 4). Similarly, in conditions of high cell crowding, lack of contact with TECs put non-lesioned clones at a disadvantage, while lesioned clones appeared to benefit (Figure 4 – Supplement 1, Figure 4 – Supplement Movie 2). This outcome might be explained by the signaling dynamics within our virtual model, where the engagement of the NOTCH1 receptor on thymocytes with the DLL4 ligand on TECs, on the one hand, promotes proliferation and, on the other hand, is a prerequisite for the differentiation process until commitment to a non-dividing fate. In our virtual model, if the NOTCH1 receptor fails to engage with DLL4 on TECs due to their low density or cell crowding effects, cells will remain in an undifferentiated state for a longer time. Closer inspection of time-normalized clones in scenario 6 (very sparse TECs) and in scenario 26 (very dense TECs) indeed confirms that the thymocyte population has a longer turnover time when TECs are sparse (almost 80h, Figure 2D middle panel), and a shorter turnover when TECs are dense (less than 60h, Figure 2D right panel), indicative of the effect of Notch signaling. Together, cells in an environment with fewer TECs and thus lower Notch signaling will experience a prolonged proliferative phase but still require pro-proliferative signals to commit to the cell cycle. Owing to the higher levels of IL7R signal intrinsic to the lesion, lesioned clones will be more easily competent to proliferate even without Notch signaling and, therefore, benefit from a lower TEC density.

The interplay between IL7R and NOTCH1 signals in the clonal expansion of thymocytes

In our virtual thymus model, ETPs simultaneously assess whether the combined sum of IL7R and NOTCH1 signals surpasses a threshold required for entry into the cell cycle ((Aghaallaei et al. 2021); Figure 5A). Besides its effect on proliferation, NOTCH1 signaling also promotes ETP differentiation (Aghaallaei et al. 2021; 2022), and we used this fact and the observations in *notch1b* medaka mutant phenotypes to include an accelerating effect of NOTCH1 on differentiation in our model (Aghaallaei et al. 2021). Thus, NOTCH1 has a dual effect in thymocytes, which we implemented in our model as an incoherent feed-forward loop (Figure 5A): NOTCH1 signaling promotes proliferation and accelerates differentiation, but differentiation inhibits proliferation by preventing further entry into the cell cycle. IL7R signaling promotes proliferation independently of NOTCH1 and has no direct or indirect effect on the differentiation process. To further explore the interplay between these two factors, we decided to compare scenarios where lesioned clones exhibited only constitutive activation of the NOTCH1 receptor (hereafter called NOTCH1^{DA}) or in combination with either IL7R^{DA} or IL7R^{HI}. The simulations predicted that lesioned clones with only NOTCH1^{DA} modification displayed a very small advantage compared to non-lesioned clones; this increase in proliferation was only marginally higher than expected for a clone expressing the highest endogenous levels of IL7R (hereafter IL7R^{WT}; Figure 5B). In contrast, lesions that specifically delay differentiation led to higher proliferation. Indeed, the addition of NOTCH1^{DA} modification to IL7R^{DA} or IL7R^{HI} lesioned clones did not show any effect on clonal expansion but combining delayed differentiation with IL7R^{DA} or IL7R^{HI} doubled proliferation. This result indicates that modulating the effect of NOTCH1 globally does not affect proliferation in our model, which is likely due to feedback inhibition, but specifically targeting the differentiation process leads to clonal expansion.

To test the outcome of the virtual model, we performed a series of *in vivo* experiments to induce NOTCH1 and IL7R signaling in thymocytes. To constitutively activate NOTCH1 signaling in thymocytes, we took advantage of the previously cloned medaka *notch1b* intracellular domain (NICD) construct (Aghaallaei et al. 2021; 2022), which was driven by a thymocyte-specific promoter (Bajoghli et al. 2015). Furthermore, the destabilizing PEST domain from the NICD was removed in this construct (hereafter referred to as NICD$^{\Delta PEST}$; Figure 5C). This genetic modification was made due to the known impact of the PEST domain on protein stability, as nonsense mutations lacking the PEST domain of the human *NOTCH1* gene have been

Figure 5 | Interplay of IL7R and NOTCH1 potentiates clonal expansion *in vivo*. (A) Schematic of Notch1 and IL7R signaling and their regulation of thymocyte cell cycle entry, proliferation, and differentiation as implemented in the virtual thymus model. (B) Simulated data; rounds of cell division of normal and lesioned clones with different types of lesions in NOTCH1 or IL7R signaling. Note that the y-axis is base-2 logarithmic. Green boxes: Condition/lesion present. Gray boxes: Condition/lesion absent. (C) Illustration of plasmids for DNA micro-injection for transient overexpression of the different genes. Note that the expression of GFP was used to select successfully injected embryos. (D) Representative image: (left) of the WT control at 9 dpf where thymocytes are labeled in green with normal thymus size; (middle) of a thymus hyperplasia phenotype at 7 dpf; (right) same embryo developed T-ALL phenotype at 9 dpf, where infiltration in other organs was detected, namely in central nervous system (CNS) and gut. (E) Percentage of phenotypes: normal thymus, thymus hyperplasia and T-ALL at 11dpf. For each construct injection, we did at least three technical replicates. For detailed statistics see Figure 5 - Supplement Table 1. Statistical data rounded to 4 decimal places. (F) Number of pH3 positive cells per thymus counted during confocal imaging. (G) Representative confocal images of immunostaining against pH3 (red) and GFP (green). Scale bar is 20μm. *N* represents the number of biological samples.

frequently observed in T-ALL patients (Breit et al. 2006; Ferrando 2009; Weng et al. 2004) (Figure 5 - Supplement 1A). To mimic the *in silico* IL7R^{HI} clones, we employed a previously developed construct wherein a thymocyte-specific promoter drives the medaka full length *il7r* cDNA (Aghaallaei et al. 2021). We then performed mutation in the extracellular juxtamembrane-transmembrane region of the medaka *il7r* cDNA (Figure 5 - Supplement 1D) to develop a dominant active IL7R form, akin to the NPC mutation found in the *IL7R* gene in some T-ALL patients (Zenatti et al. 2011; Oliveira et al. 2022), hereafter called *il7r*^{NPC}. In our experimental setup, the promoter also co-expressed GFP, which allowed us to (i) identify thymocytes expressing the oncogenes, (ii) determine the clonal expansion of cells expressing the oncogene, and (iii) assess thymus hyperplasia and infiltration into other organs, a characteristic feature for T-ALL. DNA constructs were then injected into blastomeres of embryos at one-cell stage and they were observed during their development using live imaging, with a focus on two specific phenotypes: Firstly, we assessed whether the thymus exhibited enlargement beyond the normal size for its developmental stage. In particular, thymus lymphoma was defined as hyperplasia with an increase of more than twice the organ's typical size. Secondly, we identified T-ALL by observing infiltration of GFP-co-expressing malignant cells in other organs, including the brain, intestine, or heart (Figure 5D). To align our *in vivo* experiments with the *in silico* conditions, we limited our observations to 11 days post-fertilization (dpf), concluding the experiments at the freshly hatched yolk-sac larval stage. Since thymopoiesis begins at 3 dpf in medaka embryos (Bajoghli et al. 2015; 2009), this means we monitored thymus growth for a time span of 8 days.

None of the embryos injected with the NICD$^{\Delta PEST}$ construct (N=32) displayed thymus hyperplasia at 11 dpf (Figure 5E), supporting our simulation outcome showing that global activation of NOTCH1 signaling in thymocytes alone is insufficient to result in clonal expansion within a short time frame of 8 days. The MYC oncogene is an endogenous downstream target of NOTCH1 and plays a major role in NOTCH1-induced transformation (Sanchez-Martin and Ferrando 2017). Therefore, to attempt to shift the balance of NOTCH1 action towards proliferation, we decided to introduce the medaka *mycn* cDNA into our construct. Consequently, we observed that 25% of injected embryos with NICD$^{\Delta PEST}$ and *mycn* (N=36) displayed thymus hyperplasia at 11 dpf (Figure 5E). Among them, 8% also exhibited a massive infiltration of GFP-expressing cells in other organs such as the brain and gut. Further analysis of sorted GFP-expressing thymocytes revealed ectopic *il7* expression, while the expression level of endogenous *il7r* remained unchanged (Figure 5 -

Supplement 1B). Whole-mount in situ hybridization (WISH) analysis further
confirmed a robust upregulation of *il7* expression in the thymus of these
embryos (Figure 5 - Supplement 1C). MYCN is a basic helix-loop-helix
transcription factor that is downstream of several pro-proliferative signaling
pathways (Ruiz-Pérez, Henley, and Arsenian-Henriksson 2017), including
NOTCH1 in the context of thymocytes (Sanchez-Martin and Ferrando 2017)).
Together with our experimental results, it is therefore likely that MYCN could
enact the proliferative effect of NOTCH1, shifting the balance in the
incoherent feed-forward loop towards proliferation. Interestingly, our results
also indicate that transformed thymocytes acquire the ability to release IL-
7 (Figure 5 - Supplement 1B, C), which may further promote proliferation in
an autocrine fashion.

To test whether constitutive IL7R activation could enhance the
development of thymus hyperplasia and T-ALL, we injected various
constructs to overexpress either *il7r* or the dominant active *il7r*^{NPC} alone, or
in combination with NICD$^{\Delta PEST}$ and *mycn*. Thymus hyperplasia was found in
a slightly higher frequency of 26% (N=49) or 59% (N=29) of embryos,
respectively, when only *il7r* or *il7r*^{NPC} was overexpressed in thymocytes.
Additionally, we FACS-sorted thymocytes and performed a qPCR for lck,
since this gene was shown to be upregulated in zebrafish after IL7R^{DA} T-ALL
development using RNASeq data (Oliveira et al. 2022). Indeed, we detected
a significant upregulation of *lck* upon *il7r*^{NPC} overexpression when compared
to the WT counterparts. However, upregulation was not significantly
different when the WT was compared to *il7r*^{HI} condition (Figure 5 -
Supplement 1D). Combining *il7r*^{NPC} and NICD$^{\Delta PEST}$ and *mycn* yielded a
notably elevated frequency of ~73% (N=33) for thymus hyperplasia (Figure
5E). Of this group, 33% also exhibited a T-ALL phenotype, suggesting an
additive effect of these factors in the T-ALL development (Figure 5 -
Supplement Table 1). GFP and pH3 double staining of embryos exhibiting
the T-ALL phenotype further confirmed that the combination of NICD$^{\Delta PEST}$,
mycn and *il7r* overexpression in thymocytes resulted in a higher number of
mitotically active cells within the thymus (Figure 5F). Notably, we observed
many pH3-stained cells in the inner zone of the medaka thymus (Figure 5G,
arrows), an area where WT thymocytes are mitotically quiescent (Bajoghli
et al. 2015; Aghaallaei et al. 2021).

The prediction that higher NOTCH1 signaling does not lead to increased
proliferation is likely due to the incoherent feed-forward loop downstream
of NOTCH1, which shuts down excessive proliferation by triggering
differentiation to a non-dividing cell fate (Figure 5B). We observed a similar
effect in our *in vivo* experiments, where activation of NOTCH1 alone did not

lead to thymus hyperplasia within the observed time window of 8 days (Figure 5E). In contrast, combining NOTCH1 with its downstream pro-proliferative effector *mycn* led to thymus hyperplasia. Intriguingly, *in vivo*, continuous activation of NOTCH1 signaling in thymocytes is one of the main drivers of T cell leukemogenesis and T-ALL development (Weng et al. 2004; Lin et al. 2006; Chen et al. 2007; Liu et al. 2017; Neumann et al. 2014). Several *in vivo* studies have demonstrated that constitutive activation of NOTCH1 receptor leads to leukemia development after several weeks in zebrafish (Chen et al. 2007; Blackburn et al. 2012) or months in mice (Chiang et al. 2008; Hu et al. 2009; Sharma et al. 2006; Wendorff and Ferrando 2020). Indeed, we did not observe thymus hyperplasia when upregulating NOTCH1 signaling alone, which may stem from the shorter time window of our *in vivo* experiments (i.e., 8 days). However, in the virtual thymus model, a sole upregulation of NOTCH1 cannot produce thymus hyperplasia regardless of the duration. Different possibilities could reconcile the discrepancy between *in vivo* and the computational model. Firstly, the balance between pro-proliferative and pro-differentiation effects of NOTCH1 is different from what we modeled. Secondly, there is additional negative feedback between proliferation and differentiation, which our model does not consider. Thirdly, proliferation *in vivo* can continue even past differentiation if NOTCH1 signaling is sufficiently stimulated.

Overall, our *in vivo* results support the outcomes of our simulations, showing that overexpression of IL7R alone, but not NOTCH1 alone, is sufficient to induce thymus hyperplasia in a very short time period. However, dual activation of IL7R and the NOTCH1-MYCN axis can promote clonal expansion *in vivo*, leading to the rapid development of thymus hyperplasia and T-ALL in the medaka model system within eight days.

An abundance of thymic IL-7 milieu could accelerate the T-ALL development

One of the most unexpected outcomes of our simulations was that a defective TEC network provides lesioned clones with a substantial advantage in proliferation. Given the crucial role of IL-7 in proliferation of thymocytes, and considering that TECs, and not thymocytes, serve as the primary source of IL-7 cytokine in the normal thymic milieu, we next asked whether elevated *il7* expression in TECs alone is sufficient to induce thymus hyperplasia and to enhance T-ALL development in a thymus where thymocytes express high levels of *il7r*. This question was also driven by the results of scenario 27, which showed a twofold increase in thymocyte population size when all TECs express IL-7 (Figure 2E-G). In this thymic niche

Figure 6 | IL-7 supplied by the niche can lead to thymus hyperplasia. (A) Simulated data; rounds of cell division of normal and lesioned clones for different combinations of lesions and TECs expressing IL-7 ubiquitously. Note that some conditions are reproduced from Figure 5B for easier comparison. (B) Left panel: Frequency of thymus hyperplasia and T-ALL phenotype observed in the injected embryos at 11 dpf. For detailed statistics see Figure 5 - Supplement Table 1. Statistical data rounded to 4 decimal places. Note that some data from Figure 5E are repeated here to facilitate a better comparison. *N* represents the number of biological samples. Right panel: representative images at 10 dpf after co-injection of lck:gfp-NICD$^{\Delta PEST}$-*mycn* and ccl25a:tagRFP-il-7 constructs. Arrows indicate GFP-expressing malignant thymocytes in the thymus, brain, and gut. Scale bar is 400 μm (C) We propose a hypothetical negative feedback loop from proliferation to differentiation, which could explain discrepancy between *in silico* predictions and *in vivo* observations.

enriched with IL-7, lesioned clones either with IL-7^{HI} or extended differentiation times showed a significant advantage in clonal expansion compared to their WT counterparts (Figure 6A). Further, the impact of alterations in the thymic niche on T-ALL development has not been previously addressed. Therefore, we monitored the thymus development of the TEC:IL-7^{HI} embryos until 11 dpf to mimic scenario 27 *in vivo*. We found that 12% of these embryos (N=24) displayed signs of thymus hyperplasia (Figure 6B), however, we did not observe massive infiltration of GFP-expressing cells into other organs. The frequency of thymus hyperplasia was increased to 41% (N=32) when DNA constructs designed to overexpress *il7* in TECs were co-injected with the construct overexpressing *il7r* in thymocytes (Figure 6B). Comparably, thymus hyperplasia was only observed in 26% of embryos injected with the IL7R overexpression construct in thymocytes alone (N=49). A synergistic effect was observed when *il7* was overexpressed in TEC along with NICD and *mycn* in thymocytes (Figure 6B).

Interestingly, the frequency of embryos with T-ALL development was
increased to 25% (N=48) in this group. Taken together, our *in vivo* results
reveal that an excess of IL-7 in the thymic environment, combined with
alterations that affect the differentiation status of thymocytes, triggers the
T-ALL development in a short period of time.

Overall, several of our experimental observations indicate that there might
be additional negative feedback from pro-proliferative signaling to
differentiation (Figure 6C). Most strikingly, manipulations in IL7R or in
thymic IL-7 levels are sufficient to induce hyperplasia and T-ALL (Figure 5E
and Figure 6B) – an outcome that is much more severe than the model
predicts (Figure 5B and Figure 6A). In simulations, we could only obtain such
massive overproliferation by modifying multiple parameters, including
delaying differentiation, accelerating cell divisions, and reducing TEC
density (Figure 4 – Supplement 5), as differentiation in our model is a hard
stop to proliferation regardless of pro-proliferative stimuli. Similarly, the
experimental observation that NOTCH1-MYCN upregulation leads to
thymus hyperplasia and T-ALL development also suggests that proliferation
can proceed unbridled by differentiation *in vivo*. A negative feedback loop
from proliferation (or cell cycle entry) on differentiation could explain these
discrepancies between the virtual thymus model and *in vivo* observations.
We therefore propose that continued thymocyte proliferation can
downregulate differentiation pathways and maintain cells in a constant
proliferative state.

CONCLUSION

This study shows the powerful synergy between computational modeling
and *in vivo* experimental validation in elucidating the complex cellular and
molecular interactions within an organ that contribute to the initiation and
progression of diseases. Here, we highlight a previously underexplored area
– the role of TECs and the thymic niche in the development of T-ALL –
bridging a significant gap in our understanding of the initiation phase of the
disease. By simulating over 1500 scenarios, we identified potential drivers
such as alterations in the shape, density, and network of TECs that
significantly influence the clonal expansion of lesioned clones, aspects
which are challenging to manipulate in experimental setups. Consistent
with the outcome of our simulations, our *in vivo* results reveal that an
enriched thymic IL-7 milieu can accelerate disease progression, particularly
when thymocytes display enhanced IL7R levels or continuous NOTCH1
activation and MYCN overexpression. This challenges traditional gene-
centric views of the disease and highlights the importance of spatial

structure of the thymic niche for understanding the mechanisms of T-ALL
initiation and progression. It is important to note the limitations of our cell-
based computational model, calibrated to mimic the embryonic thymus of
medaka (Bajoghli et al. 2015; 2009; Aghaallaei et al. 2021), which might not
scale accurately to different species. Therefore, further experimental studies,
particularly using murine models, are essential to comprehensively
investigate the role of TECs in T-ALL development. Despite simplifying the
biological complexity inherent in modeling T-cell development, our
integrated approach significantly enhances our understanding of
environmental influences on disease progression. It also could open new
avenues for exploring how the environment contributes to
chemoresistance and disease relapse. This could lead to innovative
therapeutic strategies targeting the leukemic-supportive niche, offering
hope for improved treatment outcomes.

MATERIALS AND METHODS

In silico model

To develop, implement, and simulate the virtual thymus model, we used the
modeling and simulation software EPISIM (Sütterlin et al. 2017; 2013). This
multiscale simulation software uses an agent-based paradigm to represent
cells as spheres or ellipsoids in three-dimensional space, enabling each of
the cells to individually perform internal processes based on flow diagrams
and logic rules (e.g. if/else statements) or differential equations, and also
implements a partial differential equation solver to simulate diffusion of
chemicals in the extracellular space. The virtual thymus model has been
comprehensively described in the supplementary material to our previous
study (Aghaallaei et al. 2021). In this work, we used this model as a baseline
to introduce a small number of additions, explained in the following.

Rate of IL-7 pathway signal transduction

866 The equation describing the IL-7 signaling rate in each cell was rescaled to
867 introduce the parameter $a_{\text{IL-7}}$ for the IL-7 signal transduction activation rate.
868 The default value of $a_{\text{IL-7}}$ was chosen such that the model behavior did not
869 change. The rescaled equation for IL-7 signal transduction reads:

$$\frac{d}{dt}\sigma_{\text{IL-7}} = [\text{IL-7R}]\langle[\text{IL-7}_{\text{ex}}]\rangle a_{\text{IL-7}} - d_{\text{IL-7}}\sigma_{\text{IL-7}} \qquad \text{Eq. (1)}$$

where $\sigma_{\text{IL-7}}$ is the IL-7 pathway signal transduction activity, [IL7R] is the IL7R
concentration in the given cell, $\langle[\text{IL-7}_{\text{ex}}]\rangle$ is the mean extracellular IL-7

concentration in the cell's microenvironment, and $d_{\text{IL-7}}$ is the signal
transduction deactivation rate.

In the scenario where we introduced a lesioned clone with dominant-active
IL7R, IL-7 signal transduction activity was calculated as

$$\frac{d}{dt}\sigma_{\text{IL-7}} = [\text{IL-7R}] - d_{\text{IL-7}}\sigma_{\text{IL-7}}$$ Eq. (2)

876 Thus, in IL7R$^{\text{DA}}$ clones, neither extracellular IL-7 nor the signal transduction
877 activation rate $a_{\text{IL-7}}$ had an impact on cells' IL-7 signaling pathway activity.

Depletion of extracellular IL-7

879 In our original model (Aghaallaei et al. 2021), IL-7 ligand is released back to
880 the extracellular space after binding to IL7R. Thus, thymocytes do not affect
881 the extracellular IL-7 gradient, which is given by the partial differential
882 equation

$$\frac{\partial}{\partial t}[\text{IL-7}_{\text{ex}}] = D_{\text{IL-7}}\nabla^2[\text{IL-7}_{\text{ex}}] - k_{\text{IL-7}}[\text{IL-7}_{\text{ex}}] + \sum_i s_i$$ Eq. (3)

where $D_{\text{IL-7}}$ is the IL-7 diffusion constant, $k_{\text{IL-7}}$ is a baseline level of extracellular
IL-7 degradation, and s_i is a source term accounting for IL-7 secretion, e.g.
by a subset of TECs. In the modeling platform we use, secretion occurs
throughout a cell's volume; more specifically, for solving the partial
differential equation, space is discretized into voxels, and all voxels i that
intersect with the ellipsoids that are used to represent a cell contribute to
the source term.

In this work, we introduce the option, controlled via a Boolean flag *DEPL*
that is set by the user before the simulation starts, to simulate a scenario
where thymocytes internalize IL-7 ligand after it is bound to IL7R. In this
scenario, the extracellular IL-7 concentration is given by:

$$\frac{\partial}{\partial t}[\text{IL-7}_{\text{ex}}] = D_{\text{IL-7}}\nabla^2[\text{IL-7}_{\text{ex}}] - k_{\text{IL-7}}[\text{IL-7}_{\text{ex}}] + \sum_i s_i - \sum_j [\text{IL-7R}]_j\langle[\text{IL-7}_{\text{ex}}]\rangle_j a_{\text{IL-7}}$$ Eq. (4)

894 Note that the additional sink term in Equation 4 represents extracellular IL-
895 7 depletion and is simply the IL-7 signal transduction activity from Equation
896 1 summed over all voxels j that intersect with thymocytes. We assume that
897 all internalized IL-7 is permanently removed from the extracellular pool.

898 For lesioned IL7R$^{\text{DA}}$ clones, we used the alternative sink term

$$-\sum_k [\text{IL-7R}]_k$$

899 which is summed over all voxels k that intersect with thymocytes of the
900 lesioned clone. This term corresponds to the IL-7 signal transduction activity
901 as shown in Equation 2. Reduction of extracellular IL-7 below zero was
902 prevented by a computational check.

Determining clonal lineages

The EPISIM software we used assigns to each cell a unique cell identifier. We
made use of this unique cell identifier to generate unique clonal identifiers
as follows. For each thymocyte that entered the simulation via migration
from outside the thymus, we used the value of their unique cell identifier to
assign the value of a new clonal identifier variable. This cell-specific value of
the clonal identifier variable was passed on to any daughter cells produced
via cell division. Thus, using the clonal identifier, we essentially generated an
in silico clonal label which we used to determine clonally related cells.

Lesioned clone

To generate a lesioned clone in a given simulation, we created a
computational label to mark a single cell selected randomly among new
thymic immigrants once the thymic cell population reached its
homeostatic size (t >= 60h). This label marked this cell and its progeny as a
"lesioned clone", such that we could specifically modulate parameters and
introduce new rules only for this single clonal lineage. We tested the effect
of several alterations specific to the lesioned clone (Figure 4 - Supplement
Table 1).

Parameter variations

922 Please refer to the supplementary material of (Aghaallaei et al. 2021) for a
923 comprehensive explanation of all parameters and the choice of the
924 reference values. Altered parameters used in this work are listed in Figure 1
925 - Supplement Table 1.

Calculation of simulated clone size

Clone size was defined as the number of terminal leaves in a cell lineage tree
(see scheme in Figure 4C); terminal leaves were counted regardless of the
ultimate fate of those cells (i.e. positive or negative selection). All clones of
the same genotype (wildtype or lesioned) belonging to the same simulated

scenario were averaged across all replicate simulations of that scenario. This
calculation gives the average number of cells per clone in a given scenario.
The number of cells was log2-transformed to represent rounds of cell
division.

In vivo model

Medaka (*Oryzias latipes*) husbandry was performed in accordance with the
German animal welfare standards (Tierschutzgesetz §11, Abs. 1, Nr. 1,
husbandry permit no. 35/9185.46/Uni TÜ.) The transgenic line (tg) lck:gfp was
described previously (Bajoghli et al. 2015). All experiments conducted in
medaka embryos were performed prior to the legal onset of animal life
stages under protection, utilizing both males and females of the wildtype
line Cab (Loosli et al. 2000).

Cloning of DNA constructs

To overexpress *il7* in TECs, we used the construct ccl25a:*il7*, which was
described previously (Aghaallaei et al. 2021). To overexpress *il7r* in
thymocytes, the full-length medaka *il7r* cDNA (Aghaallaei et al. 2021) was
cloned into vectors containing a medaka thymocyte-specific promoter
(Bajoghli et al. 2015) that drives a fluorescent protein (sfGFP or TagRFP). To
generate a dominant active form of *il7r*, we introduced three amino acids,
Asparagine (N), Proline (P), and Cysteine (C) (known as NPC mutation, as
shown in (Zenatti et al. 2011) and (Oliveira et al. 2022)), in the medaka *il7r*
extracellular juxtamembrane-transmembrane interface region after
position 266 using site-directed mutagenesis (Figure 5 - Supplement Figure
1D), this corresponds to the human NPC mutation in the position 242 of *IL7R*
gene. To activate Notch signaling in thymocytes, we utilized the medaka
notch1b intracellular domain (NICD), as described previously (Aghaallaei et
al. 2021), and removed its PEST domain to enhance protein stability,
mimicking the situation reported in most T-ALL patients harboring gain-of-
function mutations in the NOTCH1 gene (Weng et al. 2004). For
overexpressing the *mycn* oncogene, we first isolated the full-length medaka
mycn cDNA (accession number: ENSORLG00000022362) and introduced a
mutation at position 44 from Proline to Leucine (P44L) using site-directed
mutagenesis, as identified in T-ALL patients (Liu et al. 2017). The DNA
fragments were then cloned into vectors containing a medaka thymocyte-
specific promoter, either alone or in combination (Bajoghli et al. 2015).

DNA micro-injection

Plasmids at concentration 10-25 ng/µl together with I-*Sce*I meganuclease and NEB buffer (NewEngland BioLabs), were co-injected into the blastomere at one-cell stage embryos. Fluorescent signals, based on the constructs either sfGFP, tagRFP or mTurquoise, were used to select positive embryos.

Generation and genotyping of medaka *il7* mutant

The CRISPR-Cas9 approach was employed to generate medaka *il7* crispant. Crispr RNA (crRNA AGTAGACTGATGCAAAGAAG) was designed using the CCTop website (Stemmer et al. 2015; 2017) and ordered from IDT. RNP complexes were prepared as described before (Hoshijima et al. 2019) using AltR tracrRNA (IDT) and Alt-R S.p. Cas9 nuclease (IDT). The injection mixture contained a total of 25 µM crRNA:tracrRNA duplex and 25 µM Cas9. Injection was performed into the blastomere at the one-cell stage transgenic embryos carrying the lck:gfp reporter. Injected embryos were raised until 8 dpf and then fixed in 4% PFA/2xPBS + 0.1% Tween20 for whole-mount immunostaining. To correlate the phenotype with the genotype, each embryo was genotyped using PCR and Sanger sequencing

Phenotype assessment

The development of injected embryos was monitored using a NIKON SMZ18 stereo-fluorescent microscope. Only embryos with fluorescent signals in the thymus region were selected for further analysis. Thymuses of embryos before hatching or freshly hatched yolk sac larvae were examined. Thymuses with at least a two-fold increase in size compared to transgenic embryos carrying the lck:gfp reporter construct (Bajoghli et al. 2015) were considered as having thymic hyperplasia. The identification of cells expressing fluorescent proteins and oncogenes outside the thymus, such as in the brain, gut, and heart, was considered indicative of the T-ALL phenotype.

Whole-mount immunostaining

Immunostaining procedures followed established protocols (Inoue and Wittbrodt 2011; Aghaallaei et al. 2021). Briefly, mitotically active cells were identified using a rabbit anti-phosphohistone-3 antibody (Ser10, Millipore 06–570, 1:500 dilution), with a Cy3-donkey anti-rabbit immunoglobulin G secondary antibody (the Jackson laboratories, 711-165-152; 1:500 dilution). GFP expression was detected using a goat Anti-GFP antibody (Abcam, ab5450, 1:500 dilution), with Alexa488-donkey anti-goat IgG (Abcam,

1003 ab15129, 1:600 dilution) as the secondary antibody. To quantify the number
1004 of pH3-expressing cells, the entire thymus region was imaged using a
1005 LSM710 (Zeiss) confocal microscope with z-stacks (z=1 μm).

Whole-mount *in situ* hybridization (WISH)

1007 Whole-mount *in situ* hybridization in medaka embryos was carried out
1008 using digoxigenin-labeled RNA probes, as previously described (Aghaallaei
1009 et al. 2005). Probes used in this study for *il7r* and *il7* were previously
1010 described (Aghaallaei et al. 2021).

Cell sorting and quantitative RT-PCR

Injected embryos were first smashed through a 40 μM strainer (Greiner) and
cells were collected in 0.9x PBS with 500 U.I. Heparinum Natricum
(Liquemin). Sorting was then performed using the Cell Sorter MA900 (Sony
Biotechnology). WT embryos were used as negative sorting control. Only
GFP positive cells were collected and used for RNA isolation. RNA of sorted
cells was isolated by using NucleoSpin RNA XS (Macherey-Nagel) and 20 ng
carrier RNA, following the manufacturer's protocol. RNA was treated with
rDNase and eluted in 14 μl dAqua. The first-strand cDNA synthesis was
carried out with random hexamer primers and SuperScript III Reverse
Transcriptase (Thermo Fisher Scientific) by following the manufacturer's
protocol. SYBR Green Kit (Applied Biosystems) was used for Quantitative
PCR on the LightCycler 480 (Roche). The data was evaluated in Microsoft
Excel using the ΔCt method and normalized to the housekeeping gene ef1a.
The primers used for RT-PCR have been described previously (Aghaallaei et
al. 2021).

Statistical analysis

Statistical analysis was performed using RStudio version 2024.4.2.764 (Posit
team 2024) using R version 4.4.0 (R Core Team 2024), the R library jtools
(Long 2022) and GraphPad Prism version 8.0.2. The two-tailed Fisher's exact
t-test was used to compare the thymus hyperplasia phenotype with normal
thymus. Welch's t-test was used for thymus volume size comparison in WT
and *il7* crispants, multiple t-test was used for comparison of normal thymus
with thymus hyperplasia and T-ALL phenotype. Mann-Whitney test was
used to evaluate the qPCRs. p-values <0.05 were considered as statistically
significant. The number of biological samples for each experiment (*N*) is
indicated in the figures, and bar graphs present the absolute numbers with
mean ± SD, ± SEM or the percentage as indicated in the figure legends.

Acknowledgment

The authors thank Anna-Sophia Hellmuth for assistance with *il7* gRNA injection; the institute of Medical Virology and Microbiology for continuous support in confocal microscopy; Larissa Doll for support and encouragement. Initial computational work presented here was performed using the ALICE computer resources provided by Leiden University, later work was performed using the computer lab in the Theoretical Biology and Bioinformatics department of Utrecht University. This work was supported by the Madeleine Schickedanz Kinderkrebsstiftung (grant number D.30.28666), and Deutsche Forschungsgemeinschaft (BA 5766/5-1). ET is funded by the Dutch Research Council (NWO) in the NWO Talent Programme with project number VI.Veni.222.323. We thank members of the Theoretical Biology and Bioinformatics division for their constructive feedback on the manuscript.

Author contributions

E.T. designed and developed the computational model. A.M.D. performed the *in vivo* experiments and analyzed their results. B.B. designed and supervised the study. All authors jointly interpreted the data and wrote the manuscript.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data needed to evaluate the conclusion in the paper are present in the paper and/or the Supplementary Materials. The computational model is available on Zenodo: doi:10.5281/zenodo.11656320 (Tsingos 2024). Additional materials related to this paper such as plasmids or DNA constructs may be requested from the authors. Some graphs were created with BioRender.com.

REFERENCES

Aghaallaei, Narges, Baubak Bajoghli, Ingrid Walter, and Thomas Czerny. 2005. "Duplicated Members of the Groucho/Tle Gene Family in Fish." *Developmental Dynamics* 234 (1): 143–50. https://doi.org/10.1002/dvdy.20510.

Aghaallaei, Narges, Advaita M. Dick, Erika Tsingos, Daigo Inoue, Eva Hasel, Thomas Thumberger, Atsushi Toyoda, Maria Leptin, Joachim Wittbrodt, and Baubak Bajoghli. 2021. "Αβ/Γδ T Cell Lineage Outcome Is Regulated by Intrathymic Cell Localization and Environmental Signals." *Science Advances* 7 (29): eabg3613. https://doi.org/10.1126/sciadv.abg3613.

Aghaallaei, Narges, Daigo Inoue, Eva Hasel de Carvalho, Advaita M. Dick, Joachim Wittbrodt, Maria Leptin, and Baubak Bajoghli. 2022. "Notch1 Deficiency Alters the Migratory Behavior of Developing T Cells and Calcium Signaling in the Thymus of Medaka." *European Journal of Immunology* 52 (2): 261–69. https://doi.org/10.1002/eji.202149512.

Bajoghli, Baubak, Narges Aghaallaei, Isabell Hess, Immanuel Rode, Nikolai Netuschil, Boon-Hui Tay, Byrappa Venkatesh, et al. 2009. "Evolution of Genetic Networks Underlying the Emergence of Thymopoiesis in Vertebrates." *Cell* 138 (1): 186–97. https://doi.org/10.1016/j.cell.2009.04.017.

Bajoghli, Baubak, Advaita M. Dick, Annisa Claasen, Larissa Doll, and Narges Aghaallaei. 2019. "Zebrafish and Medaka: Two Teleost Models of T-Cell and Thymic Development." *International Journal of Molecular Sciences* 20 (August):4179. https://doi.org/10.3390/ijms20174179.

Bajoghli, Baubak, Paola Kuri, Daigo Inoue, Narges Aghaallaei, Marleen Hanelt, Thomas Thumberger, Matteo Rauzi, Joachim Wittbrodt, and Maria Leptin. 2015. "Noninvasive In Toto Imaging of the Thymus Reveals Heterogeneous Migratory Behavior of Developing T Cells." *The Journal of Immunology* 195 (5): 2177–86. https://doi.org/10.4049/jimmunol.1500361.

Blackburn, J. S., S. Liu, D. M. Raiser, S. A. Martinez, H. Feng, N. D. Meeker, J. Gentry, et al. 2012. "Notch Signaling Expands a Pre-Malignant Pool of T-Cell Acute Lymphoblastic Leukemia Clones without Affecting Leukemia-Propagating Cell Frequency." *Leukemia* 26 (9): 2069–78. https://doi.org/10.1038/leu.2012.116.

Bloh, Kevin, Rohan Kanchana, Pawel Bialk, Kelly Banas, Zugui Zhang, Byung-Chun Yoo, and Eric B. Kmiec. 2021. "Deconvolution of Complex DNA Repair (DECODR): Establishing a Novel Deconvolution Algorithm for Comprehensive Analysis of CRISPR-Edited Sanger Sequencing Data." *The CRISPR Journal* 4 (1): 120–31. https://doi.org/10.1089/crispr.2020.0022.

Breit, Stephen, Martin Stanulla, Thomas Flohr, Martin Schrappe, Wolf-Dieter Ludwig, Gabriele Tolle, Margit Happich, Martina U. Muckenthaler, and Andreas E. Kulozik. 2006. "Activating NOTCH1 Mutations Predict Favorable Early Treatment Response and Long-Term Outcome in Childhood Precursor T-Cell Lymphoblastic Leukemia." *Blood* 108 (4): 1151–57. https://doi.org/10.1182/blood-2005-12-4956.

Buffière, Anne, Benjamin Uzan, Romain Aucagne, François Hermetet, Manon Mas, Sandra Nassurdine, Aziza Aznague, et al. 2019. "T-Cell Acute Lymphoblastic Leukemia Displays Autocrine Production of Interleukin-7." *Oncogene*, August. https://doi.org/10.1038/s41388-019-0921-4.

Chen, Huaibin, Jinghong Ma, Wanqing Li, Anna V. Eliseenkova, Chongfeng Xu, Thomas A. Neubert, W. Todd Miller, and Moosa Mohammadi. 2007. "A Molecular Brake in the Kinase Hinge Region Regulates the Activity of Receptor Tyrosine Kinases." *Molecular Cell* 27 (5): 717–30. https://doi.org/10.1016/j.molcel.2007.06.028.

Chiang, Mark Y., Lanwei Xu, Olga Shestova, Gavin Histen, Sarah L'Heureux, Candice Romany, M. Eden Childs, Phyllis A. Gimotty, Jon C. Aster, and Warren S. Pear. 2008. "Leukemia-Associated NOTCH1 Alleles Are Weak Tumor Initiators but Accelerate K-Ras–Initiated Leukemia." *Journal of Clinical Investigation* 118 (9): 3181–94. https://doi.org/10.1172/JCI35090.

De Keersmaecker, Kim, Zeynep Kalender Atak, Ning Li, Carmen Vicente, Stephanie Patchett, Tiziana Girardi, Valentina Gianfelici, et al. 2013. "Exome Sequencing Identifies Mutation in CNOT3 and Ribosomal Genes RPL5 and RPL10 in T-Cell Acute Lymphoblastic Leukemia." *Nature Genetics* 45 (2): 186–90. https://doi.org/10.1038/ng.2508.

Efroni, Sol, David Harel, and Irun R. Cohen. 2007. "Emergent Dynamics of Thymocyte Development and Lineage Determination." *PLOS Computational Biology* 3 (1): e13. https://doi.org/10.1371/journal.pcbi.0030013.

Ferrando, Adolfo A. 2009. "The Role of NOTCH1 Signaling in T-ALL." *Hematology* 2009 (1): 353–61. https://doi.org/10.1182/asheducation-2009.1.353.

Gameiro, Jacy, Patrícia Nagib, and Liana Verinaud. 2010. "The Thymus Microenvironment in Regulating Thymocyte Differentiation" 4 (3).

Girardi, Tiziana, Carmen Vicente, Jan Cools, and Kim De Keersmaecker. 2017. "The Genetics and Molecular Biology of T-ALL." *Blood* 129 (9): 1113–23. https://doi.org/10.1182/blood-2016-10-706465.

González-García, Sara, Marina García-Peydró, Enrique Martín-Gayo, Esteban Ballestar, Manel Esteller, Rafael Bornstein, José Luis de la Pompa, Adolfo A. Ferrando, and María L. Toribio. 2009. "CSL–MAML-Dependent Notch1 Signaling Controls T Lineage–Specific IL-7Rα Gene Expression in Early Human Thymopoiesis and Leukemia." *Journal of Experimental Medicine* 206 (4): 779–91.

Hess, Isabell, and Thomas Boehm. 2012. "Intravital Imaging of Thymopoiesis Reveals Dynamic Lympho-Epithelial Interactions." *Immunity* 36 (2): 298–309. https://doi.org/10.1016/j.immuni.2011.12.016.

Hoshijima, Kazuyuki, Michael J. Jurynec, Dana Klatt Shaw, Ashley M. Jacobi, Mark A. Behlke, and David Jonah Grunwald. 2019. "Highly Efficient CRISPR-Cas9-Based Methods for Generating Deletion Mutations and F0 Embryos That Lack Gene Function in Zebrafish." *Developmental Cell* 51 (5): 645-657.e4. https://doi.org/10.1016/j.devcel.2019.10.004.

Hu, Xiaoxia, Hongmei Shen, Chen Tian, Hui Yu, Guoguang Zheng, Richard XuFeng, Zhenyu Ju, Jing Xu, Jianmin Wang, and Tao Cheng. 2009. "Kinetics of Normal Hematopoietic Stem and Progenitor Cells in a Notch1-Induced Leukemia Model." *Blood* 114 (18): 3783–92. https://doi.org/10.1182/blood-2009-06-227843.

Inoue, Daigo, and Joachim Wittbrodt. 2011. "One for All—A Highly Efficient and Versatile Method for Fluorescent Immunostaining in Fish Embryos." *PLOS ONE* 6 (5): e19713. https://doi.org/10.1371/journal.pone.0019713.

Iwanami, Norimasa, Fernando Mateos, Isabell Hess, Nico Riffel, Cristian Soza-Ried, Michael Schorpp, and Thomas Boehm. 2011. "Genetic Evidence for an Evolutionarily Conserved Role of IL-7 Signaling in T Cell Development of Zebrafish." *The Journal of Immunology* 186 (12): 7060–66. https://doi.org/10.4049/jimmunol.1003907.

Ji, Zhiwei, Ke Yan, Wenyang Li, Haigen Hu, and Xiaoliang Zhu. 2017. "Mathematical and Computational Modeling in Complex Biological Systems." *BioMed Research International* 2017:1–16. https://doi.org/10.1155/2017/5958321.

King, Jasia, Kerbaï Saïd Eroumé, Roman Truckenmüller, Stefan Giselbrecht, Ann E. Cowan, Leslie Loew, and Aurélie Carlier. 2021. "Ten Steps to Investigate a Cellular System with

Mathematical Modeling." Edited by Patricia M. Palagi. *PLOS Computational Biology* 17 (5): e1008921. https://doi.org/10.1371/journal.pcbi.1008921.

Koch, Ute, Emma Fiorini, Rui Benedito, Valerie Besseyrias, Karin Schuster-Gossler, Michel Pierres, Nancy R. Manley, Antonio Duarte, H. Robson MacDonald, and Freddy Radtke. 2008. "Delta-like 4 Is the Essential, Nonredundant Ligand for Notch1 during Thymic T Cell Lineage Commitment." *Journal of Experimental Medicine* 205 (11): 2515–23. https://doi.org/10.1084/jem.20080829.

Lin, Ying-Wei, Rebecca A. Nichols, John J. Letterio, and Peter D. Aplan. 2006. "Notch1 Mutations Are Important for Leukemic Transformation in Murine Models of Precursor-T Leukemia/Lymphoma." *Blood* 107 (6): 2540–43. https://doi.org/10.1182/blood-2005-07-3013.

Liu, Ziqing, Olivia Chen, J. Blake Joseph Wall, Michael Zheng, Yang Zhou, Li Wang, Haley Ruth Vaseghi, Li Qian, and Jiandong Liu. 2017. "Systematic Comparison of 2A Peptides for Cloning Multi-Genes in a Polycistronic Vector." *Scientific Reports* 7 (1): 1–9. https://doi.org/10.1038/s41598-017-02460-2.

Long, JA. 2022. *Jtools: Analysis and Presentation of Social Scientific Data* (version 2.2.0). https://cran.r-project.org/package=jtools.

Loosli, Felix, Reinhard W Köster, Matthias Carl, Ronald Kühnlein, Thorsten Henrich, Manuela Mücke, Annette Krone, and Joachim Wittbrodt. 2000. "A Genetic Screen for Mutations Affecting Embryonic Development in Medaka Fish (*Oryzias Latipes*)." *Mechanisms of Development* 97 (1): 133–39. https://doi.org/10.1016/S0925-4773(00)00406-8.

Madeira, Fábio, Matt Pearce, Adrian R N Tivey, Prasad Basutkar, Joon Lee, Ossama Edbali, Nandana Madhusoodanan, Anton Kolesnikov, and Rodrigo Lopez. 2022. "Search and Sequence Analysis Tools Services from EMBL-EBI in 2022." *Nucleic Acids Research* 50 (W1): W276–79. https://doi.org/10.1093/nar/gkac240.

Metzcar, John, Yafei Wang, Randy Heiland, and Paul Macklin. 2019. "A Review of Cell-Based Computational Modeling in Cancer Biology." *JCO Clinical Cancer Informatics*, no. 3 (December), 1–13. https://doi.org/10.1200/CCI.18.00069.

Neumann, Martin, Sebastian Vosberg, Cornelia Schlee, Sandra Heesch, Stefan Schwartz, Nicola Gökbuget, Dieter Hoelzer, et al. 2014. "Mutational Spectrum of Adult T-ALL." *Oncotarget* 6 (5): 2754–66.

Oliveira, Mariana L., Padma Akkapeddi, Daniel Ribeiro, Alice Melão, and João T. Barata. 2019. "IL-7R-Mediated Signaling in T-Cell Acute Lymphoblastic Leukemia: An Update." *Advances in Biological Regulation* 71 (January):88–96. https://doi.org/10.1016/j.jbior.2018.09.012.

Oliveira, Mariana L., Alexandra Veloso, Elaine G. Garcia, Sowmya Iyer, Clara Pereira, Vasco M. Barreto, David M. Langenau, and João T. Barata. 2022. "Mutant IL7R Collaborates with MYC to Induce T-Cell Acute Lymphoblastic Leukemia." *Leukemia* 36 (6): 1533–40. https://doi.org/10.1038/s41375-022-01590-5.

Posit team. 2024. *RStudio: Integrated Development Environment for R.* (version 2024.4.2.764). Boston, MA.: Posit Software, PBC. http://www.posit.co/.

R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: Foundation for Statistical Computing. https://www.R-project.org.

Robert, Philippe A., Heike Kunze-Schumacher, Victor Greiff, and Andreas Krueger. 2021. "Modeling the Dynamics of T-Cell Development in the Thymus." *Entropy* 23 (4): 437. https://doi.org/10.3390/e23040437.

Ruijtenberg, Suzan, and Sander van den Heuvel. 2016. "Coordinating Cell Proliferation and
Differentiation: Antagonism between Cell Cycle Regulators and Cell Type-Specific Gene
Expression." *Cell Cycle (Georgetown, Tex.)* 15 (2): 196–212.
https://doi.org/10.1080/15384101.2015.1120925.

Ruiz-Pérez, María Victoria, Aine Brigette Henley, and Marie Arsenian-Henriksson. 2017. "The
MYCN Protein in Health and Disease." *Genes* 8 (4): 113. https://doi.org/10.3390/genes8040113.

Sanchez-Martin, Marta, and Adolfo Ferrando. 2017. "The NOTCH1-MYC Highway toward T-
Cell Acute Lymphoblastic Leukemia." *Blood* 129 (9): 1124–33. https://doi.org/10.1182/blood-
2016-09-692582.

Sharma, Vishva Mitra, Jennifer A. Calvo, Kyle M. Draheim, Leslie A. Cunningham, Nicole
Hermance, Levi Beverly, Veena Krishnamoorthy, Manoj Bhasin, Anthony J. Capobianco, and
Michelle A. Kelliher. 2006. "Notch1 Contributes to Mouse T-Cell Leukemia by Directly
Inducing the Expression of c -*Myc*." *Molecular and Cellular Biology* 26 (21): 8022–31.
https://doi.org/10.1128/MCB.01091-06.

Shochat, Chen, Noa Tal, Obul R. Bandapalli, Chiara Palmi, Ithamar Ganmore, Geertruy te
Kronnie, Gunnar Cario, et al. 2011. "Gain-of-Function Mutations in Interleukin-7 Receptor-α
(IL7R) in Childhood Acute Lymphoblastic Leukemias." *Journal of Experimental Medicine*
208 (5): 901–8. https://doi.org/10.1084/jem.20110580.

Silva, Ana, Afonso R. M. Almeida, Ana Cachucho, João L. Neto, Sofie Demeyer, Mafalda de
Matos, Thea Hogan, et al. 2021. "Overexpression of Wild-Type IL-7Rα Promotes T-Cell Acute
Lymphoblastic Leukemia/Lymphoma." *Blood* 138 (12): 1040–52.
https://doi.org/10.1182/blood.2019000553.

Souza-e-Silva, Hallan, Wilson Savino, Raúl A. Feijóo, and Ana Tereza Ribeiro Vasconcelos.
2009. "A Cellular Automata-Based Mathematical Model for Thymocyte Development."
PLOS ONE 4 (12): e8233. https://doi.org/10.1371/journal.pone.0008233.

Stemmer, Manuel, Thomas Thumberger, Maria Del Sol Keyer, Joachim Wittbrodt, and Juan
L. Mateo. 2015. "CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target Prediction
Tool." *PloS One* 10 (4): e0124633. https://doi.org/10.1371/journal.pone.0124633.

———. 2017. "Correction: CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target
Prediction Tool." *PloS One* 12 (4): e0176619. https://doi.org/10.1371/journal.pone.0176619.

Sütterlin, Thomas, Christoph Kolb, Hartmut Dickhaus, Dirk Jäger, and Niels Grabe. 2013.
"Bridging the Scales: Semantic Integration of Quantitative SBML in Graphical Multi-Cellular
Models and Simulations with EPISIM and COPASI." *Bioinformatics (Oxford, England)* 29 (2):
223–29. https://doi.org/10.1093/bioinformatics/bts659.

Sütterlin, Thomas, Erika Tsingos, Jalil Bensaci, Georgios N. Stamatas, and Niels Grabe. 2017.
"A 3D Self-Organizing Multicellular Epidermis Model of Barrier Formation and Hydration
with Realistic Cell Morphology Based on EPISIM." *Scientific Reports* 7 (1): 43472.
https://doi.org/10.1038/srep43472.

Takahama, Yousuke. 2006. "Journey through the Thymus: Stromal Guides for T-Cell
Development and Selection." *Nature Reviews Immunology* 6 (2): 127–35.
https://doi.org/10.1038/nri1781.

Terwilliger, T, and M Abdul-Hay. 2017. "Acute Lymphoblastic Leukemia: A Comprehensive
Review and 2017 Update." *Blood Cancer Journal* 7 (6): e577–e577.
https://doi.org/10.1038/bcj.2017.53.

Thomas-Vaslin, Véronique, Hester Korthals Altes, Rob J. de Boer, and David Klatzmann.
2008. "Comprehensive Assessment and Mathematical Modeling of T Cell Population

Dynamics and Homeostasis1." *The Journal of Immunology* 180 (4): 2240–50. https://doi.org/10.4049/jimmunol.180.4.2240.

Tsingos, E. 2024. *Virtual Thymus Model* (version 2.0). https://doi.org/10.5281/zenodo.11656320.

Vadillo, Eduardo, Elisa Dorantes-Acosta, Rosana Pelayo, and Michael Schnoor. 2018. "T Cell Acute Lymphoblastic Leukemia (T-ALL): New Insights into the Cellular Origins and Infiltration Mechanisms Common and Unique among Hematologic Malignancies." *Blood Reviews* 32 (1): 36–51. https://doi.org/10.1016/j.blre.2017.08.006.

Vibert, Julien, and Véronique Thomas-Vaslin. 2017. "Modelling T Cell Proliferation: Dynamics Heterogeneity Depending on Cell Differentiation, Age, and Genetic Background." *PLoS Computational Biology* 13 (3): e1005417. https://doi.org/10.1371/journal.pcbi.1005417.

Wendorff, Agnieszka, and Adolfo Ferrando. 2020. "Modeling NOTCH1 Driven T-Cell Acute Lymphoblastic Leukemia in Mice." *BIO-PROTOCOL* 10 (10). https://doi.org/10.21769/BioProtoc.3620.

Weng, Andrew P., Adolfo A. Ferrando, Woojoong Lee, John P. Morris, Lewis B. Silverman, Cheryll Sanchez-Irizarry, Stephen C. Blacklow, A. Thomas Look, and Jon C. Aster. 2004. "Activating Mutations of NOTCH1 in Human T Cell Acute Lymphoblastic Leukemia." *Science (New York, N.Y.)* 306 (5694): 269–71. https://doi.org/10.1126/science.1102160.

Yui, Mary A., and Ellen V. Rothenberg. 2014. "Developmental Gene Networks: A Triathlon on the Course to T Cell Identity." *Nature Reviews Immunology* 14 (8): 529–45. https://doi.org/10.1038/nri3702.

Zamisch, Monica, Billie Moore-Scott, Dong-ming Su, Philip J. Lucas, Nancy Manley, and Ellen R. Richie. 2005. "Ontogeny and Regulation of IL-7-Expressing Thymic Epithelial Cells." *The Journal of Immunology* 174 (1): 60–67. https://doi.org/10.4049/jimmunol.174.1.60.

Zenatti, Priscila P., Daniel Ribeiro, Wenqing Li, Linda Zuurbier, Milene C. Silva, Maddalena Paganin, Julia Tritapoe, et al. 2011. "Oncogenic IL7R Gain-of-Function Mutations in Childhood T-Cell Acute Lymphoblastic Leukemia." *Nature Genetics* 43 (10): 932–39. https://doi.org/10.1038/ng.924.

Figure 1 | The virtual thymus model enables spatiotemporally resolved clonal analysis. (A) Schematic overview highlighting 6 key steps during T-cell development, which are integrated into the virtual thymus model: (1) ETPs enter the thymus niche; (2) they receive Dll4 and IL-7 signals from the TECs. Triggered by these signals, (3) thymocytes undergo proliferation and (4) differentiation into two distinct T-cell sublineages (5). Finally, thymocytes undergo selection, either leaving the organ or undergoing cell death (6). (B-C) Experimentally measured ("Medaka") and computationally modeled ("Virtual") cell speed (B) and cell directionality (C) for ETPs and thymocytes. Bars are mean values, and error bars standard deviation. (D) Timeline of medaka embryonic development (top) compared to simulated time (bottom). As indicated by the cartoon at the bottom, the first ETP enters the thymus roughly at 2.5 dpf (t = 0 h in the simulation). At roughly 60 h (roughly 5 dpf), the thymus reaches its homeostatic cell population, with both cell entry and exit. (E) Representative snapshots of the simulation at homeostasis showing spatial expression patterns and signaling activity of components of IL7R and Notch1 signaling pathways in TECs (top two panels) and thymocytes (bottom 4 panels). Data are shown in arbitrary units normalized to a scale from 0 to 1. (F) Left: Cartoon explaining the asynchronous arrival (homing) of new cells into the organ, their clonal expansion, and asynchronous cell death or exit. Right: Visualization of clonal diversity in a typical simulation; each color shade uniquely indicates a clonal lineage. The height of each colored patch indicates the number of cells in that clone, the length indicates simulation time. For illustrative purposes, a black arrow indicates the homing of a sample of lineages. (G) Data from the right panel in (F) was normalized to time of homing. This visualization highlights developmental phases of thymocyte lineages. Gray arrowheads indicate rounds of cell division. Scale bar 50 cells. Abbreviations: ETP = early thymic progenitor, T = Thymocyte, *N* = number of replicates (biological or computational).

1297

1298　**Figure 1 - Supplement 1 | The virtual thymus represents a slice of the organ.** Three-dimensional renders of the
1299　simulation. (A) Showing only TECs in cyan. (B) Showing both thymocytes (bright green) and TECs (cyan). (C) Detail
1300　showing clonal thymocyte lineages. Each lineage was assigned a randomly selected color, TECs in dark gray. In
1301　homeostasis, most clones consist of 1-8 cells.

Figure_1_SuppMov_1.mp4

Figure 1 - Supplement Movie 1 | Simulation time-lapse. Three-dimensional renders of a typical simulation where each thymocyte lineage was assigned a randomly selected color. TECs are shown in dark gray. Simulation runs for 48000 steps, where 1 step is 15 seconds. Note how thymocytes can crawl over and under TECs.

Figure_1_SuppMov_2.mp4

Figure 1 - Supplement Movie 2 | Simulation time-lapse with fewer cells to highlight clonal dynamics. In this simulation, a total of eight ETPs were introduced into the simulation at the same time to highlight clonal expansion and motility within the organ (00:10-02:00s), and later cell exit and cell death (starting at 02:08s). Three-dimensional renders of a simulation where each thymocyte lineage was assigned a randomly selected color. TECs are shown in dark gray. Each simulation step is 15 seconds.

Figure 1 - Supplement Table 1 - Parameters varied in this work

For a full list of parameters and their values, please refer to Aghaallaei et al. 2021.

Symbol	Values tested (reference in bold)	Description
N	2 ; **3** ; 4	Number of concentric subdivisions in algorithm to generate TEC positions. The higher the value, the larger the number of TECs, thus the larger the TEC density.
p_{TEC}	3 ; **4** ; 5	Number of protrusions initialized for each TEC.
r_{TEC}	2.0 µm ; **2.5 µm** ; 3.0 µm	Radius of the main TEC body. Also scales the radius of the spheres in the TEC protrusions.
$DEPL$	True ; **False**	Boolean flag that sets if the simulation has thymocytes that internalize IL-7 according to Equation 4 in Materials and Methods (True) or if it defaults to Equation 3 in Materials and Methods (False).
$a_{IL\text{-}7}$	120 h^{-1} ; **240 h^{-1}** ; 480 h^{-1}	IL7R signaling activation rate
$d_{IL\text{-}7}$	25 h^{-1} ; **50 h^{-1}** ; 100 h^{-1}	IL7R signaling deactivation rate
θ_{prol}	1.3 ; **1.4** ; 1.5	Threshold level of IL-7 and Notch signaling activity required to progress through G1 phase and to commit to the cell cycle.
μ	5 h ; **7 h** ; 9 h	Mean cell cycle duration.
T_{diff}	15 h ; **24 h** ; 33 h	Minimum duration of the proliferative phase before terminal differentiation.

1312

1313

Figure 2 | TEC density influences thymocyte proliferation via IL7R signaling. (A) Left: Detail of confocal section highlighting the tight spatial interaction between TECs and thymocytes (T). Scale bar 5 μm. Right: Detail from a three-dimensional render of a simulation snapshot; TECs cyan, thymocytes green. (B) Parameter permutations tested. (C) Impact of parameter permutations on average homeostatic cell population. Bars indicate means, error bars standard deviation. (D) Clonal lineages normalized to time of thymic entry for reference (left), a scenario with low average population (middle) and a scenario with high average population (right). Scale bar: 100 cells. (E) Extracellular IL-7 gradient in simulations; sum projection of all z-planes of a simulated confocal stack. Units are arbitrary. (F) Mean extracellular IL-7 concentration in the entire simulated volume. Note that these values are lower than in (E) because of averaging over the volume, including empty space (black areas in (E)). (G) Thymocyte population size over time averaged over several simulation runs. Shaded area indicates standard deviation. (H) Illustration of experimental setup for IL-7^{KO} and TEC:IL-7^{HI}. At least 3 technical replicates were done for the injection of each construct. (I) Representative images of embryos stained with GFP (green) and pH3 (red) of WT, IL-7^{KO} and TEC:IL-7^{HI}. Scale bar 20μm. (J) Numbers of pH3 positive cells per thymus normalized to mean of WT. *N* represents the number of individual fish. Data show means ± standard deviation.

1328

Figure 2 - Supplement 1 | Statistical analysis of simulated data. (A) Three-dimensional rendering of TECs in the
simulation with the indicated parameter alterations. Orange: IL-7-secreting TECs. Cyan: Non-secreting TECs.
Dotted outline indicates thymus outline in the simulation. (B) Partial residual effect plot showing linear model fit
on log2-transformed thymic population size. Shaded area is 95% confidence interval. (C) Estimated coefficients
and p-values of the linear model. Significance codes: <0.001 "***"; <0.01 "**"; <0.05 "*". All parameters have a
statistically significant effect on cell number. Parameter interactions are also statistically significant, which
indicates that changing combinations has a greater effect than the sum of individual changes.

Figure 2 - Supplement 2 | Reduced thymus volume upon IL-7^KO. (A) Experimental design: gRNA targeting the signal peptide of il-7 was co-injected with Cas9 into the 1-cell-stage blastomere of Tg(lck:gfp) medaka. The embryo was raised until 8 dpf. Whole-mount immunostaining was performed using anti-GFP and M-phase marker phospho-histone 3 (pH3). The thymus was imaged using confocal microscopy, and embryos were genotyped by sequencing. (B) Schematic of the il-7 gene structure and binding site for the crRNA for CRISPR-Cas9 gene editing in the signal peptide region of exon 2. Zoom in: Representative sequencing data analyzed with decodr.org (Bloh et al. 2021) software. (C) Representative image of immunostaining of Tg(lck:gfp) in WT and IL-7^KO and evaluation of the thymus size using the area measurement tool in Imaris software (yellow). Scale bar 20 µm. (D) Quantification of thymus size: Thymus area in µm² in WT and IL-7^KO. N represents the number of biological samples. For statistics, unpaired t-test with Welch's correction was used; Data show means ± standard deviation. Significance threshold was set to p= 0.05. ** means p < 0.01.

1348

Figure 3 | High IL7R signaling promotes proliferation. (A) Illustration of scenarios without extracellular IL-7 depletion (left) and with extracellular IL-7 depletion (right). (B) Ordinary differential equation used to model the IL7R signaling activity. Parameters that were modified are highlighted. (C) Effect of IL7R signaling-related parameter permutation. Bars indicate means and error bars standard deviation. (D) Left: Mean extracellular IL-7 concentration in the entire simulated volume. Right: Extracellular IL-7 gradient in simulations; sum projection of all z-planes of a simulated confocal stack. Units are arbitrary. The introduction of IL-7 depletion by thymocytes leads to a slight reduction in extracellular IL-7 availability.

Figure 3 - Supplement 1 | Statistical analysis of simulated data. (A) Partial residual effect plot showing linear model fit on log2-transformed thymic population size. Shaded area is 95% confidence interval. (B) Estimated coefficients and p-values of the linear model. Significance codes: <0.001 "***"; <0.01 "**"; <0.05 "*". Although the population size is slightly reduced for IL-7 depletion, the effect is not statistically significant. In contrast, IL7R signaling activation and deactivation have a statistically significant effect. None of the interactions have statistically significant effects.

	Estimate	Std. Error	p-value	
(Intercept)	6.499e+00	1.877e-01	< 2e-16	***
Depletion	-6.803e-02	2.676e-01	0.800	
Deactivation	-1.838e-02	2.933e-03	1.33e-08	***
Activation	4.726e-03	6.110e-04	1.61e-11	***
Depletion : Deactivation	1.622e-03	4.152e-03	0.697	
Depletion : Activation	-2.709e-04	8.651e-04	0.755	
Deactivation : Activation	-1.005e-05	9.280e-06	0.282	
Depletion : Activation : Deactivation	-3.729e-06	1.313e-05	0.777	

1364

Figure 4 | Systematic screen shows that lower TEC density promotes malignant expansion. (A) Schematic
representation of the major lesions to IL7R that were implemented in the model. (B) Representative clonal plot
highlighting in black the lesioned clone. (C) We defined clone size as the maximum number of terminal leaves of
a lineage, regardless of eventual fate of each cell. Clone size distribution in the reference simulation (REF),
compared to IL7R^DA and IL7R^HI. REF: N=20 simulations, with 2088 clones total: IL7R^DA: N=13 simulations with 1330
non-lesioned clones and 13 lesioned clones. IL7R^HI: N=13 simulations with 1336 non-lesioned clones and 13 lesioned
clones. (D) Schematic of tested scenarios shown in (E). (E) Simulated permutations ordered along increasing mean
lesioned clone size for the IL7R^HI condition. Expected effect on proliferation is a qualitative estimate based on
preliminary simulations. A total of 1580 permutations were tested with a varying amount of replicates per
permutation. Error bars indicate standard deviation. Note that in (C) and (E) the axis is in base-2 logarithmic scale
to better illustrate rounds of cell division.

Figure 4 - Supplement Table 1 - Clonal lesions and their effects

Lesion	Explanation
IL7R^{WT}	The concentration of IL7R receptor in the lesioned clone was set to the maximum attainable in wildtype thymocytes. Thus, this lesion is representative of an extreme example among the non-lesioned population and was included as a control.
IL7R^{DA}	This concentration of IL7R receptor in the lesioned clone was set to the maximum attainable in wildtype thymocytes. Moreover, the receptor was constitutively active, irrespective of the presence of IL-7 ligand.
IL7R^{HI}	The concentration of IL7R receptor in the lesioned clone was set to 10-fold higher than the maximum attainable in wildtype thymocytes.
IL-7 autocrine	Lesioned clones secrete IL-7 at the same rate as TECs, acting as additional sources in Equation 4.
NOTCH1^{DA}	Lesioned clones had constitutively active Notch signaling irrespective of the presence of Delta ligand. The levels of Notch signaling were set to the maximum attainable in non-lesioned thymocytes and could not be reduced.
Delayed differentiation	The lesioned clone had its rate of differentiation halved, effectively doubling the duration of the proliferative phase.
Slower cell speed	The lesioned clone's maximum speed was half of the non-lesioned population's maximum speed.

1376

	A	B	C	D	E	F	G	H
1	grouping	genotype	primary_lesion	replicate	simulation_datafile	Cell_cycle_duration	Differentiation_duration	Cell_cycle_entry_th
2	A	T-ALL	IL-7R_10x_wt	1	2339227_12	7	24	
3	A	T-ALL	IL-7R_10x_wt	2	1745039_149	7	24	
4	A	T-ALL	IL-7R_10x_wt	3	1743691_95	7	24	
5	A	T-ALL	IL-7R_10x_wt	4	1744514_122	7	24	
6	A	T-ALL	IL-7R_10x_wt	5	1745085_167	7	24	
7	A	T-ALL	IL-7R_10x_wt	6	1743581_41	7	24	
8	A	T-ALL	IL-7R_10x_wt	7	1745186_203	7	24	
9	A	T-ALL	IL-7R_10x_wt	8	1742060_14	7	24	
10	A	T-ALL	IL-7R_10x_wt	9	1745129_185	7	24	
11	A	T-ALL	IL-7R_10x_wt	10	1743664_68	7	24	
12	A	T-ALL	IL-7R_10x_wt	11	1661851772_44	7	24	
13	A	T-ALL	IL-7R_10x_wt	12	1661831619_44	7	24	
14	A	T-ALL	IL-7R_10x_wt	13	1661784880_44	7	24	

Figure 4 – Source Data 1 | Source data file for visualization in Figure 4 E. In the data file, each row corresponds to the log2-transformed average clone size for one cell genotype (normal or lesioned) from one single independent replicate simulation. Additionally, the parameter values used in that run are listed in separate columns. The "grouping" column corresponds to the simulated scenarios (parameter permutation). The "simulation_datafile" column is the unique identifier of a given simulation run.

1383

Figure 4 - Supplement Movie 1 | Simulation time-lapse with lesioned clone. Two-dimensional projections of the virtual thymus volume showing time-lapse of simulations representing conditions in Figure 4B. Each thymocyte lineage was assigned a unique color; the lesioned clone was colored in black. Representative position of TECs shown in light blue (non-secreting) and light orange (IL-7-secreting). Movies were synchronized to the time of entry of the first thymocyte.

Figure 4 - Supplement 1 | Modulation of proliferation-related parameters affects lesioned thymocytes more strongly. (A) Clones with IL7R[HI] lesion. (B) Clones with IL7R[DA] lesion. (C) Control simulations without lesioned clones. Note that the y-axis of all plots is base-2 logarithmic to better illustrate rounds of cell division. Error bars indicate standard deviation. By modulating proliferation parameters, lesioned clones gain up to 6 rounds of division, while non-lesioned clones gain only 2 rounds of division. Expected effect on proliferation is a qualitative estimate based on preliminary simulations.

Figure 4 - Supplement Movie 2 | Simulation time-lapse. Two-dimensional projections of the virtual thymus volume showing time-lapse of simulations representing conditions in Figure 4 – Supplement 1A-C. Specifically, left column represents the condition with reduced proliferation, and the right column the condition with increased proliferation. Note how, with increased proliferation, newcomer cells struggle to enter deeper into the organ due to cell crowding. Each thymocyte lineage was assigned a unique color; the lesioned clone was colored in black. Representative position of TECs shown in light blue (non-secreting) and light orange (IL-7-secreting). Movies were synchronized to the time of entry of the first thymocyte.

Figure 4 - Supplement 2 | Modulation of parameters affecting IL7-signaling have almost no effect on lesioned clones. (A) Clones with IL7R^{HI} lesion. (B) Clones with IL7R^{DA} lesion. (C) Control simulations without lesioned clones. Note that the y-axis of all plots is base-2 logarithmic to better illustrate rounds of cell division. Error bars indicate standard deviation. By modulating IL-7 signaling parameters, lesioned clones do not gain extra rounds of division, while non-lesioned clones gain up to 3 rounds of division. Expected effect on proliferation is a qualitative estimate based on preliminary simulations.

Figure 4 - Supplement 3 | A secondary lesion delaying differentiation doubles proliferative potential. (A) Clones with IL7R[HI] lesion. (B) Clones with IL7R[DA] lesion. (C) Control simulations without lesioned clones. Note that the y-axis of all plots is base-2 logarithmic to better illustrate rounds of cell division. Error bars indicate standard deviation. Secondary clonal lesions lead to a gain of 4 rounds of division in the lesioned clones, with no effect on non-lesioned clones. Expected effect on proliferation is a qualitative estimate based on preliminary simulations. (D) Mean extracellular IL-7 concentration over time for selected simulations shown in (A)-(C). The increase in extracellular IL-7 by autocrine activity of lesioned clones is not enough to promote substantial proliferation of non-lesioned cells.

1421

**Figure 4 - Supplement 4 | Parameters affecting TEC distribution have opposing effects on lesioned and non-
lesioned clones.** (A) Clones with IL7R[HI] lesion. (B) Clones with IL7R[DA] lesion. (C) Control simulations without
lesioned clones. Note that the y-axis of all plots is base-2 logarithmic to better illustrate rounds of cell division. As
TEC architecture is made denser, lesioned clones lose up to 2 rounds of cell division, while non-lesioned clones
gain 2 rounds of cell division. Expected effect on proliferation is a qualitative estimate based on preliminary
simulations.

Figure 4 - Supplement 5 | Sparse TEC distribution is among permutations that most enabled lesioned clone expansion. Shown are top 8 conditions from Figure 4 (E). (A) Clones with IL7R^{HI} lesion. (B) Clones with IL7R^{DA} lesion. (C) Control simulations without lesioned clones. Note how lesioned clones with autocrine IL-7 production favor expansion of non-lesioned clones as well. The y-axis of plots in (A)-(C) is base-2 logarithmic to better illustrate rounds of cell division. Points are mean values, error bars indicate standard deviation. Expected effect on proliferation is a qualitative estimate based on preliminary simulations. (D) Volume occupied by thymocytes (1 thymocyte ≅ 33.5 µm^3) of a hypothetical lesioned clone compared to the volume of the virtual thymus (≅ 7854 µm^3). The y-axis is scaled in millions of cubic micrometers. The volumes strongly diverge after 12 rounds of cell division indicating extremely dense cell packing that would result in hyperplasia or metastasis *in vivo*.

Figure 5 | Interplay of IL7R and NOTCH1 potentiates clonal expansion *in vivo*. (A) Schematic of Notch1 and IL7R signaling and their regulation of thymocyte cell cycle entry, proliferation, and differentiation as implemented in the virtual thymus model. (B) Simulated data; rounds of cell division of normal and lesioned clones with different types of lesions in NOTCH1 or IL7R signaling. Note that the y-axis is base-2 logarithmic. Green boxes: Condition/lesion present. Gray boxes: Condition/lesion absent. (C) Illustration of plasmids for DNA micro-injection for transient overexpression of the different genes. Note that the expression of GFP was used to select successfully injected embryos. (D) Representative image: (left) of the WT control at 9 dpf where thymocytes are labeled in green with normal thymus size; (middle) of a thymus hyperplasia phenotype at 7 dpf; (right) same embryo developed T-ALL phenotype at 9 dpf, where infiltration in other organs was detected, namely in central nervous system (CNS) and gut. (E) Percentage of phenotypes: normal thymus, thymus hyperplasia and T-ALL at 11dpf. For each construct injection, we did at least three technical replicates. For detailed statistics see Figure 5 - Supplement Table 1. Statistical data rounded to 4 decimal places. (F) Number of pH3 positive cells per thymus counted during confocal imaging. (G) Representative confocal images of immunostaining against pH3 (red) and GFP (green). Scale bar is 20µm. *N* represents the number of biological samples.

1453

Figure 5 - Supplement 1 | Increased *il7* expression upon NICD^ΔPEST-mycn overexpression.
(A) Schematic depiction of NOTCH1 signaling (left) and overexpression of NICD (right). The engagement of NOTCH1 receptor with its ligand (DLL4) leads to its cleavage, and release and translocation of the NICD in the nucleus. NICD through interaction with other co-factors and triggers the expression of the target genes. Overexpression of NICD leads to activation of NOTCH signaling independent of the ligand. (B) Relative expression of *il7r* and *il7* of isolated thymocytes from WT compared to those expressing both NICD^ΔPEST and *mycn*. Each dot represents data from an individual sorted embryo. Data are $\log_{10}$ means ± standard error of the mean. Statistics: unpaired Mann-Whitney test, significance threshold was set to 0.05. (C) WISH with the il7 probe of freshly hatched yolk sac WT and NICD^ΔPEST and *mycn* larvae. Scale bare is 100µm. (D) Pairwise sequence alignment was done using EMBOSS Needle and the Needleman-Wunsch algorithm (Madeira et al. 2022), highlighting the partial medaka IL7R protein sequence and IL7R^NPC insertion mutation of amino acids NPC at position 266 in the extracellular juxtamembrane-transmembrane interface. (E) Relative *lck* expression quantified by qPCR of sorted WT thymocytes compared to those in IL7R^HI and IL7R^NPC. The *lck* expression was normalized to the housekeeping gene *ef1a*. Each dot represents an individual sorted embryo. Data are $\log_{10}$ means ± standard error of the mean. Statistics: unpaired Mann-Whitney test. Significance threshold was set to p=0.05. * means p < 0.05.

Figure 5 - Supplement Table 1: Detailed statistics for phenotype comparison using Fisher's exact test and pairwise comparisons; adjusted p values were calculated with the Benjamini & Hochberg method. Values reported in figures are the adjusted p values. Significance threshold was set to 0.05. Data rounded to 4 decimal places.

Group 1	Group 2	p.value	significant	p.value adjusted	significant adjusted
WT	lck:NICD	1	FALSE	1	FALSE
WT	ccl25a:il7	0.0486	TRUE	0.0694	FALSE
WT	lck:NICD-MycN	0.0007	TRUE	0.0017	TRUE
WT	ccr9a:il7r	0.0002	TRUE	0.0006	TRUE
WT	ccr9a:il7r + ccl25a:il7	4.90E-06	TRUE	2.20E-05	TRUE
WT	lck:NICD-MycN + ccl25a:il7	4.15E-08	TRUE	4.67E-07	TRUE
WT	ccr9a:il7r^{NPC}	8.70E-09	TRUE	1.31E-07	TRUE
WT	lck:NICD-MycN + ccr9a:il7r	2.70E-07	TRUE	2.03E-06	TRUE
WT	lck:NICD-MycN + ccr9a:il7r^{NPC}	3.00E-12	TRUE	1.47E-10	TRUE
lck:NICD	ccl25a:il7	0.0730	FALSE	0.0996	FALSE
lck:NICD	lck:NICD-MycN	0.0040	TRUE	0.0076	TRUE
lck:NICD	ccr9a:il7r	0.0011	TRUE	0.0024	TRUE
lck:NICD	ccr9a:il7r + ccl25a:il7	5.2885E-05	TRUE	0.0002	TRUE
lck:NICD	lck:NICD-MycN + ccl25a:il7	6.92E-07	TRUE	4.42E-06	TRUE
lck:NICD	ccr9a:il7r^{NPC}	9.67E-08	TRUE	8.70E-07	TRUE
lck:NICD	lck:NICD-MycN + ccr9a:il7r	1.59E-06	TRUE	7.93E-06	TRUE
lck:NICD	lck:NICD-MycN + ccr9a:il7r^{NPC}	1.24E-10	TRUE	2.78E-09	TRUE
ccl25a:il7	lck:NICD-MycN	0.4027	FALSE	0.4531	FALSE
ccl25a:il7	ccr9a:il7r	0.2344	FALSE	0.2776	FALSE
ccl25a:il7	ccr9a:il7r + ccl25a:il7	0.0351	TRUE	0.0526	FALSE
ccl25a:il7	lck:NICD-MycN + ccl25a:il7	0.0025	TRUE	0.0049	TRUE
ccl25a:il7	ccr9a:il7r^{NPC}	0.0008	TRUE	0.0018	TRUE
ccl25a:il7	lck:NICD-MycN + ccr9a:il7r	0.0007	TRUE	0.0017	TRUE
ccl25a:il7	lck:NICD-MycN + ccr9a:il7r^{NPC}	8.65E-06	TRUE	3.54E-05	TRUE
lck:NICD-MycN	ccr9a:il7r	0.0872	FALSE	0.1155	FALSE
lck:NICD-MycN	ccr9a:il7r + ccl25a:il7	0.1314	FALSE	0.1642	FALSE
lck:NICD-MycN	lck:NICD-MycN + ccl25a:il7	0.0493	TRUE	0.0694	FALSE
lck:NICD-MycN	ccr9a:il7r^{NPC}	0.0170	TRUE	0.0283	TRUE
lck:NICD-MycN	lck:NICD-MycN + ccr9a:il7r	0.0281	TRUE	0.0436	TRUE
lck:NICD-MycN	lck:NICD-MycN + ccr9a:il7r^{NPC}	0.0003	TRUE	0.0007	TRUE
ccr9a:il7r	ccr9a:il7r + ccl25a:il7	0.2187	FALSE	0.2660	FALSE
ccr9a:il7r	lck:NICD-MycN + ccl25a:il7	0.0003	TRUE	0.0007	TRUE
ccr9a:il7r	ccr9a:il7r^{NPC}	6.7619E-05	TRUE	0.0002	TRUE
ccr9a:il7r	lck:NICD-MycN + ccr9a:il7r	8.0909E-05	TRUE	0.0003	TRUE
ccr9a:il7r	lck:NICD-MycN + ccr9a:il7r^{NPC}	7.85E-07	TRUE	4.42E-06	TRUE
ccr9a:il7r + ccl25a:il7	lck:NICD-MycN + ccl25a:il7	0.0258	TRUE	0.0415	TRUE
ccr9a:il7r + ccl25a:il7	ccr9a:il7r^{NPC}	0.0135	TRUE	0.0234	TRUE
ccr9a:il7r + ccl25a:il7	lck:NICD-MycN + ccr9a:il7r	0.0078	TRUE	0.0141	TRUE
ccr9a:il7r + ccl25a:il7	lck:NICD-MycN + ccr9a:il7r^{NPC}	0.0016	TRUE	0.0033	TRUE
lck:NICD-MycN + ccl25a:il7	ccr9a:il7rNPC	0.7902	FALSE	0.8269	FALSE
lck:NICD-MycN + ccl25a:il7	lck:NICD-MycN + ccr9a:il7r	0.6492	FALSE	0.6955	FALSE
lck:NICD-MycN + ccl25a:il7	lck:NICD-MycN + ccr9a:il7r^{NPC}	0.1243	FALSE	0.1598	FALSE
ccr9a:il7r^{NPC}	lck:NICD-MycN + ccr9a:il7r	0.9308	FALSE	0.9519	FALSE
ccr9a:il7r^{NPC}	lck:NICD-MycN + ccr9a:il7r^{NPC}	0.4821	FALSE	0.5291	FALSE
lck:NICD-MycN + ccr9a:il7r	lck:NICD-MycN + ccr9a:il7r^{NPC}	0.3705	FALSE	0.4275	FALSE

1472

Figure 6 | IL-7 supplied by the niche can lead to thymus hyperplasia. (A) Simulated data; rounds of cell division of normal and lesioned clones for different combinations of lesions and TECs expressing IL-7 ubiquitously. Note that some conditions are reproduced from Figure 5B for easier comparison. (B) Left panel: Frequency of thymus hyperplasia and T-ALL phenotype observed in the injected embryos at 11 dpf. For detailed statistics see Figure 5 - Supplement Table 1. Statistical data rounded to 4 decimal places. Note that some data from Figure 5E are repeated here to facilitate a better comparison. *N* represents the number of biological samples. Right panel: representative images at 10 dpf after co-injection of lck:gfp-NICD$^{\Delta PEST}$-*mycn* and ccl25a:tagRFP-il-7 constructs. Arrows indicate GFP-expressing malignant thymocytes in the thymus, brain, and gut. Scale bar is 400 μm (C) We propose a hypothetical negative feedback loop from proliferation to differentiation, which could explain discrepancy between *in silico* predictions and *in vivo* observations.

A

TECs
T

B

	LOW	REF	HIGH
Size (S)			
Number of protrusions (P)	3	4	5
Density (D)			

C

Scenario	S	P	D
Reference	•	•	•
1	↖	↖	•
2	↖	↖	↖
3	•	•	↖
4	↖	↗	↖
5	↖	↗	•
6	↖	↗	↖
7	↖	•	↖
8	↖	•	•
9	↖	↗	•
10	•	↖	•
11	•	↗	↖
12	↖	•	•
13	•	↗	•
14	↖	↗	↗
15	↗	↖	↖
16	↗	•	↖
17	↗	↖	↖
18	↗	•	•
19	↗	↗	•
20	↗	•	•
21	↗	↗	•
22	↗	•	•
23	↗	↗	↖
24	↗	↗	•
25	↗	•	•
26	↗	•	•

Population size: 0 100 200 300

D

Reference   Scenario 6   Scenario 26

Time since thymus entry (h)

E

Reference

Scenario 6

Scenario 26

All TECs express IL-7
(Scenario 27)

No TECs express IL-7
(Scenario 28)

Extracellular IL-7 level: 1400 1200 1000 800 600 400 200 0

F

Mean extracellular IL-7 (A.U.)

Simulation time (h)

Reference — Scenario 6
Scenario 26 — Scenario 27
Scenario 28

G

Thymocytes (#)

Simulation time (h)

H

Medaka embryonic thymus
il7+
il7neg

Injection of il7 gRNA+Cas9 → IL-7KO
il7neg

Injection of ccl25a:il7 → TEC:IL-7HI
il7high

I

WT   IL-7KO   TEC:IL-7HI

J

fold change of pH3 cells in the thymus

p<0.0001
p=0.0008

WT   IL-7KO   TEC:IL-7HI

N= 18   9   3

A

gRNA+Cas9 co-injection → Tg(lck:gfp) 8dpf → whole-mount immunostaining → confocal imaging → sequencing

B

primer · UTR · coding region

signal peptide

crRNA

362 bp + 3bp

il-7 · Exon 1 · ATG · Intron 1 · Exon 2 · Intron 2 · Exon 3 · Intron 3 · Exon 4 · Intron 4 · Exon 5

365 bp · 126bp · 90bp · 123bp · 153bp

INDEL —	% —	crRNA
G1 -4	54.1	TTGGCTGATATTTTTGTTTTTCTTCTTTTCCCCACATTGTGACTAAACTTCCAGCC- - - -|CTTTGCATCA
G1 -19	45.9	TTGGCTGATATTTTTGTTTTTCTTCTTTTCCCCACATTGTG- - - - - - - - - - - - - - - -|CTTTGCATCA

Intron 1 · Exon 2

C

WT · IL-7^KO

D

thymus area in µm²

**

WT · IL-7^KO

N= · 8 · 7

A

IL-7
Depletion (D)

off · | on ✓

B

$$\frac{d\sigma_{\text{IL-7}}}{dt} = [\text{IL-7R}] \langle [\text{IL-7}]_{\text{ex}} \rangle \boxed{a_{\text{IL-7}}} - \boxed{d_{\text{IL-7}}} \sigma_{\text{IL-7}}$$

signaling
activation rate
(SA)

signaling
deactivation rate
(SD)

C

Scenario	D	SD	SA
Reference	·	·	·
29	·	↗	↗
30	✓	↗	↗
31	✓	↗	↗
32	·	·	↗
33	✓	·	↗
34	·	↗	·
35	✓	↗	↗
36	✓	·	↗
37	·	·	↗
38	✓	↗	↗
39	·	↗	↗
40	✓	↗	↗
41	✓	↗	·
42	·	↗	·
43	·	·	↗
44	✓	↗	↗
45	·	↗	↗

Population size

D

Reference

N=19

Scenario 36

N=12

Scenario 44

N=4

Scenario 45

N=4

Mean extracellular IL-7 (A.U.)

Simulation time (h)

Extracellular
IL-7 level

A

| | IL-7 depletion | signaling deactivation rate | signaling activation rate |

B

	Estimate	Std. Error	p-value	
(Intercept)	6.499e+00	1.877e-01	< 2e-16	***
Depletion	-6.803e-02	2.676e-01	0.800	
Deactivation	-1.838e-02	2.933e-03	1.33e-08	***
Activation	4.726e-03	6.110e-04	1.61e-11	***
Depletion : Deactivation	1.622e-03	4.152e-03	0.697	
Depletion : Activation	-2.709e-04	8.651e-04	0.755	
Deactivation : Activation	-1.005e-05	9.280e-06	0.282	
Depletion : Activation : Deactivation	-3.729e-06	1.313e-05	0.777	

A

Virtual thymus

TEC — DLL-4 — IL-7

NOTCH1 — IL7R

Thymocyte

Cell cycle entry

Differentiation — Proliferation

B

Clone genotype

Lesioned

Normal

NOTCH^DA longer diff.													
IL7R^WT													
IL7R^HI													
IL7R^DA													

C

gfp-NICD^ΔPEST : gfp | *nicd*^ΔPEST | pA

gfp-NICD^ΔPEST-*mycn* : gfp | *nicd*^ΔPEST | *mycn* | pA

gfp-*il7r* : gfp | *il7r* | pA

gfp-*il7r*^NPC : gfp | *il7r*^ΔNPC | pA

D

WT — thymus, eye, yolk — 9 dpf

Thymus lymphoma — thymus, eye, yolk, gut — 7 dpf

T-ALL — thymus, CNS, eye, yolk, gut — 9 dpf

E

Normal thymus
Thymus hyperplasia
Thymus hyperplasia+ T-ALL

p=0.0017
p=0.0283
p=0.0007
p=0.0002
p=0.4275

Phenotype (%)

N=	40	32	36	49	29	19	33
GFP	+	-	-	-	-	-	-
NICD^ΔPEST	-	+	-	-	-	-	-
NICD^ΔPEST-*mycn*	-	-	+	-	-	+	+
IL7R	-	-	-	+	-	+	-
IL7R^NPC	-	-	-	-	+	-	+

F

p=0.003
p=0.003
p=0.008
p=0.03
p=0.003

pH3+ cells per thymus (#)

N=	10	3	3	3	4	5
GFP	+	-	-	-	-	-
NICD^ΔPEST-*mycn*	-	+	-	-	+	+
IL7R	-	-	+	-	+	-
IL7R^NPC	-	-	-	+	-	+

G

WT — *il7r* — *il7r*^NPC — NICD^ΔPEST+*mycn* — NICD^ΔPEST+*mycn*+*il7r* — NICD^ΔPEST+*mycn*+*il7r*^NPC

A

NOTCH1 signaling

TEC

NOTCH1

DLL4

Thymocyte

cleavage

NICD

NICD

Signaling

Signaling

NICD$^{\Delta PEST}$

NICD

Signaling

B

relative il7r expression [log10]

ns

N= 4   5

Thymocytes   NICD$^{\Delta PEST}$ and *mycn*

relative il7 expression [log10]

*

N= 4   5

Thymocytes   NICD$^{\Delta PEST}$ and *mycn*

C

WT   NICD$^{\Delta PEST}$-*mycn*

il7

eye   thymus

eye

E

relative lck expression [log10]

*

p=0.11   ns

N= 3   4   5

Thymocytes   IL7R^{HI} IL7R^{NPC}

D

```
IL7R      251 WVLPADPPVVKMEHIL---TVCLMVVVVVGVTSSAFIFWKKQ    289
              ||||||||||||||||   |||||||||||||||||||||
IL7R^NPC  251 WVLPADPPVVKMEHILNPCTVCLMVVVVVGVTSSAFIFWKKQ    292
                              ___
```

A

Clone genotype
● Lesioned
● Normal

TEC: IL-7^{HI}												
longer diff.		■	■			■	■			■	■	
IL7R^{WT}		■	■			■	■					
IL7R^{HI}					■	■	■					
IL7R^{DA}									■	■	■	■

B

■ Normal thymus
■ Thymus lymphoma
■ Thymus lymphoma+ T-ALL

N=	40	24	36	48	49	32
Control	+	-	-	-	-	-
ccl25a:*il7*	-	+	-	+	-	+
lck:NICD^{ΔPEST}-*mycn*	-	-	+	+	-	-
ccr9a:*il7r*	-	-	-	-	+	+

$p=0.0694$ $p=0.0049$ $p=0.0694$ ns

C

lck: gfp-NICD^{ΔPEST}-*mycn*

ccl25a:rfp-*il7*