Predictions of bimanual self-touch determine the temporal tuning of somatosensory perception

Noa Cemeljic[1], Xavier Job[1], Konstantina Kilteni[1,2*]

[1]Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden

[2] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

*Correspondence and requests for materials should be addressed to konstantina.kilteni@ki.se

Abstract

We effortlessly distinguish between touching ourselves with our hands and being touched by other people or objects. Motor control theories posit that this distinction is made possible by the brain predicting the somatosensory consequences of our voluntary movements based on an 'efference copy', and attenuating our responses to the predicted self-touch. However, it remains unclear how these predictions impact somatosensory perception at times other than during self-touch: for example, as our hand reaches to touch our body or moves away from it. Here participants discriminated forces applied on their passive left index finger. The forces were applied during the reaching movement of their right hand towards the left hand, including the time the reaching ended by simulating self-touch between the hands, or after the reaching movement. We observed that the forces on the left hand felt progressively weaker during the reaching phase, reached their minimum perceived intensity at the time of self-touch, and quickly recovered after the end of the reaching. All effects were replicated with a new cohort of participants that further demonstrated that this gradual attenuation of the perceived magnitude of touch vanished during similar right hand reaching movements that did not produce expectations for self-touch between the two hands. Together, our results indicate a temporal tuning of somatosensory perception during movements to self-touch and underscore the role of sensorimotor context in forming predictions that attenuate the intensity of self-generated touch.

Introduction

Self-touch occurs in a wide range of everyday activities, from rubbing our eyes with our fists to alleviate itching, to warming ourselves with our hands during shivering, and affectively stimulating our bodies for pleasure. Self-touch behaviors begin in utero (1,2) and continue after birth as a way for infants to explore their bodies (3–5). Whether spontaneous or purposeful (6), self-touch is considered fundamental for building and maintaining a sense of self through detecting contingencies between motor and sensory cues (7–10).

Besides self-touch, we also experience numerous instances of touch generated by other people or objects. Remarkably, our brain possesses the capacity to distinguish between self-touch and touch of external origin to guide behaviour; for example, we do not react to self-touch between our hands, but we might recoil from similar touch initiated by a stranger, just as we would brush away a fly that lands our hand.

How does the brain implement this distinction? Motor control theories posit that the brain uses internal models to predict the somatosensory consequences of voluntary movements (11–15) based on a copy of the motor command (*i.e.,* efference copy) and attenuates (or cancels) the responses to the self-generated touch (16–21). Accordingly, when reaching with the (active) right hand to touch the (passive) left hand, the internal model predicts and attenuates the resulting touch on both hands using the efference copy associated with the right hand's reaching movement. In contrast, a touch of external origin cannot be predicted based on the movement and is not attenuated (17,20). It has been proposed that this attenuation of self-generated touch facilitates the distinction from externally generated touch by increasing the salience of the latter relative to the former (15,16,20). In agreement with this framework, several studies have shown that self-touch elicits an attenuated perceptual response (*i.e.*, it feels less intense or ticklish) (22–39) and triggers weaker activity in somatosensory cortices and the cerebellum (40–46) than identical touch applied by another person or a machine.

Nevertheless, it remains unclear how the predictions of the internal model about the somatosensory consequences of the movement affect perception at times other than the exact moment of self-touch. For example, imagine reaching with the right hand to touch your left hand (**Figure 1a**). It remains elusive how touch on the left hand is perceived as the reaching movement of the right hand progresses, before contact between the hands occurs, and after the reaching movement has ended. This is due to several reasons. First, several earlier studies assessed somatosensory perception across time on the active hand, rather than the passive hand which served as a target of the active hand's movement (*e.g.*, (47–49)). However, it has been demonstrated that somatosensory perception on a limb that moves can be 'gated' by nonpredictive mechanisms (50–56) (see (35) for discussion). To isolate the effects of the internal model's predictions on somatosensation, other studies have focused on the perception of the passive hand, which is not influenced by non-predictive confounds because it does not move (16,17,35). Nevertheless, these studies either probed perception only *at the moment* of self-touch (22–24,28–31,33–39,57), overlooking the periods before or after the hands' contact, or otherwise focused on the time *before* the contact (58), overlooking the periods during or after self-touch. Third, other studies have introduced temporal delays to test somatosensory

perception before and/or after the predicted self-touch (26,27,32,46,59), but those probed
timings were tailored to the moment of the predicted tactile sensation and did not directly relate
to the reaching movement of the active hand. For example, one can probe somatosensory
perception at a fixed time before the anticipated self-touch, but that timing could align with
different phases (early or late) of the active hand's reaching depending on its duration. Finally,
in addition to being constrained in scope, the abovementioned research has produced
conflicting and inconclusive results. Specifically, an earlier study reported a constant increase
in the attenuation of touch on the passive hand as the active hand approached the passive (27),
while another study found facilitation, rather than attenuation, of tactile perception on the
passive hand when the active hand moved towards the passive (58).

Here, we hypothesized that the predictions of the internal model should continuously influence
somatosensory perception during the reaching movement of the right hand to touch the left
hand. Specifically, we reasoned that as the reaching movement of the right hand progresses,
the brain gains access to the latest sensory information to adjust the movement and re-estimate
the position and velocity of the right hand (15,19,60), as well as to update the predictions of
the internal model about the remainder of the movement (61–63), including the expected timing
of contact with the left hand (*i.e.*, self-touch). Consequently, the prediction about the time of
self-touch should become sharper (*i.e.* narrower probability distribution) at later timings during
the reaching than earlier ones (**Figure 1b**). Moreover, a touch received on the left hand at times
further away from the expected contact time with the right hand would elicit a larger prediction
error, compared to a touch received closer to the expected time (**Figure 1c**, arrows) (19,20,64–
67). Therefore, we expected to observe a gradually increasing attenuation of the perceived
magnitude of touch during the reaching, which recovers after the hands make contact (**Figure
1c**). That is, the attenuation of somatosensory perception should not be an on-off phenomenon
that manifests only at the time of the self-touch, but rather it should gradually build up over the
reaching time.

Figure 1. Schematic illustrations of the experimental hypothesis. (a) We investigated how participants perceive touch on their left hand during and after reaching with their right hand to touch their left hand. The reaching movement is shown from a third person perspective and the top right small window indicates the orientation of the participant. **(b)** From a theoretical perspective, during the reaching the brain estimates the state of the moving hand (*e.g.*, its position, velocity) using the predicted and received sensory input and updates its predictions about the rest of the movement accordingly. Consequently, as the action progresses, the predictions of the internal model about the timing of self-touch become narrower given that there is a better estimation based on the latest sensory input (*e.g.*, visual and proprioceptive information). **(c)** Touch received during the reaching (yellow, red, and blue) including its conclusion (green), or after the reaching (brown) would be attenuated inversely proportionally to the error between the predicted and the received time (arrows). The shape of the displayed curve depicts a schematic-only illustration of how somatosensory perception could change as a function of time, and it does not restrict the hypothesis to a specific type of function. **(a, b, c)** Phases of the reaching movement and times are colour-coded. Yellow represents the early reaching phase (acceleration), red indicates the phase around the peak velocity, blue denotes the late reaching phase (deceleration), green corresponds to the moment the reaching movement concludes upon reaching the target (self-touch), and brown refers to the phase after the reaching.

Results

Experiment 1. The perceived magnitude of touch gradually decreases during the reaching movement, attaining its minimum at the moment of the predicted self-touch.

Twenty-nine participants (15 female, 28 right-handed, aged 18 - 34 years) rested their left hand palm up with their left index finger placed inside a moulded support on a desk. Their right hand was placed palm down on top of a wooden surface placed on top of the desk (**Figure 2a**). Upon presentation of an auditory GO cue, they performed a reaching movement with their active right hand starting from a designated starting position (**Figure 2a**, *starting position*), and concluded the movement by tapping with their right index finger a force sensor placed above, but not in contact with, their passive left index finger (**Figure 2a**, *cyan force sensor*).

During or after the reaching movement, the participants received a brief force on their left index finger via a probe attached to the lever of a DC electric motor (**Figure 2a**, *lever*). This *test* force had a fixed amplitude of 2 N. We quantified how strongly participants perceived the intensity of this *test* force over time using a well-established force discrimination task (16,27–35,46). In each trial, participants received the *test* force (2 N) and a *comparison* force with an intensity that pseudorandomly varied around the intensity of the *test* force (between 1 and 3 N) (**Figure 2b**). The magnitudes of the *test* and *comparison* forces were confirmed by a second identical force sensor placed inside the probe (**Figure 2a**, *red force sensor*). As mentioned above, the *test* force was applied at a different time relative to the reaching movement of the right hand on any given trial. However, the *comparison* force was always applied 1200 ms after the *test* force – that is, when the right hand had finished the reaching and stopped moving. In other words, our experimental manipulation exclusively involved the *test* force, while the

comparison force served as a reference. After the presentation of the two forces, participants
verbally indicated which one of the *test* or the *comparison* force felt stronger (**Figure 2b**).

The participants performed a block of 560 reaching trials in which the *test* force was delivered
during one of five different phases relative to the movement of their right hand in randomized
order (**Figure 2c**): (a) the *early* reaching trials included the period between movement onset
until the time the velocity reached 85% of its peak value (early acceleration phase); (b) the *mid*
reaching trials included the period after the velocity reached 85% of its peak value until the
time it dropped below 85% of its peak value (peak velocity phase); (c) the *late* reaching trials
included the period after the velocity dropped below 85% of its peak value until before the
moment the right index finger tapped the force sensor (late deceleration phase); (d) the *target*
trials included the moment when the right index finger tapped the sensor and triggered the
touch on the left hand, thereby simulating bimanual self-touch (end of reaching); and (e) the
post-reach trials included the moment at 300 ms after the tap on the sensor and thus after the
reaching movement. Participants were aware that the *test* force could come at different times
but did not know when they would receive it on every trial. These 5 trial groups were defined
to correspond to different phases of the reaching movement (68,69), and capture the time of
peak velocity, where changes in the somatosensory perception were previously reported (58)
and the post-reach phase where changes are typically minimal (27,32). On average, participants
took 540 ± 47 ms to reach their left hand (**Figure 2d**) with a peak velocity of 95 ± 11 cm/s at
176 ± 30 ms after movement onset (**Figure 2e**). The average times at which the *test* force was
delivered in the *early*, *mid*, and *late* trials were 81 ± 17 ms, 201 ± 31 ms, and 350 ± 27 ms
relative to movement onset, corresponding to 15%, 37%, and 66% of the average reaching time
in the *early*, *mid*, and *late* trials, respectively (**Figure 2f**) (**Materials and Methods**). There
were no significant differences in the forces participants tapped on the sensor with their right
index finger between any trial type ($F(4, 112) = 1.001$, $p = 0.410$, $\eta_p^2 = 0.035$)), confirming
that the participants behaved comparably across all reaching trials (**Supplementary Figure
S1**).

Figure 2. Methods and results of Experiment 1. (a) Participants made a reaching movement with their active right hand from the *starting position* to tap the force sensor (depicted as the cyan sensor) placed above but not in contact with their left passive index finger. Another identical force sensor (depicted as the red sensor) was placed within the probe of the lever controlled by the motor to measure the forces applied on the left index finger. In every trial, a motion tracking sensor (*motion sensor*) recorded the position of the right index finger. The right hand was placed on top of the wooden table (brown surface), which was placed on top of the desk (gray surface). For illustration purposes, the wooden table appears transparent. The reaching movement is shown from a third-person perspective. **(b)** In every trial, the participants received two forces on the pulp of their left index finger – the *test* and the *comparison* force – via the lever of the motor. After receiving both forces, they verbally indicated which force felt stronger. **(c)** Example of a velocity profile of one pilot participant illustrating the different phases for a reaching trial. The various phases were determined using the 3D velocity data, but the horizontal velocity is plotted here for simplicity. In the reaching trials, participants received the *test* force on their left index finger during one of five different phases relative to their

movement: *early* (yellow), *mid* (red), *late* (blue), at the *target* of the movement (green), or 300 ms *post-reach* (brown). The *comparison* force was always delivered 1200 ms after the *test* force and when the right hand had completed the reaching and had stopped moving. In the *baseline* trials, participants kept both hands relaxed and received the *test* and the *comparison* forces at rest. **(d-e)** Individual and group reach durations **(d)** and peak velocities **(e)**. The markers represent individual data, box plots show the medians and interquartile ranges, and raincloud plots depict the data distributions. **(f)** Histograms of the times at which the *test* force was applied to the participants' left index finger in the different types of reaching trials. Times are expressed relative to the time the target was reached (force sensor tap) and binned into 25 ms bins. The vertical dashed line represents the mean time of peak velocity (v_{max}), and the grey shaded area represents the standard error of the mean (s.e.m.). **(g)** Responses and fitted logistic models for all types of trials for one participant. We extracted the PSE ($p = 0.5$) for each participant and each trial type. **(h)** Individual PSE values are shown for each type of reaching trial after being normalized to the *baseline* PSE. The individual PSEs are plotted against the mean time the *test* force occurred in each trial type relative to the target. In the figure, individual points are jittered ($\pm$ 10 ms) to avoid complete overlap. To illustrate the pattern in our data and facilitate the comparison with our hypothesis (**Figure 1b**), we overlaid a regression line based on a generalized additive model. The shaded area represents a 95% confidence interval. **(i)** Normalized group PSE values for each trial type (mean $\pm$ s.e.m.). The touch felt significantly weaker as the reaching time progressed (*late < early, target < early, target < mid, target < late*) but quickly recovered afterwards (*post-reach > target*). Asterisks denote significant pairwise comparisons of interest after corrections for multiple comparisons. **(j)** Normalized individual PSE data during reaching (*early* to *target*). Data are jittered ($\pm$ 10 ms) to avoid complete overlap. The PSEs of each participant were fitted with a linear model, and we display the group's linear fit. The shaded area depicts a 95% confidence interval. **(k)** Slope coefficients of the participants' linear models showing the change in PSEs over the reaching. The slopes were overall negative, indicating an increasing attenuation, and significantly different from zero. The markers represent the individual slope coefficients, box plots show the medians and interquartile ranges, and raincloud plots depict the coefficients' distributions (** $p < 0.01$, *** $p < 0.001$).

The participants performed an additional block of 56 *baseline* trials in which the participants remained still while performing the force discrimination task. These trials were conducted either before or after the block of reaching trials in a fully counterbalanced order and served to quantify the participants' perceived magnitude of the *test* force in the absence of action. For all types of trials (*early, mid, late, target, post-reach*, and *baseline*), we fitted the participants' responses from the force-discrimination task with logistic models (see **Supplementary Figure S2** for all individual fits). All fits were very good (McFadden $R^2 > 0.41$). Next, we extracted the point of subjective equality (PSE), which indicates the intensity at which the *test* force felt as strong as the *comparison* force ($p = 0.5$) and represents the perceived intensity of the *test* force per trial type (**Figure 2g**). A lower PSE in one trial type compared to another indicates that the *test* force felt weaker in the former than in the latter trial type.

PSE values of the *reaching* trials were first normalized to the PSEs of the *baseline* trials ($PSE_{reaching} - PSE_{baseline}$) and then subjected to a one-way repeated measures ANOVA with the

type of trial (*early*, *mid*, *late*, *target*, *post-reach*) as the within-subjects factor. The type of trial
was significant: $F(2.09, 58.44) = 15.856$, $p < 0.001$, $\eta_p^2 = 0.362$. Planned pairwise comparisons
revealed that the perceived intensity of the *test* force gradually decreased the closer the touch
was delivered to the moment of the predicted self-touch (**Figure 2h-i**): specifically, the PSEs
were significantly lower in the *late* compared to the *early* trials ($n = 29$, $t(28) = -2.956$, $p =$
0.009 *FDR-corrected*, $CI^{95} = [-0.090, -0.016]$, $d = -0.549$), in the *target* compared to the *early*
trials ($n = 29$, $W = 11$, $p < 0.001$ *FDR-corrected*, $CI^{95} = [-0.234, -0.115]$, $rrb = -0.949$), in the
target compared to the *mid* trials ($n = 29$, $W = 31$, $p < 0.001$ *FDR-corrected*, $CI^{95} = [-0.225,$
$-0.097]$, $rrb = -0.857$) and in the *target* compared to the *late* trials ($n = 29$, $W = 41$, $p < 0.001$
FDR-corrected, $CI^{95} = [-0.180, -0.064]$, $rrb = -0.811$). Differences in the PSEs between *early*
and *mid* trials ($n = 29$, $t(28) = 0.829$, $p = 0.414$ *FDR-corrected*, $CI^{95} = [-0.021, 0.050]$, $d =$
0.154) and between *mid* and *late* trials ($n = 29$, $t(28) = 1.813$, $p = 0.094$ *FDR-corrected*, $CI^{95} =$
$[-0.005, 0.083]$, $d = 0.337$) were not significant. The lowest PSEs thus occurred at the time of
contact between the two index fingers (*i.e.*, *target* trials) when the *test* force was perceived to
be significantly weaker than the *test* force at all other probed times. These PSEs quickly
recovered in the *post-reach* trials and were significantly greater than in the *target* trials ($n = 29$,
$W = 412$, $p < 0.001$ *FDR-corrected*, $CI^{95} = [0.135, 0.268]$, $rrb = 0.894$).

To further corroborate the gradual decrease in PSEs during reaching, we next fitted each
participant's normalized PSEs in *early*, *mid*, *late*, and *target* trials with a linear regression
model (**Figure 2j**). As hypothesized, the extracted slope coefficients were negative ($\beta =$
$-4.142 \times 10^{-4} \pm 4.289 \times 10^{-4}$) and significantly different from zero ($n = 29$, $W = 14$, $p < 0.001$,
$CI^{95} = [-5.056 \times 10^{-4}, -2.421 \times 10^{-4}]$, $rrb = -0.936$) (**Figure 2k**), indicating that the *test* force was
perceived weaker the closer it was delivered to the expected time of self-touch. Similar results
were obtained when fitting the PSEs from the *early*, *mid*, and *late* reaching trials only.

Taken together, the results of Experiment 1 demonstrated that the touch felt gradually weaker
as the reaching movement progressed, attained its minimum perceived intensity at the moment
of the predicted self-touch, and quickly recovered after the reaching.

Experiment 2. The gradual attenuation of touch during reaching depends on whether the
reaching is expected to produce self-touch.

The results of Experiment 1 can be attributed to a dynamic influence of the internal model's
predictions on somatosensory perception during the reaching movement. To rule out that the
observed PSE effects are due to other factors related to the movement or the task *per se* (for
example, general anticipation of tactile stimulation on the left index finger, unspecific time
effects, divided attention to both hands or feelings of agency over the action), we conducted a
control experiment. In Experiment 2, we hypothesized that if the temporal tuning of
somatosensory perception observed in Experiment 1 stems from the predictions of the internal
model, then performing a reaching movement with the right hand that does not have any tactile
consequences for the left hand should not modulate somatosensory perception in time. This
would occur because, in such a scenario, the internal model would not predict somatosensory
input on the left hand based on the efference copy (**Figure 3a**).

Two new groups of naive participants participated in Experiment 2. Participants in the *Self-*
touch group (n = 29, 15 female, 28 right-handed, aged 18-37 years) performed the same task
as those in Experiment 1: they reached with their right index finger to tap the sensor above
their left index finger and received forces on their left index finger at different times with
respect to the movement of their right hand (**Figure 3b**). In contrast, participants in the *No self-*
touch group (n = 27, 15 female, 26 right-handed, aged 21-37 years) performed the same
reaching movement with their right hand as participants in the *Self-touch* group did, but they
were instructed to never make contact between their two fingers (**Figure 3b**). Rather, they
approached a distance sensor placed on top of, but not in contact with, their left index finger
which triggered the force once the position of the right index finger became lower than a pre-
set position threshold. Then they lifted their right hand as the *Self-touch* group, simulating a
virtual tap in the air. The stimuli were delivered at the same time between the two groups, with
the *test* force in the *target* trials being applied upon triggering the distance sensor in the *No*
self-touch group and upon triggering the force sensor in the *Self-touch* group (see **Materials**
and Methods).

As in Experiment 1, participants in the *Self-touch* group were told that their hands would make
contact through the probe in some trials, simulating self-touch, but they would not know
beforehand which trials these would be. In contrast, participants in the *No self-touch* group
were instructed to stop their right hand at a certain distance above their left hand and then lift
it without ever making contact between their hands. Previous studies (27,28,30) have shown
that if participants deliberately stop their right hand movement above their left hand, the touch
applied on the left hand is not attenuated because it cannot be predicted by the motor command.
In contrast, if participants predict based on their movement that they will make contact between
their two hands but unexpectedly miss the contact, robust attenuation is still observed (28,30)
indicating that the major driver of somatosensory attenuation is the prediction of self-touch
based on the efference copy and not the hands' contact *per se*. Therefore, in Experiment 2, we
expected to replicate all the effects of Experiment 1 with the *Self-touch* group and further
observe no temporal modulation in the *No self-touch* group because the internal model would
no longer predict touch on the left hand.

The two groups performed reaching movements of comparable durations (**Figure 3c-d**). The
No-self touch group exhibited slightly higher peak velocity than the *Self-touch* group (**Figure**
3e-f), but both groups received the *test* forces at similar times (**Figure 3g-h**, **Supplementary**
Text S1). As in Experiment 1, the *Self-touch* group pressed comparable forces across the
reaching trials (**Figure S3**) and all psychometric fits of both groups were very good (McFadden
R^2 > 0.47) (**Supplementary Figures S4-S5**). Importantly, the *Self-touch* group replicated all
effects of Experiment 1 (**Supplementary Text S2, Figure S6**).

To compare the somatosensory perception of the two groups, we inserted the participants'
normalized PSEs in a mixed ANOVA with the group as a between-subjects factor and the type
of trial as a within-subjects factor. The model revealed a significant main effect of the trial type
($F(3.27, 176.33)$ = 10.044, $p < 0.001$, η_p^2 = 0.157), a significant main effect of group ($F(1, 54)$
= 8.284, p = 0.006, η_p^2 = 0.133) and a significant group-by-trial type interaction ($F(3.27,$
176.33) = 7.072, $p < 0.001$, η_p^2 = 0.116). Pairwise comparisons revealed that the PSEs were

significantly lower in the *Self-touch* group than in the *No self-touch* group, in the *mid* ($t(53.89)$
$= -2.206$, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.156, -0.007]$, $d = -0.590$), *late* ($t(53.37) =$
-2.488, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.171, -0.018]$, $d = -0.663$), *target* ($W = 129$, $p <$
0.001 *FDR-corrected*, $CI^{95} = [-0.286, -0.123]$, $rrb = -0.670$), and *post-reach* trials ($t(53.97) =$
-2.299, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.162, -0.011]$, $d = -0.614$). No difference was
detected in the *early* trials ($t(51.84) = -0.338$, $p = 0.737$ *FDR-corrected*, $CI^{95} = [-0.105, 0.075]$,
$d = -0.091$) (**Figure 3i**). In other words, although both groups perceived a comparable
magnitude of touch in the early phase of the reaching, their perception deviated at later timings
depending on whether the reaching movement had somatosensory consequences for the left
hand or not. Further control analyses indicated that the observed PSE differences between the
two groups were not due to differences in the kinematics or task demands of the reaching
movements (**Supplementary Text S3, Figure S7**).

Next, we fitted linear models to the PSEs during reaching (*early*, *mid*, *late*, and *target* trials)
and extracted the slope coefficients for every participant in each group. While the slope
coefficients from the *Self-touch* group were negative ($\beta = -2.687 \times 10^{-4} \pm 3.210 \times 10^{-4}$) and
significantly different from zero ($n = 29$, $t(28) = -4.507$, $p < 0.001$, $CI^{95} = [-3.908 \times 10^{-4}$,
$-1.466 \times 10^{-4}]$, $d = -0.837$), replicating Experiment 1, the slope coefficients from the *No self-*
touch group were positive ($\beta = 9.233 \times 10^{-5} \pm 2.913 \times 10^{-4}$) and did not significantly differ from
zero ($n = 27$, $t(26) = 1.647$, $p = 0.112$, $CI^{95} = [-2.290 \times 10^{-5}, 2.076 \times 10^{-4}]$, $d = 0.317$). As expected,
the coefficients were significantly lower in the *Self-touch group* compared to the *No self-touch*
group ($t(53.97) = -4.412$, $p < 0.001$, $CI^{95} = [-5.251 \times 10^{-4}, -1.970 \times 10^{-4}]$, $d = -1.178$) (**Figure**
3j-k). Similar results were obtained when fitting the PSEs from the *early*, *mid*, and *late*
reaching trials only.

Together, the results of Experiment 2 indicate that the gradual attenuation of the perceived
magnitude of touch on the left hand depends on whether the reaching movement of the right
hand has somatosensory consequences for the left hand (*i.e.*, self-touch).

Figure 3. Methods and results of Experiment 2. (a) When reaching with the active right hand towards the passive left hand but without planning to touch it (*i.e.*, planning to stop the movement before contact), the brain does not predict somatosensory consequences on the left hand based on the efference copy of the right hand. Consequently, somatosensory perception on the left hand should not show any modulation over time. **(b)** As in Experiment 1, participants in the *Self-touch* group made a reaching movement with their right hand from the *starting position* to tap the force sensor (cyan sensor) placed above, but not in contact with, their left index finger. Participants in the *No self-touch* group (right) also made a reaching movement with their right hand from the *starting position* to approach a distance sensor (yellow sensor) placed above, but not in contact with, their left index finger. The reaching movement is shown from a third-person perspective. **(c-f)** The durations of reaching movements were comparable between the two groups **(c-d)**. The *Self-touch* group exhibited a lower peak velocity than the

No self-touch group **(e-f)**. **(g-h)** Histograms of the times at which the *test* force was applied in
the different types of reaching trials for the **(g)** *Self-touch* group and **(h)** *No self-touch* group.
The times did not significantly differ between the two groups. Times are expressed relative to
the time the target was reached and binned into 25 ms bins. The vertical dashed line represents
the mean time of peak velocity (v_{max}), and the grey shaded area represents the standard error
of the mean (s.e.m.). **(i)** PSEs were significantly lower in the *mid*, *late*, *target*, and *post-reach*
trials when self-touch was expected (*Self-touch* group) than when it was not predicted (*No Self-*
touch group). Asterisks denote pairwise comparisons of interest after corrections for multiple
comparisons. **(j)** Individual normalized PSE data during the reaching movement only (*early* to
target) are plotted for each group. Individual points are jittered ($\pm$ 10 ms) to avoid complete
overlap. We fitted a linear model for every participant in every group separately, but for
illustration purposes, we display the group fits. The shaded area depicts a 95% confidence
interval. **(k)** In the *Self-touch* group the slopes were overall negative and significantly lower
than the slopes in the *No self-touch*, which did not differ from zero. The markers represent the
slope value for each participant, box plots show the medians and interquartile ranges, and
raincloud plots depict the data distributions. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, *n.s.*
nonsignificant)

Discussion

Determining the time course of somatosensory perception during movements to self-touch is
essential for understanding how the brain's predictions influence perception over time
(15,19,62), as well as for evaluating alterations in clinical and neurobiological conditions such
as psychosis spectrum disorders (65,67,70–75), functional movement disorders (76),
Parkinson's disease (36) and ageing (37). By combining kinematic recordings with
somatosensory psychophysics, the present study tested the time course of somatosensory
perception during a movement to self-touch; that is, how participants perceived touch on their
left hand as their right hand reached to touch the left hand, and moved away from it. Going
beyond earlier research that did not test perception during the entire movement of the active
hand (22–24,28–31,33–39,45,57,58) or tested perception at times that were not linked to the
reaching movement of the active hand (26,27,32,46,77), our study revealed the timeline of
somatosensory perception on the passive left hand during the different phases of the active
right hand's movement to touch it. In Experiment 1, we demonstrated that touch on the left
hand was perceived with gradually attenuated intensity as the reaching of the right hand
progressed: the touch felt significantly weaker in the late reaching phase than in the early
reaching phase and was significantly more attenuated at the moment of self-touch. The
minimum perceived intensity was observed at the time of self-touch between the hands, and it
quickly recovered after the reaching had ended. All these results were replicated in Experiment
2, where participants in the *Self-touch* group exhibited the same temporal tuning of
somatosensory responses. In contrast, participants in the *No self-touch* group who performed
similar right hand movements that had no somatosensory consequences for the left hand
showed no such modulation. Complementary analysis in that group (**Text S2**) showed that the
perceived intensity of touch did not change significantly at any probed time during the right
hand movement. Together, our findings indicate a temporal tuning of somatosensory perception

during movements to self-touch and highlight the role of the sensorimotor context in predicting
and attenuating the received sensory input.

Our results are in strong agreement with the computational framework of internal forward
models. Specifically, when reaching with the right hand to touch the left hand, the internal
model uses the efference copy of the right hand movement to predict the sensory consequences
for the right and left hands, including self-touch and its expected timing (15,17,19). These
predictions are combined with the received sensory input (*e.g.*, visual and proprioceptive)
during the movement to re-estimate the state of the right limb *(e.g.*, position, velocity) and
update the brain's predictions about the remainder of the movement (61–63). Because both
sensory (18,78,79) and motor signals (80–82) are noisy, the moment of the predicted touch
corresponds to a time window reflecting a probability distribution rather than a single point in
time and thereby incorporates a certain tolerance for overcoming the intrinsic noise (17). As
the movement of the right hand progresses, new sensory input is received and the predictions
of the internal model become increasingly more precise (61–63), narrowing the time window.
The received touch is attenuated proportionally to how close they were received with respect
to this predicted time window, which continuously narrows during the movement.

Notably, the gradual attenuation of somatosensory perception during movements to self-touch
cannot be attributed to task-related differences such as dividing attention between the hands,
the mere movement of one hand, a sense of agency, or general cognitive anticipation of tactile
stimulation. If this were the case, then we would have observed the same perceptual tuning in
the participants of the *No Self-touch* group, in which the same discrimination task and right
hand reaching movements were performed. In contrast, we did not observe such a temporal
modulation in those participants, and a Bayesian analysis supported the absence of gradual
attenuation effects. The direct comparison between the two groups revealed significant
differences at almost all probed time points, indicating significantly greater attenuation in the
Self-touch group than in the *No self-touch* group. This was further supported by linear
regression analyses, which showed a significant gradual increase in attenuation during the
reaching both in Experiment 1 and in the *Self-touch* group of Experiment 2 but not in the *No*
Self-touch group. Finally, further control analyses revealed that these effects cannot be
attributed to differences in kinematics or task demands between the two groups.

The results of Experiment 2 demonstrated that the differences in somatosensory perception
between the two groups were driven by the sensorimotor context within which participants
moved; that is, whether the internal model could predict touch on their left hand based on the
motor command of their right hand. Theoretically, the brain should attenuate incoming touch
that is likely to be the somatosensory feedback of the movement (17,19,63). However, the
brain needs to first determine the sensorimotor context that describes the relationship between
the movement and the predicted sensory input (*i.e.*, internal model) (83,84). This is critical
because the same motor command will be associated with different sensory consequences in
different contexts. For example, when manipulating objects, the same movement can produce
different predictions about sensory feedback in a context where both hands manipulate the
same object compared to a context where each hand manipulates a different object (85).
Similarly, when moving the right hand towards the left hand, the movement of the right hand

may predict touch on the left hand in a self-touch context but not in a context where the right
hand deliberately stops before contacting the left hand (27,28,30). Likewise, pressing the right
index finger a few centimetres away from the left index finger should not produce any
expectations about receiving touch on the left hand (17,23,39) unless in a context where the
right hand holds a tool to directly press on the left one (38). Consequently, the movements the
No Self-touch group performed were not associated with any somatosensory consequences on
their left hand, which explains why we did not observe an attenuation of the perceived
magnitude during their reaching movements. Our results therefore provide evidence that the
brain evaluates the sensorimotor context to dynamically shape somatosensory perception in
time depending on whether somatosensory input can be predicted based on the efference copy.

Our results cannot be attributed to backward or forward masking of the *test* force by the press
of the right index finger. The *test* force was presented at 184 ms and 163 ms on average for
Experiment 1 and Experiment 2 (*Self-touch* group) before the target was reached (*late* trials),
and at 300 ms afterwards (*post-reach* trials), while backward masking effects were not detected
at 150 ms (86) and forward masking effects last only up to 100-200 ms (87–90). In addition,
our findings do not support the proposal that motor predictions enhance the expected
somatosensory signals (91), as we did not observe any enhancement effects at any of the time
points probed in either of the two experiments. Instead, we consistently observed robust
attenuation effects when subjects expected self-touch based on their movement. In line with
the present findings, we recently demonstrated that the previously reported "enhancement"
effects these proposals are based on (92) stem from a biased reference condition used rather
than action prediction (30). If a bias-free condition is employed, no enhancement effects are
experimentally observed (30). Moreover, our effects contradict the proposal that attenuation
effects are due to non-predictive mechanisms related to double tactile stimulation of both
fingers (92) as we observed somatosensory perception at times *before* the two hands made
contact (*i.e.*, in *mid* and *late* trials), in line with earlier findings (27) (see also (21)). Critically,
our effects were dependent on whether participants expected self-touch (*Self-touch* group) and
extended beyond the mere influence of the movement of the right hand, the touch on the left
hand, or the discrimination task *per se* (*No self-touch* group). Consistent with the current
results, we recently demonstrated that attenuation effects are observed independently of the
double tactile stimulation of the two hands: touch is robustly attenuated if participants predict
that they will make contact between their two hands but unexpectedly miss the contact (30)
(see also (28) for similar findings). Therefore, the present study provides additional evidence
that somatosensory attenuation is driven by the prediction of the sensory consequences of
voluntary movement rather than somatosensory reafference *per se*.

A methodological strength of the present study is that it assessed how the predictions of the
internal model affect the somatosensory perception of a stationary limb. Previous studies have
shown that somatosensory perception is 'gated' on a limb that moves compared to one at rest
(47,50,86,93–98) and that this gating can be variably affected by movement (compare (58,99–
102)). However, several studies have consistently demonstrated that these gating effects are
also observed during passive movements (*i.e.*, in the absence of motor signals), indicating that
somatosensory effects on a moving limb are influenced by postdictive mechanisms related to
peripheral reafference (51–54,56,103). In contrast, somatosensory perception on a limb that

remains stationary is not subject to gating effects (54,98,101,103–105). Moreover,
somatosensory perception on a passive limb is attenuated only if the received touch results
from an active movement, not from a passive movement (34) or passive stimulation (27) (see
also (35)). Thus, our approach isolated the effects of sensorimotor predictions while avoiding
the confounding influence of peripheral mechanisms.

An earlier study reported constant tactile facilitation on the left hand as the right hand moved,
peaking at the time the right hand reached its maximum velocity (58). However, we did not
observe such an effect in either of our two experiments despite having a greater sample size
than this earlier study. A reason for this discrepancy could be that we used suprathreshold force
stimuli while that study used near-threshold vibrotactile stimuli (58). In the auditory domain,
studies using near-threshold and suprathreshold stimuli have shown different perceptual effects
(106,107). However, this remains a speculation as there is no evidence for a similar effect in
the tactile modality. Another difference between the two studies is that the earlier study (58)
was conducted in darkness, while our participants could use vision to guide their movements.
The authors of the earlier research suggested that the darkness facilitated the participants'
somatosensory perception to guide their hand movement (48,49). However, we believe this
interpretation requires further evidence given other findings reporting the opposite effect
(102,108) or null findings (47,109) regarding the interaction between vision and
somatosensation during arm movements. Finally, another possible reason for this discrepancy
is that participants in the study of (58) reached with their right hand to either their left index
finger or thumb while receiving stimulation on their little finger. Consequently, their
experimental design introduced an incongruency between the finger where touch was expected
based on the movement and the finger where touch was received. While the authors argue that
tactile perception is uniformly modulated across the whole target hand (48) other research has
shown that such incongruency can hamper somatosensory attenuation (110). Here, in all trials
of our experiments, the participants reached with their right hand towards their left index finger
while receiving stimulation on their left index finger. Consequently, this allowed us to
investigate how somatosensory attenuation on the left index finger changes when the
participants form expectations about receiving touch on the same finger.

Understanding how expectations about the somatosensory consequences of the movement (*i.e.*,
the internal model) dynamically shape somatosensory perception in time is clinically
important. Patients with schizophrenia show deficits in attenuating self-generated
somatosensory input by exhibiting perceptual and neural responses to self-touch that are
comparable to those of externally generated touch (73,74). Using the same task that was used
in the present study, we previously showed that individuals with high schizotypal personality
traits, which are considered a subclinical analogue of schizophrenia (111,112), also show a
reduced ability to attenuate self-touch compared to externally generated touch (29). In addition
to the somatosensory domain, patients with schizophrenia also show difficulty in attenuating
self-generated speech (113–115) (see also (116)) and oculomotor responses (117). Together,
these findings indicated that individuals with schizophrenia form incorrect predictions about
the sensory consequences of their actions, misattributing the received self-generated input to
external sources, producing the hallucinations and delusions commonly associated with
schizophrenia (66,72,75,118,119). Interestingly, one study showed that patients with

schizophrenia have a reduced attenuation of neural responses to self-generated auditory stimuli but a-close-to-normal attenuation for the same stimuli when delivered with a 50 ms delay (120), suggesting that the prediction of the sensory consequences of the action might be delayed in patients with schizophrenia (120,121) (see also (122)). Additionally, both patients with schizophrenia (123–125) and individuals with high schizotypal traits (126–129) tend to exhibit a perturbed temporal profile of multisensory integration, experiencing stimuli that are further apart in time as cooccurring. We therefore speculate that patients may show a different temporal tuning of somatosensory perception during self-touch (*e.g.,* flatter slopes, or shifted temporal window) than that observed in the present study. This hypothesis warrants further exploration in future experiments.

Sensory attenuation is not unique to humans but is observed across many species with research on crickets (130,131), mice (132–134), weakly electric fish (135–137), and primates (138–140) showing selective attenuation and even cancellation of neural responses to self-generated sensory inputs. Broadly, this serves to increase the salience and facilitate the interpretation of externally generated sensations (141–146). Of particular interest are recent reports of a temporally tuned cancellation of self-produced sounds in mice. Specifically, mice trained to associate tones with moving a lever to a specific position, showed attenuated responses in the auditory cortex to these self-generated sounds when presented at the expected time (*i.e.*, when the lever reached the expected position) (134). Importantly, this attenuation was not of fixed magnitude during movement: auditory responses were decreased for sounds played just before movement, or 80 ms after the predicted time or onset and reached their minimum at the expected time (134). In addition, mice exhibit attenuated activity in the primary somatosensory cortex when receiving the pulsatile stimulus on their whiskers at the predicted time, while less attenuation is observed if the temporal delays of 50 ms and 100 ms from the predicted time are introduced (147). The similarity of these findings with those of the present study suggests the presence of an evolutionarily conserved mechanism for generating precise temporal predictions to attenuate expected sensory inputs.

Materials and methods

Participants

Ninety naive participants (47 females and 43 males; 87 right-handed, 2 left-handed, and 1 ambidextrous, aged 18-37 years) were recruited to participate in Experiments 1 and 2. The sample size was set to 30 participants per experiment/group based on our earlier studies investigating somatosensory perception using the same task and equipment (30–35,46). After data collection, one participant was excluded from Experiment 1 due to missing kinematic data, and four participants were excluded from Experiment 2: one from the *Self-touch* group because the same response was given in 95% of the *target* trials resulting in an unreliable psychometric fit, and three participants from the *No self-touch* group because they performed very fast right hand movements, resulting in too few *early* trials (< 10 trials, arbitrary threshold) (see *Kinematic trials – segmentation*). Therefore, the analysis of Experiment 1 was based on data from 29 adults, and the analysis of Experiment 2 was based on data from 56 adults: 29 in the *Self-touch* group and 27 in the *No self-touch* group.

553 In both experiments, handedness was assessed using the Edinburgh Handedness Inventory
554 (148). All participants provided their informed written consent and reported not having any
555 neurological or psychiatric disorders or taking any psychoactive medication. The experiments
556 were approved by the Ethical Review Authority of Stockholm (registration number: 2021-
557 03790).

Experimental apparatus and procedures in Experiment 1

In both experiments, the participants sat at a desk and rested their left hand palm up, with the
left index finger placed inside a support (**Figure 2a**). A vacuum pillow (AB Germa) was
provided to support the participants' left arm and increase their comfort. On top of the pulp of
their left index finger laid a cylindrical probe (20 mm diameter) that was attached to a lever
controlled by a DC electric motor (Maxon EC Motor EC 90 flat; manufactured in Switzerland)
and operated by an Arduino microcontroller (**Figure 2a,** *lever*). A force sensor (FSG15N1A,
Honeywell Inc.; diameter, 5 mm; minimum resolution, 0.01 N; response time, 1 ms;
measurement range, 0–15 N) was placed within the probe to record the forces exerted on the
left index finger by the motor (**Figure 2a,** *force sensor* depicted in red). A force of 0.1 N was
constantly applied to the left index finger to maintain contact between the probe and the finger.
A second identical force sensor was placed inside an aluminium capsule (20 mm diameter) and
positioned on top of, but not in contact with, the cylindrical probe (**Figure 2a,** *force sensor*
depicted in cyan). The right forearm was comfortably placed on top of sponge padding, and the
right hand rested above a wooden table (70 cm × 25 cm × 12.5 cm). The wooden table was
placed on the desk in front of the participants covering their left hand, and thus preventing
direct visual feedback from the left hand and fingers. The right index finger was placed at a
position marked on top of the wooden table (**Figure 2a**, *starting position*), corresponding to 25
cm to the right of the left index finger.

Upon an auditory GO-cue generated by Arduino, participants were instructed to perform a
reaching movement with their right hand from the starting position towards the force sensor
located above their left hand. They were instructed to perform a fast but natural movement that
would be concluded once they tapped the force sensor with their right index finger (*i.e.*,
touching the force sensor with the right index finger and immediately lifting the finger).
Participants were allowed to freely look at their right hand and the force sensor as needed to
guide their movement. A motion tracking sensor (Polhemus Liberty, Micro Sensor 1.8™) was
attached to the right index finger with medical tape to continuously record the right index finger
position (**Figure 2a,** *motion sensor*). White noise was provided by a laptop through a pair of
headphones to mask any sounds made by the motor.

Force trials

588 In each experiment, there were 560 reaching trials and 56 *baseline* trials, resulting in a total of
589 616 trials per participant grouped in two experimental blocks. The seven possible intensities of
590 the *comparison* force (1, 1.5, 1.75, 2, 2.25, 2.5, 3 N) were tested 16 times for every type of
591 reaching trial and 8 times in the *baseline*. The intensity of the *test* force was always fixed at 2
592 N.

During the reaching trials, we experimentally manipulated the times at which the *test* force was applied on the left index finger. We applied the *test* force at five different times based on pilot experiments to correspond to different phases of a natural right hand movement reaching toward the left hand. The *test* force could be applied at three different times during the movement of the right hand (*i.e.*, at 250 ms, 370 ms, and 550 ms after the trial onset). Based on the time of the *test* force occurrence, the trials were binned in the analysis to correspond to the *early*, *mid,* and *late* phases of the right hand movement (see *Kinematic trials – segmentation*), at the time when the participants tapped the sensor (*target*), which simulated self-touch between the hands (system delay ~35 ms), and at a fixed time of 300 ms after tapping (*post-reach*). All 5 types of reaching trials were randomly interleaved and had an equal probability of occurrence (20%). During the *baseline* trials, participants remained motionless, and the *test* force was applied 100 ms after the onset of the trial, followed by the *comparison* force 1200 ms later. In every trial, after receiving both forces, participants responded verbally about which of the two forces (the first force - *test* or the second force - *comparison*) felt stronger. Participants were instructed not to balance their responses (*i.e.*, saying 50% of the time that the first or the second force is stronger), and were further instructed to make their best guess if the intensity of the two forces felt similar. No feedback was ever provided to the participants regarding their responses.

Experimental apparatus and procedures in Experiment 2

Experiment 2 was a between-group experiment. The *Self-touch* group (**Figure 3b**) performed the same task as the participants in Experiment 1. For the *No self-touch* group, the force sensor that participants approached with their right index finger (**Figure 3b**, *force sensor* depicted in cyan) was replaced by a distance sensor (Ultrasonic Distance Sensor HC-SR04 5 V Version) (**Figure 3b**, *distance sensor* depicted in yellow). The distance sensor was placed on top of, but not in contact with, the probe and was connected to an Arduino microcontroller that controlled a servo motor (Grove-Servo). Once the distance sensor detected that the position of the right index finger became lower than a pre-set position threshold, it triggered a servo motor that hit the force sensor that triggered the *test* force. The servo motor and the force sensor were hidden from the participants' view. The position threshold was set to approximately match the position participants had when pressing the force sensor with their right index finger in Experiment 1 and in the *Self-touch* group. Participants were instructed to perform a fast but natural movement with their right hand toward the distance sensor and to conclude the movement by tapping the right index finger above the sensor without touching it. That is, the participants were instructed to perform movements like those of Experiment 1, with the difference that they never experienced contact between their two fingers and thus never expected to make such contact. The end of the reaching was defined as the time point at which the distance sensor was triggered by the downward movement of the right index finger.

All probed times in Experiment 2 were identical for the *Self-touch* and *No self-touch* groups. However, forces were delivered 100 ms later in Experiment 2 than in Experiment 1. All the other details, including the analysis, remained identical to those of Experiment 1.

Kinematic trials – preprocessing

634 The position of the right index finger was recorded at a sampling rate of 240 Hz. TTL signals
635 were used to indicate the onset of the trial, as well as the *test* force and the *comparison* force,
636 thus co-registering the force trials with the kinematic trials. We smoothed the position data
637 using a moving average filter of 5 data points span ($\approx$ 21 ms).

Kinematic trials – segmentation

For each reaching trial, we calculated the movement onset as the first time point during the trial
when the velocity exceeded 5 cm/s for 100 ms sequentially (99,101,102). The end of the
reaching (*i.e.*, offset) was calculated as the first time point when the force exerted on the force
sensor exceeded a pre-set threshold (0.3 N) indicating that participants tapped the sensor with
their right index finger and thus concluded the movement. The duration of the movement was
calculated as the time difference between the movement offset and onset.

Next, we calculated the time when the 3D velocity of the right hand reached its peak value. To
capture the peak velocity of the reaching movement, which typically occurs during the first
half of the movement (68), and not the peak velocity of the index finger when tapping the
sensor (occurring at the end of the movement), we restricted the time window of the peak
velocity to within 67% of the movement duration.

Based on the calculated kinematic parameters, we binned the reaching trials into three distinct
categories depending on when the *test* force occurred within a specific trial: *early* trials (*test*
force occurring between the movement onset and the time the right hand velocity reached 85%
of its peak value), *mid* trials (*test* force occurring after the right hand velocity reached 85% of
its peak value up until it dropped below 85% of its peak value), and *late* trials (*test* force
occurring after the right hand velocity dropped below 85% of its peak value and before the
right index finger made contact with the force sensor). In addition, the trials when the *test* force
occurred simultaneously with the participant tapping the force sensor were classified as *target*
trials, and the trials when the *test* force occurred 300 ms after the participants' tap on the force
sensor were classified as *post-reach* trials (**Figure 2c**). Therefore, the final five groups of
reaching trials entering the statistical analysis were *early*, *mid*, *late*, *target*, and *post-reach*
trials.

Exclusion of trials in Experiment 1

As mentioned earlier, each participant was administered 616 trials resulting in 17864 trials in
total (29 × 616). We excluded (a) any *baseline* or reaching trials in which the intensity of the
test force (2 N) was not properly applied (*test* < 1.85 N or *test* > 2.15 N), or the participants'
force responses or kinematic data were not properly registered, (b) any reaching trials in which
the participants failed to tap the sensor or did not press/release the sensor properly, (c) any
reaching trials in which the participants did not move as instructed (*e.g.*, they moved before the
auditory GO-cue, they started their movement from a position closer to the force sensor than
the starting position (< 5 cm away from the starting position on the x-axis)), (d) any reaching
trials in which the participants received the *comparison* force while moving, and (e) any *early*,
mid, or *late* trials in which the *test* force was applied either before the participants started
moving (*i.e.*, participants were too slow in starting their movement) or after they tapped the
sensor (*i.e.*, participants were too fast in finishing their movement). Finally, since our setup

contained metallic surfaces (*e.g.*, the aluminium capsule) that can distort the electromagnetic
recordings of the motion tracking sensor when in close proximity, we rejected (f) any reaching
trials in which the sensor was inside the distortion range at times used for trial segmentation
(see below). In total, we rejected 1590 out of 17864 trials (8.9%). This resulted in an average
of 110 *early*, 84 *mid*, 92 *late*, 109 *target*, 110 *post-reach,* and 56 *baseline* trials.

Exclusion of trials in Experiment 2

We used identical criteria for excluding trials as in Experiment 1 with the exception that in the
No self-touch group we also rejected any *target* and *post-reach* trials in which the participants
did not trigger the distance sensor when moving their finger downwards. In addition, for this
group we also modified our exclusion criteria under b) (see above) to exclude any trials in
which the distance sensor was not properly triggered, instead of participants failing to press the
sensor. In total, we excluded 1411 out of 17864 trials (7.9%) in the *Self-touch* group and 2051
out of 16632 trials (12.3%) in the *No self-touch* group. This resulted in an average of 87 *early*,
101 *mid*, 108 late, 107 *target*, 108 *post-reach* trials, and 56 *baseline* trials for the *Self-touch*
group and an average of 64 *early*, 114 *mid*, 136 *late*, 85 *target*, 85 *post-reach,* and 56 *baseline*
trials for the *No self-touch* group.

Psychophysical fits

The participants' responses were fitted with psychometric curves (**Figure 2g**), using a
generalized linear model (Equation 1):

$$p = \frac{e^{\beta 0 + \beta 1 x}}{1 + e^{\beta 0 + \beta 1 x}} \text{ (Equation 1)}$$

We extracted the point of subjective equality ($PSE = -\frac{\beta 0}{\beta 1}$) that corresponds to the intensity
at which the *test* force felt equally strong to the *comparison* force ($p = 0.5$). This was our
variable of interest since it represents the perceived intensity of the *test* force. For control
analysis, we also extracted the just noticeable difference in Experiment 2 that reflects the
participants' discrimination capacity in the psychophysics task ($JND = \frac{log\ (3)}{\beta 1}$) and
corresponds to the difference between the thresholds at $p = 0.5$ and $p = 0.75$. A higher JND
indicates poorer discrimination ability for the applied forces, suggesting greater difficulty of
the task. Before fitting the responses, the *comparison* forces were rebinned to the nearest of
seven possible force intensities (1, 1.5, 1.75, 2, 2.25, 2.5, or 3 N). The goodness of fit was
assessed using McFadden's pseudo R^2. The group psychometric curves were generated using
the *PsychShape* function in R.

Statistical analyses

The data were analysed in R (version 4.2.2) (149) and JASP (version 0.17) (150). We performed
repeated measures and mixed ANOVAs to analyse the data. In the case of sphericity violations,
Greenhouse-Geisser corrections were applied. Effect sizes for the ANOVA were given by
partial eta-squared ($\eta_p{}^2$). Planned pairwise comparisons between trial types were performed
using parametric (paired or independent-sample *t*-tests) or nonparametric (Wilcoxon signed-
rank and Wilcoxon rank sum) tests depending on the normality of the variable distributions.

The normality of the data was assessed with the Shapiro–Wilk test. Corrections for multiple comparisons were performed using the false discovery rate (FDR) and are denoted as "*FDR-corrected*". For every statistical comparison, we report the corresponding statistic, the 95% confidence intervals (CI^{95}), and the effect size (Cohen's d or the matched rank-biserial correlation (rrb)). Linear regression and multiple linear regression were performed using the *lm* function of the R *stats* package. A Bayesian control analysis was performed to provide evidence in favour of the null hypothesis for comparisons of interest. The analysis was done in JASP using default Cauchy priors of 0.707, where the Bayes factor BF_{0+} represents the likelihood of the null hypothesis compared to the alternative hypothesis for the one-tailed tests. We interpret a Bayes factor from 1 to 3 as providing 'anecdotal' support for the null hypothesis, a Bayes factor from 3 to 10 as providing 'moderate' support for the null hypothesis, and a Bayes factor above 10 as providing 'strong' support for the null hypothesis (151–153). General additive model regression lines were generated using the *geom_smooth* function of the R *ggplot2* package.

Acknowledgments

The authors would like to thank Dr. Pawel Tacikowski, Dr. Antonella Maselli, Dr. Luc Selen, and Dr. Maximilian Hauser for fruitful discussions. All authors were supported by the Swedish Research Council (VR Starting Grant 2019-01909 to K.K.). Experiment costs were covered by the same project and the Åke Wibergs Foundation (M20-0038 granted to K.K.). X.J. received funding from the European Union's Horizon Europe research and innovation program under the Marie Sklodowska-Curie grant agreement number 101059348.

Author Contributions

Noa Cemeljic: Conceptualization, Data curation, Formal Analysis, Investigation, Software, Visualization, Writing – original draft, Writing – review & editing. **Xavier Job**: Conceptualization, Writing – review & editing. **Konstantina Kilteni:** Conceptualization, Data curation, Formal Analysis, Funding acquisition, Methodology, Software, Visualization, Supervision, Writing – original draft, Writing – review & editing.

Declarations of interests

The authors declare no competing interests.

References

1. Myowa-Yamakoshi M, Takeshita H. Do human fetuses anticipate self-oriented actions? A study by four-dimensional (4D) ultrasonography. Infancy. 2006;10(3):289–301.

2. Kurjak A, Azumendi G, Veček N, Kupešic S, Solak M, Varga D, et al. Fetal hand movements and facial expression in normal pregnancy studied by four-dimensional sonography. J Perinat Med. 2003;31(6).

3. Reese A. Patterns of Self-Directed Touch over the First 18 Months of Life - ProQuest [Internet]. 2023 [cited 2023 Dec 11]. Available from:

https://www.proquest.com/docview/2822959179?pq-origsite=gscholar&fromopenview=true&sourcetype=Dissertations%20&%20Theses

4. Thomas BL, Karl JM, Whishaw IQ. Independent development of the Reach and the Grasp in spontaneous self-touching by human infants in the first 6 months. Front Psychol. 2014;5(OCT):1526.

5. DiMercurio A, Connell JP, Clark M, Corbetta D. A naturalistic observation of spontaneous touches to the body and environment in the first 2 months of life. Front Psychol. 2018 Dec 18;9(DEC):2613.

6. Spille JL, Grunwald M, Martin S, Mueller SM. Stop touching your face! A systematic review of triggers, characteristics, regulatory functions and neuro-physiology of facial self touch. Vol. 128, Neuroscience and Biobehavioral Reviews. Elsevier Ltd; 2021. p. 102–16.

7. Bolanowski SJ, Verrillo RT, McGlone Francis. Passive, active and intra-active (self) touch. Somatosens Mot Res. 1999;16(4):304–11.

8. Schütz-Bosbach S, Musil JJ, Haggard P. Touchant-touché: The role of self-touch in the representation of body structure. Conscious Cogn [Internet]. 2009;18(1):2–11. Available from: http://dx.doi.org/10.1016/j.concog.2008.08.003

9. Ehrsson HH, Holmes NP, Passingham RE. Touching a rubber hand: Feeling of body ownership is associated with activity in multisensory brain areas. Journal of Neuroscience. 2005 Nov 9;25(45):10564–73.

10. Rochat P, Hespos SJ. Differential Rooting Response by Neonates: Evidence for an Early Sense of Self. Infant Child Dev. 1997;6(3–4):105–12.

11. Kawato M, Furukawa K, Suzuki R. A hierarchical neural-network model for control and learning of voluntary movement. Biol Cybern. 1987 Oct;57(3):169–85.

12. Miall RC, Wolpert DM. Forward Models for Physiological Motor Control. Neural Networks. 1996 Nov;9(8):1265–79.

13. Franklin DW, Wolpert DM. Computational Mechanisms of Sensorimotor Control. Neuron. 2011 Nov;72(3):425–42.

14. Jordan M. Forward models: Supervised learning with a distal teacher. Cogn Sci. 1992 Sep;16(3):307–54.

15. McNamee D, Wolpert DM. Internal Models in Biological Control. Annu Rev Control Robot Auton Syst. 2019;2(1):339–64.

16. Kilteni K. Methods of Somatosensory Attenuation. In: Holmes NP, editor. Somatosensory Research Methods [Internet]. New York, NY: Springer US; 2023. p. 35–53. Available from: https://doi.org/10.1007/978-1-0716-3068-6_2

17. Bays PM, Wolpert DM. Predictive attenuation in the perception of touch. In: Haggard EP, Rosetti Y, Kawato M, editors. Sensorimotor Foundations of Higher Cognition. Oxford University Press; 2008. p. 339–58.

788   18.   Bays PM, Wolpert DM. Computational principles of sensorimotor control that
789   minimize uncertainty and variability. J Physiol. 2007;578(Pt 2):387–96.

790   19.   Wolpert DM, Flanagan JR. Motor prediction. Current Biology [Internet]. 2001
791   Sep;11(18):R729–32. Available from:
792   https://linkinghub.elsevier.com/retrieve/pii/S0960982201004328

793   20.   Blakemore SJ, Wolpert D, Frith C. Why can't you tickle yourself? Neuroreport
794   [Internet]. 2000 Aug;11(11):R11–6. Available from:
795   http://journals.lww.com/00001756-200008030-00002

796   21.   Vale N, Tomic I, Girones Z, Wolpert DM, Kilteni K, Bays P. Divisive attenuation
797   based on noisy sensorimotor predictions accounts for excess variability in self-touch.
798   bioRxiv [Internet]. 2024 Jan 1;2024.06.20.599826. Available from:
799   http://biorxiv.org/content/early/2024/06/24/2024.06.20.599826.abstract

800   22.   Shergill SS, Bays PM, Frith CD, Wolpert DM. Two Eyes for an Eye: The Neuroscience
801   of Force Escalation. Science (1979). 2003 Jul 11;301(5630):187–187.

802   23.   Kilteni K, Andersson BJ, Houborg C, Ehrsson HH. Motor imagery involves predicting
803   the sensory consequences of the imagined movement. Nat Commun. 2018 Dec
804   24;9(1):1617.

805   24.   Lalouni M, Fust J, Vadenmark-Lundqvist V, Ehrsson HH, Kilteni K, Birgitta Jensen K.
806   Predicting pain: differential pain thresholds during self-induced, externally induced,
807   and imagined self-induced pressure pain. Pain. 2021 May;162(5):1539–44.

808   25.   Weiskrantz L, Elliott J, Darlington C. Preliminary Observations on Tickling Oneself.
809   Nature. 1971 Apr;230(5296):598–9.

810   26.   Blakemore SJ, Frith CD, Wolpert DM. Spatio-temporal prediction modulates the
811   perception of self-produced stimuli. J Cogn Neurosci. 1999 Sep;11(5):551–9.

812   27.   Bays PM, Wolpert DM, Flanagan JR. Perception of the consequences of self-action is
813   temporally tuned and event driven. Current Biology. 2005;15:1125–8.

814   28.   Bays PM, Flanagan JR, Wolpert DM. Attenuation of self-generated tactile sensations is
815   predictive, not postdictive. PLoS Biol. 2006;4(2):281–4.

816   29.   Asimakidou E, Job X, Kilteni K. The positive dimension of schizotypy is associated
817   with a reduced attenuation and precision of self-generated touch. Schizophrenia. 2022
818   Dec 1;8(1):57.

819   30.   Job X, Kilteni K. Action does not enhance but attenuates predicted touch. Elife
820   [Internet]. 2023 Dec 15;12. Available from: https://elifesciences.org/articles/90912

821   31.   Timar L, Job X, Orban de Xivry JJ, Kilteni K. Aging exerts a limited influence on the
822   perception of self-generated and externally generated touch. J Neurophysiol. 2023 Oct
823   1;130:871–82.

824   32.   Kilteni K, Houborg C, Ehrsson HH. Rapid learning and unlearning of predicted
825   sensory delays in self-generated touch. Elife. 2019 Nov;8:e42888.

33. Kilteni K, Engeler P, Boberg I, Maurex L, Ehrsson HH. No evidence for
somatosensory attenuation during action observation of self-touch. European Journal
of Neuroscience. 2021;54(7):6422–44.

34. Kilteni K, Engeler P, Ehrsson HH. Efference Copy Is Necessary for the Attenuation of
Self-Generated Touch. iScience [Internet]. 2020 Feb;23(2):100843. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S2589004220300262

35. Kilteni K, Ehrsson HH. Predictive attenuation of touch and tactile gating are distinct
perceptual phenomena. iScience [Internet]. 2022 Apr 15 [cited 2022 Mar
31];25(4):104077. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S2589004222003479

36. Wolpe N, Zhang J, Nombela C, Ingram JN, Wolpert DM, Rowe JB. Sensory
attenuation in Parkinson's disease is related to disease severity and dopamine dose. Sci
Rep [Internet]. 2018 Dec 23;8(1):15643. Available from:
http://www.nature.com/articles/s41598-018-33678-3

37. Wolpe N, Ingram JN, Tsvetanov KA, Geerligs L, Kievit RA, Henson RN, et al. Ageing
increases reliance on sensorimotor prediction through structural and functional
differences in frontostriatal circuits. Nat Commun [Internet]. 2016 Dec 3;7(1):13034.
Available from: http://www.nature.com/articles/ncomms13034

38. Kilteni K, Ehrsson HH. Sensorimotor predictions and tool use: Hand-held tools
attenuate self-touch. Cognition [Internet]. 2017 [cited 2017 May 9];165:1–9. Available
from: http://www.sciencedirect.com/science/article/pii/S0010027717300999

39. Kilteni K, Ehrsson HH. Body ownership determines the attenuation of self-generated
tactile sensations. Proceedings of the National Academy of Sciences [Internet]. 2017
Aug 1;114(31):8426–31. Available from:
http://www.pnas.org/content/early/2017/07/11/1703347114.full

40. Blakemore SJ, Wolpert DM, Frith CD. Central cancellation of self-produced tickle
sensation. Nat Neurosci. 1998 Nov;1(7):635–40.

41. Blakemore SJ, Wolpert DM, Frith CD. The Cerebellum Contributes to Somatosensory
Cortical Activity during Self-Produced Tactile Stimulation. Neuroimage. 1999
Oct;10(4):448–59.

42. Blakemore SJ, Frith CD, Wolpert DM. The cerebellum is involved in predicting the
sensory consequences of action. Neuroreport. 2001 Jul;12(9):1879–84.

43. Hesse MD, Nishitani N, Fink GR, Jousmaki V, Hari R. Attenuation of Somatosensory
Responses to Self-Produced Tactile Stimulation. Cerebral Cortex. 2010 Feb
1;20(2):425–32.

44. Shergill SS, White TP, Joyce DW, Bays PM, Wolpert DM, Frith CD. Modulation of
somatosensory processing by action. Neuroimage. 2013 Apr;70:356–62.

45. Kilteni K, Ehrsson HH. Functional Connectivity between the Cerebellum and
Somatosensory Areas Implements the Attenuation of Self-Generated Touch. The

Journal of Neuroscience [Internet]. 2020 Jan 22;40(4):894–906. Available from:
http://www.jneurosci.org/content/40/4/894.abstract

46. Kilteni K, Houborg C, Ehrsson HH. Brief temporal perturbations in somatosensory
reafference disrupt perceptual and neural attenuation and increase supplementary
motor area–cerebellar connectivity. The Journal of Neuroscience [Internet]. 2023 Jun
20;JN-RM-1743-22. Available from:
http://www.jneurosci.org/content/early/2023/06/16/JNEUROSCI.1743-
22.2023.abstract

47. Gertz H, Voudouris D, Fiehler K. Reach-relevant somatosensory signals modulate
tactile suppression. J Neurophysiol. 2017;117(6):2262–8.

48. Voudouris D, Fiehler K. Spatial specificity of tactile enhancement during reaching.
Atten Percept Psychophys. 2017;79(8):2424–34.

49. Voudouris D, Fiehler K. Enhancement and suppression of tactile signals during
reaching. J Exp Psychol Hum Percept Perform. 2017;43(6):1238.

50. Williams SR, Shenasa J, Chapman CE. Time course and magnitude of movement-
related gating of tactile detection in humans. I. Importance of stimulus location. J
Neurophysiol. 1998;79(2):947–63.

51. Williams SR, Chapman CE. Time course and magnitude of movement-related gating
of tactile detection in humans. III. Effect of motor tasks. J Neurophysiol.
2002;88(4):1968–79.

52. Chapman CE. Active versus passive touch: Factors influencing the transmission of
somatosensory signals to primary sornatosensory cortex. Can J Physiol Pharmacol.
1994;72(5):558–70.

53. Chapman CE, Beauchamp E. Differential controls over tactile detection in humans by
motor commands and peripheral reafference. J Neurophysiol. 2006;96(3):1664–75.

54. Chapman CE, Bushnell MC, Miron D, Duncan GH, Lund JP. Sensory perception
during movement in man. Exp Brain Res. 1987;68(3):516–24.

55. Voss M, Ingram JN, Wolpert DM, Haggard P. Mere expectation to move causes
attenuation of sensory signals. PLoS One. 2008;3(8):2–6.

56. Arikan BE, Voudouris D, Straube B, Fiehler K. Distinct role of central predictive
mechanisms in tactile suppression. Available at SSRN 4622696. 2023;

57. Walsh LD, Taylor JL, Gandevia SC. Overestimation of force during matching of
externally generated forces. J Physiol. 2011 Feb;589(3):547–57.

58. Voudouris D, Fiehler K. Dynamic temporal modulation of somatosensory processing
during reaching. Sci Rep [Internet]. 2021;11(1):1–12. Available from:
https://doi.org/10.1038/s41598-021-81156-0

59. Kilteni K, Ehrsson HH. Dynamic changes in somatosensory and cerebellar activity
mediate temporal recalibration of self-touch. Commun Biol [Internet]. 2024;7(1):522.
Available from: https://doi.org/10.1038/s42003-024-06188-4

60. Desmurget M, Grafton S. Forward modeling allows feedback control for fast reaching
movements. Trends Cogn Sci. 2000;4(11):423–31.

61. Brenner E, Smeets JBJ. Continuously updating one's predictions underlies successful
interception. REVIEW Control of Movement J Neurophysiol [Internet].
2018;120:3257–74. Available from: www.jn.org

62. Shadmehr R, Krakauer JW, Neuroanatomy AC, Motor FOR. A computational
neuroanatomy for motor control. Exp Brain Res. 2008;185(3):359–81.

63. Shadmehr R, Smith M a, Krakauer JW. Error correction, sensory prediction, and
adaptation in motor control. Annu Rev Neurosci [Internet]. 2010 Jan [cited 2014 Jul
9];33:89–108. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20367317

64. Blakemore SJ, Wolpert DM, Frith CD. Abnormalities in the awareness of action.
Trends Cogn Sci. 2002;6(6):237–242.

65. Frith CD, Blakemore SJ, Wolpert DM. Explaining the symptoms of schizophrenia:
Abnormalities in the awareness of action. Brain Res Rev [Internet]. 2000 Mar;31(2–
3):357–63. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0165017399000521

66. Leptourgos P, Corlett PR. Embodied Predictions, Agency, and Psychosis. Front Big
Data. 2020;3(August):1–13.

67. Corlett PR, Horga G, Fletcher PC, Alderson-Day B, Schmack K, Powers AR.
Hallucinations and Strong Priors. Vol. 23, Trends in Cognitive Sciences. 2019. p. 114–
27.

68. McCrea PH, Eng JJ, Hodgson AJ. Biomechanics of reaching: clinical implications for
individuals with acquired brain injury. Disabil Rehabil. 2002 Jan 7;24(10):534–41.

69. Deng CL, Geng P, Hu YF, Kuai SG. Beyond Fitts's Law: A Three-Phase Model
Predicts Movement Time to Position an Object in an Immersive 3D Virtual
Environment. Human Factors: The Journal of the Human Factors and Ergonomics
Society. 2019 Sep 26;61(6):879–94.

70. Blakemore SJ, Smith J, Steel R, Johnstone EC, Frith CD. The perception of self-
produced sensory stimuli in patients with auditory hallucinations and passivity
experiences: evidence for a breakdown in self-monitoring. Psychol Med.
2000;30(5):1131–9.

71. Frith C. The neural basis of hallucinations and delusions. C R Biol [Internet].
2005;328(2):169–75. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15771003

72. Frith CD. Can a Problem With Corollary Discharge Explain the Symptoms of
Schizophrenia? Biol Psychiatry Cogn Neurosci Neuroimaging [Internet].
2019;4(9):768–9. Available from: https://doi.org/10.1016/j.bpsc.2019.07.003

73. Shergill SS, Samson G, Bays PM, Frith CD, Wolpert DM. Evidence for sensory
prediction deficits in schizophrenia. American Journal of Psychiatry.
2005;162(12):2384–6.

74. Shergill SS, White TP, Joyce DW, Bays PM, Wolpert DM, Frith CD. Functional Magnetic Resonance Imaging of Impaired Sensory Prediction in Schizophrenia. JAMA Psychiatry [Internet]. 2014 Jan 1;71(1):28. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24196370

75. Frith C. The self in action: lessons from delusions of control. Conscious Cogn [Internet]. 2005 Dec [cited 2014 Sep 30];14(4):752–70. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16098765

76. Parees I, Brown H, Nuruki a., Adams R a., Davare M, Bhatia KP, et al. Loss of sensory attenuation in patients with functional (psychogenic) movement disorders. Brain. 2014;137:2916–21.

77. Kilteni K, Ehrsson HH. Dynamic changes in somatosensory and cerebellar activity mediate temporal recalibration of self-touch. bioRxiv [Internet]. 2023 Jan 1;2023.12.01.569520. Available from: http://biorxiv.org/content/early/2023/12/04/2023.12.01.569520.abstract

78. Faisal AA, Selen LPJ, Wolpert DM. Noise in the nervous system. Nat Rev Neurosci. 2008 Apr;9(4):292–303.

79. Jones PW, Gabbiani F. Impact of neural noise on a sensory-motor pathway signaling impending collision. J Neurophysiol. 2012 Feb 15;107(4):1067–79.

80. Harris CM, Wolpert DM. Signal-dependent noise determines motor planning. Nature. 1998 Aug;394(6695):780–4.

81. Todorov E, Jordan MI. Optimal feedback control as a theory of motor coordination. Nat Neurosci. 2002 Nov;5(11):1226–35.

82. Thorp EB, Kording KP, Mussa-Ivaldi FA. Using noise to shape motor learning. J Neurophysiol. 2017 Feb 1;117(2):728–37.

83. Imamizu H, Sugimoto N, Osu R, Tsutsui K, Sugiyama K, Wada Y, et al. Explicit contextual information selectively contributes to predictive switching of internal models. Exp Brain Res. 2007 Aug;181(3):395–408.

84. Haruno M, Wolpert DM, Kawato M. MOSAIC Model for Sensorimotor Learning and Control. Neural Comput. 2001;13(10):2201–20.

85. Blakemore SJ, Goodbody SJ, Wolpert DM. Predicting the consequences of our own actions: the role of sensorimotor context estimation. J Neurosci. 1998;18(18):7511–8.

86. Fraser LE, Fiehler K. Predicted reach consequences drive time course of tactile suppression. Behavioural Brain Research [Internet]. 2018;350(December 2017):54–64. Available from: https://doi.org/10.1016/j.bbr.2018.05.010

87. Bhattacharjee A, Ye AJ, Lisak JA, Vargas MG, Goldreich D. Vibrotactile masking experiments reveal accelerated somatosensory processing in congenitally blind braille readers. Journal of Neuroscience. 2010 Oct 27;30(43):14288–98.

88. Craig JC, Evans PM. Vibrotactile masking and the persistence of tactual features. Vol. 42, Perception & Psychophysics. 1987.

89. Coquery JM, Amblard B. Backward and forward masking in the perception of
cutaneous stimuli. Vol. 13, Perception & Psychophysics. 1973.

90. Makous JC, Gescheider GA, Bolanowski SJ. Decay in the effect of vibrotactile
masking. J Acoust Soc Am. 1996;99(2):1124–9.

91. Press C, Kok P, Yon D. The Perceptual Prediction Paradox. Trends Cogn Sci [Internet].
2020;24(1):13–24. Available from: https://doi.org/10.1016/j.tics.2019.11.003

92. Thomas ER, Yon D, de Lange FP, Press C. Action Enhances Predicted Touch. Psychol
Sci. 2021 Dec;

93. Angel RW, Malenka RC. Velocity-dependent suppression of cutaneous sensitivity
during movement. Exp Neurol. 1982;77(2):266–74.

94. Post LJ, Zompa IC, Chapman CE. Perception of vibrotactile stimuli during motor
activity in human subjects. Exp Brain Res. 1994;100(1):107–20.

95. Voudouris D, Broda MD, Fiehler K. Anticipatory grasping control modulates
somatosensory perception. J Vis. 2019;19(5):1–10.

96. Cybulska-Klosowicz A, Meftah EM, Raby M, Lemieux ML, Chapman CE. A critical
speed for gating of tactile detection during voluntary movement. Exp Brain Res.
2011;210(2):291–301.

97. Williams SR, Chapman CE. Time course and magnitude of movement-related gating
of tactile detection in humans. II. Effects of stimulus intensity. J Neurophysiol.
2000;84(2):863–75.

98. Papakostopoulos D, Cooper R, Crow HJ. Inhibition of cortical evoked potentials and
sensation by self-initiated movement in man. Nature. 1975;258(5533):321–4.

99. Colino FL, Binsted G. Time Course of Tactile Gating in a Reach-to-Grasp and Lift
Task. J Mot Behav. 2016;48(5):390–400.

100. Juravle G, Deubel H, Tan HZ, Spence C. Changes in tactile sensitivity over the time-
course of a goal-directed movement. Behavioural Brain Research [Internet].
2010;208(2):391–401. Available from: http://dx.doi.org/10.1016/j.bbr.2009.12.009

101. Colino FL, Buckingham G, Cheng DT, van Donkelaar P, Binsted G. Tactile gating in a
reaching and grasping task. Physiol Rep. 2014;2(3):1–11.

102. Colino FL, Lee JH, Binsted G. Availability of vision and tactile gating: vision
enhances tactile sensitivity. Exp Brain Res. 2017;235(1):341–8.

103. Rushton DN, Roghwell JC, Craggs MD. Gating of somatosensory evoked potentials
during different kinds of movement in man. Brain. 1981;104(3):465–91.

104. Cohen LG, Starr A. Localization, timing and specificity of gating of somatosensory
evoked potentials during active movement in man. Brain. 1987;110(2):451–67.

105. Pertovaara A, Kemppainen P, Leppänen H. Lowered cutaneous sensitivity to
nonpainful electrical stimulation during isometric exercise in humans. Exp Brain Res.
1992;89(2):447–52.

106. Reznik D, Henkin Y, Levy O, Mukamel R. Perceived Loudness of Self-Generated Sounds Is Differentially Modified by Expected Sound Intensity. PLoS One. 2015 May 18;10(5):e0127651.

107. Paraskevoudi N, SanMiguel I. Self-generation and sound intensity interactively modulate perceptual bias, but not perceptual sensitivity. Sci Rep. 2021 Dec 24;11(1):17103.

108. Juravle G, Colino FL, Meleqi X, Binsted G, Farnè A. Vision facilitates tactile perception when grasping an object. Sci Rep. 2018 Dec;8(1):15653.

109. McManus M, Schütz I, Voudouris D, Fiehler K. How visuomotor predictability and task demands affect tactile sensitivity on a moving limb during object interaction in a virtual environment. R Soc Open Sci. 2023;10(12):231259.

110. Knoetsch F, Zimmermann E. The spatial specificity of sensory attenuation for self-touch. Conscious Cogn. 2021 Jul;92:103135.

111. Kemp KC, Bathery AJ, Barrantes-Vidal N, Kwapil TR. Positive, Negative, and Disorganized Schizotypy Predict Differential Patterns of Interview-Rated Schizophrenia-Spectrum Symptoms and Impairment. Assessment. 2021 Jan 19;28(1):141–52.

112. Barrantes-Vidal N, Grant P, Kwapil TR. The Role of Schizotypy in the Study of the Etiology of Schizophrenia Spectrum Disorders. Schizophr Bull. 2015 Mar 1;41(suppl 2):S408–16.

113. Ford JM, Mathalon DH, Heinks T, Kalba S, Faustman WO, Roth WT. Neurophysiological evidence of corollary discharge dysfunction in schizophrenia. American Journal of Psychiatry. 2001;158(12):2069–71.

114. Ford JM, Gray M, Faustman WO, Roach BJ, Mathalon DH. Dissecting corollary discharge dysfunction in schizophrenia. Psychophysiology. 2007;44(4):522–9.

115. Ford JM, Mathalon DH, Roach BJ, Keedy SK, Reilly JL, Gershon ES, et al. Neurophysiological evidence of corollary discharge function during vocalization in psychotic patients and their nonpsychotic first-degree relatives. Schizophr Bull. 2013 Nov;39(6):1272–80.

116. Rummell BP, Bikas S, Babl SS, Gogos JA, Sigurdsson T. Altered corollary discharge signaling in the auditory cortex of a mouse model of schizophrenia predisposition. Nat Commun. 2023 Dec 1;14(1):7388.

117. Thakkar KN, Diwadkar VA, Rolfs M. Oculomotor Prediction: A Window into the Psychotic Mind. Vol. 21, Trends in Cognitive Sciences. Elsevier Ltd; 2017. p. 344–56.

118. Fletcher PC, Frith CD. Perceiving is believing: a Bayesian approach to explaining the positive symptoms of schizophrenia. Nat Rev Neurosci [Internet]. 2009 Jan 3;10(1):48–58. Available from: http://www.nature.com/articles/nrn2536

119. Thakkar KN, Mathalon DH, Ford JM. Reconciling competing mechanisms posited to underlie auditory verbal hallucinations: Mechanisms of Auditory Hallucinations.

Philosophical Transactions of the Royal Society B: Biological Sciences. 2021 Feb
1;376(1817):20190702.

120. Whitford TJ, Mathalon DH, Shenton ME, Roach BJ, Bammer R, Adcock RA, et al.
Electrophysiological and diffusion tensor imaging evidence of delayed corollary
discharges in patients with schizophrenia. Psychol Med. 2011;41(5):959–69.

121. Whitford TJ, Ford JM, Mathalon DH, Kubicki M, Shenton ME. Schizophrenia,
myelination, and delayed corollary discharges: A hypothesis. Schizophr Bull.
2012;38(3):486–94.

122. Okimura T, Maeda T, Mimura M, Yamashita Y. Aberrant sense of agency induced by
delayed prediction signals in schizophrenia: a computational modeling study.
Schizophrenia. 2023 Dec 1;9(1):72.

123. Haß K, Sinke C, Reese T, Roy M, Wiswede D, Dillo W, et al. Enlarged temporal
integration window in schizophrenia indicated by the double-flash illusion. Cogn
Neuropsychiatry. 2017 Mar 4;22(2):145–58.

124. Stevenson RA, Park S, Cochran C, McIntosh LG, Noel JP, Barense MD, et al. The
associations between multisensory temporal processing and symptoms of
schizophrenia. Schizophr Res. 2017 Jan;179:97–103.

125. Foucher JR, Lacambre M, Pham BT, Giersch A, Elliott MA. Low time resolution in
schizophrenia. Schizophr Res. 2007 Dec;97(1–3):118–27.

126. Ferri F, Venskus A, Fotia F, Cooke J, Romei V. Higher proneness to multisensory
illusions is driven by reduced temporal sensitivity in people with high schizotypal
traits. Conscious Cogn. 2018 Oct;65:263–70.

127. Fotia F, Cooke J, Van Dam L, Ferri F, Romei V. The temporal sensitivity to the tactile-
induced double flash illusion mediates the impact of beta oscillations on schizotypal
personality traits. Conscious Cogn. 2021 May;91:103121.

128. Marsicano G, Cerpelloni F, Melcher D, Ronconi L. Lower multisensory temporal
acuity in individuals with high schizotypal traits: a web-based study. Sci Rep. 2022
Feb 17;12(1):2782.

129. Ferri F, Nikolova YS, Perrucci MG, Costantini M, Ferretti A, Gatta V, et al. A Neural
"Tuning Curve" for Multisensory Experience and Cognitive-Perceptual Schizotypy.
Schizophr Bull. 2017 Jul;43(4):801–13.

130. Poulet JFA, Hedwig B. Corollary Discharge Inhibition of Ascending Auditory Neurons
in the Stridulating Cricket. The Journal of Neuroscience. 2003 Jun 1;23(11):4717–25.

131. Poulet JFA, Hedwig B. The Cellular Basis of a Corollary Discharge. Science (1979).
2006 Jan 27;311(5760):518–22.

132. Schneider DM, Sundararajan J, Mooney R. A cortical filter that learns to suppress the
acoustic consequences of movement. Nature. 2018 Sep 12;561(7723):391–5.

133. Chinta S, Pluta SR. Neural mechanisms for the localization of unexpected external
motion. Nat Commun. 2023 Sep 30;14(1):6112.

134. Audette NJ, Zhou WX, La Chioma A, Schneider DM. Precise movement-based
predictions in the mouse auditory cortex. Current Biology. 2022 Nov 21;32(22):4925-
4940.e6.

135. Fukutomi M, Carlson BA. A History of Corollary Discharge: Contributions of
Mormyrid Weakly Electric Fish. Front Integr Neurosci. 2020 Jul 29;14.

136. Sawtell NB. Neural Mechanisms for Predicting the Sensory Consequences of
Behavior: Insights from Electrosensory Systems. Annu Rev Physiol. 2017 Feb
10;79(1):381–99.

137. Wallach A, Sawtell NB. An internal model for canceling self-generated sensory input
in freely behaving electric fish. Neuron. 2023 Aug;111(16):2570-2582.e5.

138. Roy JE, Cullen KE. Dissociating Self-Generated from Passively Applied Head Motion:
Neural Mechanisms in the Vestibular Nuclei. The Journal of Neuroscience. 2004 Mar
3;24(9):2102–11.

139. Brooks JX, Carriot J, Cullen KE. Learning to expect the unexpected: rapid updating in
primate cerebellum during voluntary self-motion. Nat Neurosci. 2015 Sep
3;18(9):1310–7.

140. Eliades SJ, Wang X. Neural substrates of vocalization feedback monitoring in primate
auditory cortex. Nature. 2008 Jun 4;453(7198):1102–6.

141. Brooks JX, Cullen KE. Predictive Sensing: The Role of Motor Signals in Sensory
Processing. Biol Psychiatry Cogn Neurosci Neuroimaging. 2019 Sep;4(9):842–50.

142. Crapse TB, Sommer MA. Corollary discharge across the animal kingdom. Nat Rev
Neurosci. 2008 Aug;9(8):587–600.

143. Cullen KE. Sensory signals during active versus passive movement. Curr Opin
Neurobiol. 2004 Dec;14(6):698–706.

144. Cullen KE. The vestibular system: multimodal integration and encoding of self-motion
for motor control. Trends Neurosci. 2012 Mar;35(3):185–96.

145. Schneider DM, Mooney R. How Movement Modulates Hearing. Annu Rev Neurosci.
2018 Jul 8;41(1):553–72.

146. Straka H, Simmers J, Chagnaud BP. A New Perspective on Predictive Motor Signaling.
Current Biology. 2018 Mar;28(5):R232–43.

147. Gupta K, Chowdhury RR, Chakrabarti S, Schwarz C. Discerning state estimation and
sensory gating, two presumptive predictive signals in mouse barrel cortex. bioRxiv.
2023;

148. Oldfield RC. The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia. 1971 Mar;9(1):97–113.

149. R Core Team. R: A Language and Environment for Statistical Computing . Vienna,
Austria; 2022.

150. JASP Team. JASP (Version 0.17). 2023;

151.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
1137   Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series
1138   B (Methodological). 1995 Jan;57(1):289–300.

1139   152.   Quintana DS, Williams DR. Bayesian alternatives for common null-hypothesis
1140   significance tests in psychiatry: A non-technical guide using JASP. BMC Psychiatry.
1141   2018;18(1):1–8.

1142   153.   van Doorn J, van den Bergh D, Böhm U, Dablander F, Derks K, Draws T, et al. The
1143   JASP guidelines for conducting and reporting a Bayesian analysis. Vol. 28,
1144   Psychonomic Bulletin and Review. 2021.

1145

1146