Predictions of bimanual self-touch determine the temporal tuning of somatosensory perception

Noa Cemeljic[1], Xavier Job[1], Konstantina Kilteni[1,2*]

[1]Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden

[2] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

[*]Correspondence and requests for materials should be addressed to konstantina.kilteni@ki.se

Abstract

We effortlessly distinguish between touching ourselves with our hands and being touched by other people or objects. Motor control theories posit that this distinction is made possible by the brain predicting the somatosensory consequences of our voluntary movements based on an 'efference copy', and attenuating our responses to the predicted self-touch. However, it remains unclear how these predictions impact somatosensory perception at times other than during self-touch: for example, as our hand reaches to touch our body or moves away from it. Here participants discriminated forces applied on their left index finger by a motor. The forces were applied during the reaching movement of their right hand towards the left hand, including the time the reaching ended by simulating self-touch between the hands, or after the reaching movement. We observed that the forces on the left hand felt progressively weaker during the reaching phase, reached their minimum perceived intensity at the time of self-touch, and quickly recovered after the end of the reaching. All effects were replicated with a new cohort of participants that further demonstrated that this gradual attenuation of the perceived magnitude of touch vanished during similar right hand reaching movements that did not produce expectations for self-touch between the two hands. Together, our results indicate a temporal tuning of somatosensory perception during movements to self-touch and underscore the role of sensorimotor context in forming predictions that attenuate the intensity of self-generated touch.

Introduction

Self-touch occurs in a wide range of everyday activities, from rubbing our eyes with our fists to alleviate itching, to warming ourselves with our hands during shivering, and affectively stimulating our bodies for pleasure. Self-touch behaviors begin in utero [1,2] and continue after birth as a way for infants to explore their bodies [3–5]. Whether spontaneous (*e.g.*, clenching our fists when nervous) or purposeful (*e.g.*, moving the hair away from our face) [6], self-touch is considered fundamental for building and maintaining a sense of self [7–10] through detecting and maintaining contingencies between our motor (*i.e.*, moving to touch our body) and somatosensory cues (*i.e.*, receiving the corresponding touch).

Besides self-touch, we also experience numerous instances of touch generated by other people or objects. Remarkably, our brain possesses the capacity to distinguish between self-touch and touch of external origin to guide behaviour; for example, we do not react to self-touch between our hands, but we might recoil from similar touch initiated by a stranger, just as we would brush away a fly that lands our hand.

How does the brain implement this distinction? Motor control theories posit that the brain uses internal models to predict the somatosensory consequences of voluntary movements [11–15] based on a copy of the motor command (*i.e.,* efference copy) and attenuates (or cancels) the responses to the self-generated touch [16–21]. Accordingly, when reaching with the (active) right hand to touch the (passive) left hand, the internal model predicts and attenuates the resulting touch on both hands using the efference copy associated with the right hand's reaching movement. In contrast, a touch of external origin cannot be predicted based on the movement and is not attenuated [17,20]. It has been proposed that this attenuation of self-generated touch facilitates the distinction from externally generated touch by increasing the salience of the latter relative to the former [15,16,20]. In agreement with this framework, several studies have shown that self-touch elicits an attenuated perceptual response (*i.e.*, it feels less intense or ticklish) [22–39] and triggers weaker activity in somatosensory cortices and the cerebellum [40–46] than identical touch applied by another person or a machine.

Nevertheless, it remains unclear how the predictions of the internal model about the somatosensory consequences of the movement affect perception at times other than the exact moment of self-touch. For example, imagine reaching with the right hand to touch your left hand (**Figure 1a**). It remains elusive how touch on the left hand is perceived as the reaching movement of the right hand progresses, before contact between the hands occurs, and after the reaching movement has ended. This is due to several reasons. First, several earlier studies assessed somatosensory perception across time on the active hand, rather than the passive hand which served as a target of the active hand's movement (*e.g.*, [47–50]). However, it has been demonstrated that somatosensory perception on a limb that moves can be 'gated' by nonpredictive mechanisms [51–57] (see [35] for discussion). To isolate the effects of the internal model's predictions on somatosensation, other studies have focused on the perception of the passive hand, which is not influenced by non-predictive confounds because it does not move [16,17,35]. Nevertheless, these studies either probed perception only *at the moment* of self-touch [22–24,28–31,33–39,58], overlooking the periods before or after the hands' contact, or otherwise

focused on the time *before* the contact [59], overlooking the periods during or after self-touch.
Third, other studies have introduced temporal delays to test somatosensory perception before
and/or after the predicted self-touch [26,27,32,46,60], but those probed timings were tailored to the
moment of the predicted tactile sensation and did not directly relate to the reaching movement
of the active hand. For example, one can probe somatosensory perception at a fixed time before
the anticipated self-touch (*e.g.*, −300 ms), but that timing could align with the early (*i.e.*,
acceleration) phase of a short reaching movement (*e.g.*, 500 ms) or the late (*i.e.*, deceleration)
phase of a longer reaching movement (*e.g.*, 1500 ms). Related to this point, an earlier study [59]
suggested that somatosensory perception on the passive hand changes only when the active
hand reaches its peak velocity. Finally, in addition to being constrained in scope, the
abovementioned research has produced conflicting and inconclusive results. Specifically, an
earlier study reported a constant increase in the attenuation of touch on the passive hand as the
active hand approached the passive [27], while another study found facilitation, rather than
attenuation, of tactile perception on the passive hand when the active hand reached its peak
velocity [59].

Here, we hypothesized that the predictions of the internal model should continuously influence
somatosensory perception during the reaching movement of the right hand to touch the left
hand. Specifically, we reasoned that as the reaching movement of the right hand progresses,
the brain gains access to the latest sensory information to adjust the movement and re-estimate
the position and velocity of the right hand [15,19,61], as well as to update the predictions of the
internal model about the remainder of the movement [62–64], including the expected timing of
contact with the left hand (*i.e.*, self-touch). Consequently, the prediction about the time of self-
touch should become sharper (*i.e.* narrower probability distribution) at later timings during the
reaching than earlier ones (**Figure 1b**). Therefore, we expected to observe a gradually
increasing attenuation of the perceived magnitude of touch during the reaching, which recovers
after the hands make contact (**Figure 1c**). That is, the attenuation of somatosensory perception
should not be an on-off phenomenon that manifests only at the time of the self-touch, but rather
it should gradually build up over the reaching time.

Figure 1. Schematic illustrations of the experimental hypothesis. (a) We investigated how participants perceive touch on their left hand during and after reaching with their right hand to touch their left hand. The reaching movement is shown from a third person perspective and the top right small window indicates the orientation of the participant. (**b**) From a theoretical perspective, during the reaching the brain estimates the state of the moving hand (*e.g.*, its position, velocity) using the predicted and received sensory input and updates its predictions about the rest of the movement accordingly. Consequently, as the action progresses, the predictions of the internal model about the timing of self-touch become narrower given that there is a better estimation based on the latest sensory input (*e.g.*, visual and proprioceptive information). (**c**) Touch received during the reaching (yellow, red, and blue) including its conclusion (green), or after the reaching (brown) would be attenuated proportionally to its probability of occurring at that time point. (**a, b, c**) Phases of the reaching movement and times are colour-coded. Yellow represents the early reaching phase (acceleration), red indicates the phase around the peak velocity, blue denotes the late reaching phase (deceleration), green corresponds to the moment the reaching movement concludes upon reaching the target (self-touch), and brown refers to the phase after the reaching.

Results

Experiment 1. The perceived magnitude of touch gradually decreases during the reaching movement, attaining its minimum at the moment of the predicted self-touch.

Twenty-nine participants (15 female, 28 right-handed, aged 18 - 34 years) rested their left hand palm up with their left index finger placed inside a moulded support on a desk. Their right hand was placed palm down on top of a wooden surface placed on top of the desk (**Figure 2a, Supplementary Figure S1**). Upon presentation of an auditory GO cue, they performed a reaching movement finger with their active right hand starting from a designated starting position (reaction time, mean $\pm$ SD, 220 ± 42 ms) (**Figure 2a**, *starting position*), and concluded the movement by tapping with their right index finger a force sensor (force duration, mean $\pm$ SD, 129 ± 39 ms) placed above, but not in contact with, their left index finger resting on the desk (**Figure 2a**, *cyan force sensor*).

During or after the reaching movement, the participants received a brief force on their left index finger via a probe attached to the lever of a DC electric motor (**Figure 2a**, *lever*). This *test* force had a fixed amplitude of 2 N. We quantified how strongly participants perceived the intensity of this *test* force over time using a well-established force discrimination task [16,27–35,46]. In each trial, participants received the *test* force (2 N) and a *comparison* force with an intensity that pseudorandomly varied around the intensity of the *test* force (between 1 and 3 N) (**Figure 2b**). The magnitudes of the *test* and *comparison* forces were confirmed by a second identical force sensor placed inside the probe (**Figure 2a**, *red force sensor*). As mentioned above, the *test* force was applied at a different time relative to the reaching movement of the right hand on any given trial. However, the *comparison* force was always applied 1200 ms after the *test* force – that is, when the right hand had finished the reaching and stopped moving. In other words, our experimental manipulation exclusively involved the *test* force, while the *comparison* force

served as a reference. After the presentation of the two forces, participants verbally indicated
which one of the *test* or the *comparison* force felt stronger (**Figure 2b**).

The participants performed a block of 560 reaching trials in which the *test* force was delivered
during one of five different phases relative to the movement of their right hand in randomized
order (**Figure 2c**): (a) the *early* reaching trials included the period between movement onset
until the time the velocity reached 85% of its peak value (early acceleration phase); (b) the *mid*
reaching trials included the period after the velocity reached 85% of its peak value until the
time it dropped below 85% of its peak value (peak velocity phase); (c) the *late* reaching trials
included the period after the velocity dropped below 85% of its peak value until before the
moment the right index finger tapped the force sensor (late deceleration phase); (d) the *target*
trials included the moment when the right index finger tapped the sensor and triggered the
touch on the left hand, thereby simulating bimanual self-touch (end of reaching); and (e) the
post-reach trials included the moment at 300 ms after the tap on the sensor and thus after the
reaching movement. Participants were aware that the *test* force could come at different times
but did not know when they would receive it on every trial. These 5 trial groups were defined
to correspond to different phases of the reaching movement [65,66], and capture the time of peak
velocity, where changes in the somatosensory perception were previously reported [59] and the
post-reach phase where changes are typically minimal [27,32]. On average, participants took
(mean ± SD) 542 ± 46 ms to reach their left hand (within-subjects SD of movement time, 77 ±
20 ms) (**Figure 2d**) with a peak velocity of 95 ± 11 cm/s at 203 ± 31 ms after movement onset
(**Figure 2e**). The average times at which the *test* force was delivered in the *early*, *mid*, and *late*
trials were 81 ± 17 ms, 201 ± 31 ms, and 350 ± 27 ms relative to movement onset,
corresponding to 15%, 37%, and 65% of the average reaching time in the *early*, *mid*, and *late*
trials, respectively (**Figure 2f**) (**Materials and Methods**). There were no significant
differences in the forces participants tapped on the sensor with their right index finger between
any trial type ($F(4, 112) = 1.001$, $p = 0.410$, $\eta_p^2 = 0.035$)), confirming that the participants
behaved comparably across all reaching trials (**Supplementary Figure S2**).

Figure 2. Methods and results of Experiment 1. (a) Participants made a reaching movement with their active right hand from the *starting position* to tap the force sensor (depicted as the cyan sensor) placed above but not in contact with their left passive index finger. Another identical force sensor (depicted as the red sensor) was placed within the probe of the lever controlled by the motor to measure the forces applied on the left index finger. In every trial, a motion tracking sensor (*motion sensor*) recorded the position of the right index finger. The right hand was placed on top of the wooden table (brown surface), which was placed on top of the desk (gray surface). For illustration purposes, the wooden table appears transparent. The reaching movement is shown from a third-person perspective. **(b)** In every trial, the participants received two forces on the pulp of their left index finger – the *test* and the *comparison* force – via the lever of the motor. After receiving both forces, they verbally indicated which force felt stronger. **(c)** Example of a velocity profile of one pilot participant illustrating the different phases for a reaching trial. The various phases were determined using the 3D velocity data, but the horizontal velocity is plotted here for simplicity. In the reaching trials, participants received the *test* force on their left index finger during one of five different phases relative to their

movement: *early* (yellow), *mid* (red), *late* (blue), at the *target* of the movement (green), or 300 ms *post-reach* (brown). The *comparison* force was always delivered 1200 ms after the *test* force and when the right hand had completed the reaching and had stopped moving. In the *baseline* trials, participants kept both hands relaxed and received the *test* and the *comparison* forces at rest. **(d-e)** Individual and group reach durations **(d)** and peak velocities **(e)**. The markers represent individual data, boxplots show the medians and interquartile ranges, and raincloud plots depict the data distributions. **(f)** Histograms of the times at which the *test* force was applied to the participants' left index finger in the different types of reaching trials. Times are expressed as percentages relative to the reaching movement duration (*i.e.*, the time needed to tap the force sensor) and are binned into 5% intervals. The vertical dashed line represents the mean time of peak velocity (v_{max}), and the grey shaded area represents the standard error of the mean (s.e.m.). **(g)** Responses and fitted logistic models for all types of trials for one participant. We extracted the PSE ($p = 0.5$) for each participant and each trial type. **(h)** Individual PSE values are shown for each type of reaching trial after being normalized to the *baseline* PSE. The individual PSEs of the *early*, *mid*, *late*, and *target* trials are plotted against the time the *test* force occurred in each trial type relative to the reach duration (%). In the *post-reach* trials, the PSEs were computed for the forces delivered 300 ms after the end of the reach (the dotted line indicates a discontinuity in the x-axis). In the figure, individual points are jittered (± 1%) to avoid complete overlap. To illustrate the pattern in our data and facilitate the comparison with our hypothesis (**Figure 1c**), we overlaid a regression line based on a generalized additive model. The shaded area represents a 95% confidence interval. **(i)** Normalized group PSE values for each trial type (mean ± s.e.m.). The touch felt significantly weaker as the reaching time progressed (*late* < *early*, *target* < *early*, *target* < *mid*, *target* < *late*) but quickly recovered afterwards (*post-reach* > *target*). Asterisks denote significant pairwise comparisons of interest after corrections for multiple comparisons. The dotted line indicates the discontinuity in the x-axis. **(j)** Normalized individual PSE data during reaching (*early* to *target*). Data are jittered (± 1%) to avoid complete overlap. The PSEs of each participant were fitted with a linear model, and we display the group's linear fit. The shaded area depicts a 95% confidence interval. **(k)** Slope coefficients of the participants' linear models showing the change in PSEs over the reaching. The slopes were overall negative, indicating an increasing attenuation, and significantly different from zero. The markers represent the individual slope coefficients, boxplots show the medians and interquartile ranges, and raincloud plots depict the coefficients' distributions (** $p < 0.01$, *** $p < 0.001$).

The participants performed an additional block of 56 *baseline* trials in which the participants remained still while performing the force discrimination task. These trials were conducted either before or after the block of reaching trials in a fully counterbalanced order and served to quantify the participants' perceived magnitude of the *test* force in the absence of action. For all types of trials (*early*, *mid*, *late*, *target*, *post-reach*, and *baseline*), we fitted the participants' responses from the force-discrimination task with logistic models (see **Supplementary Figure S3** for all individual fits). All fits were very good (McFadden $R^2 > 0.41$). Next, we extracted the point of subjective equality (PSE), which indicates the intensity at which the *test* force felt as strong as the *comparison* force ($p = 0.5$) and represents the perceived intensity of the *test*

force per trial type (**Figure 2g**). A lower PSE in one trial type compared to another indicates
that the *test* force felt weaker in the former than in the latter trial type.

PSE values of the *reaching* trials were first normalized to the PSEs of the *baseline* trials
(PSE$_{reaching}$ − PSE$_{baseline}$) and then subjected to a one-way repeated measures ANOVA with the
type of trial (*early*, *mid*, *late*, *target*, *post-reach*) as the within-subjects factor. The type of trial
was significant: $F(2.09, 58.44) = 15.856, p < 0.001, \eta_p^2 = 0.362$. Planned pairwise comparisons
revealed that the perceived intensity of the *test* force gradually decreased the closer the touch
was delivered to the moment of the predicted self-touch (**Figure 2h-i**): specifically, the PSEs
were significantly lower in the *late* compared to the *early* trials ($n = 29, t(28) = -2.956, p =$
0.009 *FDR-corrected*, $CI^{95} = [-0.090, -0.016], d = -0.549$), in the *target* compared to the *early*
trials ($n = 29, W = 11, p < 0.001$ *FDR-corrected*, $CI^{95} = [-0.234, -0.115], rrb = -0.949$), in the
target compared to the *mid* trials ($n = 29, W = 31, p < 0.001$ *FDR-corrected*, $CI^{95} = [-0.225,$
$-0.097], rrb = -0.857$) and in the *target* compared to the *late* trials ($n = 29, W = 41, p < 0.001$
FDR-corrected, $CI^{95} = [-0.180, -0.064], rrb = -0.811$). Differences in the PSEs between *early*
and *mid* trials ($n = 29, t(28) = 0.829, p = 0.414$ *FDR-corrected*, $CI^{95} = [-0.021, 0.050], d =$
0.154) and between *mid* and *late* trials ($n = 29, t(28) = 1.813, p = 0.094$ *FDR-corrected*, $CI^{95} =$
$[-0.005, 0.083], d = 0.337$) were not significant. The lowest PSEs thus occurred at the time of
contact between the two index fingers (*i.e.*, *target* trials) when the *test* force was perceived to
be significantly weaker than the *test* force at all other probed times. These PSEs quickly
recovered in the *post-reach* trials and were significantly greater than in the *target* trials ($n = 29,$
$W = 412, p < 0.001$ *FDR-corrected*, $CI^{95} = [0.135, 0.268], rrb = 0.894$).

To further corroborate the gradual decrease in PSEs during reaching, we next fitted each
participant's normalized PSEs in *early*, *mid*, *late*, and *target* trials with a linear regression
model (**Figure 2j**). As hypothesized, the extracted slope coefficients were negative (β =
$-2.293×10^{-3} ± 2.494×10^{-3}$) and significantly different from zero ($n = 29, W = 14, p < 0.001,$
$CI^{95} = [-0.003, -0.001], rrb = -0.936$) (**Figure 2k**), indicating that the *test* force was perceived
weaker the closer it was delivered to the expected time of self-touch. Similar results were
obtained when fitting the PSEs from the *early*, *mid*, and *late* reaching trials only.

Taken together, the results of Experiment 1 demonstrated that the touch felt gradually weaker
as the reaching movement progressed, attained its minimum perceived intensity at the moment
of the predicted self-touch, and quickly recovered after the reaching.

Experiment 2. The gradual attenuation of touch during reaching depends on whether the
reaching is expected to produce self-touch.

The results of Experiment 1 can be attributed to a dynamic influence of the internal model's
predictions on somatosensory perception during the reaching movement. To rule out that the
observed PSE effects are due to other factors related to the movement or the task *per se* (for
example, general anticipation of tactile stimulation on the left index finger, unspecific time
effects, divided attention to both hands or feelings of agency over the action), we conducted a
control experiment. In Experiment 2, we hypothesized that if the temporal tuning of
somatosensory perception observed in Experiment 1 stems from the predictions of the internal
model, then performing a reaching movement with the right hand that does not have any tactile

consequences for the left hand should not modulate somatosensory perception in time. This
would occur because, in such a scenario, the internal model would not predict somatosensory
input on the left hand based on the efference copy (**Figure 3a**).

Two new groups of naive participants participated in Experiment 2. Participants in the *Self-*
touch group (*n* = 29, 15 female, 28 right-handed, aged 18-37 years) performed the same task
as those in Experiment 1: they reached with their right index finger to tap the sensor above
their left index finger (reaction time, mean ± SD, 171 ± 38ms) and received forces on their left
index finger at different times with respect to the movement of their right hand (**Figure 3b**). In
contrast, participants in the *No self-touch* group (*n* = 27, 15 female, 26 right-handed, aged 21-
37 years) performed the same reaching movement with their right hand as participants in the
Self-touch group did (reaction time, mean ± SD, 130 ± 35 ms), but they were instructed to never
make contact between their two fingers through the probe (**Figure 3b**). Rather, they approached
a distance sensor placed on top of, but not in contact with, their left index finger which triggered
the force once the position of the right index finger became lower than a pre-set position
threshold. Then they lifted their right hand as the *Self-touch* group, simulating a virtual tap in
the air. The stimuli were delivered at the same time between the two groups, with the *test* force
in the *target* trials being applied upon triggering the distance sensor in the *No self-touch* group
and upon triggering the force sensor in the *Self-touch* group (see **Materials and Methods**).

As in Experiment 1, participants in the *Self-touch* group were told that their hands would make
contact through the probe in some trials, simulating self-touch, but they would not know
beforehand which trials these would be. In contrast, participants in the *No self-touch* group
were instructed to stop their right hand at a certain distance above their left hand and then lift
it without ever making contact between their hands through the probe. Previous studies [27,28,30]
have shown that if participants deliberately stop their right hand movement above their left
hand, the touch applied on the left hand is not attenuated because it cannot be predicted by the
motor command. In contrast, if participants predict based on their movement that they will
make contact between their two hands but unexpectedly miss the contact, robust attenuation is
still observed [28,30] indicating that the major driver of somatosensory attenuation is the
prediction of self-touch based on the efference copy and not the hands' contact *per se*.
Therefore, in Experiment 2, we expected to replicate all the effects of Experiment 1 with the
Self-touch group and further observe no temporal modulation in the *No self-touch* group
because the internal model would no longer predict touch on the left hand.

The two groups performed reaching movements of comparable durations (**Figure 3c**). The *No-*
self touch group exhibited slightly higher peak velocity than the *Self-touch* group (**Figure 3d**),
but both groups received the *test* forces at similar times (**Figure 3e-f**, **Supplementary Text**
S1). As in Experiment 1, the *Self-touch* group pressed comparable forces across the reaching
trials (**Supplementary Figure S4)** and all psychometric fits of both groups were very good
(McFadden R^2 > 0.47) (**Supplementary Figures S5-S6**). Importantly, the *Self-touch* group
replicated all effects of Experiment 1 (**Supplementary Text S2, Figure S7**).

To compare the somatosensory perception of the two groups, we inserted the participants'
normalized PSEs in a mixed ANOVA with the group as a between-subjects factor and the type
of trial as a within-subjects factor. The model revealed a significant main effect of the trial type

($F(3.27, 176.33) = 10.044$, $p < 0.001$, $\eta_p^2 = 0.157$), a significant main effect of group ($F(1, 54)$
$= 8.284$, $p = 0.006$, $\eta_p^2 = 0.133$) and a significant group-by-trial type interaction ($F(3.27,$
$176.33) = 7.072$, $p < 0.001$, $\eta_p^2 = 0.116$). Pairwise comparisons revealed that the PSEs were
significantly lower in the *Self-touch* group than in the *No self-touch* group, in the *mid* ($t(53.89)$
$= -2.206$, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.156, -0.007]$, $d = -0.590$), *late* ($t(53.37) =$
-2.488, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.171, -0.018]$, $d = -0.663$), *target* ($W = 129$, $p <$
0.001 *FDR-corrected*, $CI^{95} = [-0.286, -0.123]$, $rrb = -0.670$), and *post-reach* trials ($t(53.97) =$
-2.299, $p = 0.040$ *FDR-corrected*, $CI^{95} = [-0.162, -0.011]$, $d = -0.614$). No difference was
detected in the *early* trials ($t(51.84) = -0.338$, $p = 0.737$ *FDR-corrected*, $CI^{95} = [-0.105, 0.075]$,
$d = -0.091$) (**Figure 3g**). In other words, although both groups perceived a comparable
magnitude of touch in the early phase of the reaching, their perception deviated at later timings
depending on whether the reaching movement had somatosensory consequences for the left
hand or not. Further control analyses indicated that the observed PSE differences between the
two groups were not due to differences in the kinematics or task demands of the reaching
movements (**Supplementary Text S3, Figure S8**).

Next, we fitted linear models to the PSEs during reaching (*early*, *mid*, *late*, and *target* trials)
and extracted the slope coefficients for every participant in each group. While the slope
coefficients from the *Self-touch* group were negative ($\beta = -1.431 \times 10^{-3} \pm 1.622 \times 10^{-3}$) and
significantly different from zero ($n = 29$, $t(28) = -4.750$, $p < 0.001$, $CI^{95} = [-0.002, -8.138 \times 10^{-4}]$
$^4]$, $d = -0.882$), replicating Experiment 1, the slope coefficients from the *No self-touch* group
were positive ($\beta = 5.274 \times 10^{-4} \pm 1.596 \times 10^{-3}$) and did not significantly differ from zero ($n = 27$,
$t(26) = 1.717$, $p = 0.098$, $CI^{95} = [-1.041 \times 10^{-4}, 0.001]$, $d = 0.330$). As expected, the coefficients
were significantly lower in the *Self-touch group* compared to the *No self-touch* group ($t(53.83)$
$= -4.551$, $p < 0.001$, $CI^{95} = [-0.003, -0.001]$, $d = -1.217$) (**Figure 3h-i**). Similar results were
obtained when fitting the PSEs from the *early*, *mid*, and *late* reaching trials only, suggesting
that the results are not solely driven by attenuation differences in the *target* trials. Together, the
results of Experiment 2 indicate that the gradual attenuation of the perceived magnitude of
touch on the left hand depends on whether the reaching movement of the right hand has
somatosensory consequences for the left hand (*i.e.*, self-touch).

Figure 3. Methods and results of Experiment 2. (a) When reaching with the active right hand towards the passive left hand but without planning to touch it (*i.e.*, planning to stop the movement before contact), the brain does not predict somatosensory consequences on the left hand based on the efference copy of the right hand. Consequently, somatosensory perception on the left hand should not show any modulation over time. **(b)** As in Experiment 1, participants in the *Self-touch* group made a reaching movement with their right hand from the *starting position* to tap the force sensor (cyan sensor) placed above, but not in contact with, their left index finger. Participants in the *No self-touch* group (right) also made a reaching movement with their right hand from the *starting position* to approach a distance sensor (yellow sensor) placed above, but not in contact with, their left index finger. The reaching movement is shown from a third-person perspective. **(c-d)** The durations of reaching movements were comparable

between the two groups **(c)**. The *Self-touch* group exhibited a lower peak velocity than the *No self-touch* group **(d)**. **(e-f)** Histograms of the times at which the *test* force was applied in the different types of reaching trials for the **(e)** *Self-touch* group and **(f)** *No self-touch* group. The times did not significantly differ between the two groups. Times are expressed as percentages relative to the reaching movement duration and are binned into 5% intervals. The vertical dashed line represents the mean time of peak velocity (v_{max}), and the grey shaded area represents the standard error of the mean (s.e.m.). **(g)** PSEs were significantly lower in the *mid*, *late*, *target*, and *post-reach* trials when self-touch was expected (*Self-touch* group) than when it was not predicted (*No Self-touch* group). Asterisks denote between-group comparisons of interest after corrections for multiple comparisons. The dotted line indicates the discontinuity in the x-axis. **(h)** Individual normalized PSE data during the reaching movement only (*early* to *target*) are plotted for each group. Individual points are jittered ($\pm$ 1%) to avoid complete overlap. We fitted a linear model for every participant in every group separately, but for illustration purposes, we display the group fits. The shaded area depicts a 95% confidence interval. **(i)** In the *Self-touch* group the slopes were overall negative and significantly lower than the slopes in the *No self-touch*, which did not differ from zero. The markers represent the slope value for each participant, boxplots show the medians and interquartile ranges, and raincloud plots depict the data distributions. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, *n.s.* nonsignificant)

Discussion

Determining the time course of somatosensory perception during movements to self-touch is essential for understanding how the brain's predictions influence perception over time [15,19,63]. By combining kinematic recordings with somatosensory psychophysics, the present study tested the time course of somatosensory perception during a movement to self-touch; that is, how participants perceived touch on their left hand as their right hand reached to touch the left hand, and moved away from it. Going beyond earlier research that did not test perception during the entire movement of the active hand [22–24,28–31,33–39,45,58,59] or tested perception at times that were not linked to the reaching movement of the active hand [26,27,32,46,67], our study revealed the timeline of somatosensory perception on the passive left hand during the different phases of the active right hand's movement to touch it. In Experiment 1, we demonstrated that touch on the left hand was perceived with gradually attenuated intensity as the reaching of the right hand progressed: the touch felt significantly weaker in the late reaching phase (*i.e.,* deceleration) than in the early reaching phase (*i.e.,* acceleration) and was significantly more attenuated at the moment of self-touch. The minimum perceived intensity was observed at the time of self-touch between the hands, and it quickly recovered after the reaching had ended.

All these results were replicated in Experiment 2, where participants in the *Self-touch* group exhibited the same temporal tuning of somatosensory responses. In contrast, participants in the *No self-touch* group who performed similar right hand movements that had no somatosensory consequences for the left hand showed no such modulation. Complementary analysis in that group (**Supplementary Text S2**) showed that the perceived intensity of touch did not change significantly at any probed time during the right hand movement. Together, our findings

indicate a temporal tuning of somatosensory perception during movements to self-touch and highlight the role of the sensorimotor context in predicting and attenuating the received sensory input.

Our results are in strong agreement with the computational framework of internal forward models. Specifically, when reaching with the right hand to touch the left hand, the internal model uses the efference copy of the right hand movement to predict the sensory consequences for the right and left hands, including self-touch and its expected timing [15,17,19]. These predictions are combined with the received sensory input (*e.g.*, visual and proprioceptive) during the movement to re-estimate the state of the right limb *(e.g.*, position, velocity) and update the brain's predictions about the remainder of the movement [62–64]. Because both sensory [18,68,69] and motor signals [70–72] are noisy, the moment of the predicted touch corresponds to a time window reflecting a probability distribution rather than a single point in time and thereby incorporates a certain tolerance for overcoming the intrinsic noise [17]. As the movement of the right hand progresses, new sensory input is received and the predictions of the internal model become increasingly more precise [62–64], narrowing the time window. The received touch is attenuated proportionally to how close they were received with respect to this predicted time window, which continuously narrows during the movement.

Notably, the gradual attenuation of somatosensory perception during movements to self-touch cannot be attributed to task-related differences such as dividing attention between the hands, the mere movement of one hand, a sense of agency, or general cognitive anticipation of tactile stimulation. If this were the case, then we would have observed the same perceptual tuning in the participants of the *No Self-touch* group, in which the same discrimination task and right hand reaching movements were performed. In contrast, we did not observe such a temporal modulation in those participants, and a Bayesian analysis supported the absence of gradual attenuation effects (**Supplementary Text S2**). The direct comparison between the two groups revealed significant differences at almost all probed time points, indicating significantly greater attenuation in the *Self-touch* group than in the *No self-touch* group. This was further supported by linear regression analyses, which showed a significant gradual increase in attenuation during the reaching both in Experiment 1 and in the *Self-touch* group of Experiment 2 but not in the *No Self-touch* group. Finally, further control analyses revealed that these effects cannot be attributed to differences in kinematics or task demands between the two groups.

The results of Experiment 2 demonstrated that the differences in somatosensory perception between the two groups were driven by the sensorimotor context within which participants moved; that is, whether the internal model could predict touch on their left hand based on the motor command of their right hand. Theoretically, the brain should attenuate incoming touch that is likely to be the somatosensory feedback of the movement [17,19,64]. However, the brain needs to first determine the sensorimotor context that describes the relationship between the movement and the predicted sensory input (*i.e.*, internal model) [73,74]. This is critical because the same motor command will be associated with different sensory consequences in different contexts. For example, when manipulating objects, the same movement can produce different predictions about sensory feedback in a context where both hands manipulate the same object compared to a context where each hand manipulates a different object [75]. Similarly, when

moving the right hand towards the left hand, the movement of the right hand may predict touch on the left hand in a self-touch context but not in a context where the right hand deliberately stops before contacting the left hand [27,28,30]. Likewise, pressing the right index finger a few centimetres away from the left index finger should not produce any expectations about receiving touch on the left hand [17,23,39] unless in a context where the right hand holds a tool to directly press on the left one [38]. Consequently, the movements the *No Self-touch* group performed were not associated with any somatosensory consequences on their left hand, which explains why we did not observe an attenuation of the perceived magnitude during their reaching movements. Our results therefore provide evidence that the brain evaluates the sensorimotor context to dynamically shape somatosensory perception in time depending on whether somatosensory input can be predicted based on the efference copy. This conclusion complements changes in movement strategies that humans can make to minimize movement-related suppression effects (*e.g.*, moving with slow velocity [76]) and attenuation of self-generated input (*e.g.*, applying weak forces that are subject to less attenuation [77]) when tactile feedback is beneficial for task completion, such as movement guidance [78,79] or active object exploration [80,81].

Our results cannot be attributed to backward or forward masking of the *test* force by the press of the right index finger. The test force was presented at 192 ms and 174 ms on average for Experiment 1 and Experiment 2 (*Self-touch* group) before the target was reached (*late* trials), and at 300 ms afterwards (*post-reach* trials), while backward masking effects were not detected at 150 ms [50] and forward masking effects last only up to 100-200 ms [82–85]. In addition, our findings do not support the proposal that motor predictions enhance the expected somatosensory signals [86], as we did not observe any enhancement effects at any of the time points probed in either of the two experiments. Instead, we consistently observed robust attenuation effects when subjects expected self-touch based on their movement. In line with the present findings, we recently demonstrated that the previously reported "enhancement" effects these proposals are based on [87] stem from a biased reference condition used rather than action prediction [30]. If a bias-free condition is employed, no enhancement effects are experimentally observed [30]. Moreover, our effects contradict the proposal that attenuation effects are due to non-predictive mechanisms related to double tactile stimulation of both fingers [87] as we observed somatosensory perception at times *before* the two hands made contact (*i.e.*, in *mid* and *late* trials), in line with earlier findings [27] (see also [21]). Critically, our effects were dependent on whether participants expected self-touch (*Self-touch* group) and extended beyond the mere influence of the movement of the right hand, the touch on the left hand, or the discrimination task *per se* (*No self-touch* group). Consistent with the current results, we recently demonstrated that attenuation effects are observed independently of the double tactile stimulation of the two hands: touch is robustly attenuated if participants predict that they will make contact between their two hands but unexpectedly miss the contact [30] (see also [28] for similar findings). Therefore, the present study provides additional evidence that somatosensory attenuation is driven by the prediction of the sensory consequences of voluntary movement rather than somatosensory reafference *per se*.

A methodological strength of the present study is that it assessed how the predictions of the internal model affect the somatosensory perception of a stationary limb. Previous studies have

shown that somatosensory perception is 'gated' on a limb that moves compared to one at rest
[47,50,51,76,88–92] and that this gating can be variably affected by movement (compare [59,93–96]).
However, several studies have consistently demonstrated that these gating effects are also
observed during passive movements (*i.e.*, in the absence of motor signals), indicating that
somatosensory effects on a moving limb are influenced by postdictive mechanisms related to
peripheral reafference [52–55,57,97]. In contrast, somatosensory perception on a limb that remains
stationary is not subject to gating effects [55,92,95,97–99]. Moreover, somatosensory perception on a
passive limb is attenuated only if the received touch results from an active movement, not from
a passive movement [34] or passive stimulation [27] (see also [35]). Thus, our approach isolated the
effects of sensorimotor predictions while avoiding the confounding influence of peripheral
mechanisms.

Our results indicate a scaling of attenuation magnitude as the reaching movement progresses.
In contrast, an earlier study reported no changes in somatosensory perception on the static left
hand during the early or late phases of the reaching movement of the right hand, except for
tactile facilitation—rather than attenuation—around the time the right hand reached its peak
velocity [59]. We did not observe changes in somatosensory perception depending on when the
active hand reached its peak velocity in either of our two experiments, despite having a greater
sample size than this earlier study. A reason for this discrepancy could be that we used
suprathreshold force stimuli while that study used near-threshold vibrotactile stimuli [59]. In the
auditory domain, studies using near-threshold and suprathreshold stimuli have shown different
perceptual effects [100,101]. However, this remains a speculation as there is no evidence for a
similar effect in the tactile modality. Another difference between the two studies is that the
earlier study [59] was conducted in darkness, while our participants could use vision to guide
their movements. The authors of the earlier research suggested that the darkness facilitated the
participants' somatosensory perception to guide their hand movement [48,49]. However, we
believe this interpretation requires further evidence given other findings reporting the opposite
effect [96,102] or null findings [47,103] regarding the interaction between vision and somatosensation
during arm movements. Finally, another possible reason for this discrepancy is that participants
in the study of [59] reached with their right hand to either their left index finger or thumb while
receiving stimulation on their little finger. Consequently, their experimental design introduced
an incongruency between the finger where touch was expected based on the movement and the
finger where touch was received. While the authors argue that tactile perception is uniformly
modulated across the whole target hand [48] other research has shown that such incongruency
can hamper somatosensory attenuation [104]. Here, in all trials of our experiments, the
participants reached with their right hand towards their left index finger while receiving
stimulation on their left index finger. Consequently, this allowed us to investigate how
somatosensory attenuation on the left index finger changes when the participants form
expectations about receiving touch on the same finger.

Understanding how expectations about the somatosensory consequences of the movement (*i.e.*,
the internal model) dynamically shape somatosensory perception in time is clinically
important, given previously observed alterations in psychosis spectrum disorders [105–112],
functional movement disorders [113], Parkinson's disease [36] and ageing [37]. Interestingly, patients

with schizophrenia show deficits in attenuating self-generated somatosensory input by
exhibiting perceptual and neural responses to self-touch that are comparable to those of
externally generated touch [110,111]. Using the same task that was used in the present study, we
previously showed that individuals with high schizotypal personality traits, which are
considered a subclinical analogue of schizophrenia [114,115], also show a reduced ability to
attenuate self-touch compared to externally generated touch [29]. In addition to the
somatosensory domain, patients with schizophrenia also show difficulty in attenuating self-
generated speech [116–118] (see also [119]) and oculomotor responses [120]. Together, these findings
indicated that individuals with schizophrenia form incorrect predictions about the sensory
consequences of their actions, misattributing the received self-generated input to external
sources, producing the hallucinations and delusions commonly associated with schizophrenia
[109,112,121–123]. Interestingly, one study showed that patients with schizophrenia have a reduced
attenuation of neural responses to self-generated auditory stimuli but a-close-to-normal
attenuation for the same stimuli when delivered with a 50 ms delay [124], suggesting that the
prediction of the sensory consequences of the action might be delayed in patients with
schizophrenia [124,125] (see also [126]). Additionally, both patients with schizophrenia [127–129] and
individuals with high schizotypal traits [130–133] tend to exhibit a perturbed temporal profile of
multisensory integration, experiencing stimuli that are further apart in time as cooccurring. We
therefore speculate that patients may show a different temporal tuning of somatosensory
perception during self-touch (*e.g.,* flatter slopes, or shifted temporal window) than that
observed in the present study. This hypothesis warrants further exploration in future
experiments.

Sensory attenuation is not unique to humans but is observed across many species with research
on crickets [134,135], mice [136–138], weakly electric fish [139–141], and primates [142–144] showing selective
attenuation and even cancellation of neural responses to self-generated sensory inputs. Broadly,
this serves to increase the salience and facilitate the interpretation of externally generated
sensations [145–150]. Of particular interest are recent reports of a temporally tuned cancellation of
self-produced sounds in mice. Specifically, mice trained to associate tones with moving a lever
to a specific position, showed attenuated responses in the auditory cortex to these self-
generated sounds when presented at the expected time (*i.e.,* when the lever reached the
expected position) [138]. Importantly, this attenuation was not of fixed magnitude during
movement: auditory responses were decreased for sounds played just before movement, or 80
ms after the predicted time or onset and reached their minimum at the expected time [138]. In
addition, mice exhibit attenuated activity in the primary somatosensory cortex when receiving
the pulsatile stimulus on their whiskers at the predicted time, while less attenuation is observed
if the temporal delays of 50 ms and 100 ms from the predicted time are introduced [151]. The
similarity of these findings with those of the present study suggests the presence of an
evolutionarily conserved mechanism for generating precise temporal predictions to attenuate
expected sensory inputs. We speculate that such a mechanism could be evolutionarily
beneficial by increasing the salience of external stimuli (*e.g.,* enhancing the detection of stimuli
an animal receives unexpectedly during movement), thereby improving the detection of
potential threats and, consequently, promoting survival.

Materials and methods

Participants

Ninety naive participants (47 females and 43 males; 87 right-handed, 2 left-handed, and 1 ambidextrous, aged 18-37 years) were recruited to participate in Experiments 1 and 2. The sample size was set to 30 participants per experiment/group based on our earlier studies investigating somatosensory perception using the same task and equipment [30–35,46]. After data collection, one participant was excluded from Experiment 1 due to missing kinematic data, and four participants were excluded from Experiment 2: one from the *Self-touch* group because the same response was given in 95% of the *target* trials resulting in an unreliable psychometric fit, and three participants from the *No self-touch* group because they performed very fast right hand movements, resulting in too few *early* trials (< 10 trials, arbitrary threshold) (see *Kinematic trials – segmentation*). Therefore, the analysis of Experiment 1 was based on data from 29 adults, and the analysis of Experiment 2 was based on data from 56 adults: 29 in the *Self-touch* group and 27 in the *No self-touch* group.

In both experiments, handedness was assessed using the Edinburgh Handedness Inventory (mean ± SD, 83 ± 25, range: −60 to 100) [152]. All participants provided their informed written consent and reported not having any neurological or psychiatric disorders or taking any psychoactive medication. The experiments were approved by the Ethical Review Authority of Stockholm (registration number: 2021-03790).

Experimental apparatus and procedures in Experiment 1

In both experiments, the participants sat at a desk and rested their left hand palm up, with the left index finger placed inside a support (**Figure 2a**, **Supplementary Figure S1**). A vacuum pillow (AB Germa) was provided to support the participants' left arm and increase their comfort. On top of the pulp of their left index finger laid a cylindrical probe (20 mm diameter) that was attached to a lever controlled by a DC electric motor (Maxon EC Motor EC 90 flat; manufactured in Switzerland) and operated by an Arduino Mega 2560 microcontroller using custom software written in C++ (**Figure 2a,** *lever*). A force sensor (FSG15N1A, Honeywell Inc.; diameter, 5 mm; minimum resolution, 0.01 N; response time, 1 ms; measurement range, 0–15 N) was placed within the probe to record the forces exerted on the left index finger by the motor (**Figure 2a,** *force sensor* depicted in red). A force of 0.1 N was constantly applied to the left index finger to maintain contact between the probe and the finger. A second identical force sensor was placed inside an aluminium capsule (20 mm diameter) and positioned on top of, but not in contact with, the cylindrical probe (**Figure 2a,** *force sensor* depicted in cyan). The right forearm was comfortably placed on top of sponge padding, and the right hand rested above a wooden table (70 cm × 25 cm × 12.5 cm). The wooden table was placed on the desk in front of the participants covering their left hand, and thus preventing direct visual feedback from the left hand and fingers. The right index finger was placed at a position marked on top of the wooden table (**Figure 2a**, *starting position*), corresponding to 25 cm to the right of the left index finger.

Upon an auditory GO-cue generated by the Arduino, participants were instructed to perform a
reaching movement with their right hand from the starting position towards the force sensor
located above their left hand. They were instructed to perform a fast but natural movement that
would be concluded once they tapped the force sensor with their right index finger (*i.e.*,
touching the force sensor with the right index finger and immediately lifting the finger).
Participants were allowed to freely look at their right hand and the force sensor as needed to
guide their movement. A motion tracking sensor (Polhemus Liberty, Micro Sensor 1.8™) was
attached to the right index finger with medical tape to continuously record the right index finger
position (**Figure 2a,** *motion sensor*). White noise was provided by a laptop through a pair of
headphones to mask any sounds made by the motor.

Force trials

In each experiment, there were 560 reaching trials and 56 *baseline* trials, resulting in a total of
616 trials per participant grouped in two experimental blocks. The seven possible intensities of
the *comparison* force (1, 1.5, 1.75, 2, 2.25, 2.5, 3 N) were tested 16 times for every type of
reaching trial and 8 times in the *baseline*. The intensity of the *test* force was always fixed at 2
N and its duration was set to 100 ms.

During the reaching trials, we experimentally manipulated the times at which the *test* force was
applied on the left index finger. We applied the *test* force at five different times, based on pilot
experiments, to correspond to different phases of a natural right hand movement reaching
toward the left hand. The *test* force could be applied at 250 ms, 370 ms, and 550 ms after the
go-cue, but the different trials were binned into time bins depending on how the participants
moved on every trial (*i.e.*, times of movement onset, peak velocity, movement offset) to account
for within- and between- subjects differences in movements. Specifically, based on the time at
which the *test* force occurred with respect to the participants' movement (see *Kinematic trials
– segmentation*), the trials were binned to correspond to the *early*, *mid,* and *late* phases of the
right hand movement, at the time when the participants tapped the sensor (*target*), which
simulated self-touch between the hands (system delay ~35 ms), and at a fixed time of 300 ms
after tapping (*post-reach*). All 5 types of reaching trials were randomly interleaved and had an
equal probability of occurrence (20%). During the *baseline* trials, participants remained
motionless, and the *test* force was applied 100 ms after the onset of the trial, followed by the
comparison force 1200 ms later. In every trial, after receiving both forces, participants
responded verbally about which of the two forces (the first force - *test* or the second force -
comparison) felt stronger. Participants were instructed not to balance their responses (*i.e.*,
saying 50% of the time that the first or the second force is stronger), and were further instructed
to make their best guess if the intensity of the two forces felt similar. No feedback was ever
provided to the participants regarding their responses.

Experimental apparatus and procedures in Experiment 2

Experiment 2 was a between-group experiment. The *Self-touch* group (**Figure 3b**) performed
the same task as the participants in Experiment 1. For the *No self-touch* group, the force sensor
that participants approached with their right index finger (**Figure 3b**, *force sensor* depicted in
cyan) was replaced by a distance sensor (Ultrasonic Distance Sensor HC-SR04 5 V Version)

(**Figure 3b**, *distance sensor* depicted in yellow). The distance sensor was placed on top of, but
not in contact with, the probe and was connected to an Arduino microcontroller that controlled
a servo motor (Grove-Servo). Once the distance sensor detected that the position of the right
index finger became lower than a pre-set position threshold, it triggered a servo motor that hit
the force sensor that triggered the *test* force. The servo motor and the force sensor were hidden
from the participants' view. The position threshold was set to approximately match the position
participants had when pressing the force sensor with their right index finger in Experiment 1
and in the *Self-touch* group. A visual marker was placed 2 cm above the distance sensor to
indicate the approximate vertical position where participants should perform the virtual tap to
avoid physical contact with the sensor, and the position of the pre-set threshold was adjusted
to this distance. Participants were instructed to perform a fast but natural movement with their
right hand toward the distance sensor and to conclude the movement by tapping the right index
finger above the sensor without touching it. That is, the participants were instructed to perform
movements like those of Experiment 1, with the difference that they never experienced contact
between their two fingers and thus never expected to make such contact. Similar to our previous
study [30], we demonstrated the movement to the participants prior to the experiment, and there
were practice trials to ensure that they performed the movement as requested before starting
the experiment. The end of the reaching was defined as the time point at which the distance
sensor was triggered by the downward movement of the right index finger.

All probed times in Experiment 2 were identical for the *Self-touch* and *No self-touch* groups.
However, forces were delivered 100 ms later in Experiment 2 than in Experiment 1. All the
other details, including the analysis, remained identical to those of Experiment 1.

Kinematic trials – preprocessing

The position of the right index finger was recorded at a sampling rate of 240 Hz. TTL signals
were used to indicate the onset of the trial, as well as the *test* force and the *comparison* force,
thus co-registering the force trials with the kinematic trials. We smoothed the position data
using a moving average filter of 5 data points span ($\approx$ 21 ms).

Kinematic trials – segmentation

For each reaching trial, we calculated the movement onset as the first time point during the trial
when the velocity exceeded 5 cm/s for 100 ms sequentially [93,95,96]. The end of the reaching (*i.e.*,
offset) was calculated as the first time point when the force exerted on the force sensor
exceeded a pre-set threshold (0.3 N) indicating that participants tapped the sensor with their
right index finger and thus concluded the movement. The duration of the movement was
calculated as the time difference between the movement offset and onset.

Next, we calculated the time when the 3D velocity of the right hand reached its peak value. To
capture the peak velocity of the reaching movement, which typically occurs during the first
half of the movement [65], and not the peak velocity of the index finger when tapping the sensor
(occurring at the end of the movement), we restricted the time window of the peak velocity to
within 67% of the movement duration.

Based on the calculated kinematic parameters of every reaching movement, we binned the
reaching trials into three distinct categories depending on when the *test* force occurred within
a specific trial: *early* trials (*test* force occurring between the movement onset and the time the
right hand velocity reached 85% of its peak value), *mid* trials (*test* force occurring after the
right hand velocity reached 85% of its peak value up until it dropped below 85% of its peak
value), and *late* trials (*test* force occurring after the right hand velocity dropped below 85% of
its peak value and before the right index finger made contact with the force sensor). Therefore,
this binning approach allowed us to tailor the probed times to the different phases of the
reaching movement, by taking into consideration potential changes in reaction and reaching
times from trial to trial. In addition, the trials when the *test* force occurred simultaneously with
the participant tapping the force sensor were classified as *target* trials, and the trials when the
test force occurred 300 ms after the participants' tap on the force sensor were classified as *post-*
reach trials (**Figure 2c**). Therefore, the final five groups of reaching trials entering the statistical
analysis were *early*, *mid*, *late*, *target*, and *post-reach* trials.

Exclusion of trials in Experiment 1

As mentioned earlier, each participant was administered 616 trials resulting in 17864 trials in
total (29 × 616). We excluded (a) any *baseline* or reaching trials in which the intensity of the
test force (2 N) was not properly applied (*test* < 1.85 N or *test* > 2.15 N), or the participants'
force responses or kinematic data were not properly registered, (b) any reaching trials in which
the participants failed to tap the sensor or did not press/release the sensor properly, (c) any
reaching trials in which the participants did not move as instructed (*e.g.*, they moved before the
auditory GO-cue, they started their movement from a position closer to the force sensor than
the starting position (< 5 cm away from the starting position on the x-axis)), (d) any reaching
trials in which the participants received the *comparison* force while moving, and (e) any *early*,
mid, or *late* trials in which the *test* force was applied either before the participants started
moving (*i.e.*, participants were too slow in starting their movement) or after they tapped the
sensor (*i.e.*, participants were too fast in finishing their movement). Finally, since our setup
contained metallic surfaces (*e.g.*, the aluminium capsule) that can distort the electromagnetic
recordings of the motion tracking sensor when in close proximity, we rejected (f) any reaching
trials in which the sensor was inside the distortion range at times used for trial segmentation
(see below). In total, we rejected 1590 out of 17864 trials (8.9%). This resulted in an average
of 110 *early*, 84 *mid*, 92 *late*, 109 *target*, 110 *post-reach*, and 56 *baseline* trials.

Exclusion of trials in Experiment 2

We used identical criteria for excluding trials as in Experiment 1 with the exception that in the
No self-touch group we also rejected any *target* and *post-reach* trials in which the participants
did not trigger the distance sensor when moving their finger downwards. In addition, for this
group we also modified our exclusion criteria under b) (see above) to exclude any trials in
which the distance sensor was not properly triggered, instead of participants failing to press the
sensor. In total, we excluded 1411 out of 17864 trials (7.9%) in the *Self-touch* group and 2051
out of 16632 trials (12.3%) in the *No self-touch* group. This resulted in an average of 87 *early*,
101 *mid*, 108 *late*, 107 *target*, 108 *post-reach* trials, and 56 *baseline* trials for the *Self-touch*

group and an average of 64 *early*, 114 *mid*, 136 *late*, 85 *target*, 85 *post-reach,* and 56 *baseline*
trials for the *No self-touch* group.

Psychophysical fits

The participants' responses were fitted with psychometric curves (**Figure 2g**), using the *glm*
function of the R *stats* package, with a generalized linear model (Equation 1):

$$p = \frac{e^{\beta 0 + \beta 1 x}}{1 + e^{\beta 0 + \beta 1 x}} \text{ (Equation 1)}$$

We extracted the point of subjective equality ($PSE = -\frac{\beta 0}{\beta 1}$) that corresponds to the intensity
at which the *test* force felt equally strong to the *comparison* force ($p = 0.5$). This was our
variable of interest since it represents the perceived intensity of the *test* force. For control
analysis, we also extracted the just noticeable difference in Experiment 2 that reflects the
participants' discrimination capacity in the psychophysics task ($JND = \frac{\log (3)}{\beta 1}$) and
corresponds to the difference between the thresholds at $p = 0.5$ and $p = 0.75$. A higher JND
indicates poorer discrimination ability for the applied forces, suggesting greater difficulty of
the task. Before fitting the responses, the *comparison* forces were rebinned to the nearest of
seven possible force intensities (1, 1.5, 1.75, 2, 2.25, 2.5, or 3 N). The goodness of fit was
assessed using McFadden's pseudo R^2.

Statistical analyses

The data were analysed in R (version 4.2.2) [153] and JASP (version 0.17) [154]. We performed
repeated measures and mixed ANOVAs to analyse the data. In the case of sphericity violations,
Greenhouse-Geisser corrections were applied. Effect sizes for the ANOVA were given by
partial eta-squared (η_p^2). Planned pairwise comparisons between trial types were performed
using parametric (paired or independent-sample *t*-tests) or nonparametric (Wilcoxon signed-
rank and Wilcoxon rank sum) tests depending on the normality of the variable distributions.
We limited the pairwise comparisons to those central to our hypothesis, which focused on
indicating a serial time dependence during reaching (*early* vs. *mid*, *early* vs. *late*, *early* vs.
target, *mid* vs. *late*, *mid* vs. *target*, *late* vs. *target*) and recovery after reaching (*target* vs. *post-
reach*). The normality of the data was assessed with the Shapiro–Wilk test. Corrections for
multiple comparisons were performed using the false discovery rate (*FDR*) and are denoted as
"*FDR-corrected*". For every statistical comparison, we report the corresponding statistic, the
95% confidence intervals (CI^{95}), and the effect size (Cohen's *d* or the matched rank-biserial
correlation (*rrb*)). Linear regression and multiple linear regression were performed using the
lm function of the R *stats* package. A Bayesian control analysis was performed to provide
evidence in favour of the null hypothesis for comparisons of interest. The analysis was done in
JASP using default Cauchy priors of 0.707, where the Bayes factor BF_{0+} represents the
likelihood of the null hypothesis compared to the alternative hypothesis for the one-tailed tests.
We interpret a Bayes factor from 1 to 3 as providing 'anecdotal' support for the null hypothesis,
a Bayes factor from 3 to 10 as providing 'moderate' support for the null hypothesis, and a Bayes
factor above 10 as providing 'strong' support for the null hypothesis [155–157]. General additive

model regression lines were generated using the *geom_smooth* function of the R *ggplot2* package.

Acknowledgments

The authors would like to thank Dr. Pawel Tacikowski, Dr. Antonella Maselli, Dr. Luc Selen, and Dr. Maximilian Hauser for fruitful discussions. Experiment costs were covered by the Swedish Research Council (VR Starting Grant 2019-01909) and the Åke Wibergs Foundation (M20-0038). N.C. was supported by the Swedish Research Council (VR Starting Grant 2019-01909). X.J. received funding from the European Union's Horizon Europe research and innovation program under Marie Sklodowska-Curie (101059348). K.K. was supported by the European Research Council (ERC Starting Grant 101039152).

Author Contributions

Noa Cemeljic: Conceptualization, Data curation, Formal Analysis, Investigation, Software, Visualization, Writing – original draft, Writing – review & editing. **Xavier Job**: Conceptualization, Writing – review & editing. **Konstantina Kilteni:** Conceptualization, Data curation, Formal Analysis, Funding acquisition, Methodology, Software, Visualization, Supervision, Writing – original draft, Writing – review & editing.

Declarations of interests

The authors declare no competing interests.

References

1. Myowa-Yamakoshi, M., and Takeshita, H. (2006). Do human fetuses anticipate self-oriented actions? A study by four-dimensional (4D) ultrasonography. Infancy *10*, 289–301. https://doi.org/10.1207/s15327078in1003_5.

2. Kurjak, A., Azumendi, G., Veček, N., Kupešic, S., Solak, M., Varga, D., and Chervenak, F. (2003). Fetal hand movements and facial expression in normal pregnancy studied by four-dimensional sonography. J Perinat Med *31*. https://doi.org/10.1515/JPM.2003.076.

3. Reese, A. (2023). Patterns of Self-Directed Touch over the First 18 Months of Life - ProQuest.

4. Thomas, B.L., Karl, J.M., and Whishaw, I.Q. (2014). Independent development of the Reach and the Grasp in spontaneous self-touching by human infants in the first 6 months. Front Psychol *5*, 1526. https://doi.org/10.3389/fpsyg.2014.01526.

5. DiMercurio, A., Connell, J.P., Clark, M., and Corbetta, D. (2018). A naturalistic observation of spontaneous touches to the body and environment in the first 2 months of life. Front Psychol *9*, 2613. https://doi.org/10.3389/fpsyg.2018.02613.

793 6. Spille, J.L., Grunwald, M., Martin, S., and Mueller, S.M. (2021). Stop touching your
794   face! A systematic review of triggers, characteristics, regulatory functions and neuro-
795   physiology of facial self touch. Preprint at Elsevier Ltd,
796   https://doi.org/10.1016/j.neubiorev.2021.05.030.
797

798 7. Bolanowski, S.J., Verrillo, R.T., and McGlone Francis (1999). Passive, active and
799   intra-active (self) touch. Somatosens Mot Res *16*, 304–311.
800

801 8. Schütz-Bosbach, S., Musil, J.J., and Haggard, P. (2009). Touchant-touché: The role of
802   self-touch in the representation of body structure. Conscious Cogn *18*, 2–11.
803   https://doi.org/10.1016/j.concog.2008.08.003.
804

805 9. Ehrsson, H.H., Holmes, N.P., and Passingham, R.E. (2005). Touching a rubber hand:
806   Feeling of body ownership is associated with activity in multisensory brain areas.
807   Journal of Neuroscience *25*, 10564–10573. https://doi.org/10.1523/JNEUROSCI.0800-
808   05.2005.
809

810 10. Rochat, P., and Hespos, S.J. (1997). Differential Rooting Response by Neonates:
811   Evidence for an Early Sense of Self. Infant Child Dev *6*, 105–112.
812

813 11. Kawato, M., Furukawa, K., and Suzuki, R. (1987). A hierarchical neural-network
814   model for control and learning of voluntary movement. Biol Cybern *57*, 169–185.
815   https://doi.org/10.1007/BF00364149.
816

817 12. Miall, R.C., and Wolpert, D.M. (1996). Forward Models for Physiological Motor
818   Control. Neural Networks *9*, 1265–1279. https://doi.org/10.1016/S0893-
819   6080(96)00035-4.
820

821 13. Franklin, D.W., and Wolpert, D.M. (2011). Computational Mechanisms of
822   Sensorimotor Control. Neuron *72*, 425–442.
823   https://doi.org/10.1016/j.neuron.2011.10.006.
824

825 14. Jordan, M. (1992). Forward models: Supervised learning with a distal teacher. Cogn
826   Sci *16*, 307–354. https://doi.org/10.1016/0364-0213(92)90036-T.
827

828 15. McNamee, D., and Wolpert, D.M. (2019). Internal Models in Biological Control. Annu
829   Rev Control Robot Auton Syst *2*, 339–364. https://doi.org/10.1146/annurev-control-
830   060117-105206.
831

832 16. Kilteni, K. (2023). Methods of Somatosensory Attenuation. In Somatosensory
833   Research Methods, N. P. Holmes, ed. (Springer US), pp. 35–53.
834   https://doi.org/10.1007/978-1-0716-3068-6_2.
835

17. Bays, P.M., and Wolpert, D.M. (2008). Predictive attenuation in the perception of
touch. In Sensorimotor Foundations of Higher Cognition, E. P. Haggard, Y. Rosetti,
and M. Kawato, eds. (Oxford University Press), pp. 339–358.

18.   Bays, P.M., and Wolpert, D.M. (2007). Computational principles of sensorimotor
control that minimize uncertainty and variability. J Physiol *578*, 387–396.
https://doi.org/10.1113/jphysiol.2006.120121.

19.   Wolpert, D.M., and Flanagan, J.R. (2001). Motor prediction. Current Biology *11*,
R729–R732. https://doi.org/10.1016/S0960-9822(01)00432-8.

20.   Blakemore, S.-J., Wolpert, D., and Frith, C. (2000). Why can't you tickle yourself?
Neuroreport *11*, R11–R16. https://doi.org/10.1097/00001756-200008030-00002.

21.   Vale, N., Tomic, I., Girones, Z., Wolpert, D.M., Kilteni, K., and Bays, P. (2024).
Divisive attenuation based on noisy sensorimotor predictions accounts for excess
variability in self-touch. bioRxiv, 2024.06.20.599826.
https://doi.org/10.1101/2024.06.20.599826.

22.   Shergill, S.S., Bays, P.M., Frith, C.D., and Wolpert, D.M. (2003). Two Eyes for an
Eye: The Neuroscience of Force Escalation. Science (1979) *301*, 187–187.
https://doi.org/10.1126/science.1085327.

23.   Kilteni, K., Andersson, B.J., Houborg, C., and Ehrsson, H.H. (2018). Motor imagery
involves predicting the sensory consequences of the imagined movement. Nat
Commun *9*, 1617. https://doi.org/10.1038/s41467-018-03989-0.

24.   Lalouni, M., Fust, J., Vadenmark-Lundqvist, V., Ehrsson, H.H., Kilteni, K., and
Birgitta Jensen, K. (2021). Predicting pain: differential pain thresholds during self-
induced, externally induced, and imagined self-induced pressure pain. Pain *162*, 1539–
1544. https://doi.org/10.1097/j.pain.0000000000002151.

25.   Weiskrantz, L., Elliott, J., and Darlington, C. (1971). Preliminary Observations on
Tickling Oneself. Nature *230*, 598–599. https://doi.org/10.1038/230598a0.

26.   Blakemore, S.J., Frith, C.D., and Wolpert, D.M. (1999). Spatio-temporal prediction
modulates the perception of self-produced stimuli. J Cogn Neurosci *11*, 551–559.
https://doi.org/10.1162/089892999563607.

27.   Bays, P.M., Wolpert, D.M., and Flanagan, J.R. (2005). Perception of the consequences
of self-action is temporally tuned and event driven. Current Biology *15*, 1125–1128.
https://doi.org/10.1016/j.cub.2005.05.023.

28.   Bays, P.M., Flanagan, J.R., and Wolpert, D.M. (2006). Attenuation of self-generated
tactile sensations is predictive, not postdictive. PLoS Biol *4*, 281–284.
https://doi.org/10.1371/journal.pbio.0040028.

29. Asimakidou, E., Job, X., and Kilteni, K. (2022). The positive dimension of schizotypy
is associated with a reduced attenuation and precision of self-generated touch.
Schizophrenia *8*, 57. https://doi.org/10.1038/s41537-022-00264-6.

30. Job, X., and Kilteni, K. (2023). Action does not enhance but attenuates predicted
touch. Elife *12*. https://doi.org/10.7554/eLife.90912.

31. Timar, L., Job, X., Orban de Xivry, J.-J., and Kilteni, K. (2023). Aging exerts a limited
influence on the perception of self-generated and externally generated touch. J
Neurophysiol *130*, 871–882. https://doi.org/10.1152/jn.00145.2023.

32. Kilteni, K., Houborg, C., and Ehrsson, H.H. (2019). Rapid learning and unlearning of
predicted sensory delays in self-generated touch. Elife *8*, e42888.
https://doi.org/10.7554/eLife.42888.

33. Kilteni, K., Engeler, P., Boberg, I., Maurex, L., and Ehrsson, H.H. (2021). No evidence
for somatosensory attenuation during action observation of self-touch. European
Journal of Neuroscience *54*, 6422–6444. https://doi.org/10.1111/ejn.15436.

34. Kilteni, K., Engeler, P., and Ehrsson, H.H. (2020). Efference Copy Is Necessary for the
Attenuation of Self-Generated Touch. iScience *23*, 100843.
https://doi.org/10.1016/j.isci.2020.100843.

35. Kilteni, K., and Ehrsson, H.H. (2022). Predictive attenuation of touch and tactile
gating are distinct perceptual phenomena. iScience *25*, 104077.
https://doi.org/10.1016/J.ISCI.2022.104077.

36. Wolpe, N., Zhang, J., Nombela, C., Ingram, J.N., Wolpert, D.M., and Rowe, J.B.
(2018). Sensory attenuation in Parkinson's disease is related to disease severity and
dopamine dose. Sci Rep *8*, 15643. https://doi.org/10.1038/s41598-018-33678-3.

37. Wolpe, N., Ingram, J.N., Tsvetanov, K.A., Geerligs, L., Kievit, R.A., Henson, R.N.,
Wolpert, D.M., and Rowe, J.B. (2016). Ageing increases reliance on sensorimotor
prediction through structural and functional differences in frontostriatal circuits. Nat
Commun *7*, 13034. https://doi.org/10.1038/ncomms13034.

38. Kilteni, K., and Ehrsson, H.H. (2017). Sensorimotor predictions and tool use: Hand-
held tools attenuate self-touch. Cognition *165*, 1–9.
https://doi.org/10.1016/j.cognition.2017.04.005.

39. Kilteni, K., and Ehrsson, H.H. (2017). Body ownership determines the attenuation of
self-generated tactile sensations. Proceedings of the National Academy of Sciences
114, 8426–8431. https://doi.org/10.1073/pnas.1703347114.

40. Blakemore, S.-J., Wolpert, D.M., and Frith, C.D. (1998). Central cancellation of self-
produced tickle sensation. Nat Neurosci *1*, 635–640. https://doi.org/10.1038/2870.

41. Blakemore, S.-J., Wolpert, D.M., and Frith, C.D. (1999). The Cerebellum Contributes
to Somatosensory Cortical Activity during Self-Produced Tactile Stimulation.
Neuroimage *10*, 448–459. https://doi.org/10.1006/nimg.1999.0478.

42. Blakemore, S.-J., Frith, C.D., and Wolpert, D.M. (2001). The cerebellum is involved in
predicting the sensory consequences of action. Neuroreport *12*, 1879–1884.
https://doi.org/10.1097/00001756-200107030-00023.

43. Hesse, M.D., Nishitani, N., Fink, G.R., Jousmaki, V., and Hari, R. (2010). Attenuation
of Somatosensory Responses to Self-Produced Tactile Stimulation. Cerebral Cortex *20*,
425–432. https://doi.org/10.1093/cercor/bhp110.

44. Shergill, S.S., White, T.P., Joyce, D.W., Bays, P.M., Wolpert, D.M., and Frith, C.D.
(2013). Modulation of somatosensory processing by action. Neuroimage *70*, 356–362.
https://doi.org/10.1016/j.neuroimage.2012.12.043.

45. Kilteni, K., and Ehrsson, H.H. (2020). Functional Connectivity between the
Cerebellum and Somatosensory Areas Implements the Attenuation of Self-Generated
Touch. The Journal of Neuroscience *40*, 894–906.
https://doi.org/10.1523/JNEUROSCI.1732-19.2019.

46. Kilteni, K., Houborg, C., and Ehrsson, H.H. (2023). Brief temporal perturbations in
somatosensory reafference disrupt perceptual and neural attenuation and increase
supplementary motor area–cerebellar connectivity. The Journal of Neuroscience, JN-
RM-1743-22. https://doi.org/10.1523/JNEUROSCI.1743-22.2023.

47. Gertz, H., Voudouris, D., and Fiehler, K. (2017). Reach-relevant somatosensory signals
modulate tactile suppression. J Neurophysiol *117*, 2262–2268.
https://doi.org/10.1152/jn.00052.2017.

48. Voudouris, D., and Fiehler, K. (2017). Spatial specificity of tactile enhancement during
reaching. Atten Percept Psychophys *79*, 2424–2434. https://doi.org/10.3758/s13414-
017-1385-7.

49. Voudouris, D., and Fiehler, K. (2017). Enhancement and suppression of tactile signals
during reaching. J Exp Psychol Hum Percept Perform *43*, 1238.
https://doi.org/10.1037/xhp0000373.

50. Fraser, L.E., and Fiehler, K. (2018). Predicted reach consequences drive time course of
tactile suppression. Behavioural Brain Research *350*, 54–64.
https://doi.org/10.1016/j.bbr.2018.05.010.

51. Williams, S.R., Shenasa, J., and Chapman, C.E. (1998). Time course and magnitude of
movement-related gating of tactile detection in humans. I. Importance of stimulus
location. J Neurophysiol *79*, 947–963. https://doi.org/10.1152/jn.00527.2001.

52.   Williams, S.R., and Chapman, C.E. (2002). Time course and magnitude of movement-
977   related gating of tactile detection in humans. III. Effect of motor tasks. J Neurophysiol
978   *88*, 1968–1979. https://doi.org/10.1152/jn.2002.88.4.1968.

980   53.   Chapman, C.E. (1994). Active versus passive touch: Factors influencing the
981   transmission of somatosensory signals to primary sornatosensory cortex. Can J Physiol
982   Pharmacol *72*, 558–570. https://doi.org/10.1139/y94-080.

984   54.   Chapman, C.E., and Beauchamp, E. (2006). Differential controls over tactile detection
985   in humans by motor commands and peripheral reafference. J Neurophysiol *96*, 1664–
986   1675. https://doi.org/10.1152/jn.00214.2006.

988   55.   Chapman, C.E., Bushnell, M.C., Miron, D., Duncan, G.H., and Lund, J.P. (1987).
989   Sensory perception during movement in man. Exp Brain Res *68*, 516–524.
990   https://doi.org/10.1007/BF00249795.

992   56.   Voss, M., Ingram, J.N., Wolpert, D.M., and Haggard, P. (2008). Mere expectation to
993   move causes attenuation of sensory signals. PLoS One *3*, 2–6.
994   https://doi.org/10.1371/journal.pone.0002866.

996   57.   Arikan, B.E., Voudouris, D., Straube, B., and Fiehler, K. (2024). Distinct role of
997   central predictive mechanisms in tactile suppression. iScience *27*, 110582.
998   https://doi.org/10.1016/j.isci.2024.110582.

1000   58.   Walsh, L.D., Taylor, J.L., and Gandevia, S.C. (2011). Overestimation of force during
1001   matching of externally generated forces. J Physiol *589*, 547–557.
1002   https://doi.org/10.1113/jphysiol.2010.198689.

1003

1004   59.   Voudouris, D., and Fiehler, K. (2021). Dynamic temporal modulation of
1005   somatosensory processing during reaching. Sci Rep *11*, 1–12.
1006   https://doi.org/10.1038/s41598-021-81156-0.

1007

1008   60.   Kilteni, K., and Ehrsson, H.H. (2024). Dynamic changes in somatosensory and
1009   cerebellar activity mediate temporal recalibration of self-touch. Commun Biol *7*, 522.
1010   https://doi.org/10.1038/s42003-024-06188-4.

1011

1012   61.   Desmurget, M., and Grafton, S. (2000). Forward modeling allows feedback control for
1013   fast reaching movements. Trends Cogn Sci *4*, 423–431.

1014

1015   62.   Brenner, E., and Smeets, J.B.J. (2018). Continuously updating one's predictions
1016   underlies successful interception. REVIEW Control of Movement J Neurophysiol *120*,
1017   3257–3274. https://doi.org/10.1152/jn.00517.2018.-This.

1018

1019 63.   Shadmehr, R., Krakauer, J.W., Neuroanatomy, A.C., and Motor, F.O.R. (2008). A
1020   computational neuroanatomy for motor control. Exp Brain Res *185*, 359–381.
1021   https://doi.org/10.1007/s00221-008-1280-5.
1022

1023 64.   Shadmehr, R., Smith, M. a, and Krakauer, J.W. (2010). Error correction, sensory
1024   prediction, and adaptation in motor control. Annu Rev Neurosci *33*, 89–108.
1025   https://doi.org/10.1146/annurev-neuro-060909-153135.
1026

1027 65.   McCrea, P.H., Eng, J.J., and Hodgson, A.J. (2002). Biomechanics of reaching: clinical
1028   implications for individuals with acquired brain injury. Disabil Rehabil *24*, 534–541.
1029   https://doi.org/10.1080/09638280110115393.
1030

1031 66.   Deng, C.-L., Geng, P., Hu, Y.-F., and Kuai, S.-G. (2019). Beyond Fitts's Law: A Three-
1032   Phase Model Predicts Movement Time to Position an Object in an Immersive 3D
1033   Virtual Environment. Human Factors: The Journal of the Human Factors and
1034   Ergonomics Society *61*, 879–894. https://doi.org/10.1177/0018720819831517.
1035

1036 67.   Kilteni, K., and Ehrsson, H.H. (2023). Dynamic changes in somatosensory and
1037   cerebellar activity mediate temporal recalibration of self-touch. bioRxiv,
1038   2023.12.01.569520. https://doi.org/10.1101/2023.12.01.569520.
1039

1040 68.   Faisal, A.A., Selen, L.P.J., and Wolpert, D.M. (2008). Noise in the nervous system. Nat
1041   Rev Neurosci *9*, 292–303. https://doi.org/10.1038/nrn2258.
1042

1043 69.   Jones, P.W., and Gabbiani, F. (2012). Impact of neural noise on a sensory-motor
1044   pathway signaling impending collision. J Neurophysiol *107*, 1067–1079.
1045   https://doi.org/10.1152/jn.00607.2011.
1046

1047 70.   Harris, C.M., and Wolpert, D.M. (1998). Signal-dependent noise determines motor
1048   planning. Nature *394*, 780–784. https://doi.org/10.1038/29528.
1049

1050 71.   Todorov, E., and Jordan, M.I. (2002). Optimal feedback control as a theory of motor
1051   coordination. Nat Neurosci *5*, 1226–1235. https://doi.org/10.1038/nn963.
1052

1053 72.   Thorp, E.B., Kording, K.P., and Mussa-Ivaldi, F.A. (2017). Using noise to shape motor
1054   learning. J Neurophysiol *117*, 728–737. https://doi.org/10.1152/jn.00493.2016.
1055

1056 73.   Imamizu, H., Sugimoto, N., Osu, R., Tsutsui, K., Sugiyama, K., Wada, Y., and Kawato,
1057   M. (2007). Explicit contextual information selectively contributes to predictive
1058   switching of internal models. Exp Brain Res *181*, 395–408.
1059   https://doi.org/10.1007/s00221-007-0940-1.
1060

1061 74.   Haruno, M., Wolpert, D.M., and Kawato, M. (2001). MOSAIC Model for
1062   Sensorimotor Learning and Control. Neural Comput *13*, 2201–2220.
1063   https://doi.org/10.1162/089976601750541778.
1064

1065   75.   Blakemore, S.J., Goodbody, S.J., and Wolpert, D.M. (1998). Predicting the
1066   consequences of our own actions: the role of sensorimotor context estimation. J
1067   Neurosci *18*, 7511–7518. https://doi.org/Not available.
1068

1069   76.   Cybulska-Klosowicz, A., Meftah, E.M., Raby, M., Lemieux, M.L., and Chapman, C.E.
1070   (2011). A critical speed for gating of tactile detection during voluntary movement. Exp
1071   Brain Res *210*, 291–301. https://doi.org/10.1007/s00221-011-2632-0.
1072

1073   77.   Shergill, S.S., Bays, P.M., Frith, C.D., and Wolpert, D.M. (2003). Two eyes for an eye:
1074   the neuroscience of force escalation. Science *301*, 187.
1075   https://doi.org/10.1126/science.1085327.
1076

1077   78.   Ryan, C.P., Ciotti, S., Balestrucci, P., Bicchi, A., Lacquaniti, F., Bianchi, M., and
1078   Moscatelli, A. (2024). The relativity of reaching: Motion of the touched surface alters
1079   the trajectory of hand movements. iScience *27*, 109871.
1080   https://doi.org/10.1016/j.isci.2024.109871.
1081

1082   79.   Moscatelli, A., Bianchi, M., Ciotti, S., Bettelani, G.C., Parise, C. V., Lacquaniti, F., and
1083   Bicchi, A. (2019). Touch as an auxiliary proprioceptive cue for movement control. Sci
1084   Adv *5*. https://doi.org/10.1126/sciadv.aaw3121.
1085

1086   80.   Pruszynski, J.A., Flanagan, J.R., and Johansson, R.S. (2018). Fast and accurate edge
1087   orientation processing during object manipulation. Elife *7*.
1088   https://doi.org/10.7554/eLife.31200.
1089

1090   81.   Morley, J.W., Goodwin, A.W., and Darian-Smith, I. (1983). Tactile discrimination of
1091   gratings. Exp Brain Res *49*. https://doi.org/10.1007/BF00238588.
1092

1093   82.   Bhattacharjee, A., Ye, A.J., Lisak, J.A., Vargas, M.G., and Goldreich, D. (2010).
1094   Vibrotactile masking experiments reveal accelerated somatosensory processing in
1095   congenitally blind braille readers. Journal of Neuroscience *30*, 14288–14298.
1096   https://doi.org/10.1523/JNEUROSCI.1447-10.2010.
1097

1098   83.   Craig, J.C., and Evans, P.M. (1987). Vibrotactile masking and the persistence of tactual
1099   features.
1100

1101   84.   Coquery, J.-M., and Amblard, B. (1973). Backward and forward masking in the
1102   perception of cutaneous stimuli.
1103

1104   85.   Makous, J.C., Gescheider, G.A., and Bolanowski, S.J. (1996). Decay in the effect of
1105   vibrotactile masking. J Acoust Soc Am *99*, 1124–1129.
1106

1107   86.   Press, C., Kok, P., and Yon, D. (2020). The Perceptual Prediction Paradox. Trends
1108   Cogn Sci *24*, 13–24. https://doi.org/10.1016/j.tics.2019.11.003.
1109

1110 87. Thomas, E.R., Yon, D., de Lange, F.P., and Press, C. (2021). Action Enhances
1111 Predicted Touch. Psychol Sci. https://doi.org/10.1177/09567976211017505.
1112

1113 88. Angel, R.W., and Malenka, R.C. (1982). Velocity-dependent suppression of cutaneous
1114 sensitivity during movement. Exp Neurol 77, 266–274. https://doi.org/10.1016/0014-
1115 4886(82)90244-8.
1116

1117 89. Post, L.J., Zompa, I.C., and Chapman, C.E. (1994). Perception of vibrotactile stimuli
1118 during motor activity in human subjects. Exp Brain Res 100, 107–120.
1119 https://doi.org/10.1007/BF00227283.
1120

1121 90. Voudouris, D., Broda, M.D., and Fiehler, K. (2019). Anticipatory grasping control
1122 modulates somatosensory perception. J Vis 19, 1–10. https://doi.org/10.1167/19.5.4.
1123

1124 91. Williams, S.R., and Chapman, C.E. (2000). Time course and magnitude of movement-
1125 related gating of tactile detection in humans. II. Effects of stimulus intensity. J
1126 Neurophysiol 84, 863–875. https://doi.org/10.1152/jn.2000.84.2.863.
1127

1128 92. Papakostopoulos, D., Cooper, R., and Crow, H.J. (1975). Inhibition of cortical evoked
1129 potentials and sensation by self-initiated movement in man. Nature 258, 321–324.
1130 https://doi.org/10.1038/258321a0.
1131

1132 93. Colino, F.L., and Binsted, G. (2016). Time Course of Tactile Gating in a Reach-to-
1133 Grasp and Lift Task. J Mot Behav 48, 390–400.
1134 https://doi.org/10.1080/00222895.2015.1113917.
1135

1136 94. Juravle, G., Deubel, H., Tan, H.Z., and Spence, C. (2010). Changes in tactile
1137 sensitivity over the time-course of a goal-directed movement. Behavioural Brain
1138 Research 208, 391–401. https://doi.org/10.1016/j.bbr.2009.12.009.
1139

1140 95. Colino, F.L., Buckingham, G., Cheng, D.T., van Donkelaar, P., and Binsted, G. (2014).
1141 Tactile gating in a reaching and grasping task. Physiol Rep 2, 1–11.
1142 https://doi.org/10.1002/phy2.267.
1143

1144 96. Colino, F.L., Lee, J.H., and Binsted, G. (2017). Availability of vision and tactile gating:
1145 vision enhances tactile sensitivity. Exp Brain Res 235, 341–348.
1146 https://doi.org/10.1007/s00221-016-4785-3.
1147

1148 97. Rushton, D.N., Roghwell, J.C., and Craggs, M.D. (1981). Gating of somatosensory
1149 evoked potentials during different kinds of movement in man. Brain 104, 465–491.
1150 https://doi.org/10.1093/brain/104.3.465.
1151

1152 98. Cohen, L.G., and Starr, A. (1987). Localization, timing and specificity of gating of
1153 somatosensory evoked potentials during active movement in man. Brain 110, 451–467.
1154 https://doi.org/10.1093/brain/110.2.451.
1155

1156 99.   Pertovaara, A., Kemppainen, P., and Leppänen, H. (1992). Lowered cutaneous
1157   sensitivity to nonpainful electrical stimulation during isometric exercise in humans.
1158   Exp Brain Res *89*, 447–452. https://doi.org/10.1007/BF00228261.
1159

1160 100.   Reznik, D., Henkin, Y., Levy, O., and Mukamel, R. (2015). Perceived Loudness of
1161   Self-Generated Sounds Is Differentially Modified by Expected Sound Intensity. PLoS
1162   One *10*, e0127651. https://doi.org/10.1371/journal.pone.0127651.
1163

1164 101.   Paraskevoudi, N., and SanMiguel, I. (2021). Self-generation and sound intensity
1165   interactively modulate perceptual bias, but not perceptual sensitivity. Sci Rep *11*,
1166   17103. https://doi.org/10.1038/s41598-021-96346-z.
1167

1168 102.   Juravle, G., Colino, F.L., Meleqi, X., Binsted, G., and Farnè, A. (2018). Vision
1169   facilitates tactile perception when grasping an object. Sci Rep *8*, 15653.
1170   https://doi.org/10.1038/s41598-018-33916-8.
1171

1172 103.   McManus, M., Schütz, I., Voudouris, D., and Fiehler, K. (2023). How visuomotor
1173   predictability and task demands affect tactile sensitivity on a moving limb during
1174   object interaction in a virtual environment. R Soc Open Sci *10*, 231259.
1175

1176 104.   Knoetsch, F., and Zimmermann, E. (2021). The spatial specificity of sensory
1177   attenuation for self-touch. Conscious Cogn *92*, 103135.
1178   https://doi.org/10.1016/j.concog.2021.103135.
1179

1180 105.   Blakemore, S.-J., Smith, J., Steel, R., Johnstone, E.C., and Frith, C.D. (2000). The
1181   perception of self-produced sensory stimuli in patients with auditory hallucinations and
1182   passivity experiences: evidence for a breakdown in self-monitoring. Psychol Med *30*,
1183   1131–1139.
1184

1185 106.   Corlett, P.R., Horga, G., Fletcher, P.C., Alderson-Day, B., Schmack, K., and Powers,
1186   A.R. (2019). Hallucinations and Strong Priors. Preprint,
1187   https://doi.org/10.1016/j.tics.2018.12.001 https://doi.org/10.1016/j.tics.2018.12.001.
1188

1189 107.   Frith, C. (2005). The neural basis of hallucinations and delusions. C R Biol *328*, 169–
1190   175. https://doi.org/10.1016/j.crvi.2004.10.012.
1191

1192 108.   Frith, C.D., Blakemore, S.J., and Wolpert, D.M. (2000). Explaining the symptoms of
1193   schizophrenia: Abnormalities in the awareness of action. Brain Res Rev *31*, 357–363.
1194   https://doi.org/10.1016/S0165-0173(99)00052-1.
1195

1196 109.   Frith, C.D. (2019). Can a Problem With Corollary Discharge Explain the Symptoms of
1197   Schizophrenia? Biol Psychiatry Cogn Neurosci Neuroimaging *4*, 768–769.
1198   https://doi.org/10.1016/j.bpsc.2019.07.003.
1199

1200 110. Shergill, S.S., Samson, G., Bays, P.M., Frith, C.D., and Wolpert, D.M. (2005).
1201   Evidence for sensory prediction deficits in schizophrenia. American Journal of
1202   Psychiatry *162*, 2384–2386. https://doi.org/10.1176/appi.ajp.162.12.2384.
1203

1204 111. Shergill, S.S., White, T.P., Joyce, D.W., Bays, P.M., Wolpert, D.M., and Frith, C.D.
1205   (2014). Functional Magnetic Resonance Imaging of Impaired Sensory Prediction in
1206   Schizophrenia. JAMA Psychiatry *71*, 28.
1207   https://doi.org/10.1001/jamapsychiatry.2013.2974.
1208

1209 112. Frith, C. (2005). The self in action: lessons from delusions of control. Conscious Cogn
1210   *14*, 752–770. https://doi.org/10.1016/j.concog.2005.04.002.
1211

1212 113. Parees, I., Brown, H., Nuruki, a., Adams, R. a., Davare, M., Bhatia, K.P., Friston, K.,
1213   and Edwards, M.J. (2014). Loss of sensory attenuation in patients with functional
1214   (psychogenic) movement disorders. Brain *137*, 2916–2921.
1215   https://doi.org/10.1093/brain/awu237.
1216

1217 114. Kemp, K.C., Bathery, A.J., Barrantes-Vidal, N., and Kwapil, T.R. (2021). Positive,
1218   Negative, and Disorganized Schizotypy Predict Differential Patterns of Interview-
1219   Rated Schizophrenia-Spectrum Symptoms and Impairment. Assessment *28*, 141–152.
1220   https://doi.org/10.1177/1073191119900008.
1221

1222 115. Barrantes-Vidal, N., Grant, P., and Kwapil, T.R. (2015). The Role of Schizotypy in the
1223   Study of the Etiology of Schizophrenia Spectrum Disorders. Schizophr Bull *41*, S408–
1224   S416. https://doi.org/10.1093/schbul/sbu191.
1225

1226 116. Ford, J.M., Mathalon, D.H., Heinks, T., Kalba, S., Faustman, W.O., and Roth, W.T.
1227   (2001). Neurophysiological evidence of corollary discharge dysfunction in
1228   schizophrenia. American Journal of Psychiatry *158*, 2069–2071.
1229

1230 117. Ford, J.M., Gray, M., Faustman, W.O., Roach, B.J., and Mathalon, D.H. (2007).
1231   Dissecting corollary discharge dysfunction in schizophrenia. Psychophysiology *44*,
1232   522–529. https://doi.org/10.1111/j.1469-8986.2007.00533.x.
1233

1234 118. Ford, J.M., Mathalon, D.H., Roach, B.J., Keedy, S.K., Reilly, J.L., Gershon, E.S., and
1235   Sweeney, J.A. (2013). Neurophysiological evidence of corollary discharge function
1236   during vocalization in psychotic patients and their nonpsychotic first-degree relatives.
1237   Schizophr Bull *39*, 1272–1280. https://doi.org/10.1093/schbul/sbs129.
1238

1239 119. Rummell, B.P., Bikas, S., Babl, S.S., Gogos, J.A., and Sigurdsson, T. (2023). Altered
1240   corollary discharge signaling in the auditory cortex of a mouse model of schizophrenia
1241   predisposition. Nat Commun *14*, 7388. https://doi.org/10.1038/s41467-023-42964-2.
1242

1243 120. Thakkar, K.N., Diwadkar, V.A., and Rolfs, M. (2017). Oculomotor Prediction: A
1244   Window into the Psychotic Mind. Preprint at Elsevier Ltd,
1245   https://doi.org/10.1016/j.tics.2017.02.001 https://doi.org/10.1016/j.tics.2017.02.001.

121. Fletcher, P.C., and Frith, C.D. (2009). Perceiving is believing: a Bayesian approach to explaining the positive symptoms of schizophrenia. Nat Rev Neurosci *10*, 48–58. https://doi.org/10.1038/nrn2536.

122. Thakkar, K.N., Mathalon, D.H., and Ford, J.M. (2021). Reconciling competing mechanisms posited to underlie auditory verbal hallucinations: Mechanisms of Auditory Hallucinations. Philosophical Transactions of the Royal Society B: Biological Sciences *376*, 20190702. https://doi.org/10.1098/rstb.2019.0702.

123. Leptourgos, P., and Corlett, P.R. (2020). Embodied Predictions, Agency, and Psychosis. Front Big Data *3*, 1–13. https://doi.org/10.3389/fdata.2020.00027.

124. Whitford, T.J., Mathalon, D.H., Shenton, M.E., Roach, B.J., Bammer, R., Adcock, R.A., Bouix, S., Kubicki, M., De Siebenthal, J., Rausch, A.C., et al. (2011). Electrophysiological and diffusion tensor imaging evidence of delayed corollary discharges in patients with schizophrenia. Psychol Med *41*, 959–969. https://doi.org/10.1017/S0033291710001376.

125. Whitford, T.J., Ford, J.M., Mathalon, D.H., Kubicki, M., and Shenton, M.E. (2012). Schizophrenia, myelination, and delayed corollary discharges: A hypothesis. Schizophr Bull *38*, 486–494. https://doi.org/10.1093/schbul/sbq105.

126. Okimura, T., Maeda, T., Mimura, M., and Yamashita, Y. (2023). Aberrant sense of agency induced by delayed prediction signals in schizophrenia: a computational modeling study. Schizophrenia *9*, 72. https://doi.org/10.1038/s41537-023-00403-7.

127. Haß, K., Sinke, C., Reese, T., Roy, M., Wiswede, D., Dillo, W., Oranje, B., and Szycik, G.R. (2017). Enlarged temporal integration window in schizophrenia indicated by the double-flash illusion. Cogn Neuropsychiatry *22*, 145–158. https://doi.org/10.1080/13546805.2017.1287693.

128. Stevenson, R.A., Park, S., Cochran, C., McIntosh, L.G., Noel, J.-P., Barense, M.D., Ferber, S., and Wallace, M.T. (2017). The associations between multisensory temporal processing and symptoms of schizophrenia. Schizophr Res *179*, 97–103. https://doi.org/10.1016/j.schres.2016.09.035.

129. Foucher, J.R., Lacambre, M., Pham, B.-T., Giersch, A., and Elliott, M.A. (2007). Low time resolution in schizophrenia. Schizophr Res *97*, 118–127. https://doi.org/10.1016/j.schres.2007.08.013.

130. Ferri, F., Venskus, A., Fotia, F., Cooke, J., and Romei, V. (2018). Higher proneness to multisensory illusions is driven by reduced temporal sensitivity in people with high schizotypal traits. Conscious Cogn *65*, 263–270. https://doi.org/10.1016/j.concog.2018.09.006.

1292 131. Fotia, F., Cooke, J., Van Dam, L., Ferri, F., and Romei, V. (2021). The temporal
1293   sensitivity to the tactile-induced double flash illusion mediates the impact of beta
1294   oscillations on schizotypal personality traits. Conscious Cogn *91*, 103121.
1295   https://doi.org/10.1016/j.concog.2021.103121.
1296

1297 132. Marsicano, G., Cerpelloni, F., Melcher, D., and Ronconi, L. (2022). Lower
1298   multisensory temporal acuity in individuals with high schizotypal traits: a web-based
1299   study. Sci Rep *12*, 2782. https://doi.org/10.1038/s41598-022-06503-1.
1300

1301 133. Ferri, F., Nikolova, Y.S., Perrucci, M.G., Costantini, M., Ferretti, A., Gatta, V., Huang,
1302   Z., Edden, R.A.E., Yue, Q., D'Aurora, M., et al. (2017). A Neural "Tuning Curve" for
1303   Multisensory Experience and Cognitive-Perceptual Schizotypy. Schizophr Bull *43*,
1304   801–813. https://doi.org/10.1093/schbul/sbw174.
1305

1306 134. Poulet, J.F.A., and Hedwig, B. (2003). Corollary Discharge Inhibition of Ascending
1307   Auditory Neurons in the Stridulating Cricket. The Journal of Neuroscience *23*, 4717–
1308   4725. https://doi.org/10.1523/JNEUROSCI.23-11-04717.2003.
1309

1310 135. Poulet, J.F.A., and Hedwig, B. (2006). The Cellular Basis of a Corollary Discharge.
1311   Science (1979) *311*, 518–522. https://doi.org/10.1126/science.1120847.
1312

1313 136. Schneider, D.M., Sundararajan, J., and Mooney, R. (2018). A cortical filter that learns
1314   to suppress the acoustic consequences of movement. Nature *561*, 391–395.
1315   https://doi.org/10.1038/s41586-018-0520-5.
1316

1317 137. Chinta, S., and Pluta, S.R. (2023). Neural mechanisms for the localization of
1318   unexpected external motion. Nat Commun *14*, 6112. https://doi.org/10.1038/s41467-
1319   023-41755-z.
1320

1321 138. Audette, N.J., Zhou, W.X., La Chioma, A., and Schneider, D.M. (2022). Precise
1322   movement-based predictions in the mouse auditory cortex. Current Biology *32*, 4925-
1323   4940.e6. https://doi.org/10.1016/j.cub.2022.09.064.
1324

1325 139. Fukutomi, M., and Carlson, B.A. (2020). A History of Corollary Discharge:
1326   Contributions of Mormyrid Weakly Electric Fish. Front Integr Neurosci *14*.
1327   https://doi.org/10.3389/fnint.2020.00042.
1328

1329 140. Sawtell, N.B. (2017). Neural Mechanisms for Predicting the Sensory Consequences of
1330   Behavior: Insights from Electrosensory Systems. Annu Rev Physiol *79*, 381–399.
1331   https://doi.org/10.1146/annurev-physiol-021115-105003.
1332

1333 141. Wallach, A., and Sawtell, N.B. (2023). An internal model for canceling self-generated
1334   sensory input in freely behaving electric fish. Neuron *111*, 2570-2582.e5.
1335   https://doi.org/10.1016/j.neuron.2023.05.019.
1336

142. Roy, J.E., and Cullen, K.E. (2004). Dissociating Self-Generated from Passively Applied Head Motion: Neural Mechanisms in the Vestibular Nuclei. The Journal of Neuroscience *24*, 2102–2111. https://doi.org/10.1523/JNEUROSCI.3988-03.2004.

143. Brooks, J.X., Carriot, J., and Cullen, K.E. (2015). Learning to expect the unexpected: rapid updating in primate cerebellum during voluntary self-motion. Nat Neurosci *18*, 1310–1317. https://doi.org/10.1038/nn.4077.

144. Eliades, S.J., and Wang, X. (2008). Neural substrates of vocalization feedback monitoring in primate auditory cortex. Nature *453*, 1102–1106. https://doi.org/10.1038/nature06910.

145. Brooks, J.X., and Cullen, K.E. (2019). Predictive Sensing: The Role of Motor Signals in Sensory Processing. Biol Psychiatry Cogn Neurosci Neuroimaging *4*, 842–850. https://doi.org/10.1016/j.bpsc.2019.06.003.

146. Crapse, T.B., and Sommer, M.A. (2008). Corollary discharge across the animal kingdom. Nat Rev Neurosci *9*, 587–600. https://doi.org/10.1038/nrn2457.

147. Cullen, K.E. (2004). Sensory signals during active versus passive movement. Curr Opin Neurobiol *14*, 698–706. https://doi.org/10.1016/j.conb.2004.10.002.

148. Cullen, K.E. (2012). The vestibular system: multimodal integration and encoding of self-motion for motor control. Trends Neurosci *35*, 185–196. https://doi.org/10.1016/j.tins.2011.12.001.

149. Schneider, D.M., and Mooney, R. (2018). How Movement Modulates Hearing. Annu Rev Neurosci *41*, 553–572. https://doi.org/10.1146/annurev-neuro-072116-031215.

150. Straka, H., Simmers, J., and Chagnaud, B.P. (2018). A New Perspective on Predictive Motor Signaling. Current Biology *28*, R232–R243. https://doi.org/10.1016/j.cub.2018.01.033.

151. Gupta, K., Chowdhury, R.R., Chakrabarti, S., and Schwarz, C. (2023). Discerning state estimation and sensory gating, two presumptive predictive signals in mouse barrel cortex. bioRxiv.

152. Oldfield, R.C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. Neuropsychologia *9*, 97–113. https://doi.org/10.1016/0028-3932(71)90067-4.

153. R Core Team (2022). R: A Language and Environment for Statistical Computing. Preprint.

154. JASP Team (2023). JASP (Version 0.17).

1383 155. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A
1384   Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical
1385   Society: Series B (Methodological) *57*, 289–300. https://doi.org/10.1111/j.2517-
1386   6161.1995.tb02031.x.
1387
1388 156. Quintana, D.S., and Williams, D.R. (2018). Bayesian alternatives for common null-
1389   hypothesis significance tests in psychiatry: A non-technical guide using JASP. BMC
1390   Psychiatry *18*, 1–8. https://doi.org/10.1186/s12888-018-1761-4.
1391
1392 157. van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz,
1393   A., Evans, N.J., Gronau, Q.F., Haaf, J.M., et al. (2021). The JASP guidelines for
1394   conducting and reporting a Bayesian analysis. Preprint,
1395   https://doi.org/10.3758/s13423-020-01798-5 https://doi.org/10.3758/s13423-020-
1396   01798-5.
1397
1398