KATMAP: Inferring splicing factor activity and regulatory targets from knockdown data

Michael P. McGurk[1,*], David C. McWatters[1], and Christopher B. Burge [1,*]

[1]Massachusetts Institute of Technology, Department of Biology
[*]Direct correspondence to: mmcgurk@mit.edu and cburge@mit.edu

June 25, 2024

Abstract

It is well-established that splicing factors (SFs) regulate splicing in a position-dependent manner. However, despite extensive experimental work focused on this regulation, quantifying SF regulatory activity in a form that can be readily applied to new analyses remains a challenge. Here we present an interpretable regression model, KATMAP, which explicitly models splicing changes upon SF knockdown in terms of changes in SF binding across the transcriptome and the resulting loss of regulation. This allows us to learn an SF's positional activity and predict the factor's regulatory impact on individual exons, yielding regulatory models consistent with prior literature and highly reproducible across knockdowns of the same SF in different cell types, even generalizing well to related species. We leverage this generalizability to distinguish direct targets from the indirect effects of knockdown, and identify the secondary SFs responsible for these widespread indirect effects. We further uncover evidence of cooperative regulation between RBFOX and QKI. The generality of our statistical framework means it could be extended to the study of RNA regulation and processing beyond splicing.

Main

The development of a single cell into a complete organism, the physiology that underlies life, and many of the diseases that cut it short arise from how, when, and in what combinations genes are utilized. Orchestrating these gene expression programs are regulatory factors which recognize sequence elements distributed across a genome, coordinating the usage of many genes simultaneously. The transcriptomic differences between two biological states can be large, with hundreds or thousands of genes differing in splicing or expression, but these myriad differences typically result from a much smaller set of regulatory factors. Disentangling gene expression programs in terms of these regulatory factors and their targets is critical to understanding the genomic basis of normal physiology and responding to its disruption in disease. Doing so, however, requires determining the rules by which these factors exert their activities.

Of all the steps in eukaryotic gene expression, splicing stands out in its complexity and contribution to the functional diversity of the transcriptome. Pre-mRNA splicing is a multi-step process orchestrated by a collection of ribonucleoproteins (snRNPs) through a sequence of chemical reactions and physical rearrangements (Maniatis and Reed 1987). Canonically, this requires recognition of a 5' splice site (5SS) by U1 snRNP, a 3' splice site (3SS) by U2-associated factors, and a branch point (Lerner et al. 1980; Zamore and Green 1991; Zhuang and Weiner 1986). This is further modulated by additional splicing factors (SFs) that recognize splicing regulatory elements (SREs), allowing a single gene to generate dozens of distinct splicing isoforms (spliceoforms) often with distinct functions. SFs achieve this through a variety of mechanisms, but the effect of an SF typically depends on where it is bound relative to an exon's splice sites (Fig.1a, (Wang and Burge 2008)).

Over the years a number of strategies for using sequencing data to connect binding to the regulation of transcription and RNA processing have been developed. Some cast a wide net, aiming to simultaneously assign regulatory activity to hundreds or thousands of motifs or kmers without using prior knowledge of transcription (Balwierz et al. 2014; J. D. Rubin et al. 2021), stability (Krismer et al. 2020), or splicing factors (Bak et al. 2024; Hugh-White 2021). These approaches sidestep the need to tailor the model to a specific factor, but require discovering the putative activity associated with each motif every time they are applied. Others have been more focused, integrating knockdown and binding data for a given SF to obtain visual summaries of its position-specific regulation ("RNA maps") (Ule et al. 2006; Van Nostrand et al. 2020) or incorporating expert knowledge of the factor's sequence specificity and activity to predict its targets, using crosslinking/immunoprecipitation (CLIP) data and sequence conservation to supplement predictions of binding (Han et al. 2014; Weyn-Vanhentenryck et al. 2014; C. Zhang et al. 2010). These target prediction models can separate out the factor's direct targets from indirect effects of knockdown, but their reliance on crosslinking data ties the target predictions to where the SF was detectably bound in specific experimental data. Further, the challenge of crafting these models to the specific details of the SF's binding and activity has limited their application to well-studied factors.

We present a flexible and interpretable model for learning regulatory activity from knockdown or overexpression data. Our approach, KATMAP (Knockdown Activity and Target Models from Additive regression Predictions) requires as inputs only the results of a perturbation experiment and some model of the factor's RNA sequence specificity, such as a

position-weight matrix (PWM) (fig1b). We explicitly model the effect of knockdown by assigning changes in SF occupancy at potential binding sites throughout the transcriptome, learning from the splicing changes how to translate the similarity to the canonical motif into occupancy. Integrating these with the observed splicing changes, we infer an "activity map" describing the factor's position-specific regulatory activity (Fig. 1c). Where a traditional RNA map summarizes the enrichment of binding sites, our activity map instead describes the effect of binding. KATMAP uses these to predict how the splicing regulation of each exon in the transcriptome was affected by SF perturbation. Because our inferred models require only sequence to make new predictions they are not tied to the cellular context of the experimental data from which they were learned. We show that our models can be applied across cell types and related species to interpret observed splicing changes in terms of the underlying SFs and cis-elements, disentangle direct from indirect effects, and uncover evidence of cooperative regulation.

Results

Overview of the KATMAP model

The intuition guiding our model is as follows. Prior to a perturbation in the level of a sequence-specific splicing factor (SF), sequence motifs in the transcriptome are bound by the SF at varying levels of occupancy. We assume the factor can only influence the splicing reaction if its binding sites are properly situated relative to the exon and that binding elsewhere has negligible effect. After knockdown, the free protein concentration decreases, leading to reduced occupancy across the protein's binding sites. If binding is lost from a regulatory region, then loss of regulation results in a change in exon inclusion. The greater the decrease in binding, the more likely it is to affect inclusion. The logic is similar for SF overexpression, except that free protein concentration, occupancy, and regulation increase upon perturbation.

Following this logic, KATMAP models splicing changes in terms of changes in SF binding to the transcript. As inputs, our model requires only some information about a SF's sequence-specificity and the results of a differential splicing analysis which has labelled each exon as upregulated or downregulated or "not significant." The model aims to predict whether and in which direction each exon changes in splicing by answering two questions: 1) Are there binding sites near the exon which were affected by the SF knockdown? 2) Where must the SF bind near an exon to regulate its inclusion? We frame this as a generalized additive model which represents SF binding and regulation with its nonlinear terms but accounts for other features predictive of splicing changes through additional linear terms.

Our model begins by scoring all potential binding motifs around an exon's splice sites, extending 200 nt on the intronic and 60 nt on the exonic side, using position weight matrices (PWMs), position-specific affinity matrices (PSAMs), or other affinity models (Fig. 1d). These *motif scores* are then transformed to changes-in-binding using a *binding function* based on a biophysical description of how occupancy changes after a decrease in SF availability (Fig. 1e, f). The shape of the binding function is controlled by a *most-relevant-affinity* parameter that defines which binding sites were most affected by the knockdown (i.e. shifts

the function along the X-axis) and an *affinity scale* parameter which defines the range of motif scores affected by the knockdown (i.e. widens/narrows the function); we refer to these collectively as the *binding parameters* (see Methods).

We represent the SF's position-specific regulatory activity with a set of parameters termed the *activity map* (Fig. 1g), which assigns enhancing (> 0) or suppressing (< 0) regulatory activity to different positions on the intronic and exonic sides of 3SS and 5SS. These regulatory activities are shared across all exons in the analysis. Critically, we assume that activity varies smoothly with position, with discontinuities at the exon-intron boundaries that allow exonic and intronic binding to have distinct activities; how smoothly depends on two *spatial correlation* parameters. We multiply each change-in-binding by the corresponding activity to determine its impact on the splicing outcome (Fig. 1h). The sum of these products defines the SF's *splicing impact* on the exon, making the simplifying assumption that regulation is additive.

Finally, we incorporate these splicing impacts into a regression framework which assigns each exon a probability of being downregulated, unchanged, or upregulated (Fig. 1i). In addition to SF binding, the unperturbed level of exon inclusion and gene expression impact our statistical power to detect splicing changes. Specifically, exons with inclusion levels (PSI values) near the boundaries of 0% or 100% inclusion are less likely to be detected as down- or upregulated, respectively, while power to detect splicing changes is limited for exons in low-expression genes. We include these confounding variables (e.g. unperturbed inclusionlevel, expression) and their squares as linear predictors, whose effects are controlled for by a set of *linear coefficients*.

This model architecture answers the two questions we posed in an integrated fashion. SF binding and regulatory activity of the perturbed SF are modeled simultaneously, without the need to define arbitrary cutoffs for binding or activity. The activity map provides a simple description of the SF's position-specific activity, identifying where a binding site must occur to influence an exon's inclusion positively or negatively, an intuitive notion of the SF's function, which substitutes for a traditional RNA map. Assuming that this activity changes smoothly with position (except at exon-intron boundaries) is both biologically plausible (Graveley, Hertel, and Maniatis 1998) and improves statistical power by pooling information spatially (Supplementary fig. 1a,b). How smoothly is automatically learned from the data.

The goal of KATMAP's inference algorithm is to determine which parameter values provide good explanations for the observed splicing changes. Prior target prediction models obtained point estimates of the most likely parameter values (C. Zhang et al. 2010), but we wished to quantify uncertainty by identifying the range of plausible hypotheses. We therefore used Bayesian inference to precisely evaluate the plausibility of hypotheses given observed data. We structured KATMAP's model such that posterior inference could be split into two tractable subproblems. Loosely speaking, the parameters used to denote binding (*most-relevant-affinity* and *affinity-scale*) and spatial correlation can be considered *hyperparameters*. If the hyperparameters are held constant, the model simplifies to a generalized linear model (GLM), such that the dozens of *activity* and *linear* coefficients can be identified by straightforward optimization (Rue, Martino, and Chopin 2009). To leverage this property, we couple adaptive importance sampling with integrated nested Laplace approximations (INLA) (Rue, Martino, and Chopin 2009), dividing posterior inference into: 1) an inner step that solves conditional models to learn which activity and linear coefficients best

explain the data assuming particular binding and spatial correlation parameter values to be true; and 2) an outer step that evaluates a marginal model to learn which binding and spatial correlation parameter values are consistent with the data. This approach yields samples from the joint posterior that span the range of plausible parameter values (Fig.1j), allowing us to compute point estimates of the most plausible parameters, quantify our uncertainty with credible intervals, and propagate this uncertainty forward into our predictions of regulation.

KATMAP accurately infers parameters and quantifies uncertainty

To validate our inference strategy, we simulated 21 SF knockdown datasets with known parameter values and found that KATMAP robustly recovers the true parameters (Fig. 1k, Supplementary fig. 1 c,d). Importantly, our posterior samples properly quantify the uncertainty regarding the activity and binding parameters (Fig. 1l, Supplementary fig. 1 e). Because the activity and binding parameters are accurately recovered, so are the splicing impacts used to simulate the splicing changes (Supplementary fig. 1f). Linear coefficients are slightly underestimated, likely due to the multivariate Gaussian approximation to their conditional posterior (Rue, Martino, and Chopin 2009). These parameters serve primarily to control for potential confounders, and this subtle bias is unlikely to interfere with this.

To assess performance on real data, we selected ten knockdowns with many splicing changes. For each experiment, we trained on 70% of the differentially spliced exons, holding out the remaining 30% as a test set. We evaluated our models against these held-out exons, comparing the true splicing changes to our predictions as to which exons were unchanged, up- or downregulated upon knockdown. In all ten evaluations, the predicted probabilities were well-calibrated, with the probability that an exon was affected by knockdown in a given direction corresponding to the frequency of that outcome (Fig 1m, Supplementary fig. 1g)

KATMAP robustly infers splicing regulatory activity

We applied KATMAP to knockdown data from the ENCODE project for 35 SFs for which we also had good models of the protein's sequence specificity (Supplementary Table 1) (Van Nostrand et al. 2020). We used RNA-bind-n-seq-derived position-specific affinity matrices (PSAMs) as the binding model for 23 SFs (Jens et al. 2022; Lambert et al. 2014), and RNAcompete-derived position weight matrices (PWMs) for the remaining 12 (Ray, Kazan, Chan, et al. 2009; Ray, Kazan, Cook, et al. 2013). Eighteen SFs showed significant enhancing or repressing activity maps and outperformed a reduced model (with splicing activity excluded) in at least one knockdown, with 9 displaying enhancing activity, 6 showing repressing activity, and 3 with both enhancing and repressing activity (Fig. 2a, Supplementary Data 1). Among these eighteen, 14 had available knockdown/RNA-seq data from both lymphocyte-derived K562 cells and hepatocyte-derived HepG2 cells. For all but one factor (SF1), the activity maps were consistent between the two knockdowns, highlighting the reproducibility of our inferences (Fig. 2b).

The evidence KATMAP identified for splicing activity correlated with knockdown efficiency ($r=0.45$, Supplementary fig. 1fh), with stronger knockdowns providing more evidence for splicing activity. Similarly, a sufficient number of splicing changes was required to infer activity, with evidence of enhancing or repressing activity only identified in knockdowns

Figure 1: **Illustration of inference with KATMAP's model. a)** An example of position-dependent regulation. The depicted factor (red) enhances the inclusion of an exon when bound downstream and suppresses inclusion when bound upstream. **b, c)** Schematic representations of the inputs and outputs of the KATMAP model. **d-g)** Examples of the model applied to three real exons in the *RBFOX2* (HepG2) knockdown. **d)** RBFOX2 motif scores assigned to sequences around the 3' and 5'ss. **e)** The binding function that transforms the scores into relative changes in binding. **f)** Predicted changes in binding obtained by passing the motifs scores (d) through the binding function (e). **g)** An activity map describing the effect on splicing of RBFOX2 binding at positions near the 3' and 5'ss. Discontinuities at the splice sites allow activity to differ between introns and exons. **h)** The predicted splicing impacts, computed by multiplying the binding changes (f) by the activity map (g). **i)** The predicted probabilities of being downregulated ($\downarrow$), unchanged (−), or upregulated ($\uparrow$) for these exons and the observed splicing changes. **j)** A posterior sample of 200 activity maps for the RBFOX2 (HepG2) KD. **k)** True activities used in the 20 simulated datasets versus the values (with 94%-CI) recovered by inference. Only four activities per simulation are shown to ensure independence of the plotted values. **l)** Coverage plot of how frequently credible intervals contain the true activity, for different CI-levels. **m)** Calibration plot for the PUF60 (K562) knockdown, comparing average $Pr(Downreg)$ and fraction truly downregulated for sets of 200 held out exons. The blue line is a binomial regression fit.

Supplementary Figure 1: **Assessment of KATMAP's performance. a, b)** The activity maps inferred from an RBFOX2 knockdown in HepG2 cells with priors assuming independence among the activity co-efficients (**a**) or with our full model's assumption of spatial correlation (**b**). The dots represent posterior means and the bars are 94%-credible intervals. **c, d)** The true hyperparameters (**c**) and linear coefficients (**d**) used in the simulation versus the values recovered by inference, summarized by posterior means with 94%-credible intervals. **e)** Coverage plots of how frequently credible intervals contain the true hyperparameter or linear coefficient, for different CI-levels. **f)** An example comparing splicing impacts used to simulate splicing changes and the splicing impacts recovered by inference. **g)** The slopes and intercepts of binomial regressions evaluating whether the downreg (∇), unchanged ($-$), and upreg ($\triangle$) probabilities correctly describe the splicing changes at held out exons (see Fig. 1m) for ten knockdowns. In all evaluations, the slopes are near 1 and the intercepts near 0, indicating that KATMAP's probabilities are well-calibrated and generalize out of the training set. **h)** The relationship between the efficiency of an ENCODE knockdown as shrinkage estimate of the gene's of log2 fold-change in the DEseq analysis. **i-k)** The numbers of differentially spliced exons in knockdowns where KATMAP identified enhancing (**i**) or suppressing (**j**) activity, or activity in either direction (**k**).

where at least 100 exons were down- or upregulated, respectively (Supplementary fig. 1i-k). For example. both ENCODE knockdowns the splicing activator *TRA2A* yielded < 100 downregulated exons and no clear evidence of activity–perhaps because of compensation by paralog *TRA2B*–, but applying our model to a *TRA2A/B* double knockdown experiment with > 1000 downregulated exons (Best et al. 2014) recovered strong evidence of the expected exonic enhancing activity (Fig. 2c).

Examining the significant activity maps identifies three distinct groups of splicing activators. First, we identify SRSF1, SRSF7, and TRA2A/B as exhibiting the classic exonic enhancing pattern of SR protein regulation (Fig. 2a,c). Second, several polypyrimidine-binding factors show enhancing activity directly upstream of the 3SS, consistent with polyY tract definition. For U2AF2–which canonically defines the polyY tract (Merendino et al. 1999)–and its homolog PUF60 (Page-McCaw, Amonlirdviman, and Sharp 1999) we infer intronic enhancing activity just upstream of the 3SS (Fig. 2a,d). U2AF2's inferred activity is constrained to within 20 nt of the 3SS, while PUF60's activity is slightly more upstream and extends 40-60 nt into the intron, consistent with observations of extended polyY tracts at PUF60-responsive exons (Královičová et al. 2018). Similar upstream enhancing activity is inferred for PCBP1 (Fig. 2a), but with additional activity downstream of the 5SS, consistent with prior observations of C-rich motifs up- and downstream of PCBP1- and PCBP2-activated exons (Ji et al. 2016). KATMAP failed to detect the expected enhancing activity of the branch point-binding factor SF1 and the exonic enhancing activity of SRSF5, perhaps due to indirect effects of other SFs obscuring the signal in these knockdowns.

While upstream activators display greatest activity near the 3SS, a third group of factors with downstream enhancing activity–DAZAP1, HNRNPF/H, QKI, and RBFOX–have their greatest activity dozens of nucleotides away from the 5SS and influence splicing from a wider intronic window (Fig. 2a). Among these, two pairs of factors are strikingly similar in their activity. The related poly-G binding factors HNRNPF and HNRNPH1 both enhance inclusion when bound in the downstream intron while suppressing inclusion in the exon. HNRNPH1 was one of the first SFs demonstrated to activate and repress inclusion in a position-dependent manner (Chen, Kobayashi, and Helfman 1999; Chou et al. 1999). RBFOX2 and QKI also share patterns of activity (Fig. 2e,f)–enhancing downstream while suppressing upstream–but, unlike HNRNPF/H, are not related and bind distinct motifs (YGCAUG versus ACUAAC).

In contrast to splicing activators, all inferred repressors displayed activity in multiple regions (Fig. 2a). The PCBP-related HNRNPK shows silencing activity upstream of the 3SS and in the exon (Fig. 2g), consistent with its role as splicing repressor (Cao et al. 2012) and observations that PCBP2 and HNRNPK can regulate the same exon in opposing directions (Ghanem et al. 2018). We find a similar pattern of upstream and exonic silencing activity for PTBP1 (polypyrimidine tract binding protein), as well as evidence of downstream enhancing activity (Fig. 2a), all of which is consistent with prior studies (Llorian et al. 2010). This exonic activity of HNRNPK and PTBP1 stands in contrast to the pyrimidine-binding upstream activators–which lacked exonic activity. The other repressors have even wider regions of suppression. HNRNPC's silencing activity is strongest in the ~ 100 nt up- and downstream of the exon, but has two additional regions of silencing activity at $+/-$ 200 nt (Fig. 2h). HNRNPC is know to form tetramers, with repression suggested to result from looping out the exon, which may explain the multiple regions of activity (Zarnack et al.

2013). HNRNPL activity maps also display both up- and downstream suppressing activity (Fig. 2a), but further suppresses within the exon, consistent with prior work (Heiner et al. 2010; Rothrock, House, and Lynch 2005). MATR3 (Matrin3) has even broader regions of suppression, extending throughout both up- and downstream introns in the HepG2 model, but only showing evidence of upstream repression in K562 cells (Fig. 2a). This broad silencing activity is consistent with CLIP-based enrichment maps of MATR3-repressed exons which found MATR3 uniformly distributed out to at least 500 nt in the flanking introns (Coelho et al. 2015).

The diverse position-dependent patterns of splicing regulation we uncover underscore the power and flexibility of KATMAP's approach to describing activity. Assuming only that activity varies smoothly with distance (and learning how smoothly from the data) allowed KATMAP to detect both broad and narrow ranges of activity, within, upstream, or downstream of target exons. We were able to robustly and reproducibly uncover this activity by modelling the sequence of exons and without relying on crosslinking data, avoiding the bias toward exonic signal of eCLIP-based RNA maps ((Van Nostrand et al. 2020)).

Activity maps define regulatory targets

Inferring the activity map of an SF leads to a natural definition of its direct targets as exons with motifs of appropriate strength and location for regulation. For each exon, we predict the change in regulation due to knockdown as "splicing impacts" computed by multiplying predicted changes in binding by the SF's activity map (Fig. 1f-h, Supplementary Table 2). A positive impact represents a loss of suppression and increases the model's prediction that the exon will be upregulated, while a negative impact favors a prediction of downregulation. We consider an exon to be a direct suppression or enhancement target if the splicing impact of the exon is sufficiently positive or negative to improve predictions over a reduced model that excludes all splicing activity (Fig. 3a).

Comparing these predictions to the literature recovers previously described targets. For example, PTBP1 is known to silence exons in its paralogs *PTBP2* (Boutz et al. 2007) and *PTBP3*(Spellman, Llorian, and Smith 2007) during development, as well as in *RTN4*(Cheung et al. 2009) and *FLNA*(X. Zhang et al. 2016), all of which were identified as direct targets of PTBP1 by KATMAP. To assess our target predictions more comprehensively, we leveraged HNRNPC's known role as a silencer of exons derived from the highly abundant transposable element *Alu*. These *Alu*-derived exons provide a clear ground-truth for validating our target predictions. Roughly 30% of all exons upregulated upon HNRNPC knockdown are *Alu*-derived. KATMAP's target predictions recover HNRNPC's known silencing of *Alu*-exons, strongly sorting *Alu*-derived exons into the set of targets, with few being labelled nontargets (Fig. 3b). That only 30% of exons upregulated upon knockdown are *Alu*-derived, compared to 50–60% of our predicted HNRNPC targets, highlights the extent to which measured splicing changes are diluted by the indirect effects of knockdown. That only ∼5% of the predicted nontargets are *Alu*-derived underscores the effectiveness of KATMAP at winnowing out these indirect effects from the SF's true targets.

Figure 2: **KATMAP robustly infers splicing activity a)** Significant activity maps and their motifs organized by hierarchical clustering. For ease of comparison, activity is normalized relative to the maximum absolute activity in each map. If the binding model included multiple motifs, only the top motif is shown. **b)** The correlation between activity maps for the same RBP inferred from different knockdowns. **c-h)** Activity maps for TRA2A/B (**c**), PUF60 (**d**), RBFOX2 (**e**), QKI (**f**), HNRNPK (**g**), HNRNPC (**h**). The blue dots indicate the posterior mean and and the lines covering the 98%-credible intervals.

Figure 3: **KATMAP predicts regulatory targets a)** The probabilities of downregulation predicted by the full model and a reduced model lacking splicing impacts for exons downregulated upon RBFOX2 knockdown (HepG2). Exons where the full model assigns a higher probability than the reduced model are likely enhancing targets, while those assigned the same predictions by both models are likely nontargets. The axes are in log-odds scale and depicted the range is 0.01–0.99. **b)** The fraction of exons derived from *Alu* among targets of HNRNPC's suppression, nontargets, or all exons that changed in a given direction, computed for each of the three splicing categories. **c)** The enrichment of eCLIP peaks in differentially spliced targets compared to nontargets differentially spliced in the same direction. The enrichment summarizes eCLIP peaks that span the significant regions of a given SF's activity map. **d)** The correlation in splicing impacts of exons across the human transcriptome computed from the HepG2 and K562 models for SFs with data for both cell types. **e)** Comparing RBFOX splicing impacts in the mouse transcriptome computed using the human RBFOX2 (HepG2) model and a model inferred from mouse RBFOX1/2/3 triple knockout data. **f)** The difference in average inclusion-level change between confident targets and confident nontargets among exons that were significantly altered by knockdown (red, blue) and those that were not (grey). The 94%-credible intervals were obtained via Bayesian bootstrapping. **g)** ECDF plots of the distributions of inclusion-level changes at targets and nontargets upon HNRNPC knockdown (HepG2) for exons significantly upregulated upon knockdown (left) and nonsignificant exons (right). It is the difference in the means of the target and nontarget distributions that is shown in **f**). We excluded exons where the data supported a confidently small effect of knockdown (spike at 0, $|\Delta\psi| < 10^{-3}$) to focus on those exons called nonsignificant potentially due to a lack of statistical power; the estimates with all exons included are shown in Supplementary fig. 2g

Predicted targets are bound *in vivo* and generalize across biological contexts

Knockdown and crosslinking experiments are specific to the cell types and conditions in which the experiment was performed, but if a model successfully captures regulatory activity, its predictions should generalize to other biological contexts. While our predictions are based on sequence similarity to motifs identified *in vitro* and do not incorporate crosslinking data, intersecting predicted targets with eCLIP peaks can be used to confirm binding in living cells. For the 11 SFs with credible activity maps which also had ENCODE eCLIP data, we asked whether differentially spliced target exons were enriched for *in vivo* binding compared to differentially spliced nontargets. For example, QKI is predicted to suppress inclusion upstream of the exon and enhance inclusion downstream: ∼70% and ∼80% of differentially spliced suppression and enhancing targets, respectively, had upstream and downstream QKI eCLIP peaks, compared to $< 5\%$ and $< 15\%$ of up- and downregulated nontargets (Supplementary fig. 2a). Of the 11 considered SFs, 9 showed clear enrichment for *in vivo* binding at targets, with most showing between 2- to 8-fold enrichment (Fig. 3c).

To assess whether our models can generalize across contexts, we considered SFs with knockdowns in distinct cell types and compared their predictions. For 12 of the 14 factors, models inferred from HepG2 and K562 cells yielded highly similar predictions of regulation (Fig. 3d), indicating that KATMAP consistently identifies the same underlying relationship between binding motifs and splicing regulation. To assess whether this generalizability extends across species, we applied KATMAP to mouse *Rbfox1/2/3* triple knockout data (Jacko et al. 2018). The inferred activity map (Supplementary fig. 2b) was highly consistent with those obtained from the human *RBFOX2* knockdowns (Fig. 2e, Supplementary fig.2c). Using the human RBFOX models to score exons in the mouse transcriptome predicts essentially the same regulation as the splicing impacts learned directly from mouse neuronal *Rbfox1/2/3* triple-knockout data (Jacko et al. 2018) (Fig. 3e, Supplementary fig. 2d). Together, these analyses indicate that the models KATMAP infers can be applied to different tissues and for related species, where *in vivo* binding data is absent or genes and exons not expressed in the original knockdown experiment are present.

KATMAP identifies regulatory targets missed by differential splicing analyses

Among KATMAP's predicted targets are exons that were not called significantly affected in the differential splicing analysis but have binding motifs properly situated to affect splicing. Some such exons might be true targets, for which the RNAseq data did not provide sufficient power to confidently detect a splicing change. We asked, therefore, whether those exons predicted to be targets, but not called differentially spliced, had inclusion changes in the expected direction.

Our models were trained to predict the directions–but not the magnitudes–of changes in exon inclusion, so we first assessed whether our inferred splicing impacts were also predictive of magnitudes of changes. We found that among differentially spliced exons those predicted to be direct targets had larger changes in inclusion following knockdown than nontargets (Fig. 3f). For example, among exons upregulated upon HNRNPC knockdown, those with

Supplementary Figure 2: **Additional validation of target predictions a)** The fraction of significant exons in the QKI (HepG2) analysis with QKI eCLIP peaks spanning a given position for upregulated suppression targets (blue line) and downregulated enhancing targets (red line). The regulatory activity associated with each position is indicated by the background color. **b)** The RBFOX activity map inferred from mouse *Rbfox1/2/3* knockdown data **c)** The nonsignificant human RBFOX2 activity map inferred from the K562 knockdown. **d)** Splicing impacts computed for the mouse exons using the nonsignificant human RBFOX2 (K562) model compared to those learned directly directly mouse RBFOX1/2/3 triple knockdown data. **e, f)** The relationship between the change in the $\log_2$-odds of inclusion, $\Delta\phi$, and splicing impact for exons called significantly upregulated (**e**) and those not called significantly affected (**f**) by HNRNPC knockdown in HepG2. Red and grey shaded regions indicate confident targets and nontargets respectively. **g)** The difference in average inclusion-level change between confident targets and confident nontargets among exons called nonsignificant with the spike at 0 in **f**, fig3g; $|\Delta\psi| < 10^{-3}$–included in (red) or excluded from the averages (pink). The 94%-credible intervals were obtained via Bayesian bootstrapping.

larger splicing impacts tended to have larger changes in inclusion (Fig. 3g, Supplementary fig. 2e), with the average effect of knockdown being two-fold higher at upregulated HNRNPC targets compared to upregulated nontargets.

We find this same relationship among the exons not called differentially spliced–the exons KATMAP predicted to be direct targets changed, on average, in the expected direction compared to nontargets–consistent with the presence of true regulatory targets missed by the initial differential splicing analysis (Fig. 3f,g, Supplementary fig. 2f). An exon might not be called significantly affected by knockdown for two reasons: 1) the data provided clear evidence that inclusion did not change; or 2) the data did not provide sufficient evidence to distinguish an inclusion change from noise. The first scenario can be readily identified on the basis of confidently small $\Delta\Psi$ (Fig. 3g, Supplementary fig. 2f). Among the remaining exons, KATMAP's target predictions are able to hone in on the subset most likely to be truly regulated by the SF, as evidenced by the propensity for change in the expected direction. By bringing to bear information about the SF's activity and binding sites, KATMAP's regulatory models increase the power to flag potential regulatory targets that would otherwise be ignored using conventional approaches.

Validating inferred *cis*-regulatory element activity

Implicit in KATMAP's predictions are statements about cis-regulatory elements: for any predicted target, we can examine the collection of predicted SREs. For example, QKI is predicted to enhance the inclusion of *NF2* exon 16, which is one of two stop-codon-containing exons in the gene, the skipping of which suppresses tumor growth 3- to 4-fold in schwannoma (Sherman et al. 1997). Examining the position-wise splicing impacts suggests that QKI enhances its inclusion through a single ACUAA motif 53 nt downstream of the 5' splice site (Fig. 4a). Other targets are predicted to be controlled by multiple binding sites. For example, HNRNPC is predicted to suppress the inclusion of a highly conserved exon in own 5' UTR through multiple poly-U motifs in both the up- and downstream introns (Fig. 4b).

We used this interpretability to validate selected predictions using a minigene splicing assay. We chose target exons for five splicing factors–DAZAP1, HNRNPC, PCBP1, QKI, and RBFOX2–and evaluated the effects of mutating their predicted SREs on splicing. For targets with both up- and downstream SREs, we considered the effects of mutating only the upstream or the downstream elements. We inserted the reference and mutated exons into reporters and measured inclusion (Fig. 4c). To confirm that the effects of mutation were mediated by the expected SF, we assayed reporters with and without knockdown of the expected factor.

Inclusion changed in the expected direction for seven of the eight mutant reporters (Fig. 4d). Mutation of the QKI and RBFOX2 binding sites downstream of the NF2 and EPB41 exons, respectively, lead to decreased inclusion comparable to knockdown. Similarly, disrupting the DAZAP1 binding sites upstream of the MBNL1 exon decreased inclusion as expected, but to a greater extent than the knockdown. One possibility is that the mutations introduced a binding site for a different SF; in line with this, our HNRNPC model predicts the creation of a good HNRNPC binding site in a region where it could suppress inclusion. Both the upstream and downstream HNRNPC mutations behaved as expected, with either being sufficient to increase inclusion +40% to near complete inclusion. The PCBP1

Figure 4: **Validation of selected targets by minigene reporter assay a,b)** The expected changes in regulation due to loss of QKI (**a**) and HNRNPC (**b**) binding in predict targets. The blue arrows indicate the change in regulation from the pre-knockdown levels (tail of arrow) to the post-knockdown (grey bar). **c)** Schematics of the minigene reporter design. **Top:** The expected occupancy at the predicted splice regulatory elements (i–iii from **a**) and the predicted effects of mutating these elements. The reference sequence is on the X-axes with the expected binding motif capitalized; the specific mutations are below. **Bottom:** The reference and mutated versions of predicted targets were inserted into a splicing reporter. Inclusion was assayed with and without SF knockdown. **d)** The results of the reporter assay. The large arrow depicts the direction the mutations were expected to alter inclusion. The thin arrows depict the observed effect of knockdown, mutation, and mutation+knockdown.

downstream mutations decreased inclusion as expected, but to a smaller degree than the knockdown, possibly because the three upstream PCBP1 sites were intact. The upstream PCBP1 mutations are one case where the inclusion change was opposite to our expectation. In this case, the mutations occurred very close to the 3SS and are predicted to increase the strength of the PPT, which may override any impact of the loss of PCBP1 binding.

Target predictions reveal cooperative regulation

Our model makes the simplifying assumption that SFs act independent of each other. However, we reasoned that we could uncover evidence of cooperative regulation by examining our model's predictions for deviations from this null assumption. To evaluate this, we focused on RBFOX and QKI, which have strikingly similar activity maps despite binding distinct motifs (Fig. 2e,f), raising the possibility of a functional relationship between them. Earlier analyses found that QKI and RBFOX knockdowns affect a common subset of exons (Brosseau et al. 2014; Van Nostrand et al. 2020), and further noted enrichment of QKI eCLIP signal at RBFOX-downregulated exons. However, clear evidence of both QKI and RBFOX binding at the same exons was not observed and it was concluded that the correlated effects of knockdown did not represent direct coregulation by the factors (Van Nostrand et al. 2020). These changes could arise without direct coregulation, for example perturbing RBFOX might secondarily perturb QKI and indirectly affect its targets. These analyses, however, could not distinguish between the direct and indirect effects of knockdown, a limitation that might have obscured signatures of coregulation.

We used KATMAP's ability to discriminate direct regulatory targets from indirect effects to reassess the case for direct coregulation. In the scenario of strong cooperativity, knocking down either factor should primarily affect exons bound by both factors, whereas exons bound by only one should be unaffected. As KATMAP's predictions model one SF at a time, an exon bound by the knocked down factor will be called a target, whether or not its coregulator is present. The way in which cooperative regulation deviates from our assumptions should produce a clear signature at predicted targets: knockdown-affected targets should be enriched for targeting by both coregulators compared to targets unaffected by knockdown (Fig. 5a,b). We detected this expected signature of cooperativity, with differentially spliced QKI targets being enriched for RBFOX enhancing regulation (Fig. 5c). The reciprocal relationship was also observed: RBFOX2 targets affected by *RBFOX2* knockdown are enriched for QKI binding sites (Fig. 5d). Further, this pattern is evident in mouse data where all three RBFOX paralogs (*Rbfox1/2/3*) were knocked down (Fig. 5e), supporting that coregulation by RBFOX and QKI proteins is conserved out to mice (∼90 mya).

These coregulated exons have both QKI and RBFOX binding sites (Fig. 5f)–frequently multiple of each–occurring as clusters of interspersed motifs separated by a few dozen nt (Fig. 5g). This proximity of binding sites suggests that QKI and RBFOX proteins could physically interact, which is supported by prior yeast two-hybrid (Lang et al. 2021) and pull-down mass spectrometry data (Huttlin et al. 2021). Further, reassessment of the eCLIP data focused on direct targets supports binding of both QKI and RBFOX to the same exons: QKI crosslinking is detected at both knockdown-affected and unaffected QKI targets, but RFBOX2 is primarily bound at the targets affected by QKI knockdown. Together, these observations support direct, cooperative coregulation of exons by QKI and RBFOX proteins.

Figure 5: **QKI and RBFOX cooperatively regulate a splicing subprogram a)** In the case of cooperativity, predicted targets affected knockdown should be enriched for the coregulator compared to those unaffected by knockdown. **b)** The expected distributions of the coregulator's splicing impact computed at KD-affected (orange) and KD-unaffected targets (black). **c-e)** Distributions of RBFOX2 (**c**) and QKI (**d,e**) splicing impacts at the targets of the other protein (**c**: QKI; **d,e**: RBFOX), comparing the KD-affected (orange) to KD-unaffected targets (black). **f)** Locations of predicted binding sites (|rel. change-in-binding| > 20%) for both RBFOX (red) and QKI (blue) in KD-affected QKI enhancing targets affected (**top**) and an equal number of KD-unaffected targets (**bottom**). The exons are sorted by RBFOX splicing impact and the bars to the right denote the presence of eCLIP peaks in the downstream intron. **g)** The distributions of distances between the nearest QKI and RBFOX sites in KD-affected (orange) and unaffected (black) targets. **h)** GO-analysis of confident QKI-RBFOX cotargets; red denotes GO-membership.

Our analyses identified twenty human exons with binding sites for both QKI and RBFOX proteins and responsiveness to either *QKI* or *RBFOX2* knockdown (Supplementary Table 3). Of these twenty, eleven are in genes involved in cytoskeleton binding and organization, with seven being actin-binding proteins (Fig. 5h). This represents not only strong enrichment for these categories relative to expression-matched genes, but also relative to the full set of RBFOX or QKI targets, consistent with a focused subprogram active only in cells where both QKI and RBFOX are expressed and distinct from their individual regulatory roles. Cytoskeletal remodelling and cell migration are integral to early neuronal development, and a number of these exons are in genes with known roles neuronal growth and development. In particular, NDEL1 is critical for cell migration during neural differentiation (Youn et al. 2009) and NF2 exon 16 skipping promotes glial tumor growth 3–4 fold (Sherman et al. 1997). Both RBFOX and QKI proteins are expressed in neuronal progenitor cells (NPCs), which differentiate into both neurons – where RBFOX predominates – and glia – where QKI is predominant – suggesting a potential venue for coregulation (Hardy 1998; X. Zhang et al. 2016).

Indirect effects can be explained by secondary RBPs

The set of splicing changes observed in RNAseq datasets following perturbations such as stress, drug treatments, or cell differentiation are expected to result from changes in the activity of one or more splicing factors. Identifying these causal factors would provide paths to elucidating the underlying regulatory network. While the ENCODE RBP knockdowns offer a resource to accomplish this, the splicing changes in the knockdowns are themselves mixtures of direct and indirect effects. Indeed, most knockdowns caused abundant splicing changes in both directions, even for established activators (e.g. SRSF1) and repressors (e.g. HNRNPL) of splicing, where targets should be down- and upregulated respectively. We infer that most splicing changes in the ENCODE knockdowns are indirect effects, with less than 30% of splicing changes directly caused by loss of the knocked down RBP (Fig. 6a). The high proportion of indirect effects likely results from the multi-day period of antibiotic selection common to knockdown experiments, providing considerable time for secondary effects to manifest.

KATMAP distills knockdown data into predictions about direct regulation across the transcriptome. If a given SF contributed to the splicing changes in an RNAseq dataset, the splicing impacts we learned from our analyses of its knockdown data should be predictive of which exons were affected. We express this logic with a multinomial logistic regression that predicts each exon's splicing changes based on the splicing impacts inferred from the different ENCODE knockdowns (Fig. 6b). To validate this approach, we used the splicing models inferred from K562 knockdowns to explain splicing changes in the HepG2 knockdowns (and *vice versa*). This approach correctly identified the knocked down SF as most predictive in 21 out of 23 knockdowns (Fig. 6c), demonstrating that our regulatory models can generalize beyond their training data to infer the SFs underlying transcriptomic changes.

In most knockdown experiments, this analysis additionally implicated secondary SFs as responsible for some indirect effects (Fig. 6d,e, (Supplementary Table 4)). For example, in the *HNRNPC* (K562) knockdown, while splicing changes were most consistent with a loss of HNRNPC activity, they also implied a loss of PCBP1 and DAZAP1 regulation (Fig. 6d).

This loss was likely driven by decreased levels of PCBP1 and DAZAP1, both of whose mR-NAs were significantly downregulated in the *HNRNPC* knockdown. More generally, *PCBP1* was significantly downregulated ($FDR < 0.2$) in 42 knockdowns and upregulated in 5. This variation in *PCBP1* levels drives indirect effects across the ENCODE knockdowns: the knockdowns where *PCBP1* was secondarily downregulated had splicing changes consistent with loss of PCBP1 regulation, and *vice versa* when upregulated (Fig. 6f, Supplementary fig. 3a), with a significant Pearson's correlation of 0.3 (94%-CI: 0.2–0.4). The presence of measurement errors makes Pearson's r an underestimate of the underlying association (Supplementary fig. 3a,b), so we employed an error-aware hierarchical model that properly disentangles the true association from the estimation errors. (Supplementary fig. 3b, c), uncovering a substantial correlation between expression and splicing variation of 0.55 (94%-CI: 0.41–0.69).

Discussion

Since the development of RNA sequencing, perturbation experiments have been the predominant tool for probing the rules by which regulatory factors control RNA expression and processing. But these studies have typically reported visual summaries of regulation, which are not directly useful for quantitatively interpreting regulation in new biological settings. Further, most analyses do not distinguish between the direct and indirect effects of perturbation, instead assuming that, on average, the signal from direct targets would rise above the background of indirect effects. Our approach, KATMAP, instead extracts actionable insights about splicing regulation from knockdown data by explicitly modelling how changes in SF binding upon knockdown lead to changes in splicing. This approach is supported by the reproducibility of our predictions when applied across cellular contexts and related species. We use this to not only distinguish the direct from indirect effects of SF knockdown but to further interpret these indirect effects in terms of secondary factors. These choices make KATMAP a general framework for rigorously applying insights from knockdown and other perturbation data to new types of analyses.

Central to this work was crafting a description of splicing regulation general enough to apply to most SFs. Some of the most detailed previous models were tailored to specific SFs (Nova and Rbfox) using expert knowledge of the factors' sequence specificity and activity (Weyn-Vanhentenryck et al. 2014; C. Zhang et al. 2010). For example, predefining binding sites based on similarity to a canonical motif and assuming that regulatory activity decays with distance from the splice site at a fixed rate. While reasonable for Nova and Rbfox family proteins, these assumptions will not apply to all SFs. Studying SFs in general requires an automatic approach to determining which assumptions about binding and activity are appropriate to each SF. We accomplished this with a hierarchical model spanning a continuum of assumptions, and employed an inference strategy that learns which are plausible from the knockdown data. This allowed us to make the relaxed assumption that activity varies smoothly with distance from the splice site, with the exact shape and smoothness learned from the data, contrasting with previous approaches that either assumed a predefined activity-distance relation (C. Zhang et al. 2010), fit independent models to different windows of preset size (e.g. 50 nt) (Bak et al. 2024), or derived smoothness simply from the

Figure 6: **KATMAP's regulatory models disentangle indirect effects.** **a)** The fractions targets and nontargets among differentially spliced exons (rMATS FDR < 0.05). Ambiguous target predictions ($Pr(Target)$ and $Pr(Nontarget) < 0.8$) are not included. **b)** Schematic showing how we infer underlying SF perturbations by comparing splicing changes to KATMAP's regulatory models. In this example, exons predicted to be suppressed by HNRNPC are upregulated and exons enhanced by PCBP1 are downregulated, suggesting that decreases in HNRNPC and PCBP1 regulation caused the splicing changes. There is no association between the splicing changes and QKI's predicted targets, suggest no perturbations to QKI's regulation. **c)** Predicted changes in regulation for the knocked down factors, inferred by comparing to all activity models learned from the other cell type. Knockdowns where the expected factor gave the strongest signal are indicated in blue. **d)** The predicted regulation changes in the HNRNPC (K562) knockdown. **e)** Predicted regulation changes as in **d** for knockdowns of all RBPs with a significant activity map. Significant changes in regulation are denoted with thick boxes. **f)** The predicted changes in PCBP1 splicing regulation in datasets where PCBP1's RNA expression was significantly downregulated, not significantly changed, or upregulated.

Supplementary Figure 3: **Deconvoluting true associations from measurement error. a)** Comparison across ENCODE RNAseq datasets between inferred perturbations to PCBP1's splicing regulation and PCBP1's expression variation. The PCBP1 knockdowns are in red. The error bars show the uncertainty in the point estimates (94%-ci) for select datasets. **b)** A true association (left) is obscured by measurement error (middle). Pearson's correlation computed from the noisy point estimates substantially underestimates the true association. If the degree of uncertainty associated with each point estimate (blue lines) is known, it can be incorporated into a hierarchical model to recover the true association (right). **c)** Evaluation of the hierarchical applied to simulated data with true correlations between -0.95 and 0.95 and the same measurement errors as the PCBP1 analysis. The hierarchical model recovers the true association, with a nearly one-to-one relationship between the estimates and true values. Pearson's correlation applied directly to the point-estimates, however, underestimates the association by $\sim 50\%$.

low resolution of CLIP read coverage, as in traditional RNA maps (Yee et al. 2019). While modern neural networks could similarly find implicit mappings from sequence to binding and position to activity (Gupta et al. 2024), our model has the advantage of doing so through explicit, testable statements about SF activity rather than opaque transformations.

The hierarchical architecture of KATMAP spares users from the need to tune parameters each time they begin an analysis. A particular tuning problem common to most approaches to predict regulation in terms of binding motifs is the need to define some identity threshold relating a PWM to binding sites. We solve this by avoiding the assumption of linearity common to many models used to study transcriptional regulation (Balwierz et al. 2014) and their recent extensions to splicing (Bak et al. 2024; Hugh-White 2021). Linear models are easy to fit, but cannot learn nonlinear relationships (like that between occupancy and affinity), requiring predefinition of locations of motifs in the transcriptome likely to be bound. Our additive model can instead accommodate a biophysical (sigmoid) description of occupancy as a function of affinity, allowing us to learn from the data how an SF's PWM or affinity model relates to binding changes in each knockdown dataset.

In distilling knockdown experiments into generalizable models, KATMAP provides the opportunity to leverage decades of experimental work to answer new questions. Our ability to identify secondary SFs responsible for the indirect effects of knockdown suggests that inferring SF perturbations in other settings will be feasible. One such application is determining which SFs are responsible for splicing differences observed in clinical data. Another use of KATMAP's inferences is as null models to explore violations of our simplifying assumptions, such as the cooperativity we uncovered between RBFOX and QKI proteins. Finally, our framework is sufficiently general that it should be applicable to other aspects of gene expression. The key requirement is that regulatory activity depends on the position of binding sites relative to some defined reference point. Therefore, the model is potentially applicable to the study cleavage and polyadenylation (with polyA signal or cleavage site as the reference point), mRNA stability (e.g., with the stop codon as reference point), or even transcriptional regulation.

Online Methods

Selecting the knockdown experiments

We selected ENCODE knockdown RNAseq experiments for proteins annotated as splicing factors (Van Nostrand et al. 2020) and which had a good affinity model derived from RBNS (Lang et al. 2021) using RBPamp (Jens et al. 2022) or a PWM derived from RNAcompete (Ray, Kazan, Chan, et al. 2009) obtained from the CISBP-RNA data database (Ray, Kazan, Cook, et al. 2013). We use the rMATS differential splicing (Shen et al. 2014) results for the hg19 human reference genome in our analyses.

The structure of the KATMAP regression model

The goal of KATMAP is infer a splicing factor's regulatory activity and targets from knockdown data by explaining the resulting splicing changes in terms of changes in SF binding.

Accomplishing this requires several inputs. First, we need the results of a differential splicing analysis, which assigns labels Y to exons indicating whether their inclusion increased, decreased, or was unaffected by knockdown. These are the observations KATMAP seeks to predict. The second input is a model of the SF's sequence specificity. This assigns exons in the differential splicing analysis a set of scores, S, which evaluate all potential binding sites within predefined windows around their splice sites. These scores, S, are what KATMAP will use to explain the splicing changes, Y, in terms of SF binding, and ultimately infer regulatory activity. Finally, additional information about each exon can be included using a set of optional linear predictors, X. In our analyses, we use these linear features to control for expression and pre-knockdown inclusion levels.

Defining the splicing changes, Y

Any differential splicing analysis that labels each exon as unchanged, up-, or downregulated may be used to define the observations, Y. We use one-hot encoding to define Y_i as $(1, 0, 0)$, $(0, 1, 0)$, or $(0, 0, 1)$ if exon i is labelled downregulated, unchanged, and upregulated, respectively.

In our analyses, we defined splicing changes by further processing the rMATS outputs with ashr to better account for the uncertainty in rMATS's estimates (Stephens 2017). This had two motivations. First, ashr accomplishes something like a multiple test correction on parameter estimates, but accounting for the possibility that more exons are upregulated than downregulated. Second, while KATMAP does not consider the magnitude of splicing changes, some of our downstream analyses do. The shrinkage estimates provided by ashr control for measurement error due to low read counts and yield a cleaner view of splicing differences than the raw rMATS $\Delta\psi$ summaries.

Briefly, we used the read counts and rMATS p-values to reconstruct the effect size in log-odds scale, which we denote $\Delta\phi$, and associated uncertainty (Extended Methods). We then pass these to ashr to obtain shrinkage estimates for $\Delta\phi$ and associated standard deviations. We construct 94% credible intervals based on these, calling exons differentially spliced if the 94%-CI does not overlap zero and nonsignificant otherwise.

Defining the training set

We use the rMATS skipped exon table in our analyses. If a set of skipped exons partially overlapped or shared flanking exons, we retained only the most significant to avoid double counting. We include all remaining significant exons in our training sets. For computational efficiency, we downsample the nonsignificant exons, retaining 2,000 randomly chosen exons.

Scoring motifs with a binding model

To explain splicing in terms of binding, KATMAP requires some model of the perturbed SF's binding motif. The binding model may be a traditional position-weight matrix (PWM) obtained from RNAcompete, one or more position-specific affinity matrices (PSAM) (e.g. from RBPamp), or more general affinity models. The key requirement is that the binding model approximately orders potential binding sites from weakest to strongest. In our analyses we used PSAMs derived from RBNS (Lambert et al. 2014) using RBPamp (Jens et al. 2022)

when available, and PWMs derived from RNAcompete (Ray, Kazan, Chan, et al. 2009) obtained from the CISBP-RNA database (Ray, Kazan, Cook, et al. 2013). We excluded RBNS models where the PSAMs showed strong complementarity to the sequencing adapters, which likely reflects a technical artefact rather than true sequence specificity.

To score a k-length motif x with a single 4-by-k PWM or PSAM, W, we sum the scores (in log-scale) for each nucleotide in the motif

$$S(x, W) = \sum_{i=1}^{k} \sum_{nt \in \{A,C,G,U\}} W_{nt,i}[x_i = nt]$$

where the Iverson bracket $[x = nt]$ equals 1 if true and 0 if false.

If we have multiple matrices $W_1, .., W_n$, each with associated weights $a_1, .., a_n$, then we compute the combined score as

$$S(x) = \log \left(\sum_{i}^{n} a_i exp\left[S(x, W_i) \right] \right)$$

In practice, we always define the scores such that the optimal motif is assigned a score of 0 and all other motifs receive negative scores. This is accomplished by subtracting out the score of the optimal motif.

Even for affinity models, however, we do not assume that these initial sequence-based scores are fully accurate representations of affinity. Rather, we treat them as related to log-affinity, but potentially off by some scaling factor. To account for this, we include an *affinity-scale* parameter, λ, which rescales the binding scores.

$$a(x) = \lambda S(x)$$

These rescaled scores are used as log-affinities from which occupancy and changes-in-occupancy are computed. If the scores are truly linearly related to log-affinity, this can be thought of as calibrating the scores against the observed splicing changes. If scores are related to affinity in a monotonic but not linear manner, this rescaling can be thought of widening or narrowing the interval of scores assigned appreciable changes in occupancy.

Scoring the sequences around exons

KATMAP scores sequence in windows around the splice sites of each exon. In our analyses, we scored 200 potential binding sites on the intronic and 60 on the exonic side of each cassette exon's splice sites. If desired, larger or smaller windows can be used or the regions adjacent to the splice sites of the flanking exons included.

Our model includes a parameter for each of the potential binding sites analysed per exon. For computational tractability, rather than model binding at nucleotide resolution at each exon, we aggregate neighboring binding sites into non-overlapping regions of ten potential binding sites each. We index these regions, $1..J$, where in most of our analyses $J = \frac{1}{10}(200 + 60) \times 2 = 52$. More generally, in an analysis with a different region size, different intronic or exonic window size, or number of the splice sites (e.g. including the

flanking exons)

$$J = \frac{1}{\text{RegionSize}}(\text{IntronicWindowSize} + \text{ExonicWindowSize}) \times \text{NumberOfSplicesites}$$

We then assign each subregion j in exon i a log-affinity. We denote the motif scores of the ten binding sites ending in subregion j as $\mathbf{S}_{ij}$ and compute the log-affinity of the region by first scaling the log-affinity of each binding site by λ and taking the log of the sum of their exponents:

$$\mathbf{a}_{ij} = \text{LogSumExp}(\lambda\mathbf{S}_{ij})$$

Defining the binding function

For a given free protein concentration F, the occupancy of a site with given log-affinity, a, is described by the Langmuir equation

$$\Theta(a; F) = \frac{1}{1 + \frac{1}{Fe^a}}$$

The change in occupancy, then, is determined by the free protein concentration before and after knockdown/overexpression (Supplementary fig. 4a). If the free protein concentration changes by some factor k, the change in occupancy can be computed (Supplementary fig. 4b)

$$\delta(a; F, k) = \Theta(a; kF) - \Theta(a; F)$$

However, these parameters, F and k, while biophysically meaningful, are not immediately interpretable in terms of which affinities experience large changes in binding. We therefore reparameterize the change-in-binding function in terms of its shape. First, we include a most-relevant-affinity parameter, m, which indicates which affinity experienced the greatest change in binding. Second, we describe the magnitude of the change in binding at sites with affinity m with a *greatest-change-in-binding* parameter, d (Supplementary fig. 4b,c). That is we reparameterize such that that maximum change in occupancy, d, occurs as at log-affinity m

The corresponding biophysical parameters can be computed by solving

$$\delta'(m; F, k) = 0 \qquad \text{fix the optimum at } m \text{ by setting the derivative to zero}$$
$$\delta(m; F, k) = d \qquad \text{assert that the change in occupancy at } m \text{ is } d$$

which yields

$$k(d) = \frac{(d+1)^2}{(d-1)^2}$$

$$F(m, d) = \left(e^m\sqrt{k}\right)^{-1}$$

so

$$\tilde{\delta}(a; m, d) = \delta\left(a; F(m, d), k(d)\right)$$

where $\tilde{\delta}$ denotes a reparameterization of δ.

Our model's predictions are constructed by multiplying these binding changes by other parameters we term *activity coefficients*, α. Because d changes the height of the binding function but not the relative shape (unless d is near 1 or -1, $\frac{B(s; m, d_1)}{B(s; m, d_2)} \approx \frac{d_1}{d_2}$; Supplementary fig. 4c), we cannot usually identify both the magnitude of binding changes and the magnitude of regulatory activity: most effects on the model's predictions of changing d from value d_1 to d_2 can be entirely reversed by scaling the magnitudes of the activity coefficients by $\frac{d_1}{d_2}$. To resolve this, we normalize the changes in binding by $|d|$ to compute the relative change-in-binding (Supplementary fig. 4d)

$$\tilde{\delta}_{rel}(a; m, d) = \frac{1}{|d|}\tilde{\delta}(a; m, d)$$

Further, because for all values of $|d| < 0.95$ yield essentially the same relative changes-in-binding, we fix $d = -\frac{1}{2}$ for knockdown experiments and $d = \frac{1}{2}$ for overexpression. If $|d| > 0.95$, as might be the case for knockout data, the relative change-in-binding function flattens out, but this can be approximated with small values of the affinity scale parameter, $\lambda \ll 1$ (Supplementary fig. 4e)

To recap, we score sequence around each exon's splice site using a model of the RBP's binding site. We then divide the sequence around each exon into non-overlapping 10nt regions, each containing ten potential binding sites. We denote the set of all scores as S, where S_{ij} refers to the scores of the ten potential binding sites in region j of exon i. We then scale these scores with an *affinity scale* parameter, λ, and treat the resulting values as log-affinities. We then compute the log-affinity of the regions based on the affinities of the ten binding sites contained in each. Finally, we assign each region a relative changing-in-binding, determined by the *most-relevant-affinity* parameter, m. We describe this entire procedure of transforming the motif score to relative changes-in-binding with the binding function

$$B(\mathbf{S}_{ij}; \lambda, m, d) = \tilde{\delta}_{rel}\left(\text{LogSumExp}(\lambda\mathbf{S}_{ij}); m, d\right)$$

fixing $d = -\frac{1}{2}$ for knockdown experiments.

Predicting changes in regulation from changes in binding

We describe the position-specific effect of RBP binding with a set of *activity coefficients*: $\alpha_{1..J}$. A positive value of α_j indicates that binding at region j enhances exon inclusion. A negative value instead describes suppressing activity. Critically, we assume that activity varies smoothly with position, described later when discussing our model's priors.

To compute the predicted effect of loss of binding near exon i on its splicing outcome, we multiply the change in binding at each region j, $B(S_{ij}; \lambda, m, d)$, by the corresponding activity coefficient α_j. We make the simplifying assumption that activity is additive across

Supplementary Figure 4: Binding functions. **a)** The affinity-occupancy relationship before (black) and after knockdown (blue) for different pre-knockdown free protein concentrations (F) and knockdown efficiencies $(1-k)$. **b)** The change in binding functions that result from occupancy curves in **a)**, with the corresponding values for the shape parameters m and d. Note that unlike the biophysical parameters (F and k) these shape parameters explicitly define the affinity and magnitude of the greatest binding change (open circle). **c-d)** Absolute change in binding, $\tilde{\delta}$, (**c**) and relative change in binding, $\tilde{\delta}_{rel}(\cdot)$, (**b**) with the most-relevant affinity $m = 0$ for different values of d. **e)** The best approximations of the binding function at a given d obtainable by rescaling the affinities

all binding sites, so the total change in activity due to the changes in binding across regions $1..J$ is

$$\Delta \mathbf{A}_i = \sum_{j=1}^{J} B(\mathbf{S}_{ij}; \lambda, m, d) \alpha_j$$

Connecting changes in regulation to observed splicing changes

To predict the observed splicing changes, the model must assign each exon i three probabilities, the probability of being downregulated, unchanged, or upregulated: $\hat{p}_i = \left(p_i^{(-)}, p_i^{(0)} p_i^{(+)} \right)$. We describe the log-odds ($q = log\frac{p}{1-p}$) of these probabilities with an additive model:

$$q_i^{(+)} = \Delta \mathbf{A}_i + ... = \sum_{j=1}^{J} B(S_{ij}; \lambda, m, d) \alpha_j + ...$$

$$q_i^{(-)} = -\Delta \mathbf{A}_i + ... = -\sum_{j=1}^{J} B(S_{ij}; \lambda, m, d) \alpha_j + ...$$

That is, if the activity at exon i increases, the predicted log-odds of inclusion being upregulated increases and the log-odds of being downregulated decreases. In an overexpression experiment, the changes-in-binding $B(S_{ij}; \lambda, m, d)$ are positive, and upregulation is driven by gains in enhancing activity and downregulation by gains in suppressing activity. In knockdown experiments, the changes-in-binding are negative, flipping the signs of the change in activity: upregulation is driven by loss of suppressing activity and downregulation by loss of enhancing activity.

Incorporating additional information into the predictions as linear features

Our models aims to explain the results of the differential splicing analysis in terms of SF binding, but other information unrelated to binding is likely to be predictive of which exons were called differentially spliced, for biological and/or technical reasons. We incorporate these features by adding a standard linear model into the predictions. The K linear features included in the model are organized as an I-by-K table, X. All features are mean-centered and standardized. Any quadratic terms are created using the standardized features to reduce the dependence structure of the target posterior distribution.

Because not every linear feature will be relevant to each of the three predictions–downregulated, unchanged, upregulated–we divide the linear features X into three (potentially overlapping) subsets, denoted $X^{(-)}$, $X^{(0)}$, and $X^{(+)}$. For example, the read count of an exon influences the statistical power to detect changes in splicing and so should inform the predicted probability of being unchanged, $p^{(0)}$ and so is assigned to $X^{(0)}$. The original inclusion level of an exon determines whether up- or downregulation is detectable, because there is little room for inclusion to increase or decrease when the exon was already nearly 100% or 0% included. So, the inclusion-level prior to SF perturbation is likely directly informative of $p_i^{(-)}$ and $p_i^{(+)}$,

and so is assigned to both $X^{(-)}$ and $X^{(+)}$. These feature is incorporated into the predictions as

$$q_i^{(-)} = -\sum_{j=1}^{J} B\left(S_{ij}; \lambda, m, d\right) \alpha_j + \sum_{k=1}^{K^{(-)}} \beta_k^{(-)} X_{ik}^{(-)} + \beta_0^{(-)}$$

$$q_i^{(0)} = \sum_{k=1}^{K^{(0)}} \beta_k^{(0)} X_{ik}^{(0)}$$

$$q_i^{(+)} = \sum_{j=1}^{J} B\left(S_{ij}; \lambda, m, d\right) \alpha_j + \sum_{k=1}^{K^{(+)}} \beta_k^{(+)} X_{ik}^{(+)} + \beta_0^{(+)}$$

where $\beta_{1..K^{(-)}}^{(-)}$, $\beta_{1..K^{(0)}}^{(0)}$, and $\beta_{1..K^{(+)}}^{(+)}$ denote the linear coefficients describing the effect the linear features on the predictions, and where $\beta^{(-)}$ and $\beta^{(+)}$ are the intercepts.

Connecting predictions to observed splicing changes

The probabilities for category z are computed

$$\hat{p}_i^{(z)} = \frac{exp\left(q_i^{(z)}\right)}{exp\left(q_i^{(-)}\right) + exp\left(q_i^{(0)}\right) + exp\left(q_i^{(+)}\right)}$$

We need to connect these to the observed splicing changes with a categorical distribution

$$Y_i \sim Categorical\left(p = \left[\hat{p}_i^{(-)}, \hat{p}_i^{(0)}, \hat{p}_i^{(+)}\right]\right)$$

The probability of an observation under this distribution is simply equal to the predicted probability for that category. For example if exon i is downregulated, the likelihood $P(Y_i|\hat{p}_i) = \hat{p}_i^{(-)}$.

Defining the model's priors

We represent assumptions about our model's parameters with Bayesian priors. Some of the priors we employ reflect what we consider to be plausible based on our understanding of molecular biology. Others reflect statistical considerations that aim to avoid unrealistically strong predictions or complications to statistical inference. For example, given the limitations of experimental data and modelling thereof we do not believe KATMAP should ever make extremely confident predictions on the order of million-to-one odds. In other cases, we wish to avoid identifiability issues, where large sets of distinct parameter values would yield essentially the same predictions. As we describe the model's prior, we will note for each parameter which consideration is at play.

Supplementary Figure 5: Priors on the hyperparameters and resulting spatial correlations. Prior densities on **a)** the affinity scale (in log_2-scale); **b)** the average amount of binding lost per exon (implicit prior on most-relevant-affinity); **c)** the length-scale of the Gaussian process prior (in units of 10 nt regions) on the activity coefficients; and **d)** stationary standard deviation of the Gaussian process. **e)** Examples of the spatial smoothness resulting from different length-scales.

Priors on the binding parameters: λ, m

The parameter λ scales the motif scores S to assign binding sites log-affinities $S \times \lambda$. When $\lambda < 1$ the scores are compressed, assigning a narrower range of affinities. When $\lambda > 1$ the scores are stretched, assigning a wider range of affinities. We choose a prior that asserts both to be equally plausible. Beyond this, the prior we place on λ largely reflects a practical consideration about operations in log-scale. Multiplication on log-scale corresponds to exponentiating the actual affinities:

$$affinity = e^{\lambda S} = e^{S\lambda}$$

Even modest values of λ will yield extremely large or small affinities. So we constrain λ to a relatively narrow range of values by using a unit-normal prior

$$\log_2 \lambda \sim Normal(0, 1)$$

which places about 99% of the prior probability on λ between $\frac{1}{6}$ and 6 (Supplementary fig. 5a).

The most-relevant-affinity parameter m describes the which affinity experiences the greatest change in binding. Assuming the binding site model is accurate, motifs with moderate to high affinity should experience changes in binding; motifs with very low affinity are too weak to be appreciably bound in the first place. So a biologically reasonable prior would favor values of m closer to the affinity of the optimal motif and disfavor very low affinities. But beyond this, it is difficult to precisely express how plausible a given value of m is.

However, m determines how much SF binding our model predicts was lost from near each exon. It is much easier assess the biological plausibility of these statements. Certain values of m might imply that the average exon has > 20 copies of the SF bound within a few hundred nucleotides. For SFs that bind specific sequence motifs, this is unrealistically high. So rather than explicitly define a prior on m, we instead define our prior on the average change in binding near each exon

$$\bar{B} = \left| \frac{1}{N} \sum_{i=1}^{N} \sum_{j=1}^{J} B(S_{ij}; m, \lambda, d) \right|$$

We divide this quantity by the number of potential binding sites per exon ($J = 52$ in our analyses) and place the prior on the log-odds of this fraction (adding small constants, $\epsilon = 10^{-7}$, to the numerator and denominator to avoid numerical issues due to computational precision):

$$\tilde{B} = \frac{\bar{B} + \epsilon}{J + 2\epsilon}$$

We then define our prior as

$$\log \frac{\tilde{B}}{1 - \tilde{B}} \sim Normal(\mu = -3.9, \sigma = 0.84)$$

which places 99% of the prior probability on the average change in binding per exon between 0.12 and 7.8 (Supplementary fig. 5b). To exclude extremely unrealistic values of

m, we constrain the model to only consider values between the median log-affinity and the maximum affinity per 10nt region $log(10)$ (or more generally $log(RegionSize)$)

Assuming that activity varies smoothly with distance: $\alpha_{1..J}$

A fundamental assumption of our model is that the effect of SF binding on exon-inclusion depends on where the protein is bound relative to the exon. This is described by activity coefficients $\alpha_{1..J}$. We further assume that the activities of nearby regions, say $j = 2$ and $j = 3$, are similar, but that distant positions may have very different activities. Statistically, we express this with a Gaussian process prior that assumes the activity coefficients are spatially correlated (Rasmussen and Williams 2006). Gaussian process can be conceptualized as a multivariate normal distribution, where the covariance between variables α_j and α_k is a decreasing function of how far apart they are $|j - k|$.

$$\alpha_{1..J} \sim MvNormal\left(0, \Sigma(L, \sigma)\right)$$

We choose the smooth, yet flexible, Matern-3/2 covariance function to express these spatial correlations (Rasmussen and Williams 2006). Because we further want to assume that the effects of exonic versus intronic binding may be very different, we restrict the spatial correlations to pairs of regions within the same biological window. We express this with a set labels $R_{1..J}$ that indicate whether region j is on the intronic side of the 3' splice site, the exonic side, the exonic side of 5'-splice site, or the intronic side. If a pair of regions j and k correspond to differ biological labels, the spatial correlation between α_j and α_k is set to zero

$$\Sigma_{jk}(L, \sigma) = \begin{cases} \sigma^2 \left(1 + \frac{\sqrt{3}|j-k|}{L}\right) exp\left(-\frac{\sqrt{3}|j-k|}{L}\right) & \text{if } R_j = R_k \\ 0 & \text{if } R_j \neq R_k \end{cases}$$

Assuming this spatial correlation structure requires introducing two additional parameters that must be learned. The length-scale, L, determines how far apart a pair of regions j and k must be before they are weakly correlated . The stationary standard deviation, σ, determines the range of magnitudes the activity coefficients are expected to span. To prevent the inference algorithm from wasting computation by exploring large but practically equivalent regions of parameter space, we constructed priors that avoid both large and small values of these hyperparameters (Supplementary fig. 5c,d, see Extended Methods). This places 99% of the prior probability for the length-scale on the interval 3.2 nt $< L <$ 226 nt (Supplementary fig. 5c), which ranges from near independence to correlations that span the whole intronic window (Supplementary fig. 5e). For the stationary standard deviation, 99% of the prior probability falls in the interval $0.14 < \sigma < 8.7$ (Supplementary fig. 5d).

Assumptions about the linear coefficients: $\beta_{0..K}$

Because we mean center and standardize the linear features, β_k indicates how much an increase in linear feature k by one standard deviation changes the log-odds of the prediction. Because the impact is in log-odds scale, coefficients of magnitude $|\beta| > 10$ would

have extreme impacts on the prediction: increasing the linear feature by one standard deviation would increase a predicted probability of $p = .01$ to $p > .99$. As the standardized features typically span a range of more than four standard deviations, we employed a weakly-informative prior for the linear coefficients and intercept, using a normal density with mean 0 and standard deviation 5

$$\beta_{0..K} \sim Normal(\mu = 0, \sigma = 5)$$

The model is not sensitive to this choice of prior, as it spans the range of reasonable effect sizes, both strong and weak.

High-level overview of model inference

The goal of Bayesian inference is to determine which parameters are plausible given the observed data. In practice, this is often a harder task than formulating the statistical model itself. Evaluating the exact plausibility requires applying Baye's theorem, which expresses a reasonable proposition: to determine how plausible a given hypothesis h is, you must compare it to every other hypothesis admitted by the model, $h \in \Theta$:

$$P(h|Y) = \frac{P(Y|h)P(h)}{\int_{\tilde{h} \in \Theta} P(Y|\tilde{h})P(\tilde{h})d\tilde{h}}$$

This is typically impossible to compute exactly, because it requires integrating over the entire parameter space. In the case of KATMAP's regression model, this parameter space has > 50 dimensions.

A common solution is Monte Carlo sampling. Rather than evaluate the exact posterior density $P(\theta, \phi|Y)$, we instead seek to obtain a sample of hundreds of parameter values which follow the posterior distribution

$$(\theta_1, \phi_1), .., (\theta_N, \phi_N) \sim P(\theta, \phi|Y)$$

This posterior sample provides a representative set of parameters that are plausible given the observed data from which we can estimate the posterior mean. We can also construct credible intervals describing the range of plausible parameters values. Because the model predictions for each exon are computed based on these parameters, it is further possible to obtain interval estimates for all of the exon-level predictions.

Bayesian inference with the KATMAP model

When writing the posterior for KATMAP's model it is useful to divide the model parameters into two groups. First, we have the regression coefficients $\theta = (\alpha_{1..J}, \beta_{0..K})$. Second, with slight abuse of terminology, we have the hyperparameters $\varphi = (\lambda, m, L, \sigma)$. Partitioning the parameters in this manner makes clear the hierarchical nature of the model

$$P(\theta, \varphi|Y) \propto P(Y|\theta, \varphi)P(\theta|\varphi)P(\varphi)$$

Integrated nested Laplace approximations (INLA) provide a means to leverage this structure and divide the problem of posterior inference into two manageable, nested sub-problems.

First, an inner step focuses on identifying which regression coefficients are plausible given the data and hyperparameters. This approximates the conditional posterior $P(\theta|Y,\varphi)$ for a specific hyperparameter value, φ. Second, an outer step focuses on determining which hyperparameters are plausible given the data. This approximates the marginal posterior for the hyperparameters $P(\varphi|Y)$

Approximating the conditional posterior over the regression coefficients

The inner step takes advantage of how the model simplifies if the hyperparameters are held fixed to a particular value. The known binding parameters (λ, m) would transform the table of motif scores S into a fixed table of changes-in-binding $B_{(\lambda,m)}$. The known spatial correlation parameters would yield a fixed prior on the activity coefficients. The problem reduces to a generalized linear model, albeit a Bayesian formulation with priors.

We use second-order optimization identify the optimal regression coefficients as the mode of the conditional posterior, θ_φ^*, determining the step size with a backtracking line search (Nocedal and Wright 2006). This requires evaluating the gradient ∇ and Hessian matrix H, which we compute using automatic differentiation with the jax library (Bradbury et al. 2018). How sharply the log-posterior is curved around the mode is described by the Hessian matrix H_{θ^*} and can be used to describe the uncertainty about the coefficients with a covariance matrix, $\Sigma_\varphi^* = H_{\theta_\varphi^*}^{-1}$. This provides a multivariate normal approximation to the conditional posterior

$$P(\theta|Y,\varphi) \approx \mathcal{N}(\theta; \mu = \theta_\varphi^*, \Sigma = \Sigma_\varphi^*)$$

Approximating the marginal posterior over the hyperparameters

The outer step evaluates the plausibility of specific hyperparameters φ by approximating the marginal posterior, $P(\varphi|Y)$. The hyperparameters, φ only relate to the observations through the regression coefficients θ. So evaluating the plausibility of some specific φ requires considering all possible values the regression coefficients θ could take. Rue, Martino, and Chopin (2009) accomplish this using the normal approximation to the conditional posterior obtained in the inner step evaluated at its mode, θ_φ^*

$$P(\varphi|Y) \underset{\sim}{\propto} \frac{P(Y, \theta_\varphi^*, \varphi)}{\mathcal{N}(\theta_\varphi^*; \theta_\varphi^*, \Sigma_\varphi^*)} = P(Y, \theta_\varphi^*, \varphi) \det\left((2\pi)^d \Sigma_\varphi^*\right)^{\frac{1}{2}}$$

$$= P(Y, \theta_\varphi^*, \varphi) \int_{x \in \mathbb{R}^d} exp\left(\theta_\varphi^* - x\right)^T \Sigma_\varphi^{*-1}\left(\theta_\varphi^* - x\right) dx$$

This approximation to the marginal density multiplies the joint density $P(Y, \theta_\varphi^*, \varphi)$ by the volume of regression coefficients that are plausible given φ.

Monte Carlo sampling from the joint posterior with INLA

We couple adaptive importance sampling with these integrated nested Laplace approximations (INLA) to draw samples of plausible parameters from the joint posterior. First, we use adaptive importance sampling and the INLA approximation to draw a representative

sample of plausible hyperparameters from their marginal posterior, $\varphi_1, .., \varphi_N \sim \varphi|Y$. This explores the hyperparameters in a way that iteratively hones in on the true posterior, by fitting mixture model approximations with importance-weighted expectation-maximization (Cappé et al. 2008) (Supplementary fig. 6 a, Extended Methods). Second, for each sampled hyperparameter we solve the resulting conditional model by optimization, approximating the conditional posteriors (including their correlation structure) over the regression coefficients with a multivariate Gaussian. We then sample regression coefficients from these conditional posteriors, $\theta_1, .., \theta_N \sim \theta|Y, \varphi_{1..N}$. Together, these constitute an approximate sample from the joint posterior, $(\theta_1, \varphi_1), .., (\theta_N, \varphi_N) \sim \theta, \varphi|Y$. We run this algorithm twice for each knockdown experiment, generating two separate posterior samples. If both runs converge to the same distribution, we consider inference successful and merge the distributions.

Evaluating convergence

We use two criteria to assess whether our Monte Carlo samples are reliable representations of the target posterior. First, we use the split $\hat{r}$-statistic (Vehtari, Gelman, et al. 2021) to assess whether our two inference runs converged to the same distribution. Values of $\hat{r}$ close to one indicate highly overlapping distributions and if marginal distributions of all parameters yield $\hat{r} < 1.05$, we consider the Monte Carlo sampling runs to have converged. Second, during adaptive importance sampling of the hyperparameters, we stabilize the importance weights with Pareto-smoothing. This automatically provides a diagnostic statistic, $\hat{k}$, indicating whether the resulting importance sample is reliable. We use the cutoff suggested by Vehtari, Simpson, et al. (2024), requiring $\hat{k} < 1 - \frac{1}{log_{10} N_{proposals}}$ or $\hat{k} < 0.7$–whichever is more stringent– for the posterior samples to be considered trustworthy.

Evaluating evidence of splice regulatory activity

To quantify evidence of in favor of splice regulatory activity, we compare to a reduced model with the splicing impacts removed. We compare the full and reduced models using Pareto-smoothed importance sampling leave-one-out cross-validation (PSIS-LOO) (Vehtari, Simpson, et al. 2024), a fully Bayesian approach to model comparison which uses the posterior samples from the full and reduced models to estimate whether the full model better predicts held out data. We bootstrap observations to compute a Z-score describing our confidence that including splicing activity in the model improves performance.

Determining which activity maps are significant

We consider an activity map significant based on two criteria. First, we require that at least two activity coefficients be significantly nonzero (98%-CI does not overlap zero). Second, we require a model comparison z-score > 2.

We additionally considered whether the splicing changes are well-described by the expected motif. We excluded any activity models that choose binding parameters which assigned an unreasonable number of binding changes (> 20 per exon). This indicates that the knockdown-affected exons are distinguished by some characteristic sequence composition, but it is not well described by the perturbed factor's expected motif. This excluded SRSF9

Supplementary Figure 6: **Posterior inference for the KATMAP model.** **a**) Exploration of the hyperparameters through adaptive importance sampling, showing the evaluation points (left) and the adaptive proposal distribution at successive iterations. New hyperparameters (blue) are drawn from the proposal estimated in the previous iteration, which are in turn used to refine the adaptive proposal distribution to better match the target posterior. The proposal is a student T mixture model fit by importance-weighted Expectation-Maximization. Upon termination, hyperparameters are accepted by importance sample to obtain a sample from the posterior (blue).

from our analysis, which yielded a significant map by assigning appreciable occupancy to most sites in the transcriptome.

Evaluating a reduced model without splicing activity

To evaluate evidence of splice regulatory activity, we compare to a reduced version of the model that only includes the linear predictors. The predictions under this model are computed

$$q_i^{\emptyset(-)} = \sum_{k=1}^{K^{(-)}} \beta_k^{(-)} X_{ik}^{(-)} + \beta_0^{(-)}$$

$$q_i^{\emptyset(0)} = \sum_{k=1}^{K^{(0)}} \beta_k^{(0)} X_{ik}^{(0)}$$

$$q_i^{\emptyset(+)} = \sum_{k=1}^{K^{(+)}} \beta_k^{(+)} X_{ik}^{(+)} + \beta_0^{(+)}$$

This is a standard multinomial logistic regression and has no hyperparameters that would necessitate the use of INLA. We approximate the posterior distribution over the regression coefficients using the same optimization and Laplace approximations used to evaluate the conditional posterior for the full model. We then draw a posterior sample from the reduced model, $\theta_{1..N}^{(\emptyset)}$, of the same size as the posterior sample obtained for the full model.

Defining direct targets

We consider an exon a direct target if its splicing impact is sufficiently large to improve upon a reduced model without splicing activity. To assess this, we compare the posterior distribution of predictions between the full and the reduced models, p and $p^{\emptyset}$. In a knockdown experiment, enhancing targets should be downregulated and suppression targets upregulated. So we consider an exon an enhancing target if we are confident the splicing impact, A_i, is negative enough to improve the downregulation prediction

$$P(A_i < 0|Y) > 0.9 \quad \text{and} \quad P\left(p_i^{(-)} > p_i^{\emptyset(-)}\Big|Y\right) > 0.9$$

and a suppression target if the splicing impact is positive enough to improve the upregulation prediction

$$P(A_i > 0|Y) > 0.9 \quad \text{and} \quad P\left(p_i^{(+)} > p_i^{\emptyset(+)}\Big|Y\right) > 0.9$$

This assigns each exon into one of three mutually exclusive categories: enhancing target, suppression target, and nontarget. In some analyses, we wish to evaluate the confidence of these classifications and we do so with a multinomial regression that asks what fraction of exons with a given splicing impact were assigned to each category. This assigns each exon a probability of being an enhancing target, a nontarget, or a suppression target based on its splicing impact. We assumed that fractions of direct targets differ among upregulated, downregulated, and nonsignificant exons by fitting separate intercepts for each category.

Evaluating eCLIP enrichment at targets

For SFs with significant activity maps and available eCLIP data, we downloaded the bed files representing peak calls for both replicates. We did not use the results of the irreproducible discovery rate analysis, because we believe this to represent an overly stringent criteria for combining replicate information. We considered the presence of a peak > 2-fold enriched over input in at least one of the eCLIP replicates as evidence of potential binding. The signal of differential splicing and eCLIP binding are clearest in highly expressed transcripts due to better representation in the sequencing reads; evaluating enrichment between differentially spliced and nonsignificant exons risks spurious associations due to correlated statistical power. To avoid this, we compared binding at targets to nontargets that significantly changed in the same direction. We further focused on the positions predicted by KATMAP to impact splicing. For both targets and nontargets, we counted the number of exons with evidence of binding, k_{target} and $k_{nontarget}$, and used a beta-binomial conjugate model with a uniform prior to obtain 2,000 posterior samples of the fractions of bound target and nontarget exons, f_{target} and $f_{nontarget}$

$$f_{target}|k_{target} \sim Beta(\alpha = k_{target} + 1, \beta = N_{target} - k_{target} + 1)$$

$$f_{nontarget}|k_{nontarget} \sim Beta(\alpha = k_{nontarget} + 1, \beta = N_{nontarget} - k_{nontarget} + 1)$$

where N_{target} and $N_{nontarget}$ are the total number of targets and nontargets that significantly changed in a given direction. We constructed a posterior sample of enrichment estimates as

$$log_2 \frac{f_{target}}{f_{nontarget}}$$

and computed the means and credible intervals from this.

Defining effect sizes for changes in splicing

Differential splicing analyses typically report inclusion changes as $\Delta\psi = \psi_{kd} - \psi_{ctrl}$. However, under the hood, the statistical analysis is typically performed with respect to the log odds transformation of inclusion, $\phi = \log_2 \frac{\psi}{1-\psi}$, with the effect size being the log-odds ratio of inclusion before and after knockdown.

$$\Delta\phi = \phi_{kd} - \phi_{ctrl} = \log_2 \frac{\psi_{kd}/\left(1 - \psi_{kd}\right)}{\psi_{ctrl}/\left(1 - \psi_{ctrl}\right)}$$

While our model does not aim to predict the magnitude of inclusion changes, only the direction, our analyses make use of inclusion-levels in two ways. First, we use the pre-knockdown inclusion level, ϕ_{ctrl} as a linear predictor, being informative of whether an increase or decrease in inclusion is detectable. Second, we assess whether our splicing impacts can predict the magnitudes of inclusion changes, despite not being trained on them. In both cases, we favor the log-odds representation.

The log-odds representations, ϕ and $\Delta\phi$, offer two advantages. First, ψ by definition must be between 0 and 1, and so the magnitude of $\Delta\psi$ only has meaning in the context of the original inclusion-level. $\Delta\psi = +0.1$ is valid for $\psi_{ctrl} = 0.05$ but not for $\psi_{ctrl} = 0.95$,

because $\psi_{kd} = 1.05$ is impossible. A second, related concern, is that inclusion levels near 0 or 1 likely represent more extreme splicing contexts than does intermediate inclusion. A change in inclusion from 0.5 to 0.6 corresponds to $\Delta\psi = 0.1$, as does a change from 0.01 to 0.11. However, an exon included in only 1% of transcripts likely represents a strong silencing context or has weak splice sites, and so the increase of 10% likely reflects a more substantial change in regulation than a change from 50% to 60% . The odds-ratio reflects this, treating an inclusion change of $0.5 \rightarrow 0.6$ as representing a smaller effect ($\Delta\phi \approx +0.6$) than a change of $0.01 \rightarrow .11$ ($\Delta\phi \approx +3.6$).

GO analysis of QKI-RBFOX cooperatively regulated exons

We defined confident QKI-RBFOX cotargets as predicted targets of either QKI or RBFOX that were downregulated upon knockdown and were inferred to also be regulated by the other factor based on splicing impacts. For the latter criteria, we chose a splicing impact cutoff of $<$ -0.25 by examining the eCDFs (Fig. 5c,d). We constructed expression-matched background sets from all genes expressed in the QKI and RBFOX HepG2 knockdowns using importance sampling and used GOrilla to identify enriched GO terms relative to this background (Eden et al. 2009). We also performed this analysis using all differentially splicing QKI and RBFOX targets as the background. We used all three ontologies–process, function, and component. We identified significantly enriched terms using a 10% false discovery rate threshold.

Inferring SF perturbations from splicing changes

To explain a set of splicing changes in terms of changes in SF regulation, we use the splicing impacts previously inferred from the knockdown data as the linear features in a multinomial regression. For each splicing factor, f, with a significant activity map, we use the posterior means of its activity ($\hat{\alpha}_{1..J}^{(f)}$) and binding parameters ($\hat{\lambda}^{(f)}, \hat{m}^{(f)}$) to compute splicing impacts for every considered exon i

$$\mathbf{A}_i^{(f)} = \sum_{j=1}^{J} B\left(\mathbf{S}_{ij}; \hat{\lambda}^{(f)}, \hat{m}^{(f)}, -\frac{1}{2}\right) \hat{\alpha}_j$$

We incorporate these into a regression model, including linear coefficients to account for unperturbed inclusion-levels, expression, and their squares

$$q_i^{(-)} = -\sum_{f \in SF} \mathbf{A}_i^{(f)} + \sum_{k=1}^{K^{(-)}} \beta_k^{(-)} X_{ik}^{(-)} + \beta_0^{(-)}$$

$$q_i^{(0)} = \sum_{k=1}^{K^{(0)}} \beta_k^{(0)} X_{ik}^{(0)}$$

$$q_i^{(+)} = \sum_{f \in SF} \mathbf{A}_i^{(f)} + \sum_{k=1}^{K^{(+)}} \beta_k^{(+)} X_{ik}^{(+)} + \beta_0^{(+)}$$

And describe the observed splicing changes $Y_{1..N}$ as following a categorical distribution

$$\hat{p}_i^{(z)} = \frac{exp\left(q_i^{(z)}\right)}{exp\left(q_i^{(-)}\right) + exp\left(q_i^{(0)}\right) + exp\left(q_i^{(+)}\right)}$$

$$Y_i \sim Categorical\left(p = \left[\hat{p}_i^{(-)}, \hat{p}_i^{(0)}, \hat{p}_i^{(+)}\right]\right)$$

Because all of the splicing impacts $\mathbf{A}$ are precomputed this is a standard multinomial logistic regression. We use second-order optimization to find the posterior mode and, as described previously, use the curvature at the mode to approximate the posterior with a multivariate Gaussian. To assume sparse regression coefficients *post hoc*, we use ashr to apply shrinkage to the estimates.

Simulation study of model inference

For our simulation studies, we wished to assess how reliably our inference algorithm could recover the parameters which truly generated observed splicing changes. For our true parameters, $(\theta, \varphi)^{(true)}$, we used the posterior means inferred from twenty knockdowns that yielded significant activity maps. These parameters coupled with the predictor variables (S and X) naturally assign each exon probabilities of being downregulated, unchanged, or upregulated after knockdown. So that our simulated data provides enough statistical power to make confident inferences, we adjust the intercepts to ensure around 500 up- and downregulated exons. To generate the simulated observations $Y^{(sim)}$, we sample an outcome for each exon from a categorical distribution based on these predicted probabilities. We then use KATMAP to predict these simulated observations using the original predictor variables S and X to obtain a posterior sample $(\theta, \varphi)_1, .., (\theta, \varphi)_N | Y^{(sim)}$. We computed posterior means and credible intervals and compared these to the true values $(\theta, \varphi)^{(true)}$.

Statistical analyses

We represent uncertainty in our analyses with credible intervals. For KATMAP's inferences, these are obtained by Monte Carlo sampling. With estimates based on count or presence/absence data, we employ the beta-binomial conjugate model, with uniform priors. For frequency estimates so obtained, we compute the mean and CI from the posterior beta distribution. For enrichment estimates, we sample frequencies from the posteriors of foreground and background to obtain a posterior sample of enrichments and construct the CI based on this. Otherwise, we typically employed Bayesian bootstrapping, wherein a posterior sample of estimates is obtained by repeatedly computing weighted averages, with the weights sampled from a uniform Dirichlet distribution (D. B. Rubin 1981).

There are several technical details consistent throughout our approach to inference. We use the jax scientific library to compute the gradients and Hessian matrices of our probability densities, which we need to optimize and quantify uncertainty in our model. We always work with log-probabilities for the sake of numerical stability. Despite this, loss of numerical precision occasionally results in non-positive semidefinite covariance matrices, that is matrices with eigenvalues less than zero. In this case, we follow theorem 3.2 in Cheng and Hingham

(1998) to identify the nearest positive semidefinite matrix to Σ with eigenvalues greater than 10^{-5}, which amounts to setting all eigenvalues less than 10^{-5} to this cutoff.

Reporter assay

Constructing mutant sequences

For each selected exon, we identified the 10nt region most strongly contributing to the splicing impact (Fig. 4a,b). We then treated all regions that were at least 1/2 as impactful as potential SREs. For each SRE, we identified the three single-nucleotide polymorphisms most disruptive to binding. For exons with both up and downstream SREs, we constructed separate sequences with the upstream or downstream mutations alone, as well as with both upstream and downstream mutations.

Cell culture

HepG2 cells (ATCC) were grown in Dulbecco's Modified Eagle Medium (GIBCO) supplemented with 10% bovine growth serum (GIBCO) and 1X penicillin-streptomycin (Gen Clone, Cat no. 25-512) in a 37°C incubator under 5% CO_2.

Minigene splicing reporter constructs

RHCglo minigene reporter plasmid DNA (Addgene plasmid # 80169) was digested with SalI-HF (NEB, cat no. R3138S) and XbaI (NEB, cat no. R0145S) restriction enzymes to remove synthetic exon and flanking human β-globin intronic and splice site sequence. Digests were performed in a single reaction according to the manufacturer's instructions, then purified using DNA Clean & Concentrator Kit (Zymo, Cat no. D4003). DNA constructs consisting of target exonic sequences + 250 nucleotides of flanking upstream and downstream intronic sequence, and sequence overlapping RHCglo sequence (Supplementary Table 5) were synthesized by Twist Bioscience and inserted into the linearized RHCglo pDNA using NEBuilder® HiFi DNA Assembly Master Mix (NEB, cat no. E2621S). Reactions were used to transform Mix N Go DH5 cells (Zymo). Plasmid DNA from successful transformants were purified using Qiagen Spin Miniprep kit (Qiagen, cat no. 27104). RHCglo was a gift from Thomas Cooper.

Splicing factor knockdown and minigene experiments

For siRNA knockdown minigene reporter experiments 12-well plates were seeded 24h in advance with 0.1 x 10^6 HepG2 cells. Cells were treated for 48h with 50 pmol of siRNA pool targeting the SF of interest (Supplementary Table 5) or ON-TARGETplus™ siRNA control pool using RNAiMAX reagent (Dharmacon™). At 48h media was removed and cells were transfected with 50 ng of minigene reporter pDNA using Lipofectamine 3000 reagent and incubated for 24h to allow transcript expression. Culture media was removed and cells were washed twice with 1X PBS pH 7.4 and total RNA was extracted using RNeasy® Mini kit (Cat no. 74106, Qiagen) according to the manufacturer's instructions. Each experiment was performed as three biological replicates. The PCPB1 upstream mutant reporter assay was

performed without the siRNA knockdown step as a single replicate and not used in later experiments due to the impact of mutations on the 3SS strength.

Minigene splicing quantification by semi-quantitative RT-PCR

First strand cDNA was generated from 100-150ng total extracted RNA using an Oligo d(T)$_{20}$ primer (Invitrogen, cat no. 18418020) and SuperScript™ IV reverse transcriptase (Invitrogen, cat no. 18090050) according to the manufacturer's instructions. cDNA was PCR amplified using RHCglo gene product specific primers RSV5U and RTRHC (universal to all constructs) and Q5$^{®}$ High-Fidelity 2x Master Mix (NEB, cat no. M0492S) for 28 cycles. PCR products were resolved on 3% agarose gels and band intensity for exon inclusion and exon skipping isoforms was quantified using the ImageJ software package. PSI was calculated as band intensity of the inclusion isoform divided by total intensity of both isoforms.

Confirmation of SF knockdown by Western blot

HepG2 cells were treated with each siRNA and minigene combination as described above. Cells were washed with 1 x PBS then lysed in RIPA buffer (50 mM Tris (pH 7.0), 150 mM NaCl, 0.1% SDS, 0.5% sodium deoxycholate, 1X Triton x-100). Protein from each lysate was resolved on Bis-Tris NuPAGE acrylamide gels (Invitrogen) then transfered to a iBlot® nitrocellulose membrane (Cat no. IB4301001) using the iBLOT® dry blot system. Membranes were blocked for 1h in Azure Fluorescent Blot Blocking Buffer (Azure Biosystems Inc). Rabbit primary antibodies for each SF (Supplementary Table 5) at 1:1000 dilution were added to the membranes and incubated at 4°C overnight with gentle agitation. Membranes were washed with 1X TBST then incubated with fluorescently labeled secondary antibody (see Supplementary Table 5) at a 1:10,000 dilution for 1h at room temperature, washed with 1X TBST and visualized using Azure c600 (Azure Biosystems Inc). Each membrane was then incubated with mouse anti-α-tubulin (1:2500) for 1 hour at room temperature, washed with 1 X TBST, and incubated with secondary antibody (Supplementary Table 5) at room temperature for 1 hour, washed and visualized. Band intensity was quantified using ImageJ software and KD efficiency was calculated as α-tubulin normalized intensity of the SF's band in the SF siRNA treated cells over the control siRNA treated cells (Supplementary Table 5).

Computational resources

All analyses were performed on a Dell Precision 7740 laptop with an NVIDIA Quadro RTX 3000 GPU (5.9GiB, CUDA version 11.4), 16 CPUs (Intel(R) Xeon(R) E-2286M, 2.40GHz), and 67.2 GB of RAM plus 137.4 GiB swap.

Software availability

KATMAP is available at https://gitlab.com/LaptopBiologist/katmap

Supplementary Material

Supplementary Data 1: A compressed archive containing KATMAP outputs for the activity maps used in our analyses. This includes the configuration files, training data, and motif models used to run each analysis; visual summaries of inferred activity and binding; the joint posterior samples over all model parameters; and the tables describing the inferred direct targets of each factor.

Supplementary Table 1: High-level overview of the KATMAP analyses. For each analysis performed, the source and accession of the differential splicing results and the motif models are provided, as well as descriptions of exons used to train the model. High-level summaries of the results are reported as the model comparison z-scores and numbers of significant activity coefficients, along the average number of predicted binding changes per exon. Whether an analysis yielded a significant activity map is indicated for each analysis, and if a given SF had both an RBNS-based PSAM and a PWM we indicate which model we used in our analyses.

Supplementary Table 2: Splicing impacts computed for all cassette exons in the ENCODE knockdowns for each SF with a significant activity map. Each row describes an exon, with the chromosomal coordinates denoting the first and last nucleotide of the exon, along with the splicing impacts assigned to that exon from KATMAP's activity models for the different SFs. If an SF gave a significant activity model in more than one knockdown, we include the splicing impacts from the model with strongest evidence of splicing activity.

Supplementary Table 3: Summary of exons identified as being coregulated by QKI and RBFOX based on the QKI (HepG2) and RBFOX2 (HepG2) knockdowns and the mouse *Rbfox*1/2/3 triple knockdown data.

Supplementary Table 4: Splicing factor perturbations inferred from the ENCODE RNA-binding protein (RBP) knockdown datasets (including non-SF RBPs) using KATMAP's activity models. Each row represents the results of a multinomial regression analysis using precomputed splicing impacts as predictors. The optimal coefficients, associated standard deviations, and sparse shrinkage estimates are provided in separate sheets.

Supplementary Table 5: Details on minigene splicing assays. Contains sequences for semiquantitative RT-PCR primers. Also includes siRNAs, primary and secondary antibodies used in minigene reporter experiments. Final sheet contains the name and measured knockdown efficiency for each RBP/minigene combination normalized to α-tubulin, as well as coordinates (hg19) and exonic/intronic sequences used to construct minigene reporters.

Extended methods

Adaptive importance sampling from the marginal posterior

Before describing the details of this procedure, it is important to note that we work with unconstrained representations of the hyperparameters. That is, for the length-scale L and stationary scale σ, we use their log-transformed values. If we denote the hyperparameters $\varphi_i = (m_i, \lambda_i, L_i, \sigma_i)$, then we denote unconstrained version as $\tilde{\varphi}_i = (m_i, \lambda_i, \log L_i, \log \sigma_i)$. Computing the probability density on the unconstrained hyperparameters requires account-

ing for how the transformation stretches or compresses the density, so $P(\tilde{\varphi}) = \frac{P(\varphi)}{JacDet(\varphi)}$ where $JacDet$ is the Jacobian determinant of the transform. In practice, this amounts to multiplying the constrained density by $L_i \times \sigma_i$.

We begin using unconstrained version of the prior as our first proposal distribution

$$G^{(1)}(\tilde{\varphi}_i) = P(\varphi_i) \times L_i \times \sigma_i$$

From this, we sample n unconstrained hyperparameters, $\tilde{\varphi}_1^{(1)}, .., \tilde{\varphi}_n^{(1)}$ and compute the importance weights which describe how the proposal distribution differs from the target marginal posterior $P(\tilde{\varphi}|Y)$:

$$w_i^{(1)} = \frac{P(\varphi_i^{(1)}|Y) \times L_i^{(1)} \times \sigma_i^{(1)}}{G^{(1)}(\tilde{\varphi}_i^{(1)})}$$

We use these evaluation points to estimate a new proposal distribution, $G^{(2)}$ that better matches the target posterior, with an adaptation procedure rooted in fitting a Student's T mixture model (TMM) with importance weighted expectation-maximization (IW-EM), described below. From this, we generate and evaluate a new sample of hyperparameters. We repeat this procedure, obtaining a series of proposal distributions that approach the target, until we are confident that we will draw a sufficiently large sample from the target.

At each iteration τ, this yields a set of evaluation points drawn from different proposal distributions

$$\tilde{\varphi}_1^{(t=1)}, \tilde{\varphi}_2^{(t=1)}, .., \tilde{\varphi}_n^{(t=1)} \sim G^{(t=1)}$$

$$\tilde{\varphi}_1^{(t=2)}, \tilde{\varphi}_2^{(t=2)}, .., \tilde{\varphi}_n^{(t=2)} \sim G^{(t=2)}$$

$$...$$

$$\tilde{\varphi}_1^{(t=\tau)}, \tilde{\varphi}_2^{(t=\tau)}, .., \tilde{\varphi}_n^{(t=\tau)} \sim G^{(t=\tau)}$$

Each proposal distribution contributed equally to the set of generated evaluation points, so we compute the proposal probabilities from an equally weighted mixture of all past proposal distributions

$$G_{full}^{(\tau)}(\tilde{\varphi}) = \frac{1}{\tau} \sum_{t=1}^{\tau} G^{(t)}(\tilde{\varphi})$$

Then the raw importance weights are

$$w_i^{(t)} = \frac{P\left(\tilde{\varphi}_i^{(t)}|Y\right)}{G_{full}^{(\tau)}\left(\tilde{\varphi}_i^{(t)}\right)}$$

and the raw acceptance probabilities are

$$\pi_i^{(t)} = \frac{w_i^{(t)}}{\max_{j,s} w_j^{(s)}}$$

Following Vehtari et al, we apply a stabilizing correction to these importance weights, which we describe below, to obtain $\hat{\pi}_{1..n}^{(1..\tau)}$

From these acceptance probabilities, we can determine what fraction of samples we expect to accept if we were to stop at the current iteration

$$f_{acc} = \frac{1}{n\tau} \sum_{i=1}^{n} \sum_{t=1}^{\tau} \hat{\pi}_i^{(t)}$$

which also provides an estimate of the effective sample size

$$n_{eff} = n f_{acc}$$

To determine whether we can stop iterating, we ask whether there is at least a 99% probability the number of accepted samples n_{acc} exceeds the desired sample size n_{final}. The number of samples we will accept, n_{acc}, is the sum of independent Bernoulli trials with variable success probabilities and so follows a Poisson-binomial distribution. As this distribution is never more variable than a binomial distribution with the same mean and number of trials, we use the binomial distribution to obtain a conservative estimate of the lower bound on n_{acc}.

$$P(n_{acc} \geq n_{final}) \geq .99, \qquad n_{acc} \sim Binomial(N = n\tau, p = f_{acc})$$

Adapting the proposal distribution

We first use the EM procedure described below to fit a Student's T mixture model (TMM) to the target, obtaining mixing weights $\omega_{1..C}$, means $\mu_{1..C}$, and covariance matrices $\Sigma_{1..C}$, as well as a degree-of-freedom parameter ν shared by all C component distributions. This provides an estimate of the target that can account for both its local structure and the heaviness of its tails. However, in the first few iterations of adaptive importance sampling the effective sample is typically small (< 5) and we found the TMM-based proposal to occasionally become too concentrated around the modes of the target density, requiring extra rounds of adaption to properly explore tails. To decrease the time to convergence, we employ two heuristics to stabilize the adaptation in the early rounds of importance sampling. Both heuristics aim to ensure the proposal is distributed around the target density, but always somewhat more dispersed than the true target.

First, we double the fitted the covariance matrices, defining the mixture distribution which captures the local structure of the target density as

$$TMM_{local}(x) = \sum_{c=1}^{C} \omega_c \mathcal{T}(x; \nu, \mu_c, 2\Sigma_c)$$

By intentionally making the proposal broader than the IW-EM estimate of the target density, we sacrifice somewhat the efficiency with which we obtain samples from the target

density. However, this ensures that subsequent evaluation points fall in the tails of the target, decreasing the risk that the adaptation overshoots.

However, when the effective sample from the target is small, we lack precise information about the finer details of the distribution's shape and TMM_{local} may still overfit. We therefore construct an even more conservative proposal. When n_{eff} is close to zero, IW-EM will shift the means in the direction of a mode in the target, but will learn little about the target's covariance. Given that covariance of the target remains obscure, we may as well continue to use the previous proposal distribution. We approximate this by computing the covariance of the proposals $\hat{\Sigma}_{proposal}$. If n_{eff} is somewhat larger than zero, we might have enough information to estimate the global covariance of the target, $\hat{\Sigma}_{target}$, which we can obtain by computing an importance weighted covariance from the proposals. To find a intermediate between these two extremes, we take a weighted average that approaches $\hat{\Sigma}_{target}$ as n_{eff} increases

$$\hat{\Sigma} = \frac{1}{1 + n_{eff}} \left(\hat{\Sigma}_{proposal} + 2 \times n_{eff} \times \hat{\Sigma}_{target} \right)$$

The Bayesian flavor of transitioning from an agnostic *a priori* assumption to a data-driven estimate as more information becomes available is not coincidental. This estimator can be thought of as being twice the posterior mean, Σ', of the target covariance estimated using an inverse-Wishart Gaussian conjugate model with a prior scale matrix based on the initial proposal distribution, $\Psi_0 = \frac{1}{2}\hat{\Sigma}_{proposal}$, and $\nu_0 = p + 2$ degrees of freedom, where p is the number of rows/columns in the covariance matrix,

$$\Sigma' = \frac{1}{n + \nu_0 - p - 1} (\Psi_0 + n\Sigma) = \frac{1}{1 + n} \left(\frac{1}{2}\hat{\Sigma}_{proposal} + n\Sigma \right)$$

$$\hat{\Sigma} = 2\Sigma'$$

where Σ is a covariance matrix computed from n data points.

From this, we construct our more conservative proposal distribution

$$TMM_{global}(x) = \sum_{c=1}^{C} \omega_c \mathcal{T} \left(x; \nu = 35, \mu_c, \hat{\Sigma} \right)$$

This uses the IW-EM estimates of target's location, but replaces the covariances with a global estimate of the target's covariance, to ensure new proposals explore the target density's tails.

Our final proposal overlays these as a mixture, with the mixing weights depending upon n_{eff}

$$G(x) = \frac{n_{eff}}{\rho + n_{eff}} TMM_{local}(X) + \frac{\rho}{\rho + n_{eff}} TMM_{global}(X)$$

As n_{eff} increases, the composition of the mixture shifts from the conservative TMM_{global} to TMM_{local}. We choose $\rho = 4$ such that when $n_{eff} = 1$, 80% of the new proposals come from the conservative mixture, at $n_{eff} = 20$ it is about 16% proposals, and when $n_{eff} = 100$ the proposal is largely derived from the local TMM, with less than 4% of proposal derived from the conservative mixture. Whatever n_{eff}, the updated proposal always moves towards

modes of the target density, because both TMM_{local} and TMM_{global} use the means and mixing proportions estimated from IW-EM.

Importance-weighted Expectation Maximization for a multivariate T-mixture

To construct proposals that can robustly adapt to the target, we follow Cappé et al. (2008) and fit a TMM by importance weighted Expectation-Maximization (IW-EM). One approach to conceptualizating a multivariate T distribution is a scale-mixture of multivariate Gaussians. So a mixture of Student T distributions can be described as a mixture of Gaussians with two latent variable per realization x_i

$$z_i \sim Categorical(\omega_{1..K})$$

$$\gamma_i \sim Gamma\left(\frac{\nu_{z_i}}{2}, \frac{\nu_{z_i}}{2}\right)$$

$$x_i \sim MvNormal\left(\mu_{z_i}, \frac{\Sigma_{z_i}}{\gamma_i}\right)$$

where ν is the degrees of freedom. $\omega_{1..K}$ is a vector of mixing weights describing the fraction of the full mixture each component distribution contributes. $\mu_{1..K}$ and $\Sigma_{1..K}$ are means and covariances of the K component distributions. $z_{1..N}$ is vector of unobserved variables indicating which component each observation x_i truly derives from. The second set of unobserved variables $\gamma_{1..N}$ come from the scale-mixture representation of the Student T distribution and describe how Gaussian z_i was rescaled to generate each realization x_i.

While, Cappé et al. (2008) initialize once and perform a single EM-update at each step of importance sampling, we instead reinitialize at each importance sampling iteration t and perform t EM-updates to prevent the adaptive distribution from getting trapped. We use K-means clustering of the proposals with $10 \times t$ components, to define the initial means ensure the components are well-distributed throughout the proposal distribution.

Expectation Step

For each observed x_i, there are two unobserved variables, $z_{1..N}$ and $\gamma_{1..N}$. The expectation step infers these latent variables by computing their expected values given the parameters of the mixture distribution $(\omega_{1..K}, \mu_{1..K}, \Sigma_{1..K})$.

To estimate the probability that x_i arose from a given component distribution d, we compute

$$\eta_{id} = E\left[z_i = d | X\right] = \frac{\omega_d \mathcal{T}(x_i; \nu_d, \mu_d^{(t)}, \Sigma_d^{(t)})}{\sum_{\tilde{d}} \omega_{\tilde{d}} \mathcal{T}(x_i; \nu_d, \mu_{\tilde{d}}^{(t)}, \Sigma_{\tilde{d}}^{(t)})}$$

where $\mathcal{T}$ denotes the multivariate Student-t probability density function as implemented in scipy.

Then to estimate the scale of the Gaussian from which x_i arose, assuming it belongs to the d-th Student T distribution, we compute

$$u_{id} = E\left[\gamma_i | X, z_i = d\right] = \frac{\nu_d + r}{\nu_d + \left(x_i - \mu_d^{(t)}\right)^T \left(\Sigma_d^{(t)}\right)^{-1} \left(x_i - \mu_d^{(t)}\right)}$$

where r is the dimension of the distribution (in our case, the number of hyperparameters in the additive model).

Maximization Step

The maximization step uses these estimates of the latent variables to refine the estimates of the mixture distribution's parameters. These estimates are essentially averages weighted by the component probabilities η_{id}. To fit the mixture to the target rather than the proposals, we follow Cappé et al. (2008) compute the updates as weighted averages using the importance weights $\pi_{i..N}$ to weight the observations:

$$\omega_j^{(t+1)} = \frac{\sum_i \eta_{ij} \pi_i}{\sum_{\tilde{i}} \pi_{\tilde{i}}}$$

$$\mu_j^{(t+1)} = \frac{\sum_i \eta_{ij} \pi_i x_i}{\sum_{\tilde{i}} \eta_{\tilde{i}j} \pi_{\tilde{i}}}$$

$$\Sigma_j^{(t+1)} = \frac{\sum_i \eta_{ij} u_{ij} \pi_i \left(x_i - \mu_j^{(t+1)}\right) \left(x_i - \mu_j^{(t+1)}\right)^T}{\sum_{\tilde{i}} \eta_{\tilde{i}j} \pi_{\tilde{i}}}$$

The degree-of-freedom parameter, ν, controls the tails of the distribution. There is no closed-form estimate of ν and so it must be estimated by optimization. To make this tractable, we constrain the components of the mixture to all share the same ν parameter. By reducing this to a single parameter, we can use a binary search to update $\nu^{(t+1)}$, considering values between $\nu \in (4, 100)$. The importance weighted EM scheme optimizes the entropy of the normalized importance weights we would get if we used the mixture as the proposal distribution

$$w_i = \frac{P(x_i)}{\sum_{j=1}^{K} \omega_j^{(t+1)} \mathcal{T}\left(x_i; \nu^{(t+1)}, \mu_j^{(t+1)}, \Sigma_j^{(t+1)}\right)}$$

$$\bar{w}_i = \frac{w_i}{\sum_{\tilde{i}}^{N} w_{\tilde{i}}}$$

$$\mathbf{H} = -\sum_i^{N} \bar{w}_i log(\bar{w}_i)$$

So we evaluate the entropy at twenty values of ν between 4 and 100 and identify all local maxima on this grid. We then perform binary searches between the neighbors of each maxima to hone in on the true local maxima and set $\nu^{(t+1)}$ to the value of ν that yielded the highest entropy.

Stabilizing the importance weights

The set of importance weights $\pi_{1..n}^{(1..T)}$ represent an estimate of how the proposal distribution differs from the target distribution. To properly characterize the target distribution, the importance weights must be based on a sufficient number of proposals spanning typical set of the target. If the proposal distribution is very different from the target, the computed importance weights will poorly characterize the target and importance sampling will be inefficient or untrustworthy.

Vehtari et al provide an approach for obtaining more stable estimates of the importance weights by fitting a Pareto distribution to the tail of $\pi_{1..n}^{(1..T)}$. Using the implementation of this Pareto smoothing from the arviz library (Kumar et al. 2019), we compute smoothed weights $\hat{\pi}_{1..n}^{(1..T)}$. Importantly, the shape parameter of the fitted Pareto distribution, $\hat{k}$, is diagnostic of whether importance sampling will be reliable.

We only accept posterior samples based on the Pareto-smoothed importance weights. Occasionally in the early iterations of adapting the proposal distribution, however, the range of values computed for the importance weights may be too extreme for Pareto smoothing to yield reliable importance estimates (e.g. $\hat{k} > 0.7$). In these cases, we use an older, simpler approach to stabilize the weights (Ionides 2008). This defines a cutoff based on the total number of importance weights N_π and average importance weight $\bar{\pi}$

$$\gamma = \bar{\pi}\sqrt{N_\pi}$$

and then replaces all importance weights greater than γ with γ

$$\hat{\pi}_i = \begin{cases} \pi_i & \text{if } \pi_i < \gamma \\ \gamma & \text{if } \pi_i \geq \gamma \end{cases}$$

The way in which this flattens the importance weights will bias the importance estimates by accepting some proposals which should be rejected. When used to adapt the proposal distribution, this should result in an estimate of the target distribution that is broader than the true target. This decreases the efficiency with which the proposal is adapted, but prevents it from collapsing to a subset of the target distribution.

Assumptions about the spatial correlations: L, σ

There are two identifiability issues that can arise with Gaussian process priors. If the length-scale L is near zero, it corresponds to the biologically implausible assumption of complete spatial independence of the activity coefficients. But it also introduces problems for statistical inference. As we perform inference with respect to the log-transform of L, all values of $logL \ll 0$ are essentially equivalent. Similarly, length-scales much larger than the maximum distance between regions analyzed $L \gg max(|j - k|)$ are practically indistinguishable.

To prevent the inference algorithm from wasting computation by exploring large but practically equivalent regions of parameter space, we construct a prior that avoids both large and small values of L. To avoid overly large length-scales, we use the right-tail of a generalized gamma distribution

$$GG(x; a, d, p) = \Gamma\left(\frac{d}{p}\right)\frac{p}{a^d}x^{d-1}exp\left(\left(\frac{x}{a}\right)^p\right)$$

This does not prevent L from being close to zero. We avoid these small values by convolving the generalized gamma with the left tail of an inverse-gamma distribution

$$HGG(x; a, d, p, b) = \frac{1}{Z(a, d, p, b)}exp\left(-\frac{b}{x}\right)GG(x; a, d, p)$$

where the normalizing constant is computed numerically ('scipy.integrate.quad')

$$Z(a, d, p, b) = \int_0^\infty exp\left(-\frac{b}{x}\right)GG(x; a, d, p)$$

We set the prior on the length-scale as

$$L \sim HGG(x; a = 20, d = .01, p = 5, b = 1)$$

which places 99% of the prior probability on the interval $0.32 < L < 22.6$.

We use this same distribution to exclude small values of the stationary standard deviation. If σ is close to zero, all of the activity coefficients become constrained to be close to zero solely by the prior. One way of conceptualizing the prior on σ is that it reflects different assumptions we might make about the regression coefficients in our regression analysis. We would never perform a regression analysis where we assumed *a priori* that all of the regression coefficient were zero. Whether the coefficients should be close to zero is something we wish to learn from the data, not assume in advance. So we define our prior on the stationary standard deviation as

$$\sigma \sim HGG(x; a = 2, d = .25, p = 5, b = .5)$$

which places 99% of the prior probability on the interval $0.14 < \sigma < 8.7$

Model comparison against the reduced model

To evaluate whether the data support splicing activity, we compare the full model against a reduced model without splicing activity. To verify that any improvement over the reduced model did not reflect overfitting to the training data, we sought to evaluate each model's performance on held-out data. However, to sidestep the need to rerun the inference algorithm dozens of times for each knockdown, we employ the Pareto-smoothed importance sampling approximation to leave-one-out (PSIS-LOO) cross-validation proposed by Vehtari, Simpson, et al. (2024)

PSIS-LOO uses the posterior sample from training on all data as a proposal distribution to obtain estimates regarding the leave-one-out posterior distribution that would be obtained by excluding the i-th observation. It leverages the insight that the ratio between the complete and the leave-one-out posteriors depends only on the left-out observation, with all other terms in the posteriors cancelling out.

$$r_i(\theta) \propto \frac{P(\theta|Y_{-i})}{P(\theta|Y)}$$

$$\propto \frac{P(Y_1|\theta) \times .. \times P(Y_{i-1}|\theta) \times P(Y_{i+1}|\theta) \times .. \times P(Y_N|\theta) \times P(\theta)}{P(Y_1|\theta) \times .. \times P(Y_{i-1}|\theta) \times \boldsymbol{P(Y_i|\theta)} \times P(Y_{i+1}|\theta) \times .. \times P(Y_N|\theta) \times P(\theta)}$$

$$\propto \frac{1}{P(Y_i|\theta)}$$

These importance weights are then stabilized using the aforementioned Pareto smoothing to obtain $\tilde{r}_i(\theta)$. We estimate the log expected posterior predictive probability for each held-out exon i by computing the importance-weighted average over the complete posterior sample

$$lpd_i = \log \left(\frac{1}{\sum_{k=1}^{M} \tilde{r}_i(\theta_k)} \sum_{j=1}^{M} \tilde{r}_i(\theta_j) P(Y_i|\theta_j) \right)$$

To combine this across all held-out exons, we sum the estimated log-probabilities, exactly as we would when computing the model's objective function

$$\widehat{elpd} = \sum_{i=1}^{N} lpd_i$$

We chose this estimator as it is the finite sample version of the widely-applicable information criterion (WAIC). We compute this for the full, $\widehat{elpd}_{Full}$, and reduced models, $\widehat{elpd}_{\emptyset}$, and summarize the model comparison as $\widehat{MC} = \widehat{elpd}_{Full} - \widehat{elpd}_{\emptyset}$. A positive value of $\widehat{MC}$ means the full model with splicing activity is a better explanation of the data than the reduced model without splicing activity. To quantify how confident we are in this evidence for splicing activity, we bootstrap the exons 10,000 times, estimating $\widehat{MC}$ each time, and use these obtain a standard deviation $SD(\widehat{MC})$. We then compute a Z-score

$$Z(\widehat{MC}) = \frac{\widehat{MC}}{SD(\widehat{MC})}$$

Inferring the underlying correlation of variables with known uncertainty

In some of our analyses we wanted to determine the correlation between estimated values $\hat{x}_{1..N}$ and $\hat{y}_{1..N}$, when each of the estimates was associated with known uncertainty $\sigma_{x,1..N}$ and $\sigma_{y,1..N}$. Simply computing the correlation between $\hat{x}_{1..N}$ and $\hat{y}_{1..N}$ would underestimate the true association between the true values x and y, which is obscured by the uncertainty in the estimates. We therefore employ a hierarchical model to disentangle the underlying association from the uncertainty in the individual estimates.

We first assume that the estimated values $\hat{x}_{1..N}$ and $\hat{y}_{1..N}$ are normally distributed around the true values $x_{1..N}$ and $y_{1..N}$ with standard deviations $\sigma_{x,1..N}$ and $\sigma_{y,1..N}$

$$\hat{x}_i \sim Normal(x_i, \sigma_{x,i})$$

$$\hat{y}_i \sim Normal(y_i, \sigma_{y,i})$$

we then assume that the underlying true values follow a latent multivariate normal distribution with the correlation structure of interest

$$(x, y)_{1..N} \sim MvNormal \left(\mu = \begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \Sigma = \begin{bmatrix} \sigma_X^2 & \rho \\ \rho & \sigma_Y^2 \end{bmatrix} \right)$$

Our primary goal is infer ρ, disentangled from the uncertainty. For the full hierarchical model, we place priors on the means and standard deviations of this latent Gaussian using the means and standard deviations of the noisy estimates $\hat{x}_{1..N}$ and $\hat{y}_{1..N}$. We also place a relatively uninformative prior on the correlation coefficient in logit scale.

$$(\mu_X, \mu_Y) \sim MvNormal \left(\mu = \begin{bmatrix} E[\hat{x}_{1..N}] \\ E[\hat{y}_{1..N}] \end{bmatrix}, \Sigma = \begin{bmatrix} Var[\hat{x}_{1..N}] & 0 \\ 0 & Var[\hat{y}_{1..N}] \end{bmatrix} \right)$$

$$(\sigma_X, \sigma_Y) \sim MvNormal \left(\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \Sigma = \begin{bmatrix} Var[\hat{x}_{1..N}] & 0 \\ 0 & Var[\hat{y}_{1..N}] \end{bmatrix} \right)$$

$$\tilde{\rho} \sim Normal(0, 1.4)$$

$$\rho = 1 - 2 \times InvLogit(\tilde{\rho})$$

$$(x, y)_{1..N} \sim MvNormal \left(\mu = \begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \Sigma = \begin{bmatrix} \sigma_X^2 & \rho \\ \rho & \sigma_Y^2 \end{bmatrix} \right)$$

$$\hat{x}_i \sim Normal(x_i, \sigma_{x,i})$$

$$\hat{y}_i \sim Normal(y_i, \sigma_{y,i})$$

In practice, the parameters of hierarchical models often have complicated associations with each other that prevents efficient exploration of the posterior. We follow the standard practice to resolve this by using the noncentered parameterization of the latent Gaussian and representing $(x, y)_{1..N}$ as the product of a unit normal distribution and the Cholesky factor of the covariance matrix

$$z_{1..N} \sim MvNormal \left(\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right)$$

$$LL^T = \begin{bmatrix} \sigma_X^2 & \rho \\ \rho & \sigma_Y^2 \end{bmatrix}$$

$$\begin{bmatrix} x_{1..N} & y_{1..N} \end{bmatrix} = \begin{bmatrix} \mu_X & \mu_Y \end{bmatrix}_{1..N} + zL$$

We use the No-U-Turn sampler, as implemented in PyMC3 (3.11.2), to draw 4,000 samples from the joint posterior over all parameters $(\rho, \mu_X, \mu_Y, \sigma_X, \sigma_Y, x_{1..N}, y_{1..N})$ in 4 chains with the target acceptance probability set to 0.9.

Working with differential splicing analyses

Recovering effect sizes estimates from rMATS outputs

rMATs reports the inclusion levels and inclusion level difference using direct summaries of the read counts, rather than the effect sizes inferred by the underlying model. The actual parameters of the rMATS model, however, are defined in log-odds scale. This log-odds representation allows interval estimates to be naturally defined around the effect size in a way that respects the constraint that $0 < \psi < 1$. The read count based summaries, however, may equal 0 or 1, and cannot be converted as is to log-odds scale. To resolve this, we recompute PSI with pseudocounts added to the observed reads, which is equivalent to estimating the inclusion read fraction with a beta-binomial model.

$$f_i = \frac{a + IJC_i}{a + b + IJC_i + SJC_i}$$

We identify the appropriate pseudocounts from the data itself, fitting a beta-binomial distribution to the read counts of all exons. We then estimate the inclusion read fraction for each exon and convert these to PSI

$$\psi_i = \frac{f_i L_i^{(S)}}{f_i L_i^{(S)} + (1 - f_i) L_i^{(I)}}$$

where the $L_i^{(I)}$ and $L_i^{(S)}$ account for the differing amounts of informative sequence associated with the inclusion and exclusion junctions respectively.

Reconstructing uncertainty about effect size from rMATS outputs

We sought to reconstruct the posterior standard deviation for these effect sizes from the rMATS p-values. rMATS computes its p-values using a likelihood ratio test, comparing the likelihood of the full model to that where the effect size is set to a small value near zero, $LR = \frac{Lk_\emptyset}{Lk_{full}}$. The test statistic $\Lambda_i = -2logLR$ follows a chi-squared distribution and can be recovered by inputting the rMATS p-value, p_i for exon i into the ch-squared inverse cumulative density function

$$\Lambda_i = CDF^{-1}_{\chi^2(df=1)}(1 - p_i)$$

Describing the uncertainty about the effect sizes with a mean and standard deviation implies a normal approximation to the likelihood. The likelihood-ratios, therefore, can be thought of as comparing the density of this normal distribution at its mode to a point in its tail. The likelihood of the full model is evaluated at its mode (e.g. effect size)

$$L^{full} = \frac{1}{\sigma_i \sqrt{2\pi}} exp\left[-\frac{(\delta_i - \delta_i)^2}{2\sigma_i^2}\right] = \frac{1}{\sigma_i \sqrt{2\pi}}$$

and is entirely determined by the standard deviation. The null model, however, evaluates the likelihood at a point away from the mode, $\delta_\emptyset$, assuming the effect size is close to zero

$$L^{null} = \frac{1}{\sigma_i \sqrt{2\pi}} exp\left[-\frac{(\delta_\emptyset - \delta_i)^2}{2\sigma_i^2}\right]$$

The likelihood ratio is then

$$LR_i = \frac{\frac{1}{\sigma_i \sqrt{2\pi}} exp\left[-\frac{(\delta_\emptyset - \delta_i)^2}{2\sigma_i^2}\right]}{\frac{1}{\sigma_i \sqrt{2\pi}}} = exp\left[-\frac{(\delta_\emptyset - \delta_i)^2}{2\sigma_i^2}\right]$$

and the negative log-likelihood ratio is

$$logLR = -\frac{(\delta_\emptyset - \delta_i)^2}{2\sigma_i^2}$$

To get the variance σ_i^2 of the normal approximation we can rearrange

$$\sigma_i^2 = \frac{(\delta_\emptyset - \delta_i)^2}{-2logLR}$$

The Chi2 test statistic, $\Lambda_i = -2logLR_i$, is the denominator so this can be written:

$$\sigma_i^2 = \frac{(\delta_\emptyset - \delta_i)^2}{\Lambda_i}$$

References

Bak, Maciej et al. (May 15, 2024). "MAPP unravels frequent co-regulation of splicing and polyadenylation by RNA-binding proteins and their dysregulation in cancer". In: *Nature Communications* 15.1. Publisher: Nature Publishing Group, p. 4110. ISSN: 2041-1723. DOI: 10.1038/s41467-024-48046-1.

Balwierz, Piotr J. et al. (May 1, 2014). "ISMARA: automated modeling of genomic signals as a democracy of regulatory motifs". In: *Genome Research* 24.5. Company: Cold Spring Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Harbor Laboratory Press Publisher: Cold Spring Harbor Lab, pp. 869–884. ISSN: 1088-9051, 1549-5469. DOI: 10.1101/gr.169508.113.

Best, Andrew et al. (Sept. 11, 2014). "Human Tra2 proteins jointly control a CHEK1 splicing switch among alternative and constitutive target exons". In: *Nature Communications* 5, p. 4760. ISSN: 2041-1723. DOI: 10.1038/ncomms5760.

Boutz, Paul L. et al. (July 1, 2007). "A post-transcriptional regulatory switch in polypyrimidine tract-binding proteins reprograms alternative splicing in developing neurons". In: *Genes & Development* 21.13. Company: Cold Spring Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Harbor Laboratory Press Publisher: Cold Spring Harbor Lab, pp. 1636–1652. ISSN: 0890-9369, 1549-5477. DOI: 10.1101/gad.1558107.

Bradbury, James et al. (2018). *JAX: composable transformations of Python+NumPy programs.* Version 0.3.13.

Brosseau, Jean-Philippe et al. (Feb. 1, 2014). "Tumor microenvironment–associated modifications of alternative splicing". In: *RNA* 20.2. Company: Cold Spring Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Harbor Laboratory Press Publisher: Cold Spring Harbor Lab, pp. 189–201. ISSN: 1355-8382, 1469-9001. DOI: 10.1261/rna.042168.113.

Cao, Wenguang et al. (Sept. 2012). "Control of alternative splicing by forskolin through hnRNP K during neuronal differentiation". In: *Nucleic Acids Research* 40.16, pp. 8059–8071. ISSN: 0305-1048. DOI: 10.1093/nar/gks504.

Cappé, Olivier et al. (Dec. 1, 2008). "Adaptive importance sampling in general mixture classes". In: *Statistics and Computing* 18.4, pp. 447–459. ISSN: 1573-1375. DOI: 10.1007/s11222-008-9059-x.

Chen, Charlie Degui, Ryuji Kobayashi, and David M. Helfman (Mar. 1, 1999). "Binding of hnRNP H to an exonic splicing silencer is involved in the regulation of alternative splicing of the rat -tropomyosin gene". In: *Genes & Development* 13.5, pp. 593–606. ISSN: 0890-9369.

Cheung, Hannah C. et al. (Aug. 1, 2009). "Splicing factors PTBP1 and PTBP2 promote proliferation and migration of glioma cell lines". In: *Brain* 132.8, pp. 2277–2288. ISSN: 0006-8950. DOI: 10.1093/brain/awp153.

Chou, Min-Yuan et al. (Jan. 1999). "hnRNP H Is a Component of a Splicing Enhancer Complex That Activates a c-src Alternative Exon in Neuronal Cells". In: *Molecular and Cellular Biology* 19.1, pp. 69–77. ISSN: 0270-7306.

Coelho, Miguel B et al. (Mar. 4, 2015). "Nuclear matrix protein Matrin3 regulates alternative splicing and forms overlapping regulatory networks with PTB". In: *The EMBO Journal* 34.5. Num Pages: 668 Publisher: John Wiley & Sons, Ltd, pp. 653–668. ISSN: 0261-4189. DOI: 10.15252/embj.201489852.

Eden, Eran et al. (Feb. 3, 2009). "GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists". In: *BMC Bioinformatics* 10.1, p. 48. ISSN: 1471-2105. DOI: 10.1186/1471-2105-10-48.

Ghanem, Louis R. et al. (July 30, 2018). "Poly(C)-Binding Protein Pcbp2 Enables Differentiation of Definitive Erythropoiesis by Directing Functional Splicing of the Runx1 Transcript". In: *Molecular and Cellular Biology* 38.16, e00175–18. ISSN: 0270-7306. DOI: 10.1128/MCB.00175-18.

Graveley, Brenton R., Klemens J. Hertel, and Tom Maniatis (Nov. 16, 1998). "A systematic analysis of the factors that determine the strength of pre-mRNA splicing enhancers". In: *The EMBO Journal* 17.22. Num Pages: 6756 Publisher: John Wiley & Sons, Ltd, pp. 6747–6756. ISSN: 0261-4189. DOI: 10.1093/emboj/17.22.6747.

Gupta, Kavi et al. (Jan. 16, 2024). "Improved modeling of RNA-binding protein motifs in an interpretable neural model of RNA splicing". In: *Genome Biology* 25.1, p. 23. ISSN: 1474-760X. DOI: 10.1186/s13059-023-03162-x.

Han, Areum et al. (Jan. 30, 2014). "De Novo Prediction of PTBP1 Binding and Splicing Targets Reveals Unexpected Features of Its RNA Recognition and Function". In: *PLOS Computational Biology* 10.1. Publisher: Public Library of Science, e1003442. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1003442.

Hardy, R. J. (Oct. 1, 1998). "QKI expression is regulated during neuron-glial cell fate decisions". In: *Journal of Neuroscience Research* 54.1, pp. 46–57. ISSN: 0360-4012. DOI: `10.1002/(SICI)1097-4547(19981001)54:1<46::AID-JNR6>3.0.CO;2-H`.

Heiner, Monika et al. (Jan. 1, 2010). "HnRNP L-mediated regulation of mammalian alternative splicing by interference with splice site recognition". In: *RNA Biology* 7.1. Publisher: Taylor & Francis _eprint: https://doi.org/10.4161/rna.7.1.10402, pp. 56–64. ISSN: 1547-6286. DOI: `10.4161/rna.7.1.10402`.

Hugh-White, Rupert (2021). "Analysing RNA-seq datasets to determine how pre-mRNA splicing is regulated by RNA binding proteins and cis-acting elements". PhD thesis. King's College London.

Huttlin, Edward L. et al. (May 27, 2021). "Dual proteome-scale networks reveal cell-specific remodeling of the human interactome". In: *Cell* 184.11, 3022–3040.e28. ISSN: 0092-8674. DOI: `10.1016/j.cell.2021.04.011`.

Ionides, Edward L. (June 1, 2008). "Truncated Importance Sampling". In: *Journal of Computational and Graphical Statistics*. Publisher: Taylor & Francis. DOI: `10.1198/106186008X320456`.

Jacko, Martin et al. (Feb. 21, 2018). "Rbfox Splicing Factors Promote Neuronal Maturation and Axon Initial Segment Assembly". In: *Neuron* 97.4, 853–868.e6. ISSN: 1097-4199. DOI: `10.1016/j.neuron.2018.01.020`.

Jens, Marvin et al. (Nov. 9, 2022). *RBPamp: Quantitative Modeling of Protein-RNA Interactions in vitro Predicts in vivo Binding*. Pages: 2022.11.08.515616 Section: New Results. DOI: `10.1101/2022.11.08.515616`.

Ji, Xinjun et al. (Mar. 18, 2016). "CP binding to a cytosine-rich subset of polypyrimidine tracts drives a novel pathway of cassette exon splicing in the mammalian transcriptome". In: *Nucleic Acids Research* 44.5, pp. 2283–2297. ISSN: 0305-1048. DOI: `10.1093/nar/gkw088`.

Královičová, Jana et al. (July 6, 2018). "PUF60-activated exons uncover altered 3 splice-site selection by germline missense mutations in a single RRM". In: *Nucleic Acids Research* 46.12, pp. 6166–6187. ISSN: 0305-1048. DOI: `10.1093/nar/gky389`.

Krismer, Konstantin et al. (Aug. 25, 2020). "Transite: A Computational Motif-Based Analysis Platform That Identifies RNA-Binding Proteins Modulating Changes in Gene Expression". In: *Cell Reports* 32.8, p. 108064. ISSN: 2211-1247. DOI: `10.1016/j.celrep.2020.108064`.

Kumar, Ravin et al. (Jan. 15, 2019). "ArviZ a unified library for exploratory analysis of Bayesian models in Python". In: *Journal of Open Source Software* 4.33, p. 1143. ISSN: 2475-9066. DOI: `10.21105/joss.01143`.

Lambert, Nicole et al. (June 5, 2014). "RNA Bind-n-Seq: quantitative assessment of the sequence and structural binding specificity of RNA binding proteins". In: *Molecular Cell* 54.5, pp. 887–900. ISSN: 1097-4164. DOI: `10.1016/j.molcel.2014.04.016`.

Lang, Benjamin et al. (July 9, 2021). "Matrix-screening reveals a vast potential for direct protein-protein interactions among RNA binding proteins". In: *Nucleic Acids Research* 49.12, pp. 6702–6721. ISSN: 0305-1048. DOI: `10.1093/nar/gkab490`.

Lerner, Michael R. et al. (Jan. 1980). "Are snRNPs involved in splicing?" In: *Nature* 283.5743. Publisher: Nature Publishing Group, pp. 220–224. ISSN: 1476-4687. DOI: `10.1038/283220a0`.

Llorian, Miriam et al. (Sept. 2010). "Position-dependent alternative splicing activity revealed by global profiling of alternative splicing events regulated by PTB". In: *Nature Structural & Molecular Biology* 17.9. Publisher: Nature Publishing Group, pp. 1114–1123. ISSN: 1545-9985. DOI: `10.1038/nsmb.1881`.

Maniatis, Tom and Robin Reed (Feb. 1987). "The role of small nuclear ribonucleoprotein particles in pre-mRNA splicing". In: *Nature* 325.6106. Publisher: Nature Publishing Group, pp. 673–678. ISSN: 1476-4687. DOI: `10.1038/325673a0`.

Merendino, L. et al. (Dec. 16, 1999). "Inhibition of msl-2 splicing by Sex-lethal reveals interaction between U2AF35 and the 3' splice site AG". In: *Nature* 402.6763, pp. 838–841. ISSN: 0028-0836. DOI: `10.1038/45602`.

Nocedal, Jorge and Stephen Wright (Dec. 11, 2006). *Numerical Optimization*. Google-Books-ID: VbHYoSyelFcC. Springer Science & Business Media. 686 pp. ISBN: 978-0-387-40065-5.

Page-McCaw, P. S., K. Amonlirdviman, and P. A. Sharp (Dec. 1999). "PUF60: a novel U2AF65-related splicing activity". In: *RNA (New York, N.Y.)* 5.12, pp. 1548–1560. ISSN: 1355-8382. DOI: `10.1017/s1355838299991938`.

Rasmussen, Carl Edward and Christopher K. I. Williams (2006). *Gaussian processes for machine learning*. Adaptive computation and machine learning. MIT Press. I-XVIII, 1-248. ISBN: 0-262-18253-X.

Ray, Debashish, Hilal Kazan, Esther T. Chan, et al. (July 2009). "Rapid and systematic analysis of the RNA recognition specificities of RNA-binding proteins". In: *Nature Biotechnology* 27.7. Publisher: Nature Publishing Group, pp. 667–670. ISSN: 1546-1696. DOI: `10.1038/nbt.1550`.

Ray, Debashish, Hilal Kazan, Kate B. Cook, et al. (July 2013). "A compendium of RNA-binding motifs for decoding gene regulation". In: *Nature* 499.7457. Publisher: Nature Publishing Group, pp. 172–177. ISSN: 1476-4687. DOI: `10.1038/nature12311`.

Rothrock, Caryn R, Amy E House, and Kristen W Lynch (Aug. 3, 2005). "HnRNP L represses exon splicing via a regulated exonic splicing silencer". In: *The EMBO Journal* 24.15. Num Pages: 2802 Publisher: John Wiley & Sons, Ltd, pp. 2792–2802. ISSN: 0261-4189. DOI: `10.1038/sj.emboj.7600745`.

Rubin, Donald B. (Jan. 1981). "The Bayesian Bootstrap". In: *The Annals of Statistics* 9.1. Publisher: Institute of Mathematical Statistics, pp. 130–134. ISSN: 0090-5364, 2168-8966. DOI: `10.1214/aos/1176345338`.

Rubin, Jonathan D. et al. (June 2, 2021). "Transcription factor enrichment analysis (TFEA) quantifies the activity of multiple transcription factors from a single experiment". In: *Communications Biology* 4.1. Publisher: Nature Publishing Group, pp. 1–15. ISSN: 2399-3642. DOI: `10.1038/s42003-021-02153-7`.

Rue, Håvard, Sara Martino, and Nicolas Chopin (2009). "Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations". In: *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 71.2. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1467-9868.2008.00700.x, pp. 319–392. ISSN: 1467-9868. DOI: `10.1111/j.1467-9868.2008.00700.x`.

Shen, Shihao et al. (Dec. 23, 2014). "rMATS: Robust and flexible detection of differential alternative splicing from replicate RNA-Seq data". In: *Proceedings of the National Academy of Sciences* 111.51. Publisher: Proceedings of the National Academy of Sciences, E5593–E5601. DOI: `10.1073/pnas.1419161111`.

Sherman, Larry et al. (Nov. 1997). "Interdomain binding mediates tumor growth suppression by the NF2 gene product". In: *Oncogene* 15.20. Publisher: Nature Publishing Group, pp. 2505–2509. ISSN: 1476-5594. DOI: 10.1038/sj.onc.1201418.

Spellman, Rachel, Miriam Llorian, and Christopher W.J. Smith (Aug. 3, 2007). "Cross-regulation and Functional Redundancy between the Splicing Regulator PTB and Its Paralogs nPTB and ROD1". In: *Molecular Cell* 27.3, pp. 420–434. ISSN: 1097-2765. DOI: 10.1016/j.molcel.2007.06.016.

Stephens, Matthew (Apr. 1, 2017). "False discovery rates: a new deal". In: *Biostatistics* 18.2, pp. 275–294. ISSN: 1465-4644. DOI: 10.1093/biostatistics/kxw041.

Ule, Jernej et al. (Nov. 30, 2006). "An RNA map predicting Nova-dependent splicing regulation". In: *Nature* 444.7119, pp. 580–586. ISSN: 1476-4687. DOI: 10.1038/nature05304.

Van Nostrand, Eric L. et al. (July 2020). "A large-scale binding and functional map of human RNA-binding proteins". In: *Nature* 583.7818. Publisher: Nature Publishing Group, pp. 711–719. ISSN: 1476-4687. DOI: 10.1038/s41586-020-2077-3.

Vehtari, Aki, Andrew Gelman, et al. (June 1, 2021). "Rank-normalization, folding, and localization: An improved $\widehat{R}$ for assessing convergence of MCMC". In: *Bayesian Analysis* 16.2. ISSN: 1936-0975. DOI: 10.1214/20-BA1221. arXiv: 1903.08008[stat].

Vehtari, Aki, Daniel Simpson, et al. (Mar. 13, 2024). *Pareto Smoothed Importance Sampling.* DOI: 10.48550/arXiv.1507.02646. arXiv: 1507.02646[stat].

Wang, Zefeng and Christopher B. Burge (May 2008). "Splicing regulation: from a parts list of regulatory elements to an integrated splicing code". In: *RNA (New York, N.Y.)* 14.5, pp. 802–813. ISSN: 1469-9001. DOI: 10.1261/rna.876308.

Weyn-Vanhentenryck, Sebastien M. et al. (Mar. 27, 2014). "HITS-CLIP and Integrative Modeling Define the Rbfox Splicing-Regulatory Network Linked to Brain Development and Autism". In: *Cell Reports* 6.6, pp. 1139–1152. ISSN: 2211-1247. DOI: 10.1016/j.celrep.2014.02.005.

Yee, Brian A. et al. (Feb. 2019). "RBP-Maps enables robust generation of splicing regulatory maps". In: *RNA* 25.2, pp. 193–204. ISSN: 1355-8382. DOI: 10.1261/rna.069237.118.

Youn, Yong Ha et al. (Dec. 9, 2009). "Distinct Dose-Dependent Cortical Neuronal Migration and Neurite Extension Defects in Lis1 and Ndel1 Mutant Mice". In: *The Journal of Neuroscience* 29.49, pp. 15520–15530. ISSN: 0270-6474. DOI: 10.1523/JNEUROSCI.4630-09.2009.

Zamore, P. D. and M. R. Green (Jan. 1991). "Biochemical characterization of U2 snRNP auxiliary factor: an essential pre-mRNA splicing factor with a novel intranuclear distribution". In: *The EMBO journal* 10.1, pp. 207–214. ISSN: 0261-4189. DOI: 10.1002/j.1460-2075.1991.tb07937.x.

Zarnack, Kathi et al. (Jan. 31, 2013). "Direct Competition between hnRNP C and U2AF65 Protects the Transcriptome from the Exonization of Alu Elements". In: *Cell* 152.3, pp. 453–466. ISSN: 0092-8674. DOI: 10.1016/j.cell.2012.12.023.

Zhang, Chaolin et al. (July 23, 2010). "Integrative modeling defines the Nova splicing-regulatory network and its combinatorial controls". In: *Science (New York, N.Y.)* 329.5990, pp. 439–443. ISSN: 0036-8075. DOI: 10.1126/science.1191150.

Zhang, Xiaochang et al. (Aug. 25, 2016). "Cell-Type-Specific Alternative Splicing Governs Cell Fate in the Developing Cerebral Cortex". In: *Cell* 166.5, 1147–1162.e15. ISSN: 0092-8674. DOI: 10.1016/j.cell.2016.07.025.

Zhuang, Yuan and Alan M. Weiner (1986). "A compensatory base change in U1 snRNA suppresses a 5 splice site mutation". In: *Cell* 46.6. Publisher: Elsevier, pp. 827–835.