Neural and sensory basis of homing behavior in the invasive cane toad, *Rhinella marina*

Daniel A. Shaykevich[1]*, Daniela Pareja-Mejía[1,2], Chloe Golde[1], Andrius Pašukonis[3], and Lauren A. O'Connell[1,4]*

1 Department of Biology, Stanford University, Stanford, CA, USA
2 Graduate Program in Zoology, Universidade Estadual de Santa Cruz, Bahía, Brazil
3 Institute of Biosciences, Vilnius University, Vilnius, Lithuania
4 Wu Tsai Institute for Neuroscience, Stanford University, Stanford CA, USA

Keywords: spatial cognition, navigation, amphibian, medial pallium, olfaction, magnetoreception

*To whom correspondence should be addressed:
Daniel Shaykevich
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA 94305
shaykeda@stanford.edu

Lauren O'Connell
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA 94305
loconnel@stanford.edu

Word count: Abstract: 198, Main text: 9176

Abstract

The behavioral, sensory, and neural bases of vertebrate navigation are primarily described in mammals and birds. However, we know much less about navigational abilities and mechanisms of vertebrates that move on smaller scales, such as amphibians. To address this knowledge gap, we conducted an extensive field study on navigation in the cane toad, *Rhinella marina*. First, we performed a translocation experiment to describe how invasive toads in Hawaiʻi navigate home following displacements of up to one kilometer. Next, we tested the effect of olfactory and magnetosensory manipulations on homing, as these senses are most commonly associated with amphibian navigation. We found that neither ablation alone prevents homing, suggesting that toad navigation is multimodal. Finally, we tested the hypothesis that the medial pallium, the amphibian homolog to the hippocampus, is involved in homing. By comparing neural activity across homing and non-homing toads, we found evidence supporting the involvement of the medial pallium, lateral pallium, and septum in navigation, suggesting a conservation of neural structures supporting navigation across vertebrates. Our study lays the foundation to understand the behavioral, sensory, and neural bases of navigation in amphibians and to further characterize the evolution of behavior and neural structures in vertebrates.

Introduction

The ability to process spatial information is fundamental to survival for many animals, who must navigate through their environment to locate shelter, forage, care for young, and more. Much of our understanding of spatial navigation comes from research in mammals, where the hippocampus has been recognized as the neural center for spatial representations [1–3], alongside other regions like the and entorhinal cortex [4,5]. Recent efforts have characterized similar functions of homologous hippocampal brain regions in birds [6,7], fish [8,9] and reptiles [10], suggesting conservation of the neural architectures supporting this critical behavior. However, the neural basis of amphibian spatial behavior remains understudied, especially in the context of navigation in complex natural environments. As amphibians represent the major evolutionary transition from aquatic to terrestrial habitats in the vertebrate lineage, filling this gap will provide a more complete understanding of the neural representations of spatial cognition and their evolution across vertebrates.

Research across many amphibian species suggests a wide-spread ability to navigate to fixed sites important for reproduction. For example, many temperate-region amphibians migrate to breeding sites on scales up to several kilometers, such as some salamanders (including *Taricha rivularis* and *Plethodon jordani*)[11–13] and pond breeding toads (*Bufo bufo*, *B. japonicus*) [14,15]. Some neotropical poison frog species, such as *Oophaga sylvatica*, *Dendrobates tinctorius*, and *Allobates femoralis*, move with high spatial precision within a few hundred meters to transport tadpoles to water pools [16–18]. A variety of sensory modalities facilitate these movements, with olfaction [12,14,19] and magnetoreception [20–22] being the most often implicated in migratory amphibians. In one relevant example, Sinsch (1987) demonstrated that the European common toad (*Bufo bufo*) relies on both olfaction and magnetoreception to orient towards its breeding pond [14]. Impressive navigation abilities have also been demonstrated in amphibians that do not have fixed reproduction sites, such as the

cane toad, *Rhinella marina* [23]. However, the sensory modalities used for navigation back to

terrestrial home sites, rather than to water bodies has rarely been investigated. More research is

needed on amphibian navigation behaviors outside of reproduction and what sensory modalities

are important, which will provide a more complete picture of vertebrate perception and sensory

integration, and how these lead to successful homing.

The cane toad is native to Central and South America and was introduced globally,

including in Hawai'i and Australia. In its invasive front in Australia, the cane toad can disperse

and travel long distances [24,25], sometimes moving 200-meters per night to new locations [26],

creating difficulties for species management. However, other than on the Australian invasive

front, cane toads are relatively sedentary, rarely move more than 200 meters, and often return

to the same diurnal shelters [23,27–29]. We previously showed that toads in their native French

Guiana are able to return home accurately from a full kilometer, a distance exceeding their

routine movements [23]. These navigation abilities and the toad's extreme prevalence as a

devastating invasive species make it an ideal model for studying homing behavior and probing

unanswered questions regarding the neural and sensory basis of homing.

In this study, we conduct a field experiment to test cane toad navigational ability,

quantify neural activity associated with navigational outcomes, and test which sensory systems

are required for navigation. We hypothesized that the medial pallium (the amphibian

hippocampal homolog [30–32]) is involved in navigation and predicted that homing toads would

have higher neural activity in this brain region compared to non-homing and non-translocated

toads. We next hypothesized that olfaction and magnetoreception guide homing behavior,

given their functionality in other bufonids [14,15,33], and predicted that olfactory or

magnetoreception ablations would disrupt the homing ability or accuracy. Our study identifies

behavioral characteristics involved in homing, brain regions that are more active during

navigation, and provides future directions on sensory system research in cane toad navigation.

Materials and Methods

(a)   Study sites

Experiments occurred at four sites on the island of Oʻahu in Hawaiʻi. We sampled at two sites on the southern, leeward side of the island (West Loch Golf Course: 21° 22' 04.7" N 158° 01' 31.5" W; Hawaiʻi Prince Golf Course: 21° 18' 41.5" N 158° 00' 32.7" W) and two sites on the northern, windward side of the island (Waimea Valley: 21° 38' 04.4" N 158° 03' 14.6" W; Turtle Bay Golf Course: 21° 42' 07.5" N 157° 59' 44.6" W). Waimea Valley was a forested area while the other three sites were golf courses (Figure 1B).

(b)   Animals

Toads were located visually at night (sunset was approximately 19:00 and sunrise approximately 06:30), as cane toads are nocturnal and generally do not exhibit diurnal movements. We photographed the dorsum for re-identification purposes, measured snout-vent-length (SVL) with a tape measure, and measured mass with a portable scale (KUBEI-790, KUBEI, China). Eighty-two toads (40 males, 42 females) were tagged and tracked for at least three days between 18 February and 12 May 2022 across the four sites. Their mean mass was 223.8 g (sd +/- 98.6) and mean SVL was 12.4 cm (+/- 1.6). Toad size was not evenly distributed between sites (Mass: Kruskal-Wallis, $\chi^2(3)$= 19.687, p = 0.0002; SVL: ANOVA, df = 3, F = 14.67, p = 1.15e-07). Sex was determined by size and male release call, and confirmed by gonad identification upon dissection. Females were significantly larger than males (Mass: Wilcoxon, W = 1315.5, p = 1.05e-05; SVL: t-test, t = 4.44, df = 77.41, p = 2.99e-05). Unless otherwise stated, "size" refers to SVL hereafter.

(c) Tagging and baseline tracking

Radio tagging and subsequent tracking was executed similarly to the method we previously described [23]. Briefly, radio tags (BD2, Holohil Systems Ltd, Carp, ON, Canada) were attached to the toad's waists using belts made of silicone tubing and thread (Figure 1A). The tagging process lasted 14.2 min on average (+/- 7.8) after which toads were released. A GPS point was taken in the ArcGIS Field Maps app (ESRI, Redlands, CA, USA) using a Bad Elf GPS Pro+ gps device (Bad Elf, West Hartford, CT USA) connected by Bluetooth to an iPhone (Apple, Cupertino, CA, USA). Following release, toads were located once in the evening prior to their active, nocturnal period and at least two times each night (between sunset and sunrise). Toads were tracked for natural baseline movement information (termed 'baseline' hereafter) between three and 28 days (5.701 +/- 4.04 days) prior to translocation or euthanasia.

(d) Translocation

Translocations occurred during the night generally between 20:00 and 22:00. Sixty-two toads were translocated. Toads were caught and their waists examined to ensure no abrasion from the waist belt. Toads were placed into an enclosed net cage and walked in an indirect route for 30-60 minutes to a distance approximately 500- or 1000-meters from the baseline position. At the release site, the distance from the toad's nearest baseline point was confirmed on a GPS-connected handheld device, the net cage was left at the release point for 5 minutes, and the toad was gently released from the cage and allowed to move freely.

Following release, toads were relocated approximately once per hour during the nighttime, beginning at 20:00 and continuing until tagged animals had stopped moving (06:40 at the latest). On subsequent days, toads were relocated once in the early evening to determine location before nocturnal movements began. If a toad was not active at night (e.g. not moving or making small, multidirectional movements), the frequency of observations was reduced to every 2-3 hours while tracking moving toads.

Tracking ended once the animal reached a point within 200-meters of one of its recorded

baseline positions or 72 hours had passed after translocation. Homing toads (captured within

200-meters of a baseline point) were euthanized and tissues were collected for downstream

analyses (described below) within 30 minutes of crossing the 200-meter threshold. Non-homing

toads were sacrificed after 20:00 wherever they were at that time. Two toads in the homing

group were unintentionally caught before they crossed the 200-meter threshold (B8 at 233-m

and D5 at 218-m) and two more toads were caught and euthanized approximately 300-m from

home after encountering physical obstacles following large homeward movements (C8 and F5) ,

but were included in the homing condition (both for considering behavior and neuronal activity).

One toad moved outside of the study area and was euthanized because it became impossible

to track; its behavior and brain data were not included in analysis.

158   (e)   Sensory Ablations

A subset of translocated toads (n = 43) received magnetoreception or olfactory ablations

to test the roles of these sensory systems in homing. In each group, we had experimental (i.e.

ablated) animals, as well as control animals receiving sham ablation treatments. Toads were

tracked for at least three days prior to ablation, and then were tracked for at least 48 hours post-

ablation before translocation to evaluate any effects of the ablation procedures on space use.

In the olfactory ablation group (n = 11), toads were captured and lightly anesthetized

through immersion in an MS-222 bath (1.4 g/L) for ~5 min. We performed olfactory ablations

through zinc sulfate ($ZnSO_4$) treatment, which temporarily damages the olfactory epithelium in

juvenile and adult amphibians [34,35] and other vertebrates [36,37]. Once animals were

anesthetized and not responding to toe pinches, they were held upside-down and each nostril

was washed with 1% $ZnSO_4$ dissolved in amphibian ringer solution (NaCl, KCl, $CaCl_2 \cdot 6H_2O$ and

distilled water; Electron Microscopy Services, Hatfield, PA 19440) applied in five 200 µl doses

through the internal nare with a P200 pipette. Any $ZnSO_4$ that spilled into the animal's mouth

was sucked up with the pipette. Following administration of $ZnSO_4$, the toad's mouth was

washed several times with amphibian ringer solution. Animals recovered in a net cage until fully

mobile (usually 20-30 minutes), after which they were released at the capture point. Animals

experiencing sham ablations (n = 11) underwent an identical procedure, but their nostrils were

washed with amphibian ringer solution without $ZnSO_4$.

Magnetoreception ablations were carried out by attaching a small Nickel plated

Neodymium magnet (N50, 10 mm x 6 mm x 2 mm, ~1 g; Super Magnet Man, Pelham, Alabama,

USA) with cyanoacrylate based adhesive (EM-150, Starbond, Torrance, CA, USA) to the skin

over the skulls of toads (n = 11), similarly to previous studies in anurans [14,33,38]. Toads were

captured, magnets applied, and toads were kept in a net cage for ~30 minutes before release.

Sham ablation animals had a brass rectangle (12.7 mm x 6 mm x 1.6 mm, ~ 1 g; Metals Depot,

Winchester, Kentucky, USA) attached in place of a magnet (n = 10). Two animals lost their

magnets following translocation, so their behavior and brain measurements were not included in

the magnetoreception analysis.

(f) Tissue collection

Toads were euthanized at one of three timepoints: (a) the completion of at least three

days of baseline tracking if the individual was being used as a baseline comparison for non-

homing and homing translocated toads (baseline, n = 12), (b) three days post-translocation if

the toad failed to return within 200-meters of its baseline location (non-homing, n = 27), (c) or

once the toad reached a point 200-meters from a previously observed baseline location

(homing, n = 33).

Toads were euthanized through MS-222 injection (0.7% solution, 300 mg/kg body

weight) followed by cervical transection with sharpened hedge shears. Brains were immediately

dissected, placed in 1 mL of 4% paraformaldehyde (PFA), and kept on ice until they could be

placed into a 4℃ refrigerator. Brains were fixed overnight in PFA, washed three times in 1x PBS

and then placed into 1x PBS with 0.02% sodium azide, or 1x PBS if transportation to the

laboratory site would happen in two days. Brains were kept in 1x PBS with sodium azide < 1

week and two days prior to travel to the laboratory, the brains were transferred to 1 mL of 30%

sucrose dissolved in 1x PBS for cryoprotection. Once the brains sank in sucrose, they were

transported on ice to a laboratory at the University of Hawai'i at Manoa, where they were

embedded in Tissue-Tek OCT Compound (Electron Microscopy Sciences) and frozen on dry ice

before being stored at -80℃. We collected 73 brains (61 translocated and 12 baseline). One

homing toad lost its tag within 200-meters of its home area, so its brain was not collected.

(g)   Immunohistochemistry

Brains were sectioned on a cryostat in 20 μm slices and thaw-mounted onto slides in

four series. Slides dried at room temperature for ~48 hours before being transferred to -80℃ for

storage prior to staining. We used immunohistochemistry to detect Phospho-S6 Ribosomal

Protein (pS6) as a proxy for neural activity [39] as previously described by our laboratory

[40,41], except with a 1:300 primary antibody dilution (full protocol is presented in SM1). pS6 is

a reliable activity marker in amphibians [42] and other animals [39]. Of the 73 collected brains,

two were discarded due to poor tissue quality and four were discarded for being collected at the

wrong time point, leaving 67 for analysis.

(h)   Microscopy and Cell Counting

Brains were imaged on a Leica DM6B microscope with brightfield capability at 20x

magnification. One hemisphere was selected to image for each brain. Nine sections along the

AP axis were chosen based on anatomical landmarks: three from the anterior of the

telencephalon (posterior to the olfactory bulb), three from the medial portion, and three from the

posterior of the telencephalon (anterior to the ventral hypothalamus) (Figure S1). ImageJ (NIH,

Bethesda, MD, USA) was used to count pS6-positive cells in each of five brain regions: medial

pallium (Mp), lateral pallium (Lp), dorsal pallium (Dp), septum (Sep) and the striatum (Stri) given

their role in spatial behavior in toads [43–45] and other vertebrates [3,46,47]. Delineation of

brain regions was based on Sotelo et al. (2016), which defined pallial regions by Westhoff and

Roth (2002) and ventral regions by Moreno and González (2004)[30,43,48]. ImageJ was used

to calculate the surface area of each region. Densities of pS6-positive neurons for each region

were calculated by dividing the number of pS6-positive cells by the area.

For toads in the olfaction ablation experiment, we used ImageJ to count active cells in a

300 μm by 300 μm box positioned at the lateral edge of the granule cell layer in the olfactory

bulb (Figure S2) to measure the effect of the olfactory ablation in comparison to the sham

ablation.

(i) Data analysis

(i) Toad movement data

GPS points were visualized in ArcGIS online and ArcGIS Pro 2 (ESRI). Spatial and

temporal attributes were calculated using the package 'adehabitatLT' [49] in R Studio (v

2023.09.1+494, Posit Software, PBC, Sunnyvale, CA) running R (v 4.3.1, R Foundation for

Statistical Computing, Vienna, Austria). We calculated several descriptors of toad movement

and homing: cumulative movement, daily movement, movement range, straightness index,

homing duration/speed, and active homing duration/speed (details of descriptor calculations in

SM2).

To estimate the similarity of a toad's home range before and after ablation treatment, we

plotted minimum convex polygons in ArcGIS Pro encapsulating all location points pre-ablation

and calculated the proportion of post-ablation relocations that fell inside the polygon.

Outliers were values more than three standard deviations from the mean and were

removed prior to statistical comparisons and visualization. Normality was verified using the

Shapiro-Wilks Test. Toad size (SVL) was compared between sexes using a Student's t-test and

between field sites by ANOVA with a post-hoc Tukey's test. Comparisons of mass, movement

range, and homing duration/speed between field sites were done with Kruskal-Wallis tests with

a post-hoc Dunn's test. Wilcoxon Rank Sum Tests were used to compare mass and movement

range between sexes. It was also used to compare the effects of translocation distance on

homing descriptors (straightness, duration, speed, and time to move 100 meters homeward), as

well as differences between sham and ablated individuals in home area pre/post manipulation,

homing duration, and cumulative movement. A Paired Wilcoxon test was used to compare

movement before and after ablations for sensory manipulated toads. Pearson's Chi-Squared

tests were used to compare binary homing success between translocation distances and

sham/ablated individuals. The effect of toad size on homing duration was modeled with a linear

model using the 'lm' function in the 'stats' R package [50] and with a linear mixed-effects model

using the 'lme4' R package [51], where duration was the response variable, size was the fixed

effect, and field site was the random effect.

We used the 'circular' package in R [52] to analyze the directionality of movement

angles between homing and non-homing toads. The "ltraj" objects generated using the

'adehabitatLT' package provide absolute angles (i.e. compass direction) and relative angles (i.e.

the change in angle from the previous movement) for each movement segment. In our

comparisons, we calculated the circular mean of all absolute angles of movement for each toad

and normalized with respect to the home direction. We used the Rayleigh test of circular

uniformity to test whether mean absolute angles differed from the home direction in homing and

non-homing toads. We also compared the distribution of mean angles directly between homing

and non-homing animals using a MANOVA based on trigonometric functions [53] using

'manova' functions in the 'stats' R package [50].

(ii) Cell Count Data

We used generalized linear mixed models ('glmmTMB' in R [54]) to test for differences in

brain activity and its relation with behavior. Models used a negative binomial distribution and

best model fit was confirmed using the 'DHARma' package in R [55]. Homing condition, brain

region, and their interaction were the main effects predicting the number of pS6-positive cells.

Toad ID was included as a random factor and brain region area was included as an offset

variable to account for brain area size. We modeled differences in brain activity due to ablations

separately using the same model parameters as above except that ablation was also included

as a main effect. Post hoc pairwise contrasts (between homing condition groups and ablation

treatments) were calculated with estimated marginal means using the "emmeans" package in R

[52].

To test if brain activity was related to either straightness or duration of homing, we

performed linear mixed effects models using the "lme4" package, where pS6 count was the

response variable and the homing descriptor (straightness or duration), brain region, and their

interaction were fixed effects. Toad ID was a random effect and brain region area was an offset

variable in the model. Linear mixed effects models were also performed within count data

specific to each brain region, in which the response variable was pS6 counts within a given

region, the homing descriptor was the fixed effect, Toad ID was a random effect, and region

area was an offset variable.

Results

Toads generally exhibit site fidelity

Prior to translocation, toads exhibited a mean movement range of 130.19 m (s.d. 138.75

m), mean cumulative movement of 539.12 m (s.d. 745.45), and mean daily movement of 110.48

m (s.d. 157.20). Movement range varied significantly between sites (Kruskal-Wallis, $\chi^2(3)$ =

13.82, p = 0.0032), with toads at the Waimea Valley forest site exhibiting smaller ranges than

toads at golf courses (Figure 1B). There was no difference in movement range between males

and females (Wilcoxon, W = 783, p = 0.96). All three statistical outliers were males. Two of

these individuals moved between two water sources more than 500-meters apart (Figure S3),

apparently for mating, as we observed mating calls from the destination ponds. Size did not

have a significant effect on range.

Figure 1. Movement range varies across locations. (A) Observed movement range of cane
toads (in meters) at each tracking site. Letters represent significant differences between sites
(Kruskal-Wallis, χ²(3) = 13.82, p = 0.0032). Shapes represent sex (▲ = male, ⬤ = female). Inset
shows toad with radio tag. Letters represent significant differences from post-hoc Dunn's Test
(with Benjamini-Hochberg adjustment) **(B)** Photographs depicting typical landscapes at each
tracking site.

Toads exhibited long-distance homing

Out of 62 translocated tods, 34 exhibited homing behavior (Figure 2A-D). Toads

returned home in 3.27 to 77.48 hours (25.50 +/- 22.23) with a mean straightness of 0.81 (+/-

0.16). Considering all toads (including those in ablation experiments), 19 out of 31 toads

returned home from 500-meters and 15 out of 30 toads returned from 1000-meters. There was

no effect of translocation distance on homing success (Pearson's Chi-squared, $\chi^2_{(1)}$ = 0.40, p =

0.53). Distance had no effect on homing duration (Wilcoxon, W = 147.5 , p = 0.99) (Figure S4A)

and toads translocated 1000-meters had a faster homing speed (Wilcoxon, W = 72, p = 0.014;

Figure S4B). A linear mixed-effects model accounting for the unequal distribution of size across

field sites showed no significant effect of size on homing duration (t = -0.51, df = 30.52, p =

0.61; Figure S5).

Homing speed and duration varied significantly between sites, with toads at Waimea

Valley homing slower than toads at Hawaii Prince and West Loch (Speed: Kruskal-Wallis, χ^2(3)=

10.52, p = 0.015; Duration: Kruskal-Wallis, χ^2(3)= 12.38, p = 0.0062; Figure 2E). Toads often

did not immediately move homeward, so we measured an "active homing speed" using only

movement and time after the first 100-meters of homeward movement. Active homing speed

and duration did not vary between sites (Speed: Kruskal-Wallis, χ^2(3) = 5.37, p = 0.15; Duration:

Kruskal-Wallis, χ^2(3)= 0.66, p = 0.88), suggesting that the main differences in homing time are

due to stationary periods prior to homeward movements (Figure 2F). On average, the time to

move at least 100-meters homeward was 64% of the total homing duration.

Figure 2 Toads exhibit long-distance homing. Normalized trajectories of toads translocated **(A)** 500-m and **(B)** 1000-m. Plot centers represent translocation release points and the gray circles at the top represent the 200-m buffer threshold around baseline points. 19/31 toads returned from 500-m and 15/30 from 1000-m. Distance to home over time for toads translocated **(C)** 500-m and **(D)** 1000-m. Gray rectangle at the bottom of the plot represents the 200-m buffer threshold around baseline points **(E)** Duration of homing differs between the experimental sites (Kruskal-Wallis, $\chi^2(3)$= 12.38, p = 0.0062) Letters represent significant differences from post-hoc Dunn's Test (with Benjamini-Hochberg adjustment). **(F)** Active homing time shows no difference between locations (Kruskal-Wallis, $\chi^2(3)$= 0.66, p = 0.88).

Twenty-seven of the 62 translocated toads did not exhibit homing behavior. Cumulative

movement of non-homing toads averaged 547.07 m (+/- 488.5) and the mean minimum

distance from home reached was 719.3 m (+/- 359.5). Mean movement angles of homing toads

showed a significant orientation towards the home direction (Rayleigh p = 0.024) and non-

homing toads did not (Rayleigh p = 0.83). However, there was only a marginal difference in

distribution of angles of movement between homing and non-homing toads (MANOVA, Pilliai =

0.092, F(2,58) = 2.95, p = 0.06; Figure S6).

Pallial and septal brain regions show increased neural activity in homing toads

Given the variability in homing performance, we next asked whether neural activity in

brain regions is correlated with homing success by quantifying the number of pS6-positive cells

in five brain regions (Figure 3, Table S1). We first asked whether toads, regardless of sensory

treatment, showed differences in brain activity due to homing behavior. We found a significant

effect of homing behavior on neural activity in a brain region specific manner (brain region x

homing condition: $\chi^2(8)$ = 103.742, p < 2.2e-16) (Table 1). In the medial pallium and the

septum, we found that homing animals had more pS6-positive neurons than both non-homing

and baseline toads (Mp: Homing-Baseline- z = -3.65, p = 0.0008; Homing-Non-homing- z =

4.18, p = 0.0001. Sep: Homing-Baseline - z = -3.71, p = 0.0006; Homing-Non-homing - z = 4.25,

p = 0.0001). In the lateral pallium, we also found differences in the number of pS6-positive

neurons between Homing and Non-homing toads (z = 2.77, p = 0.015) and marginal differences

between Homing and Baseline toads (z = -2.34, p = 0.051). Neither the dorsal pallium nor the

striatum showed significant differences in the number of pS6-positive cells associated with

homing and there were no differences between Non-homing and Baseline toads in any brain

region.

We next asked if neural activity was correlated with path straightness and homing

duration. There were no differences in brain activity across brain regions due to path

straightness (Tables S2 and S3). However, the medial pallium (p = 0.0098) and septum (p =

0.0034) showed significant differences in activity related to duration of homing compared to

other brain regions (Table S4) and duration showed a significant effect of homing on septal

neuronal activity (t =-2.59, df = 27.02, p = 0.015) (Table S5).

Figure 3. Pallial regions and the septum are more active due to homing behavior. (A)
pS6-positive cells per mm^2 averaged per individual for each brain region across homing
conditions. Brain activity within brain regions varied significantly due to homing conditions (χ^2(8)
= 103.742, p < 2.2e-16). Letters near box plots represent significant pairwise differences within
brain regions based on a mixed model with cell count as a response factor. Abbreviations: Dp,
dorsal pallium; Lp, lateral pallium; Mp, medial pallium; Sep, septum; Stri, striatum. **(B)**
Representative examples of stained Mp brain tissue from each homing group. Scale bars
represent 50 µm. In inset, arrows indicate pS6-positive cells.

We next restricted our analysis to toads without sensory system manipulations (Figure

S7, Tables S6 and S7) and found that differences in pS6-positive cells between Homing and

Baseline individuals in the medial and lateral pallium still hold (Mp: Homing-Baseline- $z = -$

3.553, $p = 0.0011$; Lp: Homing-Baseline- $z = -3.019$, $p = 0.0072$). The differences in the

number of pS6-positive cells between Homing and Baseline and Non-homing animals in the

septum were also consistent (Sep: Homing-Baseline- $z = -3.508$, $p = 0.0013$; Homing-Non-

homing- $z = 3.005$, $p = 0.0075$). Similar to the full sample, there were no differences between

groups in the dorsal pallium and striatum. In contrast to the full dataset, there were no

differences between Homing and Non-homing animals in the medial and lateral pallium. The

differences in the results are likely due to sample size, as the non-ablated subset includes only

28 out of 67 animals (11 Baseline, 10 Homing, and 7 Non-homing).

Table 1. Summary of main statistical effects from generalized linear mixed model for
differences in brain activity

	X^2	df	p
Homing Condition	13.075	2	0.001448
Brain Region	2529.991	4	< 2.2e-16
Homing Condition x Brain Region	103.742	8	<2.2e-16

Olfactory ablation influences the homing path

We next asked whether ablation of olfaction led to a disruption in spatial behavior and

navigation, as has been demonstrated in other amphibians [13–15]. There was no significant

difference in movement range pre- and post-ablation for sham (Paired-Wilcoxon, V = 38, p =

0.70) or ablated animals (Paired Wilcoxon V = 36, p = 0.83). However, when comparing the

space occupied pre- and post-ablation, ablated toads showed greater divergence from their pre-

ablation area than sham animals (Wilcoxon, W = 340, p = 0.021), suggesting that loss of

olfaction might disrupt routine movements and space use. There were no significant differences

of homing success ($\chi^2(1) = 0.73$, $p = 0.39$) or duration (Wilcoxon, $W = 23$, $p = 0.42$) between

sham and ablated groups (ablated homing success = 45.5%, sham homing success

63.6%)(Figure 4A). After translocation, ablated toads moved more than sham toads when

considering Homing and Non-homing toads together (Wilcoxon, $W = 93$, $p = 0.034$) (Figure 4B).

Though the sample is too small for statistical comparison, ablated toads that homed showed

more cumulative movement (831.90 m +/- 288.07) than sham toads that homed (629.79 m +/-

334.19).

We next tested whether olfactory ablations influenced brain activity during homing.

Ablated toads had significantly less pS6-positive cells in their olfactory bulbs than sham toads,

confirming reduced olfactory function from chemical ablation (Wilcoxon, $W = 93.5$, $p =$

0.0074)(Figure 4C). While there was a significant interaction effect of ablation and brain region

on pS6 count ($\chi^2(4) = 19.23$, $p = 0.0071$)(Table S8), the magnitude of the effect was insufficient

to detect significant pairwise differences in region-specific activity between ablated and sham

groups (Table S9, Figure S8).

410
411
Figure 4 Olfactory ablation did not prevent homing but may affect trajectories.
(A) Trajectories of toads in the olfactory ablation experiment translocated (i) 500-meters and (ii)
1000-meters. **(B)** Ablated toads exhibited higher cumulative movement post- translocation than
sham toads (Wilcoxon, W = 93, p = 0.036). **(C)** Sham toads exhibited more pS6-positive cells in
the olfactory bulb than ablated animals (Wilcoxon, W = 93.5, p = 0.0074). Each point represents
the mean pS6 count across three sections of olfactory bulb in the same toad. **(D)**
Representative areas of olfactory bulb (i) sham and (ii) ablated animals. Scale bars represent 25
μm. In inset, arrows indicate pS6-positive cells.

Magnetoreception ablations do not affect homing

420    Magnetoreception is a sensory modality linked to homing in several amphibians, so we

421 tested the hypothesis that cane toads also use magnetoreception for homing. Movement range

422 was significantly lower in both sham (Paired Wilcoxon, $V = 56$, $p = 0.042$) and ablated (Paired

423 Wilcoxon, $V = 62$, $p = 0.0068$) animals (Figure 5C). There was no significant difference when

424 comparing similarity in positions pre- and post-ablation (Wilcoxon, $W = 236$, $p = 0.90$). There

425 were no significant differences of homing success ($\chi^2(1) = 1.27$, $p = 0.26$) or duration (Wilcoxon,

426 $W = 5$, $p = 0.55$) between sham and ablated groups (ablated homing success = 55.6%, sham

427 homing success = 30.0%) (Figure 5A). There was no difference in cumulative movement post-

428 translocation between sham and ablated animals (Wilcoxon, $W = 55$, $p = 0.44$)(Figure 5B).

429 While there was a significant interaction effect of ablation and brain region on pS6 count ($\chi^2(4) =$

430 14.22 p = 0.0066) (Table S10), the effect was insufficient to detect significant pairwise

431 differences in activity in any region between ablated and sham groups (Figure S9, Table S11).

Figure 5 Magnetoreception ablation did not affect homing. (A) Trajectories of toads in the
magnetoreception ablation experiment translocated (i) 500 meters and (ii) 1000 meters. **(B)**
There was no difference in cumulative movement between sham and ablated toads. **(C)**
Movement range varied significantly between pre- and post- ablation periods in both sham
(Paired Wilcoxon, V = 56, p = 0.042) and ablated (Paired Wilcoxon, V = 62, p = 0.0068) toads.
(D) Example of a toad with a magnet.

Discussion

In the current study, we examined the neural correlates and sensory basis of toad

homing behavior. On a behavioral level, our results demonstrate an impressive map-like

navigation ability in cane toads that did not lessen with increasing distance. On the neural level,

we found that areas associated with navigation in mammals and birds, such as medial pallium

and the septum, show increased activity in homing toads. Contrary to our predictions, we found

that neither olfactory nor magnetic ablation alone disrupted homing, suggesting that this

behavior is multimodal or based on other sensory modalities. These results contribute to filling

in gaps in the understanding of amphibian spatial behaviors and neural architecture of

vertebrate spatial cognition.

(a) Homing in toads

We found that toads displayed robust navigation from one kilometer. We observed that

toads would often remain relatively stationary near their release site for some time before

moving home. In other studies, direct homing following a stationary period has been observed in

amphibians, including dendrobatid poison frogs [17,46], where homing abilities have been

assumed to be adaptations to the spatial challenges of parental care. However, the homing

abilities of cane toads suggest that refined navigation may not be reserved for species with

specialized reproductive behaviors but is instead more widely present across amphibians. While

our translocations exceeded the scale of routine movements of most toads, observations of

occasional longer distance movements further suggest the presence of unexplored spatial

abilities. Toad homing is even more impressive when considering their habitat and type of

locomotion. Well-studied species capable of long distance navigation, such as pigeons, bats,

and sea turtles, move through three-dimensions [57–59], allowing them to travel efficiently and

overcome obstacles. That toads, confined to the ground and moving through obstructed

landscapes, including water-crossing, demonstrate direct navigation on the scale of a kilometer

reveals a remarkable capacity for spatial accuracy.

(b) Conserved role of pallial brain regions in spatial tasks

Our study contributes to the accumulating evidence that medial pallium is functionally

homologous to the mammalian hippocampus. The hippocampus has long been studied as a

center of navigation-related activity, as it holds neurons encoding place information [1,3] and is

required in the consolidation of spatial memory [2]. In birds, hippocampal lesions impair homing

in pigeons [57] and caching behavior is associated with larger hippocampal regions [60]. Fish

and turtles require homologous regions to solve a spatial task [10]. In amphibians, the medial

pallium has been identified as a region with similar developmental origins [31], but different

topography compared to amniotes [32]. Numerous neuronal activation and lesion studies in *R.*

arenarum suggest that the medial pallium is used in spatial tasks involving environmental

geometry [43] and feature cues [44]. The medial pallium is also shown to have increased activity

in tadpole-transporting poison frogs, a parental care task that requires high spatial precision

[41]. Our work corroborates this accumulating evidence that the medial pallium is involved in the

regulation of spatial behaviors in amphibians.

In our experiment, the lateral pallium showed increased activity in homing animals, while

the dorsal pallium did not. The role of these brain regions is not well studied in amphibians, but

both regions were shown to have increases in activity in a geometry based task in *R. arenarum*

[43]. Specifically, the lateral pallium showed increased activation when a visual cue was present

in the maze [43]. Anatomically, the pallial regions are interconnected, where the dorsal pallium

receives input from the lateral pallium and sends input to both the lateral and medial palliums,

and the lateral pallium connects to the medial pallium through the dorsal association bundle

[31]. Molecular profiling in salamanders suggests that the dorsal pallium resembles the

entorhinal cortex, another mammalian region involved in spatial coding [61]. The lateral pallium

also receives olfactory projections from the lateral olfactory tract [31], but we saw no evidence of

ablation-related effects. Our experiment suggests distinct functionality may be expected from

these regions in relation to navigation and more work is needed to understand specific functions

associated with each area.

Our results showed increased activity in the septum in homing animals and increasing

activity with faster homing. The medial septum has been implicated as an important regulator of

spatial tasks in rodents, integrating the function of multiple circuits important to path integration

[62]. In mammals, the lateral septum is also innervated by the hippocampus and holds neurons

that exhibit reward-related place fields [63,64]. The role of the septum in navigation has not

been as well studied outside of mammals. In amphibians, dendrites from the region connect

with the medial pallium [31]. In *R. arenarum,* the septum showed no difference in activity in

various conditions in a geometry based arena [35], but the medial septum did show heightened

activity when animals located goal locations in relation to mating calls [45]. Further study in

different navigation scenarios will be necessary to determine the septum's spatial function.

The striatum did not exhibit a difference based on homing conditions in our study. In

mammals, the dorsal striatum has been implicated in memory and may be an integral part of

neural circuitry that allows for the integration of sensory information in goal-directed spatial

tasks [46]. For example, rats with dorsal striatum ablations could not perform a "reinforced

stimulus response task" that required association of a maze arm food with light cues [65].

Interestingly, *R. arenarum* showed no differential activity of the striatum when completing a

maze with a visual cue [43]. Future investigations of differences in region-associated spatial

functions in amphibians and other vertebrates may pinpoint evolutionary changes in neural

architecture.

One important caveat of our experiment is the relationship of the behavior and the timing

of the endpoint. Homing animals were often stationary before moving back in a straight line,

suggesting that the animals gain their bearings during this time. The snapshot of brain activity

captured by our pS6 immunohistochemistry does not account for this period, but for brain

activity happening during active homing. In future experiments, it would be useful to compare

this brain activity to that of the pre-homing period. Furthermore, we cannot conclude brain

regions with higher neural activity are required for homing behavior, as surgeries for brain

region-specific lesions in field conditions are very difficult. Future studies that incorporate

regional ablations or laboratory electrophysiology experiments would elucidate the specific roles

of these regions in amphibian navigation.

522   (c) Use of sensory systems in homing

Olfaction and magnetoreception have been implicated as important senses in navigation

studies in many amphibians [12–15,20,22,33,66] and other vertebrates [67,68]. Ablation

experiments strongly suggested that *B. bufo* relies on these senses to orient towards breeding

ponds [14]*,* while *B. japonicus* seemed to rely solely on olfaction [15]. In a variety of salamander

species (*P. jordani*, *P. cinerus*, and *Notophthalmus virideseens*), both magnetoreception and

olfaction are used to orient and return to breeding streams [12,13,20,21]. However, in our study,

ablations did not disrupt *R. marina* homing. If either is used in cane toad navigation, they likely

work in combination with other senses or can compensate for each other.

Olfaction is used in multiple ways in navigation. Birds use odors distributed by wind to

determine goal locations [67]. Mammals sample odor information through sniffing to gain

information about their environment [68]. For amphibians, cues can come directly from aquatic

navigation goals, as seems to be the case in other toad species [14,15] and poison frogs [19].

Some salamanders may smell pheromones left by themselves or conspecifics [12]. Although we

did not observe a reduced success in homing with olfactory ablations, ablated toads showed

less similarity in space occupied post-ablation and moved more post-translocation than sham

toads. These results suggest that olfactory cues may help toads move in more direct paths and

are important for recognizing home range.

Magnetic orientation has been demonstrated in several amphibians [20,69,70], including

other bufonids [14,71], but we found no evidence that magnetic field disruption impacts long-

distance homing in cane toads. Other vertebrates, such as pigeons, sea turtles, and cetaceans

exhibit impressive, long range navigation using magnetic cues [58,72,73]. However, much of the

functional understanding of magnetoreception comes from salamanders [20,22,69]. Amphibian

magnetoreception can be light dependent [21] and can be used not only as a compass, but also

as a map sense (similarly to sea turtles and birds), even though their movements are on much

smaller scales [74–76]. The effect of magnetoreception ablation on homing in other toad

species is variable: *B. bufo* showed a less accuracy in orientation following translocation [14],

while *B. japonicus* showed no response to ablation [15]. We cannot discount magnetoreception

in *R. marina* on the basis of the current experiment, as the sham group performed poorly, with

only 30% of toads returning home. We used methodology previously deployed in other

amphibians [14,38], and performed our sham ablations by attaching brass bars in the same

manner as the magnets. Both ablated and sham groups moved less following ablation,

suggesting that the ablation potentially had an unintended consequence on behavior, possibly

through physically covering the pineal gland, which contains photoreceptors [77] that may play a

role in magnetic compass orientation.

Other senses may play a role in homing behavior. For example, both male and female

cane toads exhibit phonotaxis to male display calls [78]. However, at our field sites, we

observed toads distributed over several ponds and calling not always occurring when toads

were homing, making the use of auditory cues unlikely. Vision may also play a role and is used

by toads and poison frogs in laboratory arenas [43,79]. However, it is unlikely to be the primary

sense involved in homing, as the visual information toads receive (ground-level, at night) is very

limited in relation to the scale of the translocation distance. In addition, migratory amphibians

with visual ablations are able to return home [13,14]. However, vision may be used to sense

celestial cues, which provide compass information to some amphibians [66,80]. The field of

amphibian navigation would benefit from further study of the functionality of many possible

modalities.

Conclusion

Our work combines intensive field-based telemetry with laboratory analysis of brain

activity to study the neural and sensory basis of navigation. We demonstrate that cane toads (a)

are capable of long-distance navigation to specific sites, (b) exhibit a potentially conserved

function of the pallial and septal regions in relation to spatial behavior, and (c) likely have a

multimodal basis of navigation. In the future, lesion experiments are needed to identify the

specific function of pallium brain regions in navigation. In addition, further lab-based studies

incorporating electrophysiology may characterize how neurons encode spatial information in

amphibians. The current work provides a foundation upon which to begin further examinations

of the neural and sensory basis of amphibian navigation through the integration of field and

laboratory data.

Acknowledgements

The authors respectfully and with reflection offer a Hōʻoia ʻĀina, a Land Acknowledgement,
acknowledging that this work was performed in Hawaiʻi, an indigenous space whose original
people are today identified as Native Hawaiʻians. We also acknowledge that the laboratory
portion of this research was conducted on the ancestral lands of the Muwekma Ohlone people
at Stanford. We understand the implications of the historical and present colonialism the Ohlone
people experience and celebrate their continued stewardship of their lands. We would like to
thank Dr. Amber Wright at the University of Hawaiʻi at Mānoa for providing us with lab spaces
and community. We would like to thank the representatives of our field sites who made our work
possible: George Crisologo (West Loch Golf Course), William Suckoll (Hawaiʻi Prince Golf
Course), Travis Joeger (Turtle Bay Golf Course), and Chad Middleton (Waimea Valley). We
also thank all employees at these locations who monitored our presence and assisted us. We
would like to thank Camilo Rodríguez Lopez for his counseling on statistical matters and Neil
Khosla for reading preliminary drafts of this work.

Data Availability

594   All data is included in supplementary materials.

Author contributions

Conceptualization: D.A.S., A.P., L.A.O; Methodology: D.A.S.; Formal analysis: D.A.S.;

Investigation: D.A.S., D.P.M., C.G.; Resources: L.A.O.; Data curation: D.A.S., D.P.M., C.G;

Writing – original draft: D.A.S.; Writing – review and editing: D.A.S., L.A.O., A.P., D.P.M., C.G.;

Visualization: D.A.S.; Supervision: L.A.O.; Project administration: L.A.O.; Funding acquisition:

D.A.S., L.A.O.

Funding

This work was supported by the National Science Foundation (CAREER award IOS-1845651)

and the National Institutes for Health BRAIN Initiative (R34NS127103-01) to L.A.O.. D.A.S. is

supported by a National Science Foundation Graduate Research Fellowship (2019255752) and

the National Institutes of Health (T32GM007276). LAO is a New York Stem Cell Foundation –

Robertson Investigator. AP is supported by Marius Jakulis Jason Foundation.

Competing interests

The authors declare no competing or financial interests.

Citations

1. O'Keefe J, Dostrovsky J. 1971 The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171–175. (doi:10.1016/0006-8993(71)90358-1)
2. Florian C, Roullet P. 2004 Hippocampal CA3-region is crucial for acquisition and memory consolidation in Morris water maze task in mice. *Behav. Brain Res.* **154**, 365–374. (doi:10.1016/j.bbr.2004.03.003)
3. Humphreys GW, Duncan J, Treisman A, O'Keefe J, Burgess N, Donnett JG, Jeffery KJ, Maguire EA. 1998 Place cells, navigational accuracy, and the human hippocampus. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **353**, 1333–1340. (doi:10.1098/rstb.1998.0287)
4. Fyhn M, Molden S, Witter MP, Moser EI, Moser M-B. 2004 Spatial Representation in the Entorhinal Cortex. *Science* **305**, 1258–1264. (doi:10.1126/science.1099901)
5. Moser EI, Kropff E, Moser M-B. 2008 Place Cells, Grid Cells, and the Brain's Spatial Representation System. *Annu. Rev. Neurosci.* **31**, 69–89. (doi:10.1146/annurev.neuro.31.061307.090723)

6. Sherry DF, Grella SL, Guigueno MF, White DJ, Marrone DF. 2017 Are There Place Cells in
the Avian Hippocampus? *Brain Behav Evol* **90**, 73–80.
7. Payne HL, Lynch GF, Aronov D. 2021 Neural representations of space in the hippocampus
of a food-caching bird. *Science* **373**, 343–348. (doi:10.1126/science.abg2009)
8. Vinepinsky E, Cohen L, Perchik S, Ben-Shahar O, Donchin O, Segev R. 2019
Representation of Borders and Swimming Kinematics in the Brain of Freely-Navigating Fish.
bioRxiv , 291013. (doi:10.1101/291013)
9. Cohen L, Vinepinsky E, Donchin O, Segev R. 2023 Boundary vector cells in the goldfish
central telencephalon encode spatial information. *PLOS Biol.* **21**, e3001747.
(doi:10.1371/journal.pbio.3001747)
10. Rodríguez F, López JC, Vargas JP, Gómez Y, Broglio C, Salas C. 2002 Conservation of
spatial memory function in the pallial forebrain of reptiles and ray-finned fishes. *J. Neurosci.*
Off. J. Soc. Neurosci. **22**, 2894–2903. (doi:20026211)
11. Twitty V, Grant D, Anderson O. 1967 Home Range in Relation to Homing in the Newt
Taricha rivularis (Amphibia: Caudata). *Copeia* **1967**, 649–653. (doi:10.2307/1442244)
12. Jaeger RG, Fortune D, Hill G, Palen A, Risher G. 1993 Salamander Homing Behavior and
Territorial Pheromones: Alternative Hypotheses. *J. Herpetol.* **27**, 236–239.
(doi:10.2307/1564945)
13. Madison DM. 1969 Homing behaviour of the red-cheeked salamander, Plethodon jordani.
Anim. Behav. **17**, 25–39. (doi:10.1016/0003-3472(69)90109-2)
14. Sinsch U. 1987 Orientation behaviour of toads (Bufo bufo) displaced from the breeding site.
J. Comp. Physiol. A **161**, 715–727. (doi:10.1007/BF00605013)
15. Ishii S, Kubokawa K, Kikuchi M, Nishio H. 1995 Orientation of the Toad, Bufo japonicus,
toward the Breeding Pond. *Zoolog. Sci.* **12**, 475–484. (doi:10.2108/zsj.12.475)
16. Pašukonis A *et al.* 2022 Contrasting parental roles shape sex differences in poison frog
space use but not navigational performance. *eLife* **11**, e80483. (doi:10.7554/eLife.80483)
17. Beck KB, Loretto M-C, Ringler M, Hödl W, Pašukonis A. 2017 Relying on known or
exploring for new? Movement patterns and reproductive resource use in a tadpole-
transporting frog. *PeerJ* **5**, e3745. (doi:10.7717/peerj.3745)
18. Stynoski JL. 2009 Discrimination of offspring by indirect recognition in an egg-feeding
dendrobatid frog, *Oophaga pumilio. Anim. Behav.* **78**, 1351–1356.
(doi:10.1016/j.anbehav.2009.09.002)
19. Serrano-Rojas SJ, Pašukonis A. 2021 Tadpole-transporting frogs use stagnant water odor
to find pools in the rainforest. *J. Exp. Biol.* **224**, jeb243122. (doi:10.1242/jeb.243122)
20. Phillips JB, Diego-Rasilla FJ. 2022 The amphibian magnetic sense(s). *J. Comp. Physiol. A*
Neuroethol. Sens. Neural. Behav. Physiol. **208**, 723–742. (doi:10.1007/s00359-022-01584-
9)
21. Phillips JB, Borland SC. 1992 Behavioural evidence for use of a light-dependent
magnetoreception mechanism by a vertebrate. *Nature* **359**, 142–144.
(doi:10.1038/359142a0)
22. Phillips JB. 1986 Two Magnetoreception Pathways in a Migratory Salamander. *Science* **233**,
765–767. (doi:10.1126/science.3738508)
23. Shaykevich DA, Pašukonis A, O'Connell LA. 2022 Long distance homing in the cane toad
(Rhinella marina) in its native range. *J. Exp. Biol.* **225**, jeb243048. (doi:10.1242/jeb.243048)
24. Urban MC, Phillips BL, Skelly DK, Shine R. 2007 The cane toad's (Chaunus [Bufo] marinus)
increasing ability to invade Australia is revealed by a dynamically updated range model.
Proc. R. Soc. B Biol. Sci. **274**, 1413–1419. (doi:10.1098/rspb.2007.0114)
25. Urban MC, Phillips BL, Skelly DK, Shine R. 2008 A Toad More Traveled: The
Heterogeneous Invasion Dynamics of Cane Toads in Australia. *Am. Nat.* **171**, E134–E148.
(doi:10.1086/527494)
26. Shine R *et al.* 2021 Increased rates of dispersal of free-ranging cane toads (Rhinella

marina) during their global invasion. *Sci. Rep.* **11**, 23574. (doi:10.1038/s41598-021-02828-5)

27. Zug GR, Zug PB. 1979 The marine toad, Bufo marinus : a natural history resumé of native populations.

28. DeVore JL, Shine R, Ducatez S. 2021 Spatial ecology of cane toads (Rhinella marina) in their native range: a radiotelemetric study from French Guiana. *Sci. Rep.* **11**, 11817. (doi:10.1038/s41598-021-91262-8)

29. Ward-Fear G, Greenlees MJ, Shine R. 2016 Toads on Lava: Spatial Ecology and Habitat Use of Invasive Cane Toads (Rhinella marina) in Hawai'i. *PLOS ONE* **11**, e0151700. (doi:10.1371/journal.pone.0151700)

30. Westhoff G, Roth G. 2002 Morphology and projection pattern of medial and dorsal pallial neurons in the frog Discoglossus pictus and the salamander Plethodon jordani. *J. Comp. Neurol.* **445**, 97–121. (doi:10.1002/cne.10136)

31. Northcutt RG, Kicliter E. 1980 Organization of the Amphibian Telencephalon. In *Comparative Neurology of the Telencephalon* (ed SOE Ebbesson), pp. 203–255. Boston, MA: Springer US. (doi:10.1007/978-1-4613-2988-6_8)

32. Sotelo MI, Daneri MF, Bingman VP, Muzio RN. 2024 Amphibian spatial cognition, medial pallium and other supporting telencephalic structures. *Neurosci. Biobehav. Rev.* , 105739. (doi:10.1016/j.neubiorev.2024.105739)

33. Sinsch U. 1992 Sex-biassed site fidelity and orientation behaviour in reproductive natterjack toads (Bufo calamita). *Ethol. Ecol. Evol.* **4**, 15–32. (doi:10.1080/08927014.1992.9525347)

34. Yovanovich CAM, Jungblut LD, Heer T, Pozzi AG, Paz DA. 2009 Amphibian larvae and zinc sulphate: a suitable model to study the role of brain-derived neurotrophic factor (BDNF) in the neuronal turnover of the olfactory epithelium. *Cell Tissue Res.* **336**, 1–9. (doi:10.1007/s00441-009-0752-7)

35. Smith CG. 1951 Regeneration of sensory olfactory epithelium and nerves in adult frogs. *Anat. Rec.* **109**, 661–671. (doi:10.1002/ar.1091090403)

36. Hentig JT, Byrd-Jacobs CA. 2016 Exposure to Zinc Sulfate Results in Differential Effects on Olfactory Sensory Neuron Subtypes in Adult Zebrafish. *Int. J. Mol. Sci.* **17**, 1445. (doi:10.3390/ijms17091445)

37. Keller M, Douhard Q, Baum MJ, Bakker J. 2006 Destruction of the Main Olfactory Epithelium Reduces Female Sexual Behavior and Olfactory Investigation in Female Mice. *Chem. Senses* **31**, 315–323. (doi:10.1093/chemse/bjj035)

38. Freake MJ, Borland SC, Phillips JB. 2002 Use of a Magnetic Compass for Y-Axis Orientation in Larval Bullfrogs, Rana catesbeiana. *Copeia* **2002**, 466–471. (doi:10.1643/0045-8511(2002)002[0466:UOAMCF]2.0.CO;2)

39. Knight ZA, Tan K, Birsoy K, Schmidt S, Garrison JL, Wysocki RW, Emiliano A, Ekstrand MI, Friedman JM. 2012 Molecular profiling of activated neurons by phosphorylated ribosome capture. *Cell* **151**, 1126–1137. (doi:10.1016/j.cell.2012.10.039)

40. Fischer EK, Alvarez H, Lagerstrom KM, McKinney JE, Petrillo R, Ellis G, O'Connell LA. 2020 Neural correlates of winning and losing fights in poison frog tadpoles. *Physiol. Behav.* **223**, 112973. (doi:10.1016/j.physbeh.2020.112973)

41. Fischer EK, Roland AB, Moskowitz NA, Tapia EE, Summers K, Coloma LA, O'Connell LA. 2019 The neural basis of tadpole transport in poison frogs. *Proc Biol Sci* **286**, 20191084.

42. Fischer EK, Roland AB, Moskowitz NA, Tapia EE, Summers K, Coloma LA, O'Connell LA. 2019 The neural basis of tadpole transport in poison frogs. *Proc. R. Soc. B Biol. Sci.* **286**, 20191084. (doi:10.1098/rspb.2019.1084)

43. Sotelo MI, Daneri MF, Bingman VP, Muzio RN. 2016 Telencephalic Neuronal Activation Associated with Spatial Memory in the Terrestrial Toad Rhinella arenarum: Participation of the Medial Pallium during Navigation by Geometry. *Brain. Behav. Evol.* **88**, 149–160. (doi:10.1159/000447441)

44. Inés Sotelo M, Bingman VP, Muzio RN. 2022 The medial pallium and the spatial encoding
of geometric and visual cues in the terrestrial toad, Rhinella arenarum. *Neurosci. Lett.* **786**,
136801. (doi:10.1016/j.neulet.2022.136801)
45. Sotelo MI, Bingman VP, Muzio RN. 2019 The Mating Call of the Terrestrial Toad, Rhinella
arenarum, as a Cue for Spatial Orientation and Its Associated Brain Activity. *Brain. Behav.*
Evol. **94**, 7–17. (doi:10.1159/000504122)
46. Chersi F, Burgess N. 2015 The Cognitive Architecture of Spatial Navigation: Hippocampal
and Striatal Contributions. *Neuron* **88**, 64–77. (doi:10.1016/j.neuron.2015.09.021)
47. Vorhees CV, Williams MT. 2014 Assessing spatial learning and memory in rodents. *ILAR J.*
55, 310–332. (doi:10.1093/ilar/ilu013)
48. Moreno N, González A. 2004 Localization and connectivity of the lateral amygdala in anuran
amphibians. *J. Comp. Neurol.* **479**, 130–148. (doi:10.1002/cne.20298)
49. Calenge C. 2006 The package "adehabitat" for the R software: A tool for the analysis of
space and habitat use by animals. *Ecol. Model.* **197**, 516–519.
(doi:10.1016/j.ecolmodel.2006.03.017)
50. R Core Team. 2023 Package 'stats'. *CRAN*
51. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
52. Lenth R. 2024 Package 'emmeans'. *CRAN*
53. Landler L, Ruxton GD, Malkemper EP. 2021 Advice on comparing two independent
samples of circular data in biology. *Sci. Rep.* **11**, 20337. (doi:10.1038/s41598-021-99299-5)
54. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ,
Machler M, Bolker BM. 2017 glmmTMB balances speed and flexibility among packages for
zero-inflated generalized linear mixed modeling. *R J.* **9**, 378–400. (doi:10.3929/ethz-b-
000240890)
55. Hartig F. 2022 Package 'DHARMa': Residual Diagnostics for Hierarchical (Multi-Level /
Mixed) Regression Models. *CRAN*
56. Pašukonis A, Loretto M-C, Hödl W. 2018 Map-like navigation from distances exceeding
routine movements in the three-striped poison frog (Ameerega trivittata). *J. Exp. Biol.* **221**.
(doi:10.1242/jeb.169714)
57. Bingman VP, Mench JA. 1990 Homing behavior of hippocampus and parahippocampus
lesioned pigeons following short-distance releases. *Behav. Brain Res.* **40**, 227–238.
(doi:10.1016/0166-4328(90)90079-T)
58. Lohmann K, Lohmann C. 1996 Orientation and open-sea navigation in sea turtles. *J. Exp.*
Biol. **199**, 73–81.
59. Yartsev MM, Ulanovsky N. 2013 Representation of Three-Dimensional Space in the
Hippocampus of Flying Bats. *Science* **340**, 367–372. (doi:10.1126/science.1235338)
60. Clayton NS. 1998 Memory and the hippocampus in food-storing birds: a comparative
approach. *Neuropharmacology* **37**, 441–452. (doi:10.1016/S0028-3908(98)00037-9)
61. Woych J *et al.* 2022 Cell-type profiling in salamanders identifies innovations in vertebrate
forebrain evolution. *Science* (doi:10.1126/science.abp9186)
62. Tsanov M. 2017 Speed and Oscillations: Medial Septum Integration of Attention and
Navigation. *Front. Syst. Neurosci.* **11**. (doi:10.3389/fnsys.2017.00067)
63. Takamura Y, Tamura R, Zhou TL, Kobayashi T, Tran AH, Eifuku S, Ono T. 2006 Spatial
firing properties of lateral septal neurons. *Hippocampus* **16**, 635–644.
(doi:10.1002/hipo.20196)
64. Wirtshafter HS, Wilson MA. 2020 Differences in reward biased spatial representations in the
lateral septum and hippocampus. *eLife* **9**, e55252. (doi:10.7554/eLife.55252)
65. McDonald RJ, White NM. 2013 A triple dissociation of memory systems: Hippocampus,
amygdala, and dorsal striatum. *Behav. Neurosci.* **127**, 835–853. (doi:10.1037/a0034883)
66. Sinsch U. 2006 Orientation and navigation in Amphibia. *Mar. Freshw. Behav. Physiol.* **39**,

65–71. (doi:10.1080/10236240600562794)

67. Gagliardo A. 2013 Forty years of olfactory navigation in birds. *J. Exp. Biol.* **216**, 2165–2171.
(doi:10.1242/jeb.070250)

68. Marin AC, Schaefer AT, Ackels T. 2021 Spatial information from the odour environment in
mammalian olfaction. *Cell Tissue Res.* **383**, 473–483. (doi:10.1007/s00441-020-03395-3)

69. Phillips JB. 1977 Use of the earth's magnetic field by orienting cave salamanders (Eurycea
lucifuga). *J. Comp. Physiol.* **121**, 273–288. (doi:10.1007/BF00609616)

70. Deutschlander ME, Phillips JB, Borland SC. 2000 Magnetic Compass Orientation in the
Eastern Red-Spotted Newt, Notophthalmus viridescens: Rapid Acquisition of the Shoreward
Axis. *Copeia* **2000**, 413–419. (doi:10.1643/0045-8511(2000)000[0413:MCOITE]2.0.CO;2)

71. Landler L, Gollmann G. 2011 Magnetic orientation of the Common Toad: establishing an
arena approach for adult anurans. *Front. Zool.* **8**, 6. (doi:10.1186/1742-9994-8-6)

72. Bauer GB, Fuller M, Perry A, Dunn JR, Zoeger J. 1985 Magnetoreception and
Biomineralization of Magnetite in Cetaceans. In *Magnetite Biomineralization and*
Magnetoreception in Organisms: A New Biomagnetism (eds JL Kirschvink, DS Jones, BJ
MacFadden), pp. 489–507. Boston, MA: Springer US. (doi:10.1007/978-1-4613-0313-8_24)

73. Mouritsen H, Ritz T. 2005 Magnetoreception and its use in bird navigation. *Curr. Opin.*
Neurobiol. **15**, 406–414. (doi:10.1016/j.conb.2005.06.003)

74. Fischer JH, Freake MJ, Borland SC, Phillips JB. 2001 Evidence for the use of magnetic map
information by an amphibian. *Anim. Behav.* **62**, 1–10. (doi:10.1006/anbe.2000.1722)

75. Phillips JB, Freake MJ, Fischer JH, Borland CS. 2002 Behavioral titration of a magnetic map
coordinate. *J. Comp. Physiol. A* **188**, 157–160. (doi:10.1007/s00359-002-0286-x)

76. Diego-Rasilla FJ, Phillips JB. 2021 Evidence for the use of a high-resolution magnetic map
by a short-distance migrant, the Alpine newt (Ichthyosaura alpestris). *J. Exp. Biol.* **224**,
jeb238345. (doi:10.1242/jeb.238345)

77. Adler K. 1976 EXTRAOCULAR PHOTORECEPTION IN AMPHIBIANS. In *Extraretinal*
Photoreception (ed M Menaker), pp. 275–298. Pergamon. (doi:10.1016/B978-0-08-020965-
4.50008-0)

78. Yasumiba K, Alford RA, Schwarzkopf L. 2015 Why do male and female cane toads, *Rhinella*
marina, respond differently to advertisement calls? *Anim. Behav.* **109**, 141–147.
(doi:10.1016/j.anbehav.2015.08.015)

79. Liu Y, Day LB, Summers K, Burmeister SS. 2019 A cognitive map in a poison frog. *J. Exp.*
Biol. **222**, jeb197467. (doi:10.1242/jeb.197467)

80. Ferguson DE. 1971 The Sensory Basis of Orientation in Amphibians. *Ann. N. Y. Acad. Sci.*
188, 30–36. (doi:10.1111/j.1749-6632.1971.tb13087.x)