Main Manuscript for

Left-Right Brain-Wide Asymmetry of Neuroanatomy in the Mouse
Brain.

Andrew Silberfeld[1,2], James M. Roe[3], Jacob Ellegood[4,5], Jason P. Lerch[5,6,7,8], Lily Qiu[5,7], Yongsoo
Kim[9], Jong Gwan Lee[1,2], William D. Hopkins[10], Joanes Grandjean[11,12], Yangming Ou[13], Olivier
Pourquié*[1,2].

Affiliations:
1. Department of Genetics, Harvard Medical School, Boston, MA, USA.
2. Department of Pathology, Brigham and Women's Hospital, Boston, MA, USA.
3. Center for Lifespan Changes in Brain and Cognition (LCBC), Department of Psychology,
University of Oslo, Norway.
4. Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Toronto, Ontario,
Canada.
5. Mouse Imaging Centre, Hospital for Sick Children, Toronto, Ontario, Canada.
6. Department of Medical Biophysics, The University of Toronto, Toronto, ON, Canada.
7. Department of Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, ON,
Canada.
8. Department of Preclinical Imaging, Wellcome Centre for Integrative Neuroimaging, University
of Oxford, Oxford, UK.
9. Department of Neural and Behavioral Sciences, College of Medicine, The Pennsylvania State
University, Hershey, PA, USA.
10. Department of Comparative Medicine & Michale E Keeling Center for Comparative Medicine
and Research, The university of Texas MD Anderson Cancer Center.
11. Donders Institute for Brain, Behaviour, and Cognition, Nijmegen, Netherlands.
12. Department for Medical Imaging, Radboud University Medical Center, PO Box 9101,
Nijmegen, Netherlands.
13. Boston Children's Hospital, Harvard Medical School, Harvard University, Boston, MA, USA.

* Corresponding Author: Dr. Olivier Pourquié

Email: pourquie@genetics.med.harvard.edu

Author Contributions: A.S. and O.P. designed all experiments and wrote the paper. A.S.
performed all experiments, data analysis and computer programming related to mice and non-
human primates. J.M.R. performed all FreeSurfer analysis with human subjects. A.S. analyzed
human FreeSurfer cohort summaries. J.E., J.P.L., and L.Q. provided MRI scans from Qiu et al.
2018 and contributed information critical for determining left-right orientation for the POND and Qiu
et al. 2018 cohorts. Y.K. provided left-right orientation information for the BICCN and Newmaster
et al. 2020 cohorts. W.D.H. provided the Chimpanzee MRI scans along with left-right orientation
information. J.G.L and J.G. helped with interpretation of the data. Y.O. helped with interpretation of
the data and provided critical input and feedback for study design. All authors contributed to the
editing of the manuscript.

Declaration of Interests: These authors declare no competing interests.

This PDF file includes:

Abstract

Left-right asymmetry of the human brain is widespread through its anatomy and function. However, limited microscopic understanding of it exists, particularly for anatomical asymmetry where there are few well-established animal models. In humans, most brain regions show subtle, population-average regional asymmetries in thickness or surface area, alongside a macro-scale twisting called the cerebral petalia in which the right hemisphere protrudes anteriorly past the left. Here, we ask whether neuroanatomical asymmetries can be observed in mice, leveraging 6 mouse neuroimaging cohorts from 5 different research groups (~3,500 animals). We found an anterior-posterior pattern of volume asymmetry with anterior regions larger on the right and posterior regions larger on the left. This pattern appears driven by similar trends in surface area and positional asymmetries, with the results together indicating a small brain-wide twisting pattern, similar to the human cerebral petalia. Furthermore, the results show no apparent relationship to known functional asymmetries in mice, emphasizing the complexity of the structure-function relationship in brain asymmetry. By establishing a signature of anatomical brain asymmetry in mice, we aim to provide a foundation for future studies to probe the mechanistic underpinnings of brain asymmetry seen in humans – a feature of the brain with extremely limited understanding.

Keywords: brain asymmetry, lateralization, mouse brain, hemisphere, left-right, structure function, sex differences, morphometry, laterality.

Significance Statement

The human brain shows significant left-right anatomical asymmetry. Understanding its microscopic basis has implications for studies of autism and schizophrenia, evolution, embryonic brain development, and the relationship between structure and function in the brain. One of the biggest challenges to understanding this aspect of the brain is that animal models are limited. Here we show a brain-wide twisting pattern of asymmetry in the mouse brain using over 3,500 animals from six independent cohorts. These findings provide a basis for using mice to interrogate the microscopic underpinnings of anatomical asymmetry in humans and a roadmap for exploring anatomical asymmetry in additional species.

80   **Main Text**
81

Introduction

A central feature of the vertebrate body plan is its bilaterally symmetrical left-right
organization. Some structures, such as the visceral organs break this symmetry as a result of
asymmetrical signaling during development[1]. The human brain, while generally symmetrical
between sides, exhibits distinct asymmetries in both anatomy and function. Numerous cognitive
abilities such as language and face recognition depend on circuitry that reliably exhibits greater
activation on one side of the brain than the other, and does so in a consistent direction across the
population[2–8]. Anatomically, most regions of the human brain show small population-average
asymmetries of cortical thickness (CT) or surface area (SA)[9–12], alongside a macro-scale "twisting"
of the brain called the *cerebral petalia* in which the right hemisphere is shifted anteriorly relative to
the left[13–17]. Anatomical asymmetry of the posterior temporal lobe (the *planum temporale*) is thought
to relate to functional asymmetry of human language, although studies show this relationship
between anatomical and functional asymmetries to be complex at the individual level[18–22].

Extensive research in humans on anatomical asymmetry has identified highly reproducible
patterns of CT and SA asymmetries[9,23] as well as an anterior-posterior pattern of CT asymmetry[9,23–
25]. Neurodevelopmental and cytoskeletal genes have been associated with modified anatomical
asymmetry[26], with some of these genes also implicated in a recent genome-wide association study
of handedness[27]. However, fundamental aspects of human anatomical brain asymmetry remain
poorly understood and challenging to address in human subjects. For example, the cellular basis
of anatomical asymmetries seen in humans is still unknown, as is the mechanism behind the small
shifts observed in asymmetry for individuals with autism spectrum disorder[28,29] and
schizophrenia[30,31]. Furthermore, the developmental mechanisms driving region-specific brain
asymmetries remain unclear, as does the extent to which anatomical asymmetries are even present
in other mammalian species.

Understanding of these questions would benefit substantially from animal models for
conducting cellular-level studies of anatomical asymmetry. However, existing models have
limitations which largely hinder their applicability to asymmetry in humans. In zebrafish and
Drosophila, well-defined asymmetries respectively exist in the habenula[32–34] and brainstem[35,36], but
the rest of the brain in these animals shows no other apparent morphological asymmetries. This
raises an important question about whether these isolated examples seen in distant species are
fundamentally distinct from the extensive, subtle pattern of asymmetries seen brain-wide in
humans. In chimpanzees and macaques, asymmetry has been generally reported across many
brain regions, but studies show limited reproducibility[37–43]. This is further complicated by the
difficulty of comparing studies using different metrics (volume[37,38], SA[25], CT[25,44]), and different
levels of granularity (voxel-based[43,44] vs atlas-based[25,42] results). One exception to this
reproducibility challenge is the *planum temporale*, which is widely seen to be expanded in the left
hemisphere in chimpanzees[25,37,45–48] and baboons[49–51] – similar to humans[52,53]. Yet, this represents
the only well-defined example of asymmetry within those species. As a result, it has been difficult
to establish a consensus pattern of asymmetry for the whole brain in any species outside of
humans.

Alternatively, mice, which are a commonly used species for understanding the human
brain's organization[54], are an ideal model organism for studying neuroanatomical asymmetry.
Despite its overall appearance of symmetry, the mouse brain possesses many well-described
functional asymmetries: processing of vocalizations and pitch sweeps[55–58], visceral stimuli[59], spatial
memory[60–63], fear[64–67], and handedness[68]. In addition, numerous brain-wide structural imaging
datasets and sensitive tools for morphometry analysis exist[69–72], allowing for investigation into

neuroanatomical asymmetries along with replication analyses between cohorts – something that is
largely absent from the existing non-human literature.

Three prior studies have investigated neuroanatomical asymmetry in mice using single
cohorts, but as in primates, they reached conflicting descriptions. One study by Spring et al. 2010[73]
showed an overall larger left hemisphere volume of 2.8% while a more recent study by Elkind et al.
2023[74] identified an asymmetry pattern varying regionally without any clear trend. A third
investigation by Rivera-Olvera et al. 2024[75] further identified regional volume asymmetries but
which did not resemble either of the 2 previous studies. However, this investigation also reported
an anterior-posterior pattern of asymmetry: the right side being larger in the anterior with the left
side larger in the posterior. The widespread availability of distinct imaging datasets from separate
groups now provides the opportunity to comprehensively investigate this question in mice while
accounting for the overall generalizability of the results.

Here, we investigate neuroanatomical asymmetry in mice using multiple cohorts from
different imaging modalities. We found that anatomical brain asymmetry is indeed present in the
adult mouse brain but is initially obscured by strong cohort-specific patterns of asymmetry without
obvious trends. By instead looking for what is consistent between these groups of animals, we
identified an anterior-posterior pattern of volume asymmetry where anterior regions are larger on
the right and posterior regions are larger on the left. The results together indicate a global "twisting"
of the brain in mice, similar to the cerebral petalia, rather than the distinct local asymmetries
observed in humans. We thus propose mice as a novel mammalian animal model for investigating
neuroanatomical asymmetry at the microscopic level. Future understanding of the genetic and
cellular underpinnings of this asymmetry signature may provide insight into observations seen in
humans.
151
152 **Results**
153

Pipeline Overview. We obtained brain-wide imaging data from six different published studies
performed by five different research groups (**Table S1**). These studies used mice of different
genetic backgrounds, with drastically varying numbers of animals (n = 23 to 2,992). We used three
high-resolution serial histology studies (Serial Two-Photon Tomography; STPT) and three
magnetic resonance imaging (MRI) studies. Animals of a single study are referred to here as a
"cohort". To investigate anatomical asymmetry, we used automated image analysis to measure
volume in each atlas-defined brain region (**Fig. 1A**). We used the well-established software
ANTsPy[71] to perform automated linear and non-linear image registration, which aligned each
sample image to a symmetrical, three-dimensional (3D) reference template. This produces a 3D
image (Jacobian determinant field) in which each pixel represents the local volumetric change
between the original sample and the reference template. We then obtained region-specific left-right
asymmetries by integrating over this field in combination with the pixel-wise atlas annotations
(Common Coordinate Framework version 3; CCFv3).

Reproducible Volume Asymmetries Exist Across Independent Cohorts. Our investigation of
asymmetry initially considered each cohort individually. Doing so revealed a large number of
significantly asymmetric brain regions in each of the 4 largest cohorts as well as a substantial
enrichment for significant asymmetries in the 2 smallest (**Fig. S1**). However, when we attempted
to correlate these region-wise patterns across cohorts, we found only weak consistency between
them (**Fig. 1B**). In contrast, sub-sampling 20 animals from a given cohort showed much stronger
consistency within that cohort (**Fig. 1C**). We then concentrated on the 50 most significant leftward
(left > right) and 50 most significant rightward (right > left) asymmetric regions from each cohort,
out of 615 atlas regions, asking whether the most significant asymmetries from one cohort could
predict the direction of asymmetry for those same regions in a second cohort. We observed such
a result in 73% of cross-cohort comparisons (**Fig. 1D-E**), demonstrating a degree of predictive
capability but which was generally noisy. Overall, these asymmetry patterns, despite their

significance, indicate strong cohort-specific effects, with little linearity in the relationship across cohorts.

To instead identify more consistent asymmetries, we reasoned that the average asymmetry pattern of several cohorts together is likely to provide a more reproducible set of results than any one cohort individually. We thus performed a leave-one-out cross-validation, averaging the summary patterns of asymmetry from five cohorts and comparing the resulting values to the remaining sixth cohort, repeating this procedure for each combination of five training cohorts and 1 testing cohort. In five of the six possible combinations, the top 50 leftward and top 50 rightward asymmetries, defined using the five-cohort average, significantly predicted the direction of asymmetry in the remaining sixth cohort (**Fig. 1F**). The last combination showed a non-significant trend in the expected direction (likely non-significant due to the high rate of tissue defects present in the images from that study, combined with its low number of animals). This result means that on five of six unique occasions, the top asymmetries obtained through a five-cohort average generalized well to the remaining sixth cohort. By then taking the average asymmetry of all six cohort summaries together, we obtained a final, high-confidence description of asymmetries which are generalizable across independent cohorts.

We lastly confirmed that this result extended to new animals by identifying a seventh, previously un-analyzed cohort of mice from additional public data. We found that our final six-cohort average successfully and significantly predicted the pattern of asymmetry in this seventh previously un-used cohort (**Fig. 1G**). We further found that this result could be obtained with 80% power when using only 16-animal sub-samples from this cohort, indicating that for future studies, smaller-sized datasets can be used (**Fig. S2**). Overall, the magnitude of the most significant asymmetries in the six-cohort average ranges from 0.3% to 2.8% difference in regional volume between sides (**Fig. 1H, Dataset S1**). The biological implications of this asymmetry pattern are explored in **Fig. 4-7**, after the control experiments presented next (**Fig. 2-3**).

Brain Asymmetry in Mice Is Not Due to Technical Artifacts or Biases. We first sought to validate that our morphometry analysis can recapitulate previously reported results in a non-asymmetry context. We therefore used the same volume data generated above to calculate regional male-female differences in volume (**Fig. 2A**). Previous studies in mice have defined a clear signature of male-female neuroanatomical differences – a larger bed nuclei stria terminalis, medial amygdala, and medial preoptic area in males[74,76,77] – which we utilized as a positive control analysis. We found that male-female differences are highly reproducible across all 6 original cohorts, and we identified this expected regional signature as the top result (**Fig. 2B-G**). Additionally, we quantitatively compared our data to previous results by Elkind et al. 2023[74] who used one of the same cohorts as us (AMBCA). For this cohort, our results were consistent with theirs, both for sex differences as well as for asymmetry (**Fig. 2H-I, Fig. S3**), despite our groups using independent pipelines. Overall, these results confirm that we are recapitulating established findings.

We next sought to assess whether any bias in the image processing pipeline is contributing to our observed asymmetries. To do this, we reversed the left-right orientation of 1 STPT cohort (BICCN) and 1 MRI cohort (POND), as well as artificially symmetrized them, and re-ran our analysis. In the STPT cohort, left-right flipping the original images prior to any processing almost entirely inverted the asymmetry pattern (**Fig. 3A**). The symmetrized images similarly showed much smaller asymmetries than non-symmetrized images, but which, interestingly, were still highly significant (**Fig. 3B-C; Fig. S4A**). Nonetheless, this data shows that asymmetries in our non-symmetrized STPT cohorts depend almost entirely on the image left-right orientation, an important result which demonstrates that there is no left-right bias in the pipeline to drive asymmetries. Similarly, in the MRI cohort, we found that asymmetries also depended strongly on the left-right orientation but included some noise (**Fig. S4A-D**), likely from an MRI-specific algorithm (the N4 intensity inhomogeneity correction step), which, when given a pair of identical but flipped input images, does not produce perfectly flipped output images (**Fig. S4E-F**). We tested whether this

effect is contributing to the overall asymmetry pattern by comparing the average of mouse MRI
cohorts to the average of mouse STPT cohorts (**Fig. 3D-E**) – the latter of which do not use this
MRI-specific algorithm during their processing. We found that asymmetry generalizes between
these two fundamentally distinct image modalities and thus asymmetry is not a product of this noise
in the pipeline or any MRI-specific or STPT-specific artifact – such as *ex cranio* (STPT) vs *in cranio*
(MRI) imaging. Overall, the results indicate that our observed asymmetries depend predominantly
on the image left-right orientation and not on pipeline or tissue processing biases.

Finally, we assessed whether our results generalize to a different image registration
software – Elastix[72] – which uses a different transformation model from the ANTsPy registration
package used above. We also used different loss functions between these software. Re-registering
each sample using Elastix revealed a similar pattern of asymmetry to that observed with ANTsPy
(**Fig. 3F, Fig. S5**). This result makes the prospect of an overall registration error unlikely and rules
out the possibility that asymmetry is a software-specific result. Altogether, our results argue strongly
in favor of a population-wide pattern of directional asymmetry existing in the adult mouse brain,
robust to numerous biological and technical differences between cohorts, independent pipelines,
image modalities, and registration packages.

Top Asymmetries Are Comprised of All Major Anatomical Structures. Having established that
anatomical asymmetry is indeed present in the mouse brain, we turned our attention to interpreting
the regional pattern of asymmetry. Inspecting the set of top asymmetries reveals a diverse
collection of regions spanning each major division of the mouse brain (**Fig. 4A**). We saw significant
enrichment of atlas labels within the pons and cerebellum among the top set of asymmetries (**Fig.
4B**). However, outside of these areas, we saw no clear enrichment for any other division of the
brain among our top results. Among isocortical labels, we saw no evidence of layer-specific
enrichment either (**Fig. 4C**). Thus, excluding the pons and cerebellum, our set of top asymmetries
mainly reflects the frequency of these structures expected by chance given their proportion in the
CCFv3 atlas.

No Association of Anatomical Asymmetries with Functional Asymmetries. Among the top
asymmetric regions, the most asymmetric structure is the Paraflocculus, with a 2.8% average
rightward asymmetry (**Dataset S1**). The locus coeruleus and medial pretectal area further showed
a 2.6% leftward asymmetry. Finding these regions as the top asymmetric structures was
unexpected given that no evidence for lateralized function exists at these areas. Interestingly, no
auditory cortical regions were found in the set of top asymmetric structures despite robust evidence
of functionally asymmetric auditory processing in mice[55–58]. We therefore investigated, using the
statistical enrichment approach above, whether our set of top anatomical asymmetries were indeed
distinct from known functional asymmetries in mice (**Fig. S6A-B, Table S2**). We found that areas
of the mouse brain with strong evidence for asymmetric function showed no association with the
top anatomical asymmetries identified here (**Fig. 4D, Fig. S6C, Table S2**). We further saw through
leave-one-out cross-validation that these functionally asymmetric regions don't appear to drive the
main effects seen in **Fig. 1F,** as they were less consistent across cohorts than the overall set of top
regions (consistent only for 3 of 6 cohorts; **Fig. S6C**). Overall, these analyses show that gross
anatomical asymmetries seen here are generally distinct from known functional asymmetries in
mice.

An Anterior-Posterior Pattern of Volume Asymmetry is Driven by SA but Not CT. Although
we did not identify an association with known functional asymmetries, we did observe an anterior-
posterior (AP) pattern of asymmetry in which anterior regions are larger on the right and posterior
regions are larger on the left (**Fig. 1H**). We quantified this by calculating the 3D centroid of each
region from the CCFv3 atlas and plotting the AP component of that centroid against its asymmetry.
We observe this pattern both when using the set of top regions or when using all regions in our 6-
cohort average (**Fig. S7A, Fig. 5A**). Notably, this result is not restricted to the cortex, reflecting the
global nature of the effect. This AP effect is stronger in the 6-cohort average than in any one cohort
individually (**Fig. S7B-C, Fig. S8**), demonstrating how the AP effect is generally obscured by the

cohort-specific patterns. Flipping the left-right orientation of the BICCN and POND images reverses
the directionality of the pattern while symmetrizing these images eliminates it (**Fig. S7D-E**).
Moreover, this effect is evident in both the STPT- and MRI-specific averages and is also present in
our analysis using Elastix (**Fig. S7F-H**). We see no evidence of a similar trend for the dorsal-ventral
(DV) or medial-lateral (ML) axes (**Fig. 5B, Fig. S7I**).

In humans, a CT asymmetry trend exists where anterior regions are thicker on the left and
posterior regions are thicker on the right[9,23–25]. To directly assess whether our observations in mice
match this reported trend in humans, we measured CT asymmetry directly in all samples. We did
this by propagating the CCFv3 cortical labels onto each sample through the reverse registration
and applying the Laplacian method[78,79] (**Fig. 5C**; see supplemental methods). Although our AP
volume asymmetry pattern is brain-wide, we focused on the cortex here because measurements
of thickness are well-defined and because doing so best enables a direct comparison to
observations in humans. We found no evidence of a pattern in mice for CT asymmetry with respect
to the AP axis (**Fig. 5D**), but we also measured SA asymmetry in parallel and found a highly
significant AP gradient in the same direction as our volume asymmetry pattern (**Fig. 5E**).
Interestingly this gradient was in the opposite direction of the human CT pattern.

We validated these separate CT and SA measurements by using the hemisphere-specific
CT and SA to re-derive volume asymmetry for each cortical territory (layers 1-6b combined into a
single label per cortical area). This re-derived volume asymmetry was strongly correlated with our
original volume asymmetries measured from the forward registration (**Fig. 5F**), confirming the
accuracy of our measurements. Altogether, this data indicates that an AP volume asymmetry
pattern exists and is driven by SA asymmetries, but not CT.

Positional Asymmetries but No Cerebral Petalia Exists in Mice. Given the AP pattern of
asymmetry observed in mice, we wondered whether this indicated a cerebral petalia-like effect. In
humans, the right hemisphere protrudes ahead of the left in a brain-wide "twisting" or "warping"
pattern[13–17]. Roughly two-thirds of individuals show this trend while the remaining one-third show a
reversal or near-symmetry[17]. To examine this in mice, we applied an automated approach based
on recent large-scale papers measuring the petalia in humans and chimpanzees[13,14]. For reference
and validation, we also analyzed 4 cohorts of human subjects (positive controls) and 6 cohorts of
non-human primates (NHPs; negative controls) consisting of chimpanzees, macaques, and
marmosets (2 cohorts each; **Table S3**). We note that while the presence of a significant petalia in
NHPs is controversial, studies agree that such an effect, if present, is much smaller in magnitude
and of a weaker frequency in that species' population than in humans[13,17,80,81].

In humans, we found a significant right-frontal protrusion of 0.24% and a significant left-
occipital protrusion of 0.70%. Both are normalized by the length of the left hemisphere and are
consistent with established findings[13–16] (**Fig. 6A-B**). In NHPs, we saw a significant right-frontal
protrusion as well, but which was significantly smaller than in humans by approximately 3-fold
(0.07% overall; **Fig. 6A**). In contrast, we saw no significant occipital protrusion in NHPs (**Fig. 6B**).
In mice, we saw no significant protrusion for either cortical pole, with average measurements similar
to the NHPs (anterior = 0.06%, posterior = 0.07%; **Fig. 6A-B**). Thus, our measurements suggest
that mice do not exhibit an AP protrusion of one hemisphere with respect to the other (e.g. a
cerebral petalia).

Despite this, we considered whether a more sensitive approach might better reveal such
an effect, leveraging the statistical power of many brain regions rather than focusing on single
cortical poles. To explore this, we measured positional asymmetries across all CCFv3-annotated
regions within each mouse brain, using the 3D centroids of these regions as mapped onto that
animal during the petalia analysis. Cross-validation across different cohorts allowed us to assess
whether the position of sets of brain regions was shifted reproducibly between sides in an AP
direction. Indeed, we found highly reproducible left-right asymmetries of AP position in mice across
all 6 of our cohorts (**Fig. 6C**). Regions within the cerebral hemisphere were consistently shifted
anteriorly on the right side while regions within the brainstem and cerebellum were shifted anteriorly

on the left side (**Fig. 6D-E**). Thus, while no significant AP protrusion asymmetry exists at the mouse
cortical poles, mice do exhibit reproducible brain-wide shifts in the tissue that align with the AP
volume and SA gradients.

Evolutionary Context of Mouse Brain Volume Asymmetries. We lastly examined to what
degree, if any, our observed volume asymmetries in mice were consistent with those of humans
and NHPs. Initial inspection of the overall asymmetry pattern in each species revealed sharp
interspecies differences (**Fig. 7A**), along with cohort-specific asymmetries for marmosets and
macaques, similar to those seen in mice (**Fig. 7B, Fig. S9**). To quantitatively compare asymmetry
across species, we utilized the cross-species atlas annotations from Garrin et al. 2022[82], the
marmoset and macaque cohorts used earlier, and human volume asymmetry data from Roe et al.
2023[9]. The two chimpanzee cohorts from the previous section were excluded for technical reasons
(see methods).

We found weak correlations in the region-wise asymmetry pattern between any of the
species analyzed, using either cytoarchitecturally matched (Brodmann) regions or whole cortical
lobes with composite subcortical structures (**Fig. 7C-D**). We additionally found no evidence of an
AP pattern for volume asymmetry in marmosets or macaques, but interestingly, we found an AP
pattern in humans in the opposite direction of mice (**Fig. S10**) – consistent with the known CT
asymmetry pattern in humans. Overall, our data indicates anatomical asymmetry to be species-
specific, with little overlap between asymmetry patterns in mice, humans, and NHPs.

351
Discussion

Anatomical asymmetry is a significant feature of the human brain, yet there is a lack of
mammalian animal models in which to investigate its microscopic basis. Through our work, we
identified an anterior-posterior pattern of volume and SA asymmetry in mice, for which anterior
regions are larger and have more surface area on the right, while posterior regions are larger and
have more surface area on the left. The cerebral hemisphere as a whole is shifted anteriorly on the
right while the brainstem and cerebellum are shifted anteriorly on the left. Cross-cohort validation
and extensive control experiments demonstrate our results, though small in magnitude, are
consistent in 6 of 7 groups of animals, are free of pipeline bias, and are independent of image
modality and analysis software. Our data collectively indicates that a genuine brain-wide signature
of anatomical asymmetry exists in the adult mouse brain.

Considering our results in the context of humans, our findings best align with the idea of a
cerebral petalia-like effect in mice, rather than the distinct local asymmetries reported in humans.
Our set of 100 top asymmetries largely reflects the above anterior-posterior pattern. In this context,
it makes intuitive sense that our top anatomical asymmetries lack an obvious overlap with known
functionally asymmetric brain regions. The subtle asymmetries captured by our analysis appear to
represent a twisting pattern manifesting across the entire brain, while region-specific asymmetries
like those seen in humans or the zebrafish habenula appear to be entirely absent in mice. This
apparent decoupling of anatomical and functional asymmetry in mice underscores the complexity
of the relationship between anatomical and functional asymmetries reported in humans[18–22]. To-
date, the basis of functional asymmetries in mice has been identified at the levels of the
synapse[55,58], protein[83], and gene expression[55,56,59], but our data does not support the presence of
a gross anatomical substrate.

One possible explanation for the emergence of macroscale asymmetries is the molecular
chirality of cytoskeletal proteins. A recent GWAS study identified enrichment for tubulin genes as
genetic modifiers of anatomical asymmetry in humans[26]. In *Drosophila*, the actin cytoskeleton
shows an intrinsic handedness that is sufficient to polarize a cell's planarity and yield whole-organ
and whole-organism helical-twisting[84]. Neurite growth cones in rodents have been similarly
observed *in vitro* to twist in a right-screw (clock-wise) direction in a manner dependent on the actin

cytoskeleton[85–87], although whether this occurs *in vivo* is unknown. Our observations here are
consistent with such a mechanism occurring in mice.

An alternative explanation could be that asymmetry emerges through a subtle timing
difference in development between the left and right hemispheres. In humans, cortical folding in
the temporal lobe begins 1-2 weeks sooner on the right than on the left[88–90], and two recent studies
found that transcriptome-wide changes occurring with age in the early fetal brain were slightly more
advanced on one side than the other[91,92]. Anatomical asymmetries are theorized to be a "record"
of this developmental timing delay[93]. Whether such an effect is occurring in mice remains unknown
as does the potential mediator of such a timing delay. However, having mice as a novel animal
model of anatomical brain asymmetry can be a powerful tool to investigate this and other
hypotheses.

A longstanding area of interest in humans is the relationship between asymmetry of the
visceral organs and asymmetry of the brain. While most evidence points toward separate
mechanisms conferring asymmetry in these contexts[94–99], one potential exception is the cerebral
petalia. In humans, individuals with reversed visceral organ asymmetry (*situs inversus*) show a high
frequency of a reversed cerebral petalia[95–97,100], although the number of subjects studied in this
context remains low (45 total over four studies). This emerging idea is particularly interesting in
light of our observation of a similar torsion-like effect in the mouse brain. Two prior studies in mice
that reversed visceral organ asymmetry observed the loss of a functional asymmetry in the
hippocampus[62,101]. Thus, we wonder whether the signature of neuroanatomical asymmetry we
observe here would be reversed in mice with *situs inversus*. The rarity of such individuals in humans
makes this relationship challenging to interrogate (hence the low number of subjects thus far), but
research using our novel signature of neuroanatomical asymmetry in mice would provide a critical
datapoint for better understanding this relationship.

Toward the goal of building consensus in the mouse asymmetry literature, our study makes
significant strides by drawing together the seemingly unrelated findings of Elkind et al. 2023[74] and
Rivera-Olvera et al. 2024[75]. We quantitatively replicate the volume asymmetry results of Elkind et
al. while we observe the same AP asymmetry trend as Rivera-Olvera et al. Our multi-cohort
analysis also clarifies the interpretation of both studies by demonstrating that apparent region-
specific asymmetries in mice should be understood as part of a global pattern, rather than isolated
expansions of specific structures. However, our study cannot account for the left-hemisphere bias
reported in Spring et al. 2010[73]; perhaps it stems from a residual global intensity or distortion bias
from the MRI. In our work, we observed a global bias in positional asymmetries of the Guma2021
cohort, with a global trend toward regions shifting anteriorly on the left, but the relative difference
in asymmetry between anterior and posterior regions remained evident. Such a true global bias in
the brain appears doubtful to us when considering the other cohorts in our analysis.

For future studies attempting to replicate our results, the asymmetry for the top 100 regions
can serve as a direct test of replicability by asking the following question: when measured in a new
set of animals, do the leftward and rightward asymmetries defined in this study vary significantly
from each other in the predicted direction? (e.g. **Fig. 1F**). A significant difference in this context in
the predicted direction provides evidence of replication. This approach can also be used for our
reported positional asymmetries.

Our focus here on sets of asymmetries highlights the probabilistic nature of our results.
The sets of leftward and rightward asymmetries we define are repeatable across independent
cohorts, however, individual regions within these sets cannot be considered significantly
asymmetric on their own. Nonetheless, defining sets of asymmetric regions ultimately ends up
being a strength when applied to future studies: an investigation into the cellular properties of
asymmetric regions is likely to be more powerful if the results are applicable over many regions –
even with some added noise – rather than in only a few focused regions.

A surprising finding from our study is the significant asymmetries present in most mouse
imaging cohorts but which tend to be cohort-specific. This variability contrasts with studies of sex

differences, where effects in one cohort are consistently reproduced in another. Since this cross-
cohort variability is specific to left-right comparisons, we believe it is unlikely to arise from factors
affecting both hemispheres equally, such as image modality, number of subjects, or original image
resolution. Similarly, biological influences – like age, genetic variation, the male-female ratio –
appear to have only minor impacts on human brain asymmetry and would not account for the
dramatic differences seen between cohorts[9,23,26,102]. Instead, we suspect technical factors, which
do not impact both sides equally, may be responsible, although further research is needed to
confirm this. For STPT samples, causal factors could be the intensity vignetting in the tiled images.
For MRI samples, this could be residual shading biases or residual geometric distortion, among
other possibilities. In principle, these technical issues could create cohort-specific local minima
during the registration's optimization process that could trap a more "aggressive" image registration
– especially given the small magnitude of the asymmetries. An alternative explanation could involve
husbandry-specific factors driving environment-dependent asymmetries. However, this seems
unlikely given that the two marmoset cohorts we analyzed showed significant and largely distinct
asymmetry patterns, despite being raised in the same environment at the same institution (personal
communication, acknowledged individual J.Hata). Ultimately, our cross-cohort approach proved
advantageous to a single-cohort analysis, allowing us to identify this critical issue that should be
carefully considered in future studies building consensus asymmetry patterns beyond humans.

Although we observe no meaningful correlations in asymmetry between species,
interpretation of these results should be made cautiously. Cross-species comparisons in general
are not well-defined and this represents a broader challenge facing the field of neuroimaging. To
find similar regions between species, we implemented a curated, atlas-based approach, however,
alignment of templates between species using white-matter estimation maps[103] or a common
transcriptional space[104] may provide an improvement. Nonetheless, our results align with a
previous study which did not observe a major cross-species signature of asymmetry between
humans and chimpanzees[25]. Similarly, a study of male-female differences in humans and mice
observed weak cross-species agreement, despite robust sex differences in both species[77]. Going
forward, additional methods should be explored to further interrogate the question of cross-species
comparisons, but at present, major consistency across species seems absent.

As brain-wide mapping studies become easier to perform in small vertebrates, we
encourage authors to publish their raw images in open-source repositories like the brain image
library[105] or Zenodo.org, along with documentation to identify the left and right hemispheres. The
presence in mice of a brain-wide signature of anatomical asymmetry will further enable research
into the question of developmental and cellular underpinnings of asymmetry in humans, and
publicly available datasets like those used in our study will be invaluable toward this effort.
467
468 Overall, we find extensive anatomical asymmetry in the adult mouse brain, manifesting as
469 a brain-wide, anterior-posterior pattern. Our work provides a foundation for future studies to
470 investigate the microscopic basis of this cerebral petalia-like effect in mice.
471
472
473 **Materials and Methods**
474
Overall Morphometry Analysis. Our data collection was performed using a series of python, Fiji,
and R scripts. Images for each animal were first gathered into a z-stack (if not already) and
converted to .tif file format from one of various starting formats (.jpg, .Jp2, .nii, .mnc, .tif). From
here, each z-stack was resampled from its original resolution to 50 μm isotropic resolution. MRI
samples were skull-stripped followed by N4 bias correction. We then performed image registration
(rigid, affine, and non-linear) to align the sample images toward a population-average template
(sample image as the moving image; reference template as the fixed image). All templates and
atlases were completely left-right symmetrical. From these registrations, we collected the Jacobian
determinant field and measured its average value within each CCFv3 atlas region for each sample.
The Jacobian determinant field represents how much the local volume changes at each voxel when

transforming one image to align with another. For example, field values of 1.10 or 0.85 at 2 different voxels in the template image indicate that the corresponding volumes in the sample image were 10% larger or 15% smaller respectively than they are in the template image. After calculating average field values for each regions, we then scaled the average field value by the physical size of that region in the original atlas to obtain a volume measurement in μm^3. Extensive details about each cohort, registration settings, processing steps, additional analyses, left-right orientation, and more are provided in **Supplemental Materials and Methods**.

Acknowledgments

We thank numerous members of the neuroimaging community for their generous feedback throughout this project. In this context, we would like to acknowledge Doug Greve, Elisa Guma, Stefan Klein, Yael Balbastre, and Clyde Francks. We would also like to thank individuals who provided critical information related to the left-right orientation or demographics of different cohorts: Gabriel Devenyi, John Sled, Rasmus Birn, Jonathan Oler, Junichi Hata, and Matthew Rushworth. We are thankful to Amit Zeisel for sharing supplemental data from their publication (Elkind et al. 2023[74]). We thank the Harvard Research Computing department for providing early access to the O2portal. We also thank members of the Pourquie lab for thoughtful feedback on the data and manuscript throughout the project. Finally, we are thankful for funding which this project received from a Harvard Brain Science Initiative Bipolar Disorder Seed Grant.

References

1. Hamada H. Molecular and cellular basis of left-right asymmetry in vertebrates. *Proc Japan Acad Ser B Phys Biol Sci*. 2020;96(7):273-296. doi:10.2183/PJAB.96.021

2. Wade DT, Hewer RL, David RM, Enderby P. Aphasia after stroke: Natural history and associated deficits. *J Neurol Neurosurg Psychiatry*. 1986;49(1):11-16. doi:10.1136/jnnp.49.1.11

3. Packheiser J, Schmitz J, Arning L, Beste C, Güntürkün O, Ocklenburg S. A large-scale estimate on the relationship between language and motor lateralization. *Sci Rep*. 2020;10(1):1-10. doi:10.1038/s41598-020-70057-3

4. Becker E, Karnath HO. Incidence of visual extinction after left versus right hemisphere stroke. *Stroke*. 2007;38(12):3172-3174. doi:10.1161/STROKEAHA.107.489096

5. Çiçek M, Deouell LY, Knight RT. Brain activity during landmark and line bisection tasks. 2009;3(May):1-8. doi:10.3389/neuro.09.007

6. Hecaen H, Angelergues R. Agnosia for Faces (Prosopagnosia). *Arch Neurol*. 1962;7(2):92-100. doi:10.1001/archneur.1962.04210020014002

7. Hilliard RD. Hemispheric Laterality Effects on a Facial Recognition Task in Normal Subjects. *Cortex*. 1968;9(3):246-258. doi:10.1016/S0010-9452(73)80002-4

8. Rossion B, Hanseeuw B, Dricot L. Defining face perception areas in the human brain: A large-scale factorial fMRI face localizer analysis. *Brain Cogn*. 2012;79(2):138-157. doi:10.1016/j.bandc.2012.01.001

9. Roe JM, Vidal-Piñeiro D, Amlien IK, et al. Tracing the development and lifespan change of population-level structural asymmetry in the cerebral cortex. *Elife*. 2023;12:1-32. doi:10.7554/eLife.84685

531   10.   Goldberg E, Roediger D, Kucukboyaci NE, et al. Hemispheric asymmetries of cortical
532   volume in the human brain. *Cortex*. 2013;49(1):200-210. doi:10.1016/j.cortex.2011.11.002

533   11.   Chi JG, Dooling EC, Gilles FH. Left-Right Asymmetries of the Temporal Speech Areas of
534   the Human Fetus. *Arch Neurol*. 1977;34(6):346-348.
535   doi:10.1001/archneur.1977.00500180040008

536   12.   Kertesz A, Geschwind N. Patterns of Pyramidal Decussation and Their Relationship to
537   Handedness. *Arch Neurol*. 1971;24(4):326-332.
538   doi:10.1001/archneur.1971.00480340058006

539   13.   Xiang L, Crow T, Roberts N. Cerebral torque is human specific and unrelated to brain
540   size. *Brain Struct Funct*. 2019;224(3):1141-1150. doi:10.1007/s00429-018-01818-0

541   14.   Zhao L, Matloff W, Shi Y, Cabeen RP, Toga AW. Mapping Complex Brain Torque
542   Components and Their Genetic Architecture and Phenomic Associations in 24,112
543   Individuals. *Biol Psychiatry*. 2022;91(8):753-768. doi:10.1016/j.biopsych.2021.11.002

544   15.   Bear D, Schiff D, Saver J, Greenberg M, Freeman R. Quantitative Analysis of Cerebral
545   Asymmetries: Fronto-occipital Correlation, Sexual Dimorphism and Association with
546   Handedness. *Arch Neurol*. 1986;43(6):598-603.
547   doi:10.1001/archneur.1986.00520060060019

548   16.   Balzeau A, Gilissen E, Grimaud-Hervé D. Shared pattern of endocranial shape
549   asymmetries among great apes, anatomically modern humans, and fossil hominins. *PLoS
550   One*. 2012;7(1):1-10. doi:10.1371/journal.pone.0029581

551   17.   Kong XZ, Postema M, Schijven D, et al. Large-Scale Phenomic and Genomic Analysis of
552   Brain Asymmetrical Skew. *Cereb Cortex*. 2021;31(9):4151-4168.
553   doi:10.1093/cercor/bhab075

554   18.   Greve DN, Van der Haegen L, Cai Q, et al. A Surface-based Analysis of Language
555   Lateralization and Cortical Asymmetry. *J Cogn Neurosci*. 2013;25(9):1477-1492.
556   doi:10.1162/jocn_a_00405

557   19.   Guadalupe T, Kong XZ, Akkermans SEA, Fisher SE, Francks C. Relations between
558   hemispheric asymmetries of grey matter and auditory processing of spoken syllables in
559   281 healthy adults. *Brain Struct Funct*. 2022;227(2):561-572. doi:10.1007/s00429-021-
560   02220-z

561   20.   Josse G, Kherif F, Flandin G, Seghier ML, Price CJ. Predicting language lateralization
562   from gray matter. *J Neurosci*. 2009;29(43):13516-13523. doi:10.1523/JNEUROSCI.1680-
563   09.2009

564   21.   Jansen A, Liuzzi G, Deppe M, et al. Structural correlates of functional language
565   dominance: A voxel-based morphometry study: Experimental laboratory research. *J
566   Neuroimaging*. 2010;20(2):148-156. doi:10.1111/j.1552-6569.2009.00367.x

567   22.   Chormai P, Pu Y, Hu H, Fisher SE, Francks C, Kong XZ. Machine learning of large-scale
568   multimodal brain imaging data reveals neural correlates of hand preference. *Neuroimage*.
569   2022;262:0-28. doi:10.1016/j.neuroimage.2022.119534

570   23.   Kong XZ, Mathias SR, Guadalupe T, et al. Mapping cortical brain asymmetry in 17,141
571   healthy individuals worldwide via the ENIGMA consortium. *Proc Natl Acad Sci U S A*.

572     2018;115(22):E5154-E5163. doi:10.1073/pnas.1718418115

573 24. Plessen KJ, Hugdahl K, Bansal R, Hao X, Peterson BS. Sex, age, and cognitive correlates
574     of asymmetries in thickness of the cortical mantle across the life span. *J Neurosci*.
575     2014;34(18):6294-6302. doi:10.1523/JNEUROSCI.3692-13.2014

576 25. Xiang L, Crow TJ, Hopkins WD, Roberts N. Comparison of Surface Area and Cortical
577     Thickness Asymmetry in the Human and Chimpanzee Brain. Published online 2020:1-15.
578     doi:10.1093/cercor/bhaa202

579 26. Sha Z, Schijven D, Carrion-Castillo A, et al. The genetic architecture of structural left–right
580     asymmetry of the human brain. *Nat Hum Behav*. Published online 2021.
581     doi:10.1038/s41562-021-01069-w

582 27. Cuellar-Partida G, Tung JY, Eriksson N, et al. Genome-wide association study identifies
583     48 common genetic variants associated with handedness. *Nat Hum Behav*. Published
584     online 2020. doi:10.1038/s41562-020-00956-y

585 28. Postema MC, van Rooij D, Anagnostou E, et al. Altered structural brain asymmetry in
586     autism spectrum disorder in a study of 54 datasets. *Nat Commun*. 2019;10(1):1-12.
587     doi:10.1038/s41467-019-13005-8

588 29. Sha Z, Rooij D, Anagnostou E, et al. Subtly altered topological asymmetry of brain
589     structural covariance networks in autism spectrum disorder across 43 datasets from the
590     ENIGMA consortium. 2022;(August 2021). doi:10.1038/s41380-022-01452-7

591 30. Schijven D, Postema MC, Fukunaga M, et al. Large-scale analysis of structural brain
592     asymmetries in schizophrenia via the ENIGMA consortium. 2023;120(14):1-12.

593 31. Sommer I, Aleman A, Ramsey N, Bouma A, Kahn R. Handedness, language lateralisation
594     and anatomical asymmetry in schizophrenia: Meta-analysis. *Br J Psychiatry*.
595     2001;178(APR.):344-351. doi:10.1192/bjp.178.4.344

596 32. Concha ML, Russell C, Regan JC, et al. Local Tissue Interactions across the Dorsal
597     Midline of the Forebrain Establish CNS Laterality to the parapineal precursors. Ablation of
598     cells in the left ventral epithalamus can reverse laterality in wild-type embryos and impose
599     the direction of CNS asym-. *Neuron*. 2003;39:423-438.

600 33. Roussigné M, Bianco IH, Wilson SW, Blader P. Nodal signalling imposes left-right
601     asymmetry upon neurogenesis in the habenular nuclei. *Development*. 2009;136(9):1549-
602     1557. doi:10.1242/dev.034793

603 34. Lekk I, Duboc V, Faro A, Nicolaou S, Blader P, Wilson SW. Sox1A mediates the ability of
604     the parapineal to impart habenular left-right asymmetry. *Elife*. 2019;8:1-23.
605     doi:10.7554/eLife.47376

606 35. Pascual A, Huang KL, Neveu J, Préat T. Brain asymmetry and long-term memory. *Nature*.
607     2004;427(6975):605-606. doi:10.1038/427605a

608 36. Lapraz F, Boutres C, Fixary-Schuster C, et al. Asymmetric activity of NetrinB controls
609     laterality of the Drosophila brain. *Nat Commun*. 2023;14(1). doi:10.1038/s41467-023-
610     36644-4

37. Hopkins WD, Taglialatela JP, Meguerditchian A, Nir T, Schenker NM, Sherwood CC. Gray
matter asymmetries in chimpanzees as revealed by voxel-based morphometry.
Neuroimage. 2008;42(2):491-497. doi:10.1016/j.neuroimage.2008.05.014

38. Vickery S, Hopkins WD, Sherwood CC, et al. Chimpanzee brain morphometry utilizing
standardized mri preprocessing and macroanatomical annotations. *Elife*. 2020;9:1-20.
doi:10.7554/eLife.60136

39. Cantalupo C, Hopkins WD. Asymmetric Broca's area in great apes. *Nature*.
2001;414(6863):505-505. doi:10.1038/35107134

40. Keller SS, Roberts N, Hopkins W. A comparative magnetic resonance imaging study of
the anatomy, variability, and asymmetry of Broca's area in the human and chimpanzee
brain. *J Neurosci*. 2009;29(46):14607-14616. doi:10.1523/JNEUROSCI.2892-09.2009

41. Hopkins WD, Misiura M, Pope SM, Latash EM. Behavioral and brain asymmetries in
primates: a preliminary evaluation of two evolutionary hypotheses. *Ann N Y Acad Sci*.
2015;1359(1):65-83. doi:10.1111/nyas.12936

42. Kim J, Jung Y, Barcus R, et al. Rhesus Macaque Brain Developmental Trajectory: A
Longitudinal Analysis Using Tensor-Based Structural Morphometry and Diffusion Tensor
Imaging. *Cereb Cortex*. 2020;30(8):4325-4335. doi:10.1093/cercor/bhaa015

43. Xia J, Wang F, Wu Z, et al. Mapping hemispheric asymmetries of the macaque cerebral
cortex during early brain development. *Hum Brain Mapp*. 2020;41(1):95-106.
doi:10.1002/hbm.24789

44. Hopkins WD, Avants BB. Regional and hemispheric variation in cortical thickness in
chimpanzees (Pan troglodytes). *Ann Intern Med*. 2013;158(6):5241-5248.
doi:10.1523/JNEUROSCI.2996-12.2013

45. Gannon PJ, Holloway RL, Broadfield DC, Braun AR. Asymmetry of chimpanzee planum
temporale: Humanlike pattern of Wernicke's brain language area homolog. *Science (80-)*.
1998;279(5348):220-222. doi:10.1126/science.279.5348.220

46. Hopkins WD, Marino L, Rilling JK, MacGregor LA. Planum temporale asymmetries in
great apes as revealed by magnetic resonance imaging (MRI). *Neuroreport*.
1998;9(12):2913-2918. doi:10.1097/00001756-199808240-00043

47. Cantalupo C, Pilcher DL, Hopkins WD. Are planum temporale and sylvian fissure
asymmetries directly related? A MRI study in great apes. *Neuropsychologia*.
2003;41(14):1975-1981. doi:10.1016/S0028-3932(02)00288-9

48. Hopkins WD, Nir TM. Planum temporale surface area and grey matter asymmetries in
chimpanzees (Pan troglodytes): The effect of handedness and comparison with findings in
humans. *Behav Brain Res*. 2010;208(2):436-443. doi:10.1016/j.bbr.2009.12.012

49. Marie D, Roth M, Lacoste R, et al. Left Brain Asymmetry of the Planum Temporale in a
Nonhominid Primate: Redefining the Origin of Brain Specialization for Language. *Cereb
Cortex*. 2018;28(5):1808-1815. doi:10.1093/cercor/bhx096

50. Becker Y, Sein J, Velly L, et al. Early Left-Planum Temporale Asymmetry in newborn
monkeys (Papio anubis): A longitudinal structural MRI study at two stages of
development. *Neuroimage*. 2021;227(July 2020):117575.

652   doi:10.1016/j.neuroimage.2020.117575

653   51.   Becker Y, Meguerditchian A. Structural Brain Asymmetries for Language: A Comparative
654   Approach across Primates. *Symmetry (Basel)*. 2022;14(5):1-21.
655   doi:10.3390/sym14050876

656   52.   Geschwind N, Levitsky W. Human Brain : Left-Right Asymmetries in Temporal Speech
657   Region Published by : American Association for the Advancement of Science Stable URL :
658   http://www.jstor.org/stable/1723941 Human Brain : Left-Right Asymmetries in Temporal
659   Speech Region. 1968;161(3837):186-187.

660   53.   Steinmetz H, Volkmann J, Jäncke L, Freund H -J. Anatomical left-right asymmetry of
661   language-related temporal cortex is different in left- and right-handers. *Ann Neurol*.
662   1991;29(3):315-319. doi:10.1002/ana.410290314

663   54.   Ellenbroek B, Youn J. Rodent models in neuroscience research: Is it a rat race? *DMM Dis*
664   *Model Mech*. 2016;9(10):1079-1087. doi:10.1242/dmm.026120

665   55.   Levy RB, Marquarding T, Reid AP, Pun CM, Renier N, Oviedo H V. Circuit asymmetries
666   underlie functional lateralization in the mouse auditory cortex. *Nat Commun*. 2019;10(1).
667   doi:10.1038/s41467-019-10690-3

668   56.   Marlin BJ, Mitre M, D'Amour JA, Chao M V., Froemke RC. Oxytocin enables maternal
669   behaviour by balancing cortical inhibition. *Nature*. 2015;520(7548):499-504.
670   doi:10.1038/nature14402

671   57.   Ehret G. Left hemisphere advantage in the mouse brain for recognizing ultrasonic
672   communication calls. *Nature*. 1987;325(6101):249-251. doi:10.1038/325249a0

673   58.   Neophytou D, Arribas DM, Arora T, Levy RB, Park IM, Oviedo H V. Differences in
674   temporal processing speeds between the right and left auditory cortex reflect the strength
675   of recurrent synaptic connectivity. *PLoS Biol*. 2022;20(10):1-20.
676   doi:10.1371/journal.pbio.3001803

677   59.   Wu Y, Chen C, Chen M, et al. The anterior insular cortex unilaterally controls feeding in
678   response to aversive visceral stimuli in mice. *Nat Commun*. 2020;11(1).
679   doi:10.1038/s41467-020-14281-5

680   60.   El-Gaby M, Reeve HM, Lopes-dos-Santos V, et al. An emergent neural coactivity code for
681   dynamic memory. *Nat Neurosci*. 2021;24(5):694-704. doi:10.1038/s41593-021-00820-w

682   61.   Shipton OA, El-Gaby M, Apergis-Schoute J, et al. Left-right dissociation of hippocampal
683   memory processes in mice. *Proc Natl Acad Sci U S A*. 2014;111(42):15238-15243.
684   doi:10.1073/pnas.1405648111

685   62.   Goto K, Kurashima R, Gokan H, Inoue N, Ito I, Watanabe S. Left-right asymmetry defect
686   in the hippocampal circuitry impairs spatial learning and working memory in IV mice. *PLoS*
687   *One*. 2010;5(11):1-7. doi:10.1371/journal.pone.0015468

688   63.   Song D, Wang D, Yang Q, et al. The lateralization of left hippocampal CA3 during the
689   retrieval of spatial working memory. *Nat Commun*. 2020;11(1):1-13. doi:10.1038/s41467-
690   020-16698-4

64. Scicli AP, Petrovich GD, Swanson LW, Thompson RF. Contextual Fear Conditioning Is
Associated with Lateralized Expression of the Immediate Early Gene c-fos in the Central
and Basolateral Amygdalar Nuclei. *Behav Neurosci*. 2004;118(1):5-14. doi:10.1037/0735-
7044.118.1.5

65. Kim S, Mátyás F, Lee S, Acsády L, Shin HS. Lateralization of observational fear learning
at the cortical but not thalamic level in mice. *Proc Natl Acad Sci U S A*.
2012;109(38):15497-15501. doi:10.1073/pnas.1213903109

66. Zhang MM, Geng AQ, Chen K, et al. Glutamatergic synapses from the insular cortex to
the basolateral amygdala encode observational pain. *Neuron*. 2022;110(12):1993-
2008.e6. doi:10.1016/j.neuron.2022.03.030

67. Kim SW, Kim M, Baek J, et al. Hemispherically lateralized rhythmic oscillations in the
cingulate-amygdala circuit drive affective empathy in mice. *Neuron*. 2023;111(3):418-
429.e4. doi:10.1016/j.neuron.2022.11.001

68. Collins RL. On the inheritance of Laterality. *J Hered*. 1968;1(59):9-12.
doi:10.1136/bmj.1.2317.1175-a

69. Wang Q, Ding SL, Li Y, et al. The Allen Mouse Brain Common Coordinate Framework: A
3D Reference Atlas. *Cell*. 2020;181(4):936-953.e20. doi:10.1016/j.cell.2020.04.007

70. Avants BB, Epstein CL, Grossman M, Gee JC. Symmetric diffeomorphic image
registration with cross-correlation: evaluating automated labeling of elderly and
neurodegenerative brain. *Med Image Anal*. 2008;12(1):26-41.
doi:10.1016/j.media.2007.06.004

71. Tustison NJ, Cook PA, Holbrook AJ, et al. The ANTsX ecosystem for quantitative
biological and medical imaging. *Sci Rep*. 2021;11(1):9068. doi:10.1038/s41598-021-
87564-6

72. Klein S, Staring M, Murphy K, Viergever MA, Pluim JPW. elastix : A Toolbox for Intensity-
Based Medical Image Registration. 2010;29(1):196-205.

73. Spring S, Lerch JP, Wetzel MK, Evans AC, Henkelman RM. Cerebral asymmetries in 12-
week-old C57Bl/6J mice measured by magnetic resonance imaging. *Neuroimage*.
2010;50(2):409-415. doi:10.1016/j.neuroimage.2009.12.043

74. Elkind D, Hochgerner H, Aloni E, Shental N, Zeisel A. Sex, strain, and lateral differences
in brain cytoarchitecture across a large mouse population. *Elife*. 2023;12:1-21.
doi:10.7554/eLife.82376

75. Rivera-Olvera A, Houwing DJ, Ellegood J, et al. The universe is asymmetric, the mouse
brain too. *Mol Psychiatry*. 2024;(September 2023):1-8. doi:10.1038/s41380-024-02687-2

76. Qiu LR, Fernandes DJ, Szulc-Lerch KU, et al. Mouse MRI shows brain areas relatively
larger in males emerge before those larger in females. *Nat Commun*. 2018;9(1).
doi:10.1038/s41467-018-04921-2

77. Guma E, Beauchamp A, Liu S, et al. Comparative neuroimaging of sex differences in
human and mouse brain anatomy. Published online 2023.
https://doi.org/10.1101/2023.08.23.554334

78.   Jones SE, Buchbinder BR, Aharon I. Three-dimensional mapping of cortical thickness
732   using Laplace's equation. *Hum Brain Mapp*. 2000;11(1):12-32. doi:10.1002/1097-
733   0193(200009)11:1<12::AID-HBM20>3.0.CO;2-K

734   79.   Lerch JP, Carroll JB, Dorr A, et al. Cortical thickness measured from MRI in the YAC128
735   mouse model of Huntington's disease. *Neuroimage*. 2008;41(2):243-251.
736   doi:10.1016/j.neuroimage.2008.02.019

737   80.   Holloway RL, De La Costelareymondie MC. Brain endocast asymmetry in pongids and
738   hominids: Some preliminary findings on the paleontology of cerebral dominance. *Am J
739   Phys Anthropol*. 1982;58(1):101-110. doi:10.1002/ajpa.1330580111

740   81.   Fears SC, Scheibel K, Abaryan Z, et al. Anatomic brain asymmetry in vervet monkeys.
741   *PLoS One*. 2011;6(12):1-10. doi:10.1371/journal.pone.0028243

742   82.   Garin CM, Garin M, Silenzi L, Jaffe R, Constantinidis C. Multilevel atlas comparisons
743   reveal divergent evolution of the primate brain. *Proc Natl Acad Sci U S A*. 2022;119(25):1-
744   9. doi:10.1073/pnas.2202491119

745   83.   Kawakami R, Shinohara Y, Kato Y, Sugiyama H, Shigemoto R, Ito I. Asymmetrical
746   allocation of NMDA receptor ε2 subunits in hippocampal circuitry. *Science (80-)*.
747   2003;300(5621):990-994. doi:10.1126/science.1082609

748   84.   Lebreton G, Géminard C, Lapraz F, et al. Molecular to organismal chirality is induced by
749   the conserved myosin 1D. *Science (80-)*. 2018;362(6417):949-952.
750   doi:10.1126/science.aat8642

751   85.   Tamada A, Kawase S, Murakami F, Kamiguchi H. Autonomous right-screw rotation of
752   growth cone filopodia drives neurite turning. *J Cell Biol*. 2010;188(3):429-441.
753   doi:10.1083/jcb.200906043

754   86.   Heacock AM, Agranoff BW. Clockwise growth of neurites from retinal explants. *Science*.
755   1977;198(4312):64-66. doi:10.1126/science.897684

756   87.   Romijn HJ, Mud MT, Wolters PS, Corner MA. Neurite formation in dissociated cerebral
757   cortex in vitro: evidence for clockwise outgrowth and autotopic contacts. *Brain Res*.
758   1980;192(2):575-580. doi:10.1016/0006-8993(80)90910-5

759   88.   Chi JG, Dooling EC, Gilles FH. Gyral development of the human brain. *Ann Neurol*.
760   1977;1(1):86-93. doi:10.1002/ana.410010109

761   89.   Dubois J, Benders M, Cachia A, et al. Mapping the early cortical folding process in the
762   preterm newborn brain. *Cereb Cortex*. 2008;18(6):1444-1454. doi:10.1093/cercor/bhm180

763   90.   Kasprian G, Langs G, Brugger PC, et al. The prenatal origin of hemispheric asymmetry:
764   An in utero neuroimaging study. *Cereb Cortex*. 2011;21(5):1076-1083.
765   doi:10.1093/cercor/bhq179

766   91.   de Kovel CGF, Lisgo S, Karlebach G, et al. Left–Right Asymmetry of Maturation Rates in
767   Human Embryonic Neural Development. *Biol Psychiatry*. 2017;82(3):204-212.
768   doi:10.1016/j.biopsych.2017.01.016

769   92.   de Kovel CGF, Lisgo SN, Fisher SE, Francks C. Subtle left-right asymmetry of gene

770   expression profiles in embryonic and foetal human brains. *Sci Rep*. 2018;8(1):1-13.
771   doi:10.1038/s41598-018-29496-2

93. Best CT. The emergence of cerebral asymmetries in early human development: A literature review and a neuroembryological model. *Brain Lateralization Child Dev Implic*. 1988;88:5-34.

94. Vingerhoets G, Li X, Hou L, et al. Brain structural and functional asymmetry in human situs inversus totalis. *Brain Struct Funct*. 2018;223(4):1937-1952. doi:10.1007/s00429-017-1598-5

95. Kennedy DN, O'Craven KM, Ticho BS, Goldstein AM, Makris N, Henson JW. Structural and functional brain asymmetries in human situs inversus totalis. *Neurology*. 1999;53(6):1260-1265. doi:10.1212/wnl.53.6.1260

96. Schuler AL, Kasprian G, Schwartz E, et al. Mens inversus in corpore inverso? Language lateralization in a boy with situs inversus totalis. *Brain Lang*. 2017;174:9-15. doi:10.1016/j.bandl.2017.06.005

97. Ihara A, Hirata M, Fujimaki N, et al. Neuroimaging study on brain asymmetries in situs inversus totalis. *J Neurol Sci*. 2010;288(1-2):72-78. doi:10.1016/j.jns.2009.10.002

98. Gerrits R, Vingerhoets G. Brain functional segregation, handedness and cognition in situs inversus totalis: A replication study. *Neuropsychologia*. 2023;191:108731. doi:10.1016/j.neuropsychologia.2023.108731

99. Tanaka S, Kanzaki R, Yoshibayashi M, Kamiya T, Sugishita M. Dichotic listening in patients with situs inversus:Brain asymmetry and situs asymmetry. *Neuropsychologia*. 1999;37(7):869-874. doi:10.1016/S0028-3932(98)00144-4

100. Vingerhoets G, Gerrits R, Karlsson EM. Brain (Yakovlevian) torque direction is associated with volume asymmetry of the intracranial transverse sinuses : evidence from situs inversus totalis. Published online 2024:1461-1470.

101. Kawakami R, Dobi A, Shigemoto R, Ito I. Right isomerism of the brain in inversus viscerum mutant mice. *PLoS One*. 2008;3(4):1-6. doi:10.1371/journal.pone.0001945

102. Guadalupe T, Mathias SR, vanErp TGM, et al. Human subcortical brain asymmetries in 15,847 people worldwide reveal effects of age and sex. *Brain Imaging Behav*. 2017;11(5):1497-1514. doi:10.1007/s11682-016-9629-z

103. Eichert N, Robinson EC, Bryant KL, et al. Cross-species cortical alignment identifies different types of anatomical reorganization in the primate temporal lobe. *Elife*. 2020;9:1-30. doi:10.7554/eLife.53232

104. Beauchamp A, Yee Y, Darwin BC, Raznahan A, Mars RB, Lerch JP. Whole-brain comparison of rodent and human brains using spatial transcriptomics. *Elife*. 2022;11:1-24. doi:10.7554/eLife.79418

105. Kenney M, Vasylieva I, Hood G, et al. The Brain Image Library: A Community-Contributed Microscopy Resource for Neuroscientists. *bioRxiv Prepr Serv Biol*. Published online 2024. doi:10.1101/2023.12.22.573024

809   106.   Oh SW, Harris JA, Ng L, et al. A mesoscale connectome of the mouse brain. *Nature*.
810   2014;508(7495):207-214. doi:10.1038/nature13186

811   107.   Hawrylycz M, Martone ME, Ascoli GA, et al. A guide to the BRAIN Initiative Cell Census
812   Network data ecosystem. *PLoS Biol*. 2023;21(6 June):1-30.
813   doi:10.1371/journal.pbio.3002133

814   108.   Newmaster KT, Nolan ZT, Chon U, et al. Quantitative cellular-resolution map of the
815   oxytocin receptor in postnatally developing mouse brains. *Nat Commun*. 2020;11(1):1-12.
816   doi:10.1038/s41467-020-15659-1

817   109.   Ellegood J, Anagnostou E, Babineau BA, et al. Clustering autism: using neuroanatomical
818   differences in 26 mouse models to gain insight into the heterogeneity. *Mol Psychiatry*.
819   2015;20(1):118-125. doi:10.1038/mp.2014.98

820   110.   Guma E, Bordignon P do C, Devenyi GA, et al. Early or Late Gestational Exposure to
821   Maternal Immune Activation Alters Neurodevelopmental Trajectories in Mice: An
822   Integrated Neuroimaging, Behavioral, and Transcriptional Study. *Biol Psychiatry*.
823   2021;90(5):328-341. doi:10.1016/j.biopsych.2021.03.017

824   111.   Nugent AC, Thomas AG, Mahoney M, et al. The NIMH intramural healthy volunteer
825   dataset: A comprehensive MEG, MRI, and behavioral resource. *Sci Data*. 2022;9(1):1-10.
826   doi:10.1038/s41597-022-01623-9

827   112.   Nárai Á, Hermann P, Auer T, et al. Movement-related artefacts (MR-ART) dataset of
828   matched motion-corrupted and clean structural MRI brain scans. *Sci Data*. 2022;9(1):1-6.
829   doi:10.1038/s41597-022-01694-8

830   113.   Wei D, Zhuang K, Ai L, et al. Data Descriptor: Structural and functional brain scans from
831   the cross-sectional Southwest University adult lifespan dataset. *Sci Data*. 2018;5:1-10.
832   doi:10.1038/sdata.2018.134

833   114.   Noonan MAP, Sallet J, Mars RB, et al. A Neural Circuit Covarying with Social Hierarchy in
834   Macaques. *PLoS Biol*. 2014;12(9). doi:10.1371/journal.pbio.1001940

835   115.   Hata J, Nakae K, Tsukada H, et al. Multi-modal brain magnetic resonance imaging
836   database covering marmosets with a wide age range. *Sci Data*. 2023;10(1):1-8.
837   doi:10.1038/s41597-023-02121-2

838   116.   Skibbe H, Rachmadi MF, Nakae K, et al. *The Brain/MINDS Marmoset Connectivity*
839   *Resource: An Open-Access Platform for Cellular-Level Tracing and Tractography in the*
840   *Primate Brain*. Vol 21.; 2023. doi:10.1371/journal.pbio.3002158

841   117.   Kirst C, Skriabine S, Vieites-Prado A, et al. Mapping the Fine-Scale Organization and
842   Plasticity of the Brain Vasculature. *Cell*. 2020;180(4):780-795.e25.
843   doi:10.1016/j.cell.2020.01.028

844   118.   Huang CC, Rolls ET, Feng J, Lin CP. An extended Human Connectome Project
845   multimodal parcellation atlas of the human cortex and subcortical areas. *Brain Struct*
846   *Funct*. 2022;227(3):763-778. doi:10.1007/s00429-021-02421-6

847   119.   Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and
848   surface reconstruction. *Neuroimage*. 1999;9(2):179-194. doi:10.1006/nimg.1998.0395

849   120.   Glasser MF, Coalson TS, Robinson EC, et al. A multi-modal parcellation of human
850   cerebral cortex. *Nature*. 2016;536(7615):171-178. doi:10.1038/nature18933
851
852
853

Figures and Tables

Figure 1. Reproducible Region-wise Asymmetry Patterns in the Adult Mouse Brain.

Figure 1. **A)** Experimental workflow. **B)** Matrix of region-wise asymmetry comparisons between cohorts for all brain regions. Cell value is Pearson correlation between the 2 cohorts for asymmetry of all regions. Second line in cell indicates significance of the correlation. **C)** Distribution of Pearson R values from sub-sampling 20 animals and comparing the sub-sampled asymmetry pattern with either the full cohort (AMBCA,BICCN,POND), or the remaining 21 animals (Guma2021). Dashed line is largest non-diagonal value in B). **D)** Matrix of region-wise asymmetry comparisons between datasets using top 50 leftward and top 50 rightward regions. Cell value is the median of differences (Hodges-lehmann estimator). Top regions are defined in the cohort along the row and then evaluated using the cohort in the column. Gold rectangle is example shown in E). **E)** Example of top regions categorical scatter plot. Each point is 1 brain region. Top regions were defined using the POND dataset and evaluated using the BICCN dataset. Grey value indicates the median asymmetry in the evaluation cohort for that set of brain regions. **F)** Leave-one-out analysis where title of plot indicates which cohort was left-out and used as the evaluation cohort. Top regions were defined using each 5-cohort average. **G)** Evaluation of 6-cohort average in previously un-tested Lopes2023 cohort (n = 48 animals). **H)** Montage of top 100 regional asymmetries in the 6-cohort average pattern. **D-G)** Statistics are Wilcoxon rank-sum test. * is p < 0.05; ** is p < 0.01; *** is p < 0.001; **** is p < 0.0001.

Figure 2. Same Jacobian Maps Reveal Highly Reproducible and Established Sex Differences.

Figure 2. A) Experimental overview. **B)** Matrix of region-wise sex-difference comparisons between cohorts for all regions. **C)** Matrix of region-wise sex-difference comparisons between cohorts using top 50 male-larger and top 50 female-larger regions. Value is median of differences (Hodges-lehmann estimator). Top regions are defined from the cohort along the row and are then evaluated using the cohort in the column. Gold rectangle is example in D. **D)** Example of top regions categorical scatter plot highlighted by gold rectangle in C**.** Each point is 1 brain region. Top regions were defined using the POND dataset and evaluated using the BICCN dataset. Values indicates the median sex difference in the evaluation cohort for that set of brain regions. **E)** Leave-one-out analysis where title of plot indicates which cohort was left-out and used as the evaluation cohort. Top regions were defined using each 5-cohort average. **F)** Volcano plot of significant sex differences using 6-cohort average. **G)** Montage of top regional sex differences in the 6-cohort average pattern. **H-I)** Comparison of this publication's AMBCA **H)** sex differences and **I)** asymmetries to results of Elkind et al. 2023, who used the AMBCA as well. **C-E, H-I)** Statistics are Wilcoxon rank-sum test. * is p < 0.05; ** is p < 0.01; *** is p < 0.001; **** is p < 0.0001.

Figure 3. Asymmetries Are Not Due to Technical Artefacts or Biases.

Figure 3. A) Comparison of BICCN region-wise asymmetry pattern between original and left-right reversed images. Each point is a brain region where the value is the mean asymmetry for that region across all animals in each analysis. **B-C)** Volcano plot of asymmetry for BICCN cohort where each point is a brain region. **B)** Analysis using normal images (same as **Fig S1B**). **C)** Analysis using synthetically symmetrized samples. **D)** Comparison of STPT-specific average asymmetry pattern (X-axis) in the MRI-specific average (Y-axis) using top regions. **E)** Comparison of MRI-specific average asymmetry pattern (X-axis) in STPT-specific average (Y-axis) using top regions. **D-E)** Statistics are Wilcoxon rank-sum test. *** is p < 0.001. **F)** Comparison of region-wise asymmetry pattern in 6-cohort average between ANTsPy and Elastix.

Figure 4. Characterization of Volume Asymmetry Pattern in the Adult Mouse Brain.

B   Enrichment of Anatomical Labels in Top Regions

Division	Odds-Ratio in Top 100	Nominal Composite Asymmetry (%)
Olfactory Areas	1.49	0.11
Isocortex	1.17	0.04
Hippocampal formation	0.42	0.13
Striatum	1.11	0.5
Thalamus	0.61	0.25
Hypothalamus	0.33	0.06
Midbrain	0.68	0.10
Pons	7.42*	-0.59
Medulla	0.92	-0.54
Cerebellum	5.07*	-0.30
White Matter	0.49	-0.06
Ventricular system	0.86	0.80

C   Enrichment of Isocortical Layers in Top Regions

Isocortical Layers	Odds-Ratio in Top 100	Nominal Composite Asymmetry (%)
Isocortex Layer 1	0.77	-0.03
Isocortex Layer 2/3	0.77	0.10
Isocortex Layer 4	0.56	0.07
Isocortex Layer 5	2.32	0.11
Isocortex Layer 6a	1.22	-0.11
Isocortex Layer 6b	1.22	-0.29

D   Enrichment of Functionally Asymmetric Regions

	Odds-Ratio in Top 100	Nominal Composite Asymmetry (%)
Functionally Asymmetric Regions	1.03	N/A

Figure 4. A) Bar plot of the full Allen CCFv3 where the height of each bar along the X-axis corresponds to the nominal asymmetry in the 6-cohort average. Row of sparse ticks at approximately Y = -3 indicates whether the structure is present in set of top regions derived from the 6-cohort average: presence of tick mark indicates presence of region in top asymmetries, absence of tick mark indicates absence from top asymmetries. Color bar at approximately Y = -4.5 illustrates the broader division of tissue to which that region belongs. **B-D)** Statistical enrichment of regions within **B)** various broad anatomical divisions, **C)** Isocortical layers, or **D)** regions with established functional asymmetries in the set of the top 100 asymmetric brain regions. Nominal asymmetry of the composite structure is included as well. * indicates a significant enrichment of pons and cerebellum labels in the set of top regions at FDR < 0.05. All others are non-significant. N/A indicates not applicable.

Figure 5. Anterior-Posterior Pattern of Volume Asymmetry is Driven by SA But Not CT.

Figure 5. A-B) Average asymmetry of each brain region derived from the 6-cohort average plotted against the **A)** anterior-posterior or **B)** dorsal-ventral position of that region in the atlas. **C)** Illustration of Laplacian Method for calculating cortical thickness, during which surface area can also be collected from the "Isocortex Surface". The cortex is modeled as 3 surfaces: the pial surface ("isocortex surface") or the white matter-grey matter interface (blue line overlaying corpus callosum), and a special cortical geometry surface (yellow line by piriform cortex). A smooth Laplacian field is generated between the pial surface and white matter-grey matter interface. The steepest descent path through the field (streamlines) is generated from each surface pixel to the white matter-grey matter interface. **D)** Average cortical thickness per cortical territory from 6-cohort average (all layers collapsed) calculated using method in C). **E)** Average pial surface area per cortical territory from 6-cohort average. **F)** Comparison of 6-cohort average volume asymmetry per cortical territory (all layers collapsed) measured using original Jacobian-derived volume analysis with volume asymmetry calculated through separate cortical thickness and surface area measurements. In all graphs, each point is 1 brain region and the value is the mean asymmetry for the appropriate metric calculated in the 6-cohort average. All statistics are the significance of the Pearson correlation.

Figure 6. Positional Asymmetries But No Cerebral Petalia in Mice.

Top Regions: Leave-one-out Cross-Validation

Figure 6. A-B) Cohort-specific petalia measurements for the **A)** frontal (anterior) or **B)** occipital (posterior) poles in humans (n = 4 cohorts, ~1400 individuals), mice (n = 6 cohorts, ~3500 individuals), or non-human primates (NHPs; n = 6 cohorts; 2 cohorts each for macaques, chimpanzees, and marmosets; ~1000 individuals). Each point on the graph is the average protrusion of subjects in that cohort, normalized against the length the left-hemisphere. A value of 0.2% means that the protrusion asymmetry is 0.2% the length of the left hemisphere on average. **C)** Leave-one-out cross-validation for the top positional asymmetries in all 6 cohorts. **D)** Anterior-posterior positional asymmetries derived from the mouse 6-cohort average, using the top 100 positional asymmetries. Regions that are shifted anteriorly on the right (positive values) tend to be found in the more anterior portions of the brain ("hemisphere"; red), while regions that are shifted anteriorly on the left (negative values) tend to be found in more posterior portions of the brain ("caudal brain"; black). Each point is a brain region and its color correspond to the definitions in E). **E)** Summary of positional asymmetries detected in C). "Caudal Brain" (black) is defined as midbrain, hindbrain, and cerebellum while everything anterior or dorsal to the midbrain is "hemisphere" (red). The "hemisphere" appears shifted anteriorly on the right while this trend appears opposite in the "caudal brain". Value in parentheses are the Z-axis coordinates in pixels for anterior (0) and posterior (12900) regions. *** is p < 0.001.

Figure 7. Cross-Species Comparisons of Volume Asymmetry

Figure 7. **A)** Top: 3D renderings of mouse, marmoset, macaque, and human brains from their respective 3D average templates. Middle: depiction of nominal volume asymmetry pattern for all regions in each atlas (regardless of significance), averaged across each cohort for that species. A minor brightness boost is added to better visualize the images. Bottom: display range of asymmetries shown for each species. **B)** The value in each cell and its color are the average cross-cohort consistency seen within a given species. Humans are the average of all 7 possible between-cohort combinations. Mice are similar but for 6 cohorts. Macaques and Marmosets have only 2 cohorts each, so the value indicates the lone Pearson correlation between the 2 cohorts in those species. **C-D)** Scatter plots of matched brain regions between species using the Garin et al. 2022 cross-species atlas. **C)** Cytoarchitecturally-matched (Brodmann) regions or **D)** summary values for whole cortical lobes and broad subcortical structures. Each species-pair has a different number of annotated regions which can be matched, hence the different number of data-points for each plot. Each point is a brain region. The value of the X- and Y-axes respectively indicate the cross-cohort average percent volume asymmetry observed in that species at that region. Human asymmetry data is only available for cortical regions.