Chromosome structural rearrangements in invasive haplodiploid ambrosia beetles

revealed by the genomes of *Euwallacea fornicatus* and *Euwallacea similis* (Coleoptera,

Curculionidae, Scolytinae)

Authors: James R. M. Bickerstaff[1]*, Tom Walsh[2], Leon Court[2], Gunjan Pandey[2], Kylie Ireland[3, 4,]

David Cousins[3], Valerie Caron[5,6], Thomas Wallenius[7], Adam Slipinski[1], Rahul Rane[5], Hermes E.

Escalona[1]*

[1] Australian National Insect Collection, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[2] Environment, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[3] Department of Primary Industries and Regional Development, 3 Baron-Hay Court, South Perth,

Western Australia 6151, Australia

[4] Department of Biodiversity, Conservation and Attractions, Perth Western Australia 6151, Australia

[5] Health and biosecurity, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[6] Faculty of Science and Technology, University of Canberra, Bruce, ACT 2615

[7] Department of Agriculture, Fisheries and Forestry, GPO Box 858, Canberra ACT 2601, Australia

Emails: James Bickerstaff: james.bickerstaff@csiro.au

Tom Walsh: tom.walsh@csiro.au

Leon Court: leon.court@csiro.au

Gunjan Pandey: gunjan.pandey@csiro.au

Kylie Ireland: kylie.ireland@dbca.wa.gov.au

David Cousins: david.cousins@dpird.wa.gov.au

Valerie Caron: Valerie.caron@csiro.au

Thomas Wallenius: Thomas.Wallenius@aff.gov.au

Adam Slipinski: adam.slipinski@csiro.au

Rahul Rane: Rahul.rane@csiro.au

Hermes Escalona: Hermes.escalona@csiro.au

*Correspondence address: James Bickerstaff, E-mail: james.bickerstaff@csiro.au; Hermes Escalona:

hermes.escalona@csiro.au

Abstract

Background. Bark and ambrosia beetles are among the most ecologically and economically damaging introduced plant pests worldwide, with life history traits including polyphagy, haplodiploidy, inbreeding polygyny and symbiosis with fungi contributing to their dispersal. Species vary in host tree ecologies, with many species attacking stressed or recently dead trees, such as the globally distributed *E. similis* (Ferrari). At the other end of the spectrum, the Polyphagous Shot Hole Borer (PSHB), *Euwallacea fornicatus* (Eichhoff), attacks over 680 host plants and is causing considerable economic damage in several countries worldwide. Despite their notoriety, publicly accessible genomic resources for *Euwallacea* Hopkins species are scarce, hampering better understanding of their invasive capabilities as well as modern control measures, surveillance and management.

Findings. Using a combination of long read and short read sequencing we assembled and annotated high quality (BUSCO > 98% complete) chromosome level genomes for these species. Comparative macro-synteny analysis showed that chromosome number increases in haplodiploid inbreeding species of *Euwallacea* due to fission events, reducing synteny with outbred diploid species. This suggests that life history traits can impact chromosome structure. Further, the genome of *E. fornicatus* had a higher relative proportion of repetitive elements, up to 17% more, than *E. similis*. We used the genome assemblies to identify the haplotype of the invasive population in Perth, Western Australia. Additionally, metagenomic analyses identified microbiota associated with both species, with fewer taxa found in association with *E. fornicatus* compared to *E. similis.*

Conclusions. Comparative analyses between closely related species with varying host tree ecologies and invasive capabilities will shed light into the molecular underpinning of species invasion. Moreover, these haplodiploid genomes will also contribute to the understanding of how life history traits contribute to their genome evolution.

Keywords: Invasive species, bark beetles, biosecurity, weevils, insect genomics, Polyphagous Shot Hole Borer

Background

Bark and ambrosia beetles (Curculionidae, Scolytinae) are some of the most ecologically damaging and economically important insect plant pests, contributing to tree mortality through infestation and the spread of pathogenic fungi (Biedermann et al., 2019; Raffa et al., 2008). Most bark beetles are phloeophagous and obtain nutrition from the phloem tissue while some lineages, termed ambrosia beetles, are xylomycetophagous and consume farmed symbiotic ambrosia fungi (Hulcr & Stelinski, 2017; Jordal, 2014). Many species are commonly transported to novel areas through human mediated

movement of wood and plant products and over one hundred species have established outside of their native range (Brockerhoff et al., 2006; Lantschner et al., 2020; Meurisse et al., 2019; Rassati et al., 2015). Life history traits including polyphagy, haplodiploidy, inbreeding polygyny and symbiosis with ambrosia fungi has been noted to contribute to the establishment of introduced species (Jordal et al., 2001; Lantschner et al., 2020; Vilardo et al., 2022), all of which characterise members of the tribe Xylebroini. Consequently, many such species have a global native or introduced distribution and are high priority pests in many regions of the world.

The genus *Euwallacea* Hopkins include several well studied polyphagous tree pests (Cognato et al., 2015; Kasson et al., 2013; O'Donnell et al., 2016). The genus is composed of over 80 species which are mostly circumtropical in their distribution (Storer et al., 2015). Species vary in their ecological relationships with host trees, with many species attacking stressed, cut and recently dead hosts. One such example is *E. similis* (Ferrari) (Figure 1b), a species native to South East Asia and possibly to Australia (Beaver et al., 2014), that has been introduced to the Americas and tropical African countries (Gomez et al., 2018) with 145 known host plants (Browne, 1961; Ruzzier et al., 2023). *Euwallacea fornicatus* (Eichhoff) (Figure 1a)*, also known as the Polyphagous Shot Hole Borer (PSHB) is considered one the most damaging Xyleborini species as it can infest healthy live host trees (Cook & Broughton, 2023; De Wit et al., 2022). It has been recorded to attack over 680 tree species across several plant families (Department of Primary Industries and Regional Development, Government of Western Australia, 2024; Gomez et al., 2019; Mendel et al., 2021) and is able to successfully reproduce in 168 species (Department of Primary Industries and Regional Development, Government of Western Australia, 2024 (and references therein), Mendel et al., 2021). *Euwallacea fornicatus* is widely distributed and has been introduced to California (Eskalen et al., 2012, 2013), Hawaii (Rugman-Jones et al., 2020), South Africa (Paap et al., 2018), Israel (Mendel et al., 2012), Europe (Schuler et al., 2023) and, recently, in the city of Perth, Western Australia (WA) where it is currently subject to an eradication campaign (Department of Agriculture, Fisheries and Forestry, 2024). This ambrosia beetle species is associated with several pathogenic *Fusarium* fungi, including *F. euwallaceae* Freeman *et al*. 2013 and *F. sp.* AF-18, and these fungi can induce wilt and dieback in host trees leading to tree mortality (Freeman et al., 2013). Introduced populations of *E. fornicatus* cause significant economic and ecological damage to managed and native ecosystems, and agricultural areas (Cook & Broughton, 2023; De Wit et al., 2022; Smith & Hulcr, 2015). Economic modelling has estimated the damages caused by this species to \$18.45 billion over a ten-year period in South Africa (De Wit et al., 2022), while in Perth WA the cost of eradicating this species is estimated to be AUD\$45 million and management of the species could reach AUD\$6.8 million per annum in 30 years (Cook & Broughton, 2023).

The species *E. fornicatus* has been associated with three other species (Gomez, et al., 2018; Smith et al., 2019) representing a species complex with similar morphology, which include *E. perbrevis* (Schedl), the Tea Shot Hole Borer (TSHBa), *E. fornicatior* (Eggers), also the Tea Shot Hole Borer (TSHBb) and *E. kuroshio* (Gomez and Hulcr), the Kuroshio Shot Hole Borer (KSHB) (Smith et al., 2019). Originally, these four species were all considered to be *E. fornicatus*, however, recent molecular phylogenetic and taxonomic works have stabilised their species status (Gomez, et al., 2018; Smith et al., 2019; Stouthamer et al., 2017). Few genomic resources exist for *E. fornicatus* and other Xyleborini, with much of the data available restricted to mitochondrial *cytochrome oxidase I* (*COI*) and a handful of genes for phylogenetic analyses (Bierman et al., 2022; Gomez et al., 2018; Stouthamer et al., 2017). Mitochondrial *COI* phylogenies have identified 21 haplotypes of *E. fornicatus* with much of the diversity found in Vietnam and Taiwan (Stouthamer et al., 2017). One haplotype, H33, has been identified in California, South Africa and Israel and is thought to have originated from Vietnam (Bierman et al., 2022; Stouthamer et al., 2017), however, other haplotypes, H35 and H38, have also been found in California and South Africa, respectively, indicating that the species could have been introduced multiple times to both countries (Bierman et al., 2022; Stouthamer et al., 2017). To date, the only sequence data available for *E. similis* is restricted to *COI* barcodes in its native South East Asian distribution (Lynn et al., 2021) Beyond this, little genomic data exists for these widespread pests and further molecular resources are vital to understand the invasion pattern of *Euwallacea* species, their invasive capabilities, and finding novel ways of controlling outbreaks and mitigating future harm.

Access to high quality genomes has led to advancements in many fields, including conservation management (Formenti et al., 2022; Hogg, 2023; Theissinger et al., 2023), biodiversity genomics and monitoring (Breed et al., 2019; Cordier et al., 2021; Mc Cartney et al., 2023; The Darwin Tree of Life Project Consortium et al., 2022), and species invasion research and management (McGaughran et al., 2021, 2024; North et al., 2021). In the management of invasive species, genomic resources are facilitating accurate and time-efficient diagnostics and surveillance (Chua et al., 2023; Frey et al., 2022; Gardy & Loman, 2018; Matheson & McGaughran, 2022; McGaughran et al., 2024), understanding the origin of invasive species, invasion pathways (Atsawawaranunt et al., 2023; Liu et al., 2022; Rane et al., 2023; Zhang et al., 2023), identifying genetic (Jin et al., 2023; Parvizi et al., 2023; Yin et al., 2021) and genome structural (Stapley et al., 2015; Tepolt et al., 2022) variation related to invasion potential (Camus et al., 2024; McGaughran et al., 2024; Sillero et al., 2020). These resources are also accelerating the ways in which invasive species are controlled, such as through chemical management (Hopkinson et al., 2023; Tay et al., 2022; Thia et al., 2023), biocontrol agents (Morrow & Riegler, 2021; Nikolouli et al., 2018; Ross et al., 2022), genome-informed population control (Burgess et al., 2022; Kennedy et al., 2018; Maselko et al., 2017), or gene-editing and gene drive (Devos et al., 2022; Hammond et al., 2016, 2021).

In this study we present chromosome-scale, annotated genomes of *E. fornicatus* and *E. similis* using

newly developed long read sequencing technologies and performed comparative analysis of

macrosynteny, transposable elements, metagenomics and nuclear and mitochondrial phylogenetics.

These genomes will be valuable assets for tracking the invasion pathway, developing novel

biosecurity management plans for these invasive species and will provide essential insight into

comparative of invasive species genomics.

Methods

Sample collection, DNA extraction and sequencing

Specimens of *E. fornicatus* were collected by the Department of Primary Industries and Regional

Development (DPIRD) WA from felled *Acer negundo* trees in Perth WA in October 2022. Male and

females were flash frozen by storing them on dry ice and shipped to CSIRO Black Mountain Canberra

Australian Capital Territory and stored at -80°C. *Euwallacea similis* larvae and adult females were

collected from a fallen *Ficus* sp. branch in Eureka, northern New South Wales in April 2023.

Specimens were transported from the field at 10°C and flash frozen by storing them at -80°C.

Vouchers of both species have been deposited in Australian National Insect Collection (ANIC) under

the collection codes ANIC-25-086063 – ANIC-25-086073.

Female specimens were used for DNA extraction due to their larger size compared to haploid males.

First, genomic DNA (for short read sequencing) of both species was extracted using a standard

Qiagen DNEasy kit approach. We then used a single specimen from each species for high molecular

weight DNA extraction with the Qiagen MagAttract kit following a modified protocol. Samples were

incubated overnight at 56°C and lysates were agitated at 300 rpm. Following this, DNA was

suspended in two elutions of 100 μl of Buffer EB. The concentration of all extracts were assessed

using a DeNovix spectrophotometer and a Qubit Fluorometer with the Qubit dsDNA High Sensitivity

Assay kit. Fragment size distribution and genomic DNA was assessed on an Aligent Fragment

Analyzer system with the high sensitivity genomic DNA kit DNF-488.

For short read sequencing, sequencing libraries with 150 bp paired end reads were prepared and

sequenced on the Illumina NovaSeq 6000 system at Genewiz (Suzhou, China). For long read

sequencing, high molecular weight (HMW) DNA extracts were sequenced on the PacBio HiFi

circular consensus Revio system at the AGRF in Brisbane, Queensland. For two HMW extracts of

each species, gDNA was mechanically sheared, followed by SMRTbell Express 2.0 library

construction. Short DNA fragments were removed using BluePippin. Barcoded libraries were then

sequenced using one SMRT cell for each species on the PacBio Revio system; the libraries for *E.*

fornicatus are herein referred to as EFF26 and EFF41, and the libraries for *E. similis* are herein

referred to as ESF13 and ESF15.

Genome size estimation, assembly and scaffolding

First, genome size and heterozygosity estimates were performed with all sequence reads; kmers were

counted with jellyfish (Marçais & Kingsford, 2011) and genome size and heterozygosity were

estimated with GenomeScope v2 (Ranallo-Benavidez et al., 2020; Vurture et al., 2017). HiFi reads

were then filtered for HiFi adapters with HiFiAdapterFilt (Sim et al., 2022) and subsampled with

seqtk into randomly sampled subsets, ranging from 12x, 15x, 20x, 35x and 50x, of the estimated

genome size, subsets for all four libraries. Contigs were assembled with these subsets and the full

dataset using HiCanu v2.1.1 (Nurk et al., 2020) with the genome size estimates of 190 Mb for both

EFF26 and EFF42, and 170 Mb for both ESF13 and ESF15. Contigs were also assembled with

hifiasm v0.19.6-r595 (Cheng et al., 2021) using the -l0 flag to disable purge duplication as these

genomes are highly homozygous and presumably inbred. Misassembled contigs were assessed and

corrected with CRAQ v1.0.9a (Li et al., 2023). Haplotigs and contig overlaps were removed with

purge_dups v1.2.5 (Guan et al., 2020). The quality of the corrected contig assemblies were assessed

with QUAST v5.2.0 (Mikheenko et al., 2018) and BUSCO v5.5.0 (Manni et al., 2021) using the

endopterygota_odb10 dataset and the tribolium2012 dataset was specified for Augustus (Stanke et al.,

2008) training.

Proximity ligation information was generated with Arima HiC 2.0 kits using tissue from a single

individual from both species at the JCMR BRF ANU. In short, tissues were fixed in formaldehyde,

DNA was extracted and then digested with Arima restriction enzymes. Overhangs of digested DNA

were then end repaired and marked with biotinylated nucleotides. The ends were then ligated together,

and DNA strands were de-crosslinked and sheared for Illumina PE 150bp sequencing. The *E.*

fornicatus library was sequenced on an Illumina NovaSeq 6000 for 300 cycles and the library for *E.*

similis was sequenced on an Illumina NextSeq 2000 for 300 cycles.

The Hi-C reads were mapped to draft contig assemblies following the Arima Genomics Mapping

Pipeline v03. First, 5' ends of the Hi-C reads were trimmed by 5 base pairs to remove over-

represented 3bp barcode UMI sequences and 2 dark bases from the 5' end. Trimmed reads were then

mapped independently (as single-ended reads) using BWA-MEM2 (Vasimuddin et al., 2019) with the

-5SP flag. Only the 5' ends were retained using the "filter_five_end.pl" script provided by the Arima

Genomics Mapping Pipeline. As the ends of the DNA were ligated together prior to sequencing, 3'

ends can originate from the same contiguous strands of DNA as the 5' end of the mate-pair, which can

manifest a ligation function and form chimeric reads after mapping. Following this, single-ended

mapped reads were then paired using the "two_read_bam_combiner.pl" script provided by the Arima

Genomics Mapping Pipeline and mapped reads with a mapping quality (MAPQ) < 10 were discarded

and sorted with samtools (Danecek et al., 2021). Lastly, PCR-duplicates were removed with Picard

Tools v3. Assemblies were then scaffolded using YahS v1.1a (Zhou et al., 2023) and assessed with

QUAST v5.2.0, and BUSCO v5.5.0 (Manni et al., 2021). We then compared the best scaffolded

assembly across the subset reads with an analogous scaffolding run using SALSA2 v2.3 (Ghurye et

al., 2019) and selected the best scaffolds for downstream curation using QUAST and BUSCO metrics,

and Hi-C mapping with the Juicebox Assembly Tools (JBAT) v1.b (Dudchenko et al., 2018; Durand

et al., 2016). Contaminant scaffolds, defined as contigs identified not to be of Arthropoda origin

through BLAST (Altschul, 1997), sequences with >15x coverage, and GC <0.3 and >0.45 were

removed using the BlobToolKit v4.2.1 (Challis et al., 2020). Hi-C reads were then mapped to the

decontaminated scaffolds using BWA-MEM2 following the Arima Genomics Mapping Pipeline v03

and were reviewed with JBAT v1.b (Dudchenko et al., 2018; Durand et al., 2016). Illumina short read

and PacBio HiFi reads were also mapped to the decontaminated scaffolds and all three mapped

datasets were inspected using JBrowse v2.10.3 to identify any potential misassembles (identified by

poor read alignments and regions of low coverage). We assessed the completeness of the final

scaffolded assemblies again with QUAST v5.2.0 (Mikheenko et al., 2018) and BUSCO v5.5.0 (Manni

et al., 2021). Lastly, the scaffolds of each library of the same species (i.e. both *E. fornicatus* and *E.*

similis assemblies) were renamed and reoriented to be syntenic with one another using BUSCOs to

define synteny and orientation. We identified pseudo-chromosome scaffolds by scaffolds that were >

500Kb in length and had an average Hi-C coverage > 30x. Pairs of syntenic pseudo-chromosomes

were defined by length and by BUSCO composition and these pseudo-chromosomes were named in

order of length by the EFF26 assembly for *E. fornicatus* and the ESF13 assembly for *E. similis*.

Lastly, assemblies were assessed for any remaining contamination with FCS-GX v0.5 (Astashyn et

al., 2024) and any contaminants identified were removed.

Genome annotation

Total RNA was extracted from ten adult males and females of each species, with a further eight *E.*

forniacatus females (for long read sequencing). RNA was extracted using the Zymo Research Quick-

RNA Miniprep Plus kit following the manufacturers tough-to-lyse sample preparation procedure.

RNA quantity and quality was then assessed using a DeNovix spectrophotometer and Aligent

Fragment Analyzer system with the HS RNA kit DNF-472. For one extract of each species and sex

strand-specific libraries were prepared by GENEWIZ from Azenta Life Sciences and sequenced on an

Illumina NovaSeq 6000 PE150. Libraries of the pooled eight *E. fornicatus* female RNA extract were

prepared for long-read sequencing following the ONT PCR-cDNA Barcoding Kit SQK-PCB111.24

by the BRF-JCMSR ANU. Libraries were sequenced on an ONT PromethIon across two FLO-

PRO002 R9.4.1 flow-cells. Basecalling of the fast5 reads to fastq was performed with the Guppy

basecaller (RRID:SCR_023196) v7.1.4.

Repeat annotation and masking was performed with EarlGreyTE (Baril et al., 2024). This pipeline

runs RepeatModeler (Flynn et al., 2020) to identify *de novo* Transposable Element (TE) identification

and automated TE curation pipelines are implemented to refine repeat element annotation. TEstrainer

is used to generate maximum length *de novo* TE consensus sequences following a BLAST, Extract,

Align, Trim process. This *de novo* TE library is then used to annotate the genome with RepeatMasker

(Smit et al., 2013) using sensitive search settings and then full length LTRs are identified with

LTR_finder (Xu & Wang, 2007). The TE annotations from the RepeatMasker run are then

defragmented and combined with results from LTR_finder using RepeatCraft (Wong & Simakov,

2019). Overlapping TE annotations are then removed using the EarlGreyTE custom R script.

Following repeat annotation, kimura-2 divergences between TE classes were calculated using the

parseRM.pl scrip with the -l flag.

RNA-seq reads were filtered for low quality reads and adapters were removed with fastp (Chen et al.,

2018) and the cleaned reads were mapped to the masked genome with hisat2 (Kim et al., 2019).

Mapped reads were then sorted and indexed with samtools (Danecek et al., 2021). A protein database

was constructed with sequences from the OrthoDB v11 Arthropoda dataset and annotations of the

Dendroctonus ponderosae (Hopkins)*, D. valens* (LeConte)*, Hypothenemus hampei* (Ferrari), and

Taphrorychus bicolor (Herbst) genome assemblies. Mapped RNA reads and the protein database were

used to predict genes in repeat masked assemblies using BRAKER3 (Gabriel et al., 2023).

Completeness of the annotated gene models was then assessed with BUSCO v5.5.0 (Manni et al.,

2021), run in protein mode using the Endopterygota_obd10 dataset and the tribolium2012 dataset was

specified for Augustus (Stanke et al., 2008) training. Lastly, genes were characterised for their

putative function using eggNOG-mapper v2.1.6 (Cantalapiedra et al., 2021).

Transposable Element and chromosomal comparisons across Scolytinae

We explored repeat elements, macro-syntenic linkages and chromosomal rearrangements in an

evolutionary framework across the Scolytinae. For this purpose, the, chromosome level genomes of *T.*

bicolor (NCBI), *D. valens* (Liu et al., 2022), *D. ponderosae* (Keeling et al., 2022)*, Ips nitidus* (Eggers)

(Z. Wang et al., 2023) and the Entiminae outgroup *Polydrsus cervinus* (L.) (Barclay et al., 2023) were

downloaded from NCBI or ensemble. To compare the variation of repeat elements and macro-synteny

associated with reproductive ecology within genera we included both *Dendroctonus* Erichson

assemblies to directly compare with *Euwallacea* assemblies. First, repeat annotation of downloaded

genomes were performed with EarlGreyTE (Baril et al., 2024) as above. We then calculated the

relative proportion of repetitive elements within the total assembly and pseudochromosomal regions

to compare TEs across assemblies in R (R Core Team, 2013) with dplyr and visualised with ggplot2

(Wickham et al., 2019).

Following this, the composition of TEs were compared across Scolytinae genomes downloaded and

the *Euwallacea* reference genomes (EFF26 & ESF13). We used BUSCO to call 2,108 single copy

sequences which were aligned with MAFFT v7.490 (Katoh, 2002) and poorly aligned bases were

removed with TrimAl v1.4.1r22 (Capella-Gutiérrez et al., 2009). Alignments were then concatenated

with phykit v1.19.0 (Steenwyk et al., 2021) and phylogenetic analyses were conducted in IQ-tree

v2.3.2 (Minh et al., 2020). Substitution models were estimated with ModelFinder Plus using a greedy

strategy with gene merging (Chernomor et al., 2016; Lanfear et al., 2012) and bootstrap support was

calculated with nonparametric bootstrap (Felsenstein, 1985) with 100 replicates and the phylogeny

drawn in ggtree (Xu et al., 2022).

Lastly, odp (Schultz et al., 2023) was used to explore the macrosynteny and chromosomal variation

across Scolytinae genomes downloaded and the *Euwallacea* reference genomes (EFF26 & ESF13).

Previously used BUSCO sequences were used to describe the relationship between linkage groups and

chromosomes between species. Reciprocal best hits between BUSCOs were identified with diamond

blastp and the homology of scaffolds between species were inferred with pairwise p-values calculated

with Fisher's exact test. Pairwise species dot plots syntenic linkage ribbon plots for all species were

generated using scripts available in odp. Linkage colours in the ribbon plot follows the Bilaterian-

Cnidarian-Sponge Linkage Groups (BCnS LGs) as in (Schultz et al., 2023).

Mitochondrial phylogenetics

To identify the haplotype of *E. fornicatus* present in Perth WA, previously published *COI* genes

(Bierman et al., 2022; Gomez et al., 2018; Stouthamer et al., 2017) of PSHB, TSHBa&b and KSHB

were downloaded from NCBI. The mitochondrial genome for all libraries were assembled using the

MitoHiFi v3.2.1 pipeline (Uliano-Silva et al., 2023) which implements hifiasm (Cheng et al., 2021) to

assemble reads that are mapped to a reference mitogenome (the *E. fornicatus* NC_059702 and *E.*

similis ON169990 mitogenome assemblies), MitoFinder (Allio et al., 2020) to annotate mitochondrial

genes and ARWEN for tRNA annotation (Laslett & Canbäck, 2008). Full length *COI* genes from

these assemblies were then aligned with the sequences from NCBI using MAFFT (Katoh, 2002).

Third codon positions were partitioned in the alignment and IQ-Tree v2.3.2 (Minh et al., 2020) was

used to estimate the phylogeny, substitution models were identified with ModelFinderPlus (Flynn et

al., 2020) and bootstrap support was calculated with UFBoot (Hoang et al., 2018) with 1000

replicates.

Metagenomics

To identify the microbial communities associated with *E. fornicatus* and *E. similis*, metagenomes

were assembled from the total hifi sequence read dataset for each library. Metagenome-assembled

genomes (MAGs) were assembled using hifiasm-meta (Feng et al., 2022) with default settings.

BLAST was used to identify contigs and MAG identity was explored and visualised with the

BlobToolKit v4.2.1 (Challis et al., 2020). Host contigs and those that were unidentified were filtered

out of the dataset. A closed *Wolbachia* contig in was found in both *E. similis* libraries, and so the

strain was identified using MLST approaches implemented in the PubMLST database (Jolley et al.,

2018).

Results

Genome assemblies and evaluation

In total 98.23Gb of high-quality long sequence reads (PacBio HiFi) with a mean N50 of 11.6Kb were

obtained to assemble four genomes of two *Euwallacea* species (Table S1). The estimated genome size

for *E. fornicatus* was between 192.7 Mb and 208.9 Mb, while estimates of 170.7 Mb and 181.8 Mb

were made for *E. similis* and all are predicted to be highly homozygous (het: 0.012% – 0.112%), and

all short read and long read libraries consistently predicted similar results (Figure S1). The four

chromosome-level genomes that were selected for downstream analysis for *E. fornicaus* and *E. similis*

were assembled with HiCanu, as we found HiCanu yielded better assemblies than hifiasm. Genomes

were assembled with, post sequence read filtering and subsampling, 6.5Gb for EFF26 (29.58x, 0.35

subsample), 7Gb for EFF42 (24.07x, 0.35 subsample), 5.63Gb for ESF13 (30.95x, not subsampled)

and 8.09Gb for ESF15 (44.78x, 0.15 subsample) of PacBio HiFi reads and 3.02Gb (*E. fornicatus*) and

3.3Gb (*E. similis*) of Arima Hi-C paired-end reads (Table S1).

Both pipelines for scaffolding, YahS and SALSA2, identified a similar number of pseudochromosome

scaffolds for the *E. fornicatus* assemblies but not for *E. similis*. In the *E. similis* scaffolds, SALSA

failed to break incorrectly joined contigs identified through Hi-C read mapping compared to YahS

(Figures S2 – S5). Scaffolding, decontamination and manual curation anchored 75% and 69% of the

total assembly sequences into 39 pseudo-chromosomes for EFF26 (Figure S6) and EFF42 (Figure

S7), respectively. While the total number of assembly sequences decreased post scaffolding and

manual curation, N50 also decreased by 0.3Mb – 1.3Mb. This was attributed to the breakup of larger

contig sequences that were not supported with Hi-C alignments during manual curation of assemblies.

Both EFF26 and EFF42 assemblies were larger than the estimated genome sizes, at 219.83 Mb and

239.6 Mb, respectively. Scaffolding, decontamination and manual curation resulted in 93% and 91%

of the total assembly sequences successfully anchored in 37 pseudochromosomes groups for ESF13

(Figure S8) and ESF15 (Figure S9), respectively. The N50 of the assembly increased post scaffolding

by 1.44 – 1.54 Mb and the total number of assembly sequences decreased. The ESF13 assembly was

smaller than the estimated genome size, at 177.6 Mb, while the ESF15 genome was larger than

estimated at 180.07 Mb. All assemblies were found to be highly complete with BUSCO, and all

genomes contained over 98% complete single copy sequences (Table 1, Figure S10).

Genome annotation

Genome annotation with BRAKER3 predicted, on average 13,570 transcripts and an average of

10,845 genes for the *E. fornicatus* genomes. Annotation for the *E. similis* genomes resulted in an

average 16,188 transcripts and 12,141 genes predicted. Protein coding genes occupied, on average, 16

Mb of the *E. fornicatus* genome (6 – 7.9% of total assembly lengths) and 18 Mb of *E. similis* (10% of

total assembly lengths). Gene predictions were largely complete, with >90% of predicted genes

assigned putative homologs with functional annotation for all annotations. Moreover, most

annotations were largely complete with >97% complete single copy BUSCOs (Table 3). The number

of genes annotated for both *Euwallacea* species were slightly lower than those annotated in

Dendroctonus assemblies (Copeland et al., 2024; Liu et al., 2022) and much lower than the

annotations of *H. hampei* (Navarro-Escalante et al., 2021), *I. nitidus* (Wang et al., 2023)*, I.*

typographus (Powell et al., 2021) and *T. bicolor*.

Mitochondrial and nuclear phylogenetics

Phylogenetic analysis of mitochondrial haplotypes identified that the population of *E. fornicatus* in

Perth WA belong to haplotype H38, with the *COI* barcode identical to native individuals in China and

Vietnam, and also to the invasive population in South Africa. The H38 haplotype clustered with

Haplotypes 37, 39 – 42, and were found to be phylogenetically distinct from the more prevalent

invasive H33 haplotype (Figure S11). Phylogenomic analysis using BUSCOs of published Scolytinae

genomes and those of the reference *Euwallacea* yielded a topology similar to that reported by Pistone

et al. (Pistone et al., 2018) and all nodes were strongly supported (Figure 2).

Repeat element and chromosomal variation across Scolytinae

Roughly half (47 – 50%) of the *E. fornicatus* assemblies and over one third (33 - 34%) of the *E.*
similis assemblies were masked by the EarlGreyTE pipeline (Table 3, Figure 2, Figure S12). DNA
transposons and long terminal repeats (LTR) were the most abundant repeat elements identified, with
unclassified elements only accounting for a small proportion of all annotations (Table 3). Few rolling
circle elements were identified in all assemblies and SINEs were only found in the *E. similis*
assemblies (Table S1). Kimura 2 divergence landscapes of repetitive elements indicate a recent burst
of DNA transposons and LTR elements, while LINE and rolling circle elements are experiencing
ongoing proliferation in the *Euwallacea* assemblies (Figures S13 – 16). The relative proportion of
annotated TEs varied throughout Scolytinae assemblies, with over 60% of the *T. bicolor* assembly
occupied by TEs, and the assembly of *D. ponderosae* possessed the lowest relative proportion of
annotated TEs (25%). The relative proportions of the types of TEs also varied significantly throughout
Scolytinae as well, and TE annotations were largely composed of unclassified elements, DNA
transposons and LINEs. All published Scolytinae assemblies possessed greater amounts of Satellite
sequences than the *Euwallacea* assemblies (Figure 2). Lastly, the Entiminae outgroup *P. cervinus* had
a larger proportion of the genome of annotated TEs compared to Scolytinae.
414
The assemblies of *E. fornicatus* and *E. similis* were more syntenic with one another than other
Scolytinae. Largely, synteny was retained throughout the Scolytinae assemblies with several
translocations of genomic material. However, extensive translocations, rearrangements, and fissions
of syntenic blocks were identified between the assemblies of *Euwallacea* and other Scolytinae.
Further, the X chromosome was identified to be synentic throughout all assemblies, however, blocks
corresponding to these scaffolds were broken apart in the *Euwallacea* assemblies (Figure 3).
421
422   *Metagenomics*
423
The *E. similis* libraries produced more MAG contigs (730 and 1,200) than the *E. fornicatus* libraries
(103 and 109). Across all libraries, many more Bacteria were identified than other microbial
Kingdoms, with Actinomycetota and Pseudomonadota highly prevalent (Figure S17). Several
complete circularised Bacteria assemblies were also identified across all libraries (Table S2),
including *Wolbachia* for both *E. similis* libraries. MLST profiling identified the *Wolbachia* strain as
closely related to *w*Ei (Kawasaki et al., 2016), and unique SNPs were found in the *hcpA* and *gatB* loci.
Consequently, this represents a novel strain, herein referred to as *w*Es. Several fungal taxa were
identified for both species (Figure S17), and mitogenomic contigs of *Fusarium* belonging to the
Euwallacea Fusarium Clade (EFC) were identified by BLAST for all libraries except for ESF13. The
only fungi found in *E. fornicatus* MAGs were *Fusarium*, while a richer diversity of fungi were found
in association with *E. similis*, with 13 genera classified including *Leptographium, Ophiostoma,*

Ceratocystis and *Sporothrix* (Table S1). Several viral contigs were also identified in both species, all

belonging to Caudoviricetes.

Discussion

Genome synteny across Scolytinae

We sequenced, assembled and annotated the genomes of *Euwallacea fornicatus* and *E. similis*. These

genomes are highly contiguous, complete and accurate as we sequenced the libraries of two female

specimens per species and employed multiple assembly and scaffolding pipelines to validate their

quality. The *E. fornicatus* assemblies presented herein were accepted and published in the NCBI

GenBank on the same day as two other contig-level assemblies of the same species from South

Africa, GCA_040114445 & GCA_040114495, from a female and male, respectively. The assembly

sizes of these contig-level genome assemblies are similar to our chromosome-scaffold assemblies,

further validating the quality of our assemblies. These genomes represent the first chromosome

scaffold genomes for these pests and also the first genomes of diploid females from a haplodiploid

inbreeding beetle lineage. They are among the smallest Scolytinae genomes published and contain the

highest number of pseudo-chromosomal scaffolds. Additionally, 17 – 81% fewer protein coding genes

were predicted for these *Euwallacea* assemblies compared to previously published Scolytinae

genomes (Copeland et al., 2024; Liu et al., 2022; Navarro-Escalante et al., 2021; Powell et al., 2021;

Wang et al., 2023).

The number of pseudo-chromosomal scaffolds identified for both *Euwallacea* species (n = 37 – 39)

were higher than the typical number of chromosomes in Scolytinae (n = 7 – 16) (Virkki & Smith,

1978). Chromosome number variation has previously been identified across and within genera of

scolytines. For example, the karyotype of all 19 *Dendroctonus* species have been characterised, with

chromosome numbers ranging from 2n = 12 – 30, and chromosomal fusion resulting in chromosome

number reduction (Keeling et al., 2022; Lanier, 1981; Six & Bracewell, 2015). Similarly,

chromosome number variation has also been identified across 32 species of *Ips*, with numbers ranging

from 2n = 16 – 32 (Virkki & Smith, 1978). In Curculionidae, a larger variation in chromosome

numbers have been characterised in those that are parthenogenetic due to polyploidy (Blackmon &

Demuth, 2015; Virkki & Smith, 1978). For example, parthenogenetic pseudogamy resulting in

triploid females have been characterised in *Ips acuminatus* Gyllenhal and *I. tridens* Mannerheim

(Lanier & Kirkendall, 1986). Similar observations have been made for several species of obligate

inbreeding haplodiploid Xyleborini (Takenouchi & Takagi, 1967; Virkki & Smith, 1978). Species of

the genus *Xylosandrus* Reitter have fewer chromosomes than other Scolytinae, with *X. germanus*

(Blandford) 2n = 8 and *X. compactus* (Eichhoff) 2n = 10 (Takenouchi & Takagi, 1967). In contrast,

the chromosomes of *Anisandrus dispar* (F.) are small in size and numerous, n = 40 (Virkki & Smith,
1978), similar to the *Euwallacea* species genomes.

We identified extensive chromosomal rearrangements between haplodiploid inbreeding and diploid
outbreeding Scolytinae. The synteny of autosomal chromosomes and the ancestrally conserved X
chromosome between diploid outbreeding species are largely conserved across genera. Most
chromosomes shared 1:1 conserved synteny and several blocks are either translocated within or across
scaffolds. Further, despite the difference in chromosomal numbers between *D. ponderosae* (2n = 12)
and *D. valens* (2n = 14) (Six & Bracewell, 2015), synteny across assemblies are largely conserved,
identified herein and previously (Copeland et al., 2024; Liu et al., 2022). Between the two
haplodiploid inbreeding *Euwallacea* species, the assemblies are less syntenic than the diploid
outbreeding species. Few 1:1 conserved syntenic chromosomes exist due to widespread translocations
and inversions of orthologous genes. Furthermore, extensive chromosomal rearrangement
characterised the variation in genomic architecture between *Euwallacea* and diploid outbreeding
species, with widespread translocation events shuffling syntenic blocks and fission events breaking up
chromosomes (including the ancestrally conserved X chromosome). Comparative analyses of
Coleopteran karyotypes have identified that small effective population sizes and genetic drift greatly
impact chromosome number variation across species (Blackmon et al., 2024). As *Euwallacea,*
obligately engage in inbreeding and haplodiploidy, species are highly homozygous, as identified
herein and in Storer et al., 2017; Urvois et al., 2023. As such, the reproductive ecology of Xylebroini
species may contribute to the increased chromosome number identified in *Euwallacea* and extensive
rearrangements across assemblies.

Comparative analyses indicate that the *E. fornicatus* genome has one of the highest proportions of
repeat and transposable elements relative to total assembly among the Scolytinae assessed, with
roughly 50% of the assembly composed by TEs. While the relative proportion of these elements in the
genome of *E. similis* only occupy about 30% of the total assembly*,* DNA transposons are among the
most prevalent transposable elements in both *Euwallacea* species. DNA transposons have been
identified to play a significant role in Eukaryotic genome and chromosomal evolution (Feschotte &
Pritham, 2007)*.* Moreover, it has been identified in plants (Ren et al., 2018) that genomic
rearrangement and recombination can result in net sequence loss at newly formed telomeric regions in
the chromosome. Given that we have identified recent bursts in the proliferation of DNA transposons
in the genomes of *Euwallacea*, these repetitive elements could be involved in the widespread
structural variation, high number of chromosomes and smaller genome size and content in this genus.
Further karyotyping and chromosome level assemblies of haplodiploid inbreeding Scolytinae are
needed to explore these genomic rearrangements further.

Co-evolution with symbiotic ambrosia fungi could also contribute to chromosomal rearrangements in

Xyleborini, as observed in *Euwallacea* assemblies. Genomic erosion is particularly prevalent in the

genomes of symbionts that provide nutrition to their hosts, and genes of biosynthesis pathways

producing amino acids obtained from the host are commonly lost (Beasley-Hall et al., 2023;

McCutcheon & Moran, 2012; Sloan & Moran, 2012). This has been identified to occur in host

animals as well; in the deep-sea bone-eating worm, *Osedax frankpressi* Rouse, Goffredi & Vrjenhoek

(Annelida: Polychaeta), endosymbionts provide key nutritional biosynethetic pathways which has

subsequently resulted in host genome reduction and gene loss (Moggioli et al., 2023). Similarly, in

socially parasitic ant species, gene loss of olfactory receptor repertoires corresponding with reduced

social behavioural has evolved repeatedly (Schrader et al., 2021). Host genome rearrangements have

also been reported in ants, with high levels of structural rearrangements, loss of synteny across

assembled genomes, gene family contraction, expansion and a gain of novel genes identified in the

Atta fungal farming ant lineage (Nygaard et al., 2011, 2016). Similar processes could have acted on

the genomes of fungal farming Scolytinae in the coevolution with their symbionts, leading to the

observed small genome size, genomic rearrangements and loss of synenty with other Scolytinae in

Euwallacea genomes. It should be noted that the evolution of obligate inbreeding and haplodiploidy,

in Dryocoetini, seems to preceded the development of the ambrosia fungal symbiosis in Xyleborini

(Jordal & Cognato, 2012), and as such, further chromosome scale assemblies of non-fungus farming

inbreeding haplodiploid species (Dryocoetini) are needed to assess the impact of the ambrosia

symbiosis on chromosome structure.

Genomic resources for biosecurity

The genomes assembled for *Euwallacea* will be essential resources for the management of these pests.

A vital component in managing an invasive species is understanding where the species originates and

how it arrived. Through mitochondrial haplotyping, we have identified that the population of *E.*

fornicatus introduced to Perth, Western Australia corresponds to the H38 haplotype, which has also

been introduced to California and South Africa. While further markers are needed for population

genomic analyses in order to track the invasion pathway from the native to the introduced range (Rane

et al., 2023), the assembled genomes will be vital in understanding population differentiation at the

gene and genomic level. These genomes will also be a vital resource in the diagnostics and detection

of these pests. While the mitochondrial genome for both species have already been published (Guo et

al., 2023; Wang et al., 2020), the ability to identify nuclear markers or sequences will be important for

the diagnostics and detection (such as through eDNA techniques) of the pests. Diagnostic protocols

exist for the detection of *E. fornicatus* through identification of the symbiont *Fusarium euwallaceae*

(De Jager & Roets, 2022)*,* however, the protocol operates on the detection of fungus in host plant

woody tissues, which indicates that the species is already present and likely reproducing in a given

locality. Using eDNA techniques and these reference genomes, these pests could be detected early, increasing the chances of eradication or containment, and reducing the time and resources necessary to achieve this.

Beyond tracking, detection and diagnostics, these genomes will also provide needed information for genome guided management of these pest. The endosymbiont gram-negative bacteria *Wolbachia* has been successfully deployed as a control agent for several insect pests (Morrow & Riegler, 2021; Nikolouli et al., 2018; Ross et al., 2022) and may prove useful in the control of *Euwallacea* pests. While strains have not yet been identified to be associated with *E. fornicatus*, further sequencing and MLST identification could identify naturally present strains, or strains from closely related species (such as *w*Es identified in this study) that could be artificially transferred (Grenier et al., 1998; Xi et al., 2005). However, as noted in previous studies, the impacts of *Wolbachia* on haplodiploid inbreeding Scolytinae are not yet fully understood (Bickerstaff et al., 2023; Kawasaki et al., 2016). Gene drive through CRISPR gene editing may be an avenue to control populations of invasive *E. fornicatus* and *E similis*. As demonstrated in invasive wasps (Lester et al., 2020; Meiborg et al., 2023), the ability to control population growth with this technique is limited due to haplodiploidy, with only a single gene targeting female fertility identified as useful and infertile individuals were not as competitive as fertile counterparts. Outbreeding is rare in the inbreeding Scolytinae lineages and comes at a fitness cost (Peer & Taborsky, 2005), and as such the penetration of gene drive through populations may be limited. Further insights are needed to first identify candidate genes in these genomes and then test the suitability of this management approach.

These genomes will also be of value in understanding mechanisms driving species invasion and pest impact from a genomic perspective. Preliminary comparative analyses of transposable element variation indicate that the relative proportion of genome content is similar to other beetle genomes (Gilbert et al., 2021; Petersen et al., 2019), however, a higher number of TEs are present in the *E. fornicatus* genome than in the *E. similis* genome. Transposable elements have been identified to be adaptive in species (Gilbert et al., 2021) and promote genetic diversity and rapid adaption in invasive species. The higher accumulation of TEs present in *E. fornicatus* may explain why the species can reproduce in a wider breadth of host trees and successfully establish galleries in healthy live hosts, unlike *E. similis*. Further comparative genomic analysis, such as gene family evolution (McKenna et al., 2016), structural variation (North et al., 2021; Stapley et al., 2015), and pangenomics (Gerdol et al., 2020) of both beetles and their symbionts may yield greater insights into why these species are such successful invaders compared to other haplodiploid inbreeding ambrosia beetles, understanding variation in host tree ecology and why *E. fornicatus* is more destructive than other members *Euwallacea*.

583 **Conclusions**
584

Pest species of *Euwallacea* are globally invasive and can cause significant ecological and economic
impacts in the ranges where they are established. Our study presents the first chromosome-level
genome assemblies of these pests and also for haplodiploid inbreeding species. These assemblies will
not only offer a deeper understanding of the genome structural variation that due to species ecology
but also serve as essential resources for biosecurity management and control strategies. The
comparative analyses of repeat and transposable elements, as well as the macro-synteny of
chromosome assemblies, shed light on the potential impacts of haplodiploidy and inbreeding on
genome evolution in these beetles. Our findings underscore the importance of high-quality genome
assemblies in advancing our knowledge of invasive species and contributing to their effective
management and control.
595

596 **Data Availability**
597

598 All genome and raw sequencing reads produced in this have been deposited with the NCBI, under the
599 BioProject: PRJNA1086994. Specific accession numbers for raw sequence reads and genome
600 assemblies are available in Table S1 and Table 1, respectively.
601

602 **Additional Files**
603

604 **Supplementary File 1.** The supplementary figures produced through assembly QC and analyses that
605 complement the main text.
606 **Supplementary Table S1.** Sequence read output of all libraries sequenced.
607 **Supplementary Table S2.** Classes of Repeat and Transposable Elements identified by EarlGreyTE.
608

609 **Abbreviations**
610

ANIC: Australian National Insect Collection
ANU: Australian National University
BRF-JCSMR: Biomolecular Resource Facility – John Curtin School of Medical Research
BLAST: Basic Local Alignment Search Tool
BUSCO: Benchmarking Universal Single Copy Orthologs
COI: Cytochrome Oxidase I
CSIRO: Commonwealth Scientific Industrial Research Organisation
DPIRD WA: Department of Primary Industries and Regional Development, Western Australia
EFC: *Euwallacea Fusarium* Clade

EFF: *Euwallacea fornicatus* Female

ESF: *Euwallacea similis* Female

Hi-C: High-throughput chromosome conformation capture

HiFi: High Fidelity

HMW: High Molecular Weight

MAG: Metagenome-assembled genomes

MLST: Multi-locus Sequence Typing

NCBI: National Centre for Biotechnology Information

ONT: Oxford Nanopore

PacBio: Pacific Biosciences

PSHB: Polyphagous Shot-Hole Borer

REs: Repeat Elements

TEs: Transposable Elements

Competing Interests

The authors declare no competing interests.

Funding

The work was supported by the ResearchPlus CSIRO Early Research Career Fellowship funding, the

Zimmerman Trust and ANIC-NRCA-CSIRO.

Author's Contributions

JB, HE, TW, RR, VC, AS and TW conceptualised the research. DPIRD and KI provided PSHB

specimens. TW, RR, LC and GP provided advice for HWM DNA extraction and genome assembly

pipelines. JB collected the data. JB and HE analysed and interpreted the data. JB wrote the manuscript

with input from all authors. All authors agree to the submitted manuscript.

Acknowledgements

We are grateful to the Polyphagous Shot Hole Borer response team from the Department of Primary

Industries Regional Development Western Australia for providing samples, including Basman Al-

Jalely and Emad Alshwany from the reproductive biology laboratory. Bonnie Koopmans and Debbie

Jennings for their help during fieldwork. We would like to thank Andreas Zwick, James Nicholls,

Diana Hartley and Andrew Gock (CSIRO Black Mountain) for access to laboratory facilities, Peter

Holland (Oxford University, UK) and Mara Lawniczak (Darwin Tree of Life, UK) for advice on

laboratory protocols. We thank Maxim Nekrasov and Carolina Correa Ospina (Australian National

University Biomolecular Resource Facility), and Trent Peters and Oliver Distler (Australian Genome

Research Facility), for their technical support. The authors thank John G Oakeshott for generous

advice and David Yeates (CSIRO Black Mountain) for his support of the project. We also would like

to thank Andy Bachler for advice on genome scaffolding and Ondrej Hlinka (CSIRO) for assistance

with High Performance Computing.

References

Allio, R., Schomaker-Bastos, A., Romiguier, J., Prosdocimi, F., Nabholz, B., & Delsuc, F.

(2020). MitoFinder: Efficient automated large-scale extraction of mitogenomic data in

target enrichment phylogenomics. *Molecular Ecology Resources*, *20*(4), 892–905.

https://doi.org/10.1111/1755-0998.13160

Altschul, S. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database

search programs. *Nucleic Acids Research*, *25*(17), 3389–3402.

https://doi.org/10.1093/nar/25.17.3389

Astashyn, A., Tvedte, E. S., Sweeney, D., Sapojnikov, V., Bouk, N., Joukov, V., Mozes, E.,

Strope, P. K., Sylla, P. M., Wagner, L., Bidwell, S. L., Brown, L. C., Clark, K., Davis,

E. W., Smith-White, B., Hlavina, W., Pruitt, K. D., Schneider, V. A., & Murphy, T.

D. (2024). Rapid and sensitive detection of genome contamination at scale with FCS-

GX. *Genome Biology*, *25*(1), 60. https://doi.org/10.1186/s13059-024-03198-7

Atsawawaranunt, K., Ewart, K. M., Major, R. E., Johnson, R. N., Santure, A. W., & Whibley,

A. (2023). Tracing the introduction of the invasive common myna using population

genomics. *Heredity*, *131*(1), 56–67. https://doi.org/10.1038/s41437-023-00621-w

Barclay, M. V. L., Natural History Museum Genome Acquisition Lab, Darwin Tree of Life

Barcoding collective, Wellcome Sanger Institute Tree of Life programme, Wellcome

Sanger Institute Scientific Operations: Sequencing Operations collective, Tree of Life

Core Informatics collective, & Darwin Tree of Life Consortium. (2023). The genome

sequence of a weevil, *Polydrusus cervinus* (Linnaeus, 1758). *Wellcome Open*

Research, *8*, 563. https://doi.org/10.12688/wellcomeopenres.20414.1

Baril, T., Galbraith, J., & Hayward, A. (2024). Earl Grey: A Fully Automated User-Friendly

Transposable Element Annotation and Analysis Pipeline. *Molecular Biology and*

Evolution, *41*(4), msae068. https://doi.org/10.1093/molbev/msae068

Beasley-Hall, P. G., Kinjo, Y., Rose, H. A., Walker, J., Foster, C. S. P., Kovacs, T. G. L.,

Bourguignon, T., Ho, S. Y. W., & Lo, N. (2023). Shrinking in the dark: Parallel

endosymbiont genome erosions are associated with repeated host transitions to an

underground life. *Insect Science*. https://doi.org/10.1101/2023.09.11.557272

Beaver, R. A., Sittichaya, W., & Liu, L.Y. (2014). A Synopsis of the Scolytine Ambrosia

Beetles of Thailand (Coleoptera: Curculionidae: Scolytinae). *Zootaxa*, *3875*(1).

https://doi.org/10.11646/zootaxa.3875.1.1

Bickerstaff, J. R. M., Jordal, B. H., & Riegler, M. (2023). Two sympatric lineages of

Australian *Cnestus solidus* share *Ambrosiella* symbionts but not *Wolbachia*. *Heredity*,

132(1), Article 1. https://doi.org/10.1038/s41437-023-00659-w

Biedermann, P. H. W., Müller, J., Grégoire, J.C., Gruppe, A., Hagge, J., Hammerbacher, A.,

Hofstetter, R. W., Kandasamy, D., Kolarik, M., Kostovcik, M., Krokene, P., Sallé, A.,

Six, D. L., Turrini, T., Vanderpool, D., Wingfield, M. J., & Bässler, C. (2019). Bark

Beetle Population Dynamics in the Anthropocene: Challenges and Solutions. *Trends*

in Ecology & Evolution, *34*(10), 914–924. https://doi.org/10.1016/j.tree.2019.06.002

Bierman, A., Roets, F., & Terblanche, J. S. (2022). Population structure of the invasive

ambrosia beetle, *Euwallacea fornicatus*, indicates multiple introductions into South

Africa. *Biological Invasions*, *24*(8), 2301–2312. https://doi.org/10.1007/s10530-022-

02801-x

Blackmon, H., & Demuth, J. P. (2015). Coleoptera Karyotype Database. *The Coleopterists*

Bulletin, *69*(1), 174–175. https://doi.org/10.1649/0010-065X-69.1.174

Blackmon, H., Jonika, M. M., Alfieri, J. M., Fardoun, L., & Demuth, J. P. (2024). Drift drives

the evolution of chromosome number I: The impact of trait transitions on genome

evolution in Coleoptera. *Journal of Heredity*, *115*(2), 173–182.

https://doi.org/10.1093/jhered/esae001

Breed, M. F., Harrison, P. A., Blyth, C., Byrne, M., Gaget, V., Gellie, N. J. C., Groom, S. V.

C., Hodgson, R., Mills, J. G., Prowse, T. A. A., Steane, D. A., & Mohr, J. J. (2019).

The potential of genomics for restoring ecosystems and biodiversity. *Nature Reviews*

Genetics, *20*(10), 615–628. https://doi.org/10.1038/s41576-019-0152-0

Brockerhoff, E. G., Bain, J., Kimberley, M., & Knížek, M. (2006). Interception frequency of

exotic bark and ambrosia beetles (Coleoptera: Scolytinae) and relationship with

establishment in New Zealand and worldwide. *Canadian Journal of Forest Research*,

36(2), 289–298. https://doi.org/10.1139/x05-250

Browne, F. G. (1961). The biology of Malayan Scolytidae and Platypodidae. *Malayan Forest*

Records, *22*(22), 1–255.

Burgess, B. T., Irvine, R. L., & Russello, M. A. (2022). Population genomics of Sitka black-

tailed deer supports invasive species management and ecological restoration on

islands. *Communications Biology*, *5*(1), 223. https://doi.org/10.1038/s42003-022-

03159-5

Camus, L., Gautier, M., & Boitard, S. (2024). Predicting species invasiveness with genomic

data: Is Genomic Offset related to establishment probability? *bioRxiv*.

https://doi.org/10.1101/2024.02.20.581132

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021).

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Molecular Biology and Evolution*, *38*(12),

5825–5829. https://doi.org/10.1093/molbev/msab293

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*,

25(15), 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Challis, R., Richards, E., Rajan, J., Cochrane, G., & Blaxter, M. (2020). BlobToolKit –

Interactive quality assessment of genome assemblies. *G3 Genes|Genomes|Genetics*,

10(4), 1361–1374. https://doi.org/10.1534/g3.119.400908

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics*, *34*(17), i884–i890.

https://doi.org/10.1093/bioinformatics/bty560

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de

novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2),

170–175. https://doi.org/10.1038/s41592-020-01056-5

Chernomor, O., Von Haeseler, A., & Minh, B. Q. (2016). Terrace Aware Data Structure for

Phylogenomic Inference from Supermatrices. *Systematic Biology*, *65*(6), 997–1008.

https://doi.org/10.1093/sysbio/syw037

Chua, P. Y. S., Bourlat, S. J., Ferguson, C., Korlevic, P., Zhao, L., Ekrem, T., Meier, R., &

Lawniczak, M. K. N. (2023). Future of DNA-based insect monitoring. *Trends in*

Genetics, *39*(7), 531–544. https://doi.org/10.1016/j.tig.2023.02.012

Cognato, A. I., Hoebeke, E. R., Kajimura, H., & Smith, S. M. (2015). History of the Exotic

Ambrosia Beetles *Euwallacea interjectus* and *Euwallacea validus* (Coleoptera:

Curculionidae: Xyleborini) in the United States. *Journal of Economic Entomology*,

108(3), 1129–1135. https://doi.org/10.1093/jee/tov073

Cook, D. C., & Broughton, S. (2023). Economic impact of polyphagous shot hole borer

Euwallacea fornicatus (Coleoptera: Curculionidae: Scolytinae) in Western Australia.

Agricultural and Forest Entomology, *25*(3), 449–457.

https://doi.org/10.1111/afe.12566

Copeland, M., Landa, S., Owoyemi, A., Jonika, M. M., Alfieri, J., Sylvester, T., Hoover, Z.,

Hjelmen, C. E., Johnston, J. S., Kyre, B. R., Rieske, L. K., Blackmon, H., & Casola,

C. (2024). Genome assembly of the southern pine beetle (Dendroctonus frontalis

Zimmerman) reveal the origins of gene content reduction in Dendroctonus. *bioRxiv*.

https://doi.org/10.1101/2024.05.08.592785

Cordier, T., Alonso-Sáez, L., Apothéloz-Perret-Gentil, L., Aylagas, E., Bohan, D. A.,

Bouchez, A., Chariton, A., Creer, S., Frühe, L., Keck, F., Keeley, N., Laroche, O.,

Leese, F., Pochon, X., Stoeck, T., Pawlowski, J., & Lanzén, A. (2021). Ecosystems

monitoring powered by environmental genomics: A review of current strategies with

an implementation roadmap. *Molecular Ecology*, *30*(13), 2937–2958.

https://doi.org/10.1111/mec.15472

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham,

A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of

SAMtools and BCFtools. *GigaScience*, *10*(2), giab008.

https://doi.org/10.1093/gigascience/giab008

De Jager, M. M., & Roets, F. (2022). Rapid and cost-effective detection of *Fusarium*

euwallaceae from woody tissues. *Plant Pathology*, *71*(8), 1712–1720.

https://doi.org/10.1111/ppa.13600

De Wit, M. P., Crookes, D. J., Blignaut, J. N., De Beer, Z. W., Paap, T., Roets, F., Van Der

Merwe, C., Van Wilgen, B. W., & Richardson, D. M. (2022). An Assessment of the

Potential Economic Impacts of the Invasive Polyphagous Shot Hole Borer

784   (Coleoptera: Curculionidae) in South Africa. *Journal of Economic Entomology*,

785    *115*(4), 1076–1086. https://doi.org/10.1093/jee/toac061

786 Department of Agriculture, Fisheries and Forestry. "Polyphagous Shot-Hole Borer

787    (*Euwallacea Fornicatus)* | Outbreak," January 25, 2024.

788    https://www.outbreak.gov.au/current-outbreaks/polyphagous-shot-hole-borer.

789 Department of Primary Industries and Regional Development, Government of Western

790    Australia. "Polyphagous Shot-Hole Borer (PSHB) Australian Host List." May 31, 2024.

791    https://www.agric.wa.gov.au/sites/gateway/files/PSHB-WA-Host-List.pdf.

792 Devos, Y., Mumford, J. D., Bonsall, M. B., Camargo, A. M., Firbank, L. G., Glandorf, D. C.

793    M., Nogué, F., Paraskevopoulos, K., & Wimmer, E. A. (2022). Potential use of gene

794    drive modified insects against disease vectors, agricultural pests and invasive species

795    poses new challenges for risk assessment. *Critical Reviews in Biotechnology*, *42*(2),

796    254–270. https://doi.org/10.1080/07388551.2021.1933891

797 Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R.,

798    Pham, M., Glenn St Hilaire, B., Yao, W., Stamenova, E., Hoeger, M., Nyquist, S. K.,

799    Korchina, V., Pletch, K., Flanagan, J. P., Tomaszewicz, A., McAloose, D., Pérez

800    Estrada, C., Novak, B. J., … Aiden, E. L. (2018). The Juicebox Assembly Tools

801    module facilitates *de novo* assembly of mammalian genomes with chromosome-

802    length scaffolds for under $1000. *bioRxiv*. https://doi.org/10.1101/254797

803 Durand, N. C., Robinson, J. T., Shamim, M. S., Machol, I., Mesirov, J. P., Lander, E. S., &

804    Aiden, E. L. (2016). Juicebox Provides a Visualization System for Hi-C Contact Maps

805    with Unlimited Zoom. *Cell Systems*, *3*(1), 99–101.

806    https://doi.org/10.1016/j.cels.2015.07.012

807 Eskalen, A., Gonzalez, A., Wang, D. H., Twizeyimana, M., Mayorquin, J. S., & Lynch, S. C.

808    (2012). First Report of a *Fusarium* sp. And Its Vector Tea Shot Hole Borer

(*Euwallacea fornicatus*) Causing *Fusarium* Dieback on Avocado in California. *Plant*

Disease, *96*(7), 1070–1070. https://doi.org/10.1094/PDIS-03-12-0276-PDN

Eskalen, A., Stouthamer, R., Lynch, S. C., Rugman-Jones, P. F., Twizeyimana, M., Gonzalez,

A., & Thibault, T. (2013). Host Range of Fusarium Dieback and Its Ambrosia Beetle

(Coleoptera: Scolytinae) Vector in Southern California. *Plant Disease*, *97*(7), 938–

951. https://doi.org/10.1094/PDIS-11-12-1026-RE

Felsenstein, J. (1985). Confidence Limits on Phylogenies: An Approach Using the Bootstrap.

Evolution, *39*(4).

Feng, X., Cheng, H., Portik, D., & Li, H. (2022). Metagenome assembly of high-fidelity long

reads with hifiasm-meta. *Nature Methods*, *19*(6), 671–674.

https://doi.org/10.1038/s41592-022-01478-3

Feschotte, C., & Pritham, E. J. (2007). DNA Transposons and the Evolution of Eukaryotic

Genomes. *Annual Review of Genetics*, *41*(1), 331–368.

https://doi.org/10.1146/annurev.genet.40.110405.090448

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.

(2020). RepeatModeler2 for automated genomic discovery of transposable element

families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.

https://doi.org/10.1073/pnas.1921046117

Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C.,

Crottini, A., Godoy, J. A., Höglund, J., Malukiewicz, J., Mouton, A., Oomen, R. A.,

Paez, S., Palsbøll, P. J., Pampoulie, C., Ruiz-López, M. J., Svardal, H.,

Theofanopoulou, C., … Zammit, G. (2022). The era of reference genomes in

conservation genomics. *Trends in Ecology & Evolution*, *37*(3), 197–202.

https://doi.org/10.1016/j.tree.2021.11.008

Freeman, S., Sharon, M., Maymon, M., Mendel, Z., Protasov, A., Aoki, T., Eskalen, A., &

O'Donnell, K. (2013). *Fusarium euwallaceae* sp. nov.—A symbiotic fungus of

Euwallacea sp., an invasive ambrosia beetle in Israel and California. *Mycologia*,

105(6), 1595–1606. https://doi.org/10.3852/13-066

Frey, J. E., Frey, B., Frei, D., Blaser, S., Gueuning, M., & Bühlmann, A. (2022). Next

generation biosecurity: Towards genome based identification to prevent spread of

agronomic pests and pathogens using nanopore sequencing. *PLOS ONE*, *17*(7),

e0270897. https://doi.org/10.1371/journal.pone.0270897

Gabriel, L., Brůna, T., Hoff, K. J., Ebel, M., Lomsadze, A., Borodovsky, M., & Stanke, M.

(2023). BRAKER3: Fully automated genome annotation using RNA-seq and protein

evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. *bioRxiv*.

https://doi.org/10.1101/2023.06.10.544449

Gardy, J. L., & Loman, N. J. (2018). Towards a genomics-informed, real-time, global

pathogen surveillance system. *Nature Reviews Genetics*, *19*(1), 9–20.

https://doi.org/10.1038/nrg.2017.88

Gerdol, M., Moreira, R., Cruz, F., Gómez-Garrido, J., Vlasova, A., Rosani, U., Venier, P.,

Naranjo-Ortiz, M. A., Murgarella, M., Greco, S., Balseiro, P., Corvelo, A., Frias, L.,

Gut, M., Gabaldón, T., Pallavicini, A., Canchaya, C., Novoa, B., Alioto, T. S., …

Figueras, A. (2020). Massive gene presence-absence variation shapes an open pan-

genome in the Mediterranean mussel. *Genome Biology*, *21*(1), 275.

https://doi.org/10.1186/s13059-020-02180-3

Ghurye, J., Rhie, A., Walenz, B. P., Schmitt, A., Selvaraj, S., Pop, M., Phillippy, A. M., &

Koren, S. (2019). Integrating Hi-C links with assembly graphs for chromosome-scale

assembly. *PLOS Computational Biology*, *15*(8), e1007273.

https://doi.org/10.1371/journal.pcbi.1007273

Gilbert, C., Peccoud, J., & Cordaux, R. (2021). Transposable Elements and the Evolution of

Insects. *Annual Review of Entomology*, *66*(1), 355–372.

https://doi.org/10.1146/annurev-ento-070720-074650

Gomez, D. F., Lin, W., Gao, L., & Li, Y. (2019). New host plant records for the *Euwallacea*

fornicatus (Eichhoff) species complex (Coleoptera: Curculionidae: Scolytinae) across

its natural and introduced distribution. *Journal of Asia-Pacific Entomology*, *22*(1),

338–340. https://doi.org/10.1016/j.aspen.2019.01.013

Gomez, D. F., Rabaglia, R. J., Fairbanks, K. E. O., & Hulcr, J. (2018). North American

Xyleborini north of Mexico: A review and key to genera and species (Coleoptera,

Curculionidae, Scolytinae). *ZooKeys*, *768*, 19–68.

https://doi.org/10.3897/zookeys.768.24697

Gomez, D. F., Skelton, J., Steininger, M. S., Stouthamer, R., Rugman-Jones, P., Sittichaya,

W., Rabaglia, R. J., & Hulcr, J. (2018). Species Delineation Within the *Euwallacea*

fornicatus (Coleoptera: Curculionidae) Complex Revealed by Morphometric and

Phylogenetic Analyses. *Insect Systematics and Diversity*, *2*(6), 2.

https://doi.org/10.1093/isd/ixy018

Grenier, S., Bernard, P., Heddi, A., Lassabliére, F., Jager, C., Louis, C., & Khatchadourian,

C. (1998). Successful horizontal transfer of *Wolbachia* symbionts between

Trichogramma wasps. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *265*(1404), 1441–1445. https://doi.org/10.1098/rspb.1998.0455

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying

and removing haplotypic duplication in primary genome assemblies. *Bioinformatics*,

36(9), 2896–2898. https://doi.org/10.1093/bioinformatics/btaa025

Guo, Q., Huang, W., Sang, W., Chen, X., & Wang, X. (2023). Characterization, comparative

analyses, and phylogenetic implications of mitochondrial genomes among bark and

ambrosia beetles (Coleoptera: Curculionidae, Scolytinae). *Frontiers in Ecology and*

Evolution, *11*, 1191446. https://doi.org/10.3389/fevo.2023.1191446

Hammond, A., Galizi, R., Kyrou, K., Simoni, A., Siniscalchi, C., Katsanos, D., Gribble, M.,

Baker, D., Marois, E., Russell, S., Burt, A., Windbichler, N., Crisanti, A., & Nolan, T.

(2016). A CRISPR-Cas9 gene drive system targeting female reproduction in the

malaria mosquito vector *Anopheles gambiae*. *Nature Biotechnology*, *34*(1), 78–83.

https://doi.org/10.1038/nbt.3439

Hammond, A., Pollegioni, P., Persampieri, T., North, A., Minuz, R., Trusso, A., Bucci, A.,

Kyrou, K., Morianou, I., Simoni, A., Nolan, T., Müller, R., & Crisanti, A. (2021).

Gene-drive suppression of mosquito populations in large cages as a bridge between

lab and field. *Nature Communications*, *12*(1), 4589. https://doi.org/10.1038/s41467-

021-24790-6

Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018).

UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and*

Evolution, *35*(2), 518–522. https://doi.org/10.1093/molbev/msx281

Hogg, C. J. (2023). Translating genomic advances into biodiversity conservation. *Nature*

Reviews Genetics. https://doi.org/10.1038/s41576-023-00671-0

Hopkinson, J., Balzer, J., Fang, C., & Walsh, T. (2023). Insecticide resistance management of

Bemisia tabaci (Hemiptera: Aleyrodidae) in Australian cotton – pyriproxyfen,

spirotetramat and buprofezin. *Pest Management Science*, *79*(5), 1829–1839.

https://doi.org/10.1002/ps.7361

Hulcr, J., & Stelinski, L. L. (2017). The Ambrosia Symbiosis: From Evolutionary Ecology to

Practical Management. *Annual Review of Entomology*, *62*(1), 285–303.

https://doi.org/10.1146/annurev-ento-031616-035105

Jin, M., North, H. L., Peng, Y., Liu, H., Liu, B., Pan, R., Zhou, Y., Zheng, W., Liu, K., Yang,

B., Zhang, L., Xu, Q., Elfekih, S., Valencia-Montoya, W. A., Walsh, T., Cui, P.,

Zhou, Y., Wilson, K., Jiggins, C., … Xiao, Y. (2023). Adaptive evolution to the

natural and anthropogenic environment in a global invasive crop pest, the cotton

bollworm. *The Innovation*, *4*(4), 100454. https://doi.org/10.1016/j.xinn.2023.100454

Jolley, K. A., Bray, J. E., & Maiden, M. C. J. (2018). Open-access bacterial population

genomics: BIGSdb software, the PubMLST.org website and their applications.

Wellcome Open Research, *3*, 124. https://doi.org/10.12688/wellcomeopenres.14826.1

Jordal, B. (2014). 3.7. 12 Scolytinae Latreille, 1806. *Coleoptera, Beetles. Volume 3:*

Morphology and Systematics (Phytophaga), 633–642.

Jordal, B. H., Beaver, R. A., & Kirkendall, L. R. (2001). Breaking taboos in the tropics:

Incest promotes colonization by wood-boring beetles. *Global Ecology and*

Biogeography, *10*(4), 345–357. https://doi.org/10.1046/j.1466-822X.2001.00242.x

Jordal, B. H., & Cognato, A. I. (2012). Molecular phylogeny of bark and ambrosia beetles

reveals multiple origins of fungus farming during periods of global warming. *BMC*

Evolutionary Biology, *12*(1), 133. https://doi.org/10.1186/1471-2148-12-133

Kapusta, Aurelie. (2017). *Parsing-RepeatMasker-Outputs* (Version 1) [Perl].

Kasson, M. T., O'Donnell, K., Rooney, A. P., Sink, S., Ploetz, R. C., Ploetz, J. N., Konkol, J.

L., Carrillo, D., Freeman, S., Mendel, Z., Smith, J. A., Black, A. W., Hulcr, J.,

Bateman, C., Stefkova, K., Campbell, P. R., Geering, A. D. W., Dann, E. K., Eskalen,

A., … Geiser, D. M. (2013). An inordinate fondness for *Fusarium*: Phylogenetic

diversity of fusaria cultivated by ambrosia beetles in the genus *Euwallacea* on

avocado and other plant hosts. *Fungal Genetics and Biology*, *56*, 147–157.

https://doi.org/10.1016/j.fgb.2013.04.004

Katoh, K. (2002). MAFFT: A novel method for rapid multiple sequence alignment based on

fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066.

https://doi.org/10.1093/nar/gkf436

Kawasaki, Y., Schuler, H., Stauffer, C., Lakatos, F., & Kajimura, H. (2016). *Wolbachia*

endosymbionts in haplodiploid and diploid scolytine beetles (Coleoptera:

Curculionidae: Scolytinae). *Environmental Microbiology Reports*, *8*(5), 680–688.

https://doi.org/10.1111/1758-2229.12425

Keeling, C. I., Campbell, E. O., Batista, P. D., Shegelski, V. A., Trevoy, S. A. L., Huber, D.

P. W., Janes, J. K., & Sperling, F. A. H. (2022). Chromosome-level genome assembly

reveals genomic architecture of northern range expansion in the mountain pine beetle,

Dendroctonus ponderosae Hopkins (Coleoptera: Curculionidae). *Molecular Ecology*

Resources, *22*(3), 1149–1167. https://doi.org/10.1111/1755-0998.13528

Kennedy, P. A., Meyer, K. A., Schill, D. J., Campbell, M. R., & Vu, N. V. (2018). Survival

and Reproductive Success of Hatchery YY Male Brook Trout Stocked in Idaho

Streams. *Transactions of the American Fisheries Society*, *147*(3), 419–430.

https://doi.org/10.1002/tafs.10060

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*,

37(8), 907–915. https://doi.org/10.1038/s41587-019-0201-4

Lanfear, R., Calcott, B., Ho, S. Y. W., & Guindon, S. (2012). PartitionFinder: Combined

Selection of Partitioning Schemes and Substitution Models for Phylogenetic

Analyses. *Molecular Biology and Evolution*, *29*(6), 1695–1701.

https://doi.org/10.1093/molbev/mss020

Lanier, G. (1981). Cytotaxonomy of *Dendroctonus*. *Forest, Wildlife, and Range Experiment*

Station, University of Idaho.

Lanier, G. N., & Kirkendall, L. R. (1986). Karyology of pseudogamous *Ips* bark beetles.

Hereditas, *105*(1), 87–96. https://doi.org/10.1111/j.1601-5223.1986.tb00646.x

Lantschner, M. V., Corley, J. C., & Liebhold, A. M. (2020). Drivers of global Scolytinae

invasion patterns. *Ecological Applications*, *30*(5), e02103.

https://doi.org/10.1002/eap.2103

Laslett, D., & Canbäck, B. (2008). ARWEN: A program to detect tRNA genes in metazoan

mitochondrial nucleotide sequences. *Bioinformatics*, *24*(2), 172–175.

https://doi.org/10.1093/bioinformatics/btm573

Lester, P. J., Bulgarella, M., Baty, J. W., Dearden, P. K., Guhlin, J., & Kean, J. M. (2020).

The potential for a CRISPR gene drive to eradicate or suppress globally invasive

social wasps. *Scientific Reports*, *10*(1), 12398. https://doi.org/10.1038/s41598-020-

69259-6

Li, K., Xu, P., Wang, J., Yi, X., & Jiao, Y. (2023). Identification of errors in draft genome

assemblies at single-nucleotide resolution for quality assessment and improvement.

Nature Communications, *14*(1), 6556. https://doi.org/10.1038/s41467-023-42336-w

Liu, Z., Xing, L., Huang, W., Liu, B., Wan, F., Raffa, K. F., Hofstetter, R. W., Qian, W., &

Sun, J. (2022). Chromosome-level genome assembly and population genomic

analyses provide insights into adaptive evolution of the red turpentine beetle,

Dendroctonus valens. *BMC Biology*, *20*(1), 190. https://doi.org/10.1186/s12915-022-

01388-y

Lynn, K. M. T., Wingfield, M. J., Durán, A., Oliveira, L. S. S., De Beer, Z. W., & Barnes, I.

(2021). Novel *Fusarium* mutualists of two *Euwallacea* species infesting *Acacia*

crassicarpa in Indonesia. *Mycologia*, *113*(3), 536–558.

https://doi.org/10.1080/00275514.2021.1875708

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper

Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.

Molecular Biology and Evolution, *38*(10), 4647–4654.

https://doi.org/10.1093/molbev/msab199

Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting

of occurrences of *k* -mers. *Bioinformatics*, *27*(6), 764–770.

https://doi.org/10.1093/bioinformatics/btr011

Maselko, M., Heinsch, S. C., Chacón, J. M., Harcombe, W. R., & Smanski, M. J. (2017).

Engineering species-like barriers to sexual reproduction. *Nature Communications*,

8(1), 883. https://doi.org/10.1038/s41467-017-01007-3

Matheson, P., & McGaughran, A. (2022). Genomic data is missing for many highly invasive

species, restricting our preparedness for escalating incursion rates. *Scientific Reports*,

12(1), 13987. https://doi.org/10.1038/s41598-022-17937-y

Mc Cartney, A. M., Formenti, G., Mouton, A., De Panis, D., Marins, L. S., Leitao, H. G.,

Diedericks, G., Kirangwa, J., Morselli, M., Salces, J., Escudero, N., Iannucci, A.,

Natali, C., Svardal, H., Fernandez, R., De Pooter, T., Joris, G., Strazisar, M., Wood,

J., … Mazzoni, C. J. (2023). *The European Reference Genome Atlas: Piloting a*

decentralised approach to equitable biodiversity genomics.

https://doi.org/10.1101/2023.09.25.559365

McCutcheon, J. P., & Moran, N. A. (2012). Extreme genome reduction in symbiotic bacteria.

Nature Reviews Microbiology, *10*(1), 13–26. https://doi.org/10.1038/nrmicro2670

McGaughran, A., Dhami, M. K., Parvizi, E., Vaughan, A. L., Gleeson, D. M., Hodgins, K.

A., Rollins, L. A., Tepolt, C. K., Turner, K. G., Atsawawaranunt, K., Battlay, P.,

Congrains, C., Crottini, A., Dennis, T. P. W., Lange, C., Liu, X. P., Matheson, P.,

1005 North, H. L., Popovic, I., … Wilson, J. (2024). Genomic Tools in Biological

1006 Invasions: Current State and Future Frontiers. *Genome Biology and Evolution*, *16*(1),

1007 evad230. https://doi.org/10.1093/gbe/evad230

1008 McGaughran, A., Mori, E., & Quilodrán, C. S. (2021). Editorial: The Genomics of Biological

1009 Invasion. *Frontiers in Ecology and Evolution*, *9*, 810817.

1010 https://doi.org/10.3389/fevo.2021.810817

1011 McKenna, D. D., Scully, E. D., Pauchet, Y., Hoover, K., Kirsch, R., Geib, S. M., Mitchell, R.

1012 F., Waterhouse, R. M., Ahn, S.-J., Arsala, D., Benoit, J. B., Blackmon, H., Bledsoe,

1013 T., Bowsher, J. H., Busch, A., Calla, B., Chao, H., Childers, A. K., Childers, C., …

1014 Richards, S. (2016). Genome of the Asian longhorned beetle (*Anoplophora*

1015 *glabripennis*), a globally significant invasive species, reveals key functional and

1016 evolutionary innovations at the beetle–plant interface. *Genome Biology*, *17*(1), 227.

1017 https://doi.org/10.1186/s13059-016-1088-8

1018 Meiborg, A. B., Faber, N. R., Taylor, B. A., Harpur, B. A., & Gorjanc, G. (2023). The

1019 suppressive potential of a gene drive in populations of invasive social wasps is

1020 currently limited. *Scientific Reports*, *13*(1), 1640. https://doi.org/10.1038/s41598-023-

1021 28867-8

1022 Mendel, Z., Lynch, S. C., Eskalen, A., Protasov, A., Maymon, M., & Freeman, S. (2021).

1023 What Determines Host Range and Reproductive Performance of an Invasive

1024 Ambrosia Beetle *Euwallacea fornicatus*; Lessons From Israel and California.

1025 *Frontiers in Forests and Global Change*, *4*, 654702.

1026 https://doi.org/10.3389/ffgc.2021.654702

1027 Mendel, Z., Protasov, A., Sharon, M., Zveibil, A., Yehuda, S. B., O'Donnell, K., Rabaglia,

1028 R., Wysoki, M., & Freeman, S. (2012). An Asian ambrosia beetle *Euwallacea*

1029 *fornicatus* and its novel symbiotic fungus *Fusarium* sp. Pose a serious threat to the

Israeli avocado industry. *Phytoparasitica*, *40*(3), 235–238.

https://doi.org/10.1007/s12600-012-0223-7

Meurisse, N., Rassati, D., Hurley, B. P., Brockerhoff, E. G., & Haack, R. A. (2019). Common

pathways by which non-native forest insects move internationally and domestically.

Journal of Pest Science, *92*(1), 13–27. https://doi.org/10.1007/s10340-018-0990-0

Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., & Gurevich, A. (2018). Versatile

genome assembly evaluation with QUAST-LG. *Bioinformatics*, *34*(13), i142–i150.

https://doi.org/10.1093/bioinformatics/bty266

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von

Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods

for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*,

37(5), 1530–1534. https://doi.org/10.1093/molbev/msaa015

Moggioli, G., Panossian, B., Sun, Y., Thiel, D., Martín-Zamora, F. M., Tran, M., Clifford, A.

M., Goffredi, S. K., Rimskaya-Korsakova, N., Jékely, G., Tresguerres, M., Qian, P.-

Y., Qiu, J.-W., Rouse, G. W., Henry, L. M., & Martín-Durán, J. M. (2023). Distinct

genomic routes underlie transitions to specialised symbiotic lifestyles in deep-sea

annelid worms. *Nature Communications*, *14*(1), 2814. https://doi.org/10.1038/s41467-

023-38521-6

Morrow, J. L., & Riegler, M. (2021). Genome analyses of four *Wolbachia* strains and

associated mitochondria of *Rhagoletis cerasi* expose cumulative modularity of

cytoplasmic incompatibility factors and cytoplasmic hitchhiking across host

populations. *BMC Genomics*, *22*(1), 616. https://doi.org/10.1186/s12864-021-07906-6

Navarro-Escalante, L., Hernandez-Hernandez, E. M., Nuñez, J., Acevedo, F. E., Berrio, A.,

Constantino, L. M., Padilla-Hurtado, B. E., Molina, D., Gongora, C., Acuña, R.,

Stuart, J., & Benavides, P. (2021). A coffee berry borer (*Hypothenemus hampei*)

genome assembly reveals a reduced chemosensory receptor gene repertoire and male-specific genome sequences. *Scientific Reports*, *11*(1), 4900. https://doi.org/10.1038/s41598-021-84068-1

Nikolouli, K., Colinet, H., Renault, D., Enriquez, T., Mouton, L., Gibert, P., Sassu, F., Cáceres, C., Stauffer, C., Pereira, R., & Bourtzis, K. (2018). Sterile insect technique and *Wolbachia* symbiosis as potential tools for the control of the invasive species *Drosophila suzukii*. *Journal of Pest Science*, *91*(2), 489–503. https://doi.org/10.1007/s10340-017-0944-y

North, H. L., McGaughran, A., & Jiggins, C. D. (2021). Insights into invasive species from whole-genome resequencing. *Molecular Ecology*, *30*(23), 6289–6308. https://doi.org/10.1111/mec.15999

Nurk, S., Walenz, B. P., Rhie, A., Vollger, M. R., Logsdon, G. A., Grothe, R., Miga, K. H., Eichler, E. E., Phillippy, A. M., & Koren, S. (2020). HiCanu: Accurate assembly of segmental duplications, satellites, and allelic variants from high-fidelity long reads. *Genome Research*, *30*(9), 1291–1305. https://doi.org/10.1101/gr.263566.120

Nygaard, S., Hu, H., Li, C., Schiøtt, M., Chen, Z., Yang, Z., Xie, Q., Ma, C., Deng, Y., Dikow, R. B., Rabeling, C., Nash, D. R., Wcislo, W. T., Brady, S. G., Schultz, T. R., Zhang, G., & Boomsma, J. J. (2016). Reciprocal genomic evolution in the ant–fungus agricultural symbiosis. *Nature Communications*, *7*(1), 12233. https://doi.org/10.1038/ncomms12233

Nygaard, S., Zhang, G., Schiøtt, M., Li, C., Wurm, Y., Hu, H., Zhou, J., Ji, L., Qiu, F., Rasmussen, M., Pan, H., Hauser, F., Krogh, A., Grimmelikhuijzen, C. J. P., Wang, J., & Boomsma, J. J. (2011). The genome of the leaf-cutting ant *Acromyrmex echinatior* suggests key adaptations to advanced social life and fungus farming. *Genome Research*, *21*(8), 1339–1348. https://doi.org/10.1101/gr.121392.111

O'Donnell, K., Libeskind-Hadas, R., Hulcr, J., Bateman, C., Kasson, M. T., Ploetz, R. C.,

Konkol, J. L., Ploetz, J. N., Carrillo, D., Campbell, A., Duncan, R. E., Liyanage, P. N.

H., Eskalen, A., Lynch, S. C., Geiser, D. M., Freeman, S., Mendel, Z., Sharon, M.,

Aoki, T., … Rooney, A. P. (2016). Invasive Asian *Fusarium – Euwallacea* ambrosia

beetle mutualists pose a serious threat to forests, urban landscapes and the avocado

industry. *Phytoparasitica*, *44*(4), 435–442. https://doi.org/10.1007/s12600-016-0543-

0

Paap, T., De Beer, Z. W., Migliorini, D., Nel, W. J., & Wingfield, M. J. (2018). The

polyphagous shot hole borer (PSHB) and its fungal symbiont *Fusarium euwallaceae*:

A new invasion in South Africa. *Australasian Plant Pathology*, *47*(2), 231–237.

https://doi.org/10.1007/s13313-018-0545-0

Parvizi, E., Dhami, M. K., Yan, J., & McGaughran, A. (2023). Population genomic insights

into invasion success in a polyphagous agricultural pest, *Halyomorpha halys*.

Molecular Ecology, *32*(1), 138–151. https://doi.org/10.1111/mec.16740

Peer, K., & Taborsky, M. (2005). Outbreeding depression, but no inbreeding depression in

haplodiploid ambrosia beetles with regular sibling mating. *Evolution*, *59*(2), 317–323.

https://doi.org/10.1111/j.0014-3820.2005.tb00992.x

Petersen, M., Armisén, D., Gibbs, R. A., Hering, L., Khila, A., Mayer, G., Richards, S.,

Niehuis, O., & Misof, B. (2019). Diversity and evolution of the transposable element

repertoire in arthropods with particular reference to insects. *BMC Ecology and*

Evolution, *19*(1), 11. https://doi.org/10.1186/s12862-018-1324-9

Pistone, D., Gohli, J., & Jordal, B. H. (2018). Molecular phylogeny of bark and ambrosia

beetles (Curculionidae: Scolytinae) based on 18 molecular markers. *Systematic*

Entomology, *43*(2), 387–406. https://doi.org/10.1111/syen.12281

Powell, D., Groβe-Wilde, E., Krokene, P., Roy, A., Chakraborty, A., Löfstedt, C., Vogel, H., Andersson, M. N., & Schlyter, F. (2021). A highly-contiguous genome assembly of the Eurasian spruce bark beetle, *Ips typographus*, provides insight into a major forest pest. *Communications Biology*, *4*(1), 1059. https://doi.org/10.1038/s42003-021-02602-3

R Core Team, R. (2013). *R: A language and environment for statistical computing*.

Raffa, K. F., Aukema, B. H., Bentz, B. J., Carroll, A. L., Hicke, J. A., Turner, M. G., & Romme, W. H. (2008). Cross-scale Drivers of Natural Disturbances Prone to Anthropogenic Amplification: The Dynamics of Bark Beetle Eruptions. *BioScience*, *58*(6), 501–517. https://doi.org/10.1641/B580607

Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2020). GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nature Communications*, *11*(1), 1432. https://doi.org/10.1038/s41467-020-14998-3

Rane, R., Walsh, T. K., Lenancker, P., Gock, A., Dao, T. H., Nguyen, V. L., Khin, T. N., Amalin, D., Chittarath, K., Faheem, M., Annamalai, S., Thanarajoo, S. S., Trisyono, Y. A., Khay, S., Kim, J., Kuniata, L., Powell, K., Kalyebi, A., Otim, M. H., … Tay, W. T. (2023). Complex multiple introductions drive fall armyworm invasions into Asia and Australia. *Scientific Reports*, *13*(1), 660. https://doi.org/10.1038/s41598-023-27501-x

Rassati, D., Faccoli, M., Petrucco Toffolo, E., Battisti, A., & Marini, L. (2015). Improving the early detection of alien wood-boring beetles in ports and surrounding forests. *Journal of Applied Ecology*, *52*(1), 50–58. https://doi.org/10.1111/1365-2664.12347

Ren, L., Huang, W., Cannon, E. K. S., Bertioli, D. J., & Cannon, S. B. (2018). A Mechanism for Genome Size Reduction Following Genomic Rearrangements. *Frontiers in Genetics*, *9*, 454. https://doi.org/10.3389/fgene.2018.00454

Ross, P. A., Robinson, K. L., Yang, Q., Callahan, A. G., Schmidt, T. L., Axford, J. K.,

Coquilleau, M. P., Staunton, K. M., Townsend, M., Ritchie, S. A., Lau, M.-J., Gu, X.,

& Hoffmann, A. A. (2022). A decade of stability for *w*Mel *Wolbachia* in natural

Aedes aegypti populations. *PLOS Pathogens*, *18*(2), e1010256.

https://doi.org/10.1371/journal.ppat.1010256

Rugman-Jones, P. F., Au, M., Ebrahimi, V., Eskalen, A., Gillett, C. P. D. T., Honsberger, D.,

Husein, D., Wright, M. G., Yousuf, F., & Stouthamer, R. (2020). One becomes two:

Second species of the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae)

species complex is established on two Hawaiian Islands. *PeerJ*, *8*, e9987.

https://doi.org/10.7717/peerj.9987

Ruzzier, E., Ortis, G., Vallotto, D., Faccoli, M., Martinez-Sañudo, I., & Marchioro, M.

(2023). The first full host plant dataset of Curculionidae Scolytinae of the world:

Tribe Xyleborini LeConte, 1876. *Scientific Data*, *10*(1), 166.

https://doi.org/10.1038/s41597-023-02083-5

Schrader, L., Pan, H., Bollazzi, M., Schiøtt, M., Larabee, F. J., Bi, X., Deng, Y., Zhang, G.,

Boomsma, J. J., & Rabeling, C. (2021). Relaxed selection underlies genome erosion

in socially parasitic ant species. *Nature Communications*, *12*(1), 2918.

https://doi.org/10.1038/s41467-021-23178-w

Schuler, H., Witkowski, R., Van De Vossenberg, B., Hoppe, B., Mittelbach, M.,

Bukovinszki, T., Schwembacher, S., Van De Meulengraaf, B., Lange, U., Rode, S.,

Andriolo, A., Bełka, M., Mazur, A., & Battisti, A. (2023). Recent invasion and

eradication of two members of the *Euwallacea fornicatus* species complex

(Coleoptera: Curculionidae: Scolytinae) from tropical greenhouses in Europe.

Biological Invasions, *25*(2), 299–307. https://doi.org/10.1007/s10530-022-02929-w

Schultz, D. T., Haddock, S. H. D., Bredeson, J. V., Green, R. E., Simakov, O., & Rokhsar, D.

S. (2023). Ancient gene linkages support ctenophores as sister to other animals.

Nature, *618*(7963), Article 7963. https://doi.org/10.1038/s41586-023-05936-6

Sillero, N., Huey, R. B., Gilchrist, G., Rissler, L., & Pascual, M. (2020). Distribution

modelling of an introduced species: Do adaptive genetic markers affect potential

range? *Proceedings of the Royal Society B: Biological Sciences*, *287*(1935),

20201791. https://doi.org/10.1098/rspb.2020.1791

Sim, S. B., Corpuz, R. L., Simmonds, T. J., & Geib, S. M. (2022). HiFiAdapterFilt, a

memory efficient read processing pipeline, prevents occurrence of adapter sequence

in PacBio HiFi reads and their negative impacts on genome assembly. *BMC*

Genomics, *23*(1), 157. https://doi.org/10.1186/s12864-022-08375-1

Six, D. L., & Bracewell, R. (2015). *Dendroctonus*. In *Bark Beetles* (pp. 305–350). Elsevier.

https://doi.org/10.1016/B978-0-12-417156-5.00008-3

Sloan, D. B., & Moran, N. A. (2012). Genome Reduction and Co-evolution between the

Primary and Secondary Bacterial Symbionts of Psyllids. *Molecular Biology and*

Evolution, *29*(12), 3781–3792. https://doi.org/10.1093/molbev/mss180

Smit, A., Hubley, R., & Green, P. (2013). *RepeatMasker Open-4.0.* [Computer software].

http://www.repeatmasker.org

Smith, S. M., F. Gomez, D., A. Beaver, R., Hulcr, J., & I. Cognato, A. (2019). Reassessment

of the Species in the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae)

Complex after the Rediscovery of the "Lost" Type Specimen. *Insects*, *10*(9), 261.

https://doi.org/10.3390/insects10090261

Smith, S. M., & Hulcr, J. (2015). *Scolytus* and other Economically Important Bark and

Ambrosia Beetles. In *Bark Beetles* (pp. 495–531). Elsevier.

https://doi.org/10.1016/B978-0-12-417156-5.00012-5

Stanke, M., Diekhans, M., Baertsch, R., & Haussler, D. (2008). Using native and syntenically mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics*, *24*(5), 637–644. https://doi.org/10.1093/bioinformatics/btn013

Stapley, J., Santure, A. W., & Dennis, S. R. (2015). Transposable elements as agents of rapid adaptation may explain the genetic paradox of invasive species. *Molecular Ecology*, *24*(9), 2241–2252. https://doi.org/10.1111/mec.13089

Steenwyk, J. L., Buida, T. J., Labella, A. L., Li, Y., Shen, X.-X., & Rokas, A. (2021). PhyKIT: A broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic data. *Bioinformatics*, *37*(16), 2325–2331. https://doi.org/10.1093/bioinformatics/btab096

Storer, C. G., Breinholt, J. W., & Hulcr, J. (2015). *Wallacellus* is *Euwallacea*: Molecular phylogenetics settles generic relationships (Coleoptera: Curculionidae: Scolytinae: Xyleborini). *Zootaxa*, *3974*(3), 391–400. https://doi.org/10.11646/zootaxa.3974.3.6

Storer, C., Payton, A., McDaniel, S., Jordal, B., & Hulcr, J. (2017). Cryptic genetic variation in an inbreeding and cosmopolitan pest, *Xylosandrus crassiusculus*, revealed using dd RAD seq. *Ecology and Evolution*, *7*(24), 10974–10986. https://doi.org/10.1002/ece3.3625

Stouthamer, R., Rugman-Jones, P., Thu, P. Q., Eskalen, A., Thibault, T., Hulcr, J., Wang, L., Jordal, B. H., Chen, C., Cooperband, M., Lin, C., Kamata, N., Lu, S., Masuya, H., Mendel, Z., Rabaglia, R., Sanguansub, S., Shih, H., Sittichaya, W., & Zong, S. (2017). Tracing the origin of a cryptic invader: Phylogeography of the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) species complex. *Agricultural and Forest Entomology*, *19*(4), 366–375. https://doi.org/10.1111/afe.12215

Takenouchi, Y., & Takagi, K. (1967). A chromosome study of two parthenogenetic scolytid beetles. *Annotationes Zoologicae Japonenses*, *40*(2).

Tay, W. T., Rane, R. V., James, W., Gordon, K. H. J., Downes, S., Kim, J., Kuniata, L., &

Walsh, T. K. (2022). Resistance Bioassays and Allele Characterization Inform

Analysis of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Introduction Pathways

in Asia and Australia. *Journal of Economic Entomology*, *115*(6), 1790–1805.

https://doi.org/10.1093/jee/toac151

Tepolt, C. K., Grosholz, E. D., De Rivera, C. E., & Ruiz, G. M. (2022). Balanced

polymorphism fuels rapid selection in an invasive crab despite high gene flow and

low genetic diversity. *Molecular Ecology*, *31*(1), 55–69.

https://doi.org/10.1111/mec.16143

The Darwin Tree of Life Project Consortium, Blaxter, M., Mieszkowska, N., Di Palma, F.,

Holland, P., Durbin, R., Richards, T., Berriman, M., Kersey, P., Hollingsworth, P.,

Wilson, W., Twyford, A., Gaya, E., Lawniczak, M., Lewis, O., Broad, G., Howe, K.,

Hart, M., Flicek, P., & Barnes, I. (2022). Sequence locally, think globally: The

Darwin Tree of Life Project. *Proceedings of the National Academy of Sciences*,

119(4), e2115642118. https://doi.org/10.1073/pnas.2115642118

Theissinger, K., Fernandes, C., Formenti, G., Bista, I., Berg, P. R., Bleidorn, C., Bombarely,

A., Crottini, A., Gallo, G. R., Godoy, J. A., Jentoft, S., Malukiewicz, J., Mouton, A.,

Oomen, R. A., Paez, S., Palsbøll, P. J., Pampoulie, C., Ruiz-López, M. J., Secomandi,

S., … Zammit, G. (2023). How genomics can help biodiversity conservation. *Trends*

in Genetics, *39*(7), 545–559. https://doi.org/10.1016/j.tig.2023.01.005

Thia, J. A., Korhonen, P. K., Young, N. D., Gasser, R. B., Umina, P. A., Yang, Q., Edwards,

O., Walsh, T., & Hoffmann, A. A. (2023). The redlegged earth mite draft genome

provides new insights into pesticide resistance evolution and demography in its

invasive Australian range. *Journal of Evolutionary Biology*, *36*(2), 381–398.

https://doi.org/10.1111/jeb.14144

Uliano-Silva, M., Ferreira, J. G. R. N., Krasheninnikova, K., Darwin Tree of Life

Consortium, Blaxter, M., Mieszkowska, N., Hall, N., Holland, P., Durbin, R.,

Richards, T., Kersey, P., Hollingsworth, P., Wilson, W., Twyford, A., Gaya, E.,

Lawniczak, M., Lewis, O., Broad, G., Martin, F., … McCarthy, S. A. (2023).

MitoHiFi: A python pipeline for mitochondrial genome assembly from PacBio high

fidelity reads. *BMC Bioinformatics*, *24*(1), 288. https://doi.org/10.1186/s12859-023-

05385-y

Urvois, T., Perrier, C., Roques, A., Sauné, L., Courtin, C., Kajimura, H., Hulcr, J., Cognato,

A. I., Auger-Rozenberg, M. -A., & Kerdelhué, C. (2023). The worldwide invasion

history of a pest ambrosia beetle inferred using population genomics. *Molecular*

Ecology, *32*(15), 4381–4400. https://doi.org/10.1111/mec.16993

Vasimuddin, Md., Misra, S., Li, H., & Aluru, S. (2019). Efficient Architecture-Aware

Acceleration of BWA-MEM for Multicore Systems. *2019 IEEE International*

Parallel and Distributed Processing Symposium (IPDPS), 314–324.

https://doi.org/10.1109/IPDPS.2019.00041

Vilardo, G., Faccoli, M., Corley, J. C., & Lantschner, M. V. (2022). Factors driving historic

intercontinental invasions of European pine bark beetles. *Biological Invasions*, *24*(9),

2973–2991. https://doi.org/10.1007/s10530-022-02818-2

Virkki, N., & Smith, S. G. (1978). *Animal Cytogenetics: Vol. Vol 3: Insecta 5*. Schweizerbart

Science Publishers.

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., &

Schatz, M. C. (2017). GenomeScope: Fast reference-free genome profiling from short

reads. *Bioinformatics*, *33*(14), 2202–2204.

https://doi.org/10.1093/bioinformatics/btx153

Wang, L.J., Hsu, M.H., Liu, T.Y., Lin, M.Y., & Sung, C.H. (2020). Characterization of the

complete mitochondrial genome of *Euwallacea fornicatus* (Eichhoff, 1868)

(Coleoptera: Curculionidae: Scolytinae) and its phylogenetic implications.

Mitochondrial DNA Part B, *5*(3), 3502–3504.

https://doi.org/10.1080/23802359.2020.1827070

Wang, Z., Liu, Y., Wang, H., Roy, A., Liu, H., Han, F., Zhang, X., & Lu, Q. (2023). Genome

and transcriptome of *Ips nitidus* provide insights into high-altitude hypoxia adaptation

and symbiosis. *iScience*, *26*(10), 107793. https://doi.org/10.1016/j.isci.2023.107793

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund,

G., Hayes, A., Henry, L., & Hester, J. (2019). Welcome to the Tidyverse. *Journal of*

Open Source Software, *4*(43), 1686.

Wong, W. Y., & Simakov, O. (2019). RepeatCraft: A meta-pipeline for repetitive element de-

fragmentation and annotation. *Bioinformatics*, *35*(6), 1051–1052.

https://doi.org/10.1093/bioinformatics/bty745

Xi, Z., Dean, J. L., Khoo, C., & Dobson, Stephen. L. (2005). Generation of a novel

Wolbachia infection in *Aedes albopictus* (Asian tiger mosquito) via embryonic

microinjection. *Insect Biochemistry and Molecular Biology*, *35*(8), 903–910.

https://doi.org/10.1016/j.ibmb.2005.03.015

Xu, S., Li, L., Luo, X., Chen, M., Tang, W., Zhan, L., Dai, Z., Lam, T. T., Guan, Y., & Yu,

G. (2022). *Ggtree*: A serialized data object for visualization of a phylogenetic tree and

annotation data. *iMeta*, *1*(4), e56. https://doi.org/10.1002/imt2.56

Xu, Z., & Wang, H. (2007). LTR_FINDER: An efficient tool for the prediction of full-length

LTR retrotransposons. *Nucleic Acids Research*, *35*(Web Server), W265–W268.

https://doi.org/10.1093/nar/gkm286

Yin, X., Martinez, A. S., Sepúlveda, M. S., & Christie, M. R. (2021). Rapid genetic

adaptation to recently colonized environments is driven by genes underlying life

history traits. *BMC Genomics*, *22*(1), 269. https://doi.org/10.1186/s12864-021-07553-

x

Zhang, Y., Liu, S., De Meyer, M., Liao, Z., Zhao, Y., Virgilio, M., Feng, S., Qin, Y., Singh,

S., Wee, S. L., Jiang, F., Guo, S., Li, H., Deschepper, P., Vanbergen, S., Delatte, H.,

Van Sauers-Muller, A., Syamsudin, T. S., Kawi, A. P., … Li, Z. (2023). Genomes of

the cosmopolitan fruit pest *Bactrocera dorsalis* (Diptera: Tephritidae) reveal its global

invasion history and thermal adaptation. *Journal of Advanced Research*, *53*, 61–74.

https://doi.org/10.1016/j.jare.2022.12.012

Zhou, C., McCarthy, S. A., & Durbin, R. (2023). YaHS: Yet another Hi-C scaffolding tool.

Bioinformatics, *39*(1), btac808. https://doi.org/10.1093/bioinformatics/btac808

1288

1289   **Figures**

1290

1291

1292   **Figure 1.** The ambrosia beetles A) *Euwallacea fornicatus* (PSHB) and B) *Euwallacea similis* whose

1293   genomes were sequenced herein. Photo taken by Hermes Escalona.

1294

1295

1296

Figure 2. Maximum-likelihood phylogenetic tree of the concatenated 2,108 BUSCO single copy
sequences of available pseudo-chromosome level assemblies for Scolytinae based using 2,108 single
copy complete BUSCO sequences. All nodes were strongly supported with 100 bootstrap support.
Scale bar indicated 0.03 substitutions per nucleotide position. To the right of the tips are the relative
proportion of repetitive element composition as annotated in the species' genome.

Figure 3. Ribbon plot showing the synteny and structural and chromosomal rearrangement in Scolytinae chromosomal assemblies. Syntenic blocks of single copy complete BUSCO sequences link pseudo-chromosome scaffolds between assemblies. Black bars represent individual pseudo-chromosome scaffolds and are labelled according to the name provided in the assembly. Scaffolds are ordered following the *Dendroctonus ponderosae* assemblies and the lengths of the bars representing scaffolds indicate total number of BUSCOs and not scaffold length. Coloured lines connect orthologous genes across scaffolds and the colour code follows the Bilaterian-Cnidarian-Sponge Linkage Groups (BCnS LGs) as in (Schultz et al., 2023).

1312 **Table 1.** Contig and scaffold summary statistics for the assemblies produced. Subsampled library refers to the level of subsampling and sequence statistics

1313 therein post subsampling with seqtk. Number of assembly seqs refers to the number of contigs (contig assembly) or scaffolds (scaffold assembly) present the

1314 assemblies. Mean HiFi coverage was calculated by dividing the number of base pairs in the filtered subsampled HiFi libraries by the total assembly length.

1315 Mean Hi-C coverage was calculated by dividing the number of base pairs in the filtered Hi-C libraries by the total assembly length.

1316

Species	*Euwallacea fornicatus*				*Euwallacea similis*			
Library	EFF26		EFF42		ESF13		ESF15	
Assembly	Contig	Scaffold	Contig	Scaffold	Contig	Scaffold	Contig	Scaffold
NCBI Accession								
Estimated Genome Size (Mb)	192.7	192.7	197.7	197.7	181.83	181.83	170.77	170.77
Library subsampling strategy	0.35	0.35	0.35	0.35	1	1	0.15	0.15
Subsampled seqs (K)	609.62	609.62	578.21	578.21	0.54	0.54	659.71	659.71
Subsampled bases (Gb)	6.5	6.5	7	7	5.63	5.63	8.09	8.09
Total assembly Length (Mb)	268.98	219.83	290.46	239.6	210.3	177.6	210.22	180.07
Longest assembly seq (Mb)	9.93	8.67	10.78	8.89	7.16	10.09	7.19	10.04
Number of assembly seqs	530	185	422	179	654	128	480	140
Number of pseudo-chromosomes		39		39		37		37
% assembly anchored in pseudo-chromosomes		75%		69%		93%		91%
N50 (Mb)	4.1	3.8	4.9	3.6	3.6	5.14	3.7	5.14
L50	22	20	21	23	22	13	21	13

N's per 100 kpb	0	0.36	0	0.42	0	2.49	0	1.9
Mean HiFi coverage (x)	24.18	29.58	24.07	31.80	26.79	30.95	38.50	44.78
Mean Hi-C coverage (x)	11.90	14.56	11.02	13.36	15.69	18.12	15.70	18.26
BUSCO: genome (BUSCO v5.5.0; endopterygota_odb10, n = 2124)								
Complete single copy	2080 (97.9%)	2080 (97.9%)	2082 (98%)	2079 (97.9%)	2062 (97.1%)	2084 (98.1%)	2096 (97.4%)	2084 (98.1%)
Complete duplicated	7 (0.3%)	8 (0.4%)	8 (0.4%)	8 (0.4%)	28 (1.3%)	9 (0.4%)	19 (0.9%)	9 (0.4%)
Fragmented	11 (0.5%	15 (0.7%)	12 (0.6%)	15 (0.7%)	10 (0.5%)	7 (0.3%)	9 (0.4%)	8 (0.4%)
Missing	26 (1.3%)	21 (1%)	22 (1%)	22 (1%)	24 (1.1%)	24 (1.2%)	27 (1.3%)	23 (1.1%)

1317

1318 **Table 2.** Protein and repeat annotation statistics for the assemblies produced. Proteins were identified and annotated by BRAKER3 and completeness

1319 assessed with BUSCO. Repeats were identified by EarlGreyTE.

1320

Species	*Euwallacea fornicatus*		*Euwallacea similis*	
Library	EFF26	EFF42	ESF13	ESF15
Predicted transcripts	16,390	10,749	16,170	16,206
Predicted genes	11,589	10,101	12,128	12,153
Total coding region (bp) (%)	17,388,659 (7.9%)	14,946,768 (6%)	18,078,735 (10%)	18,139,032 (10%)
Functionally annotated transcripts	14,771	9,584	14,600	14,600
Functionally annotated genes	10,767	9,375	11,327	11,285
BUSCO: protein (BUSCO v5.5.0; endopterygota_odb10, n = 2124)				
Complete single copy	2089 (97.7%)	1889 (88%)	2075 (97.7)	2075 (97.7)
Complete duplicated	14 (0.7%)	19 (0.9%)	18 (0.8%)	17 (0.8%)
Fragmented	10 (0.5%)	16 (0.8%)	4 (0.2%)	4 (0.2%)
Missing	25 (1.1%)	219 (10.3%)	27 (1.3%)	28 (1.3%)
Repeats				
DNA (bp) (%)	68,089,859 (30.9)	31,487,152 (13.1)	18,861,021 (10.6)	18,580,216 (10.3)
LINE (bp) (%)	5,329,747 (2.4)	7,068,963 (3.0)	3,325,503 (1.9)	3,673,824 (2.0)
LTR (bp) (%)	7,594,481 (3.5)	9,293,361 (3.9)	13,488,481 (7.6)	16,621,346 (9.1)
Unclassified (bp) (%)	18,446,822 (8.4)	19,202,615 (8.0)	21,972,489 (12.4)	21,763,152 (12.0)
Total masked (bp) (%)	103,661,639 (47.1)	122,116,844 (50.9)	58,944,136 (33.2)	61778298 (34.2)

1321