Chromosome structural rearrangements in invasive haplodiploid ambrosia beetles revealed by the genomes of *Euwallacea fornicatus* and *Euwallacea similis* (Coleoptera, Curculionidae, Scolytinae)

Authors: James R. M. Bickerstaff[1]*, Tom Walsh[2], Leon Court[2], Gunjan Pandey[2], Kylie Ireland[3, 4,] David Cousins [3], Valerie Caron[5,6], Thomas Wallenius[7], Adam Slipinski[1], Rahul Rane[5], Hermes E. Escalona[1]*

[1] Australian National Insect Collection, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[2] Environment, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[3] Department of Primary Industries and Regional Development, 3 Baron-Hay Court, South Perth, Western Australia 6151, Australia

[4] Department of Biodiversity, Conservation and Attractions, Perth Western Australia 6151, Australia

[5] Health and biosecurity, CSIRO, GPO Box 1700, Canberra, ACT 2601, Australia

[6] Faculty of Science and Technology, University of Canberra, Bruce, ACT 2615

[7] Department of Agriculture, Fisheries and Forestry, GPO Box 858, Canberra ACT 2601, Australia

Emails: James Bickerstaff: james.bickerstaff@csiro.au

Tom Walsh: tom.walsh@csiro.au

Leon Court: leon.court@csiro.au

Gunjan Pandey: gunjan.pandey@csiro.au

Kylie Ireland: kylie.ireland@dbca.wa.gov.au

David Cousins: david.cousins@dpird.wa.gov.au

Valerie Caron: Valerie.caron@csiro.au

Thomas Wallenius: Thomas.Wallenius@aff.gov.au

Adam Slipinski: adam.slipinski@csiro.au

Rahul Rane: Rahul.rane@csiro.au

Hermes Escalona: Hermes.escalona@csiro.au

*Correspondence address: James Bickerstaff, E-mail: james.bickerstaff@csiro.au; Hermes Escalona: hermes.escalona@csiro.au

Abstract

Bark and ambrosia beetles are among the most ecologically and economically damaging introduced

plant pests worldwide, with life history traits including polyphagy, haplodiploidy, inbreeding

polygyny and symbiosis with fungi contributing to their dispersal and impact. Species vary in host

tree ecologies, with many attacking stressed or recently dead trees, such as the globally distributed *E.*

similis (Ferrari). Other species, like the Polyphagous Shot Hole Borer (PSHB) *Euwallacea fornicatus*

(Eichhoff)*,* can attack over 680 host plants and is causing considerable economic damage in several

countries worldwide. Despite their notoriety, publicly accessible genomic resources for *Euwallacea*

Hopkins species are scarce, hampering better understanding of their invasive capabilities as well as

modern control measures, surveillance and management. Using a combination of long and short read

sequencing platforms we assembled and annotated high quality (BUSCO > 98% complete)

chromosome level genomes for these species. Comparative macro-synteny analysis showed an

increased number of chromosomes in the haplodiploid inbreeding species of *Euwallacea* compared to

diploid outbred species, due to fission events. This suggests that life history traits can impact

chromosome structure. Further, the genome of *E. fornicatus* had a higher relative proportion of

repetitive elements, up to 17% more, than *E. similis*. Additionally, metagenomic assembly pipelines

identified microbiota associated with both species including *Fusarium* fungal symbionts and a novel

Wolbachia strain*.* These novel genomes of haplodiploid inbreeding species will contribute to the

understanding of how life history traits are related to their evolution and will contribute to the

management of these invasive pests.

Keywords: Invasive species, bark beetles, biosecurity, weevils, insect genomics, Polyphagous Shot

Hole Borer

Significance

Scolytinae are significant forestry pests around the world and commonly translocated due to human

trade of wood and plant products. Life history traits including inbreeding and haplodiploidy are

attributed to their successful establishment in novel environments. *Euwallacea fornicatus* is widely

distributed and attacks a wide variety of live host trees. This study reports the genome of this species

and for, *E. similis*, which colonises dead host trees. The genome assemblies presented herein are

highly complete and scaffolded to pseudo-chromosomal level. Comparative analyses of these

genomes and of other Scolytinae highlight significant chromosomal rearrangements between

haplodiploid inbreeding *Euwallacea* and diploid outbreeding scolytinae species. Higher relative

proportions of transposable elements were identified *E. fornicatus*, which may promote the species'

ability to attack live host trees. These genomes are the first for haplodiploid beetles and will

contribute to the understanding of evolution of life history traits and the management of invasive

insects.

Introduction

Bark and ambrosia beetles (Curculionidae, Scolytinae) are some of the most ecologically damaging

and economically important insect plant pests, contributing to tree mortality through infestation and

the spread of pathogenic fungi (Hulcr & Stelinski 2017; Biedermann et al. 2019). Most bark beetles

are phloeophagous and obtain nutrition from the phloem tissue while some lineages, termed ambrosia

beetles, are xylomycetophagous and consume farmed symbiotic ambrosia fungi (Jordal 2014; Hulcr &

Stelinski 2017). Many species are commonly transported to novel areas through human mediated

movement of wood and plant products and over one hundred species have established outside of their

native range (Lantschner et al. 2020). Life history traits including polyphagy, haplodiploidy,

inbreeding polygyny and symbiosis with ambrosia fungi has been noted to contribute to the

establishment of introduced species (Jordal et al. 2001; Lantschner et al. 2020; Vilardo et al. 2022), all

of which characterise members of the tribe Xylebroini. Consequently, many such species have a

global native or introduced distribution and are high priority pests in many regions of the world.

The genus *Euwallacea* Hopkins include several well studied polyphagous tree pests (O'Donnell et al.

2016). The genus is composed of over 80 species which are mostly circumtropical in their distribution

(Storer et al. 2015). Species vary in their ecological relationships with host trees, with many species

colonising stressed, cut and recently dead hosts. One such example is *E. similis* (Ferrari) (Figure 1b),

a species native to South East Asia and possibly to Australia (Beaver et al. 2014), that has been

introduced to the Americas and tropical African countries (Gomez et al. 2018) with 145 known host

plants (Ruzzier et al. 2023). *Euwallacea fornicatus* (Eichhoff) (Figure 1a)*,* also known as the

Polyphagous Shot Hole Borer (PSHB) is considered one the most damaging Xyleborini species. It can

attack live hosts and is known to attack over 680 tree species (Gomez et al. 2019; Mendel et al. 2021;

Ruzzier et al. 2023) and is able to successfully reproduce in 168 species (Mendel et al. 2021; Ruzzier

et al. 2023). *Euwallacea fornicatus* is widely distributed and has been introduced to California

(Eskalen et al. 2012), Hawaii (Rugman-Jones et al. 2020), South Africa (Paap et al. 2018), Israel

(Mendel et al. 2012), Europe (Schuler et al. 2023) and, recently, in the city of Perth, Western

Australia (WA) where it is currently subject to an eradication campaign. This ambrosia beetle species

is associated with several pathogenic *Fusarium* fungi, including *F. euwallaceae* Freeman *et al.* (2013)

and *F. sp.* AF-18, and these fungi can induce wilt and dieback in host trees leading to tree mortality

(Freeman et al. 2013). Economic modelling has estimated the damages caused by this species to

$18.45 billion over a ten-year period in South Africa (De Wit et al. 2022) while in Perth WA the cost

of eradicating this species is estimated to be AUD$45 million and management of the species could

reach AUD$6.8 million per annum in 30 years (Cook & Broughton 2023).

The species *E. fornicatus* has been associated with the species *E. perbrevis* (Schedl), the Tea Shot

Hole Borer (TSHBa), *E. fornicatior* (Eggers), also the Tea Shot Hole Borer (TSHBb) and *E.*

kuroshiro (Gomez and Hulcr), the Kuroshio Shot Hole Borer (KSHB) representing a species complex

with similar morphology (Gomez et al. 2018; Smith et al. 2019) . Originally, these four species were

all considered to be *E. fornicatus*, however, recent molecular phylogenetic and taxonomic works have

stabilised their species status (Stouthamer et al. 2017; Gomez et al. 2018; Smith et al. 2019). Few

genomic resources exist for *E. fornicatus* and other Xyleborini, with much of the data available

restricted to mitochondrial *cytochrome oxidase I* (*COI*) and a handful of genes for phylogenetic

analyses (Stouthamer et al. 2017; Gomez et al. 2018; Bierman et al. 2022). Mitochondrial *COI*

phylogenies have identified 21 haplotypes of *E. fornicatus* with much of the diversity found in

Vietnam and Taiwan (Stouthamer et al. 2017). One haplotype, H33, has been identified in California,

South Africa and Israel and is thought to have originated from Vietnam (Stouthamer et al. 2017;

Bierman et al. 2022) however, other haplotypes, H35 and H38, have also been found in California and

South Africa, respectively, indicating that the species could have been introduced multiple times to

both countries (Stouthamer et al. 2017; Bierman et al. 2022). To date, the only sequence data

available for *E. similis* is restricted to *COI* barcodes in its native South East Asian distribution (Lynn

et al. 2021).

Fig 1. The ambrosia beetles A, *Euwallacea fornicatus* (PSHB) and B, *Euwallacea similis* whose genomes were sequenced herein. Photo taken by Hermes E. Escalona.

Genomic data are scarce for these widespread pests and further molecular resources are vital to understand the invasion patterns of *Euwallacea* species, their invasive capabilities, and exploring novel ways of controlling outbreaks and mitigating future harm. Access to high quality genomes has led to advancements in the management of invasive species, through facilitating accurate and time-efficient diagnostics and surveillance (Frey et al. 2022; McGaughran et al. 2024), understanding the origin of invasive species and invasion pathways (Rane et al. 2023) and identifying genetic and genomic variation related to invasion potential (Sillero et al. 2020; McGaughran et al. 2024). In this study we present chromosome-scale, annotated genomes of *E. fornicatus* and *E. similis* using newly developed long read sequencing technologies and performed comparative analysis of macrosynteny,

transposable elements, metagenomics and nuclear and mitochondrial phylogenetics. These genomes

will be valuable assets for tracking the invasion pathway, developing novel biosecurity management

plans and will provide essential insight into comparative of invasive species genomics.

Results

Genome assemblies and evaluation

In total 98.23Gb of high-quality Pacific Biosciences High Fidelity (PacBio HiFi) long sequence reads

with a mean N50 of 11.6Kb were obtained to assemble four genomes of two *Euwallacea* species

(Table S1). Two libraries of both species were sequenced and are herein referred to as EFF26 and

EFF42 (*E. fornicatus*) and ESF13 and ESF15 (*E. similis*). The estimated genome size for *E. fornicatus*

was between 192.7 Mb and 208.9 Mb, while estimates of 170.7 Mb and 181.8 Mb were made for *E.*

similis and all are predicted to be highly homozygous (het: 0.012% – 0.112%), and all libraries

consistently predicted similar results (Figure S1). All genomes were assembled with HiCanu (Nurk et

al. 2020) as these assemblies closer to the estimated genome size compared with hifiasm (Cheng et al.

2021), and assembly quality metrics were similar between assemblers. Genomes were assembled

with, post sequence read filtering and subsampling, 6.5Gb for EFF26 (29.58x, 0.35x subsample), 7Gb

for EFF42 (24.07x, 0.35x subsample), 5.63Gb for ESF13 (30.95x, not subsampled) and 8.09Gb for

ESF15 (44.78x, 0.15x subsample) of PacBio HiFi reads and 3.02Gb (*E. fornicatus*) and 3.3Gb (*E.*

similis) of Arima Hi-C paired-end reads (Table S1).

Scaffolding was performed with both YahS (Zhou et al. 2023) and SALSA2 (Ghurye et al. 2019).

While both scaffolders identified a similar number of pseudochromosome scaffolds for the *E.*

fornicatus assemblies, SALSA2 failed to break incorrectly joined contigs identified through Hi-C read

mapping for the *E. simils* scaffolds (Figures S2 – S5). As such, we present the pseudo-chromosomal

scaffolded assemblies produced through YahS followed by decontamination with the BlobToolKit

(Challis et al. 2020) and manual curation. In total, 39 pseudo-chromosomes were identified for the *E.*

Table 1. Genome summary statistics for the assemblies produced. Library subsampling strategy refers

to the level of subsampling. Mean HiFi coverage was calculated by dividing the number of base pairs

in the filtered subsampled HiFi libraries by the total assembly length. Mean Hi-C coverage was

calculated by dividing the number of base pairs in the filtered Hi-C libraries by the total assembly

length.

Species	*Euwallacea fornicatus*		*Euwallace similis*	
Library	EFF26	EFF42	ESF13	ESF15
NCBI Accession	GCA_04011564 5.1	GCA_04011565 5.1	GCA_03988120 5.1	GCA_03988121 5.1

Estimated Genome Size (Mb)	192.7	197.7	181.83	170.77
Library subsampling strategy	0.35	0.35	1	0.15
Subsampled seqs (K)	609.62	578.21	0.54	659.71
Subsampled bases (Gb)	6.5	7	5.63	8.09
Total assembly Length (Mb)	219.83	239.6	177.6	180.07
Longest assembly seq (Mb)	8.67	8.89	10.09	10.04
Scaffold number	185	179	128	140
Number of pseudo-chromosomes	39	39	37	37
N50 (Mb)	3.8	3.6	5.14	5.14
L50	20	23	13	13
N's per 100 kpb	0.36	0.42	2.49	1.9
Mean HiFi coverage (x)	29.58	31.80	30.95	44.78
Mean Hi-C coverage (x)	14.56	13.36	18.12	18.26
BUSCO				
Complete single copy	2080 (97.9)	2079 (97.9)	2084 (98.1)	2084 (98.1)
Complete duplicated	8 (0.4)	8 (0.4)	9 (0.4)	9 (0.4)
Fragmented	15 (0.7)	15 (0.7)	7 (0.3)	8 (0.4)
Missing	21 (1)	22 (1)	24 (1.2)	23 (1.1)

fornicatus (EFF26, Figure S6; and EFF42, Figure S7) assemblies. Both EFF26 and EFF42 assemblies

were larger than the estimated genome sizes, at 219.83 Mb and 239.6 Mb, respectively. For both *E.*

similis assemblies 37 pseudo-chromosomes were identified (ESF13, Figure S8; and ESF15, Figure

SS9). The ESF13 assembly was smaller than the estimated genome size, at 177.6 Mb, while the

ESF15 genome was larger than estimated at 180.07 Mb. All assemblies were found to be highly

complete and contained over 98% complete single copy BUSCO sequences (Table 1, Figure S10).

Genome annotation

Genome annotation with the NCBI Eukaryotic Annotation Pipeline predicted 24,417 transcripts and

12,360 genes for the EFF26 *E. fornicatus* genome. Annotation for the ESF13 *E. similis* genome

resulted in 19,838 transcripts and 11,978 genes predicted. Protein coding genes occupied 46 Mb of the

E. fornicatus genome and 38 Mb of *E. similis,* 21% of both assembly lengths, and their density varied

across chromosomes for both species (Figure 2). Annotations were largely complete with >97%

complete single copy BUSCOs (Table 2). The number of genes annotated for both *Euwallacea*

species were lower than those annotated for all other Scolytinae species, roughly 17 – 81% fewer

(Navarro-Escalante et al. 2021; Powell et al. 2021; Liu et al. 2022; Wang et al. 2023).

Fig 2. Distribution and density of annotated protein-coding genes along the pseudo-chromosome

scaffolds for A, *Euwallacea fornicatus* and B, *Euwallacea similis*.

Mitochondrial and nuclear phylogenetics

Phylogenetic analysis of mitochondrial haplotypes identified that the population of *E. fornicatus* in

Perth WA belong to haplotype H38, with the *COI* barcode identical to populations native in China and

Vietnam, and also to the invasive population in South Africa. The H38 haplotype clustered with

Haplotypes 37, 39 – 42, and were found to be phylogenetically distinct from the more prevalent

invasive H33 haplotype (Figure S11). Phylogenomic analysis using BUSCOs of published Scolytinae

genomes and those of the reference *Euwallacea* yielded a topology similar to that reported by Pistone

et al. (Pistone et al. 2018) and all nodes were strongly supported (Figure 3).

Table 2. Protein and repeat annotation statistics for the assemblies produced. Proteins were identified

and annotated by the NCBI Eukaryotic Annotation Pipeline and completeness assessed with BUSCO.

Repeats were identified by EarlGreyTE.

Species	*Euwallacea fornicatus*	*Euwallacea similis*
Library	EFF26	ESF13
NCBI Accession	GCF_040115645.1-RS_2024_07	GCF_039881205.1-RS_2024_07
Predicted transcripts	24,417	19,838
Predicted genes	12,369	11,978
Mean gene length (bp)	12,012	11,389
Total CDS length (bp)	46,577,826	38,293,762
% genome covered by genes	21%	21%
Mean transcripts per gene	2.02	1.67
Mean exon number per transcript	9.31	8.99
BUSCO		
Complete single copy (bp) (%)	2,079 (97.8)	2,083 (98)
Complete duplicated (bp) (%)	17 (0.8)	12 (0.6)
Fragmented (bp) (%)	9 (0.4)	11 (0.5)
Missing (bp) (%)	19 (0.9)	17 (0.8)
Repeats		
DNA (bp) (%)	68,089,859 (30.9)	18,861,021 (10.6)
LINE (bp) (%)	5,329,747 (2.4)	3,325,503 (1.9)
LTR (bp) (%)	7,594,481 (3.5)	13,488,481 (7.6)
Unclassified (bp) (%)	18,446,822 (8.4)	21,972,489 (12.4)
Other (bp) (%)	4,200,730 (1.9)	1,296,582 (0.7)
Total masked (bp) (%)	103,661,639 (47.1)	58,944,136 (33.2)

209 *Repeat element and chromosomal variation across Scolytinae*

Comparative analyses of repeat and transposable elements found that their relative proportion in the genomes of Scolytinae varied across lineages. Roughly half (47 – 50%) of the *E. fornicatus* assemblies and over one third (33 - 34%) of the *E. similis* assemblies were masked by the EarlGreyTE pipeline (Table 2, Figure 2, Figure S12). DNA transposons and long terminal repeats (LTR) were the most abundant repeat elements identified, with unclassified elements only accounting for a small proportion of all annotations (Table 2). Few rolling circle elements were identified in all assemblies and SINEs were only found in the *E. similis* assemblies. Kimura 2 divergence landscapes of repetitive elements indicate a recent burst of DNA transposons and LTR elements, while LINE and rolling circle elements are experiencing ongoing proliferation in the *Euwallacea* assemblies (Figures S13 – 16). The relative proportion of annotated TEs varied throughout Scolytinae assemblies, with over 60% of the *T. bicolor* assembly occupied by TEs, and the assembly of *D. ponderosae* possessed the lowest relative proportion of annotated TEs (25%). The relative proportions of the types of TEs also varied significantly throughout Scolytinae as well, and TE annotations were largely composed of unclassified elements, DNA transposons and LINEs. All published Scolytinae assemblies possessed greater amounts of Satellite sequences than the *Euwallacea* assemblies (Figure 3). Lastly, the Entiminae outgroup *P. cervinus* had a larger proportion of the genome of annotated TEs compared to Scolytinae.

Fig 3. Maximum-likelihood phylogenetic tree pseudo-chromosome level assemblies for Scolytinae based using 2,108 single copy complete BUSCO sequences. All nodes were strongly supported with 100 bootstrap support. Scale bar indicated 0.03 substitutions per nucleotide position. To the right of the tips are the relative proportion of repetitive element composition as annotated in the species' genome.

The chromosome number and synteny of haplodiploid inbreeding *Euwallacea* species greatly varied compared to diploid outbreed Scolytinae species. The assemblies of *E. fornicatus* and *E. similis* were more syntenic with one another than other Scolytinae (Figure S17). Largely, synteny was retained throughout the Scolytinae assemblies with several translocations of genomic material. However, extensive translocations, rearrangements, and fissions of syntenic blocks were identified between the assemblies of *Euwallacea* and other Scolytinae. Further, the X sex-chromosome was identified to be synentic throughout all assemblies, however, blocks corresponding to these scaffolds were broken apart in the *Euwallacea* assemblies (Figure 4).

Fig 4. Ribbon plot showing the synteny and structural and chromosomal rearrangement in Scolytinae chromosomal assemblies. Syntenic blocks of single copy complete BUSCO sequences link pseudo-chromosome scaffolds between assemblies. Black bars represent individual pseudo-chromosome scaffolds and are labelled according to the name provided in the assembly. Scaffolds are ordered following the *Dendroctonus ponderosae* assembly order and the lengths of the bars representing scaffolds indicate total number of BUSCOs and not scaffold length. Pseudo-chromosome scaffold labels follow how they were published either on NCBI or their respective publications, the X-chromosome label in parentheses is inferred based on syntenic linkages. Coloured lines connect orthologous genes across scaffolds and the colour code follows the Bilaterian-Cnidarian-Sponge Linkage Groups (BCnS LGs) as in (Schultz et al. 2023).

Metagenomics

The microbial diversity and composition varied between the two *Euwallacea* species. The two *E.*

similis libraries produced more Metagenome-assembled genomes (MAG) contigs (730 and 1,200)

than the two *E. fornicatus* libraries (103 and 109). Across all libraries, many more Bacteria were

identified than other microbial Kingdoms, with Actinomycetota and Pseudomonadota highly

prevalent (Figure S18). Several complete circularised Bacteria assemblies were also identified across

all libraries (Table S2), including *Wolbachia* for both *E. similis* libraries. Multi-locus Sequence

Typing (MLST) profiling identified this *Wolbachia* strain closely matching *w*Ei, which infects *E.*

interjectus (Kawasaki et al. 2016), however there were SNPs in the *hcpA* and *gatB* loci. Consequently,

this represents a novel strain, herein referred to as *w*Es. Several fungal taxa were identified for both

species (Figure S18), and mitogenomic contigs of *Fusarium* belonging to the *Euwallacea Fusarium*

Clade (EFC) were identified by BLAST for all libraries except for ESF13. The only fungi found in *E.*

fornicatus MAGs were *Fusarium*, while a richer diversity of fungi were found in association with *E.*

similis, with 13 genera classified including *Leptographium, Ophiostoma, Ceratocystis* and *Sporothrix*

(Table S2). Several viral contigs were also identified in both species, all belonging to Caudoviricetes.

Discussion

Genome synteny across Scolytinae

We sequenced, assembled and annotated the genomes of *Euwallacea fornicatus* and *E. similis*. These

genomes are highly contiguous, complete and accurate as we sequenced the libraries of two female

specimens per species and employed multiple assembly and scaffolding pipelines to validate their

quality. The *E. fornicatus* assemblies presented herein were accepted and released on the NCBI

GenBank on the same day as two other contig-level assemblies of the same species from South Africa

(GCA_040114445 & GCA_040114495), from a female and male, respectively. The assembly sizes of

these contig-level genome assemblies are similar to the chromosome-scaffold assemblies presented

here, further validating the quality of the assemblies presented herein. These genomes represent the

first chromosome scaffold genomes for these pests and for a haplodiploid inbreeding beetle lineage.

They are among the smallest Scolytinae genomes published and contain the highest number of

pseudo-chromosomal scaffolds. Additionally, about twelve thousand protein coding genes were

predicted for these *Euwallacea* assemblies, representing 17 – 81% fewer proteins compared to

previously published Scolytinae genomes (Navarro-Escalante et al. 2021; Powell et al. 2021; Liu et al.

2022; Wang et al. 2023).

The number of pseudo-chromosomal scaffolds identified for both *Euwallacea* species (n = 37 – 39)

were higher than the typical number of chromosomes in Scolytinae (n = 7 – 16) (Virkki & Smith

1978). Chromosome number variation has previously been identified across and within genera of

scolytines. For example, the karyotype of all 19 *Dendroctonus* species have been characterised, with

chromosome numbers ranging from 2n = 12 – 30, and chromosomal fusion resulting in chromosome

number reduction (Lanier 1981; Six & Bracewell 2015). Similarly, chromosome number variation has

also been identified across 32 species of *Ips*, with numbers ranging from 2n = 16 – 32 (Virkki &

Smith 1978). In Curculionidae, a larger variation in chromosome numbers have been characterised in

those that are parthenogenetic due to polyploidy (Virkki & Smith 1978; Blackmon & Demuth 2015).

For example, parthenogenetic pseudogamy resulting in triploid females have been characterised in *Ips*

acuminatus Gyllenhal and *I. tridens* Mannerheim (Lanier & Kirkendall 1986). Similar observations

have been made for several species of obligate inbreeding haplodiploid Xyleborini (Takenouchi &

Takagi 1967; Virkki & Smith 1978). Species of the genus *Xylosandrus* Reitter have fewer

chromosomes than other Scolytinae, with *X. germanus* (Blandford) 2n = 8 and *X. compactus*

(Eichhoff) 2n = 10 (Takenouchi & Takagi 1967). In contrast, the chromosomes of *Anisandrus dispar*

(F.) are small in size and numerous, n = 40 (Virkki & Smith 1978), similar to the *Euwallacea* species

genomes.

We identified extensive chromosomal rearrangements between haplodiploid inbreeding and diploid

outbreeding Scolytinae. The synteny of autosomal chromosomes and the ancestrally conserved X

chromosome between diploid outbreeding species are largely conserved across genera, with syntenic

blocks either translocated within or across scaffolds. Further, despite the difference in chromosomal

numbers between *D. ponderosae* (2n = 12) and *D. valens* (2n = 14) (Six & Bracewell 2015), synteny

across autosomal and X-chromosomes are largely conserved, identified herein and previously (Liu et

al. 2022). Furthermore, extensive chromosomal rearrangement characterised the variation in genomic

architecture between *Euwallacea* and diploid outbreeding species, with widespread translocation

events shuffling syntenic blocks and fission events breaking up autosomal chromosomes. The

conserved X chromosome was also found to be broken up through fission in the *Euwallacea*

assemblies. This may relate to the apparent lack of sex chromosomes in other Xyleborini species as

observed by Takenouchi & Takagi (Takenouchi & Takagi 1967) in male germ cells. Comparative

analyses of Coleopteran karyotypes have identified that small effective population sizes and genetic

drift greatly impact chromosome number variation across species (Blackmon et al. 2024). We

speculate that the low genetic diversity and high homozygosity due to obligate inbreeding in

Euwallacea may contribute to the increased chromosome number identified in these assemblies.

Comparative analyses indicate that the *E. fornicatus* genome has one of the highest proportions of

repeat and transposable elements relative to total assembly among sequenced Scolytinae species, with

roughly 50% of the assembly composed by TEs. While the relative proportion of these elements in the

genome of *E. similis* only occupy about 30% of the total assembly*,* DNA transposons are among the

most prevalent transposable elements in both *Euwallacea* species. DNA transposons have been

identified to play a significant role in Eukaryotic genome and chromosomal evolution (Feschotte &
Pritham 2007). Moreover, it has been identified in plants (Ren et al. 2018) that genomic
rearrangement and recombination can result in net sequence loss at newly formed telomeric regions in
the chromosome. Given that we have identified bursts in the proliferation of DNA transposons in the
genomes of *Euwallacea*, these repetitive elements could be involved in the widespread structural
variation, high number of chromosomes and smaller genome size and content in this genus.

Co-evolution with symbiotic ambrosia fungi could also contribute to chromosomal rearrangements in
Xyleborini, as observed in *Euwallacea* assemblies. Genomic erosion is particularly prevalent in the
genomes of symbionts that provide nutrition to their hosts, and genes of biosynthesis pathways
producing amino acids obtained from the host are commonly lost (McCutcheon & Moran 2012; Sloan
& Moran 2012). This has been identified to occur in host animals as well; in the deep-sea bone-eating
worm, *Osedax frankpressi* Rouse, Goffredi & Vrjenhoek (Annelida: Polychaeta), endosymbionts
provide key nutritional biosynethetic pathways which has subsequently resulted in host genome
reduction and gene loss (Moggioli et al. 2023). Similarly, in socially parasitic ant species, gene loss of
olfactory receptor repertoires corresponding with reduced social behavioural has evolved repeatedly
(Schrader et al. 2021). Host genome rearrangements have also been reported in ants, with high levels
of structural rearrangements, loss of synteny across assembled genomes, gene family contraction,
expansion and a gain of novel genes identified in the *Atta* fungal farming ant lineage (Nygaard et al.
2011, 2016). Similar processes could have acted on the genomes of fungal farming Scolytinae in the
coevolution with their symbionts, leading to the observed small genome size, genomic rearrangements
and loss of synenty with other Scolytinae. It should be noted that the evolution of obligate inbreeding
and haplodiploidy, in Dryocoetini, seems to precede the development of the ambrosia fungal
symbiosis in Xyleborini (Jordal & Cognato 2012), and as such, further chromosome scale assemblies
of non-fungus farming inbreeding haplodiploid species (Dryocoetini) are needed to assess the impact
of the ambrosia symbiosis on chromosome structure.

Genomic resources for biosecurity

The genomes assembled for *Euwallacea* will be essential resources for the management of these pests.
A vital component in managing an invasive species is understanding where the species originates and
how it arrived. Through mitochondrial haplotyping, we have identified that the population of *E.*
fornicatus introduced to Perth, Western Australia corresponds to the H38 haplotype, which has also
been introduced to California and South Africa. While further markers are needed for population
genomic analyses in order to track the invasion pathway from the native to the introduced range (Rane
et al. 2023), the assembled genomes will be vital in understanding population differentiation at the
gene and genomic level. These genomes will also be a vital resource in the diagnostics and detection

of these pests. While the mitochondrial genome for both species have already been published (Wang

et al. 2020; Guo et al. 2023), the ability to identify nuclear markers or sequences will be important for

the diagnostics and detection (such as through eDNA techniques) of the pests. Diagnostic protocols

exist for the detection of *E. fornicatus* through identification of the symbiont *Fusarium euwallaceae*

(De Jager & Roets 2022), however, the protocol operates on the detection of fungus in host plant

woody tissues, which indicates that the species is already present and likely reproducing in a given

locality. Using eDNA techniques and these reference genomes, these beetle pests could be detected

early, increasing the chances of eradication or containment, and reducing the time and resources

necessary to achieve this. Additionally, such methodologies could also demonstrate the absence of

this pest in sampled areas allowing for cost effective determination of management and eradication

outcomes.

Beyond tracking, detection and diagnostics, these genomes will also provide needed information for

genome guided management of these pest. The endosymbiont gram-negative bacteria *Wolbachia* has

been successfully deployed as a control agent for insect pests (Ross et al. 2022) and may prove useful

in the control of *Euwallacea* pests. While strains have not yet been identified to be associated with *E.*

fornicatus, further sequencing and MLST identification could identify naturally present strains, or

strains from closely related species (such as *w*Es identified in this study) that could be artificially

transferred (Xi et al. 2005). However, as noted in previous studies, the impacts of *Wolbachia* on

haplodiploid inbreeding Scolytinae are not yet fully understood (Kawasaki et al. 2016; Bickerstaff et

al. 2023). Gene drive through CRISPR gene editing may be an avenue to control populations of

invasive *E. fornicatus* and *E similis.* As demonstrated in invasive wasps (Lester et al. 2020; Meiborg

et al. 2023), the ability to control population growth with this technique is limited due to

haplodiploidy, with only a single gene targeting female fertility identified as useful and infertile

individuals were not as competitive as fertile counterparts. Outbreeding is rare in the inbreeding

Scolytinae lineages and comes at a fitness cost (Peer & Taborsky 2005), and as such the penetration

of gene drive through populations may be limited. Further insights are needed to first identify

candidate genes in these genomes and then test the suitability of this management approach.

These genomes will also be of value in understanding mechanisms driving species invasion and pest

impact from a genomic perspective. Preliminary comparative analyses of transposable element

variation indicate that the relative proportion of genome content is similar to other beetle genomes

(Petersen et al. 2019; Gilbert et al. 2021) however, a higher number of TEs are present in the *E.*

fornicatus genome than in the *E. similis* genome. Transposable elements have been identified to be

adaptive in species (Gilbert et al. 2021) and promote genetic diversity and rapid adaption in invasive

species. The larger diversity of TEs present in the *E. fornicatus* genome may be associated with the

species ability to infest a wide range of live host trees. Further comparative genomic analysis, such as

gene family evolution (Powell et al. 2021), structural variation (Stapley et al. 2015; North et al. 2021),

and pangenomics (Gerdol et al. 2020) of both species and their symbionts may yield greater insights

into why these species are such successful invaders compared to other haplodiploid inbreeding

ambrosia beetles, understanding variation in host tree ecology and why *E. fornicatus* is more

destructive than other members *Euwallacea*.

Conclusions

Pest species of *Euwallacea* are globally invasive and can cause significant ecological and economic

impacts in the ranges where they are established. Our study presents the first chromosome-level

genome assemblies of these pests and also for haplodiploid inbreeding beetles. These assemblies offer

a deeper understanding of the genome structural variation related to species ecology and also serve as

essential resources for biosecurity management and control strategies. The comparative analyses of

repeat and transposable elements, as well as the macro-synteny of chromosome assemblies, shed light

on the potential impacts of haplodiploidy and inbreeding on genome evolution in these beetles. Our

findings underscore the importance of high-quality genome assemblies in advancing our knowledge

of invasive species and contributing to their effective management and control.

Material and methods

Sample collection, DNA extraction and sequencing

Male and female specimens of *E. fornicatus* were collected by the Department of Primary Industries

and Regional Development (DPIRD) WA from felled *Acer negundo* trees in Perth WA in October

2022. Male and females were flash frozen by storing them on dry ice and shipped to CSIRO Black

Mountain Canberra Australian Capital Territory and stored at -80°C. *Euwallacea similis* larvae and

adult females were collected from a fallen *Ficus* sp. branch in Eureka, northern New South Wales in

April 2023. Specimens were transported from the field at 10°C and flash frozen by storing them at -

80°C. Vouchers of both species have been deposited in Australian National Insect Collection (ANIC)

under the collection codes ANIC-25-086063 – ANIC-25-086073.

Female specimens were used for DNA extraction due to their larger size compared to haploid males.

First, genomic DNA (for short read sequencing) of both species was extracted using a standard

Qiagen DNEasy kit approach. We then used a single specimen from each species for high molecular

weight DNA extraction with the Qiagen MagAttract kit following a modified protocol. Samples were

incubated overnight at 56°C and lysates were agitated at 300 rpm. Following this, DNA was

suspended in two elutions of 100 µl of Buffer EB. The concentration of all extracts were assessed

using a DeNovix spectrophotometer and a Qubit Fluorometer with the Qubit dsDNA High Sensitivity

Assay kit. Fragment size distribution and genomic DNA was assessed on an Aligent Fragment

Analyzer system with the high sensitivity genomic DNA kit DNF-488.

For short read sequencing, sequencing libraries with 150 bp paired end reads were prepared and

sequenced on the Illumina NovaSeq 6000 system at Genewiz (Suzhou, China). For long read

sequencing, high molecular weight (HMW) DNA extracts were sequenced on the PacBio HiFi

circular consensus Revio system at the AGRF in Brisbane, Queensland. For two HMW extracts of

each species, gDNA was mechanically sheared, followed by SMRTbell Express 2.0 library

construction. Short DNA fragments were removed using BluePippin. Barcoded libraries were then

sequenced using one SMRT cell for each species on the PacBio Revio system; the libraries for *E.*

fornicatus are herein referred to as EFF26 and EFF41, and the libraries for *E. similis* are herein

referred to as ESF13 and ESF15.

Genome size estimation, assembly and scaffolding

Genome size and heterozygosity estimates were performed with all sequence reads; kmers were

counted with jellyfish (Marçais & Kingsford 2011) and genome size and heterozygosity were

estimated with GenomeScope v2 (Ranallo-Benavidez et al. 2020). HiFi reads were then filtered for

HiFi adapters with HiFiAdapterFilt (Sim et al. 2022) and subsampled with seqtk into randomly

sampled subsets, ranging from 12x, 15x, 20x, 35x and 50x, of the estimated genome size, for all four

libraries. Contigs were assembled with these subsets and the full long read dataset using HiCanu

v2.1.1 (Nurk et al. 2020), with the genome size estimates of 190 Mb for both EFF26 and EFF42, and

170 Mb for both ESF13 and ESF15. Contigs were also assembled with hifiasm v0.19.6-r595 (Cheng

et al. 2021) using the -l0 flag to disable purge duplication as these genomes are highly homozygous

and presumably inbred. Misassembled contigs were assessed and corrected with CRAQ v1.0.9a (Li et

al. 2023) and haplotigs and contig overlaps were removed with purge_dups v1.2.5 (Guan et al. 2020).

The quality of the corrected contig assemblies were assessed with QUAST v5.2.0 (Mikheenko et al.

2018) and BUSCO v5.5.0 (Manni et al. 2021) using the endopterygota_odb10 dataset and the

tribolium2012 dataset was specified for Augustus (Stanke et al. 2008) training.

Proximity ligation information was generated with Arima HiC 2.0 kits using tissue from a single

individual from both species at the Biomolecular Resource Facility – John Curtin School of Medical

Research (BRF-JCSMR) Australian National University (ANU). In short, tissues were fixed in

formaldehyde, DNA was extracted and then digested with Arima restriction enzymes. Overhangs of

digested DNA were then end repaired and marked with biotinylated nucleotides. The ends were then

ligated together, and DNA strands were de-crosslinked and sheared for Illumina PE 150bp

sequencing. The *E. fornicatus* library was sequenced on an Illumina NovaSeq 6000 for 300 cycles and

the library for *E. similis* was sequenced on an Illumina NextSeq 2000 for 300 cycles.

The Hi-C reads were mapped to draft contig assemblies following the Arima Genomics Mapping

Pipeline v03. First, 5' ends of the Hi-C reads were trimmed by 5 base pairs to remove over-

represented 3bp barcode UMI sequences and 2 dark bases from the 5' end. Trimmed reads were then

mapped independently (as single-ended reads) using BWA-MEM2 (Vasimuddin et al. 2019) with the

-5SP flag. Only the 5' ends were retained using the "filter_five_end.pl" script provided by the Arima

Genomics Mapping Pipeline, as the ends of the DNA were ligated together prior to sequencing, 3'

ends can originate from the same contiguous strands of DNA as the 5' end of the mate-pair, which can

manifest a ligation function and form chimeric reads after mapping. Following this, single-ended

mapped reads were then paired using the "two_read_bam_combiner.pl" script provided by the Arima

Genomics Mapping Pipeline and mapped reads with a mapping quality (MAPQ) < 10 were discarded

and sorted with samtools (Danecek et al. 2021). Lastly, PCR-duplicates were removed with Picard

Tools v3. Assemblies were then scaffolded using YahS v1.1a (Zhou et al. 2023) and assessed with

QUAST v5.2.0, and BUSCO v5.5.0 (Manni et al. 2021). We then compared the best scaffolded

assembly with an analogous scaffolding run using SALSA2 v2.3 (Ghurye et al. 2019) and selected the

best scaffolded assembly using QUAST and BUSCO metrics for downstream curation with the

Juicebox Assembly Tools (JBAT) v1.b (Durand et al. 2016). Contaminant scaffolds, defined as

contigs identified not to be of Arthropoda origin through BLAST, sequences with >15x coverage, and

GC <0.3 and >0.45 were removed using the BlobToolKit v4.2.1 (Challis et al. 2020). Hi-C reads were

then mapped to the decontaminated scaffolds using BWA-MEM2 following the Arima Genomics

Mapping Pipeline v03 and were reviewed with JBAT v1.b (Durand et al. 2016). Illumina short read

and PacBio HiFi reads were also mapped to the decontaminated scaffolds and all three mapped

datasets were inspected using JBrowse v2.10.3 to identify any potential misassembles (identified by

poor read alignments and regions of low coverage). We assessed the completeness of the final

scaffolded assemblies again with QUAST v5.2.0 (Mikheenko et al. 2018) and BUSCO v5.5.0 (Manni

et al. 2021). Lastly, the scaffolds of each library of the same species (i.e. both *E. fornicatus* and *E.*

similis assemblies) were renamed and reoriented to be syntenic with one another using BUSCOs to

define synteny and orientation. We identified pseudo-chromosome scaffolds by scaffolds that were >

500Kb in length and had an average Hi-C coverage > 30x. Pairs of syntenic pseudo-chromosomes

were defined by length and by BUSCO composition and these pseudo-chromosomes were named in

order of length by the EFF26 assembly for *E. fornicatus* and the ESF13 assembly for *E. similis*.

Lastly, assemblies were assessed for any remaining contamination with FCS-GX v0.5 (Astashyn et al.

2024) and any contaminants identified were removed prior to submission to the NCBI GenBank.

Genome annotation

Total RNA was extracted from ten adult males and females of each species. RNA was extracted using

the Zymo Research Quick-RNA Miniprep Plus kit following the manufacturers tough-to-lyse sample

preparation procedure. RNA quantity and quality was then assessed using a DeNovix

spectrophotometer and Aligent Fragment Analyzer system with the HS RNA kit DNF-472. For one

extract of each species and sex strand-specific libraries were prepared by GENEWIZ from Azenta

Life Sciences and sequenced on an Illumina NovaSeq 6000 PE150.

Genomes were annotated by The NCBI Eukaryotic Annotation Pipeline. In short, the NCBI GenBank

reference assemblies of *E. fornicatus* (EFF26) and *E. similis* (ESF13) were masked with

WindowMasker (Morgulis et al. 2006), and available transcripts, RNA-seq (which includes data

produced herein), and protein data from RefSeq were mapped to masked genomes. Transcripts and

proteins from the RefSeq and GenBank databases were aligned with BLAST and the short-read RNA-

seq data was aligned with STAR (Dobin et al. 2013). The final set of genes were built by evaluation

against known RefSeq transcripts, curated RefSeq genomic alignments and the most highly supported

models that were predicted by Gnomon. Protein naming, locus type and GeneID assignment were

made by the NCBI Eukaryotic Annotation Pipeline. Completeness of the annotated gene models was

then assessed with BUSCO v5.5.0 (Manni et al. 2021) run in protein mode using the

Endopterygota_obd10 dataset and the tribolium2012 dataset was specified for Augustus (Stanke et al.

2008) training. The distribution and density of genes along chromosomes of EFF26 and ESF13 were

visualised with RIdeogram (Hao et al. 2020).

Transposable Elements

Chromosome level genomes of *Taphrorychus bicolor* (Telfer et al. 2024), *Dendroctonus ponderosae*

(Keeling et al. 2022)*, D. valens* (Liu et al. 2022)*, Ips nitidus* (Wang et al. 2023) and the outgroup

Polydrsus cervinus (Entiminae) (Barclay et al. 2023) were downloaded from NCBI or ensemble. First,

repeat annotation was performed with EarlGreyTE (Baril et al. 2024) to identify, classify and mask

repeat regions across published genomes and the reference assemblies of *E. fornicatus* (EFF26) and *E.*

similis (ESF13). Following repeat annotation, kimura-2 divergences between TE classes were

calculated using the parseRM.pl scrip with the -l flag (Kapusta 2017).

The composition of TEs were compared phylogenetically across Scolytinae genomes downloaded and

the *Euwallacea* reference genomes (EFF26 & ESF13). We used BUSCO to call 2,108 single copy

sequences which were aligned with MAFFT v7.490 (Katoh 2002) and poorly aligned bases were

removed with TrimAl v1.4.1r22 (Capella-Gutiérrez et al. 2009). Alignments were then concatenated

with phykit v1.19.0 (Steenwyk et al. 2021) and phylogenetic analyses were conducted in IQ-tree

v2.3.2 (Minh et al. 2020). Substitution models were estimated with ModelFinder Plus

(Kalyaanamoorthy et al. 2017) using a greedy strategy with gene merging and bootstrap support was

calculated with nonparametric bootstrap with 100. The relative proportions of TEs were then

annotated on the tips of the phylogeny.

Chromosomal synteny across Scolytinae

Macrosynteny and chromosomal variation across published Scolytinae genomes and the *Euwallacea*

reference genomes (EFF26 & ESF13) was performed with the Oxford Dot Plots (opd) pipeline

(Schultz et al. 2023). Previously used BUSCO sequences were used to describe the relationship

between linkage groups and chromosomes between species. Reciprocal best hits between BUSCOs

were identified with diamond blastp and the homology of scaffolds between species were inferred

with pairwise p-values calculated with Fisher's exact test. Pairwise species dot plots syntenic linkage

ribbon plots for all species were generated using scripts available in odp. Linkage colours in the

ribbon plot follows the Bilaterian-Cnidarian-Sponge Linkage Groups (BCnS LGs) as in (Schultz et al.

2023).

Mitochondrial phylogenetics

To identify the haplotype of *E. fornicatus* present in Perth WA, previously published *COI* genes

(Stouthamer et al. 2017; Gomez et al. 2018; Bierman et al. 2022) of PSHB, TSHBa&b and KSHB

were downloaded from NCBI. The mitochondrial genome for all libraries were assembled using the

MitoHiFi v3.2.1 pipeline (Uliano-Silva et al. 2023). Full length *COI* genes from these assemblies

were then aligned with the sequences from NCBI using MAFFT (Katoh 2002). Third codon positions

were partitioned in the alignment and IQ-Tree v2.3.2 (Minh et al. 2020) was used to estimate the

phylogeny, substitution models were identified with ModelFinderPlus (Kalyaanamoorthy et al. 2017)

and bootstrap support was calculated with UFBoot (Hoang et al. 2018) with 1000 replicates.

Metagenomics

To identify the microbial communities associated with *E. fornicatus* and *E. similis*, metagenomes

were assembled from the total hifi sequence read dataset for each library. Metagenome-assembled

genomes (MAGs) were assembled using hifiasm-meta (Feng et al. 2022) with default settings.

BLAST was used to identify contigs and MAG identity was explored and visualised with the

BlobToolKit v4.2.1 (Challis et al. 2020). Host contigs and those that were unidentified were filtered

out of the dataset. A closed *Wolbachia* contig in was found in both *E. similis* libraries, and so the

strain was identified using MLST approaches implemented in the PubMLST database (Jolley et al.

2018).

Data Availability

All genome and raw sequencing reads produced in this have been deposited with the NCBI, under the

BioProject: PRJNA1086994. Specific accession numbers for raw sequence reads and genome

assemblies are available in Table S1 and Table 1, respectively. Vouchers of the specimens are

deposited in the ANIC, CSIRO under accession numbers: ANIC-25-086063 – ANIC-25-086073.

Additional Files

Supplementary File 1. The supplementary figures produced through assembly QC and analyses that

complement the main text.

Supplementary Table S1. Sequence read output of all libraries sequenced.

Supplementary Table S2. Classes of Repeat and Transposable Elements identified by EarlGreyTE.

Competing Interests

The authors declare no competing interests.

Funding

The work was supported by the ResearchPlus CSIRO Early Research Career Fellowship funding, the

Zimmerman Trust and ANIC-NRCA-CSIRO.

Author's Contributions

JB, HE, TW, RR, VC, AS and TW conceptualised the research. DPIRD and KI provided PSHB

specimens. TW, RR, LC and GP provided advice for HWM DNA extraction and genome assembly

pipelines. JB and HE interpreted the results. JB conducted the field and laboratory work, outlined the

bioinformatic pipelines, collected and analysed the data, and wrote the manuscript with input from all

authors. All authors agree to the submitted manuscript.

Acknowledgements

We are grateful to the Polyphagous Shot Hole Borer response team from the Department of Primary

Industries Regional Development Western Australia for providing samples, including Basman Al-

Jalely and Emad Alshwany from the Reproductive Biology Laboratory. Bonnie Koopmans and

Debbie Jennings for their help during fieldwork. We would like to thank Andreas Zwick, James

Nicholls, Diana Hartley and Andrew Gock (CSIRO Black Mountain) for access to laboratory

facilities, Peter Holland (Oxford University, UK) and Mara Lawniczak (Darwin Tree of Life, UK) for

advice on laboratory protocols. We thank Maxim Nekrasov and Carolina Correa Ospina (Australian

National University Biomolecular Resource Facility), and Trent Peters and Oliver Distler (Australian

Genome Research Facility), for their technical support. The authors thank John G. Oakeshott

(CSIRO) and Robert Waterhouse (University of Lausanne) for their generous advice and David

Yeates (CSIRO Black Mountain) for his support of the project. We would like to thank Andy Bachler

for advice on genome scaffolding and Ondrej Hlinka (CSIRO) for assistance with High Performance

Computing. We also thank Cecile Gueidan (CSIRO) for comments on an earlier version of this

manuscript. Thanks to CSIRO research Office and the Zimmerman Trust for supporting this research.

References

Astashyn A, et al. 2024. Rapid and sensitive detection of genome contamination at scale with
FCS-GX. Genome Biol. 25(60). https://doi.org/10.1186/s13059-024-03198-7.

Barclay MVL, et al. 2023. The genome sequence of a weevil, *Polydrusus cervinus* (Linnaeus,
1758). Wellcome Open Res. 8(563). https://doi.org/10.12688/wellcomeopenres.20414.1.

Baril T, Galbraith J, Hayward A. 2024. Earl Grey: A fully automated user-friendly
transposable element annotation and analysis pipeline. Molecular Biology and Evolution.
41(4):msae068. https://doi.org/10.1093/molbev/msae068.

Beaver RA, Sittichaya W, Liu L-Y. 2014. A Synopsis of the scolytine ambrosia beetles of
Thailand (Coleoptera: Curculionidae: Scolytinae). Zootaxa. 3875(1):1-82.
https://doi.org/10.11646/zootaxa.3875.1.1.

Bickerstaff JRM, Jordal BH, Riegler M. 2023. Two sympatric lineages of Australian *Cnestus*
solidus share *Ambrosiella* symbionts but not *Wolbachia*. Heredity. 132(1):43–53.
https://doi.org/10.1038/s41437-023-00659-w.

Biedermann PHW, et al. 2019. Bark beetle population dynamics in the Anthropocene:
Challenges and solutions. Trends in Ecology & Evolution. 34(1):914–924.
https://doi.org/10.1016/j.tree.2019.06.002.

Bierman A, Roets F, Terblanche JS. 2022. Population structure of the invasive ambrosia
beetle, *Euwallacea fornicatus*, indicates multiple introductions into South Africa. Biol
Invasions. 24(8):2301–2312. https://doi.org/10.1007/s10530-022-02801-x.

Blackmon H, Demuth JP. 2015. Coleoptera karyotype database. The Coleopterists Bulletin.
69(1):174–175. https://doi.org/10.1649/0010-065X-69.1.174.

Blackmon H, Jonika MM, Alfieri JM, Fardoun L, Demuth JP. 2024. Drift drives the
evolution of chromosome number I: The impact of trait transitions on genome evolution in
Coleoptera. Journal of Heredity. 115(2):173–182. https://doi.org/10.1093/jhered/esae001.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25(15):1972–1973.
https://doi.org/10.1093/bioinformatics/btp348.

Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. 2020. BlobToolKit – Interactive
quality assessment of genome assemblies. G3 Genes|Genomes|Genetics. 10(4):1361–1374.
https://doi.org/10.1534/g3.119.400908.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. Nat Methods. 18(2):170–175.
https://doi.org/10.1038/s41592-020-01056-5.

Cook DC, Broughton S. 2023. Economic impact of polyphagous shot hole borer *Euwallacea*
fornicatus (Coleoptera: Curculionidae: Scolytinae) in Western Australia. Agri and Forest
Entomology. 25(3):449–457. https://doi.org/10.1111/afe.12566.

Danecek P, et al. 2021. Twelve years of SAMtools and BCFtools. GigaScience.
10(2):giab008. https://doi.org/10.1093/gigascience/giab008.

De Jager MM, Roets F. 2022. Rapid and cost-effective detection of *Fusarium euwallaceae*
from woody tissues. Plant Pathology. 71(8):1712–1720. https://doi.org/10.1111/ppa.13600.

De Wit MP, et al. 2022. An Assessment of the Potential Economic Impacts of the Invasive
Polyphagous Shot Hole Borer (Coleoptera: Curculionidae) in South Africa. Journal of
Economic Entomology. 115(4):1076–1086. https://doi.org/10.1093/jee/toac061.

Dobin A, et al. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15–
21. https://doi.org/10.1093/bioinformatics/bts635.

Durand NC, et al. 2016. Juicebox provides a visualization system for Hi-C contact maps with
unlimited zoom. Cell Systems. 3(1):99–101. https://doi.org/10.1016/j.cels.2015.07.012.

Eskalen A, et al. 2012. First report of a *Fusarium* sp. and its vector Tea Shot Hole Borer
(*Euwallacea fornicatus*) causing *Fusarium* dieback on avocado in California. Plant Disease.
96(7):1070–1070. https://doi.org/10.1094/PDIS-03-12-0276-PDN.

Feng X, Cheng H, Portik D, Li H. 2022. Metagenome assembly of high-fidelity long reads
with hifiasm-meta. Nat Methods. 19(6):671–674. https://doi.org/10.1038/s41592-022-01478-
3.

Feschotte C, Pritham EJ. 2007. DNA transposons and the evolution of Eukaryotic genomes.
Annu. Rev. Genet. 41(1):331–368. https://doi.org/10.1146/annurev.genet.40.110405.090448.

Freeman S, et al. 2013. *Fusarium euwallaceae* sp. nov.—a symbiotic fungus of *Euwallacea*
sp., an invasive ambrosia beetle in Israel and California. Mycologia. 105(6):1595–1606.
https://doi.org/10.3852/13-066.

Frey JE, et al. 2022. Next generation biosecurity: Towards genome based identification to
prevent spread of agronomic pests and pathogens using nanopore sequencing. PLoS ONE.
17(7):e0270897. https://doi.org/10.1371/journal.pone.0270897.

Gerdol M, et al. 2020. Massive gene presence-absence variation shapes an open pan-genome
in the Mediterranean mussel. Genome Biol. 21(1):275. https://doi.org/10.1186/s13059-020-
02180-3.

Ghurye J, et al. 2019. Integrating Hi-C links with assembly graphs for chromosome-scale
assembly. PLoS Comput Biol. 15(8):e1007273. https://doi.org/10.1371/journal.pcbi.1007273.

Gilbert C, Peccoud J, Cordaux R. 2021. Transposable elements and the evolution of insects.
Annu. Rev. Entomol. 66(1):355–372. https://doi.org/10.1146/annurev-ento-070720-074650.

Gomez DF, et al. 2018. Species delineation within the *Euwallacea fornicatus* (Coleoptera:
Curculionidae) complex revealed by morphometric and phylogenetic analyses. Insect
Systematics and Diversity. 2:(6)2-11. https://doi.org/10.1093/isd/ixy018.

Gomez DF, Lin W, Gao L, Li Y. 2019. New host plant records for the *Euwallacea fornicatus*
(Eichhoff) species complex (Coleoptera: Curculionidae: Scolytinae) across its natural and
introduced distribution. Journal of Asia-Pacific Entomology. 22(1):338–340.
https://doi.org/10.1016/j.aspen.2019.01.013.

Guan D, et al. 2020. Identifying and removing haplotypic duplication in primary genome
assemblies. Bioinformatics. 36(9):2896–2898.
https://doi.org/10.1093/bioinformatics/btaa025.

Guo Q, Huang W, Sang W, Chen X, Wang X. 2023. Characterization, comparative analyses,
and phylogenetic implications of mitochondrial genomes among bark and ambrosia beetles
(Coleoptera: Curculionidae, Scolytinae). Front. Ecol. Evol. 11:1191446.
https://doi.org/10.3389/fevo.2023.1191446.

Hao Z, et al. 2020. *RIdeogram*: drawing SVG graphics to visualize and map genome-wide
data on the idiograms. PeerJ Computer Science. 6:e251. https://doi.org/10.7717/peerj-cs.251.

Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving
the ultrafast bootstrap approximation. Molecular Biology and Evolution. 35(2):518–522.
https://doi.org/10.1093/molbev/msx281.

Hulcr J, Stelinski LL. 2017. The ambrosia symbiosis: From evolutionary ecology to practical
management. Annu. Rev. Entomol. 62(1):285–303. https://doi.org/10.1146/annurev-ento-
031616-035105.

Jolley KA, Bray JE, Maiden MCJ. 2018. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. Wellcome Open Res.
3:124. https://doi.org/10.12688/wellcomeopenres.14826.1.

Jordal B. 2014. 3.7. 12 Scolytinae Latreille, 1806. Coleoptera, Beetles. Volume 3:
Morphology and Systematics (Phytophaga). 633–642. Walter de Gruyter Berlin and New
York.

Jordal BH, Beaver RA, Kirkendall LR. 2001. Breaking taboos in the tropics: incest promotes colonization by wood-boring beetles. Global Ecology and Biogeography. 10(4):345–357. https://doi.org/10.1046/j.1466-822X.2001.00242.x.

Jordal BH, Cognato AI. 2012. Molecular phylogeny of bark and ambrosia beetles reveals multiple origins of fungus farming during periods of global warming. BMC Evol Biol. 12(1):133. https://doi.org/10.1186/1471-2148-12-133.

Kalyaanamoorthy S, Minh BQ, Wong TKF, Von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods. 14(6):587–589. https://doi.org/10.1038/nmeth.4285.

Kapusta. 2017. Parsing-RepeatMasker-Outputs. https://github.com/4ureliek/Parsing-RepeatMasker-Outputs.

Katoh K. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research. 30(14):3059–3066. https://doi.org/10.1093/nar/gkf436.

Kawasaki Y, Schuler H, Stauffer C, Lakatos F, Kajimura H. 2016. *Wolbachia* endosymbionts in haplodiploid and diploid scolytine beetles (Coleoptera: Curculionidae: Scolytinae). Environ Microbiol Rep. 8(5):680–688. https://doi.org/10.1111/1758-2229.12425.

Keeling CI, et al. 2022. Chromosome-level genome assembly reveals genomic architecture of northern range expansion in the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Curculionidae). Molecular Ecology Resources. 22(3):1149–1167. https://doi.org/10.1111/1755-0998.13528.

Lanier GN. 1981. Cytotaxonomy of *Dendroctonus*. Forest, Wildlife, and Range Experiment Station, University of Idaho, Moscow.

Lanier GN, Kirkendall LR. 1986. Karyology of pseudogamous *Ips* bark beetles. Hereditas. 105(1):87–96. https://doi.org/doi: 10.1111/j.1601-5223.1986.tb00646.x.

Lantschner MV, Corley JC, Liebhold AM. 2020. Drivers of global Scolytinae invasion patterns. Ecological Applications. 30(5):e02103. https://doi.org/10.1002/eap.2103.

Lester PJ, et al. 2020. The potential for a CRISPR gene drive to eradicate or suppress globally invasive social wasps. Sci Rep. 10(1):12398. https://doi.org/10.1038/s41598-020-69259-6.

Li K, Xu P, Wang J, Yi X, Jiao Y. 2023. Identification of errors in draft genome assemblies at single-nucleotide resolution for quality assessment and improvement. Nat Commun. 14(1):6556. https://doi.org/10.1038/s41467-023-42336-w.

Liu Z et al. 2022. Chromosome-level genome assembly and population genomic analyses provide insights into adaptive evolution of the red turpentine beetle, *Dendroctonus valens*. BMC Biol. 20(1):190. https://doi.org/10.1186/s12915-022-01388-y.

Lynn KMT, et al. 2021. Novel *Fusarium* mutualists of two *Euwallacea* species infesting *Acacia crassicarpa* in Indonesia. Mycologia. 113(3):536–558. https://doi.org/10.1080/00275514.2021.1875708.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel
and streamlined workflows along with broader and deeper phylogenetic coverage for scoring
of Eukaryotic, Prokaryotic, and Viral genomes. Molecular Biology and Evolution.
38(10):4647–4654. https://doi.org/10.1093/molbev/msab199.

Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of
occurrences of k -mers. Bioinformatics. 27(6):764–770.
https://doi.org/10.1093/bioinformatics/btr011.

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. Nat Rev
Microbiol. 10(1):13–26. https://doi.org/10.1038/nrmicro2670.

McGaughran A, et al. 2024. Genomic tools in biological invasions: Current state and future
frontiers. Genome Biology and Evolution. 16(1):evad230.
https://doi.org/10.1093/gbe/evad230.

Meiborg AB, Faber NR, Taylor BA, Harpur BA, Gorjanc G. 2023. The suppressive potential
of a gene drive in populations of invasive social wasps is currently limited. Sci Rep.
13(1):1640. https://doi.org/10.1038/s41598-023-28867-8.

Mendel Z, et al. 2012. An Asian ambrosia beetle *Euwallacea fornicatus* and its novel
symbiotic fungus *Fusarium* sp. pose a serious threat to the Israeli avocado industry.
Phytoparasitica. 40(3):235–238. https://doi.org/10.1007/s12600-012-0223-7.

Mendel Z, et al. 2021. What determines host range and reproductive performance of an
invasive ambrosia beetle *Euwallacea fornicatus*; Lessons from Israel and California. Front.
For. Glob. Change. 4:654702. https://doi.org/10.3389/ffgc.2021.654702.

Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome
assembly evaluation with QUAST-LG. Bioinformatics. 34(13):i142–i150.
https://doi.org/10.1093/bioinformatics/bty266.

Minh BQ, et al. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. Molecular Biology and Evolution. 37(5):1530–1534.
https://doi.org/10.1093/molbev/msaa015.

Moggioli G, et al. 2023. Distinct genomic routes underlie transitions to specialised symbiotic
lifestyles in deep-sea annelid worms. Nat Commun. 14(1):2814.
https://doi.org/10.1038/s41467-023-38521-6.

Morgulis A, Gertz EM, Schäffer AA, Agarwala R. 2006. WindowMasker: window-based
masker for sequenced genomes. Bioinformatics. 22(2):134–141.
https://doi.org/10.1093/bioinformatics/bti774.

Navarro-Escalante L, et al. 2021. A coffee berry borer (*Hypothenemus hampei*) genome
assembly reveals a reduced chemosensory receptor gene repertoire and male-specific genome
sequences. Sci Rep. 11(1):4900. https://doi.org/10.1038/s41598-021-84068-1.

North HL, McGaughran A, Jiggins CD. 2021. Insights into invasive species from whole-
genome resequencing. Molecular Ecology. 30(23):6289–6308.
https://doi.org/10.1111/mec.15999.

Nurk S, et al. 2020. HiCanu: accurate assembly of segmental duplications, satellites, and
allelic variants from high-fidelity long reads. Genome Res. 30(9):1291–1305.
https://doi.org/10.1101/gr.263566.120.

Nygaard S, et al. 2011. The genome of the leaf-cutting ant *Acromyrmex echinatior* suggests
key adaptations to advanced social life and fungus farming. Genome Res. 21(8):1339–1348.
https://doi.org/10.1101/gr.121392.111.

Nygaard S, et al. 2016. Reciprocal genomic evolution in the ant–fungus agricultural
symbiosis. Nat Commun. 7(1):12233. https://doi.org/10.1038/ncomms12233.

O'Donnell K, et al. 2016. Invasive Asian *Fusarium – Euwallacea* ambrosia beetle mutualists
pose a serious threat to forests, urban landscapes and the avocado industry. Phytoparasitica.
44(4):435–442. https://doi.org/10.1007/s12600-016-0543-0.

Paap T, De Beer ZW, Migliorini D, Nel WJ, Wingfield MJ. 2018. The polyphagous shot hole
borer (PSHB) and its fungal symbiont *Fusarium euwallaceae*: a new invasion in South
Africa. Australasian Plant Pathol. 47(2):231–237. https://doi.org/10.1007/s13313-018-0545-
0.

Peer K, Taborsky M. 2005. Outbreeding depression, but no inbreeding depression in
haplodiploid ambrosia beetles with regular sibling mating. Evolution. 59(2):317–323.
https://doi.org/10.1111/j.0014-3820.2005.tb00992.x.

Petersen M, et al. 2019. Diversity and evolution of the transposable element repertoire in
arthropods with particular reference to insects. BMC Evol Biol. 19(1):11.
https://doi.org/10.1186/s12862-018-1324-9.

Pistone D, Gohli J, Jordal BH. 2018. Molecular phylogeny of bark and ambrosia beetles
(Curculionidae: Scolytinae) based on 18 molecular markers. Systematic Entomology.
43(2):387–406. https://doi.org/10.1111/syen.12281.

Powell D, et al. 2021. A highly-contiguous genome assembly of the Eurasian spruce bark
beetle, *Ips typographus*, provides insight into a major forest pest. Commun Biol. 4(1):1059.
https://doi.org/10.1038/s42003-021-02602-3.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. Nat Commun. 11(1):1432.
https://doi.org/10.1038/s41467-020-14998-3.

Rane R et al. 2023. Complex multiple introductions drive fall armyworm invasions into Asia
and Australia. Sci Rep. 13(1):660. https://doi.org/10.1038/s41598-023-27501-x.

Ren L, Huang W, Cannon EKS, Bertioli DJ, Cannon SB. 2018. A mechanism for genome
size reduction following genomic rearrangements. Front. Genet. 9:454.
https://doi.org/10.3389/fgene.2018.00454.

Ross PA, et al. 2022. A decade of stability for *w*Mel Wolbachia in natural *Aedes aegypti*
populations. PLoS Pathog. 18(2):e1010256. https://doi.org/10.1371/journal.ppat.1010256.

Rugman-Jones PF, et al. 2020. One becomes two: second species of the *Euwallacea*
fornicatus (Coleoptera: Curculionidae: Scolytinae) species complex is established on two
Hawaiian Islands. PeerJ. 8:e9987. https://doi.org/10.7717/peerj.9987.

Ruzzier E, et al. 2023. The first full host plant dataset of Curculionidae Scolytinae of the
world: tribe Xyleborini LeConte, 1876. Sci Data. 10(1):166. https://doi.org/10.1038/s41597-
023-02083-5.

Schrader L, et al. 2021. Relaxed selection underlies genome erosion in socially parasitic ant
species. Nat Commun. 12(1):2918. https://doi.org/10.1038/s41467-021-23178-w.

Schuler H, et al. 2023. Recent invasion and eradication of two members of the *Euwallacea*
fornicatus species complex (Coleoptera: Curculionidae: Scolytinae) from tropical
greenhouses in Europe. Biol Invasions. 25(2):299–307. https://doi.org/10.1007/s10530-022-
02929-w.

Schultz DT, et al. 2023. Ancient gene linkages support ctenophores as sister to other animals.
Nature. 618(7963):110–117. https://doi.org/10.1038/s41586-023-05936-6.

Sillero N, Huey RB, Gilchrist G, Rissler L, Pascual M. 2020. Distribution modelling of an
introduced species: do adaptive genetic markers affect potential range? Proc. R. Soc. B.
287(1935):20201791. https://doi.org/10.1098/rspb.2020.1791.

Sim SB, Corpuz RL, Simmonds TJ, Geib SM. 2022. HiFiAdapterFilt, a memory efficient
read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and
their negative impacts on genome assembly. BMC Genomics. 23(1):157.
https://doi.org/10.1186/s12864-022-08375-1.

Six DL, Bracewell R. 2015. *Dendroctonus*. In: Bark Beetles. Elsevier pp. 305–350.
https://doi.org/10.1016/B978-0-12-417156-5.00008-3.

Sloan DB, Moran NA. 2012. Genome reduction and co-evolution between the primary and
secondary bacterial symbionts of psyllids. Molecular Biology and Evolution. 29(12):3781–
3792. https://doi.org/10.1093/molbev/mss180.

Smith SM, F. Gomez D, A. Beaver R, Hulcr J, I. Cognato A. 2019. Reassessment of the
species in the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) complex after
the rediscovery of the "lost" type specimen. Insects. 10(9):261.
https://doi.org/10.3390/insects10090261.

Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically mapped
cDNA alignments to improve *de novo* gene finding. Bioinformatics. 24:637–644. doi:
10.1093/bioinformatics/btn013.

Stapley J, Santure AW, Dennis SR. 2015. Transposable elements as agents of rapid
adaptation may explain the genetic paradox of invasive species. Molecular Ecology.
24(9):2241–2252. https://doi.org/10.1111/mec.13089.

Steenwyk JL et al. 2021. PhyKIT: a broadly applicable UNIX shell toolkit for processing and
analyzing phylogenomic data. Bioinformatics. 37(16):2325–2331.
https://doi.org/10.1093/bioinformatics/btab096.

Storer CG, Breinholt JW, Hulcr J. 2015. *Wallacellus* is *Euwallacea*: molecular phylogenetics
settles generic relationships (Coleoptera: Curculionidae: Scolytinae: Xyleborini). Zootaxa.
3974(3):391–400. https://doi.org/10.11646/zootaxa.3974.3.6.

Stouthamer R, et al. 2017. Tracing the origin of a cryptic invader: phylogeography of the
Euwallacea fornicatus (Coleoptera: Curculionidae: Scolytinae) species complex. Agri and
Forest Entomology. 19(4):366–375. https://doi.org/10.1111/afe.12215.

Takenouchi Y, Takagi K. 1967. A chromosome study of two parthenogenetic scolytid
beetles. Annotationes zoologicae japonenses. 40(2): 105-110.

Telfer MG, et al. 2024. The genome sequence of the beech bark beetle, *Taphrorychus bicolor*
(Herbst, 1793). Wellcome Open Res. 9:213.
https://doi.org/10.12688/wellcomeopenres.21265.1.

Uliano-Silva M, et al. 2023. MitoHiFi: a python pipeline for mitochondrial genome assembly
from PacBio high fidelity reads. BMC Bioinformatics. 24(1):288.
https://doi.org/10.1186/s12859-023-05385-y.

Vasimuddin Md, Misra S, Li H, Aluru S. 2019. Efficient architecture-aware acceleration of
BWA-MEM for multicore systems. In: 2019 IEEE International Parallel and Distributed
Processing Symposium (IPDPS). pp. 314–324. https://doi.org/10.1109/IPDPS.2019.00041.

Vilardo G, Faccoli M, Corley JC, Lantschner MV. 2022. Factors driving historic
intercontinental invasions of European pine bark beetles. Biol Invasions. 24(9):2973–2991.
https://doi.org/10.1007/s10530-022-02818-2.

Virkki N, Smith SG. 1978. *Animal Cytogenetics*. Schweizerbart Science Publishers: Stuttgart,
Germany

Wang L-J, Hsu M-H, Liu T-Y, Lin M-Y, Sung C-H. 2020. Characterization of the complete
mitochondrial genome of *Euwallacea fornicatus* (Eichhoff, 1868) (Coleoptera:
Curculionidae: Scolytinae) and its phylogenetic implications. Mitochondrial DNA Part B.
5(3):3502–3504. https://doi.org/10.1080/23802359.2020.1827070.

Wang Z, et al. 2023. Genome and transcriptome of *Ips nitidus* provide insights into high-
altitude hypoxia adaptation and symbiosis. iScience. 26(10):107793.
https://doi.org/10.1016/j.isci.2023.107793.

Xi Z, Dean JL, Khoo C, Dobson SL. 2005. Generation of a novel *Wolbachia* infection in
Aedes albopictus (Asian tiger mosquito) via embryonic microinjection. Insect Biochemistry
and Molecular Biology. 35(8):903–910. https://doi.org/10.1016/j.ibmb.2005.03.015.

Zhou C, McCarthy SA, Durbin R. 2023. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics. 39(1):btac808. https://doi.org/10.1093/bioinformatics/btac808.

928