1

Multi-dimensional oscillatory activity of mouse GnRH neurons in vivo

Su Young Han[1], Shel-Hwa Yeo[1], Jae-Chang Kim[2], Ziyue Zhou[1] and Allan E. Herbison[1]

[1]Department of Physiology, Development and Neuroscience, Downing site, University of
Cambridge, Cambridge CB2 3EG, UK and [2]Zurich Center for Neuroeconomics, Department of
Economics, University of Zurich, 8006 Zurich, Switzerland

Corresponding author: Prof. Allan Herbison aeh36@cam.ac.uk

Disclosures: The authors have nothing to disclose and no competing interests.

Abstract

The gonadotropin-releasing hormone (GnRH) neurons represent the key output cells of the
neural network controlling mammalian fertility. We used GCaMP fiber photometry to record the
population activity of the GnRH neuron distal projections in the ventral arcuate nucleus where
they merge before entering the median eminence to release GnRH into the portal vasculature.
Recordings in freely behaving intact male and female mice revealed abrupt 5-min duration
increases in activity that correlated perfectly with the appearance of a subsequent pulse of
luteinizing hormone (LH). In addition, GnRH neuron dendrons exhibited a low level of clustered,
rapidly fluctuating baseline activity in both sexes that did not change through the estrous cycle.
In female mice, a gradual increase in basal activity that exhibited ~80-min oscillations began in
the afternoon of proestrus and lasted for 12 h. This was associated with the onset of the LH
surge that ended several hours before the fall in GCaMP signal. Abrupt episodes of GCaMP
activity continued to occur on top of the rising surge baseline before ceasing in estrus. These
observations provide the first description of GnRH neuron activity in freely behaving animals.
They demonstrate three distinct patterns of oscillatory activity occur in GnRH neurons. These
are comprised of low-level rapid baseline activity, abrupt short-duration oscillations that drive
pulsatile gonadotropin secretion and, in females, a gradual and prolonged oscillating increase in
activity responsible for the relatively short-lived preovulatory LH surge.

Introduction

The founding concepts of reproductive neuroendocrinology arose from the recognition by Harris and colleagues of the neurohumoral basis of the hypophyseal pituitary system (Harris 1955, Fink 2015) and the subsequent discovery by Schally and Guillemin and co-workers of the decapeptide gonadotropin-releasing hormone (GnRH) (Amoss, Burges et al. 1971, Schally, Arimura et al. 1971). This was soon followed by an appreciation that GnRH neurons had a very unusual topography in which cell bodies scattered throughout the basal forebrain sent projections to converge on the median eminence to release GnRH into the portal system (Barry 1979). The quest to understand the patterns of GnRH released into the portal system were largely solved by the portal bleeding approach in sheep (Clarke and Cummins 1982, Caraty and Locatelli 1988). Notably, this revealed that abrupt increments of GnRH secretion evoked pulsatile gonadotropin secretion whereas, in females, prolonged increases in constant or episodic GnRH secretion were responsible the preovulatory luteinizing hormone (LH) surge (Karsch, Bowen et al. 1997, Clarke 2018).

Establishing the activity patterns of GnRH neurons *in vivo* has been hampered considerably by their widely dispersed topography and, five decades on, remains an elusive goal (Herbison 2015, Constantin 2017). The development of transgenic GnRH-GFP mouse lines greatly facilitated efforts to record the electrical properties of individual GnRH neurons in the acute brain slice (Constantin, Moenter et al. 2021). However, the patterns of firing exhibited by GnRH neurons under these conditions have not been found to align well with predicted pulse or surge patterns of GnRH secretion (Herbison 2015, Constantin, Moenter et al. 2021). Similarly, the only report of GnRH neuron firing *in vivo* to date described a range of irregular firing patterns inconsistent with known hormone secretion (Constantin, Iremonger et al. 2013).

The episodic pattern of GnRH secretion driving pulsatile LH release is generated by the intermittent synchronised activity of the arcuate kisspeptin neuron population that projects to and controls the distal processes of GnRH neurons (Herbison 2018, Goodman, Herbison et al. 2022). These processes, termed dendrons in rodents, converge in the ventrolateral aspects of the arcuate nucleus (ARN) before turning into short axons that run into the median eminence (Herbison 2021). The GnRH neuron dendron appears to operate both as an autonomous site for GnRH pulse secretion while also carrying action potentials driving the GnRH surge that are initiated upstream at the level of the GnRH neuron cell bodies (Herbison 2020).

We have previously documented that the expression of GCaMP in GnRH neuron cell bodies and dendrons can be used to faithfully record GnRH neuron electrical activity *in vitro* (Iremonger, Porteous et al. 2017). As the ventrolateral ARN provides a unique location of concentrated GnRH neuron dendrons, we reasoned that it may be possible to record their activity in freely behaving mice at this location using GCaMP fiber photometry. We now demonstrate that this is indeed

73 possible and describe here the patterns of GnRH neuron activity underlying episodic and surge
74 patterns of hormone secretion in male and female mice.
75
76

77 **Results**

78 *Characterization of Gnrh1-GCaMP mouse*
79

The Gnrh1-Cre mouse line (JAX stock #021207) (Yoon, Enquist et al. 2005) was crossed with the
Ai162 (TIT2L-GC6s-ICL-tTA2)-D Cre-dependent GCaMP6s line (JAX stock #031562) (Daigle,
Madisen et al. 2018) to generate Gnrh1-GCaMP6 mice. Dual immunofluorescence in four female
mice demonstrated that 90±2% of GnRH neuron cell bodies expressed GCaMP within the rostral
preoptic area (Fig.1A) as did the majority of GnRH dendrons and fibers within the ME (Fig.1B). In
successfully recorded mice, optic fibers were found to be located immediately above or within
the mid-caudal ARN (Fig.1C). Mice with unsuccessful recordings (n=4), that displayed no
variation in GCaMP signal over 24h recordings, were found to have optic fibers located either
mostly within the third ventricle or >200μM lateral to the ARN.
89

90 *GnRH neuron activity in male mice*
91

The GnRH neuron population activity at the level of distal dendron was measured for 24h
starting between 10.00-11.00am in freely behaving Gnrh1-GCaMP6 male mice (n=7). Mice
exhibited intermittent, abrupt increases in GCaMP signal termed here "dendron-synchronization
episodes" (dSEs; Fig.2A). These episodes had a total duration of 658±5 s comprised of a rapid
increase (full width at half maximum (FWHM) value of 30±4 s) and a slower decline (FWHM
102±7 s) (Fig.2C). The dSEs occurred on average every 93.3±7.4 min but had a large inter-
interval range of 3.9 to 346.6 min and showed no clear modal distribution pattern (Fig.2D). The
24-h recordings also revealed another low amplitude, higher frequency baseline pattern activity
(Fig.2A). Control recordings from mice with misplacement of fibers exhibited almost no baseline
activity (Fig.2B).
102

The characteristics of dSEs strongly resembled those exhibited by arcuate nucleus kisspeptin
neurons that drive pulsatile luteinizing hormone (LH) secretion (Han et al., 2019). Therefore, we
assessed the relationship of GnRH neuron dSEs to LH pulses by taking 5- to 10-minute tail-tip
blood samples for up to 240-min period while monitoring GCaMP fluorescence. This showed a
perfect association between dSE occurrence and LH pulses (Fig.2E). Analysis of the temporal
relationship between dSEs and LH pulses revealed that the peak of dSE consistently preceded LH
pulses by 5.0±0.9 min (N=6 episodes in 5 mice; Fig.2F).
110
111

112
113
GnRH neuron activity in female mice
115

To examine patterns of GnRH neuron activity across the estrous cycle, female Gnrh1-GCaMP6
mice were recorded for 6 h periods between 10am-4pm on each day of the cycle determined by
vaginal cytology on the morning of the recording. Similar to males, freely behaving female mice
exhibited a variable baseline pattern of high-frequency activity upon which intermittent, abrupt
increases in GCaMP signal occurred (Fig.3A). Each dSE showed a total duration of 463±22 s with
a rapid increase (FWHM 31±3 s) and a slower decline (FWHM 81±6 s) (Fig.3B). The frequency of
dSEs varied across the cycle in the same manner as kisspeptin neuron SEs (McQuillan, Han et al.
2019) with a dramatic slowing during estrus (dSE interval of 151.0±35.3 min; P=0.01 vs diestrus
and 0.001 vs proestrus, Dunn's posthoc; n=5) (Fig.3C). The inter-dSE intervals for metestrus,
diestrus and proestrus were not significantly different with mean±SEM of 70.2±6.7 min (n=8),
46.2±4.2 min (n=8), and 38.8±3.4 min (n=6), respectively (Fig.3C). Unlike in males, the overall
inter-dSE interval in females showed a clear right-skewed modal distribution with the inter-dSE
interval of 20 min being the most frequent mode (Fig.3D).
129
130   To examine the relationship of dSEs to LH pulses, tail-tip blood sampling was performed and
131   dSEs found to have a perfect correlation with LH pulses wherein the peak of a dSE occurred
132   6.1±0.2 min (N=6 episodes in 4 mice) before the peak of the following LH pulse (Fig.3E,F).

133
Low amplitude, clustered baseline activity in GnRH neurons in both male and female mice
135

A low-amplitude, high-frequency GCaMP signal was observed in both male and female mice
(Fig.2A, 3A, 4A,B). This activity occurred in a cluster-like manner with mean cluster durations of
22.5±2.8 min (males), 30.0±3.9 min (metestrus), 31.9±6.7 min (diestrus) 24.8±2.2 min
(proestrus) and 35.3±6.8 min (estrus) in female mice. The intra-cluster frequency (range of
0.006-0.018 Hz) was not different between males and females (Fig.4D). Neither the mean
duration nor intra-cluster frequency of the baseline activity were different between males and
females or across the estrous cycle (Fig.4C,D). The amplitude of baseline signal was 10-12%
(males) and 5-12% (females) of the mean amplitude of dSEs.

144
GnRH neuron activity in proestrous female mice
146

To examine patterns of GnRH neuron activity at the time of the preovulatory surge, female
Gnrh1-GCaMP6 mice (n=7) were recorded for a 24-hour period beginning on the morning of
proestrus. Initially, the signals were identical to those observed on metestrus and diestrus with a
low level of irregular baseline activity upon which abrupt, short increases in activity occurred

(Fig.3A,5A). A gradual increase in baseline activity was observed to begin 4.0±0.5 h (range 2-6.5
h) before "lights-off" (19:00h, 12:12 lighting) that peaked 5.9±0.5 h later and declined to
baseline by 12.6±0.8 h (range: 10.3-15.0 h)(Fig.5A). The FWHM period of incline and decline of
this slow activity was 3.1±0.4 and 4.7±0.5 h respectively, confirming the slow dynamics of the
signal. The slow increment in calcium signal was not smooth but consisted of multiple slow
oscillations (Fig.5A). The mean duration of these slow oscillations was 78±4 min (range 32 to 150
min). In addition, dSEs were observed to occur superimposed upon the slow oscillating signal
until around the time of peak baseline increment at which time they stopped for several hours
(Fig5A). Similar duration 24-hour recordings from metestrus-diestrous female mice found only
dSEs without any baseline shifts.

To assess the relationship of the slow oscillating signal to the LH surge, tail-tip bleeding with a 3-
hour interval was undertaken (Fig.5B). This showed that the LH surge began at the time of
baseline GCaMP change and peaked at 9.3±2.8 ng/ml close to when baseline GCaMP level was
at its highest (ΔF/F of 20-40%) (n=4). Notably, GCaMP activity greatly outlasted the duration of
the LH surge (Fig.5B).

To clarify the different components of GCaMP activity occurring on proestrus, we used a
customized Matlab code to separate out the three signals occurring at this time (Fig.6). Firstly, a
30-min rolling average was used to extract the slow oscillatory rise and decline in GCaMP
activity (Fig.6ii). The subtraction of this 'surge signal' from the original signal (Fig.6A) shows the
stereotypical dSEs and baseline fluctuations more clearly (Fig.6iii). Next, the peaks of individual
dSEs were selected and the signals from 60 s before and 360 s after each peak of dSE were
extracted. The remaining signals are considered to be 'the residual signal' comprising the high-
frequency basal activity described above (Fig.6iv).

Discussion

We reveal here three distinct oscillatory patterns of GnRH neuron activity in freely behaving
male and female mice. This includes a baseline of variable high-frequency cluster-like activity
upon which short abrupt episodes occur in both males and females and a prolonged, slowly
oscillating rise in activity on proestrus evening in females. These patterns of GnRH neuron
activity are sufficient to explain the pulse and surge patterns of LH secretion in females and
pulses in males.

We demonstrate that the GCaMP fiber photometry approach used previously for assessing cell
body population activity in this system (Han, Kane et al. 2019, McQuillan, Han et al. 2019) can be
adapted to assess activity in neuronal processes. This has been greatly facilitated by the unusual
topography of the GnRH neurons in which GnRH neuron projections become highly
concentrated in the ventrolateral ARN before passing into the ME. The dendrons in this area

display substantial rises in intracellular calcium in response to propagating action potentials as
well as to the local application of kisspeptin (Iremonger, Porteous et al. 2017, Liu, Yeo et al.
2021). The locations of successful optic fibers immediately above and sometimes lateral to the
ARN makes it very unlikely that recordings were made of GnRH neuron activity at the level of
their terminals within the ME.
194
The GnRH neuron dendrons exhibited clustered high-frequency (approximately 0.01Hz) baseline
activity that did not change in profile across 24h in males or throughout the estrous cycle. This
would be predicted to give rise to fluctuating, low-level inter-pulse GnRH secretion perhaps
compatible with observations made using high-frequency portal sampling in the ewe (Evans,
Dahl et al. 1995). High-frequency dopamine release also occurs within the ME between events
determining prolactin secretion (Romano, Guillou et al. 2017). The physiological significance of
inter-pulse GnRH release at the gonadotroph is unclear (Le Tissier, Campos et al. 2017). For
example, the self-priming action of GnRH at the gonadotroph is well characterized for the surge
but may also exist at other times of the cycle (Fink 1995). Nevertheless, optogenetic
experiments driving GnRH neuron firing at high-frequency burst-like intervals in ovariectomized
mice were not found to generate any appreciable change in LH secretion *in vivo* (Campos and
Herbison 2014).
207
The origin of fluctuating baseline activity in male and female GnRH neuron dendrons is unknown
but it is tempting to speculate that it may arise from episodic electrical activity of the GnRH
neurons themselves. Brain slice electrophysiological studies have consistently documented
repetitive burst firing in GnRH neuron cell bodies (Herbison 2015, Constantin, Moenter et al.
2021) and similar patterns of activity were observed in GnRH neurons in anesthetized mice
(Constantin, Iremonger et al. 2013). However, there appears to be no mechanism for
synchronization between individual GnRH neuron perikarya with dual or multiple recordings of
GnRH neuron cell bodies in brain slices never showing any coordinated activity (Constantin,
Jasoni et al. 2012, Chen and Moenter 2023). Hence, it is also possible that the rapid baseline
activity arises from elsewhere and, for example, the ARN kisspeptin pulse generator is known to
exhibit variable inter-synchronization activity (Han, Morris et al. 2023). This baseline pattern of
GnRH neuron activity may give rise to fluctuating, low-level inter-pulse GnRH secretion. While
not detectable with current LH assays in mice, this is compatible with observations made using
high-frequency portal GnRH sampling in the ewe (Evans, Dahl et al. 1995).
222
We find abrupt 6-10 min-duration increases in GnRH neuron activity that precede each LH pulse.
These events have a dynamic very similar to that of the ARN kisspeptin neuron synchronizations
measured using the same technique (Han, Kane et al. 2019, McQuillan, Han et al. 2019).
Kisspeptin released from terminals in the ventrolateral ARN operates through volume
transmission to rapidly activate and synchronize GnRH neuron dendrons (Iremonger, Porteous
et al. 2017, Liu, Yeo et al. 2021). Alongside evidence that pulsatile LH secretion is abolished in

the absence of kisspeptin signaling (Liu, Yeo et al. 2021), it is almost certain that the abrupt
episodic GnRH neuron activation recorded here arises from up-stream synchronized firing of
ARN kisspeptin neurons. What remains curious, however, is that experimentally, the effects of
kisspeptin on the dendron are typically prolonged lasting for 10s of minutes *in vitro* (Iremonger,
Porteous et al. 2017, Liu, Yeo et al. 2021). Clearly, this is not the case *in vivo* (Fig.6) and it is
possible that the termination of GnRH neuron dendron firing to provide a relatively constrained
secretory signal involves other local mechanisms such as nitric oxide signaling (Constantin,
Reynolds et al. 2021).

Investigators using a variety of pituitary portal GnRH assays in rodent, sheep and primate
models have suggested that GnRH neurons generate the LH surge through either a constant
increase in activity or by increasing their normal pulsatile pattern of release (Sarkar, Chiappa et
al. 1976, Caraty, Locatelli et al. 1989, Moenter, Brand et al. 1992, Xia, Van Vugt et al. 1992,
Clarke 1993, Pau, Berria et al. 1993, Sisk, Richardson et al. 2001). We demonstrate here that,
both predictions were correct with GnRH neuron activity changing in a multi-faceted manner
comprised of a slowly oscillating baseline increase with superimposed episodic activity.
Retrospectively, this is now readily discernable in many of those portal bleeding data (Sarkar,
Chiappa et al. 1976, Caraty, Locatelli et al. 1989, Moenter, Brand et al. 1992, Xia, Van Vugt et al.
1992, Clarke 1993, Pau, Berria et al. 1993, Sisk, Richardson et al. 2001). The slow oscillating
increase in baseline activity observed here is most probably the key event driving the LH surge.
It initiates at the time the LH surge begins and has a duration of approximately 12 hours which is
very similar to that of GnRH release in the portal system of rats, sheep, and monkeys (Sarkar,
Chiappa et al. 1976, Pau, Berria et al. 1993, Karsch, Bowen et al. 1997). The reason for the
extremely prolonged period of GnRH neuron activity and secretion that long outlasts the LH
surge remains unknown but could be related to sexual behavior (Pfaff 1973, Skinner and Caraty
2002). It has been very surprising to find that this gradual increase in activity itself has a slow
hourly oscillatory rhythm. While a circadian contribution to the onset of the surge is well
established (Tonsfeldt, Mellon et al. 2022), it had not been appreciated that an ultradian rhythm
might also exist. It is very likely that the slow increase in GnRH activity recorded here is driven
by the preoptic surge generator (Wang, Guo et al. 2020), and it will be interesting in future
studies to establish whether ultradian rhythms exist within this circuit and what function they
may play in surge generation.

The "two compartment model" of GnRH neurons involves independent pulse and surge
generators that operate on different parts of the GnRH neuron to bring about pulsatile and
surge patterns of gonadotropin secretion (Herbison 2020). We have recently shown that
selective suppression of preoptic area kisspeptin expression abolishes the surge generator but
has no impact on pulsatile LH secretion (Clarkson, Yip et al. 2023). This model predicts that, on
the afternoon of proestrus, the surge generator would operate coincidently with the pulse
generator until such time as post-ovulatory progesterone secretion suppresses the pulse
generator (Herbison 2020). This model is in accord with our present *in vivo* recordings where we

find abrupt episodic pulse activity to continue on top of the rising phase of the baseline increase
until the early hours of estrus when it stops. This highly stereotypical pattern of episodic activity
across the surge is identical to that of the ARN kisspeptin pulse generator (McQuillan, Han et al.
2019), further confirming the origin and nature of these abrupt increases in GnRH neuron
dendron activity.

An important question is whether pulse generator activity at the time of the surge may
contribute to ovulation. While pulse generator events are short in duration, they nevertheless
increase the amplitude of total GnRH neuron activity during the rising phase of the surge and
presumably also GnRH secretion within the portal system. Mice and sheep with a knockdown of
ARN kisspeptin neurons have recently been reported to have reduced LH surge amplitude
(Aerts, Griesgraber et al. 2023, Velasco, Franssen et al. 2023) consistent with the concept
proposed here that the pulse generator contributes to the LH surge directly through activation
of the GnRH neuron dendron. Intriguingly, toxin-induced knockdown of ARN kisspeptin neurons
in rats has the opposite effect of increasing LH surge amplitude (Helena, Toporikova et al. 2015,
Mittelman-Smith, Krajewski-Hall et al. 2016).

In summary, we provide here the first direct recordings of GnRH neuron activity *in vivo*. These
observations demonstrate that alongside a baseline pattern of activity of unknown function,
sharp episodic activity underlies pulse generation whereas a slow and prolonged ultradian
oscillation on proestrus is responsible for the preovulatory gonadotropin surge. The
neurobiological mechanisms underlying this unexpected pattern of surge activity remain to be
discovered.

Materials and Methods
292
Animals
Male and female 129S6Sv/Ev C57BL/6 *Gnrh1*$^{Cre/+}$ mice (Yoon, Enquist et al. 2005) crossed on to
the Ai162 (TIT2L-GC6s-ICL-tTA2)-D Cre-dependent GCaMP6s line (JAX stock #031562) (Daigle,
Madisen et al. 2018) were group-housed in conventional cages with environmental enrichment
under conditions of controlled temperature (22 ± 2^0C) and lighting (12-hour light/12-hour dark
cycle; lights on at 07:00) with *ad libitum* access to food (RM1-P, SDS, UK) and water. All animal
experimental protocols were approved by the University of Cambridge Animal Welfare and
Ethics Review Body under UK Home Office license, P174441DE.
301
Stereotaxic implantation of optic fibers
Adult mice were anaesthetized with 2% isoflurane and placed in a stereotaxic frame under
buprenorphine (0.05mg/kg, s.c.) and meloxicam (5 mg/kg, s.c.) analgesia. Dexamethasone (10
mg/kg, s.c.) was used to prevent cranial swelling. A single 400-μm diameter optic fiber (0.48 NA,
Doric lenses, QC, Canada) was implanted into the brain with the tip placed immediately above

the dorsomedial part of ARN (A-P to bregma, -2.0mm; D-V 5.9mm, M-L ±0.2mm to sagittal
sinus). Following one week of surgery, all animals were handled daily and habituated to a
photometry recording setup for at least three weeks.
310
GCaMP6 fiber photometry and blood sampling
Over a period of 4 to 12 weeks after surgery, fiber photometry experiments were undertaken to
record GCaMP fluorescence signal in freely behaving mice for 6 or 24 h periods using previously
described methodology (Han, Kane et al. 2019, Han, Morris et al. 2023). This included a custom-
built photometry system using Doric components (Doric Lenses, QC, Canada) and National
Instrument data acquisition board (TX, USA) based on a previous design (Lerner, Shilyansky et al.
2015). Blue (465-490 nm) and violet (405 nm) LED lights were sinusoidally modulated at
frequencies of 531 and 211 Hz respectively and were focused onto a single fiber optic connected
to the mouse. The light intensity at the tip of the fiber was 30-80 microwatts. Emitted
fluorescence signal from the brain was collected via the same fiber, passed through a 500-550
nm emission filter and focused onto a fluorescence detector (Doric, QC, Canada). The emissions
were collected at 10 Hz and the two GCaMP6 emissions were recovered by demodulating the
465-490 nm signals (calcium-dependent) and 405 nm (calcium-independent) signals. Signals
were either recorded in a continuous mode or a scheduled 5s on/10s off mode.
Analysis was performed in MATLAB with the subtraction of 405 signal from 465-490 signal to
extract the calcium-dependent signal followed by an exponential fit algorithm used to correct
for baseline shift. The signal was converted to $\Delta F/F$ (%) values using the equation $\Delta F/F=(F_{recorded}-$
$F_{baseline})/ F_{baseline}) \times 100$. The Findpeaks algorithm was used to detect dSEs, and the duration and
the time of SE to the half width full maximum was determined. For deconvolution of signals
from 24-h proestrus recordings, the movmean algorithm was used to extract 30-min rolling
average followed by separating dSE signals (-60 to 360 s around each peak) to visualize the
remaining low amplitude 'residual' signal. A threshold of 5% above baseline was used to extract
the residual signal. Residual signals with peaks occurring > 420s from the preceding peak were
considered to represent separate clusters.
335
To examine the relationship between calcium episodes with LH pulses, freely behaving mice
were attached to the fiber photometry system, and 4-μL blood samples were obtained every 5
to 10 minutes from the tail tip over a period of 120 to 240 minutes. To assess the relationship
between the long calcium increment during proestrus evening and the LH surge, female mice
were attached to the fiber photometry system in the morning of proestrus and blood samples
(3-μL) collected every 3 hours for 18 hours until the morning of estrus. Levels of LH were
measured by in-house LH ELISA (Steyn, Wan et al. 2013) with an assay sensitivity of 0.04 ng/mL
and intra-assay coefficient of variation of 8.2%.
344
Immunohistochemistry

Adult GnRH-Cre,GCaMP6s mice were given a lethal overdose of pentobarbital (3mg/100μL, i.p.)
and perfused transcardially with 4% paraformaldehyde. Brains were processed for dual GFP and
GnRH immunofluorescence. For GFP immunostaining, anti-chicken GFP (1:5000, Aves Lab) was
used followed by AlexaFluor 488-conjugated goat-anti-chicken (1:1000). For GnRH cell body
immunostaining, GA2 guinea pig anti-GnRH antisera (1:3000, gift from G.Anderson, New
Zealand) was used in combination with AlexaFluor 647-conjugated goat anti-guinea pig
immunoglobulin (1:500, Thermo Fisher Scientific, USA). For GnRH dendron immunostaining,
rabbit anti-GnRH (1:20,000, LR1, gift of R.Benoit, Montreal) antisera was used followed by
biotinylated goat anti-rabbit immunoglobulin (1:1000, Jackson Immunoresearch) and AlexaFluor
568-conjugated Streptavidin (1:400, Thermo Fisher Scientific, USA). Imaging was performed
using a Leica SP8 Laser Scanning Confocal Microscope (Leica Microsystems) at the Cambridge
Advanced Imaging Center and analyzed using ImageJ.
358
359 *Statistical Analysis*
360 All statistical analyses were performed in Prism 10 (GraphPad software Inc.). All values given in
361 this study are mean ± SEM, and significance is defined as $P < 0.05^*$ or $P<0.01^{**}$. For inter-peak
362 interval analysis in females across the estrous cycle and the residual cluster activity analysis,
363 Kruskal Wallis ANOVA followed by Dunn's post-hoc tests was used.
364
365

Acknowledgments
This work was supported by the Wellcome Trust (212242/Z/18/Z). JCK received a Doc.Mobility
Fellowship (P1ZHP1_184166) from the Swiss National Science Foundataion. We thank Ms. Maria
Pardo-Navarro for technical assistance. For the purpose of open access, the author has applied a
Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version
arising from this submission.

Author Contributions: SYH, SHY and ZZ undertook investigations; SYH, SHY and JCK
performed analysis; SYH and AEH designed experiments and wrote the manuscript: AEH
generated funding.

Figure Legends

Figure 1. *GCaMP expression in GnRH neurons*. (A) Photomicrographs at the level of organum
vasculosm of the laminae terminalis (OVLT) showing GFP immunofluorescence representing
GCaMP6 (green), GnRH (purple) and merged channels showing double labelled cells (white). (B)
Photomicrographs of four mice at the level of arcuate nucleus showing the location of the end
of the optic fiber (dotted line) in relation to GnRH neuron dendrons labeled with GnRH (red),
GFP representing GCaMP6 (green) and merged channels showing double labelled (yellow). 3V,
third ventricle.

Figure 2. *GnRH neuron dendron activity in male mice*. (A) A representative example of 24-h GCaMP6 photometry recording showing abrupt dendron synchronized episodes (dSEs) occurring on lower amplitude baseline activity in a male mouse. Below, expanded view of the trace indicated by the horizontal line. (B) Example of 8-h GCaMP6 photometry recording in a male mouse with a misplaced fiber optic. All recorded signals are under 3% of baseline. Below, expanded view of the trace indicated by the horizontal line. (C) Average high-resolution profile of an dSE in male mice (N=7 mice) showing a rapid onset followed by a gradual decrease back to baseline. 'a' and 'b' give values at full-width half maximum (FWHM). (D) Inter-peak intervals combined from all recordings (n= 96 dSEs) displayed as a percentage of all intervals occurring in 5-min bins. (E) Representative example showing the relationship of dSEs (black) to pulsatile LH secretion (red). (F) Normalized increase in LH plotted against the dSEs, with the time 0 being the peak of dSE.

Figure 3. *GnRH neuron dendron activity in female mice*. (A) Representative example of 6-h GCaMP6 photometry recording from female mice in (i) metestrus, (ii) diestrus, (iii) proestrus, and (iv) estrus. (B) Average high-resolution profile of dSEs in female mice showing a rapid onset followed by a gradual decrease in the signal. 'a' and 'b' show the values at full-width half maximum (FWHM). (C) Histograms showing inter-peak intervals of dSEs across the estrous cycle. M = metestrus, D = diestrus, P = proestrus E = estrus. *P<0.05, **P<0.01, Kruskal Wallis followed by Dunn's post-hoc test. (D) Inter-peak intervals combined from all recordings (n= 245 dSEs) across metestrus, diestrus, proestrus and estrus displayed as a percentage of all intervals occurring in 5-min bins. (E) A representative example showing the relationship of dSEs (black) to pulsatile LH secretion (red). (F) Normalized increase in LH plotted against the dSEs, with the time 0 being the peak of dSE.

Figure 4. *Baseline high-frequency cluster activity of GnRH neuron dendrons in male and female mice.* Representative 4-h GCaMP recordings in (A) two male mice and (B) a female mouse across the different stages of the estrous cycle showing high frequency baseline activity and larger increases in calcium activity representing dendron synchronization episodes (dSEs) indicated by red asterisks. The peaks of dSEs reaching >30% dFF were cut off to optimize the display of the low amplitude activities. Histograms showing the duration (C) and the intra-cluster frequency (D) of the baseline activity in male and female mice. No significance was found across the groups. M= metestrus, D = diestrus, P = proestrus, E= estrus.

Figure 5. *Slow oscillating increases in calcium activity on the afternoon of proestrus in female mice*. (A) Examples of 21-h GCaMP recordings from four proestrous female mice. Note the prolonged (>10h) gradual oscillating increase in calcium signal beginning on the afternoon of proestrus. The sharp dSEs continue until approximately the plateau phase of the increased baseline at which time they stop or slow. (B) A representative example showing the relationship of the slow increase in baseline calcium activity (black) with LH surge (red). Note that the rise in

LH occurs in alongside the initial rise in calcium activity but returns to baseline several hours
before the calcium signal.

Figure 6. *Deconvolution of the GCaMP signal across the proestrous surge reveals multimodal*
activity patterns of GnRH neuron dendron. (A-B) Representative examples of 24-h photometry
recordings from two female mice starting at proestrus showing (i) the original recording, (ii) 30-
min rolling average highlighting the LH surge signal (red), (iii) the LH surge signal subtracted
from the original recording, displaying the dSEs (blue), and (iv) the residual baseline signal (pink)
after subtracting both the surge and pulse profiles.

References

Aerts, E. G., M. J. Griesgraber, S. L. Shuping, E. C. Bowdridge, S. L. Hardy, R. L. Goodman, C. C.
Nestor and S. M. Hileman (2023). "The effect of NK3-Saporin injection within the arcuate
nucleus on puberty, the LH surge, and the response to Senktide in female sheep." Biol Reprod.
Amoss, M., R. Burges, R. Blackwell, W. Vale, R. Fellows and R. Guillemin (1971). "Purification,
amino acid composition and N-terminus of the hypothalamic luteinizing hormone releasing
hormone factor (LRF) of ovine origin." Biochemical Biophysical Research Communication. **44**:
205-210.
Barry, J. (1979). "Immunohistochemistry of luteinizing hormone-releasing hormone-producing
neurons of the vertebrates." Int Rev Cytol **60**: 179-221.
Campos, P. and A. E. Herbison (2014). "Optogenetic activation of GnRH neurons reveals minimal
requirements for pulsatile luteinizing hormone secretion." Proc Natl Acad Sci U S A **111**(51):
18387-18392.
Caraty, A. and A. Locatelli (1988). "Effect of time after castration on secretion of LHRH and LH in
the ram." J Reprod Fertil **82**(1): 263-269.
Caraty, A., A. Locatelli and G. B. Martin (1989). "Biphasic response in the secretion of
gonadotrophin-releasing hormone in ovariectomized ewes injected with oestradiol." Journal of
Endocrinology **123**: 375-382.
Chen, X. and S. M. Moenter (2023). "Gonadal Feedback Alters the Relationship between Action
Potentials and Hormone Release in Gonadotropin-Releasing Hormone Neurons in Male Mice." J
Neurosci **43**(40): 6717-6730.
Clarke, I. J. (1993). "Variable patterns of gonadotropin-releasing hormone secretion during the
estrogen-induced luteinizing hormone surge in ovariectomized ewes." Endocrinology **133**: 1624-
1632.
Clarke, I. J. (2018). Generation of the GnRH surge and LH surge by the positive feedback effect of
estrogen. The GnRH neuron and its control. A. E. Herbison and T. M. Plant, John Wiley & Sons**:**
325-356.
Clarke, I. J. and J. T. Cummins (1982). "The temporal relationship between gonadotropin
releasing hormone (GnRH) and luteinizing hormone (LH) secretion in ovariectomized ewes."
Endocrinology **111**(5): 1737-1739.

Clarkson, J., S. H. Yip, R. Porteous, A. Kauff, A. K. Heather and A. E. Herbison (2023). "CRISPR-Cas9 knockdown of ESR1 in preoptic GABA-kisspeptin neurons suppresses the preovulatory surge and estrous cycles in female mice." eLife.90959.1

Constantin, S. (2017). "Progress and Challenges in the Search for the Mechanisms of Pulsatile Gonadotropin-Releasing Hormone Secretion." Front Endocrinol (Lausanne) **8**: 180.

Constantin, S., K. J. Iremonger and A. E. Herbison (2013). "In vivo recordings of GnRH neuron firing reveal heterogeneity and dependence upon GABAA receptor signaling." J Neurosci **33**(22): 9394-9401.

Constantin, S., C. Jasoni, N. Romano, K. Lee and A. E. Herbison (2012). "Understanding calcium homeostasis in postnatal gonadotropin-releasing hormone neurons using cell-specific Pericam transgenics." Cell Calcium **51**(3-4): 267-276.

Constantin, S., S. M. Moenter and R. Piet (2021). "The electrophysiologic properties of gonadotropin-releasing hormone neurons." J Neuroendocrinol: e13073.

Constantin, S., D. Reynolds, A. Oh, K. Pizano and S. Wray (2021). "Nitric oxide resets kisspeptin-excited GnRH neurons via PIP2 replenishment." PNAS **118**: e2012339118.

Daigle, T. L., L. Madisen, T. A. Hage, M. T. Valley, U. Knoblich, R. S. Larsen, M. M. Takeno, L. Huang, H. Gu, R. Larsen, M. Mills, A. Bosma-Moody, L. A. Siverts, M. Walker, L. T. Graybuck, Z. Yao, O. Fong, T. N. Nguyen, E. Garren, G. H. Lenz, M. Chavarha, J. Pendergraft, J. Harrington, K. E. Hirokawa, J. A. Harris, P. R. Nicovich, M. J. McGraw, D. R. Ollerenshaw, K. A. Smith, C. A. Baker, J. T. Ting, S. M. Sunkin, J. Lecoq, M. Z. Lin, E. S. Boyden, G. J. Murphy, N. M. da Costa, J. Waters, L. Li, B. Tasic and H. Zeng (2018). "A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality." Cell **174**(2): 465-480 e422.

Evans, N. P., G. E. Dahl, D. Mauger and F. J. Karsch (1995). "Estradiol induces both qualitative and quantitative changes in the pattern of gonadotropin-releasing hormone secretion during the presurge period in the ewe." Endocrinology **136**: 1603-1609.

Fink, G. (1995). "The self-priming effect of LHRH: a unique servomechanism and possible cellular model for memory." Front Neuroendocrinol **16**(2): 183-190.

Fink, G. (2015). "60 YEARS OF NEUROENDOCRINOLOGY: MEMOIR: Harris' neuroendocrine revolution: of portal vessels and self-priming." J Endocrinol **226**(2): T13-24.

Goodman, R. L., A. E. Herbison, M. N. Lehman and V. M. Navarro (2022). "Neuroendocrine control of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion." J Neuroendocrinol **34**(5): e13094.

Han, S. Y., G. Kane, I. Cheong and A. E. Herbison (2019). "Characterization of GnRH Pulse Generator Activity in Male Mice Using GCaMP Fiber Photometry." Endocrinology **160**(3): 557-567.

Han, S. Y., P. G. Morris, J. C. Kim, S. Guru, M. Pardo-Navarro, S. H. Yeo, H. J. McQuillan and A. E. Herbison (2023). "Mechanism of kisspeptin neuron synchronization for pulsatile hormone secretion in male mice." Cell Rep **42**(1): 111914.

Harris, G. W. (1955). Neural control of the pituitary gland. London, Edward Arnold.

Helena, C. V., N. Toporikova, B. Kalil, A. M. Stathopoulos, V. V. Pogrebna, R. O. Carolino, J. A. Anselmo-Franci and R. Bertram (2015). "KNDy Neurons Modulate the Magnitude of the Steroid-Induced Luteinizing Hormone Surges in Ovariectomized Rats." Endocrinology **156**(11): 4200-4213.

Herbison, A. E. (2015). Physiology of the adult GnRH neuronal network. Knobil and Neill's Physiology of Reproduction. T. M. Plant and A. J. Zeleznik, Academic Press. **1:** 399-467.

Herbison, A. E. (2018). "The gonadotropin-releasing hormone pulse generator." Endocrinology
159(11): 3723-3736.

Herbison, A. E. (2020). "A simple model of estrous cycle negative and positive feedback
regulation of GnRH secretion." Front Neuroendocrinol **57**: 100837.

Herbison, A. E. (2021). "The dendron and episodic neuropeptide release." J Neuroendocrinol:
e13024.

Iremonger, K. J., R. Porteous and A. E. Herbison (2017). "Spike and neuropeptide-dependent
mechanisms control GnRH neuron nerve terminal Ca2+ over diverse time scales." J Neurosci
37(12): 3342-3351.

Karsch, F. J., J. M. Bowen, A. Caraty, N. P. Evans and S. M. Moenter (1997). "Gonadotropin-
releasing hormone requirements for ovulation." Biol Reprod **56**: 303-309.

Le Tissier, P., P. Campos, C. Lafont, N. Romano, D. J. Hodson and P. Mollard (2017). "An updated
view of hypothalamic-vascular-pituitary unit function and plasticity." Nat Rev Endocrinol **13**(5):
257-267.

Lerner, T. N., C. Shilyansky, T. J. Davidson, K. E. Evans, K. T. Beier, K. A. Zalocusky, A. K. Crow, R.
C. Malenka, L. Luo, R. Tomer and K. Deisseroth (2015). "Intact-Brain Analyses Reveal Distinct
Information Carried by SNc Dopamine Subcircuits." Cell **162**(3): 635-647.

Liu, X., S. H. Yeo, H. J. McQuillan, M. K. Herde, S. Hessler, I. Cheong, R. Porteous and A. E.
Herbison (2021). "Highly redundant neuropeptide volume co-transmission underlying episodic
activation of the GnRH neuron dendron." Elife **10**.

McQuillan, H. J., S. Y. Han, I. Cheong and A. E. Herbison (2019). "GnRH Pulse Generator Activity
Across the Estrous Cycle of Female Mice." Endocrinology **160**(6): 1480-1491.

Mittelman-Smith, M. A., S. J. Krajewski-Hall, N. T. McMullen and N. E. Rance (2016). "Ablation of
KNDy Neurons Results in Hypogonadotropic Hypogonadism and Amplifies the Steroid-Induced
LH Surge in Female Rats." Endocrinology **157**(5): 2015-2027.

Moenter, S. M., R. C. Brand and F. J. Karsch (1992). "Dynamics of gonadotropin-releasing
hormone (GnRH) secretion during the GnRH surge: insights into the mechanism of GnRH surge
induction." Endocrinology **130**(5): 2978-2984.

Pau, K. F., M. Berria, D. L. Hess and H. G. Spies (1993). "Preovulatory gonadotropin-releasing
hormone surge in ovarian-intact rhesus macaques." Endocrinology **133**: 1650-1656.

Pfaff, D. W. (1973). "Luteinizing hormone-releasing factor potentiates lordosis behavior in
hypophysectomized ovariectomized female rats." Science **182**(117): 1148-1149.

Romano, N., A. Guillou, D. J. Hodson, A. O. Martin and P. Mollard (2017). "Multiple-scale
neuroendocrine signals connect brain and pituitary hormone rhythms." Proc Natl Acad Sci U S A
114(9): 2379-2382.

Sarkar, D. K., S. A. Chiappa, G. Fink and N. M. Sherwood (1976). "Gonadotropin-releasing
hormone surge in pro-oestrous rats." Nature **264**: 461-463.

Schally, A. V., A. Arimura, Y. Baba, R. M. Nair, H. Matsuo, T. W. Redding and L. Debeljuk (1971).
"Isolation and properties of the FSH and LH-releasing hormone." Biochem Biophys Res Commun
43(2): 393-399.

Sisk, C. L., H. N. Richardson, P. E. Chappell and J. E. Levine (2001). "In vivo gonadotropin-
releasing hormone secretion in female rats during peripubertal development and on proestrus."
Endocrinology **142**(7): 2929-2936.

Skinner, D. C. and A. Caraty (2002). "Measurement and possible function of GnRH in
cerebrospinal fluid in ewes." Reprod Suppl **59**: 25-39.

Steyn, F. J., Y. Wan, J. Clarkson, J. D. Veldhuis, A. E. Herbison and C. Chen (2013). "Development
of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and
adult male mice." Endocrinology **154**(12): 4939-4945.
Tonsfeldt, K. J., P. L. Mellon and H. M. Hoffmann (2022). "Circadian Rhythms in the Neuronal
Network Timing the Luteinizing Hormone Surge." Endocrinology **163**(2).
Velasco, I., D. Franssen, S. Daza-Duenas, K. Skrapits, S. Takacs, E. Torres, E. Rodriguez-Vazquez,
M. Ruiz-Cruz, S. Leon, K. Kukoricza, F. P. Zhang, S. Ruohonen, D. Luque-Cordoba, F. Priego-
Capote, F. Gaytan, F. Ruiz-Pino, E. Hrabovszky, M. Poutanen, M. J. Vazquez and M. Tena-
Sempere (2023). "Dissecting the KNDy hypothesis: KNDy neuron-derived kisspeptins are
dispensable for puberty but essential for preserved female fertility and gonadotropin
pulsatility." Metabolism **144**: 155556.
Wang, L., W. Guo, X. Shen, S. Yeo, H. Long, Z. Wang, Q. Lyu, A. E. Herbison and Y. Kuang (2020).
"Different dendritic domains of the GnRH neuron underlie the pulse and surge modes of GnRH
secretion in female mice." Elife **9**.
Xia, L., D. Van Vugt, E. J. Alston, J. Luckhaus and M. Ferin (1992). "A surge of gonadotropin-
releasing hormone accompanies the estradiol-induced gonadotropin surge in the rhesus
monkey." Endocrinology **131**: 2812-2820.
Yoon, H., L. W. Enquist and C. Dulac (2005). "Olfactory inputs to hypothalamic neurons
controlling reproduction and fertility." Cell **123**(4): 669-682.