1   RUNNING TITLE: Fluctuating selection on Monkeyflowers

4   TITLE: Short-term fluctuating and long-term divergent selection on sympatric Monkeyflowers:

5   insights from decade-spanning reciprocal transplants

Caroline M. Dong[1,2], Bolívar Aponte Rolón[1], Juj K. Sullivan[1], Diana Tataru[1], Max Deleon[1],

Rachael Dennis[1], Spencer Dutton[1], Fidel J. Machado Perez[1,3], Lissette Montano[1] & Kathleen G.

Ferris[1]

[1]Tulane University, Department of Ecology and Evolutionary Biology, New Orleans, LA

[2]Grinnell College, Department of Biology, Grinnell, IA

[3]University of California Merced, Life and Environmental Sciences Department, Merced, CA

Corresponding author: cdong1@tulane.edu

KEYWORDS (5-8): adaptation, phenology, speciation, divergence, environmental variation,

evolution, natural selection, temporal variation

SUMMARY:

• Sympatric species often adapt to distinct microhabitats, leading to reproductive isolation
and phenotypic diversity. However, temporal variation in selection may cause local
maladaptation and species boundary breakdown, especially in years with large climatic
events leading to episodic selection. Repeated reciprocal transplants can reveal short and
long-term patterns of natural selection.

• To examine evolutionary trajectories of sympatric Monkeyflowers adapted to different
niches, *Mimulus guttatus* and *M. laciniatus,* we performed three replicated transplants
and combined them with previous experiments to leverage a dataset of five transplants
spanning 10 years. We performed phenotypic selection analyses on parents and hybrids
in parental habitats in Yosemite NP, CA during years of differing snowpack. If there is
ecological isolation, then we predicted local adaptation and divergent phenotypic
selection between habitats in line with species' differences.

• We found interannual fluctuations in selection, often not in predicted directions. Episodic
selection due to extreme high snowpack caused a reversal of local adaptation and
contributed to overall maladaptation of *M. guttatus*. However, a combined-year analysis
detected longer-term divergent selection on flowering time, a key temporally isolating
and adaptative trait, in agreement with species' differences.

• In conclusion, even with annual fluctuations, longer-term divergent selection may still
promote species boundaries.

INTRODUCTION

Closely related sympatric species often occupy distinct habitats within areas of geographic

overlap. Adaptation to differing environments may generate pre- and/or post-zygotic isolation

via selection against immigrants (e.g. habitat isolation) or intermediate hybrids (e.g. extrinsic

post-zygotic isolation; Rundle & Nosil, 2005). Divergent adaptation also increases the chance of

species coexistence and persistence in sympatry (Coyne & Orr, 2004; Rundle & Nosil, 2005;

Schluter, 2009). Given the importance of natural selection in the generation and maintenance of

species, it follows that much of phenotypic diversity is likely due to environmental adaptation

(Schluter, 2000, 2009). As speciation progresses, multiple forms of reproductive isolation

accumulate between divergent lineages hindering measurement of individual barriers (Coyne &

Orr, 2004). Once species are partially isolated, neutral processes also cause the accumulation of

phenotypic divergence. Experimental manipulation is necessary to determine whether habitat

isolation is a key barrier and which traits are involved in adaptive divergence (Ramsey *et al.*,

2003; Hall & Willis, 2006; Lowry *et al.*, 2008). Measuring phenotypic selection on closely

related sympatric species in a reciprocal transplant between native environments could identify

whether divergent adaptation is important for reproductive isolation and speciation (Lowry *et al.*,

2008; Anderson *et al.*, 2015).

Natural populations are subject to spatial and temporal variation in selection (Siepielski *et al.*,

2009) which may influence the strength of reproductive isolation and species boundaries (Tataru

et al., 2023). Additionally, current environmental conditions may be different than the conditions

during original divergence (Coyne & Orr, 2004; Rundle & Nosil, 2005). If environmental

adaptation is important for reproductive isolation and maintenance of species boundaries, then

we expect each species to have higher fitness in its habitat and to detect divergent phenotypic

selection in the direction of species' differences (Anderson *et al.*, 2015), indicating habitat

isolation and local adaptation to current conditions. However, if species are no longer adapted to

their niches due to climate change, then we expect to find local maladaptation (Crespi, 2000) and

selection for the non-local species' phenotype (Wilczek *et al.*, 2014; Kooyers *et al.*, 2019;

Anderson & Wadgymar, 2020). In this scenario, spatially varying selection is acting contrary to

reproductive isolation and could erode species boundaries by driving each towards the other's

phenotypic optimum. Despite divergent spatial selection within one generation, temporal

fluctuations could cause decreased reproductive isolation and eventual breakdown of species

boundaries.

A great deal of temporal fluctuation in the direction and strength of natural selection across taxa

has been found in nature, largely due to climatic factors (Siepielski *et al.*, 2009). Variation in

selection in longitudinal studies is usually measured on a seasonal or annual timescale, a short-

term measurement of population evolution. This leads to the question of how inter-annual

fluctuations affect long-term evolutionary change in species and populations (Wadgymar *et al.*,

2017). A synthesis of phenotypic selection studies by Kingsolver & Diamond (2011) found that

while temporal variation in selection is common, it does not significantly inhibit long-term

directional selection. Therefore, even with fluctuations in the direction of selection, there may be

an overall positive or negative trend over a long enough time period. Wadgymar *et al.* (2017)

found that interannual fluctuations in the direction of selection on the perennial plant *Boechera*

stricta gave rise to long-term patterns consistent with stabilizing selection and local adaptation.

How short versus long-term patterns of variable natural selection compare and impact species'

divergence and reproductive isolation is not currently well understood.

Another understudied agent of evolutionary change is episodic selection resulting from sudden

and extreme environmental events (e.g. drought, heatwave, precipitation events, fire, pest

outbreak). Long-term studies which capture the evolutionary responses to these dramatic

ecological changes are critical for understanding the maintenance of species boundaries. Through

their long-term study of the Galapagos finches, Grant & Grant (1993) were able to illustrate how

an extreme El Niño year altered patterns of selection on beak size and rates of interspecific gene

flow. Strong episodes of selection such as this may drive evolutionary change by affecting the

long-term trajectory of species evolution and biodiversity (Grant & Grant, 1993). Extreme

climatic events are predicted to become more frequent from anthropogenic climate change

(Bailey & van de Pol, 2016).

The *Mimulus guttatus* species complex, a group of closely related wildflowers, is well-suited for

studies of natural selection on species boundaries and reproductive isolation (Wu *et al.*, 2008).

Mimulus guttatus and *M. laciniatus* (syn. *Erythranthe guttata* and *E. laciniata*) are locally

adapted to contrasting microhabitats throughout the central and southern Sierra Nevada, CA and

are incompletely reproductively isolated (Vickery, 1964; Tataru *et al.*, 2024). *Mimulus guttatus*

occurs in moist meadows and seeps while *M. laciniatus* occurs on rocky granite outcrops (Ferris

et al., 2014; Ferris & Willis, 2018). Granite habitats are harsh with shallow soils, high light

intensity, extreme temperatures, and a short growing season (Ferris *et al.*, 2014). *Mimulus*

laciniatus has adapted with a smaller plant size, lobed leaves, self-fertilizing mating system,

reduced floral size, and earlier flowering phenology (DeMarche *et al.*, 2013; Ferris & Willis,

2018). Typical *M. guttatus* populations are largely outcrossing with entire leaves, larger flowers

and plant sizes (Awadalla & Ritland, 1997; Ferris *et al.*, 2014). Flowering time is a key

premating barrier and contributor to local adaptation in the *Mimulus guttatus* species complex

(Hall & Willis, 2006; Lowry *et al.*, 2008; Ferris *et al.*, 2017; Mantel & Sweigart, 2019).

Here, we build on previous work investigating spatial and temporal variation in natural selection

on interspecific divergence between *Mimulus guttatus* and *M. laciniatus* to provide five years of

repeated reciprocal transplant experiments under dramatically varying abiotic conditions (Figure

1). Previously, hybrids and parents were transplanted into each species' habitat in Yosemite

National Park, CA during years of historically low (Ferris & Willis, 2018) and high snowpack

(Tataru et al. 2023). Weaker divergent selection on a key isolating barrier, flowering time, was

found during the high snowpack year which could erode species differentiation. We repeated

these transplants during similarly low snowpack years (2021, 2022) and a year of exceptionally

high snowpack (2023; Figure 1a,b). In this study, we address the following questions: (1) Are *M.*

guttatus and *M. laciniatus* locally adapted? (2) Does the strength and direction of selection on

adaptive and reproductively isolating phenotypes fluctuate? How do short versus long-term

patterns of divergence compare? (3) How does episodic selection affect the trajectory of species

divergence? By examining both interannual variation and overarching signatures of selection

over a 10 year period, we can better elucidate the effects of a changing climate on the

maintenance of biodiversity.

MATERIALS AND METHODS

Repeated reciprocal transplant design

The reciprocal transplants used hybrids created from inbred *M. guttatus* (YVO 18; 37.723366, -

119.746433; 5,395 ft elevation) and *M. laciniatus* lines (WLF 47; 37.841533, -119.59385; 7993

ft elevation; Figure 1c). Due to seed availability, we used a different line from the same *M.*

guttatus population than was used previously (YVO 6; Ferris & Willis, 2018; Tataru *et al.*, 2023).

Parents were crossed to generate F_1 hybrids (WLF 47 maternal x YVO 18 paternal), which were

self-fertilized to generate the F_2 seeds used in 2021. 500 F_2 hybrids were self-fertilized and

propagated with single seed descent to produce the F_3 seeds used in 2022 and to develop the 300

F_6 lines used in 2023. In 2021 and 2022, we pooled seeds from 200 hybrid maternal families for

use in the transplant experiment. To prevent inbreeding depression in parental genotypes in 2023,

we intercrossed inbred lines within each parental population (WLF and YVO).

Our repeated reciprocal transplants took place in 2021, 2022, and 2023 in Yosemite National

Park, CA, USA. We replicated the experiments performed in 2013 and 2019 using the same four

study sites (Figure 1c): two granite outcrops (Granite 1 [37.810700, -119.485200; 8,436 ft

elevation], Granite 2 [37.843702, -119.573120; 8,117 ft elevation]) and two meadows (Meadow

1 [37.767781, -119.772720; 6,605 ft elevation], Meadow 2 [37.755968, -119.803031; 6,163 ft

elevation]. In 2021 and 2022, each site comprised of 50 randomized blocks of 18 and 24

individuals, respectively, in a 4×6 formation with one inch spacing between individuals. In 2021,

blocks contained 6 F_2 hybrids and 6 of each parent and Granite 2 comprised 75 instead of 50

blocks. The 2021 experiment totaled 600 individuals of each genotype (*M. guttatus*, *M.*

laciniatus, F_2 hybrids) in meadow habitat and 750 of each genotype in granite habitat with 4,050

total plants. In 2022, each randomized block contained 16 F_3 hybrids with 4 of each parent,

totaling 1,600 F_3 hybrids and 400 parents per habitat type and 4,800 overall experimental

plants. In 2023, local parental genotypes originating from each site were planted only into their

origin site, to compare the fitness of local *M. guttatus* and *M. laciniatus* parents with the parents

of experimental hybrids originating from nearby sites (WLF and YVO). We used 2–3

intrapopulation parental crosses and original inbred lines, distributed evenly across blocks and

sites, which did not differ in mean seed count and were combined for downstream analyses

(Table S1). Each site had 60 blocks of 35 individuals in a 5×7 formation, composed of 3–4 of

parental genotypes (YVO, WLF, and local), and 16–23 F_6 hybrids for 200 of each parent and

1,500 F_6 hybrids (5 replicates of 300 F_6 lines) at each site. There were a total of 1,200 of each

parent and 3,000 F_6 hybrids in each habitat type with 8,400 overall experimental plants.

We cold-stratified parental *M. laciniatus* and hybrid seeds for 10 days and parental *M. guttatus*

seeds for 5 days at 4°C to break dormancy and synchronize germination (Sunshine Mix #4

Professional Growing Mix, Sun Gro) for one week in greenhouses at the University of California

(UC) Davis (2021) and UC Merced (2022–2023). Following germination, cotyledons were

planted into block designs described above. Mortality within the first week was attributed to

transplant shock and dead germinates were replaced.

Data collection

To gain an understanding of microhabitat variation between and within sites, environmental

measurements were taken weekly from experimental blocks. We measured soil moisture (%)

using the SM150T Soil Moisture Sensor (Dynamax), soil surface temperature (°F) using a laser

thermometer, and light intensity (μmol m-2 s-1) using a MQ-200X Sunlight Quantum Meter

(Apogee Instruments). We recorded survival, herbivory (presence or absence), and phenotypes

every other day until the end of the growing season. At first flower emergence, we recorded the

date and measured plant height (to apical meristem; mm), stigma and anther lengths (mm), and

corolla width (mm). Flowering time was determined as the number of days from transplant to the

first open flower. Herkogamy (i.e. stigma-anther separation) was calculated by subtracting

stigma length from longest anther length, as a metric of outcrossing ability. To assess leaf shape,

the first true leaf was collected one week after flowering. Leaves were digitally scanned and

analyzed to calculate leaf lobing index, a measure of degree of lobing, using ImageJ (Schneider

et al., 2012; Ferris *et al.*, 2015).

Assessing fitness metrics

We collected fruits and counted fruit and seed number, and calculated their correlation using

Kendall's correlation analysis. Plants that produced flowers but did not produce fruit or seed

were indicated with a value of zero. To assess local adaptation, we used three metrics of fitness

for parental genotypes: survival to flowering ('survival'), mean seed number per reproductive

individual ('fecundity'), and mean seed number per planted seed ('total fitness'). To assess long

term patterns of fitness and local adaptation, we averaged survival, fecundity, and total fitness

across years (2013, 2019, 2021, 2022, 2023) for each habitats and species. Data from 2013 was

omitted from averaged fecundity and total fitness calculations because only fruit number was

counted.

Environmental data analysis

To find associations between environmental variables and plant survival in each habitat (granite

or meadow), we used linear mixed effects models with survival as the dependent variable and

soil moisture, time, soil temperature and light levels as fixed independent variables, and block as

a random effect. Survival in this analysis was calculated as the proportion surviving in each

block during each environmental survey. We tested for interactions between independent

variables and performed model selection for the best-fit model using AIC selection criteria using

R packages *nlme* v3.1 (Pinheiro & Bates, 2000; Pinheiro *et al.*, 2023) and *MuMIn* v1.47.5

(Bartoń, 2018). To test whether foliar herbivory and habitat are independent, we used chi-

squared tests to test for correlation and created contingency tables. To understand the effect of

herbivory on plant fitness, we used ANOVA models using seed count as the dependent variable,

and herbivory presence/absence, habitat, site, and block as independent variables and interactions

between them, and performed model selection using the R packages *stats* v3.6.2 (R Core Team,

2023) and *MuMIn* v1.47.5 (Bartoń, 2018).

Phenotypic selection analyses

To investigate plasticity between habitats in parental genotypes, we compared trait means using

t-tests and calculated the coefficient of variation (CV; standard deviation divided by mean;

(Pélabon *et al.*, 2020) using the R package *stats* v3.6.2 (R Core Team, 2023). Although hybrid

composition may differ genetically between habitats, we compared trait means of hybrids

(aggregated) for a putative indication of plasticity. To investigate linkage between traits, we

created a phenotypic trait correlation matrix using hybrid individuals, grouped separately by

habitat and year, using R packages *Hmisc* v5.1-1 (Harrell Jr, 2023) and *corrplot* v0.92 (Simko,

2021). We found generally weak correlations ($r < 0.50$) between traits, except between leaf area

and plant height or flower width in 2022 and 2023 (Figure S2, S3, S4). Therefore, leaf area was

removed from following phenotypic selection analyses.

To understand the strength and direction of selection on traits, we used multivariate linear and

quadratic phenotypic selection analyses on hybrid datasets. Quantitative traits (leaf lobing, plant

height, outcrossing, flower width, flowering time) were standardized to a mean of 0 and s.d. of 1.

We used seed count as the dependent variable with plant height, days to flowering, flower width,

herkogamy, leaf lobing, and their interactions as fixed independent variables, and block nested in

site as a random effect. We used two types of models to account for overdispersion of zeros in

seed count using the R package *glmmTMB* v1.1.8 (Brooks *et al.*, 2017; Magnusson *et al.*, 2017).

Multivariate negative binomial models were used to assess whether a trait is associated with the

production of any seeds. Additionally, we used zero-truncated Poisson analysis to assess whether

traits are associated with the number of seeds produced (i.e. omitting zero values). We performed

model selection for the best-fit model using the R package *MuMIn* v1.47.5 (Bartoń, 2018).

Phenotypes and interactions included in the top model were considered to be under selection. If a

trait did not appear in the model, we consider there to have been no selection on that phenotype.

To extend the comparison to the 2013 experiment which used only fruit number, we repeated all

analyses using fruit number as the dependent variable. Lastly, we repeated phenotypic selection

analyses on a combined dataset of hybrid genotypes from all years (2013–2023) using fruit

number as the dependent variable with plant height, days to flowering, flower width, leaf lobing,

and their interactions as fixed independent variables, and site and year as random effects. To

visualize the fitness-phenotype relationships we plotted selection gradients using the R package

visreg (Breheny & Burchett, 2017).

RESULTS

Soil moisture always influences survival

Fine-scale environmental variation was significantly tied to survival trends in both habitats

across years (Figure 2, Table S2). Soil moisture and time (day) were significantly associated with

survival across all years and habitats ($p < 0.001$). Best-fit models for *M. guttatus*'s meadows in

2021 and 2022 also included light intensity ($p < 0.001$), and 2022 included soil temperature. In

our extreme snowpack year (2023), meadow habitat differed from other years where temperature,

but not light, significantly affected survival ($p < 0.001$; Table S2). In *M. laciniatus's* granite

habitat in 2021 and 2022, time, soil moisture, light and temperature all significantly impacted

survival ($p < 0.0001$; Table S2). In 2023, light and temperature were in the best fit model but

light was not significant ($p = 0.5672$). We found that temperature was significantly associated

with survival in meadow habitat in high, but not low snowpack years. In granite, light

significantly affected survival in low, but not high snowpack years. Across habitats in 2023, light

was not associated with survival. Further, 2021–2023 differed from 2013 and 2019 where soil

moisture was the only significant environmental factor; however, light and temperature were not

measured in 2013.

The extreme high snowpack year (2023) differed from other years in several aspects. Soil

moisture declined gradually in both habitats while light and surface temperatures were lower

(Figure 2). In granite in 2023, there was no characteristic temperature spike usually coinciding

with mortality (Figure 2b). The growing season was delayed by six weeks or more in both

habitats due to the snowpack, with surveys not beginning in granite until previous years' surveys

had ended. Unlike in previous years plants were still alive in meadows at the end of the growing

season: 41.7% (Meadow 1) and 50.7% (Meadow 2) alive at the last survey before snowfall

(Figure 2;). Of the individuals still alive in Meadow 1, many had not flowered (0% of local

genotype, 31.0% of hybrids, 44.1% of *M. guttatus*, 70.7% of *M. laciniatus*). Meadow 2 was

similar (0% of local genotype, 47.3% of hybrids, 40.4% of *M. guttatus*) except most individuals

of *M. laciniatus* flowered (92.0 %). The 2023 growing season was longer in granite (Granite

2013-2022: 33-87 days; 2023: 112-113 days) but comparable to previous years in meadow

(Meadow 2013-2022: 74-160 days; 2023: 119-146 days).

Herbivory is always higher in meadows

As in previous studies (Ferris & Willis, 2018; Tataru *et al.*, 2023) herbivory pressure differed

significantly between habitats in all years with greater herbivory in *M. guttatus'*s meadows :

2021 (meadow 22%, granite 1.97%; χ^2=552.3, d.f. = 1, p < 0.0001), 2022 (meadow 37.75%,

granite 2.58%; χ^2=921.78, d.f. = 1, p < 0.0001), and 2023 (meadow 23.17%, granite 8.74%;

χ^2=325.91, d.f. = 1, p < 0.0001). Only in 2021 did the best-fit ANOVA model of the effect of

herbivory on total fitness include herbivory, site, and their (Table S3). Overall, herbivory

pressure was spatially stable but its effect on fitness varied temporally.

M. guttatus appears maladapted to its meadows

Mimulus laciniatus had higher survival to flowering than *M. guttatus* in both habitats in all three

years, similar to 2013 (Figure 3a). In 2019 when there was extremely low overall survival of

parental genotypes. With all five years combined, *M. laciniatus* had higher average survival in

both habitats (Figure 3a). Patterns of fecundity differed between years with parental species

having higher fecundity in their native habitats in 2021 indicating local adaptation and selection

against immigrants (Figure 3b). However, in 2022, *M. laciniatus* had higher fecundity in both

habitats, and in 2023, we found reciprocal local maladaptation where *M. guttatus* had higher

fecundity than *M. laciniatus* in granite and vice versa in meadows. With years combined, we

found overall evidence of local maladaptation in mean fecundity (Figure 3b); however, if the

extreme snowpack of 2023 is excluded then local adaptation in fecundity is seen (Figure S1). For

total fitness, *M. laciniatus* and *M. gutta*tus were equal in granite in high snowpack years (2019

and 2023), but *M. laciniatus* had the home-site advantage in low snowpack years (2013, 2021,

2022). Total fitness in meadow was not as predictable with *M. laciniatus* having higher total

fitness for all three recent years but lower or similar fitness in 2013 and 2019. Overall, *M.*

laciniatus had higher total fitness in both habitats (Figure 3c, S1).

Phenotypes are adaptively plastic between habitats

Parental genotypes exhibited plasticity in phenotypic traits between habitats, with the non-native

species exhibited phenotypic plasticity in the direction of the local species' phenotype. In 2021 in

its nonnative granite habitat, *M. guttatus* had significantly shorter mean flowering time and

smaller plant height, more similar to the native *M. laciniatus* trait values (Table 2). In 2022 in

granite habitat, *M. guttatus* had significantly smaller leaf area and increased leaf lobing which

are both *M. laciniatus*-like traits (Table 3). In 2023, phenotypic plasticity moved *M. guttatus* to

more *M. laciniatus*-like values in all traits with shorter flowering time, smaller plant height,

smaller flowers, smaller leaves with increased lobing in granite habitat (Table 4). In all three

years, *M. laciniatus* grew more *M. guttatus*-like when growing in its non-native meadow habitat

with significantly longer flowering time, taller plant height, larger leaf area (Tables 3, 4, 5).

Additionally in 2022 it had reduced leaf lobing like *M. guttatus*, but not in 2021 and 2023 (Table

3). These results are similar to 2013 with plasticity in every trait except *M. laciniatus*'s flower

size, in the direction of the local species' phenotype (Ferris & Willis 2018).

Similarly, mean hybrid phenotypes were different between habitats with shorter flowering time,

smaller plant height, smaller leaf area in hybrids grown in the granite habitat. In 2021, hybrids in

granite exhibited reduced herkogamy, in the direction of the *M. laciniatus*, but reduced leaf

lobing unlike *M. laciniatus* (Table 2). For hybrids in 2022, we found reduced herkogamy,

smaller flower width and increased leaf lobing in granite, all in the *M. laciniatus* direction (Table

3). In 2023, we found that hybrids had smaller flowers but increased herkogamy in granite,

(Table 4). In 2019, hybrids in granite differed by having larger flowers and larger leaf area

relative to meadow (Tataru et al. 2023). In contrast, there were no significant differences

between hybrids in granite and meadow habitats in 2013 (Ferris & Willis 2018). Overall, an

earlier mean flowering time in granite is common throughout years (2013–2023) across hybrids

and parents.

Interannual fluctuation but long-term directional selection

We found temporal fluctuation in the strength and direction of selection on each of our

quantitative traits. Based on observed trait differences between species, we predicted selection in

meadow habitat for later flowering time, increased flower width, decreased leaf lobing, increased

plant height, and increased herkogamy, with the inverse direction of selection in granite habitat

(Figure 4a). In our comparison of low snowpack years, we found less divergent selection

between habitats in the direction of species differences in 2021 relative to 2013. We were not

able to assess selection between habitats in 2022 due to insufficient seed production in granite.

We generally found weaker selection in *M. guttatus*'s meadow habitat in 2022 relative to 2021,

contrary to expectations as 2022 was slightly drier. For certain traits, we found a very strong

signature of selection in 2023 relative to other years (e.g. flower width, leaf lobing, plant height).

In the negative binomial selection analysis, measuring whether a trait influenced the production

of any versus zero seeds, we found consistent selection for earlier flowering in both habitats

across years, except in 2022 where flowering time was absent in the best-fit model (Figure 4b;

Table 5, S4). In all years except 2021, there was stronger selection for earlier flowering in

granite, matching predictions. We found similar selection for smaller flowers in 2021 in both

granite and meadow habitats, but strong selection for larger flowers in 2022 and 2023 which

echoed results in 2013 (granite) and 2019 (meadow). There was selection for reduced leaf lobing

in meadows across years, matching predictions. In granite habitat, there was selection for

increased leaf lobing in 2013, matching predictions, however leaf shape did not impact fitness in

other years. As predicted we found selection for taller plants in every year in meadows. There

was selection for taller plants in granite in 2021 and 2023 which was unexpected, but selection

for smaller plants in 2013. There was selection for increased herkogamy as predicted from the

outcrossing mating system of *M. guttatus* in 2023 in meadow habitat.

In the zero-truncated Poisson selection analysis, which measures the influence of a trait on seed

number excluding plants with zero seeds, we found divergent selection in the opposite direction

of predictions in many years. In contrast to our predictions and 2013 results, we found stronger

selection for earlier flowering in meadow habitats than granite in 2021, 2019, and 2023 (Figure

4c; Table 6, S4). In the high snowpack years (2019 and 2023), selection was actually reversed

from our predictions with later flowering increasing fitness in granite and earlier flowering

advantageous in meadows. Across all years and habitats, we found that having larger flowers and

taller plant height resulted in higher fitness, except for weak selection for smaller flowers in 2023

in meadows and strong selection for shorter plants in granite in 2023. Smaller plants were never

selected for in meadows. There was selection against leaf lobing in both habitats in 2021 and

strong selection against lobing in granite in 2023, and this is different from our predictions

(Figure 4a) and previous results. We found divergent selection on herkogamy in 2021 and 2023

with stronger selection in granite which matches the native *M. lacinaitus's* self-fertilizing mating

system. Also in line with our predictions, there was selection for reduced herkogamy in 2022 and

2023 in meadow. We found that β values derived from fruit number were similar to those from

seed number in both strength and direction (Table S5 and S6). This is likely attributed to a strong

positive correlation between seed and fruit number (Table S7).

Although we see interannual fluctuations in the strength and direction of selection on all traits,

when we combined data across years we found long term patterns of directional selection (Table

7, Figure 5, S5). In both zero-truncated Poisson and negative binomial analyses, we found long

term selection for earlier flowering time and larger plant sizes in both habitats, with stronger

selection in granite. Further, we found selection for smaller flowers in both habitats which does

not match our predictions. However, we found selection for more lobed leaves in granite and

negligible selection on leaf shape in meadow habitat. Herkogamy was not included in the

combined dataset due to its absence in 2013. Therefore, while annual fluctuations make selection

on our traits seem inconsistent, we do find longer term divergent selection in the direction of

species' differences in both flowering time and leaf shape.

DISCUSSION

In this study, we investigated spatial and temporal variation in selection on two sympatric

Monkeyflower species, *M. guttatus* and *M. laciniatus*. We compared multiple large-scale

replicated reciprocal transplants conducted in low (2013, 2021, 2022) and high (2019, 2023)

snowpack years including one extreme year resulting in episodic selection (Figure 1a,b). Using

experimental hybrids, we found fluctuating selection on phenotypic traits involved in

reproductive isolation and environmental adaptation. Our results suggest that *M. guttatus* is

maladapted to its meadow habitat across years and that habitat isolation may therefore be

asymmetric. Within each year, phenotypic selection was often contrary to our predicted

directions but we saw a longer-term trend of divergent selection in the direction of species'

differences in both flowering time, a key reproductive isolating trait, and leaf shape. Finally, we

find that episodic selection, caused by an extreme snowfall event, may disrupt local adaptation

and impact species boundaries.

M. guttatus is locally maladapted

Organisms are expected to have a fitness advantage in their home environment because spatial

heterogeneity should favor the evolution of local adaptation (Hastings, 1983). However,

temporal changes in environmental conditions are theorized to constrain adaptation if there are

opposing selection pressures over time (Stearns, 1992) which may negatively impact

reproductive isolation and species boundaries. We predicted that if habitat isolation is an

important reproductive isolating barrier between our sympatric *Mimulus*, then there would be a

fitness advantage of the local species in each habitat. Instead, we found evidence of local

maladaptation of *M. guttatus* with *M. laciniatus* having higher total fitness in both habitats across

time (Figure 3c). Annual fluctuations in snowpack did influence the relative fitness of the parent

species in granite with equal fitness of *M. laciniatus* and *M. gutta*tus in high snowpack years, but

in low years *M. laciniatus* had the home-site advantage (Figure 3c).

Although local adaptation is thought to be ubiquitous in natural populations (Hereford, 2009),

fitness trade-offs are not always found in reciprocal transplants (Bennett & Lenski, 2007; Lowry

et al., 2009). It is possible that our reciprocal transplant did not detect existing trade-offs or that

adaptation has evolved in *M. laciniatus* without fitness trade-offs across environments (Fry, 1996;

Leimu & Fischer, 2008). A failure to detect local adaptation of *M. guttatus* to its native habitat

could be attributed to germinating seeds in greenhouse conditions and transplanting as

cotyledons, and/or the reduction of competition in experimental blocks. Transplanting as

cotyledons may not account for species differences in seed survival, dormancy, and germination.

Parental genotypes may have been negatively impacted by inbreeding, affecting *M. guttatus*

more than the highly self-fertilizing *M. laciniatus* (93–97%; Ferris *et al.*, 2014). However, this

seems unlikely because outbred and inbred parental genotypes performed similarly in 2023

(Table S1).

Asymmetry in local adaptation or a lack of trade-offs, has been found in other transplant

experiments (Hereford, 2009; Gosden *et al.*, 2015; Latreille & Pichot, 2017; Toll & Willis, 2018).

The universal fitness advantage of *M. laciniatus* could be due to adaptive phenotypic plasticity

(Ghalambor *et al.*, 2015). We found that both *M. guttatus* and *M. laciniatus* exhibit plasticity in

the direction of the local species' phenotype in each habitat (Tables 2, 3, & 4). *Mimulus*

laciniatus's occupation of a harsh, highly variable environment may have driven the evolution of

adaptive plasticity (Via & Lande, 1985) allowing it to thrive in both habitats. However, *M.*

guttatus was more plastic than *M. laciniatus* in all phenotypes except fitness indicating that

plasticity, in the traits we measured, is unlikely to account for the lack of fitness trade-offs.

Alternatively, *M. guttatus* may no longer be at its adaptive optima. Local adaptation is predicted

to evolve in an annual outcrossing plant with large effective population sizes like *M. guttatus*,

but may be constrained by natural selection, gene flow, genetic drift, low genetic diversity, or the

genetic architecture of underlying traits (Kawecki & Ebert, 2004; Anderson *et al.*, 2011). Our

results could indicate an adaptive lag where, due to recent rapid climate change, the adaptive

landscape of *M. guttatus's* habitat has changed to be more *M. laciniatus*-like and *M. guttatus*'s

evolutionary response has not kept pace (Lane *et al.*, 2012; Mills *et al.*, 2013; Kooyers *et al.*,

2019). This pattern of adaptive lag and local maladaptation has been found in other populations

of *M. guttatus* (Kooyers *et al.*, 2019). Over time this trend of adaptive lag could erode species

boundaries by allowing colonization of *M. guttatus's* meadow by *M. laciniatus* potentially

increasing hybridization or worse, lead to population extinction.

Annual fluctuation in natural selection does not erase signatures of long-term divergence

Temporal and spatial variation in the strength and direction of selection has been well

characterized as a mechanism that maintains genetic variation within species (Hereford, 2009;

Siepielski *et al.*, 2009, 2017; Bergland et al. 2014; Bitter et al. 2023). However, few studies have

empirically tested how fluctuating selection impacts the strength of reproductive isolation

between sympatric species (but see Campbell & Powers, 2015). We uncovered annually

fluctuating selection on traits between low and high snowpack years, with more consistent

patterns of selection on plant height, leaf lobing, flowering time and plant height. There was

selection for earlier flowering time in granite across most years (Figure 4b,c). However, in high

snowpack years, later flowering was associated with higher seed number in *M. laciniatus's*

granite habitat which is in the opposite direction of species' differences (Figure 4c). This reflects

a relaxation of divergent selection in high snowpack years. *Mimulus guttatus* that flower later in

the season tend to produce higher seed (Mojica & Kelly, 2010; Mojica *et al.*, 2012), particularly

during a longer, high snowpack growing season with lowered desiccation risk. Selection on leaf

lobing also fluctuated across years but typically with selection patterns opposite predictions

(Figure 4c).

Annual selection fluctuated unpredictably for flower width with combined selection for smaller

flowers (Figure 4, Table 7), counter to species differences where *M. guttatus* has larger flowers

than *M. laciniatus*. Flower size affects reproductive success and can contribute to population

differentiation and reproductive isolation (Bradshaw *et al.*, 1995; Brunet, 2009; Schiestl &

Schluter, 2009; Venail *et al.*, 2010; Krizek & Anderson, 2013). Flower size is often correlated

with plant size indicating that early flowering, small stature, and small flowers are linked (Troth

et al., 2018). An antagonistic pleiotropy has been demonstrated in *M. guttatus* where later

flowering alleles also increase plant and flower size (Mojica *et al.*, 2012; Monnahan & Kelly,

2015). Although plants with larger flowers have increased fecundity, they also lower viability,

resulting in overall selection for smaller flowers (Mojica & Kelly, 2010). Although we saw

selection for early flowering and small flowers, we find that selection consistently favored larger

plants in both habitats across years suggesting that in our species' these traits are genetically

distinct (Table 7). Although we initially predicted selection for smaller plant sizes in granite

based on *M. laciniatus's* short stature, our findings suggest that larger plants always have higher

fecundity. Size at reproduction is a key determinant of fecundity across taxa (Aarseen & Taylor,

1992; Aarssen & Jordan, 2001; Kingsolver & Pfennig, 2004; Troth *et al.*, 2018). Despite

selection for larger sizes, average plant height was still smaller in granite than meadow habitats

(Tables 2, 3, & 4), likely constrained by limited water and nutrients in rocky outcrops. We found

weak selection on herkogamy perhaps indicating that selection on stigma-anther distance is no

longer as strong as during the earlier stages of speciation (Harder & Johnson, 2009).

Even with interannual fluctuations, we found longer-term divergent selection on flowering time

and leaf shape between habitats when we combined data across years (Table 7, Figures 5, S5). In

the combined analysis, there was stronger selection for early flowering in granite than meadow

habitat which aligns with our predictions based on species' differences. We also found a distinct

signature of divergent selection on leaf lobing in predicted directions. Lobed leaf shape is

thought to be involved in adaptation to dry, exposed habitats like *M. laciniatus's* rocky outcrop

environment (Nicotra *et al.*, 2011; Ferris, 2019) and therefore may also be involved in habitat

adaptation and isolation (Ferris & Willis, 2018; Tataru *et al.*, 2023). A signature of overall

divergent selection on these key traits in across transplants suggests ongoing adaptive

differentiation and the potential for long-term maintenance of species boundaries (Chapurlat *et*

al., 2020).

Episodic selection weakens species divergence

The high Sierra Nevada snowpack of 2023 resulted in the delayed onset of spring in mid to high

elevations by approximately 4–6 weeks and dramatically impacted patterns of soil moisture, light

intensity, and surface temperature (Figure 2). In shifting the growing season the extreme

snowpack contributed to local maladaptation between parental species (Figure 3b), as evidenced

by a signature of local adaptation in fecundity reappearing when 2023 was excluded (Figure S1).

Our results indicate that in *M. guttatus's* meadow habitat, this delay was detrimental to hybrids

and *M. guttatus* by delaying flowering phenology which negatively impacted fitness.

Additionally, none of the experimental local meadow genotypes flowered before snowfall,

indicating a negative effect on native populations. However, many meadow plants remained

alive at the first snowfall indicating potential facultative perenniality. Shorter day lengths were

experienced in the late shifted season and critical photoperiod thresholds may not have been met

to initiate the transition flowering (Friedman & Willis, 2013; Fishman *et al.*, 2014; Kenney &

Sweigart, 2016). End-of-season drought stress may also typically act as a flowering cue in these

populations (Kooyers *et al.*, 2015; Mantel & Sweigart, 2019), however in 2023 soil moisture in

the meadows remained higher than in previous years (Figure 2). The seasonal delay did not

appear to negatively affect the fitness of *M. laciniatus,* likely because *M. laciniatus* flowers

rapidly under shorter daylengths (Friedman & Willis, 2013; Ferris & Willis, 2018).

In granite, the delayed season in 2023 fostered a more favorable environment for the non-native

M. guttatus with high water availability, lower light intensity and surface temperatures, and a

longer growing season than is typical in the harsh habitat (Figure 2). This led to higher *M.*

guttatus survival in granite than usual (Table 1). Although *M. guttatus* still had lower survival

than *M. laciniatus*, it had remarkably high fecundity (Figure 3b) resulting in similar total fitness

between the two species in granite which was unprecedented in previous years (Figure 3c). In

contrast to the meadow habitat, all plants had senesced and many had set seed by the first

snowfall in granite habitat in 2023. This suggests that drought stress is more important than

critical photoperiod as a flowering cue in *M. lacinaitus's* habitat. It has also been shown that

high-elevation populations of *M. laciniatus* have increased plasticity in critical photoperiod

(Love & Ferris, 2024).

Our results demonstrate how an extreme climatic event may disrupt species boundaries. This

atypical year produced unusually strong selection opposite of predicted directions on key life-

history and morphology traits involved in adaptation and reproductive isolation: flowering time,

flower size and leaf lobing (Figure 4). In the long term, episodic selection could expand or

contract *M. guttatus* or *M. laciniatus'* ranges, encourage the emergence of hybrid swarms or a

new hybrid species (Grant & Grant, 1993), or shift their adaptive potential (Campbell-Staton *et*

al., 2017). The rarity and unpredictability of extreme climatic events complicates predictions

about the evolutionary response to climate change. Surveys of plant populations after extreme

climatic events can monitor changes in the selective landscape and responses (Anderson, 2016;

Bailey & van de Pol, 2016). In species with limited ranges, like the Sierra Nevada endemic *M.*

laciniatus, will these events influence extinction risk? More longitudinal field studies are needed

to understand long-term evolutionary consequences of extreme climatic events and changing

climatic patterns on native plant species. For instance, do these events drive evolutionary change

and does this outweigh selection acting during 'normal' periods? For species with a limited range,

such as the Sierra Nevada endemic *M. laciniatus*, will these events influence extinction risk of

populations?

Conclusions

Our decade-spanning repeated reciprocal transplants provide greater insight into the impacts of

fluctuating selection on species' boundaries. The foreign advantage of *M. laciniatus* in the non-

native meadow habitat across years suggests either a low cost of adaptation to the granite habitat

or recent change in the shape of the adaptive landscape of the meadow habitat leading to current

maladaptation of *M. guttatus*. Furthermore, longer-term divergent selection on flowering time

found in our combined phenotypic selection analysis should reinforce temporal and habitat

isolation between these species. However, the lack of consistent divergent selection on other

traits involved in reproductive isolation and local adaptation could lead to eventual species

fusion. Finally, a bout of episodic natural selection due to extreme snowfall shifted population

dynamics in a way that may have long term impacts on species' range, abundance, and gene flow.

Our study illustrates the strength of integrating manipulative field experiments across multiple

spatial and temporal scales to gain a deeper understanding of the maintenance of biodiversity in

the face of environmental change (Wadgymar *et al.*, 2017; Oakley *et al.*, 2023).

ACKNOWLEDGEMENTS

This work was financially supported by the National Institute of General Medicinal Sciences of

the National Institute of Health (NIH) under award number R35GM138224 to KGF. We thank

the Yosemite National Park Service for permit support (YOSE-2021-SCI-0033, YOSE-2022-

SCI-0051, YOSE-2023-SCI-0011), and Breeanne Jackson at Yosemite Field Station

(doi:10.21973/n3v36c) for logistical support and providing accommodation. We thank Johanna

Schmitt at the University of California (UC) Davis, and Danielle Edwards and Jason Sexton at

UC Merced for providing cold-stratification and greenhouse resources. We also thank members

of the Ferris Lab who assisted the project, Natalie Gonzalez and Jessica Scales. Lastly, we

acknowledge the original inhabitants of the unceded land on which our research was conducted,

the Southern Sierra Miwuk Nation, Bishop Paiute Tribe, Bridgeport Indian Colony, Mono Lake

Kutzadikaa, North Fork Rancheria of Mono Indians of California, Picayune Rancheria of the

Chukchansi Indians and the Tuolumne Band of Me-Wuk Indians.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

KGF conceived the study and CMD, DT, BAR, and JKS contributed to experimental design.

CMD, BAR, JKS, MD, RD, SD, FMP, and LM performed the reciprocal transplant experiments.

CMD completed data analysis and drafted the manuscript with input from KGF. All authors

contributed to and approved the final version of the manuscript.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study will be openly available in [repository name] at

http://doi.org/[doi], reference number [reference number], upon acceptance of the manuscript.

REFERENCES

Aarseen LW, Taylor DR. **1992**. Fecundity allocation in herbaceous plants. *Oikos* **65**: 225–232.

Aarssen LW, Jordan CY. **2001**. Between-species patterns of covariation in plant size, seed size,

and fecundity in monocarpic herbs. *Ecoscience* **8**: 471–477.

Anderson JT. **2016**. Plant fitness in a rapidly changing world. *New Phytologist* **210**: 81–87.

Anderson JT, Eckhart VM, Geber MA. **2015**. Experimental studies of adaptation in *Clarkia*

xantiana: III: Phenotypic selection across a subspecies border. *Evolution* **69**: 2249–2261.

Anderson JT, Wadgymar SM. **2020**. Climate change disrupts local adaptation and favours

upslope migration. *Ecology Letters* **23**: 181–192.

Anderson JT, Willis JH, Mitchell-Olds T. **2011**. Evolutionary genetics of plant adaptation.

Trends in Genetics **27**: 258–266.

Awadalla P, Ritland K. **1997**. Microsatellite variation and evolution in the *Mimulus guttatus*

species complex with contrasting mating systems. *Molecular Biology and Evolution* **14**: 1023–

1034.

Bailey LD, van de Pol M. **2016**. Tackling extremes: Challenges for ecological and evolutionary

research on extreme climatic events. *Journal of Animal Ecology* **85**: 85–96.

Bartoń K. **2018**. MuMIn: Multi-Model Inference. R package version 1.42.1. https://CRAN.R-

project.org/package=MuMIn.

Bennett AF, Lenski RE. **2007**. An experimental test of evolutionary trade-offs during

temperature adaptation. *Proceedings of the National Academy of Sciences of the United States of*

America **104**: 8649–8654.

Bradshaw HD, Wilbert SM, Otto KG, Schemske DW. **1995**. Genetic mapping of floral traits

associated with reproductive isolation in monkeyflowers (*Mimulus*). *Nature* **376**: 762–765.

Breheny P, Burchett W. **2017**. Visualization of Regression Models Using visreg. *The R Journal*

9: 56–71.

Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, Skaug HJ,

Maechler M, Bolker BM. **2017**. Modeling Zero-Inflated Count Data With glmmTMB. bioRxiv

preprint. *bioRxiv*.

Brunet J. **2009**. Pollinators of the Rocky Mountain columbine: Temporal variation, functional

groups and associations with floral traits. *Annals of Botany* **103**: 1567–1578.

Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards S V. **2017**.

Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard.

Science **357**: 495–498.

Campbell DR, Powers JM. **2015**. Natural selection on floral morphology can be influenced by

climate. *Proceedings of the Royal Society B: Biological Sciences* **282**.

Chapurlat E, Le Roncé I, Ågren J, Sletvold N. **2020**. Divergent selection on flowering

phenology but not on floral morphology between two closely related orchids. *Ecology and*

Evolution **10**: 5737–5747.

Coyne JA, Orr HA. **2004**. *Speciation*. Sunderland, Massachusetts: Sinauer Associates.

Crespi BJ. **2000**. The evolution of maladaptation. *Heredity* **84**: 623–629.

DeMarche ML, Rice KJ, Sexton JP. **2013**. Niche partitioning between close relatives suggests

trade-offs between adaptation to local environments and competition. *Ecology and Evolution* **3**:

512–522.

Ferris KG. **2019**. Endless forms most functional: uncovering the role of natural selection in the

evolution of leaf shape. *American Journal of Botany* **106**: 1532–1535.

Ferris KG, Barnett LL, Blackman BK, Willis JH. **2017**. The genetic architecture of local

adaptation and reproductive isolation in sympatry within the *Mimulus guttatus* species complex.

Molecular Ecology **26**: 208–224.

Ferris KG, Rushton T, Greenlee AB, Toll K, Blackman BK, Willis JH. **2015**. Leaf shape

evolution has a similar genetic architecture in three edaphic specialists within the *Mimulus*

guttatus species complex. *Annals of Botany* **116**: 213–223.

Ferris KG, Sexton JP, Willis JH. **2014**. Speciation on a local geographic scale: The evolution

of a rare rock outcrop specialist in *Mimulus*. *Philosophical Transactions of the Royal Society B:*

Biological Sciences **369**: 27–29.

Ferris KG, Willis JH. **2018**. Differential adaptation to a harsh granite outcrop habitat between

sympatric *Mimulus* species. *Evolution* **72**: 1225–1241.

Fishman L, Sweigart AL, Kenney AM, Campbell S. **2014**. Major quantitative trait loci control

divergence in critical photoperiod for flowering between selfing and outcrossing species of

monkeyflower (*Mimulus*). *New Phytologist* **201**: 1498–1507.

Friedman J, Willis JH. **2013**. Major QTLs for critical photoperiod and vernalization underlie

extensive variation in flowering in the *Mimulus guttatus* species complex. *New Phytologist* **199**:

571–583.

Fry JD. **1996**. The Evolution of Host Specialization: Are Trade-Offs Overrated? *The American*

Naturalist **148**: S84–S107.

Ghalambor CK, Hoke KL, Ruell EW, Fischer EK, Reznick DN, Hughes KA. **2015**. Non-

adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature* **525**:

372–375.

Gosden TP, Waller JT, Svensson EI. **2015**. Asymmetric isolating barriers between different

microclimatic environments caused by low immigrant survival. *Proceedings of the Royal Society*

B: Biological Sciences **282**.

Grant BR, Grant PR. **1993**. Evolution of Darwin's Finches Caused by a Rare Climatic Event.

Proceedings of the Royal Society B: Biological Sciences **251**: 111–117.

Hall MC, Willis JH. **2006**. Divergent Selection on Flowering Time Contributes To Local

Adaptation in *Mimulus Guttatus* Populations. *Evolution* **60**: 2466.

Harder LD, Johnson SD. **2009**. Darwin's beautiful contrivances: evolutionary and functional

evidence for floral adaptation. *New Phytologist* **183**: 530–545.

Harrell Jr F. **2023**. Hmisc: Harrell Miscellaneous. R package version 5.1-2.

Hastings A. **1983**. Can spatial variation alone lead to selection for dispersal? *Theoretical*

Population Biology **24**: 244–251.

Hereford J. **2009**. A quantitative survey of local adaptation and fitness trade-offs. *American*

Naturalist **173**: 579–588.

Kawecki TJ, Ebert D. **2004**. Conceptual issues in local adaptation. *Ecology Letters* **7**: 1225–
1241.

Kenney AM, Sweigart AL. **2016**. Reproductive isolation and introgression between sympatric
Mimulus species. *Molecular ecology* **25**: 2499–2517.

Kingsolver JG, Diamond SE. **2011**. Phenotypic selection in natural populations: What limits
directional selection? *American Naturalist* **177**: 346–357.

Kingsolver JG, Pfennig DW. **2004**. Individual-level selection as a cause of cope's rule of
phyletic size increase. *Evolution* **58**: 1608–1612.

Kooyers NJ, Colicchio JM, Greenlee AB, Patterson E, Handloser NT, Blackman BK. **2019**.
Lagging adaptation to climate supersedes local adaptation to herbivory in an annual
monkeyflower. *American Naturalist* **194**: 541–557.

Kooyers NJ, Greenlee AB, Colicchio JM, Oh M, Blackman BK. **2015**. Replicate altitudinal
clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual
Mimulus guttatus. *New Phytologist* **206**: 152–165.

Krizek BA, Anderson JT. **2013**. Control of flower size. *Journal of Experimental Botany* **64**:
1427–1437.

Lane JE, Kruuk LEB, Charmantier A, Murie JO, Dobson FS. **2012**. Delayed phenology and
reduced fitness associated with climate change in a wild hibernator. *Nature* **489**: 554–557.

Latreille AC, Pichot C. **2017**. Local-scale diversity and adaptation along elevational gradients
assessed by reciprocal transplant experiments: lack of local adaptation in silver fir populations.
Annals of Forest Science **74**.

Leimu R, Fischer M. **2008**. A meta-analysis of local adaptation in plants. *PLoS ONE* **3**: 1–8.

Love JM, Ferris KG. **2024**. Local adaptation to an altitudinal gradient: The interplay between
mean phenotypic trait variation and phenotypic plasticity in *Mimulus laciniatus*. *Perspectives in*
Plant Ecology, Evolution and Systematics **63**: 125795.

Lowry DB, Hall MC, Salt DE, Willis JH. **2009**. Genetic and physiological basis of adaptive
salt tolerance divergence between coastal and inland *Mimulus guttatus*. *New Phytologist* **183**:
776–788.

Lowry DB, Rockwood RC, Willis JH. **2008**. Ecological reproductive isolation of coast and
inland races of *Mimulus guttatus*. *Evolution* **62**: 2196–2214.

Magnusson A, Skaug HJ, Nielsen A, Berg CW, Kristensen K, Maechler M, Bentham KJ

van, Bolker BM, Brooks ME. **2017**. glmmTMB: Generalized Linear Mixed Models using

Template Model Builder. R package version 0.1.3.

Mantel SJ, Sweigart AL. **2019**. Divergence in drought-response traits between sympatric

species of *Mimulus*. *Ecology and Evolution* **9**: 10291–10304.

Mills LS, Zimova M, Oyler J, Running S, Abatzoglou JT, Lukacs PM. **2013**. Camouflage

mismatch in seasonal coat color due to decreased snow duration. *Proceedings of the National*

Academy of Sciences of the United States of America **110**: 7360–7365.

Mojica JP, Kelly JK. **2010**. Viability selection prior to trait expression is an essential

component of natural selection. *Proceedings of the Royal Society B: Biological Sciences* **277**:

2945–2950.

Mojica JP, Lee YW, Willis JH, Kelly JK. **2012**. Spatially and temporally varying selection on

intrapopulation quantitative trait loci for a life history trade-off in *Mimulus guttatus*. *Molecular*

Ecology **21**: 3718–3728.

Monnahan PJ, Kelly JK. **2015**. Naturally segregating loci exhibit epistasis for fitness. *Biology*

Letters **11**.

Nicotra AB, Leigh A, Boyce CK, Jones CS, Niklas KJ, Royer DL, Tsukaya H. **2011**. The

evolution and functional significance of leaf shape in the angiosperms. *Functional Plant Biology*

38: 535–552.

Oakley CG, Schemske DW, McKay JK, Ågren J. **2023**. Ecological genetics of local

adaptation in *Arabidopsis*: An 8-year field experiment. *Molecular Ecology* **32**: 4570–4583.

Pélabon C, Hilde CH, Einum S, Gamelon M. **2020**. On the use of the coefficient of variation

to quantify and compare trait variation. *Evolution Letters* **4**: 180–188.

Pinheiro J, Bates D. **2000**. *Mixed-Effects Models in S and S-PLUS*. ew York: Springer.

Pinheiro J, Bates D, R Core Team. **2023**. nlme: Linear and Nonlinear Mixed Effects Models. R

package version 3.1-163.

Ramsey J, Bradshaw HD, Schemske DW. **2003**. Components of reproductive isolation

between the monkeyflowers *Mimulus lewisii* and *M*. *cardinalis* (Phrymaceae). *Evolution* **57**:

1520–1534.

Rundle HD, Nosil P. **2005**. Ecological speciation. *Ecology Letters* **8**: 336–352.

Schiestl FP, Schluter PM. **2009**. Floral isolation, specialized pollination, and pollinator

behavior in orchids. *Annual Review of Entomology* **54**: 425–446.

Schluter D. **2000**. *The Ecology of Adaptive Radiation*. Oxford Series in Ecology and Evolution.

Schluter D. **2009**. Evidence for ecological speciation and its alternative. *Science* **323**: 737–741.

Schneider CA, Rasband WS, Eliceiri KW. **2012**. NIH Image to ImageJ: 25 years of Image

Analysis. *Nature Methods* **9**: 671–675.

Siepielski AM, Dibattista JD, Carlson SM. **2009**. It's about time: The temporal dynamics of

phenotypic selection in the wild. *Ecology Letters* **12**: 1261–1276.

Siepielski AM, Morrissey MB, Buoro M, Carlson SM, Caruso CM, Clegg SM, Coulson T,

Dibattista J, Gotanda KM, Francis CD, *et al*. 2017. Precipitation drives global variation in

natural selection. *Science* **355**: 959–962.

Simko WT. **2021**. R package 'corrplot': Visualization of a Correlation Matrix. (Version 0.92).

Stearns S. **1992**. *The evolution of life histories*. Oxford, UK: Oxford University Press.

Tataru D, De Leon M, Dutton S, Machado-Perez F, Rendahl A, Ferris KG. **2024**.

Fluctuating selection in a Monkeyflower hybrid zone. *bioRxiv*: 1–33.

Tataru D, Wheeler EC, Ferris KG. **2023**. Spatially and temporally varying selection influence

species boundaries in two sympatric *Mimulus*. *Proceedings of the Royal Society B: Biological*

Sciences **290**.

Toll K, Willis JH. **2018**. Hybrid inviability and differential submergence tolerance drive habitat

segregation between two congeneric monkeyflowers. *Ecology* **99**: 2776–2786.

Troth A, Puzey JR, Kim RS, Willis JH, Kelly JK. **2018**. Selective trade-offs maintain alleles

underpinning complex trait variation in plants. *Science* **361**: 475–478.

Venail J, Dell'Olivo A, Kuhlemeier C. **2010**. Speciation genes in the genus *Petunia*.

Philosophical Transactions of the Royal Society B: Biological Sciences **365**: 461–468.

Via S, Lande R. **1985**. Genotype-environment interaction and the evolution of phenotypic

plasticity. *Evolution* **39**: 505–522.

Vickery RK. **1964**. Barriers to gene exchange between members of the *Mimulus guttatus*

complex (Scrophulariaceae). *Evolution* **18**: 52.

Wadgymar SM, Daws SC, Anderson JT. **2017**. Integrating viability and fecundity selection to

illuminate the adaptive nature of genetic clines. *Evolution Letters* **1**: 26–39.

Wilczek AM, Cooper MD, Korves TM, Schmitt J. **2014**. Lagging adaptation to warming

climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United*

States of America **111**: 7906–7913.

Wu CA, Lowry DB, Cooley AM, Wright KM, Lee YW, Willis JH. **2008**. *Mimulus* is an

emerging model system for the integration of ecological and genomic studies. *Heredity* **100**:

220–230.

Figure 1. Temporal and spatial scale of the experimental design. Annual changes in a) snow water content and b) accumulated precipitation during experimental years of low snowpack (2013, 2021, 2022) and comparatively higher snowpack (2019 and 2023) with the c) spatial layout of experimental sites: meadow 1 and 2 (filled circles) and granite 1 and 2 (open circles). Nearby parental populations are shown: *M. guttatus* YVO (filled square) and *M. laciniatus* WLF (open triangle). Representative meadow and granite habitats are shown in the insets. Water data are averaged from the California Department of Water Resources data stations GIN, WHW, and TUM, located nearby experimental sites along Tioga Road in Yosemite National Park (cdec.water.ca.gov).

787 **Figure 2.** Seasonal changes in survival (%), soil moisture (%), light intensity (μmol m-2 s-1),

788 and soil surface temperature (°F) in each block per site a) meadow and b) granite habitats in low

789 snowpack years (2013, 2021, 2022) and high snowpack years (2019, 2023). Data from 2013 and

790 2019 are published in Ferris and Willis 2018 and Tataru *et al*. 2023 and re-analyzed here.

Figure 3. a) Survival to flowering, b) fecundity (mean seed number per reproductive individual), and c) total fitness (mean seed number per planted seed) of *Mimulus guttatus* (filled square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite habitat, meadow habitat, and averaged across years. Calculations of fecundity and total fitness for 2013 are based on fruit number because seed number was not recorded; 2013 was omitted from averaged fecundity and total fitness. Values from 2013 and 2019 are adapted from Ferris and Willis 2018 and Tataru *et al*. 2023.

Figure 4. Visual representation of selection gradients (β) measured for hybrids during experimental years in granite (open circles) and meadow (filled circles) habitats. Shown are a) predicted values in each habitat based on native species trait differences and estimated values from b) negative binomial models and c) zero-truncated Poisson models. Significance level is indicated with asterisks and negative graph areas are shaded grey for clarity. The absence of a value during an experimental year indicates that the trait was not present in the best-fit model. All β values are based on seed number, except for values from 2013 which are based on fruit number, as seed number was not recorded. Values from 2013 and 2019 are adapted from Ferris and Willis 2018 and Tataru *et al.* 2023.

Figure 5. Patterns of cumulative selection with all years combined in a) meadow (filled circles)

and b) granite (open circles) habitat in a zero-truncated Poisson analysis on flowering time (days

from planting), width of first flower (mm), leaf lobing index, and plant height at flowering (mm

to apical meristem). Y-axes show adjusted relative fitness (fruit number), statistically corrected

for other variables included in the models, and the x-axes show unstandardized trait values for

biological relevance. Data from 2013 and 2019 are published in Ferris and Willis 2018 and

Tataru *et al.* 2023 and re-analyzed here.

	2021			2022			2023			
	F_2	*M. guttatus* (YVO18)	*M. laciniatus* (WLF47)	F_3	*M. guttatus* (YVO18)	*M. laciniatus* (WLF47)	F_6	*M. guttatus* (YVO)	*M. laciniatus* (WLF)	"Local"
Meadow	63%	26%	69%	21%	10%	44%	32%	33%	72%	0%
Granite	46%	2%	35%	1%	0%	23%	8%	12%	47%	0%

828 **Table 1**. Percentage surviving to first flower of hybrids, *M. guttatus* (YVO), *M. laciniatus*

829 (WLF), and local site native genotypes (2023 only).

Genotype	Trait	Meadow			Granite		
		Mean	SE	CV	Mean	SE	CV
F$_2$	Fl. time	31.58***	0.61	0.45	14.57***	0.31	0.43
	Height	57.52***	1.40	0.57	28.43***	0.63	0.45
	Fl. Width	5.89	0.11	0.37	6.09	0.12	0.36
	Leaf area	48864.77**	3838.85	1.57	33811.16**	2769.86	1.52
	Lobing	0.08***	0.00	0.45	0.07***	0.00	0.37
	Herkogamy	-1.34***	0.05	-0.73	-0.86***	0.07	-1.36
M. guttatus (YVO 18)	Fl. time	44.59*	0.99	0.34	35.08*	4.23	0.42
	Height	74.65***	2.56	0.46	20.56***	3.50	0.51
	Fl. Width	7.49	0.22	0.33	5.88	0.81	0.36
	Leaf area	50215.94***	6314.38	1.47	11519.75***	1912.33	0.94
	Lobing	0.08*	0.00	0.68	0.06*	0.00	0.33
	Herkogamy	-1.63	0.10	-0.76	-0.94	0.31	-0.94
M. laciniatus (WLF 47)	Fl. time	26.7***	0.49	0.45	19.21***	0.54	0.48
	Height	34.04***	0.75	0.49	20.41***	0.72	0.57
	Fl. Width	1.96***	0.05	0.49	2.55***	0.15	0.73
	Leaf area	16642.84**	1268.08	1.46	12059.12**	1095.45	1.07
	Lobing	0.08***	0.00	0.47	0.07***	0.03	0.40
	Herkogamy	-0.92	0.10	-1.17	-0.74	0.08	-1.01

832 **Table 2**. Trait means, standard error (SE), standard deviation (SD), and coefficient of variation

833 (CV) from the 2021 experiment. Significant variation in trait means between habitat is indicated

834 by asterisks (p-value < 0.001***; 0.01**; 0.05*).

Genotype	Trait	Meadow			Granite		
		Mean	SE	CV	Mean	SE	CV
F$_3$	Fl. time	59.92***	0.70	0.21	44.19***	3.10	0.32
	Height	54.89***	2.07	0.69	13.14***	1.52	0.53
	Fl. Width	5.52***	0.13	0.41	3.34***	0.30	0.41
	Leaf area	1013.85***	77.72	1.31	208.62***	27.30	1.09
	Lobing	0.07***	0.00	0.33	0.09***	0.00	0.24
	Herkogamy	-1.35*	0.05	-0.67	-0.92*	0.17	-0.77
M. guttatus (YVO 18)	Fl. time	72.66	1.89	0.17	-	-	-
	Height	69.15	5.22	0.48	-	-	-
	Fl. Width	7.73	0.36	0.29	-	-	-
	Leaf area	1788.03***	220.71	0.71	169.69***	22.08	0.47
	Lobing	0.05**	0.00	0.27	0.08**	0.00	0.22
	Herkogamy	-2.40	0.13	-0.33	-	-	-
M. laciniatus (WLF 47)	Fl. time	44.53***	0.83	0.25	34.65***	1.08	0.30
	Height	15.61***	0.99	0.84	7.16***	0.42	0.57
	Fl. Width	1.25	0.08	0.74	1.22	0.14	0.97
	Leaf area	294.09***	22.64	0.97	97.56***	4.31	0.41
	Lobing	0.08*	0.00	0.30	0.09*	0.00	0.29
	Herkogamy	-0.85	0.05	-0.65	-1.01	0.11	-0.71

839 **Table 3**. Trait means, standard error (SE), standard deviation (SD), and coefficient of variation

840 (CV) from the 2022 experiment. Significant variation in trait means between habitat is indicated

841 by asterisks (p-value < 0.001***; 0.01**; 0.05*)

Genotype	Trait	Meadow				Granite			
		Mean	SE	SD	CV	Mean	SE	SD	CV
F$_6$	Fl. time	43.04***	0.28	15.3	0.36	35.77***	0.36	19.67	0.55
	Height	91.71***	1.01	55.14	0.60	25.22***	0.27	14.81	0.59
	Fl. Width	5.62***	0.04	2.23	0.40	4.68***	0.04	2.35	0.50
	Leaf area	14342.24***	287.00	15722.42	1.10	6183.81***	173.97	9528.94	1.54
	Lobing	0.08	0.00	0.05	0.63	0.08	0.00	0.06	0.75
	Herkogamy	-0.34*	0.03	1.4	-4.12	-0.4	0.02	1.09	-2.73
M. guttatus (YVO)	Fl. time	52.94**	0.61	12.14	0.23	44.6**	0.78	15.49	0.35
	Height	76.17***	1.75	34.94	0.46	34.56***	0.84	16.78	0.49
	Fl. Width	9.92*	0.13	2.68	0.27	8.69*	0.16	3.1	0.36
	Leaf area	18876.1***	942.86	18833.71	1.00	11853.5***	655.40	13091.55	1.10
	Lobing	0.07**	0.00	0.03	0.43	0.08**	0.00	0.02	0.25
	Herkogamy	-0.66	0.08	1.5	-2.27	-1.11	0.07	1.41	-1.27
M. laciniatus (WLF)	Fl. time	24.28***	0.71	14.19	0.58	20.68***	0.92	18.36	0.89
	Height	58.72***	1.72	34.45	0.59	21.16***	0.56	11.16	0.53
	Fl. Width	3.01	0.08	1.59	0.53	3.37	0.07	1.44	0.43
	Leaf area	17699.42***	850.63	16991.34	0.96	8788.1***	327.31	6537.97	0.74
	Lobing	0.15**	0.00	0.08	0.53	0.11**	0.00	0.07	0.64
	Herkogamy	-0.36	0.05	0.98	-2.72	0.12	0.05	0.92	7.67

844 **Table 4**. Trait means, standard error (SE), standard deviation (SD), and coefficient of variation

845 (CV) from the 2023 experiment. Significant variation in trait means between habitat is indicated

846 by asterisks (p-value < 0.001***; 0.01**; 0.05*).

Trait	Meadow			Granite	
	2021	**2022**	**2023**	**2021**	**2023**
	β F2	**β F3**	**β F6**	**β F2**	**β F6**
Flowering time (FT)	-0.900***	-	-0.358**	-0.632*	-0.730**
Flower width (FW)	-0.239	0.551***	0.213	-0.288	0.893***
Leaf lobing (LL)	-0.143	-0.028	-0.062	-	-
Herkogamy (H)	-	-	0.060	-	-
Plant height (PH)	0.903***	0.137	0.662***	1.461**	2.571**
FW x PH	-0.291*	-	-0.251*	-1.057	-
FT x H	-	-	0.237*	-	-
LL x PH	-0.232	0.096*	-	-	-
FT x PH	-	-	-	-	-
FW x LL	-	-	-0.483**	-	-
H x PH	-	-	-	-	-
FT x FW	-	-	-	-	-
n =	200	258	775	166	214

Table 5. Negative binomial analysis on whether or not a plant produced seeds. Values indicate the selection gradient (β), or strength of selection. Asterisks indicate significance of selection in a trait: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Missing values indicate traits which were not included in the best-fit model.

Trait	Meadow			Granite	
	2021	**2022**	**2023**	**2021**	**2023**
	β F2	β F3	β F6	β F2	β F6
Flowering time (FT)	-1.827***	-0.289***	-0.195***	-0.432	0.173
Flower width (FW)	0.957***	0.178*	-0.097*	0.415***	1.254***
Leaf lobing (LL)	-0.374***	0.094**	-0.216***	-0.794***	-4.978***
Herkogamy (H)	-0.436***	0.167***	0.195***	-1.610***	-0.800***
Plant height (PH)	0.615***	0.563***	0.577***	2.15***	-4.056***
FT x FW	1.964***	0.542***	0.317***	-	-0.854***
FT x H	-0.956***	-0.314***	0.311***	-2.022***	-
FT x PH	-0.651***	-0.420***	-0.097*	1.110*	-
FW x LL	0.715***	-	-0.423***	-	1.667***
FW x PH	-0.601***	-0.154***	-0.126***	-	-1.700***
LL x PH	-0.186***	-	0.241***	-1.517***	-8.017***
H x PH	0.401***	0.072**	-0.072***	0.769***	-1.978***
n =	128	134	202	87	51

858 **Table 6**. Zero-truncated Poisson analysis on number of seeds produced if any were produced.

859 Values indicate the selection gradient (ß), or strength of selection. Asterisks indicate significance

860 of selection in a trait: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Missing values indicate traits were

861 not included in the best-fit model.

Trait	Negative binomial		Zero-truncated Poisson	
	Meadow	**Granite**	**Meadow**	**Granite**
Flowering time (FT)	-0.428***	-0.807***	-1.082***	-1.122***
Flower width (FW)	-0.069*	-0.091**	-0.239***	-0.192*
Leaf lobing (LL)	-	-	0.056	0.198
Plant height (PH)	0.602***	0.853***	0.962***	1.056***
FT x FW	-0.093*	-	-0.218***	-0.163
FT x PH	-0.150***	-0.570***	-0.114	-0.348**
FT x LL	-	-	-	-0.305**
FW x LL	-	-	-0.059	0.096
FW x PH	-	0.118**	0.076	0.195*
LL x PH	-	-	0.040	0.472
FT x FW x LL	-	-	-	0.277**
FT x LL x PH	-	-	-	-0.223*
FT x FW x PH	-	-	0.109*	-
FW x LL x PH	-	-	-0.162*	-0.304***
n =	2,014	992	998	547

864
Table 7. Phenotypic selection analyses on a combined dataset of 2013, 2019, 2021, 2022, and
2023 using fruit number as a metric of fitness. Values indicate the selection gradient (ß), or
strength of selection. Asterisks indicate significance of selection in a trait: ***p < 0.001; **p <
0.01; *p < 0.05. Missing values indicate traits were not included in the best-fit model.
Herkogamy was excluded due to its absence from the 2013 dataset. Data from 2013 and 2019 are
published in Ferris and Willis 2018 and Tataru *et al.* 2023 and re-analyzed here.
871
872

Supplementary Information

Fig S1 Combined survival to flowering, fecundity, and total fitness excluding 2023.

Fig S2 Trait correlation matrices for F2 hybrids in 2021 in meadow and granite habitats.

Fig S3 Trait correlation matrices for F3 hybrids in 2022 in meadow and granite habitats.

Fig S4 Trait correlation matrices for F6 hybrids in 2023 in meadow and granite habitats.

Fig S5 Patterns of cumulative selection with all years combined in meadow and granite habitats in a negative binomial analysis.

Table S1 Intrapopulation parental crosses used in the 2023 experiment.

Table S2 Model selection for best fit models of plant survival and environmental variables.

Table S3 Model selection for best fit models of hybrid traits.

Table S4 Model selection for best fit models of herbivory.

Table S5 Negative binomial analysis on whether or not a plant produced fruits.

Table S6 Zero-truncated Poisson analysis on number of fruit produced if any were produced.

Table S7 Strength of correlations between fruit number and seed number.

890
891
Figure S1. Combined survival to flowering, fecundity (mean seed number per reproductive

individual), and total fitness (mean seed number per planted seed) of *Mimulus guttatus* (filled

square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite and meadow habitat

from 2013-2022 (excluding 2023). 2013 was omitted from fecundity and total fitness

calculations because seed number was not recorded, only fruit number. Values from 2013 and

2019 are adapted from Ferris and Willis 2018 and Tataru *et al.* 2023.

898
899

900 **Figure S2.** Trait correlation matrices for F2
901 hybrids in 2021 in meadow (left) and granite (right) habitats.

903
904
905 **Figure S3.** Trait correlation matrices for F3 hybrids in 2022 in meadow (left) and granite (right)

906 habitats.

908
Figure S4. Trait correlation matrices for F6 hybrids in 2023 in meadow (left) and granite (right)

habitats.

Figure S5. Patterns of cumulative selection with all years combined in a) meadow (filled circles)

and b) granite habitats (open circles) in a negative binomial analysis on flowering time (days

from planting), width of first flower (mm), leaf lobing index, and plant height at flowering (mm

to apical meristem). Y-axes show adjusted relative fitness (fruit number), statistically corrected

for other variables included in the models, and the x-axes show unstandardized trait values for

biological relevance. Data from 2013 and 2019 are published in Ferris and Willis 2018 and

Tataru *et al.* 2023 and re-analyzed here.

920
921

Habitat	Genotype	Survival to flowering	%Survival	Seed count	Mean Seed Count	Seed SD	N
Granite	WLF60xWLF47	81	44	585	7.31	21.27	184
	WLF61xWLF47	108	66	784	7.26	12.78	164
	WLF47 inbred	0	0	0	NA	NA	51
						Total:	399
	YVO18xYVO26	0	0	0	NA	NA	20
	YVO25xYVO18	12	8	417	32.08	67.69	159
	YVO26xYVO18	35	18	961	27.46	51.83	200
	YVO18 inbred	0	0	0	NA	NA	20
						Total:	399
Meadow	WLF60xWLF47	97	58	1760	17.25	32.41	166
	WLF61xWLF47	156	83	3201	20.13	33.95	187
	WLF47inbred	35	76	156	4.46	8.43	46
						Total:	399
	YVO18xYVO26	26	47	208	8	16.78	55
	YVO25xYVO18	26	23	378	14.54	47.43	114
	YVO26xYVO18	55	32	687	12.27	27.66	171
	YVO18 inbred	26	44	21	0.81	2.77	59
						Total:	399

922
923 **Table S1**. Intrapopulation parental crosses used in the 2023 experiment. Due to insufficient seed

924 stock of single crosses of YVO and WLF, we used 2-3 crosses and original inbred lines,

925 distributed evenly across blocks and sites. With habitats combined, WLF genotypes did not

926 differ significantly in total seed count ($F = 2.324$, $df = 2$, $p = 0.099$). Similarly, YVO genotypes

927 also did not differ significantly ($F = 1.944$, $df = 3$, $p = 0.124$).

Year	Habitat	Model	df	Log(L)	AIC	delta(i)	wi
2021	meadow	survival ~ time + light + soil + time:soil	7	1338.307	-2662.6	4.56	0.869
2021	granite	survival ~ time + light + soil + temp + time:light + light:temp	9	127.115	-235.9	0.38	0.546
2021	both	survival ~ time + light + soil + temp + time:light + time:soil + light:temp + soil:temp	11	1232.41	-2442.8	13.76	0.997
2022	meadow	survival ~ time + light + soil + temp + time:temp	8	608.991	-1201.9	1.12	0.608
2022	granite	survival ~ time + light + soil + temp + time:temp + light:soil + soil:temp	10	625.324	-1230.4	1.4	0.668
2022	both	survival ~ time + light + soil + temp + time:temp + light:soil + soil:temp	10	882.345	-1744.6	22.24	1
2023	meadow	survival ~ time + soil + temp + soil:temp	7	1580.3	-3146.6	2.4	0.666
2023	granite	survival ~ time + light + soil + temp + time:light + time:soil + time:temp + soil:temp	11	656.63	-1291.1	1.5	0.502
2023	both	survival ~ time + soil + temp + time:soil + time:temp + soil:temp + time:soil:temp	10	1190.095	-2360.1	4.69	0.886

Table S2. Best-fit models of plant survival and environmental variables. For each analysis, we report k, the number of parameters; Log(L), the likelihood score; AIC score, delta(i), the difference between that model's AIC and the top model's AIC; and wi, the model weight.

Year	Term	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
2021	Herbivory	1	50597	50597	25.25	5.43E-07	***
	Site	3	604179	201393	100.52	< 2e-16	***
	Herbivory*Site	3	75224	25075	12.52	4.10E-08	***
2022	Block	1	153	152.6	0.803	0.37046	
	Site	1	5	4.5	0.024	8.78E-01	**
	Block*Site	1	1961	1961.3	10.323	0.00139	*
2023	Block	1	1048	1048	2.193	0.13887	
	Site	1	4412	4412	9.23E+00	0.00242	**
	Block*Site	1	1832	1832	3.832	0.05049	

937 **Table S3**. Best-fit models of plant fitness and herbivory.

Genotype	Habitat	Metric	Best-fit model	df	Log(L)	AIC	delta(i)	wi
colspan Zero-truncated Poisson (seed)								
F2	granite	seed	seed1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:H + FT:PH + LL:PH + H:PH	12	-343.824	715.9	1.98	0.338
F2	meadow	seed	seed1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:FW + FT:H + FT:PH + FW:LL + FW:PH + LL:PH + H:PH	15	-1944.783	3923.9	28.56	1
F3	meadow	seed	seed1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:FW + FT:H + FT:PH + FW:PH + H:PH	13	-633.098	1295.2	0.85	0.329
F6	meadow	seed	seed1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:FW + FT:H + FT:PH + FW:LL + FW:PH + LL:PH + H:PH	15	-1332.565	2697.7	2.47	0.771
F6	granite	seed	seed1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:FW + FW:LL + FW:PH + LL:PH + H:PH	13	-349.91	735.7	0.2	
colspan Negative binomial (seed)								
F2	granite	seed	seed ~ FT + FW + PH + (1 | Site/Block) + FW:PH	9	-412.503	844.2	0.04	
F2	meadow	seed	seed ~ FT + FW + LL + PH + (1 | Site/Block) + FW:PH + LL:PH	11	-744.987	1513.4	0.05	0.044
F3	meadow	seed	seed ~ FW + LL + PH + (1 | Site/Block) + LL:PH	9	-663.336	1345.4	0.86	

F6	meadow	seed	seed ~ FT + FW + LL + H + PH + (1 | Site/Block) + FT:H + FW:LL + FW:PH	13	-1188.134	2402.7	0.5	
F6	granite	seed	seed ~ FT + FW + PH + (1 | Site/Block)	8	-299.661	616	0.8	
Zero-truncated Poisson (fruit)								
F2	granite	fruit	fruit1 ~ FT + PH + (1 | Site/Block)	5	-126.834	264.3	0.17	0.05
F2	meadow	fruit	fruit1 ~ FT + FW + LL + H + PH + (1 | Site/Block) + LL:PH + H:PH	10	-301.67	624.9	1.13	0.104
F3	meadow	fruit	fruit1 ~ PH + (1 | Site/Block)	4	-101.927	212.1	0.64	0.055
F6	meadow	fruit	fruit1 ~ FT + FW + H + PH + (1 | Site/Block) + H:PH	8	-302.086	620.7	0.44	0.085
F6	granite	fruit	fruit1 ~ FT + FW + PH + (1 | Site/Block) + FT:PH	7	-66.555	148.7	0	
Combined	meadow	fruit	fruit1 ~ FT + FW + LL + PH + (1 | Site) + (1 | year) + FT:FW + FT:PH + FW:LL + FW:PH + LL:PH + FT:FW:PH + FW:LL:PH	14	-1265.383	2559.2	1.13	0.093
Combined	granite	fruit	fruit1 ~ FT + FW + LL + PH + (1 | Site) + (1 | year) + FT:FW + FT:LL + FT:PH + FW:LL + FW:PH + LL:PH + FT:FW:LL + FT:LL:PH + FW:LL:PH	16	-593.575	1220.2	0.35	0.115
Negative binomial (fruit)								
F2	granite	fruit	fruit ~ PH + (1 | Site/Block)	6	-242.524	497.6	0.31	
F2	meadow	fruit	fruit ~ FT + FW + H + PH + (1 | Site/Block) + FT:FW + FW:PH + H:PH	12	-399.2	824.1	0.07	0.065

				df	Log(L)	AIC	delta(i)	wi
F3	meadow	fruit	fruit ~ FW + LL + (1 | Site/Block)	7	-289.138	592.7	0	
F6	meadow	fruit	fruit ~ FT + FW + LL + PH + (1 | Site/Block) + FT:PH	10	-811.743	1643.8	0.71	0.09
F6	granite	fruit	fruit ~ FT + FW + PH + (1 | Site/Block) + FT:PH	9	-194.445	407.8	0.77	
Combined	meadow	fruit	fruit ~ FT + FW + PH + (1 | Site) + (1 | year) + FT:FW + FT:PH	10	-2567.225	5154.6	0.2	0.063
Combined	granite	fruit	fruit ~ FT + FW + PH + (1 | Site) + (1 | year) + FT:PH + FW:PH	10	-1220.19	2460.6	0.41	

940
941
942 **Table S4.** Model selection for best-fit models of hybrid traits in 2021 (F2), 2022 (F3), 2023 (F6),

943 and with years combined (2013, 2019, 2021, 2022, 2023). For each analysis, we report df,

944 degrees of freedom; Log(L), the likelihood score; AIC score, delta(i), the difference between that

945 model's AIC and the top model's AIC; and wi, the model weight. FT, flowering time; FW,

946 flower width; LL, leaf lobing; PH, height; H, for herkogamy (stigma-anther separation).

947
948

949
950
951

Trait	Meadow			Granite	
	2021	2022	2023	2021	2023
	β F2	β F3	β F6	β F2	β F6
Flowering time (FT)	-0.90524***	-	-0.28166**	-	-0.7680*
Flower width (FW)	-0.07251	0.31918***	-0.22248*	-	0.5890***
Leaf lobing (LL)	-	0.12841	0.18433*	-	1.5173***
Herkogamy (H)	-0.04269	-	-	-	-
Plant height (PH)	0.7464***	-	0.39770***	0.5202*	-
FW x PH	-0.20885*	-	-	-	-
LL x PH	-	-	-	-	-
FT x PH	-	-	-0.25262**	-	-0.5164
FW x LL	-	-	-	-	-
H x PH	0.23229*	-	-	-	-
FT x FW	0.25901	-	-	-	-
n =	201	258	776	165	214

952
953
954 **Table S5**. Negative binomial analysis on whether or not a plant produced fruits. Values indicate

955 the selection gradient (β), or strength of selection. Asterisks indicate significance of selection in a

956 trait: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Missing values indicate traits which were not included

957 in the best-fit model.

| Trait | Meadow | | | Granite | |
| | 2021 | 2022 | 2023 | 2021 | 2023 |
	β F2	β F3	β F6	β F2	β F6
Flowering time (FT)	-0.92254***	-	-1.3223**	-0.8667**	-0.7924***
Flower width (FW)	-0.31538***	-	0.7348***	-	-0.40193***
Leaf lobing (LL)	0.11745	-	-	-	-
Herkogamy (H)	-0.04592	-	-	-	0.22461*
Plant height (PH)	0.62646***	0.4410***	1.6214**	1.4328***	0.49728***
FT x FW	-	-	-	-	-
FT x H	-	-	-	-	-
FT x PH	-	-	-1.0061	-	-
FW x LL	-	-	-	-	-
FW x PH	-	-	-	-	-
LL x PH	-0.15017*	-	-	-	-
H x PH	0.24256**	-	-	-	-0.15321**
n =	150	171	79	107	274

960 **Table S6.** Zero-truncated Poisson analysis on number of fruit produced if any were produced.

961 Values indicate the selection gradient (ß), or strength of selection. Asterisks indicate significance

962 of selection in a trait: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Missing values indicate traits were

963 not included in the best-fit model.

Year	Habitat	τ	p	n
2021	Meadow	0.56	0.001	1296
	Granite	0.59	0.001	809
2022	Meadow	0.64	0.001	545
2023	Meadow	0.75	0.0001	1393
	Granite	0.69	0.0001	480

967 **Table S7.** Strength of correlations between fruit number and seed number calculated using

968 Kendall's correlation analysis.

969
970