1 RUNNING TITLE: Fluctuating selection on Monkeyflowers

4 TITLE: Short-term fluctuating and long-term divergent selection on sympatric Monkeyflowers:

5 insights from repeated reciprocal transplants

Caroline M. Dong[1,2], Bolívar Aponte Rolón[1,3], Juj K. Sullivan[1], Diana Tataru[1,4], Max De Leon[1],

Rachael Dennis[1], Spencer Dutton[1], Fidel J. Machado Perez[1,5], Lissette Montano[1] & Kathleen G.

Ferris[1]

[1]Tulane University, Department of Ecology and Evolutionary Biology, New Orleans, LA

[2]Grinnell College, Department of Biology, Grinnell, IA

[3]Iowa State University, Department of Agricultural and Biosystems Engineering, Ames, IA

[4]Utah State University, Department of Biology, Logan, UT

[5]University of California Merced, Life and Environmental Sciences Department, Merced, CA

Corresponding author: dongca@grinnell.edu

KEYWORDS (5-8): adaptation, phenology, speciation, divergence, environmental variation,

evolution, natural selection, temporal variation

ABSTRACT:

Sympatric species are often locally adapted to distinct microhabitats. However, temporal

variation may cause local maladaptation and species boundary breakdown, especially during

extreme climatic events. Repeated reciprocal transplants can reveal the interplay between

spatially and temporally varying patterns of natural selection. To examine long-term patterns of

selection between sympatric Monkeyflowers occupying dramatically different niches, *Mimulus*

guttatus and *M. laciniatus,* we performed three replicated transplants and combined them with

previous experiments to leverage a dataset of five single-year transplants spanning a decade. We

performed phenotypic selection analyses on parents and hybrids in each species' habitat in

Yosemite NP, CA during years of drastically differing snowpack. If there is ecological isolation,

then we predicted divergent phenotypic selection between habitats in line with species'

differences and local adaptation. We found interannual fluctuations in phenotypic selection,

sometimes in unpredicted directions. However, a combined-year analysis detected longer-term

differences in the magnitude of selection between habitats on flowering time, a key temporally

isolating and adaptative trait, suggesting that selection may reinforce species boundaries despite

short-term fluctuations. Finally, we found temporal variation in local adaptation with *M.*

laciniatus locally adapted in low snowpack years, while an extreme snowfall event contributed to

overall local maladaptation of *M. guttatus*.

INTRODUCTION

Closely related species often occupy distinct habitats within areas of geographic overlap (Coyne

and Orr 2004; Rundle and Nosil 2005; Schluter 2009). Species boundaries in sympatry can be

maintained by adaptation to differing environments that generate pre- and post-zygotic isolation

via selection against immigrants (e.g. habitat isolation) and intermediate hybrids (e.g. extrinsic

post-zygotic isolation; Rundle & Nosil, 2005). However, identifying the individual traits

contributing to reproductive isolation is difficult because as speciation progresses, multiple forms

of reproductive isolation accumulate between divergent lineages which hinders the measurement

of individual barriers (Coyne and Orr 2004). Once species are partially isolated, neutral

processes also cause the accumulation of phenotypic divergence. Thus, experimental

manipulation is necessary to determine whether habitat isolation is a key barrier, and which traits

are involved in divergent adaptation (Ramsey et al. 2003; Hall and Willis 2006; Lowry et al.

2008). Measuring phenotypic selection on closely related sympatric species in a reciprocal

transplant between native environments can identify whether divergent adaptation is important

for reproductive isolation and speciation (Lowry et al. 2008; Anderson et al. 2015. If so, we

expect each species to have higher fitness in its habitat and phenotypic selection in the direction

of species' differences (Anderson et al. 2015*a*), indicating habitat isolation and local adaptation

to current conditions.

Despite these predictions of local adaptation, maladaptation to native habitats is often observed

in reciprocal transplants (Hereford 2009; Kooyers et al. 2019; Dittmar and Schemske 2023). This

could be due to the prevalence of temporal fluctuations in selection (Siepielski et al. 2009) which

can influence the strength of reproductive isolation and species boundaries (Tataru et al. 2023,

2024). Additionally, current environmental conditions may have changed from the conditions of

original divergence (Coyne and Orr 2004; Rundle and Nosil 2005). If species are no longer

adapted to their current niches due to climate change, then we expect to find local maladaptation

(Crespi 2000) and selection for the non-local species' phenotype (Wilczek et al. 2014; Kooyers

et al. 2019; Anderson and Wadgymar 2020). In this scenario, spatially varying selection may act

contrary to reproductive isolation and in the absence of strong intrinsic barriers, could erode

species boundaries by driving each towards the other's phenotypic optimum.

Variation in selection in longitudinal studies is usually measured on a seasonal or inter-annual

timescale, which is a short-term measurement of population evolution. This leads to the question

of how short-term fluctuations affect long-term evolutionary change in species and populations

(Wadgymar et al. 2017; Kelly 2022; Dittmar and Schemske 2023; Oakley et al. 2023). A

synthesis of phenotypic selection studies by Kingsolver & Diamond (2011) found that while

temporal variation in selection is common, it does not significantly inhibit long-term directional

selection. Therefore, even with fluctuations in the direction of selection, there may be an average

directional or stabilizing trend over a longer timescale. For example, Wadgymar et al. (2017)

found that interannual fluctuations in the direction of selection on the perennial plant *Boechera*

stricta gave rise to long-term patterns consistent with stabilizing selection and local adaptation.

In *Arabidopsis thaliana*, Lee et al. (2024) found long-term divergent selection on cold tolerance

between populations in Italy and Sweden despite interannual fluctuations. However, due to the

arduousness of temporally replicating reciprocal transplant experiments, it is not currently well

understood how short versus longer-term patterns of natural selection compare and impact

species' divergence and reproductive isolation.

Another understudied agent of evolutionary change is episodic selection stemming from sudden

and extreme environmental events. Long-term studies capturing the evolutionary responses to

these dramatic ecological changes are rare yet critical to understand their impact on species

boundaries (Coltman et al. 1999; Pemberton et al. 2022). For instance, through their long-term

study of the Galapagos finches, Grant and Grant (1993) were able to illustrate how an extreme El

Niño year altered selection on beak size and rates of interspecific gene flow. Strong episodes of

selection such as this can impact biodiversity by affecting the evolutionary trajectory of species

adaptation and reproductive isolation (Grant and Grant 1993). Studies such as this are vital

because extreme climatic events are predicted to become more frequent with continuing

anthropogenic climate change (Bailey and van de Pol 2016).

The *Mimulus guttatus* species complex is well-suited for studying how natural selection shapes

species boundaries and reproductive isolation (Wu et al. 2008). *Mimulus guttatus* and *M.*

laciniatus (syn. *Erythranthe guttata* and *E. laciniata*) occur in contrasting microhabitats

throughout central and southern Sierra Nevada, CA. *Mimulus guttatus* inhabits moist meadows

and seeps while *M. laciniatus* occupies rocky outcrops characterized by shallow soils, intense

light, extreme temperatures, and a short growing season (Ferris et al. 2014; Ferris and Willis

2018). *Mimulus laciniatus*, an obligate summer annual, has a smaller stature, lobed leaves, self-

fertilization with reduced floral size, and early rapid flowering (DeMarche et al. 2013; Ferris and

Willis 2018). In contrast, *M. guttatus* populations are facultatively annual, later-flowering,

predominantly outcrossing, round-leaved, and larger in size and flower morphology (Awadalla

and Ritland 1997; Ferris et al. 2014). Despite ecological divergence, reproductive isolation is

incomplete (Vickery 1964; Ferris et al. 2017; Tataru et al. 2024).

Previous studies show evidence of local adaptation to contrasting microhabitats, but it remains

unclear how environmental variation alters the strength or direction of selection over time (Ferris

and Willis 2018). Early flowering, a drought escape strategy, functions as a key pre-mating

barrier in the *Mimulus guttatus* species complex (Hall and Willis 2006; Lowry et al. 2008; Ferris
et al. 2017; Mantel and Sweigart 2019). Larger plant sizes may be advantageous in meadows
where herbivory is higher. Lobed leaves may improve temperature and water regulation through
a reduced boundary layer (Nicotra et al. 2011*a*) and are unique to *Mimulus* species found in
rocky habitat (Ferris et al. 2014; Ferris and Willis 2018; Ferris 2019). Yet how selection on these
and other traits (e.g. mating system and flower morphology) fluctuates through time is unknown.
Both species exhibit phenotypic plasticity with *M. guttatus* exhibiting greater plasticity in plant
size and flowering time than *M. laciniatus* (Ferris & Willis 2018). Such plasticity may be
adaptive in heterogenous environments like *M. laciniatus'* patchy granite outcrops (Via & Lande
1985).
130
Here, we investigate spatial and temporal variation in natural selection between *M. guttatus* and
M. laciniatus using single-year reciprocal transplants repeated across five years under contrasting
climatic conditions (Figure 1). In this study, we add three years of new results to two previously
published years which spanned years of historically low (Ferris and Willis 2018) and high
snowpack (Tataru et al., 2023). Weaker divergent selection on a key isolating barrier, flowering
time, was found during the high snowpack year, potentially eroding species differentiation. We
replicated transplants during low snowpack years (2021, 2022) and an exceptionally high
snowpack year (2023; Figure 1a,b). We address the following questions: (1) How do
environmental variables affect fitness in each species' habitat and induce phenotypic plasticity?
(2) Does the strength and direction of selection on adaptive and reproductively isolating
phenotypes fluctuate? (3) How do short versus long-term patterns of selection compare? (4) Are
M. guttatus and *M. laciniatus* locally adapted to their respective microhabitats? (5) How does
episodic selection affect the trajectory of species divergence? By examining both interannual
variation and overarching signatures of selection, we were able to elucidate the effects of a
changing climate on the maintenance of biodiversity in this system.
146
147   MATERIALS AND METHODS
148   *Repeated reciprocal transplant design*
149   Reciprocal transplants used hybrids created from inbred lines of *M. guttatus* (YVO 18;
150   37.723366, -119.746433; 5,395 ft elevation) and *M. laciniatus* (WLF 47; 37.841533, -

119.59385; 7993 ft elevation; Figure 1c). Advanced generation hybrids were used because recombination over multiple generations breaks up parental haplotypes, increasing phenotypic variation and enabling inferences about selection on individual traits (Figure S1; Nagy, 1997; Hall & Willis, 2006; Groh & Coop, 2024). F_1 hybrids (WLF 47 maternal x YVO 18 paternal) were self-fertilized to generate F_2 seeds used in 2021. Five hundred F_2 hybrids were propagated with single-seed descent to produce F_3 seeds used in 2022 and 300 F_6 lines used in 2023, with some hybrid lines lost likely due to inbreeding depression. In 2021 and 2022, seeds from 200 hybrid maternal families were pooled for transplants. To avoid inbreeding depression in parental genotypes in 2023, inbred lines were intercrossed within each parental population, as *M. guttatus* is susceptible to inbreeding depression (Willis 1999), which may impact inferences about local adaptation. The 2013 and 2019 experiments used F_4 hybrids from the same parental populations (see Ferris & Willis 2018; Tataru et al 2023).

Our transplants were conducted in 2021, 2022, and 2023 in Yosemite National Park, CA, replicating experiments from 2013 and 2019 at four sites (Figure 1c): two granite outcrops (Granite 1 [37.810700, -119.485200; 8,436 ft elevation], Granite 2 [37.843702, -119.573120; 8,117 ft elevation]) and two meadows (Meadow 1 [37.767781, -119.772720; 6,605 ft elevation], Meadow 2 [37.755968, -119.803031; 6,163 ft elevation]). We used precisely the same microhabitats within each site, experimental design, and parental source populations. Sample sizes, maternal families, and hybrid generation differed annually within logistical constraints (Table S1). Seedlings were transplanted in small blocks to capture fine-scale environmental variation and mimic granite habitat patchiness. In 2021 and 2022, each site had 50 randomized blocks (4×6 formation, 1-inch spacing) of 18 and 24 individuals, respectively. In 2021, blocks contained six F_2 hybrids and six of each parent, with Granite 2 having 75 blocks, totaling 4,050 plants. In 2022, blocks contained 16 F_3 hybrids with four of each parent, totaling 4,800 plants. In 2023, local parental genotypes originating from each site were planted only in their origin site, to compare fitness with the nearby source parents (WLF and YVO). We used 2–3 intrapopulation parental crosses and original inbred lines, distributed evenly across blocks and sites, which did not differ in mean seed count and were combined for downstream analyses (Table S2). Each site had 60 blocks (5×7 formation) of 35 individuals, composed of 3–4 of parental genotypes (YVO,

WLF, and local), and 16–23 F_6 hybrids per block, with 1,200 of each parent and 3,000 F_6 hybrids

per habitat totaling 8,400 plants.

We cold-stratified parental *M. laciniatus* and hybrid seeds for 10 days and parental *M. guttatus*

seeds for 5 days at 4°C in the dark in plastic flats filled with soil (Sunshine Mix #4 Professional

Growing Mix, Sun Gro) to break dormancy and synchronize germination for one week in

greenhouses at the University of California (UC) Davis (2021) and UC Merced (2022–2023).

Following germination upon the emergence of cotyledons, individuals were planted into block

designs described above. Mortality within the first week was attributed to transplant shock, dead

germinants were replaced and not included in subsequent analyses.

Environmental & phenotypic data collection

To assess microhabitat variation, environmental measurements were taken weekly from

experimental blocks. We measured soil moisture (%) using the SM150T sensor (Dynamax), soil

surface temperature (°F) using a laser thermometer, and light intensity (μmol m-2 s-1) using a

MQ-200X Sunlight Quantum Meter (Apogee Instruments). Light and temperature were not

measured in 2013 (Ferris and Willis 2018). We recorded survival, herbivory (presence/absence),

and putatively adaptive phenotypes under selection: flowering time, plant height, herkogamy,

flower size, leaf shape (Ferris et al. 2014; Ferris and Willis 2018; Tataru et al. 2023). At first

flowering, we recorded the date and measured plant height (mm), stigma and anther lengths

(mm), and corolla width (mm). Flowering time was calculated as days from transplant to first

flower. Herkogamy (i.e. stigma-anther separation) was calculated by subtracting stigma length

from longest anther length, a metric of outcrossing ability. To assess leaf shape, the first true leaf

was collected one-week post-flowering. Leaves were scanned and analyzed for leaf lobing index

using ImageJ (Schneider et al. 2012; Ferris et al. 2015).

Testing for effects of environmental factors on fitness

To find associations between environmental variables and plant survival within each habitat, we

pooled data across sites and used binomial mixed models with survival as the dependent variable

and soil moisture, time, soil temperature and light levels with their interactions as fixed

independent variables, and block nested within site as a random effect. Survival was structured

as the number of survivors in each block at a time point relative to the number of individuals

originally planted in each block. We performed model selection for the best-fit model using AIC

selection criteria using R packages *nlme* v3.1 (Pinheiro and Bates 2000; Pinheiro et al. 2023) and

MuMIn v1.47.5 (Bartoń 2018). To test whether foliar herbivory differs between habitats, we used

a generalized linear model with a binomial distribution with herbivory as the dependent variable

and habitat as the fixed independent variable with block nested within site as a random effect. To

understand the effect of herbivory on plant fitness, we used generalized linear models with a

Poisson distribution and fitness (i.e. seed count) as the dependent variable while herbivory and

habitat and their interactions were fixed independent variables with block nested within site as a

random effect using the R package *lme4* v1.1.32 (Bates et al. 2015).

Phenotypic plasticity analyses

We used parental inbred lines to investigate each species' phenotypically plastic response to the

environmental variation between species' habitats within individual years. These spatial

contrasts provide a coarse-grained test (Levins 1968) of how genotypes may respond to differing

environmental conditions. Plasticity was analyzed using generalized linear mixed models with

phenotype as the dependent variable, habitat as a fixed effect and block nested within site as a

random effect. Further, we examined phenotypic plasticity between habitats and years (2013-

2023) using generalized linear mixed models with phenotype as the dependent variable, habitat

and year with their interaction as fixed effects and block nested within site as a random effect.

We also compared hybrid trait means between habitats which are not true measures of plasticity

since they represent both genetic and environmental differences.

Phenotypic selection analyses

To measure lifetime fitness for our phenotypic selection analyses, we counted fruit and seed

number. Plants that produced flowers but did not produce fruit or seed were indicated with a

value of zero. For annual plants, fruit and seed number are a good approximation of fecundity

which we use in our selection analyses. We calculated the correlation between fruit and seed

number using Kendall's correlation analysis. We found that selection gradients (β values)

derived from fruit number (2013) were similar to those from seed number (all other years) in

both strength and direction (Table S4, Table S5) likely due to a positive correlation between seed

and fruit number (Table S6). Therefore, we compare fruit and seed number derived selection

gradients across years.

To understand temporal fluctuation in the strength and direction of selection on traits in each

habitat, we used multivariate phenotypic selection analyses on datasets of hybrid individuals

across all five years of transplants. To investigate linkage between traits, we calculated

correlations between traits from hybrid individuals, grouped separately by habitat and year, using

linear mixed models with traits as dependent and fixed variables with block nested within site as

a random effect. We found generally weak correlations ($r < 0.50$) between traits, except between

plant height & leaf area and leaf area & flower width (Table S3). Therefore, leaf area was

removed from the following phenotypic selection analyses.

To properly compare selection gradients from all years, data published in 2013 and 2019 were

re-analyzed using the same multivariate linear selection model as the recent transplants (Ferris

and Willis, 2018; Tataru *et al*. 2023). Quantitative traits measured in each experiment (leaf

lobing, plant height, outcrossing, flower width, flowering time) were standardized to a normal

distribution with a mean of 0 and standard deviation of 1 to enable comparison of selection on

qualitatively different traits and with previously published studies (Lande & Arnold 1983). Traits

were standardized within each year for all analyses except the combined analysis (described

below). In our phenotypic selection model, seed count (i.e. fecundity) was the dependent variable

with plant height, flowering time, flower width, herkogamy, and leaf lobing as fixed independent

variables, with block nested in site and habitat as a random effect. We did not include quadratic

or correlational selection terms in our models to simplify our interpretation of the results

(Anderson et al. 2015*b*). We did not conduct phenotypic selection analysis of fitness combined

across life history stages using an ASTER model approach because our dataset did not include

trait data on individuals that died before flowering. To account for overdispersion of zeros in our

fecundity data we divided it into two components: the probability of setting any seed

(reproducing) vs. the number of seeds set by plants that fruited (fecundity). We used two types of

generalized linear mixed models using the R package *glmmTMB* v1.1.8 (Brooks et al. 2017;

Magnusson et al. 2017) to analyze these two fitness components separately: the probability of

setting seed with a binomial distribution, and a zero-truncated Poisson distribution to assess

whether trait values were associated with the number of seeds produced. Sites were pooled

within habitats. To extend the comparison to the 2013 experiment which used only fruit number,

we repeated all analyses using fruit number as the dependent variable.

We repeated phenotypic selection analyses on a combined dataset of hybrid genotypes pooled

across all years (2013–2023) using fruit number as the dependent variable with plant height,

flowering time, flower width, and leaf lobing as fixed independent variables, and block nested in

site and year as random effects. Herkogamy was omitted from the combined dataset due to its

absence in 2013. Raw trait data were combined across years and then standardized. To visualize

the fitness-phenotype relationships, we plotted selection gradients using the R package *visreg*

(Breheny and Burchett 2017). To assess whether slopes were statistically different for individual

traits between meadow and granite habitats, we compared slopes of models using a bootstrap

approach (1000 iterations) to derive 95% confidence intervals and *p*-values. To assess whether

temporally fluctuating phenotypic selection is related to environmental fluctuation, we used the

combined dataset to run generalized linear models including interactions between traits and

snowpack. We used the snow water content accumulated (inches) by April 1st of each year

(Figure 1a). The models included fruit as the dependent variable, phenotypic traits (flowering

time, flower width, leaf lobing, plant height) and their interaction with snowpack as fixed

independent variables and then block nested in site and year as random effects.

Assessing local adaptation of M. laciniatus vs. M. guttatus

To assess parental species' local adaptation, we used three metrics of fitness for parental

genotypes: survival to flowering ('survival'), mean seeds per reproductive individual

('fecundity'), and mean seeds per planted individual ('total fitness'). For long term patterns, we

calculated arithmetic means of each metric across years (2013, 2019, 2021, 2022, 2023) for each

habitat and species. Data from 2013 was omitted from averaged fecundity and total fitness

because only fruit number was counted. Because it is not known whether these species have seed

banks, impacting long-term fitness, we also calculated geometric means across years (Figure S5).

To test for genotype by environment (GxE) interactions, we fit linear mixed-effects models for

fecundity and total fitness, and generalized linear mixed-effects models for survival, with

genotype, habitat, and their interaction as fixed effects and block nested within site as a random

effect. Finally, to assess whether the extreme high snowpack of 2023 altered long-term patterns,

we compared mean parental fitness in each habitat with (Figure 5) and without data from 2023

(Figure S4). All models described above are summarized in Table S7.

RESULTS

Environmental variation causes spatial and temporal variation in survival

Fine-scale environmental variation was significantly associated with survival trends in both

habitats across years and fluctuated both spatially and temporally with snowpack (Figure 2,

Table S8). Soil moisture availability throughout the season varies significantly between *M.*

laciniatus's granite and *M. guttatus's* meadow habitat in each year with moisture in granite

outcrops declining rapidly after an early season plateau, while in meadows there is a gradual,

linear decline throughout the growing season. The amount of soil moisture available and the

shape of the soil moisture curve fluctuated from year to year with higher snowpack years (2019,

2023) showing shallower declines in soil moisture over time in each habitat (Figure 2). Light

intensity and temperature were higher on average in *M. laciniatus's* granite environment except

in the extreme high snowpack year of 2023 where temperatures were cooler than usual and

similar between habitats throughout the growing season (Figure 2; Table S8).

Across 2021-2023 in both meadow and granite habitats, time (days; $p < 0.0001$), soil moisture (p

< 0.0001), and soil surface temperature ($p < 0.0001$), were almost always significantly associated

with survival, with decreasing soil moisture and increasing temperature over time associated

with decreased survival (Table S8). Interestingly in 2023, time ($p > 0.05$) was not significant

associated with survival in meadow habitat. This is likely because in this exceptionally high

snowpack year, approximately half of meadow individuals were alive until first snowfall at the

end of the growing season (Figure 2a). In *M. laciniatus's* granite habitat, we found that light

intensity ($p < 0.0001$) also significantly affected survival in 2021-2023. Soil moisture and soil

surface temperature impacted survival over time in the same directions as described for *M.*

guttatus's meadow habitat in all three recent years. In 2023, soil moisture ($p > 0.05$) was not

significantly associated with survival in granite habitat. In summary, we found that soil moisture

and temperature most significantly impacted survival in both habitats and light intensity

additionally impacted survival in granite habitat. Further, 2021–2023 differed from 2019 where

soil moisture was the only significant environmental factor (Tataru *et al.*, 2023).

Similar to 2013 and 2019, herbivory pressure differed significantly between habitats with greater

herbivory in *M. guttatus*'s meadows in 2021 (meadow 22%, granite 1.97%; $p = 0.042$), 2022

(meadow 37.75%, granite 2.58%; $p < 0.01$), and 2023 (meadow 23.17%, granite 8.74%; $p =$

0.032). In 2021 and 2023, herbivory and the interaction between herbivory and habitat

significantly affected total fitness but did not in 2022. Overall, herbivore pressure varied

spatially, but its effect on fitness varied temporally.

Episodic selection: an extreme climatic event dramatically shifted the growing season

The extreme high snowpack year (2023) differed from other years in several aspects. The

growing season was delayed by six weeks or more in both habitats, with surveys not beginning

in granite until previous years' surveys had ended (Figure 2a,b). Soil moisture declined more

gradually in both habitats in 2023 than any previous year (Figure 2c,d), while light intensity and

surface temperatures were lower (Figure 2e-h). In granite, there was no characteristic

temperature spike in 2023 that usually coincided with mortality (Figure 2h). Unlike previous

years, many plants had not senesced in meadows at the end of the growing season: 41.7%

(Meadow 1) and 50.7% (Meadow 2) alive at the last survey before snowfall (Figure 2a). Of the

individuals remaining in Meadow 1, many had not flowered indicating a significant life-history

delay (0% local genotype, 31.0% hybrids, 44.1% *M. guttatus*, 70.7% *M. laciniatus*). Meadow 2

was similar (0% local genotype, 47.3% hybrids, 40.4% *M. guttatus*) except most individuals of

M. laciniatus flowered (92.0%). The 2023 growing season was 36-70 days longer than previous

years in granite habitats (Granite 2013-2022: 33-87 days; 2023: 112-113 days) but comparable to

other years in meadows (Meadow 2013-2022: 74-160 days; 2023: 119-146 days).

Phenotypes are adaptively plastic between habitats

Parental genotypes exhibited consistent phenotypic plasticity between habitats within years, with

the non-native species generally shifting toward the phenotype of the local species. In nearly all

three years, *M. laciniatus* grew more *M. guttatus*-like in meadow habitat with significantly

longer flowering time, taller plant height, larger leaf area (Tables S9, S10, S11). This is similar

to patterns from 2013 except for flower size (Ferris & Willis 2018). Conversely, in its nonnative

granite habitat, *M. guttatus* shifted toward *M. laciniatus*-like traits with significantly shorter

flowering time and smaller plant height (2021 and 2023) and smaller leaf area and increased leaf

lobing (2022 and 2023; Tables S9, S10, S11). Furthermore, flowering time, plant height, flower

width, leaf area displayed significant habitat by year interactions (all $p < 0.000$), indicating

plasticity over time, whereas leaf lobing and outcrossing did not (Table S12). Hybrid means also

differed between habitats with shorter flowering time, smaller plant height, smaller leaf area in

granite habitat, in the direction of the *M. laciniatus* (Tables S9, S10, S11). In 2019, hybrids in

granite differed by having larger flowers and larger leaf area relative to meadow (Tataru et al.

2023). However, we do not interpret hybrid differences as plasticity as they may instead reflect

differences in habitat-dependent selection resulting in differing genetic composition.

Interannual fluctuation versus long-term directional selection

We found temporal and spatial fluctuations in the strength and direction of selection on each

quantitative trait (Tables 1, 2; Figure 3). Our binomial models revealed how phenotypes affected

whether plants reproduced whereas the zero-truncated Poisson models revealed how selection

acted among individuals that did reproduce (number of seeds). Based on observed trait

differences between species, we predicted selection in *M. laciniatus's* harsh granite habitat for

early flowering time, smaller flowers, increased leaf lobing, shorter plants, and decreased stigma-

anther distance (herkogamy), with the inverse direction of selection in *M. guttatus's* meadow

habitat (Figure 3a-e). We found that selection fluctuated across years in habitats and was at times

in the opposite direction of our expectations. Selection on the probability of reproduction

(binomial models) was typically stronger and more consistent in direction with predictions than

selection on fecundity (zero-truncated Poisson models), suggesting that selection acting early in

the life cycle on survival and flowering may be more vital than selection acting on fecundity

(Figure 3).

Across all years, selection favored earlier flowering time in both habitats; with early flowering

time often under stronger negative selection in the granite habitat as expected in the binomial

analysis (Figure 3f,k). Selection on flower width fluctuated but was often in the direction

expected with larger flowers more favorable in meadow than in granite in the binomial analysis

(Figure 3g). Zero-truncated models of fecundity selection on flowering time and flower width

varied but, in most years, there was still stronger selection for early flowering in granite habitats

and stronger selection for larger flowers in meadow as expected (Figure 3k, l). Plants with more

lobed leaves were more likely to reproduce in granite habitat in low snowpack years, aligning

with predicted species differences (Figure 3h). In high snowpack years, the selection was

reversed with more lobed leaves being advantageous in meadow habitat (Figure 3h). Selection on

leaf lobing also fluctuated temporally in fecundity selection, where there was often selection for

more lobed leaves in meadow, but not granite, contrary to predictions. Taller plants were

favorable across habitats and years but contrary to predictions there was often stronger positive

selection for height in granite (Figure 3i,n). For herkogamy, we found selection in the direction

predicted in high snowpack years (2019, 2023), with shorter stigma-anther separation associated

with higher fecundity in granite and greater stigma-anther separation favorable in meadow, but

reversed in the low snowpack year 2021 (Figure 3o). Phenotypic selection analyses where

models excluded herkogamy in all years were largely similar (Figure S2). We found that these

temporal fluctuations in phenotypic selection were statistically associated with variation in

snowpack. In meadow habitat, there was a significant plant height by snowpack interaction in the

zero-truncated Poisson model ($p < 0.0001$), and snowpack significantly interacted with both

plant height and leaf lobing in the binomial analysis (both $p < 0.05$). In granite habitat, flowering

time and flower width both had significant interactions with snowpack in the zero-truncated

Poisson (both $p < 0.001$) while there was a plant height by snowpack interaction in the binomial

model ($p < 0.0001$).

Although we found interannual fluctuations in the strength and direction of selection on all traits,

when we combined data across years, we saw long term patterns of directional selection (Table

1, Table 2; Figure 4, Figure S3). In both zero-truncated Poisson and binomial analyses, we found

long-term selection for earlier flowering time, smaller flowers, more lobed leaves and taller

plants in both habitats. Selection was significantly stronger in the granite habitat for early

flowering time (binomial: $p < 0.0001$), lobed leaf shape (Poisson: $p < 0.0001$), and larger plant

sizes (Poisson: $p < 0.0001$). Therefore, while inter-annual fluctuations in the strength and

direction of selection are common in this system, we still find longer-term differences in the

strength of selection in the direction of species' differences in both flowering time and leaf

shape.

Local adaptation fluctuates temporally with long-term maladaptation of M. guttatus

We found temporal fluctuation in the signature of local adaptation in both habitats. In 2013, a

drought year, we found a clear signature of local adaptation in total fitness, but this was not true

for other years in our study (Figure 5). There was also often a difference in local adaptation

between the individual components of fitness: survival and fecundity. *Mimulus laciniatus* had

higher survival to flowering than *M. guttatus* in both habitats in all years, except in 2019 when

there was low overall survival of parental genotypes (Figure 5a-e). With all five years combined,

M. laciniatus had higher average survival in both habitats (Figure 5f). Patterns of fecundity

differed between years with parental species having higher fecundity in their native habitats in

2021, like in 2013, indicating local adaptation and selection against immigrants (Figure 5g-k).

However, in 2022, *M. laciniatus* had higher fecundity in both habitats (Figure 5j), and in 2023,

we found reciprocal local maladaptation where *M. guttatus* had higher fecundity than *M.*

laciniatus in granite and vice versa in meadows (Figure 5k).

With all years combined, we found overall evidence of local maladaptation in mean fecundity

(Figure 5l); however, removing the extreme snowpack year of 2023 reveals local adaptation in

fecundity (Figure S4). A significant genotype by environment interaction GxE was detected for

total fitness in every year except 2019, and for survival and total fitness when all years were

combined (Table S13). Detection of GxE in survival and fecundity varied more across individual

years (Table S13). For total fitness, *M. laciniatus* had higher fitness and home-site advantage in

low snowpack years (2013, 2021, 2022; Figure 5m,o,d), but *M. laciniatus* and *M. guttatus*

performed equally well in granite during high snowpack years (2019, 2023; Figure 5n,q). Total

fitness in meadow was not as predictable based on snowpack with *M. laciniatus* having higher

total fitness in recent years (2021-2023) but lower fitness in 2013 and 2019. Overall, *M.*

laciniatus had higher total fitness in both habitats (Figure 5r, Figure S4). Geometric mean

fitness analyses reinforced these patterns. A single year of very low or zero fitness would drive

extinction in the granite habitat in the absence of a seed bank (Figure S5), and *M. laciniatus*

would still maintain higher geometric-mean total fitness than the native *M. guttatus* in meadow

habitat (Figure S5).

DISCUSSION

In this study, we investigated spatial and temporal variation in selection on two sympatric

Monkeyflower species, *M. guttatus* and *M. laciniatus*. We compared multiple large-scale

replicated reciprocal transplants conducted in low (2013, 2021, 2022) and high (2019, 2023)

snowpack years including one extreme year resulting in episodic selection (Figure 1a, b). We

found that key environmental variables such as soil moisture, temperature, and light intensity

fluctuated over space and time, impacting fitness and phenotypic plasticity. Using experimental

hybrids, we found fluctuating selection on phenotypic traits involved in reproductive isolation

and microhabitat adaptation. Within each year, phenotypic selection was sometimes contrary to

our predicted directions, but we saw a longer-term trend of differences in the magnitude of

selection in the direction of species' differences in both flowering time, a key reproductive

isolating trait, and leaf shape. Our results also indicate that local adaptation of the parental

species fluctuates across years but on average, *M. guttatus* is maladapted to its meadows and that

habitat isolation may therefore be asymmetric and temporally dependent. Finally, we find that

episodic selection, caused by an extreme snowfall event, may disrupt local adaptation and impact

species boundaries.

Inter-annual fluctuation in natural selection does not erase signatures of long-term divergence

Temporal and spatial variation in selection has been well studied for its role in maintaining

genetic variation within species (Keep et al. 2021; Leidinger et al. 2021; Kelly 2022; Johnson et

al. 2023). However, few studies have empirically tested how fluctuating selection influences the

strength of reproductive isolation between sympatric species (but see Campbell and Powers

2015). Here, we consider this by comparing the relative strengths and direction of phenotypic

selection on experimental hybrids between native *Mimulus* habitats and examine how these

patterns changed over five repeated transplant experiments.

Although there were interannual fluctuations in selection (Figure 3), we detected interesting

longer-term patterns aligned with species differences, particularly for leaf lobing and flowering

time, a key pre-zygotic isolating barrier in the *M. guttatus* species complex (Lowry et al. 2008;

Figure 4). Previous studies have found that later-flowering *M. guttatus* tend to exhibit higher

fecundity (Mojica and Kelly 2010; Mojica et al. 2012), especially in a longer, high snowpack

growing season with lowered desiccation risk. In our study, flowering time selection was

negative in both habitats, yet in three out of four years with comparable phenotypic selection

data between habitats, there was stronger selection for earlier flowering time in *M. laciniatus'*

granite habitat as predicted by species differences (Figure 3k). The high snowpack year of 2019

was the exception with stronger fecundity selection for early flowering in *M. guttatus'* meadows

than in granite habitat (Figure 3k), indicating a temporary relaxation of divergent selection.

Selection on leaf lobing also fluctuated across years, where selection favored the direction of

species differences in our binomial analyses in low snowpack years whereas in high snowpack

years the direction reversed to be opposite our predictions (Figure 3h). This pattern of the

direction of selection on leaf shape switching with snowpack is supported by the significant

interaction between leaf lobing and snowpack in our across-year model (Table S7). The precise

adaptive function of leaf shape variation in this system is not known but may be related to

temperature or drought tolerance as leaf shape affects the leaf boundary layer and the rates of

heat and gas exchange with the air (Nicotra et al. 2011). Two other rocky outcrop species in the

M. guttatus complex have also evolved lobed leaves, *M. filicifolius* and some populations of

serpentine adapted *M. guttatus* (Ferris et al. 2015). Further, repeated altitudinal clines in leaf

shape have been found in *M. laciniatus* populations (Love and Ferris 2023). These repeated

evolutionary patterns, along with evidence of selection for lobed leaves in granite in some years,

suggest that lobed leaf shape is indeed adaptive in these species, even though the direction of

selection is not stable across time.

Annual selection on flower width fluctuated unpredictably across years and our combined

analysis revealed weak selection for smaller flowers (Figure 3, Figure 4, Table 1, Table 2). This

runs counter to species differences as *M. guttatus* typically has larger flowers than *M. laciniatus*.

Flower size affects reproductive success and can contribute to population differentiation and

reproductive isolation (Bradshaw et al. 1995; Brunet 2009; Schiestl and Schluter 2009; Venail et

al. 2010; Krizek and Anderson 2013). Flower size is often correlated with plant size indicating

that early flowering, small stature, and small flowers are linked (Troth et al. 2018). Previous

work has demonstrated antagonistic pleiotropy in *M. guttatus* where later flowering alleles also

increase plant and flower size (Mojica et al. 2012; Monnahan and Kelly 2015). Although plants

with larger flowers have increased fecundity, they simultaneously reduce viability, resulting in

net selection for smaller flowers (Mojica and Kelly 2010). In contrast, our findings of selection

for early flowering, small flowers, and larger plants in both habitats across years suggests that in

our species' these traits are genetically distinct (Tables 1, 2). Although we initially predicted

selection for smaller plant sizes in granite based on *M. laciniatus's* short stature, our findings

suggest that larger plants always have higher fecundity. This is in line with broader evidence that

body size at reproduction is a key determinant of fecundity across taxa (Aarseen and Taylor

1992; Aarssen and Jordan 2001; Kingsolver and Pfennig 2004). Despite selection for larger

sizes, average plant height remained smaller in granite than in meadow habitats (Table S9, Table

S10, Table S11), likely reflecting ecological constraints imposed by limited water and nutrient

availability in rocky outcrops. Finally, we found weak selection on herkogamy, suggesting that

selection on stigma-anther separation is no longer as strong as during the earlier stages of

speciation (Harder and Johnson 2009).

Across these traits, our combined dataset revealed longer-term differences in patterns of

selection on flowering time, plant size, and leaf shape between habitats (Table 1, Table 2; Figure

4, Figure S3). We detected stronger selection for early flowering time in granite than meadow

habitat which aligns with our predictions based on species' differences, likely contributing to

temporal reproductive isolation as seen in other *Mimulus* species (Hall and Willis 2006; Lowry et

al 2008). This pattern of spatial variation in selection is consistent with findings from other

systems; for example, Ågren et al. (2017) demonstrated that selection on flowering time varies

temporally in Italian and Swedish *A. thaliana* populations but that overall, there is stronger

selection for earlier flowering in Italy. Further, we found a distinct signature of differential

selection on leaf lobing in predicted directions with positive selection for lobing in the granite

outcrops, but nearly no selection on leaf shape in *M. guttatus*'s meadow habitat. Lobed leaf

shape is thought to be involved in adaptation to dry, exposed habitats like *M. laciniatus's* rocky

outcrop environment (Nicotra et al. 2011; Ferris et al 2015; Ferris 2019) and therefore appears to

be involved in habitat isolation between these species (Ferris and Willis 2018; Tataru et al.

2023). Taken together, the signatures of divergent selection on flowering time and leaf shape

across multiple transplants suggests ongoing adaptive differentiation and that fluctuating

selection has not erased long-term maintenance of species boundaries (Chapurlat et al. 2020).

M. guttatus is locally maladapted

Organisms are expected to have a fitness advantage over other genotypes in their home

environment because spatial heterogeneity should favor the evolution of local adaptation

(Hastings 1983). However, temporal changes in environmental conditions are theorized to

constrain adaptation if there are opposing selection pressures over time (Stearns 1992) which

may negatively impact reproductive isolation and species boundaries. We predicted that if

habitat isolation is an important reproductive isolating barrier between our sympatric *Mimulus*,

then there would be a fitness advantage of the local species in each habitat. Instead, we found

that local adaptation fluctuated temporally (Figure 5m-q) with evidence of average local

maladaptation of *M. guttatus* (Figure 5r). Annual fluctuations in snowpack were predictive of

relative fitness of the parent species in granite with *M. laciniatus* and *M. gutta*tus performing

equally well in high snowpack years (5n,q), but in low snowpack years the drought adapted *M.*

laciniatus had the home-site advantage (Figure 5m,o,p).

Although local adaptation is thought to be ubiquitous in natural populations (Hereford 2009),

fitness trade-offs are not always found in reciprocal transplants (Bennett and Lenski 2007;

Lowry et al. 2009). For example, in his classic metanalysis Hereford (2009) found evidence of

weak trade-offs associated with adaptation of populations suggesting that local adaptation is

often not costly. It is possible that our reciprocal transplant did not detect existing trade-offs due

to logistical constraints or that there are weak to no fitness trade-offs associated with adaptation

for *M. laciniatus* across environments (Fry, 1996; Leimu & Fischer, 2008). It is also possible that

M. guttatus lines used in the transplant were negatively affected by inbreeding depression, as a

largely outcrossing species. However, this is unlikely as intrapopulation crosses and inbred lines

performed similarly in the 2023 experiment (Table S3). Alternatively, an increase in dramatic

interannual environmental fluctuations due to anthropogenic climate change may mean that *M.*

guttatus is no longer at its adaptive optima. Our results could indicate an adaptive lag where, due

to recent rapid climate change and increasing drought, the adaptive landscape of *M. guttatus's*

habitat has changed to be more *M. laciniatus*-like and *M. guttatus*'s evolutionary response has

not kept pace (Lane et al. 2012; Mills et al. 2013; Kooyers et al. 2019). This pattern of adaptive

lag and local maladaptation has been found in other populations of *M. guttatus* (Kooyers et al.

2019). Over time, this could erode species boundaries by allowing colonization of *M. guttatus's*

meadow by *M. laciniatus* potentially increasing hybridization or lead to population extinction.

Asymmetry in local adaptation, or a lack of trade-offs, has been found in other transplant

experiments (Hereford 2009; Gosden et al. 2015; Latreille and Pichot 2017; Toll and Willis

2018). In a repeated reciprocal transplant between serpentine and sandstone adapted populations

of *Leptosiphon parviflorus,* Dittmar & Schemske (2023) found that while serpentine populations

had a temporally consistent local fitness advantage, sandstone populations only had a local

advantage in two out of four years. While stable spatial heterogeneity should favor the evolution

of local adaptation, temporal fluctuation in the environment should select against local

specialization and for phenotypic plasticity (Kawecki & Ebert 2004). The average fitness

advantage of *M. laciniatus* across environments could be due to adaptive plasticity, specifically

adaptation to a history of environmental heterogeneity in the rocky outcrop habitat (Ghalambor

et al. 2015). We found that both *M. guttatus* and *M. laciniatus* exhibit plasticity in the direction

of the local species' phenotype in each habitat (Table S9, Table S10, Table S11). *Mimulus*

laciniatus's occupation of a harsh, highly variable environment (Figure 2) may have driven the

evolution of adaptive plasticity (Via and Lande 1985) allowing it to thrive in both habitats.

However, *M. guttatus* was more plastic than *M. laciniatus* in all phenotypes except fitness

indicating that plasticity is unlikely to account for *M. laciniatus's* performance advantage across

habitats and the lack of fitness trade-offs.

Episodic selection weakens species divergence

The high snowpack of 2023 delayed the onset of spring in mid to high elevations by 4–6 weeks

and altered soil moisture, light intensity, and surface temperature regimes (Figure 2). This shift

contributed to local maladaptation between parental species (Figure 5j), as signatures of local

adaptation in fecundity reappeared when 2023 was excluded (Figure S2). In *M. guttatus's*

meadow habitat, the delay was detrimental to hybrids and *M. guttatus* by delaying flowering

phenology and reducing fitness. Additionally, no local meadow genotypes flowered before

snowfall, suggesting negative impacts on native populations. However, many individuals

remained alive at the first snowfall indicating potential facultative perenniality. Delayed

flowering may have resulted from shorter day lengths failing to meet critical photoperiod

thresholds (Friedman and Willis 2013; Fishman et al. 2014; Kenney and Sweigart 2016), the

absence of end-of-season drought stress due to persistently high soil moisture (Kooyers et al.

2015; Mantel and Sweigart 2019), and unusually low light intensity (Figure 2). In contrast, *M.*

laciniatus was less affected*,* likely because it flowers rapidly under shorter daylengths (Friedman

and Willis 2013; Ferris and Willis 2018; Love and Ferris 2024).

In granite habitats, the delayed season in 2023 fostered unusually favorable environments for the

non-native *M. guttatus* with high water availability, lower light intensity and surface

temperatures, and a longer growing season than is typical in granite (Figure 2). This led to higher

M. guttatus survival in granite (12% of total planted; 24 individuals) compared to previous years

(0-2% of total planted). Although *M. guttatus* still had lower survival than *M. laciniatus*, it had

remarkably high fecundity (Figure 5e,k) producing total fitness comparable to *M. laciniatus* for

the first time (Figure 5q). Unlike the meadow habitat, all plants had senesced and many set seed

before snowfall. This suggests that drought stress, rather than photoperiod, is more important as

a flowering cue in *M. lacinaitus's* habitat. High-elevation populations of *M. laciniatus* may also

possess increased plasticity in critical photoperiod (Love and Ferris 2024).

Our results demonstrate how extreme climatic events may disrupt species boundaries. The

atypical conditions of 2023 produced unusually strong selection opposite of predicted directions

on key traits involved in adaptation and reproductive isolation: flowering time, flower size and

leaf lobing (Figure 3). In the long term, such episodic selection may shift parental ranges,

promote hybrid swarms or a new hybrid species (Grant and Grant 1993), or alter adaptive

potential (Campbell-Staton et al. 2017). The rarity and unpredictability of extreme climatic

events complicate predictions about the evolutionary response to climate change. More

longitudinal field studies are needed to understand long-term evolutionary consequences of

extreme climatic events and changing climatic patterns on plant species on their native ranges

(Anderson 2016; Bailey and van de Pol 2016). For instance, do these events drive evolutionary

change and does this outweigh selection acting during 'normal' periods? For species with limited

ranges, such as the Sierra Nevada endemic *M. laciniatus*, will these events influence extinction

risk of populations?

Conclusions

Our repeated reciprocal transplants reveal how fluctuating selection influences species'

boundaries. The foreign advantage of *M. laciniatus* in the non-native meadow habitat across

years suggests either low adaptation costs to the granite habitat or recent maladaptation of *M.*

guttatus due to environmental shifts. Furthermore, longer-term divergent selection on flowering

time and leaf shape should reinforce temporal and habitat isolation between these species.

However, the lack of consistent divergent selection on other key traits could lead to eventual

species fusion. Finally, a bout of episodic natural selection due to extreme snowfall shifted

population dynamics with potential long-term impacts on species' range, abundance, and gene

flow. Our study illustrates the strength of integrating field experiments across multiple spatial

and temporal scales to gain a deeper understanding of the maintenance of biodiversity under

environmental change (Wadgymar et al. 2017; Dittmar and Schemske 2023; Oakley et al. 2023).

COMPETING INTERESTS

None declared.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study will be openly available in [repository name] at

http://doi.org/[doi], reference number [reference number], upon acceptance of the manuscript.

REFERENCES

Aarseen, L. W., and D. R. Taylor. 1992. Fecundity allocation in herbaceous plants. Oikos
65:225–232.
Aarssen, L. W., and C. Y. Jordan. 2001. Between-species patterns of covariation in plant size,
seed size, and fecundity in monocarpic herbs. Ecoscience 8:471–477.
Ågren, J., C. G. Oakley, S. Lundemo, and D. W. Schemske. 2017. Adaptive divergence in
flowering time among natural populations of *Arabidopsis thaliana* : Estimates of selection and
QTL mapping. Evolution 71:550–564.
Anderson, J. T. 2016. Plant fitness in a rapidly changing world. New Phytologist 210:81–87.
Anderson, J. T., V. M. Eckhart, and M. A. Geber. 2015*a*. Experimental studies of adaptation in
Clarkia xantiana: III: Phenotypic selection across a subspecies border. Evolution 69:2249–2261.

———. 2015*b*. Experimental studies of adaptation in Clarkia xantiana: III: Phenotypic selection across a subspecies border. Evolution 69:2249–2261.

Anderson, J. T., and S. M. Wadgymar. 2020. Climate change disrupts local adaptation and favours upslope migration. Ecology Letters 23:181–192.

Anderson, J. T., J. H. Willis, and T. Mitchell-Olds. 2011. Evolutionary genetics of plant adaptation. Trends in Genetics 27:258–266.

Awadalla, P., and K. Ritland. 1997. Microsatellite variation and evolution in the Mimulus guttatus species complex with contrasting mating systems. Molecular Biology and Evolution 14:1023–1034.

Bailey, L. D., and M. van de Pol. 2016. Tackling extremes: Challenges for ecological and evolutionary research on extreme climatic events. Journal of Animal Ecology 85:85–96.

Bartoń, K. 2018. MuMIn: Multi-Model Inference. R package version 1.42.1. https://CRAN.R-project.org/package=MuMIn.

Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67:1–48.

Bennett, A. F., and R. E. Lenski. 2007. An experimental test of evolutionary trade-offs during temperature adaptation. Proceedings of the National Academy of Sciences of the United States of America 104:8649–8654.

Bradshaw, H. D., S. M. Wilbert, K. G. Otto, and D. W. Schemske. 1995. Genetic mapping of floral traits associated with reproductive isolation in monkeyflowers (Mimulus). Nature 376:762–765.

Breheny, P., and W. Burchett. 2017. Visualization of Regression Models Using visreg. The R Journal 9:56–71.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J. Skaug, et al. 2017. Modeling Zero-Inflated Count Data With glmmTMB. bioRxiv preprint. bioRxiv.

Brunet, J. 2009. Pollinators of the Rocky Mountain columbine: Temporal variation, functional groups and associations with floral traits. Annals of Botany 103:1567–1578.

Campbell, D. R., and J. M. Powers. 2015. Natural selection on floral morphology can be influenced by climate. Proceedings of the Royal Society B: Biological Sciences 282.

Campbell-Staton, S. C., Z. A. Cheviron, N. Rochette, J. Catchen, J. B. Losos, and S. V. Edwards. 2017. Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard. Science 357:495–498.

Chapurlat, E., I. Le Roncé, J. Ågren, and N. Sletvold. 2020. Divergent selection on flowering phenology but not on floral morphology between two closely related orchids. Ecology and Evolution 10:5737–5747.

Coltman, D. W., J. A. Smith, D. R. Bancroft, J. Pilkington, A. D. C. MacColl, T. H. Clutton-Brock, and J. M. Pemberton. 1999. Density-dependent variation in lifetime breeding success and natural and sexual selection in Soay rams. American Naturalist 154:730–746.

Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinauer Associates. Sinauer Associates, Sunderland, Massachusetts.

Crespi, B. J. 2000. The evolution of maladaptation. Heredity 84:623–629.

DeMarche, M. L., K. J. Rice, and J. P. Sexton. 2013. Niche partitioning between close relatives suggests trade-offs between adaptation to local environments and competition. Ecology and Evolution 3:512–522.

Dittmar, E. L., and D. W. Schemske. 2023. Temporal Variation in Selection Influences
Microgeographic Local Adaptation. American Naturalist 202:471–485.
Ferris, K. G. 2019. Endless forms most functional: uncovering the role of natural selection in the
evolution of leaf shape. American Journal of Botany 106:1532–1535.
Ferris, K. G., L. L. Barnett, B. K. Blackman, and J. H. Willis. 2017*a*. The genetic architecture of
local adaptation and reproductive isolation in sympatry within the Mimulus guttatus species
complex. Molecular Ecology 26:208–224.
———. 2017*b*. The genetic architecture of local adaptation and reproductive isolation in
sympatry within the Mimulus guttatus species complex. Molecular Ecology 26:208–224.
Ferris, K. G., T. Rushton, A. B. Greenlee, K. Toll, B. K. Blackman, and J. H. Willis. 2015. Leaf
shape evolution has a similar genetic architecture in three edaphic specialists within the Mimulus
guttatus species complex. Annals of Botany 116:213–223.
Ferris, K. G., J. P. Sexton, and J. H. Willis. 2014. Speciation on a local geographic scale: The
evolution of a rare rock outcrop specialist in Mimulus. Philosophical Transactions of the Royal
Society B: Biological Sciences 369:27–29.
Ferris, K. G., and J. H. Willis. 2018. Differential adaptation to a harsh granite outcrop habitat
between sympatric Mimulus species. Evolution 72:1225–1241.
Fishman, L., A. L. Sweigart, A. M. Kenney, and S. Campbell. 2014. Major quantitative trait loci
control divergence in critical photoperiod for flowering between selfing and outcrossing species
of monkeyflower (Mimulus). New Phytologist 201:1498–1507.
Friedman, J., and J. H. Willis. 2013. Major QTLs for critical photoperiod and vernalization
underlie extensive variation in flowering in the Mimulus guttatus species complex. New
Phytologist 199:571–583.
Fry, J. D. 1996. The Evolution of Host Specialization: Are Trade-Offs Overrated? The American
Naturalist 148:S84–S107.
Galen, C. 2000. High and dry: Drought stress, sex-allocation trade-offs, and selection on flower
size in the alpine wildflower Polemonium viscosum (Polemoniaceae). American Naturalist
156:72–83.
Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, and K. A. Hughes.
2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature.
Nature 525:372–375.
Gosden, T. P., J. T. Waller, and E. I. Svensson. 2015. Asymmetric isolating barriers between
different microclimatic environments caused by low immigrant survival. Proceedings of the
Royal Society B: Biological Sciences 282.
Grant, B. R., and P. R. Grant. 1993. Evolution of Darwin's Finches Caused by a Rare Climatic
Event. Proceedings of the Royal Society B: Biological Sciences 251:111–117.
Groh, J. S., and G. Coop. 2024. The temporal and genomic scale of selection following
hybridization. Proceedings of the National Academy of Sciences of the United States of America
121:1–11.
Hall, M. C., and J. H. Willis. 2006. Divergent Selection on Flowering Time Contributes To
Local Adaptation in Mimulus Guttatus Populations. Evolution 60:2466.
Harder, L. D., and S. D. Johnson. 2009. Darwin's beautiful contrivances: evolutionary and
functional evidence for floral adaptation. New Phytologist 183:530–545.
Hastings, A. 1983. Can spatial variation alone lead to selection for dispersal? Theoretical
Population Biology 24:244–251.

Hereford, J. 2009. A quantitative survey of local adaptation and fitness trade-offs. American Naturalist 173:579–588.

Johnson, O. L., R. Tobler, J. M. Schmidt, and C. D. Huber. 2023. Fluctuating selection and the determinants of genetic variation. Trends in Genetics 39:491–504.

Kawecki, T. J., and D. Ebert. 2004. Conceptual issues in local adaptation. Ecology Letters 7:1225–1241.

Keep, T., S. Rouet, J. L. Blanco-Pastor, P. Barre, T. Ruttink, K. J. Dehmer, M. Hegarty, et al. 2021. Inter-annual and spatial climatic variability have led to a balance between local fluctuating selection and wide-range directional selection in a perennial grass species. Annals of Botany 128:357–369.

Kelly, J. K. 2022. The genomic scale of fluctuating selection in a natural plant population. Evolution Letters 6:506–521.

Kenney, A. M., and A. L. Sweigart. 2016. Reproductive isolation and introgression between sympatric Mimulus species. Molecular ecology 25:2499–2517.

Kingsolver, J. G., and S. E. Diamond. 2011. Phenotypic selection in natural populations: What limits directional selection? American Naturalist 177:346–357.

Kingsolver, J. G., and D. W. Pfennig. 2004. Individual-level selection as a cause of cope's rule of phyletic size increase. Evolution 58:1608–1612.

Kooyers, N. J., J. M. Colicchio, A. B. Greenlee, E. Patterson, N. T. Handloser, and B. K. Blackman. 2019. Lagging adaptation to climate supersedes local adaptation to herbivory in an annual monkeyflower. American Naturalist 194:541–557.

Kooyers, N. J., A. B. Greenlee, J. M. Colicchio, M. Oh, and B. K. Blackman. 2015. Replicate altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual Mimulus guttatus. New Phytologist 206:152–165.

Krizek, B. A., and J. T. Anderson. 2013. Control of flower size. Journal of Experimental Botany 64:1427–1437.

Lande, R., and S. J. Arnold. 1983. The Measurement of Selection on Correlated Characters. Evolution 37:1210–1226.

Lane, J. E., L. E. B. Kruuk, A. Charmantier, J. O. Murie, and F. S. Dobson. 2012. Delayed phenology and reduced fitness associated with climate change in a wild hibernator. Nature 489:554–557.

Latreille, A. C., and C. Pichot. 2017. Local-scale diversity and adaptation along elevational gradients assessed by reciprocal transplant experiments: lack of local adaptation in silver fir populations. Annals of Forest Science 74.

Lee, G., B. J. Sanderson, T. J. Ellis, B. P. Dilkes, J. K. McKay, J. Ågren, and C. G. Oakley. 2024. A large-effect fitness trade-off across environments is explained by a single mutation affecting cold acclimation. Proceedings of the National Academy of Sciences 121.

Leidinger, L., D. Vedder, and J. S. Cabral. 2021. Temporal environmental variation may impose differential selection on both genomic and ecological traits. Oikos 130:1100–1115.

Leimu, R., and M. Fischer. 2008. A meta-analysis of local adaptation in plants. PLoS ONE 3:1–8.

Levins, R. 1968. Evolution in Changing Environments. Princeton University Press, Princeton, New Jersey.

Love, J. M., and K. G. Ferris. 2024. Local adaptation to an altitudinal gradient: The interplay between mean phenotypic trait variation and phenotypic plasticity in Mimulus laciniatus. Perspectives in Plant Ecology, Evolution and Systematics 63:125795.

Lowry, D. B., M. C. Hall, D. E. Salt, and J. H. Willis. 2009. Genetic and physiological basis of
adaptive salt tolerance divergence between coastal and inland Mimulus guttatus. New
Phytologist 183:776–788.
Lowry, D. B., R. C. Rockwood, and J. H. Willis. 2008. Ecological reproductive isolation of coast
and inland races of Mimulus guttatus. Evolution 62:2196–2214.
Magnusson, A., H. J. Skaug, A. Nielsen, C. W. Berg, K. Kristensen, M. Maechler, K. J. van
Bentham, et al. 2017. glmmTMB: Generalized Linear Mixed Models using Template Model
Builder. R package version 0.1.3.
Mantel, S. J., and A. L. Sweigart. 2019. Divergence in drought-response traits between sympatric
species of Mimulus. Ecology and Evolution 9:10291–10304.
Mills, L. S., M. Zimova, J. Oyler, S. Running, J. T. Abatzoglou, and P. M. Lukacs. 2013.
Camouflage mismatch in seasonal coat color due to decreased snow duration. Proceedings of the
National Academy of Sciences of the United States of America 110:7360–7365.
Mojica, J. P., and J. K. Kelly. 2010. Viability selection prior to trait expression is an essential
component of natural selection. Proceedings of the Royal Society B: Biological Sciences
277:2945–2950.
Mojica, J. P., Y. W. Lee, J. H. Willis, and J. K. Kelly. 2012. Spatially and temporally varying
selection on intrapopulation quantitative trait loci for a life history trade-off in Mimulus guttatus.
Molecular Ecology 21:3718–3728.
Monnahan, P. J., and J. K. Kelly. 2015. Naturally segregating loci exhibit epistasis for fitness.
Biology Letters 11.
Nagy, E. S. 1997. Selection for native characters in hybrids between two locally adapted plant
subspecies. Evolution 51:1469–1480.
Nicotra, A. B., A. Leigh, C. K. Boyce, C. S. Jones, K. J. Niklas, D. L. Royer, and H. Tsukaya.
2011*a*. The evolution and functional significance of leaf shape in the angiosperms. Functional
Plant Biology 38:535–552.
———. 2011*b*. The evolution and functional significance of leaf shape in the angiosperms.
Functional Plant Biology 38:535–552.
Oakley, C. G., D. W. Schemske, J. K. McKay, and J. Ågren. 2023. Ecological genetics of local
adaptation in Arabidopsis: An 8-year field experiment. Molecular Ecology 32:4570–4583.
Pemberton, J. M., L. E. B. Kruuk, and T. Clutton-Brock. 2022. The Unusual Value of Long-
Term Studies of Individuals: The Example of the Isle of Rum Red Deer Project. Annual Review
of Ecology, Evolution, and Systematics 53:327–351.
Pinheiro, J., and D. Bates. 2000. Mixed-Effects Models in S and S-PLUS. Springer, ew York.
Pinheiro, J., D. Bates, and R Core Team. 2023. nlme: Linear and Nonlinear Mixed Effects
Models. R package version 3.1-163.
Ramsey, J., H. D. Bradshaw, and D. W. Schemske. 2003. Components of reproductive isolation
between the monkeyflowers Mimulus lewisii and M. cardinalis (Phrymaceae). Evolution
57:1520–1534.
Rundle, H. D., and P. Nosil. 2005. Ecological speciation. Ecology Letters 8:336–352.
Schiestl, F. P., and P. M. Schluter. 2009. Floral isolation, specialized pollination, and pollinator
behavior in orchids. Annual Review of Entomology 54:425–446.
Schluter, D. 2009. Evidence for ecological speciation and its alternative. Science 323:737–741.
Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of
Image Analysis. Nature Methods 9:671–675.

Siepielski, A. M., J. D. Dibattista, and S. M. Carlson. 2009. It's about time: The temporal
dynamics of phenotypic selection in the wild. Ecology Letters 12:1261–1276.
Stearns, S. 1992. The evolution of life histories. Oxford University Press, Oxford, UK.
Tataru, D., M. De Leon, S. Dutton, F. Machado-Perez, A. Rendahl, and K. G. Ferris. 2024.
Fluctuating selection in a Monkeyflower hybrid zone. bioRxiv 1–33.
Tataru, D., E. C. Wheeler, and K. G. Ferris. 2023. Spatially and temporally varying selection
influence species boundaries in two sympatric Mimulus. Proceedings of the Royal Society B:
Biological Sciences 290.
Toll, K., and J. H. Willis. 2018. Hybrid inviability and differential submergence tolerance drive
habitat segregation between two congeneric monkeyflowers. Ecology 99:2776–2786.
Troth, A., J. R. Puzey, R. S. Kim, J. H. Willis, and J. K. Kelly. 2018. Selective trade-offs
maintain alleles underpinning complex trait variation in plants. Science 361:475–478.
Venail, J., A. Dell'Olivo, and C. Kuhlemeier. 2010. Speciation genes in the genus Petunia.
Philosophical Transactions of the Royal Society B: Biological Sciences 365:461–468.
Via, S., and R. Lande. 1985. Genotype-environment interaction and the evolution of phenotypic
plasticity. Evolution 39:505–522.
Vickery, R. K. 1964. Barriers to gene exchange between members of the Mimulus guttatus
complex (Scrophulariaceae). Evolution 18:52.
Wadgymar, S. M., S. C. Daws, and J. T. Anderson. 2017. Integrating viability and fecundity
selection to illuminate the adaptive nature of genetic clines. Evolution Letters 1:26–39.
Wilczek, A. M., M. D. Cooper, T. M. Korves, and J. Schmitt. 2014. Lagging adaptation to
warming climate in Arabidopsis thaliana. Proceedings of the National Academy of Sciences of
the United States of America 111:7906–7913.
Willis, J. 1999. The role of genes of large effect on inbreeding depression in Mimulus guttatus.
Evolution 53:1678–1691.
Wright, S. I., S. Kalisz, and T. Slotte. 2013. Evolutionary consequences of self-fertilization in
plants. Proceedings of the Royal Society B: Biological Sciences 280.
Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008.
Mimulus is an emerging model system for the integration of ecological and genomic studies.
Heredity 100:220–230.
888
889

Figure 1. Temporal and spatial scale of the experimental design. Annual changes in a) snow

water content and b) accumulated precipitation during experimental years of low snowpack

(2013, 2021, 2022) and comparatively higher snowpack (2019 and 2023). Data are averaged

from the California Department of Water Resources data stations GIN, WHW, and TUM, located

nearby experimental sites along Tioga Road in Yosemite National Park (cdec.water.ca.gov).

The c) spatial layout of experimental sites: meadow 1 and 2 (M1 and M2; filled circles) and

granite 1 and 2 (G1 and G2; open circles). Nearby parental source populations for experimental

crosses are shown: *M. guttatus* YVO (filled square) and *M. laciniatus* WLF (open triangle).

Photos of representative meadow and granite habitats are shown in the insets.

Figure 2. Seasonal changes in (a,b) survival (%), (c,d) soil moisture (%), (e,f) light intensity (μmol m-2 s-1), and (g,h) soil surface temperature (°F) in each block per site in meadow and granite habitats in low snowpack years (2013, 2021, 2022) and high snowpack years (2019, 2023). Data from 2013 and 2019 are published in Ferris and Willis 2018 and Tataru *et al*. 2023 and re-analyzed here.

Figure 3. Visual representation of selection gradients (β) of phenotypic traits measured for

hybrids during experimental years in granite (open circles) and meadow (filled circles) habitats.

Shown are (a-e) predicted values in each habitat based on native species trait differences and

estimated values from (f-j) binomial models and (k-o) zero-truncated Poisson models.

Significance of each β value is indicated with asterisks (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$)

and negative graph areas are shaded grey for clarity. All β values are based on seed number,

except for values from 2013 which are based on fruit number, as seed number was not recorded.

Data from 2013 and 2019 are published in Ferris and Willis 2018 and Tataru *et al*. 2023 and re-

analyzed here.

Figure 4. Patterns of cumulative selection with all years combined in meadow (left) and granite

(right) habitat in a zero-truncated Poisson analysis on (a,b) flowering time (days from planting),

(c,d) width of first flower (mm), (e,f) leaf lobing index, and (g,h) plant height at flowering (mm

to apical meristem). Y-axes show adjusted relative fitness (fruit number), statistically corrected

for other variables included in the models, and the x-axes show unstandardized trait values for

biological relevance. Data from 2013 and 2019 are published in Ferris and Willis 2018 and

Tataru *et al*. 2023 and re-analyzed here.

Figure 5. Survival to flowering (a-f), fecundity (mean seed number per reproductive individual; g-l), and total fitness (mean seed number per planted seed; m-r) of *Mimulus guttatus* (filled square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite and meadow habitats, from each experimental year and the average (arithmetic mean) across years (f, l, r). Calculations of fecundity and total fitness for 2013 are based on fruit number because seed number was not recorded; 2013 was omitted from averaged fecundity and total fitness. Values from 2013 and 2019 are from Ferris and Willis 2018 and Tataru *et al*. 2023.

Table 1. Binomial analysis relating phenotypes to fitness, with fitness defined as whether or not a plant produced fruit (2013) or seeds (2019, 2021, 2022, 2023). A combined dataset of all years used fruit number for fitness. The strengths of selection for linear selection gradients are represented as β values. Asterisks indicate significance of selection in a trait: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Herkogamy was excluded from the combined analysis due to its absence from the 2013 dataset. Phenotypic data from 2013 and 2019 are published in Ferris and Willis 2018 and Tataru *et al.* 2023 and re-analyzed here.

Trait	Meadow					
	2013	2019	2021	2022	2023	Combined
Flowering time	-0.054	-0.252	-0.622	-0.314*	-0.291**	-0.018
Flower width	-0.238	0.118	0.429	0.125	-0.041	0.044
Leaf lobing	-0.217	0.050	-0.005	0.025	0.123	-0.025
Herkogamy	NA	-0.006	-0.143	0.086	0.015	NA
Plant height	0.173	0.587***	0.863**	0.355*	0.121	0.290 ***
Trait	Granite					
	2013	2019	2021	2022	2023	Combined
Flowering time	-1.226***	-1.938***	-0.431	NA	-0.385*	-0.438 **
Flower width	-0.008	-0.482**	-0.424*	NA	-0.204	-0.223 **
Leaf lobing	-0.004	-0.182	0.165	NA	0.070	0.062
Herkogamy	NA	0.331	-0.083	NA	-0.340	NA
Plant height	0.038	0.557**	-0.109	NA	1.946**	0.144

Table 2. Zero-truncated Poisson analysis relating phenotypes to fitness, with fitness defined as

the number of seeds produced, on individual experimental years of 2013, 2019, 2021, 2022, and

2023 and on a combined dataset of all years. The strengths of selection for linear selection

gradients are represented as β values. Asterisks indicate significance of selection in a trait: * $p <$

0.05, ** $p < 0.01$, *** $p < 0.001$. Herkogamy was excluded to from the combined analysis due to

its absence from the 2013 dataset. Phenotypic data from 2013 and 2019 are published in Ferris

and Willis 2018 and Tataru *et al.* 2023 and re-analyzed here.

Trait	Meadow					
	2013	2019	2021	2022	2023	Combined
Flowering time	-1.132	-1.479***	-1.333***	-0.110*	-0.168***	-1.018***
Flower width	-0.276	0.021	0.075	0.088	-0.096	-0.059
Leaf lobing	0.613	0.609***	-0.362***	0.051	-0.182***	0.058
Herkogamy		-0.053*	0.089***	0.133**	0.110***	
Plant height	0.669*	2.022***	0.708***	0.749***	0.858***	0.912***
Trait	Granite					
	2013	2019	2021	2022	2023	Combined
Flowering time	-5.183***	-0.609	-1.419***		-0.590***	-0.918***
Flower width	-0.219	0.174	-0.030		-0.120	-0.100
Leaf lobing	-0.229	-0.209*	0.170		-0.421***	0.043
Herkogamy		-0.196*	0.295***		-0.203***	
Plant height	0.927***	1.179**	2.526***		2.457***	0.786***

Supplementary Information

Fig S1 Density histograms of hybrid and parental values for phenotypic traits in 2021, 2022, and 2023.

Fig S2 Visual representation of selection gradients (β) of phenotypic traits measured for hybrids during experimental years in granite (open circles) and meadow (filled circles) habitats. The models do not include herkogamy. Shown are values from (f-j) binomial models and (k-o) zero-truncated Poisson models. Significance is indicated with asterisks (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) and negative graph areas are shaded grey for clarity. All β values are based on seed number, except for values from 2013 which are based on fruit number, as seed number was not recorded. Data from 2013 and 2019 are published in Ferris and Willis 2018 and Tataru *et al*. 2023 and re-analyzed here.

Fig S3 Patterns of cumulative selection with all years combined in meadow and granite habitats in a binomial analysis.

Fig S4 Combined survival to flowering, fecundity, and total fitness excluding 2023.

Fig S5 The average (geometric mean) of a) survival, b) fecundity, and c) total fitness across experimental years for *Mimulus guttatus* (filled square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite and meadow habitats. 2013 was omitted from averaged fecundity and total fitness. Values from 2013 and 2019 are from Ferris and Willis 2018 and Tataru *et al*. 2023.

Table S1 Hybrid genotypes and generations used each experimental year, with sample sizes of each genotype planted at each site with total experimental sample sizes.

Table S2 Intrapopulation parental crosses used in the 2023 experiment.

Table S3 Hybrid trait correlations in meadow and granite habitats for 2021-2023.

Table S4 Binomial analysis on whether or not a plant produced fruits.

Table S5 Zero-truncated Poisson analysis on number of fruit produced if any were produced.

Table S6 Strength of correlations between fruit number and seed number.

Table S7. Summary of models described in methods.

Table S8 Model selection for best fit models of plant survival and environmental variables.

Table S9 Trait means, standard error, standard deviation, and coefficient of variation from 2021.

Table S10 Trait means, standard error, standard deviation, and coefficient of variation from

2022.

Table S11. Trait means, standard error, standard deviation, and coefficient of variation from

2023.

Table S12. Generalized linear mixed models examining phenotypic plasticity between years and

habitat. Models included the phenotypic trait as the dependent variable, habitat and year with

their interaction as fixed effects and block nested within site as a random effect.

Table S13. Significance of genotype (G), environment (E), and genotype by environment (GxE)

interactions for each fitness term.

998
999

1000
1001 **Figure S1.** Density histograms of hybrid and parental values for phenotypic traits (flowering
1002 time, flower width, leaf lobing, herkogamy, plant height) in a) 2021, b) 2022, and c) 2023.
1003
1004
1005

1006

Figure S2. Visual representation of selection gradients (β) of phenotypic traits measured for
hybrids during experimental years in granite (open circles) and meadow (filled circles) habitats.
Models exclude herkogamy. Shown are (a-d) predicted values in each habitat based on native
species trait differences and estimated values from (e-h) binomial models and (i-l) zero-truncated
Poisson models. Significance of each β value is indicated with asterisks (* $p < 0.05$, ** $p < 0.01$,
*** $p < 0.001$) and negative graph areas are shaded grey for clarity. All β values are based on
seed number, except for values from 2013 which are based on fruit number, as seed number was
not recorded. Data from 2013 and 2019 are published in Ferris and Willis 2018 and Tataru et al.
2023 and re-analyzed here.

1016

1017
1018

Figure S3. Patterns of cumulative selection with all years combined in meadow andgranite habitats in a binomial analysis on flowering time (days from planting), width of first flower (mm), leaf lobing index, and plant height at flowering (mm to apical meristem). Y-axes show adjusted relative fitness (fruit number), statistically corrected for other variables included in the models, and the x-axes show unstandardized trait values for biological relevance. Data from 2013 and 2019 are from Ferris and Willis 2018 and Tataru *et al.* 2023 and re-analyzed here.

1025
Figure S4. Combined survival to flowering, fecundity (mean seed number per reproductive
individual), and total fitness (mean seed number per planted seed) of *Mimulus guttatus* (filled
square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite and meadow habitat
from 2013-2022 excluding 2023. 2013 was omitted from fecundity and total fitness calculations
because seed number was not recorded, only fruit number. Values from 2013 and 2019 are from
Ferris and Willis 2018 and Tataru *et al.* 2023.
1032

1033

Figure S5. Combined survival to flowering, fecundity (mean seed number per reproductive
individual), and total fitness (mean seed number per planted seed) of *Mimulus guttatus* (filled
square, solid line) and *M. laciniatus* (open triangle, dashed line) in granite and meadow habitat
from experimental years. These values represent geometric means. 2013 was omitted from
fecundity and total fitness calculations because seed number was not recorded, only fruit
number. Data from 2013 and 2019 are from Ferris and Willis 2018 and Tataru *et al.* 2023 and re-
analyzed here.

1041

1042

1043

1044 **Table S1.** Hybrid genotypes and generations used each experimental year, with sample sizes of
1045 each genotype planted at each site with total experimental sample sizes. The maternal genotype
1046 (WLF 47) is *M. laciniatus*, and the paternal genotypes (YVO 6 and 18) are *M. guttatus*. Details
1047 of the 2013 and 2019 experiments are published in Ferris and Willis 2018 and Tataru *et al.* 2023.
1048

Year	Hybrid genotype	Granite 1	Granite 2	Meadow 1	Meadow 2	Total
2013	F4 WLF 47 maternal x YVO 6 paternal	750 hybrids, 100 maternal, 100 paternal	750 hybrids, 100 maternal, 100 paternal	750 hybrids, 100 maternal, 100 paternal	750 hybrids, 100 maternal, 100 paternal	3,800
2019	F4 WLF 47 maternal x YVO 6 paternal	600 hybrids, 60 maternal, 60 paternal	660 hybrids, 44 maternal, 88 paternal	750 hybrids, 71 maternal, 100 paternal	750 hybrids, 50 maternal, 100 paternal	3,333
2021	F2 WLF 47 maternal x YVO 18 paternal	750 hybrids, 750 maternal, 750 paternal	750 hybrids, 750 maternal, 750 paternal	600 hybrids, 600 maternal, 600 paternal	600 hybrids, 600 maternal, 600 paternal	4,050
2022	F3 WLF 47 maternal x YVO 18 paternal	800 hybrids, 200 maternal, 200 paternal	800 hybrids, 200 maternal, 200 paternal	800 hybrids, 200 maternal, 200 paternal	800 hybrids, 200 maternal, 200 paternal	4,800
2023	F6 WLF 47 maternal x YVO 18 paternal	1500 hybrids, 200 maternal, 200 paternal, 200 local parent	1500 hybrids, 200 maternal, 200 paternal, 200 local parent	1500 hybrids, 200 maternal, 200 paternal, 200 local parent	1500 hybrids, 200 maternal, 200 paternal, 200 local parent	8,400

1049
1050

1051 **Table S2**. Intrapopulation parental crosses used in the 2023 experiment. Due to insufficient seed
1052 stock of single crosses of YVO and WLF, we used 2-3 crosses and original inbred lines,
1053 distributed evenly across blocks and sites. With habitats combined, WLF genotypes did not
1054 differ significantly in total seed count (F = 2.324, df = 2, p = 0.099). Similarly, YVO genotypes
1055 also did not differ significantly (F = 1.944, df = 3, p = 0.124).
1056

Habitat	Genotype	Survival to flowering	%Survival	Seed count	Mean Seed Count	Seed SD	N
Granite	WLF60xWLF47	81	44	585	7.31	21.27	184
	WLF61xWLF47	108	66	784	7.26	12.78	164
	WLF47 inbred	0	0	0	NA	NA	51
						Total:	**399**
	YVO18xYVO26	0	0	0	NA	NA	20
	YVO25xYVO18	12	8	417	32.08	67.69	159
	YVO26xYVO18	35	18	961	27.46	51.83	200
	YVO18 inbred	0	0	0	NA	NA	20
						Total:	**399**
Meadow	WLF60xWLF47	97	58	1760	17.25	32.41	166
	WLF61xWLF47	156	83	3201	20.13	33.95	187
	WLF47inbred	35	76	156	4.46	8.43	46
						Total:	**399**
	YVO18xYVO26	26	47	208	8	16.78	55
	YVO25xYVO18	26	23	378	14.54	47.43	114
	YVO26xYVO18	55	32	687	12.27	27.66	171
	YVO18 inbred	26	44	21	0.81	2.77	59
						Total:	**399**

1057

1058

1059 **Table S3.** Hybrid phenotypic correlations in meadow (grey shaded; lower triangle) and granite
1060 (upper triangle) habitats for a) 2021, b) 2022, and c) 2023. Asterisks indicate significant (* = p <
1061 0.05, ** = p < 0.01. *** = p < 0.001).

1062 (a)

	Flowering Time	Leaf Area	Leaf Lobing	Flower Width	Plant Height	Herkogamy
Flowering Time		-0.21***	-0.04	-0.08*	0.04	-0.08**
Leaf Area	-0.38**		-0.04	0.30***	0.75***	0.02
Leaf Lobing	0.02	0.11**		0.15*	0.35*	0.10
Flower Width	0.15**	0.07	0.03		0.62***	0.14**
Plant Height	0.50***	0.30**	0.10	0.58***		0.01
Herkogamy	0.06	-0.08	0.07	-0.07	-0.07	

1063

1064 (b)

	Flowering Time	Leaf Area	Leaf Lobing	Flower Width	Plant Height	Herkogamy
Flowering Time		-0.13	-0.12	0.24	-0.14	0.03
Leaf Area	0.01		-0.04	0.01	0.02	-0.01
Leaf Lobing	0.00	-0.06		-0.38	0.36	0.44*
Flower Width	-0.01	0.27***	-0.08		-0.31	-0.06
Plant Height	0.06	0.52***	-0.05	0.65***		-0.04
Herkogamy	-0.07	-0.10	0.13	-0.10	-0.13	

1065

1066 (c)

	Flowering Time	Leaf Area	Leaf Lobing	Flower Width	Plant Height	Herkogamy
Flowering Time		-0.14	0.04	0.29	0.88***	-0.32
Leaf Area	-0.16		0.10***	0.24	0.58***	-0.08
Leaf Lobing	-0.02	0.13		0.11	0.44	-0.07
Flower Width	0.07*	0.17	0.03		0.69***	-0.17
Plant Height	0.15***	0.33***	0.11**	0.39***		-0.07
Herkogamy	-0.06	0.12	0.03	-0.06	-0.06	

Table S4. Binomial analysis on whether or not a plant produced fruits on individual experimental years of 2019, 2021, 2022, and 2023. The strengths of selection for linear selection gradients are represented as β values. Asterisks indicate significance of selection in a trait: ***p < 0.001; **p < 0.01; *p < 0.05. Data from 2019 are published in Tataru *et al.* 2023 and re-analyzed here.

	Meadow			
Trait	**2019**	**2021**	**2022**	**2023**
Flowering time	-0.189	-0.413	-0.180	0.019
Flower width	0.126	0.391	0.170	0.004
Leaf lobing	-0.086	-0.162	0.038	0.160
Herkogamy	0.113	-0.024	0.056	-0.059
Plant height	0.144	0.879*	0.209	0.116
	Granite			
Trait	**2019**	**2021**	**2022**	**2023**
Flowering time	-0.534	-0.304		-0.177
Flower width	-0.319	-0.497*		0.107
Leaf lobing	-0.255	0.230		0.181
Herkogamy	0.388	-0.026		-0.129
Plant height	0.255	-0.210		2.461***

1076 **Table S5.** Zero-truncated Poisson analysis on number of fruit produced if any were produced on
1077 individual experimental years of 2013, 2019, 2021, 2022, and 2023. The strengths of selection
1078 for linear selection gradients are represented as β values. Asterisks indicate significance of
1079 selection in a trait: ***p < 0.001; **p < 0.01; *p < 0.05.
1080

	Meadow			
Trait	**2019**	**2021**	**2022**	**2023**
Flowering time	-0.551***	-0.802***	-0.153	-0.691***
Flower width	-0.072	-0.020	0.060	-0.261*
Leaf lobing	0.117	-0.011	0.126	0.049
Herkogamy	-0.027	0.044	-0.146	-0.006
Plant height	0.422***	0.431***	0.521***	0.332**
	Granite			
Trait	**2019**	**2021**	**2022**	**2023**
Flowering time	-0.649***	-0.819**		-0.401*
Flower width	-0.072	0.114		-0.396*
Leaf lobing	0.135	0.009		-0.122
Herkogamy	-0.090	-0.142		0.002
Plant height	0.167***	1.333***		2.678***

1081

1082

1083 **Table S6.** Strength of correlations between fruit number and seed number calculated using
1084 Kendall's correlation analysis.
1085

Year	Habitat	τ	p	n
2021	Meadow	0.56	0.001	1296
	Granite	0.59	0.001	809
2022	Meadow	0.64	0.001	545
2023	Meadow	0.75	0.0001	1393
	Granite	0.69	0.0001	480

1086

1087
1088

1089 **Table S7.** Summary of models described in methods.

Description of analysis	Model	Reference
plasticity	glmmTMB(trait ~ Habitat + (1|Site/Block))	Table S9, S10, S11
trait correlations	lmer(trait1 ~ trait2 + (1|Site/Block))	Table S3
survival and plant survival	glmmTMB(cbind(successes, failures) ~ soil*days*light*temp +(1|Site/Block), family = binomial)	Figure 2, Table S8
herbivory between habitats	glmer(herbivory ~ Habitat + (1|Site/Block), family = binomial)	Results text (environmental variation causes spatial and temporal variation in survival)
herbivory and fitness	glmer(seed ~ herbivory*Habitat + (1|Site/Block), family = poisson)	Results text (environmental variation causes spatial and temporal variation in survival)
phenotypic (2019-2023), with family=binomial or truncated_poisson	glmmTMB(seed ~ lobingz2 + plheightz2 + flwidthz2 + DaysToz2 + outcrossz2 + (1|Site/Block))	Figure 3, Tables 1 and 2
phenotypic (2013), with family=binomial or truncated_poisson	glmmTMB(fruit ~ lobingz2 + plheightz2 + flwidthz2 + DaysToz2 + (1|Site/Block))	Figure 3, Table 1, 2
phenotypic without outcrossing (2019-2023), with family=binomial or truncated_poisson	glmmTMB(seed ~ lobingz2 + plheightz2 + flwidthz2 + DaysToz2 + (1|Site/Block))	Figure S2
phenotypic with fruit (2019-2023), with family=binomial or truncated_poisson	glmmTMB(fruit ~ lobingz2 + plheightz2 + flwidthz2 + DaysToz2 + outcrossz2 + (1|Site/Block))	Table S4, S5
phenotypic (years combined), with family=binomial or truncated_poisson	glmmTMB(fruit ~ DaysToz + flwidthz + lobingz + plheightz + (1|Site/Block) + (1|year))	Figure 4, S3
snowpack and phenotypic traits, with family=binomial or truncated_poisson	glmmTMB(fruit ~ (DaysToz + flwidthz + lobingz + plheightz)*swc + (1|Site/Block) + (1|year))	Results text (Interannually fluctuation versus long-term directional selection)
GxE (survival)	glmer(survival ~ genotype*Habitat + (1| Site/Block), family - binomial(ink="logit"))	Figure 5
GxE (fecundity)	lmer(fecundity ~ genotype* Habitat + (1|Site/Block))	Figure 5
GxE (total fitness)	lmer(total_Fitness ~ genotype* Habitat + (1|Site/Block))	Figure 5

1090 **Table S8.** Best-fit models of plant survival and environmental variables. For each best-fit model,
1091 we report the df, Log(L), the likelihood score; AIC score, and delta(i), the difference between
1092 that model and the next top model's AIC.

Year	Habitat	Model	df	Log(L)	AIC	delta(i)
2021	meadow	survival ~ days + light + soil + temp + (1 | Site/Plot) + days:light + days:soil + light:soil + light:temp + soil:temp	12	-7994.438	16013	1.17
2021	granite	survival ~ days + light + soil + temp + (1 | Site/Plot) + days:soil + days:temp + light:temp + soil:temp	11	-1231.031	2484.5	0.12
2022	meadow	survival ~ days + light + soil + temp + (1 | Site/Plot) + days:light + days:temp + light:soil + light:temp	11	-4057.269	8136.7	0.22
2022	granite	survival ~ days + light + soil + temp + (1 | Site/Plot) + days:light + days:temp + light:temp + soil:temp	11	-1708.705	3439.7	1.82
2023	meadow	survival ~ days + light + soil + temp + (1 | Site/Plot) + days:soil + days:temp + light:temp + soil:temp	11	-4467.387	8956.8	0.15
2023	granite	survival ~ days + light + temp + (1 | Site/Plot)	6	-4348.601	8709.3	0.9

1093

1094

1095

1096

1097 **Table S9**. Trait means, standard error (SE), standard deviation (SD), and coefficient of variation
1098 (CV) from the 2021 experiment. Significant variation in trait means between habitat is indicated
1099 by asterisks (p-value < 0.001***; 0.01**; 0.05*).

1100

Genotype	Trait	Meadow			Granite		
		Mean	**SE**	**CV**	**Mean**	**SE**	**CV**
F$_2$	Fl. time	31.58***	0.61	0.45	14.57***	0.31	0.43
	Height	57.52***	1.40	0.57	28.43***	0.63	0.45
	Fl. Width	5.89	0.11	0.37	6.09	0.12	0.36
	Leaf area	48864.77**	3838.85	1.57	33811.16**	2769.86	1.52
	Lobing	0.08	0.00	0.45	0.07	0.00	0.37
	Herkogamy	-1.34***	0.05	-0.73	-0.86***	0.07	-1.36
M. guttatus **(YVO 18)**	Fl. time	44.59	0.99	0.34	35.08	4.23	0.42
	Height	74.65**	2.56	0.46	20.56**	3.50	0.51
	Fl. Width	7.49	0.22	0.33	5.88	0.81	0.36
	Leaf area	50215.94	6314.38	1.47	11519.75	1912.33	0.94
	Lobing	0.08	0.00	0.68	0.06	0.00	0.33
	Herkogamy	-1.63	0.10	-0.76	-0.94	0.31	-0.94
M. laciniatus **(WLF 47)**	Fl. time	26.7	0.49	0.45	19.21	0.54	0.48
	Height	34.04**	0.75	0.49	20.41**	0.72	0.57
	Fl. Width	1.96**	0.05	0.49	2.55**	0.15	0.73
	Leaf area	16642.84	1268.08	1.46	12059.12	1095.45	1.07
	Lobing	0.08	0.00	0.47	0.07	0.03	0.40
	Herkogamy	-0.92	0.10	-1.17	-0.74	0.08	-1.01

1101

1102 **Table S10**. Trait means, standard error (SE), standard deviation (SD), and coefficient of
1103 variation (CV) from the 2022 experiment. Significant variation in trait means between habitat is
1104 indicated by asterisks (p-value < 0.001***; 0.01**; 0.05*).
1105

Genotype	Trait	Meadow			Granite		
		Mean	SE	CV	Mean	SE	CV
F₃	Fl. time	59.92*	0.70	0.21	44.19*	3.10	0.32
	Height	54.89*	2.07	0.69	13.14*	1.52	0.53
	Fl. Width	5.52***	0.13	0.41	3.34***	0.30	0.41
	Leaf area	1013.85	77.72	1.31	208.62***	27.30	1.09
	Lobing	0.07***	0.00	0.33	0.09***	0.00	0.24
	Herkogamy	-1.35	0.05	-0.67	-0.92	0.17	-0.77
M. guttatus (YVO 18)	Fl. time	72.66	1.89	0.17	-	-	-
	Height	69.15	5.22	0.48	-	-	-
	Fl. Width	7.73	0.36	0.29	-	-	-
	Leaf area	1788.03*	220.71	0.71	169.69*	22.08	0.47
	Lobing	0.05*	0.00	0.27	0.08*	0.00	0.22
	Herkogamy	-2.40	0.13	-0.33	-	-	-
M. laciniatus (WLF 47)	Fl. time	44.53*	0.83	0.25	34.65*	1.08	0.30
	Height	15.61**	0.99	0.84	7.16**	0.42	0.57
	Fl. Width	1.25	0.08	0.74	1.22	0.14	0.97
	Leaf area	294.09*	22.64	0.97	97.56*	4.31	0.41
	Lobing	0.08*	0.00	0.30	0.09*	0.00	0.29
	Herkogamy	-0.85	0.05	-0.65	-1.01	0.11	-0.71

1106

1107 **Table S11**. Trait means, standard error (SE), standard deviation (SD), and coefficient of
1108 variation (CV) from the 2023 experiment. Significant variation in trait means between habitat is
1109 indicated by asterisks (p-value < 0.001***; 0.01**; 0.05*).
1110

Genotype	Trait	Meadow				Granite			
		Mean	SE	SD	CV	Mean	SE	SD	CV
F₆	Fl. time	43.04***	0.28	15.3	0.36	35.77***	0.36	19.67	0.55
	Height	91.71*	1.01	55.14	0.60	25.22*	0.27	14.81	0.59
	Fl. Width	5.62**	0.04	2.23	0.40	4.68**	0.04	2.35	0.50
	Leaf area	14342.24	287.00	15722.42	1.10	6183.81	173.97	9528.94	1.54
	Lobing	0.08	0.00	0.05	0.63	0.08	0.00	0.06	0.75
	Herkogamy	-0.34	0.03	1.4	-4.12	-0.4	0.02	1.09	-2.73
M. guttatus (YVO)	Fl. time	52.94***	0.61	12.14	0.23	44.6***	0.78	15.49	0.35
	Height	76.17***	1.75	34.94	0.46	34.56***	0.84	16.78	0.49
	Fl. Width	9.92	0.13	2.68	0.27	8.69	0.16	3.1	0.36
	Leaf area	18876.1*	942.86	18833.71	1.00	11853.5*	655.40	13091.55	1.10
	Lobing	0.07***	0.00	0.03	0.43	0.08***	0.00	0.02	0.25
	Herkogamy	-0.66	0.08	1.5	-2.27	-1.11	0.07	1.41	-1.27
M. laciniatus (WLF)	Fl. time	24.28	0.71	14.19	0.58	20.68	0.92	18.36	0.89
	Height	58.72*	1.72	34.45	0.59	21.16*	0.56	11.16	0.53
	Fl. Width	3.01*	0.08	1.59	0.53	3.37*	0.07	1.44	0.43
	Leaf area	17699.42	850.63	16991.34	0.96	8788.1	327.31	6537.97	0.74
	Lobing	0.15**	0.00	0.08	0.53	0.11**	0.00	0.07	0.64
	Herkogamy	-0.36**	0.05	0.98	-2.72	0.12**	0.05	0.92	7.67

1111
1112
1113

1114   **Table 12.** Generalized linear mixed models examining phenotypic plasticity between years and
1115   habitat. Models included the phenotypic trait as the dependent variable, habitat and year with
1116   their interaction as fixed effects and block nested within site as a random effect.
1117

Trait	Term	Estimate	SE	t-value	p-value	
Flowering time	(Intercept)	7921.79	527.25	15.025	2.28E-48	***
Flowering time	habitat	2587.42	639.39	4.047	5.39E-05	***
Flowering time	year	-3.9	0.26	-14.955	5.81E-48	***
Flowering time	habitat*year	-1.28	0.32	-4.060	5.10E-05	***
Plant height	(Intercept)	-3433.89	809.76	-4.241	2.33E-05	***
Plant height	habitat	5925.78	978.14	6.058	1.65E-09	***
Plant height	year	1.72	0.4	4.301	1.79E-05	***
Plant height	habitat*year	-2.95	0.48	-6.087	1.38E-09	***
Flower width	(Intercept)	105.18	122.4	0.859	0.39	
Flower width	habitat	652.45	144.51	4.515	7.10E-06	***
Flower width	year	-0.05	0.06	-0.823	0.41	
Flower width	habitat*year	-0.32	0.07	-4.519	6.95E-06	***
Leaf area	(Intercept)	-3350437.73	345581.02	-9.695	1.01E-21	***
Leaf area	habitat	1923215.21	415529.75	4.628	3.93E-06	***
Leaf area	year	1663.58	170.92	9.733	7.10E-22	***
Leaf area	habitat*year	-954.01	205.58	-4.641	3.71E-06	***
Leaf lobing	(Intercept)	-370.59	81.29	-4.559	5.47E-06	***
Leaf lobing	habitat	65.65	96.23	0.682	0.495	
Leaf lobing	year	0.18	0.04	4.559	5.47E-06	***
Leaf lobing	habitat*year	-0.03	0.05	-0.682	0.495	
Herkogamy	(Intercept)	-569.84	73.52	-7.751	2.12E-14	***
Herkogamy	habitat	-102.43	164.4	-0.623	0.533	
Herkogamy	year	0.28	0.04	7.739	2.33E-14	***
Herkogamy	habitat*year	0.05	0.08	0.625	0.532	

1118
1119
1120

1121 **Table S13**. Significance of genotype (G), environment (E), and genotype by environment (GxE)
1122 interactions for each fitness term. Significant p-values are in bold ($p < 0.05$). Details of models
1123 are in Table S7.
1124

Year	Fitness term	G	E	G*E
2013	**Survival**	Mean sq = 17.8400 F = 17.84 p = 0.706524	Mean sq = 2.1827 F = 2.1827 p = 0.763166	Mean sq = 10.8925 F = 10.8925 **p = 0.001112**
	Fecundity	df = 144.23 F = 0.0072 p = 0.93259	df = 106.14 F = 3.8757 p = 0.5160	df = 143.49 F = 6.6423 **p = 0.01097**
	Total Fitness	df = 595.92 F = 45.096 **p < 0.0001**	df = 196.74 F = 54.167 **p < 0.0001**	df = 595.92 F = 62.473 **p < 0.0001**
2019	**Survival**	NA NA NA	NA NA NA	NA NA NA
	Fecundity	df = 10.1845 F = 0.4372 p = 0.5232	df = 5.6865 F = 0.2081 p = 0.6652	NA NA NA
	Total Fitness	df = 572 F = 0.8102 p = 0.3685	df = 572 F = 1.6206 p = 0.2035	df = 572 F = 0.7691 p = 0.3809
2021	**Survival**	Mean sq = 259.032 F = 259.032 **p < 0.0001**	Mean sq = 121.217 F = 121.217 **p < 0.0001**	Mean sq = 22.394 F = 22.394 **p < 0.0001**
	Fecundity	df = 828.26 F = 0.0846 p = 0.771185	df = 687.43 F = 7.4232 **p < 0.001**	df = 828.26 F = 0.5004 p = 0.479517
	Total Fitness	df = 2469.72 F = 63.472 **p < 0.0001**	df = 223.04 F = 39.951 **p < 0.0001**	df = 2469.72 F = 16.277 **p < 0.0001**
2022	**Survival**	Mean sq = 118.612 F = 118.6124 **p < 0.0001**	Mean sq = 5.397 F = 5.3965 p = 0.984	Mean sq = 0 F = 0.0003 p = 0.986
	Fecundity	df = 299.16 F = 7.3132 **p < 0.001**	df = 153.12 F = 15.9022 **p < 0.0001**	NA NA NA

	Total Fitness	df = 1396.10 F = 32.493 **p < 0.0001**	df = 196.92 F = 24.333 **p < 0.0001**	df = 1396.10 F = 26.408 **p < 0.0001**
2023	**Survival**	Mean sq = 118.612 F = 246.0353 **p < 0.0001**	Mean sq = 8.625 F = 8.6250 **p < 0.0001**	Mean sq = 1.173 F = 1.1729 p = 0.281073
	Fecundity	df = 642.33 F = 4.0579 **p = 0.04438**	df = 244.77 F = 2.4746 p = 0.11699	df = 640.32 F = 26.7453 **p < 0.0001**
	Total Fitness	df = 1340.93 F = 23.236 **p < 0.0001**	df = 219.93 F = 12.279 **p < 0.0001**	df = 1340.93 F = 23.790 **p < 0.0001**
Average	**Survival**	Mean sq = 357.18 F = 357.1792 **p < 0.0001**	Mean sq = 0.14 F = 0.1353 p = 0.2814	Mean sq = 32.78 F = 32.7822 **p < 0.0001**
	Fecundity	df = 877.52 F = 0.105 p = 0.746005	df = 716.72 F = 7.0909 **p = 0.007921**	df = 883.41 F = 0.4224 p = 0.515888
	Total Fitness	df = 3034.37 F = 47.8337 **p < 0.0001**	df = 290.96 F = 46.0596 **p < 0.0001**	df = 3034.37 F = 8.1155 **p < 0.001**

1125