Control of innate olfactory valence by segregated cortical amygdala circuits

James R. Howe[1,2,3,9], Chung-Lung Chan[1,9], Donghyung Lee[1], Marlon Blanquart[1], Haylie K. Romero[2,3,4], Abigail N. Zadina[5,†], Mackenzie E. Lemieux[6], Fergil Mills[6,††], Paula A. Desplats[3,4,7], Kay M. Tye[1,6,8], & Cory M. Root[1,*]

[1] Department of Biological Sciences, Section of Neuroscience, University of California, San Diego, La Jolla, CA 92093, USA

[2] Neurosciences Graduate Program, University of California, San Diego, La Jolla, CA 92093, USA

[3] Department of Neurosciences, University of California, San Diego, La Jolla, CA 92093, USA

[4] Center for Circadian Biology, University of California, San Diego, La Jolla, CA 92093, USA

[5] Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, 10027, USA

[6] Salk Institute for Biological Sciences, La Jolla, CA 92037, USA

[7] Department of Pathology, University of California, San Diego, La Jolla, CA 92093, USA

[8] Howard Hughes Medical Institute, La Jolla, CA 92037, USA

[9] These authors contributed equally

[†] Current affiliation: Department of Neuroscience and Behavior, Barnard College of Columbia University, New York City, NY 10027, USA

[††] Current affiliation: Department of Neurobiology, University of Utah School of Medicine, Salt Lake City, UT 84112, USA

* To whom correspondence should be addressed:

Cory M. Root, PhD
University of California, San Diego
9500 Gilman Drive
3119 Pacific Hall
La Jolla, CA 92093-0357
cmroot@ucsd.edu

SUMMARY

Animals perform innate behaviors that are stereotyped responses to specific evolutionarily
relevant stimuli in the absence of prior learning or experience. These behaviors can be reduced
to an axis of valence, whereby specific odors evoke approach or avoidance. The cortical
amygdala (plCoA) mediates innate attraction and aversion to odor. However, little is known
about how this brain area gives rise to behaviors of opposing motivational valence. Here, we
sought to define the circuit features of plCoA that give rise to innate olfactory behaviors of
valence. We characterized the physiology, gene expression, and projections of this structure,
identifying a divergent, topographic organization that selectively controls innate attraction and
avoidance to odor. First, we examined odor-evoked responses in these areas and found sparse
encoding of odor identity, but not valence. We next considered a topographic organization and
found that optogenetic stimulation of the anterior and posterior domains of plCoA elicits
attraction and avoidance, respectively, suggesting a functional axis for valence. Using single cell
and spatial RNA sequencing, we identified the molecular cell types in plCoA, revealing an
anteroposterior gradient in cell types, whereby anterior glutamatergic neurons preferentially
express *Slc17a6* and posterior neurons express *Slc17a7*. Activation of these respective cell
types recapitulates appetitive and aversive valence behaviors, and chemogenetic inhibition
reveals partial necessity for valence responses to innate appetitive or aversive odors. Finally,
we identified topographically organized circuits defined by projections, whereby anterior neurons
preferentially project to medial amygdala, and posterior neurons preferentially project to nucleus
accumbens, which are respectively sufficient and necessary for innate negative and positive
olfactory valence. Together, these data advance our understanding of how the olfactory system
generates stereotypic, hardwired attraction and avoidance, and supports a model whereby
distinct, topographically distributed plCoA populations direct innate olfactory valence responses
by signaling to divergent valence-specific targets, linking upstream olfactory identity to
downstream valence behaviors, through a population code. This represents a novel circuit motif
in which valence encoding is represented not by the firing properties of individual neurons, but
by population level identity encoding that is routed through divergent targets to mediate distinct
valence.

INTRODUCTION

Innate behaviors are ubiquitous across the animal kingdom, allowing specific sensory stimuli to yield stereotypical behavioral responses even in the absence of learning or past experience. These behaviors include feeding, fighting, fleeing, and mating, among others, and many can be simplified onto an axis of positive or negative valence representing approach and avoidance. Innate behaviors are the result of evolutionary selection, guiding initial behaviors that can be updated by future experiences. Given that innate behaviors are genetically hardwired, it is thought they should be mediated by simple circuits with specified connections between layers of the nervous system. Innate behaviors are common across sensory modalities but are especially prominent in olfaction, whereby diverse chemical signals, critical to survival and reproduction, must be quickly and robustly detected and processed, in the absence of prior experience. For instance, predator odors represent a potentially imminent threat and necessitate a quick, decisive, aversive response.[1] Conversely, innately appetitive odors represent potentially rewarding stimuli like food or heterospecifics, inducing attraction.[2–4] These odors comprise a small subset of perceptible chemical space, and the detection of specific odorants is both species-specific and under genetic control.[5–7]

Valence is a fundamental perceptual feature of olfaction. Motivational valence can be defined as seeking or avoiding specific stimuli, and it is observed across sensory stimuli in both innate and learned responses. Multiple circuit motifs have been proposed to mediate such valence responses.[8] In the simplest form, labeled line motifs segregate information from sensation to action throughout the nervous system. This has been observed in the taste and somatosensory systems, as well as hints of labeled lines in the olfactory system, where individual glomeruli are necessary and sufficient for innate responses consistent with this model.[9–11] Many circuits are organized along divergent path motifs, where a region receives the same sensory input but instead acts akin to a switchboard, processing and sorting its output to distinct downstream targets to convey positive or negative signals. This motif is most associated with the BLA and associated learning processes.[8,12] Still other circuits contain opposing components motifs, in which opposing inputs target a single effector region to control the balance of one target. It remains unclear which, if any, of these generalized circuit motifs are present in the olfactory system.

Olfactory sensation begins with olfactory sensory neurons (OSNs) in the olfactory epithelium (OE) that express a single receptor, projecting to spatially stereotyped glomeruli in the olfactory bulb (OB). Postsynaptic mitral/tufted cells within the OB project in parallel to third-order olfactory areas, including the posterolateral cortical amygdala (plCoA). Unlike in other third-order olfactory areas, such as piriform cortex, projections from individual glomeruli from the OB to plCoA are spatially restricted and stereotyped, consistent with genetically hardwired circuits.[13–16] Past work has demonstrated that plCoA is necessary and sufficient for innate olfactory responses, with spatially ordered labeling of responsive neurons via immediate early gene labeling.[3] However, other work suggests that there is no spatial organization to odor responses or valence encoding in plCoA.[17] Thus, a further investigation of plCoA organization is necessary to understand how this structure controls innate olfactory responses.

Distinct brain areas employ distinct coding strategies to represent information. Neuronal
ensembles within all other major olfactory regions observed thus far, such as the anterior
olfactory nucleus, OB, OE, olfactory tubercle (OT), piriform cortex (PIR), and tenia tecta all
generally perform sparse population encoding of odor identity, despite the major differences in
neuronal composition, organization, and function between the six regions.[12,17–30] On the other
hand, ensembles within amygdala subnuclei, most notably the basolateral amygdala (BLA),
instead tend to represent the valence of stimuli instead, with considerable heterogeneity based
on a given population's projection target, molecular identity, and topography.[12,18,19] A complete
investigation of the encoding properties and organization of plCoA is needed to determine how
innate olfactory valence emerges from this region.

To identify the plCoA circuitry that underlies innate valence, we investigated multiple
intersecting scales of organization, from single cell transcriptomes and spatial gene expression,
to circuit mapping, manipulation, and physiology. First, we used 2-photon calcium imaging to
find that plCoA ensembles encode odor identity, but not valence. Next, we identify a functional
gradient in plCoA where activation of anterior and posterior neurons drives responses of
opposing valence. Next, we characterized the cell types within plCoA, identifying novel,
molecularly defined populations specific to each domain of plCoA, which are respectively
sufficient and partially necessary for innate olfactory valence. Further, we perform
comprehensive projection mapping to identify downstream projection targets of plCoA,
identifying projections to the medial amygdala (MeA) and nucleus accumbens (NAc) that are
enriched based on molecular and topographic identity. Finally, manipulations of neuronal
ensembles projecting to these targets are sufficient and necessary to control innate olfactory
valence responses. Together, these findings identify a novel topographically distributed circuit
from plCoA to MeA and NAc that controls innate olfactory aversion and attraction, respectively,
consistent with a hybrid model mixing features of labeled lines and divergent path circuit motifs.

RESULTS

Population Encoding of Odor Identity in plCoA

To better understand how plCoA circuitry mediates innate olfactory valence, we first decided to examine the relationship between its spatial organization and odor-evoked activity. Prior analysis of immediate early gene expression following minutes-long odor exposure suggests that activity in the anterior and posterior domains of plCoA could respectively mediate innate aversive and appetitive odor responses.[3] However, *in vivo* electrophysiology with high-density electrode arrays found no evidence for spatial organization or valence encoding in plCoA.[17] These two studies propose contradictory models of plCoA encoding properties that are both plausible, given that the former is common in the extended amygdala and insular cortex, and the latter is ubiquitous across olfactory regions.[22,31,32] We speculated that technical differences could be responsible for these opposing findings. Immediate early gene labeling has extremely low temporal resolution and likely requires a high amount of neural activity to activate gene expression. On the other hand, the recording sites in the latter study appear biased towards the middle of plCoA, and odor was given for 2 seconds in interleaved trials, whereas olfactory valence behavior evolves on a minutes-long timescale.[3,33] We wondered if the apparent contradictions in these studies could be resolved by applying a longer odor delivery protocol that both matches the timescale of behavior and balances spatial and temporal resolution.

Therefore, we developed an approach to image neural activity in plCoA with a modified odor delivery schedule, expressing GCaMP8s targeted towards either the anterior or posterior subsection of plCoA and implanting a gradient-index relay (GRIN) lens above to allow *in vivo* imaging of calcium transients via head-fixed two-photon microscopy (**Figure 1A-C, S1A-B**). We then examined calcium responses in these mice during a long odor exposure, where odors were presented repeatedly in 20-trial blocks for 5 seconds each in counterbalanced order (**Figure 1D**). We chose two odorants of each innate valence: the appetitive odors 2-phenylethanol (2PE) and peanut oil (PEA), the neutral odors heptanol (HEP) and isoamyl acetate (IAA), and the aversive odors trimethylthiazoline (TMT) and 4-methylthiazoline (4MT).[3,17] In total, we recorded Ca^{2+} signals from 345 neurons across 13 mice.

First, we pooled anterior and posterior plCoA neurons together and performed hierarchical clustering of their trial-averaged responses to the 6 odors to categorize odor responses in an unbiased manner (**Figure 1E**). Consistent with the previous *in vivo* electrophysiology study, we found that the majority of plCoA neurons did not seem to selectively respond to odors of one valence group (**Figure 1E-F**). Across mice, a majority of plCoA neurons did not reliably respond to any of the six odors, and activity was sparse: only 34.5% of plCoA neurons responded to 1 of the 6 odors and 10% to 2 odors, while a much smaller portion of plCoA neurons responded to 3 or more odors (**Figure 1G**). Further, we found no significant relationship between the valence of odor and proportion of responsive neurons, and no difference in the proportion responsive to the different odors across anterior and posterior plCoA, suggesting a lack of bias in responsiveness to aversive or appetitive odors across the anterior-posterior axis (**Figure 1H, S1C**).

Figure 1. The plCoA encodes innately-valenced odor identity using a population code.

(A) Schematic representation of virus injection and GRIN lens implantation into aplCoA or
pplCoA for two-photon microscopy.
(B-C) Representative images (**B**) and traces (**C**) of fluorescence changes in individual neurons
over an approximately 20-minute period.
(D) Schematic of odor exposure paradigm. Each trial presented 5 seconds of odor followed by a
variable inter-trial interval (20-30s). Odors were present in blocks of 20 trials per odor, with 2
counterbalanced block schedules (1 & 2). Six odors were used: the appetitive odors 2-
phenylethanol (2PE) and peanut oil (PEA), the neutral odors heptanol (HEP) and isoamyl
acetate (IAA), and the aversive odors trimethylthiazoline (TMT) and 4-methylthiazoline (4MT).
(E) Heatmap of trial-averaged and Z-scored odor-evoked activity over time from pooled plCoA
neurons. Responses are grouped by hierarchical clustering, with the dendrogram (right). Odor
delivery marked by vertical red lines.
(F) Average of trial-averaged and Z-scored odor-evoked activity for each cluster concatenated.
The order of color-coded blocks corresponds to the order of clusters in (**E**).
(G) Proportion of neurons responsive to different numbers of odors. Bars represent the mean
across 13 animals and the error bars show SEM.
(H) Proportion responsive to each odor for aplCoA (red) or pplCoA (blue).
(I) Valence scores of individual neurons. White circles show the median of each distribution,
whereas the gray rectangle shows the 25th-75th percentile range.
(J) Proportion of neurons with significant valence scores calculated as a function of trial number.
Calculated with a 10-trial moving window. Top half shows those with significant positive valence
scores, the bottom half shows those with significant negative valence scores.
(K) Left, MNR accuracies for all pooled plCoA neurons (data) and a control distribution where
the training labels are shuffled (shuffled) in a violin plot. Right, proportion of neurons in each
animal that have MNR accuracy greater than the 95th percentile of the shuffled MNRs.
(**L**) Ranked sub-accuracies for single-neuron MNR classifiers in violin plots for those trained on
real (top) or shuffled (bottom) data. Black lines connect the median across each rank.
(**M**) Scatter plot of percentage of sub-accuracies above the 95th percentile of the shuffled
controls for each animal plotted against the rank of the sub-accuracy. Red and blue lines show
the linear and exponential fit, respectively, and a gray dashed line connects data averages.
(**N**) Cross-validated average accuracies of multinomial SVM's plotted as a function of the
number of neurons used for training during the odor period. Circles represent the mean across
100 iterations of random sampling of neurons and error bars show the standard deviation.
(**O**) An example confusion matrix for a multinomial SVM trained with 200 neurons.
(P) Comparison of inter-valence and intra-valence confusion across number of neurons used in
training the classifiers. Filled circles show the average of the data across 100 iterations, open
circles show shuffled controls.
(Q) The normalized average distance between odor pairs that have different valence (inter) or
same valence (intra).
Across panels, ns, not significant. Additional specific details of statistical tests can be found in
Supplemental Table 1.

We next attempted to quantify valence encoding by calculating a valence score for each
neuron by subtracting the average integrated aversive odor response from the average
integrated appetitive odor response and dividing this resulting difference in valence by the
integrated odor response ($V = \frac{\Sigma_{appetitive} - \Sigma_{aversive}}{\Sigma_{total}}$). Although the valence score was more
negatively skewed amongst anterior neurons than posterior ones, there was no significant
difference in their distributions, failing to support a model in which anterior and posterior plCoA
neurons broadly encode opposing innate olfactory valence (**Figure 1I**). Given that the innate
aversive and appetitive behaviors evolve over time in freely moving animals, we wanted to
assess if odor responses changed between early and late trials. Using a moving window of 5
trials, we examined if the proportion of neurons with significant aversive or appetitive valence
scores differed over time in anterior or posterior plCoA but found no significant interaction
between time and imaging location (**Figure 1J**).

We further investigated odor encoding at the single neuron level using multinomial
regression (MNR) 6-odor classifiers trained on single neuron data and found that they perform
only marginally better than chance, indicating that the majority of single plCoA neurons do not
broadly encode discriminatory information about odor identity (**Figure 1K**). Across mice, only
43.1% have overall classification rates above the 95th percentile of shuffled controls, indicating
that the majority of single plCoA neurons do not broadly encode discriminatory information
about odor identity. Interestingly, when comparing the ranked sub-accuracies for single odors
compared to shuffled controls, we found the discrepancy between data and shuffled controls to
be accentuated in the 2 highest performing sub-accuracies (**Figure 1L**). Quantifying the
proportion of neurons with sub-accuracies higher than the 95th percentile of the shuffled across
biological replicates, we found that the decrease of sub-accuracies down the rank falls
exponentially, rather than linearly (**Figure 1M**). This, paired with the observation that most
plCoA neurons respond to one or none of the six odors tested, indicates that individual plCoA
neurons are sparsely responsive and carry little information about identity or valence.

Surprisingly, we saw no evidence of valence encoding at the single neuron level in either
the anterior or posterior plCoA. Instead, the plCoA neurons appear to primarily encode odor
identity in a sparse manner much like the other higher-order olfactory areas, known to use a
population code. Thus, we next considered identity and valence encoding at the population
level. In contrast to the poor single neuron classification, SVMs trained on population-level data
vastly outperform the shuffled data, indicating good encoding of identity at the population level
(**Figure 1N**). We further used a confusion matrix to ask whether training the classifier with a
given odor could accurately predict the identity of another odor. In the matrix each column
corresponds to a predicted class and each column corresponds to the actual class (**Figure 1O**).
If the population level activity is similar between odor pairs, we expect the confusion to be
higher than less similar pairs. The output of a confusion matrix yields a proportion of true class
labels, whereby a high proportion indicates good prediction. We found no difference between
the confusion rates for intra-valence classification or inter-valence classification, suggesting a
lack of valence encoding at the population level (**Figure 1P**). The similarity between inter-
valence and intra-valence confusion was mirrored when quantifying the range-normalized
pairwise Euclidean distance across biological replicates (**Figure 1Q, S1D**). Thus, plCoA

appears to encode odor identity in a population code like other higher order olfactory regions,
with no apparent encoding of valence.

A Functional Axis for Valence in plCoA

Since we did not observe clear evidence for valence encoding, we considered other
organizational principles that could support behaviors of opposing valence. Spatial organization
for valence processing has been previously observed in the insular cortex and BLA, whereby
discrete subsections of the region contain neurons wired to preferentially signal positive or
negative valence.[18,19,34,35] Given plCoA's spatially-ordered afferent projections from OB, we next
hypothesized that plCoA circuitry could still be organized along the anterior-posterior (AP) axis
to generate innate olfactory valence. If true, it follows that activation of small ensembles of
neurons along the AP axis should generate behavioral responses along a corresponding axis of
valence. We tested this prediction by expressing channelrhodopsin (ChR2) in subsets of
neurons at different positions along this axis and photostimulating them during behavioral
testing in the previously established four-quadrant open field arena.[3]

Based on cytoarchitecture, we more formally parcellated plCoA into three domains: the
anterior plCoA (aplCoA), a two-layered region on the ventral surface lateral to the anterior
cortical amygdala, the posterior plCoA (pplCoA), a three-layered region on the ventrolateral
surface lateral to the posteromedial cortical amygdala, and a middle transition zone (mplCoA)
between them (**Figure 2A**).[3,36] To determine the potential relationship between position on the
anterior-posterior axis of plCoA and evoked valence, we performed optogenetic stimulation at
points along this entire axis, expressing channelrhodopsin (ChR2) and implanting fibers into
each zone (**Figures 2B-C, S2A**). Valence using the four-quadrant open field assay, where mice
freely explored a chamber with or without stimulation, and valence was scored with a
performance index measuring quadrant occupancy relative to chance, as well as the mean
distance to the corner port. Mice experience a 10 minute baseline period followed by 15 minutes
of closed loop optogenetic (470nm, increasing up to a maximum of 10 Hz as the mouse
proceeds closer to the corner port) stimulation in one quadrant (**Figure 2D**).[3]

Throughout the trial period, we observed a negative linear relationship between the
anterior-posterior position of the photostimulation site and the evoked change in performance
index and distance to the corner port in ChR2, but not eYFP-infected mice, where evoked
valence shifted from positive to negative as stimulation became more anterior. (**Figure 2E-F**).
We grouped these responses to determine whether these bidirectional responses were specific
to the identified plCoA zones. We found that photostimulation in aplCoA significantly reduced
time spent in the 'on' quadrant and increased the average distance to the corner port during the
treatment period, indicating activation of aplCoA neurons is aversive and leads to avoidance of
the quadrant paired with stimulation (**Figure 2G-I**). We also found the opposite was true in
pplCoA, where stimulation in that zone instead increased the time in the 'on' quadrant and
decreased average distance to the corner port, indicating pplCoA neuron activation instead is
appetitive and leads to attraction to the stimulation quadrant (**Figure 2J-L**).

Fig. 2. The plCoA has topographic organization capable of driving approach and avoidance behaviors.

(A) Schematic of plCoA domains divided into anterior (aplCoA), middle (mplCoA), and posterior (pplCoA) regions based on histology, positioning, and gradients observed in past observations.[3,15]

(B) Strategy to activate anterior-posterior topographical ensembles via optogenetics.

(**C**) Representative histology and fiber/virus placement for aplCoA and pplCoA ChR2 animals.

(**D**) Schematic of four-quadrant open field behavioral assay with closed-loop photostimulation.

(E-F) Linear-fit of change in performance index (**E**) or mean port distance (**F**) as a function of anterior-posterior position along plCoA for optical stimulation.

(G) Paths traveled during the stimulus period for a representative mouse (left) and baseline-normalized collective heatmaps (right) from both the ChR2- and eYFP-infected groups with aplCoA-localized fiber implants. Lower right stimulus quadrant indicated in blue.

(H-I) Mean effect of Photostimulation of aplCoA neurons on time spent in stimulated quadrant (performance index) (**K**) and distance from the corner (port distance) (**I**).

(J) Paths traveled during the treatment period for a representative mouse (left) and baseline-normalized collective heatmaps (right) from both the ChR2- and eYFP-infected groups with pplCoA-localized fiber implants. Lower right stimulus quadrant marked in blue.

(**K-L**) Mean effect of photostimulation of pplCoA neurons infected with ChR2, but not eYFP, is sufficient to increase time spent in the stimulation quadrant (**K**) and reduce its average distance from the stimulation port during the stimulation period (**L**).

Abbreviations: aplCoA, anterior zone of posterolateral cortical amygdala; mplCoA, middle zone of posterolateral cortical amygdala; pplCoA, posterior zone of posterolateral cortical amygdala.

Across panels, ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Specific details of statistical tests can be found in Supplemental Table 1.

We further examined the effects of anterior-posterior plCoA stimulation on other
behaviors to determine whether these effects were specific to valence, or if they extended to
other affective or motor phenomena. Using the elevated plus maze, we found no change in
anxiety based on open arm time or entries, across both the length of plCoA or within either
aplCoA or pplCoA (**Figure S3A-C, S3E-F, S3H-I**). In the open field test, we similarly found no
changes to thigmotaxis, based on time spent in corners of the open field, or exploration, based
on time spent in the center of the open field (**Figure S3K-M, S3EO-P, S3R-S**). Further,
locomotion remained constant during stimulation across both assays and the entirety of plCoA
(**Figure S3D, S3G, S3J, S3N, S3Q, S3T**). Together, these data indicate that the effects of
plCoA neuron activation across the entire anterior-posterior axis are specific to valence, with
few other behavioral effects. Overall, we find activation of plCoA neurons is sufficient to drive
valence behaviors in a topographically organized and bidirectional manner, where aplCoA
drives aversion and pplCoA drives attraction.

Molecular Diversity of Transcriptomic plCoA Cell Types Along the Anteroposterior Axis

Having identified a functional axis sufficient to produce valence behaviors (**Figure 2**) that
does not appear to encode valence (**Figure 1**), we next considered if the plCoA could be
topographically organized by molecular cell type to support behaviors of opposing valence.
Thus, we sought to determine if there is an axis of molecular cell types along the anteroposterior
domains of plCoA. To investigate this phenomenon, we performed single-nucleus RNA
sequencing (snRNA-seq) to determine the cell type composition and its relationship to the
anterior-posterior axis of plCoA.[37] To simultaneously profile these cell types and identify
domain-specific patterns, we separately extracted tissue samples from aplCoA and pplCoA by
microdissection, verified accurate dissection by histology, and pooled qualifying samples from
the selected plCoA domain for each sequencing run (**Figure 3A, S4A-D**). We also confirmed
there were few region- or batch-specific differences in sequencing depth or nuclear quality
markers (**Figure S4E-M**). Clustering of sequenced nuclei by gene expression allowed us to
initially identify all major canonical neuronal and glial cell types in plCoA based on known
marker genes identified in past scRNA-seq studies (**Figure 3B-C**).[38,39] Neurons in plCoA are
80% glutamatergic, while 20% of neurons are GABAergic (**Figure 3D**). We also identified large
numbers of vascular leptomeningeal cells (VLMCs) and arachnoid barrier cells (ABCs), two
fibroblast-like meningeal cell types that interface with vasculature and form a barrier between
the brain and CSF, likely due to meningeal presence on the cortical surface during extraction.[39–
41]

We further characterized the heterogeneity of glutamatergic and GABAergic neurons
within plCoA by re-processing and subclustering both major neuronal cell types. Within
glutamatergic neurons, we identified 14 distinct subtypes by gene expression, with largely
continuous variation between glutamatergic subtypes (**Figure 3E**). However, when examining
the relationships between these subtypes, we identified two broader groups of glutamatergic
neurons via hierarchical clustering, where subtypes within each group displayed a lower
Euclidean distance from one another in high-dimensional gene expression space (**Figure 3F**).

Fig. 3. Transcriptomic heterogeneity of plCoA molecular cell types.

(A) Schematic of freeze and-re-pool strategy for snRNA-seq.
(B) Two-dimensional UMAP (n = 47,132 nuclei, see also **Figure S4.3**), colored by broad cellular
identity assigned by graph-based clustering of neuronal and non-neuronal nuclei.
(**C**) Cell-type-specific expression of canonical marker genes indicating broad cellular identity in
the brain. Dot size is proportional to percentage of nuclei expressing the marker, with color
scale representing normalized expression level.
(**D**) Total proportion of cells of each identified type in each domain of plCoA.
(E) Two-dimensional UMAP of glutamatergic neurons, colored by molecular cell type.
(F) Clustered heatmap showing Euclidean distance between averages of each subtype
positioned based on hierarchical clustering (left), and dot plot of marker genes for all
glutamatergic subtypes (right).
(G) Two-dimensional UMAP of GABAergic neurons, colored by molecular cell type, like in (**E**).
(H) Clustered heatmap showing Euclidean distance between averages of each subtype
positioned based on hierarchical clustering (left), and dot plot of marker genes for all GABAergic
subtypes (right), like in (**F**).

Each of these two broader groups had a marker for every type within either group, where the
larger Group 1 of glutamatergic neurons express *Slc17a7 (VGLUT1)*, and the smaller Group 2
expresses *Slc17a6 (VGLUT2)*. Within each of these glutamatergic groups, most observed
marker genes are non-canonical in the amygdala and cortex, suggesting unique glutamatergic
ensembles or patterns of gene expression within glutamatergic neurons in plCoA compared to
other regions previously described. Two subtypes did fall outside of either broad glutamatergic
group, varying in a more discrete manner than most plCoA glutamatergic neuron subtypes.
Interestingly, examination of data from the Allen ISH Atlas for their respective marker genes
Etv1 and *Sim1* showed these two groups fall into adjacent regions outside of plCoA, where
Glut.Etv1 neurons localize to the posterior basomedial amygdala and Glut.Sim1 neurons
localize to the nucleus of the lateral olfactory tract (**Figure S5A-B**). However, gene expression
patterns for GABAergic neurons displayed an opposing form of heterogeneity, where subtypes
are far more discrete, without broad groups linking related subtypes (**Figure 3G**). Marker genes
for GABAergic neurons are also more canonical than those in glutamatergic neurons, whereby
most GABAergic neurons in plCoA have interneuron-like identities, expressing canonical marker
genes such as *Vip, Sst,* and *Cck* (**Figure 3H**).

We hypothesized that differences in these populations' abundance could potentially be
responsible for the difference observed between different plCoA domains, and thus examined
potential domain-specific enrichment of certain cell types within plCoA. Visualization of these
nuclei showed little clear region-specific structure in dimension-reduced space for any major cell
types (**Figure 4A**). This lack of structure was broadly confirmed quantitatively, where a few low-
abundance glial cell types showed significant domain-specific enrichment, but the high-
abundance major cell types did not (**Figure 4A-B**). When looking to other regions, however,
heterogeneity is usually more pronounced between subsets of the major cell types, instead of
by the quantities of the major cell types themselves.[42,43] We then examined abundance of
differentially expressed genes (DEGs) between plCoA domains for each major cell type. Here,
we found that both major neuronal cell types had more abundant DEGs than all major glial cell
types (**Figure 4C**). Glutamatergic neurons DEGs exceeded all other major cell types by a factor
of 4, making it likely that differences between the anterior and posterior domains are most likely
to be observed via variation in glutamatergic neurons.

Upon examination of domain-specific variation in plCoA glutamatergic neurons, we
initially observed a greater degree of domain-specific clustering in dimension-reduced space
(**Figure 4D**). Glutamatergic neuron subtypes correspondingly displayed domain-specific
enrichment, where more than half of glutamatergic neuron subtypes were significantly enriched
in either the anterior or posterior plCoA domain (**Figure 4E**). Upon closer examination, we found
that every glutamatergic subtype in the *Slc17a6*-expressing Group 1 was enriched in anterior
plCoA, while *Slc17a7*-expressing Group 2 subtypes are evenly distributed across fields or
biased towards the posterior, with one exception, Glut.Fign, which likely derives from the
aplCoA-adjacent cortex-amygdala transition area CxA, based on Allen ISH data of *Fign*
expression (**Figures 4F, S5A**). In contrast to glutamatergic neurons, we did not observe
significant plCoA domain-specific variation for any GABAergic neuron subtypes (**Figure S5D-E**).
We also found additional heterogeneity within glial cell types, including additional subtypes in

Fig. 4. Glutamatergic neurons subtypes in plCoA are spatially distributed along an
anteroposterior molecular gradient.

(**A**) UMAP of all plCoA nuclei colored by zone of origin, with dotted outlines and labels denoting
the major cell types.
(**B**) Relative proportion of nuclei from each domain within each broad identity class. Dotted line
indicates chance level for all plCoA nuclei.
(**C**) Top, abundance of domain-specific DEGs for each major cell type, either enriched in
aplCoA nuclei (top) or pplCoA nuclei (bottom). Bottom, volcano plots for domain-specific DEGs
for glutamatergic (left) and GABAergic neurons (right), the two cell types with the greatest
degree of domain specific gene expression, where negative log-fold changes indicate
enrichment in pplCoA and positive log-fold changes indicate enrichment in aplCoA.
(**D**) UMAP of plCoA glutamatergic neurons colored by domain of origin, with dotted outlines and
labels denoting the subtypes on the graph. Groups of glutamatergic neuron types identified
previously via Euclidean distance and hierarchical clustering are overlaid on top of the neuron
types of interest.
(**E**) Relative proportion of molecular subtypes from each domain within glutamatergic neurons,
where relevant subtypes are outlined according to their glutamatergic neuron group. Dotted line
indicates chance level for plCoA glutamatergic neuron nuclei.
(**F**) UMAP of all glutamatergic neuron nuclei, colored by expression levels of *Slc17a6* (top) or
Slc17a7 (bottom).
(**G**) Left, representative images of *in situ* RNAscope labeling of *Slc17a6* RNA (red) and *Slc17a7*
RNA (green) across plCoA domains. Right, proportions of glutamatergic neurons expressing
Slc17a6, Slc17a7, or both. Scale bars, 500 µm (main image), 50 µm (inset).
(**H**) UMAP of all plCoA-overlapping Visium capture spots, colored by cluster. Broad spatial
position of groups of clusters are overlaid on top of the capture spots of interest.
(**I**) UMAP of all plCoA-overlapping Visium capture spots, colored by expression levels of
Slc17a6 (top) or *Slc17a7* (bottom).
(**J**) Representative plCoA-overlapping region of one section on a Visium slide capture area, with
capture spots colored by cluster.
(**K**) Representative plCoA-overlapping region of one section on a Visium slide capture area, with
capture spots colored by expression levels of *Slc17a6* (top) or *Slc17a7* (bottom).
(**L**) Prediction scores for representative glutamatergic neuron subtypes within Group 1 (left) and
Group 2 (right), shown on a UMAP of all plCoA-overlapping capture spots across all sections
(top) and on a representative plCoA-overlapping region of one section (bottom).
(**M**) Prediction scores for a representative GABAergic neuron subtype, shown on a UMAP of all
plCoA-overlapping capture spots across all sections (top) and on a representative plCoA-
overlapping region of one section (bottom).
Across panels: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns, not significant. Additional specific
details of statistical tests can be found in Supplemental Table 1.

astrocytes and VLMCs (**Figure S5G-J, O-R**). We also observed high-dimensional domain-
specific divergence in gene expression for OPCs and astrocytes corresponding to domain-
specific DEG differences, though they did not correspond to any observable differences in
subtype enrichment (**Figure S5F, K-N**).

To confirm these findings and validate our snRNA-seq data, we directly examined spatial
RNA expression patterns. First, we used RNAscope labelling to examine the expression of
Slc17a6 and *Slc17a7 in situ* in the plCoA, quantifying the number of nuclei expressing these
genes. We found that anterior plCoA had a much greater proportion of RNAscope-labelled
Slc17a6+ nuclei (88%) than *Slc17a7*+ nuclei (11%) compared to the rest of plCoA, where
Slc17a7+ neurons predominate; posterior plCoA nuclei were almost entirely *Slc17a7*+ (97%)
expressing (**Figure 4G**). It is noteworthy that these numbers were generally consistent with
those in our sequencing data. Next, we analyzed spatial gene expression in the plCoA from an
existing Visium spatial transcriptomics dataset that contained sagittal sections bisecting the
plCoA along the midline (Romero et al., in preparation; **Figure S5S**). We asked if the domain-
specific molecular cell type composition can be recovered directly from spatial information,
without depending on inference from dissection histology. All sections used were of similarly
high quality and did not display any clearly observable batch effects, with all but one having
more than 100 spots covering the plCoA (**Figure S5T-W**). When clustering directly on spatial
data, we observed significant heterogeneity separating into three broad groups (**Figure 4H**).
Like in scRNA-seq, we found highly specific expression of *Slc17a6* and *Slc17a7* to two of the
three broad spot groups (**Figure 4I**). When examining the spatial configuration of these groups,
we found the *Slc17a6*-expressing group of clusters was in aplCoA, while the *Slc17a7*-
expressing group was in pplCoA**,** with the third intermediate group corresponding to layer 1
(**Figure 4J-K**). When computationally projecting transcriptomic cell type identities onto spatial
data, we observed that Group 1 glutamatergic neuron types would project onto pplCoA spots
and Group 2 glutamatergic neuron types would project onto aplCoA spots, while negligible
anteroposterior bias could be observed when projecting GABAergic neuron types onto plCoA
spots (**Figure 4L-M**). Overall, these data demonstrate that plCoA contains an extremely diverse
population of numerous glutamatergic and GABAergic neuron subtypes, whereas glutamatergic
neuron subtypes vary significantly along the anteroposterior axis, such that aplCoA-enriched
subtypes express *Slc17a6* and pplCoA-enriched subtypes express *Slc17a7*.

Molecularly Defined plCoA Glutamatergic Neuron Populations Are Sufficient to Drive
Valence Behavior

Given this spatial distribution bias of plCoA[Slc17a6+] neurons into aplCoA and plCoA[Slc17a7+]
neurons into pplCoA, we further hypothesized that these glutamatergic neuron subtypes could
be responsible for the opposing valence responses observed during topographic plCoA
stimulation (**Figure 2**). If distinct molecular cell types mediate opposing valence, the
topography-independent activation should elicit opposing responses. To investigate this, we
expressed ChR2 in a non-spatially-biased, cell type-specific manner using a Cre-dependent
viral construct in *Slc17a6::Cre* and *Slc17a7::Cre* transgenic mice, targeting AAV-DIO-hSyn-
ChR2 into mplCoA and implanting optic fibers above the injection site (**Figures 5A-B, S2B**).[44,45]
Using the prior four-quadrant open field task, we found that photostimulation of plCoA[Slc17a6+]

Fig. 5. Glutamatergic pICoA$^{Slc17a6+}$ and pICoA$^{Slc17a7+}$ neurons contribute to innate olfactory
behaviors of opposing valence.

(**A**) Schematic for selective photostimulation of distinct glutamatergic cell type. *Slc17a6::Cre* and
Slc17a7::Cre animals were injected with Cre-dependent viral vectors into mpICoA with a fiber
optic implant placed just above the injection site.
(**B**) Representative histology from ChR2 viral injection and fiber implantation site in an
Slc17a6::Cre animal (top) and an *Slc17a7::Cre* animal (bottom).
(**C**) Baseline-normalized collective heatmaps from both the ChR2- and eYFP-infected groups in
Slc17a6::Cre and *Slc17a7::Cre* animals with pICoA-localized fiber implants. Lower right stimulus
quadrant marked in blue.
(**D-G**) Effect of photostimulation of pICoA$^{Slc17a6+}$ neurons (**D-E**) or pICoA$^{Slc17a7+}$ neurons (**F-G**) on
time spent in the stimulation quadrant (**D, F**) and distance from the corner (**E, G**).
(**H**) Behavioral paradigm to assess innate valence responses to odor. Left, schematic of four-
quadrant open field behavioral assay for spatially-specific odor delivery. Upper right, within-trial
timeline. Lower right, odors delivered and their associated innate valence.[3]
(**I**) Schematic of strategy for selective chemoinhibition of molecularly defined glutamatergic
pICoA neurons.
(**J-M**) Effect of chemoinhibition of pICoA $^{Slc17a7+}$ neurons on time spent in the odor quadrant (**J, L**)
or decrease in mean port distance (**K, M**) in response to 2PE (**J-K**) or TMT (**L-M**).
(**N-Q**) Effect of chemoinhibition of pICoA $^{Slc17a6+}$ neurons on time spent in the odor quadrant (**N,**
P) or decrease in mean port distance (**O, Q**) in response to 2PE (**N-O**) or TMT (**P-Q**).
Across panels, ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, **** $p < 0.0001$.
Additional specific details of statistical tests can be found in Supplemental Table 1.

neurons significantly reduced time spent in the 'on' quadrant and increased the average
distance to the corner port during the treatment period , indicating activation of plCoA$^{Slc17a6+}$
neurons is aversive and leads to avoidance of the quadrant when paired with stimulation
(**Figure 5C-E**). In contrast, photostimulation of the plCoA$^{Slc17a7+}$ neurons instead increased the
time in the 'on' quadrant and decreased average distance to the corner port, indicating
plCoA$^{Slc17a7+}$ neuron activation instead is appetitive and leads to attraction to the stimulation
quadrant (**Figure 5C,F-G**). These data indicate that the divergent domain-specific valence
effects of plCoA activity are likely due to the divergent molecularly defined neuronal ensembles
predominant in each topographical field of plCoA.

Next, we sought to determine whether these two glutamatergic populations are
respectively required for innate olfactory aversion and/or attraction. We used the above
transgenic mouse lines to drive expression of a viral Cre-dependent hM4D(Gi) construct to
selectively inhibit these neurons' activity via chemogenetics (**Figure 5I, Figure S6A-C**).[46] We
administered clozapine-N-oxide (CNO) or a vehicle control and used the four-quadrant open
field assay to deliver odorants in a quadrant-specific manner to assess their behavioral
responses to the innately-attractive 2PE or the innately-aversive TMT to determine the
difference in the magnitude of temporally-counterbalanced valence responses when the
respective populations are chemogenetically silenced.[3]

We observed that while both transgenic mouse lines preserved the baseline attraction
and aversion to both 2PE and TMT following administration of the vehicle control, silencing
plCoA$^{Slc17a6+}$ neurons did not affect the response to either odorant (**Figure 5J-Q**). However,
silencing plCoA$^{Slc17a7+}$ neurons abolished the positive valence response to 2PE, without having
any effect on TMT-evoked aversion (**Figure 5J-Q**). In other words, neither group of plCoA
glutamatergic neurons are necessary for TMT aversion, but plCoA$^{Slc17a7+}$ neurons are required
for expression of 2PE attraction behaviors. Further, silencing of either population did not lead to
any broader non-olfactory behavior effects as measured by the EPM and OFT assays, including
anxiety, exploration, and motility, showing the effects of silencing these neurons are likely
limited to valence and/or olfaction alone, instead of exploratory or defensive behaviors (**Figure
S5C-N**). The necessity of plCoA$^{Slc17a7+}$ neurons for 2PE attraction, combined with their ability to
drive positive valence responses indicates that these of plCoA$^{Slc17a7+}$ neurons control innate
olfactory attraction, but plCoA$^{Slc17a6+}$ neurons do not completely control innate olfactory aversion,
instead proving sufficient but not necessary.

A Topographic Organization of plCoA Defined by Limbic Projection Targets

While comprehensively characterizing the circuit features of plCoA, we noted that the
anatomical connectivity of the plCoA has not yet been defined in the mouse brain. Further, we
posited that the differences in necessity of plCoA cell types could be due to divergent
downstream connections instead of divergent molecular features, which may partially, but not
completely overlap. Thus, we next sought to identify distinct downstream outputs of plCoA that
could explain the bidirectional valence effects of its topography. We first characterized the
downstream outputs of plCoA by co-injecting the anterograde viral tracer AAV-DIO-hSyn-
mRuby2-T2A-synaptophysin-EGFP and a constitutive AAV-Cre virus into mplCoA to label

presynaptic terminals with EGFP (**Figure S7A-B**). We observed a high amount of terminal
fluorescence within pICoA itself, suggesting the presence of recurrent connections within the
region (**Figure S6B**). In addition, we observed long-range projections to a diverse set of
regions, including surrounding extended amygdala subregions, such as MeA and the amygdalo-
hippocampal transition area (AHi), regions controlling valence and emotion, like the NAc and the
bed nucleus of the stria terminalis (BNST), and regions involved in olfactory processing,
primarily PIR and OT (**Figure 6A-D**).

540   Among these outputs, we hypothesized that the NAc and the MeA could be responsible
for the behavioral divergence between anterior and posterior pICoA, given their known
involvement in reward expectation and aversion, respectively.[47,48] We sought to confirm these
differences using retrograde tracing from MeA or NAc, where red retrobeads were injected into
MeA or NAc and labeled neurons were quantified along the anterior-posterior axis (**Figure 6E**).
For both downstream targets, we observed opposing gradients of retrobead projector labeling
throughout the entire pICoA anteroposterior axis (**Figure 6F**). MeA-projecting neurons are
enriched in apICoA, and NAc-projecting neurons are enriched in ppICoA, with each having a
frequency of around chance level in mpICoA (**Figure 6G**). Further, the majority of labeled
pICoA-MeA projection neurons were in apICoA, while the majority of labeled pICoA-NAc
projection neurons were in ppICoA (**Figure 6H**). To further confirm the spatial bias in projection
targets, we performed anterograde tracing from the apICoA and ppICoA by injecting viruses
expressing either eYFP or mCherry into either domain of pICoA in a counterbalanced manner
(**Figure S6D-E**). Anterograde projection strength from apICoA and ppICoA revealed that
projections to MeA were most dense from apICoA, and projections to NAc were most dense
from ppICoA (**Figure 6I-K**). Proportionally, apICoA sent a significantly higher proportion of its
projections to MeA than ppICoA, whereas ppICoA sent a significantly higher proportion of its
projections to NAc (**Figure 6L**).

558   We reasoned that the topographical valence sufficiency we observed could be explained
by cell type-specific divergence in projection target, whereby the topographical biases in
downstream targets are recapitulated by their underlying molecular cell type. To determine the
relationship between cell types and projection targets, we injected Cre-dependent eYFP into
mpICoA in *Slc17a6::Cre* and *Slc17a7::Cre* transgenic mice, targeting the middle to ensure
differences result from cell type, instead of simply redundant topography (**Figure S7F-G**).
Interestingly, the relationship was not as simple as one cell type, one primary projection target.
Instead, both cell types project to both structures in different proportions. The pICoA$^{Slc17a6+}$
neurons primarily project to MeA with a significant bias for that target over NAc, while
pICoA$^{Slc17a7+}$ neurons project to both MeA and NAc, with a statistically insignificant bias toward
NAc (**Figure 6M,P**). These findings demonstrate that NAc primarily receives projections
pICoA$^{Slc17a7+}$ neurons, whereas the MeA receives input from both populations. Moreover,
projection of both cell types to the MeA may explain why neither population was selectively
required for innate aversion (**Figure 5J-Q**).

572   Given that both cell types project to both MeA and NAc, we sought to determine the
573   extent of collateralization in neurons composing the two pathways. To test whether pICoA-MeA
574   and pICoA-NAc projection neurons also project to multiple or overlapping downstream targets,

Fig. 6. Projections to MeA and NAc from plCoA are Topographically Organized

(**A**) Right, whole-hemisphere view at AP = 0.98 mm from bregma. Scale bar, 500 µm. Left, Magnified images of the areas highlighted inside white dashed lines. Scale bar, 200 µm.

(**B**) Right, whole-hemisphere view at AP = -1.06 mm from bregma. Scale bar, 500 µm. Left, Magnified images of the areas highlighted inside white dashed lines. Scale bar, 200 µm.

(**C**) Other plCoA projections not found in cross-sections of the brain found in (**A**) and (**B**). Scale bar, 200 µm.

(**D**) Magnitude of anterograde synaptophysin-eYFP fluorescence in primary downstream targets of plCoA projection neurons ordered by total output strength, colored based on each region's function.

(**E**) Schematic for topographic retrograde mapping strategy from MeA and NAc into plCoA. Red retrobeads are injected into MeA or NAc and topographical projection bias is examined along the anterior-posterior axis.

(**F**) Representative images (top) for injection into MeA (left) or NAc (right) and number of neurons labeled along the anterior-posterior axis as distance (mm) from bregma (bottom). Gray lines denote individual replicates, where colored lines indicate mean $\pm$ s.e.m.

(**G**) Proportion of retrobead-labeled neurons projecting to MeA or NAc for each 100 µm segment as a function of distance from bregma. Dashed line indicates overall balance of all retrobead-labeled neurons across entire plCoA.

(**H**) Proportion of retrobead-labeled neurons from either target within each plCoA zone. MeA-labeled neurons are significantly enriched in aplCoA compared to NAc-labeled neurons, while NAc-labeled neurons are significantly enriched in pplCoA compared to those labeled from MeA.

(**I**) Representative histological images for the injection sites in aplCoA (left) and pplCoA (right) from a representative animal. Scale bar, 200 µm.

(**J**) Representative histological images for MeA from the animal in (**J**). Scale bar, 200 µm.

(**K**). Representative histological images for NAc from the animal in (**J**). Scale bar, 200 µm.

(**L**) Output strength as a proportion of total fluorescence from aplCoA and pplCoA to MeA and NAc.

(**M**) Representative histological images for the injection site in plCoA from a representative *Slc17a7::Cre* and *Slc17a6::Cre* animal. Scale bar, 200 µm.

(**N**) Representative histological images from MeA and NAc from a representative animal of either genotype. Scale bar, 200 µm.

(**O**) Left, output strength as a proportion of total fluorescence from plCoA$^{Slc17a6+}$ and plCoA$^{Slc17a7+}$ neurons to MeA and NAc. Right, comparison of same data, but by target region within genotype.

(**P**) Same data as (**O**), but by target region within genotype.

(**Q-V**) Mapping collateral projections from NAc- and MeA projecting neurons.

(**Q**) Representative histological images for the injection site in plCoA from a representative animal receiving retrograde virus into MeA or NAc. Scale bar, 200 µm.

(**R**) Representative histological images of NAc and MeA retro-Cre targeting (red) and outputs (green).

(**S**) Comparison of absolute integrated fluorescence intensities in MeA and NAc when retroAAV was injected into NAc (top) or MeA (bottom).

(**T**) Quantification of fluorescence in selected downstream brain regions from plCoA originating
from plCoA-NAc neurons proportional to eYFP fluorescence in NAc (top) or MeA (bottom).
Abbreviations: NAc, nucleus accumbens; BNST, bed nucleus of stria terminalis; MeA, medial
amygdala; Pir, piriform cortex; BLA, basolateral amygdala; Ahi, amygdalo-hippocampal
transition area; pmCoA, posteromedial cortical amygdala; Str, striatum; OT, olfactory tubercle;
EA, extended amygdala; IPAC, inferior peduncle of the anterior commissure; AA, anterior
amygdala; LA, lateral amygdala; HDB, horizontal limb of the diagonal band; VP, ventral
pallidum; AIC, anterior insular cortex; mfb, medial forebrain bundle; MO, medial orbitofrontal
cortex; LOT, lateral olfactory tract; ACo, anterior cortical amygdala; AOA, anterior olfactory area;
DG, dentate gyrus; Rt, reticular nucleus; LPO, lateral preoptic area; VMH, ventromedial
hypothalamus; DEn, dorsal endopiriform claustrum; LH, lateral hypothalamus; IL, infralimbic
cortex; DP, dorsal peduncular cortex; LS, lateral septum; CxA, cortex-amygdala transition area;
sox, supraoptic decussation; StHy, striohypothalamic nucleus; GP, globus pallidus; PLH,
perirhinal cortex; ZI, zona incerta.
Across panels, ns, not significant; * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$. Additional specific
details of statistical tests can be found in Supplemental Table 1.

we employed a combination of retrograde Cre and Cre-dependent anterograde tracer viral
vectors. A retroAAV-hSyn-Cre-mCherry virus was into either MeA or NAc, and AAV-DIO-ChR2-
eYFP was injected into plCoA to label outputs of MeA- or NAc-projecting neurons (**Figure 6Q-R,**
S7H-I). We focused on MeA and NAc, as well as the ancillary primary downstream targets
implicated in valence or olfaction.[25,49–51] We found different collateralization patterns for both
populations, where NAc-projecting neurons did not collateralize to MeA, but very strongly
collateralized to OT. In contrast, MeA-projecting neurons minimally collateralized to NAc and
most strongly collateralized to pmCoA (**Figure 6S-T**). Notably, neither projection of interest
significantly collateralized to the other. These data indicate that plCoA-MeA and plCoA-NAc
projection neurons are non-overlapping, spatially biased populations that output to different
downstream subnetworks.

Projections from plCoA NAc to MeA and Respectively Control Innate Olfactory Attraction
and Aversion to Odor

The topographic separation of MeA- and NAc-projecting neurons are consistent with a
model of divergence valence that could support the observed topographic divergence in valence
behaviors. To investigate the behavioral contributions of these projections, we first determined
whether the neurons projecting to the MeA and NAc are able to drive behavior with optogenetic
stimulation. We expressed ChR2 in a non-spatially biased manner by injecting AAV-hSyn-ChR2
into mplCoA, and we placed an optic fiber above MeA or NAc for selective optogenetic
stimulation at plCoA axon terminals (**Figure S1C**). We found that photostimulation of the plCoA-
MeA circuit in the four-quadrant open field task significantly reduced time spent in the 'on'
quadrant and increased the average distance to the corner port during the treatment period,
indicating activation of the plCoA-MeA circuit is aversive and leads to avoidance of the quadrant
paired with stimulation (**Figure 7A-D**). The opposite was true for the plCoA-NAc projection,
where stimulation in that zone instead increased the time in the 'on' quadrant and decreased
average distance to the corner port, indicating activation of the plCoA-NAc circuit is instead
appetitive and leads to attraction to the stimulation quadrant (**Figure 7B,E-F**). We next asked if
the effects of stimulating these circuits affected other non-valence behaviors by testing the mice
in the EPM and OFT. Using the EPM, we found no change in anxiety based on open arm time
or entries when stimulating either projection to the MeA or NAc (**Figure S8A-C,E-F,H-I**).
Similarly, stimulation in the OFT did not cause any change to thigmotaxis, based on time spent
in corners of the open field, or exploration, based on time spent in the center of the open field
(**Figure S8K-M,O-P,R-S**). Further, locomotion remained constant during stimulation across
both assays (**Figure S8D, S8G, S8J, S8N, S8Q, S8T**). These data indicate that the divergent
projections from plCoA to the MeA and the NAc evoke valence-specific behaviors without
modulating anxiety.

Finally, we sought to determine whether plCoA-MeA or plCoA-NAc projections are
necessary for the expression of odor-evoked appetitive or aversive behaviors. To target these
projection neurons for chemogenetic silencing, we injected a retroAAV bearing an hSyn-EBFP-
Cre construct into MeA or NAc, along with an AAV in plCoA bearing a Cre-dependent hM4D(Gi)
construct (**Figure 7G,H**). We then tested the innate responses of these animals to 2PE or TMT

Fig. 7. Projections from plCoA to NAc and MeA control innate olfactory attraction and
aversion.

(**A**) Schematic for optogenetic MeA terminal stimulation in plCoA neurons. Strategy to activate
MeA-projecting plCoA neuron terminals via optogenetics (top) and representative histology from
ChR2 viral injection and fiber implantation site (bottom).
(**B**) Baseline-normalized collective heatmaps from both the ChR2- and eYFP-infected plCoA
groups with MeA- and NAc-localized fiber implants. Lower right stimulus quadrant marked in
blue.
(**C-D**) Optogenetic MeA terminal stimulation of plCoA neurons infected with ChR2, but not
eYFP, is sufficient to reduce time spent in the stimulation quadrant (**C**) and increase its average
distance from the port (**D**) during the stimulation period.
(**E-F**) Optogenetic NAc terminal stimulation of plCoA neurons infected with ChR2, but not eYFP
is sufficient to increase time spent in the stimulation quadrant (**E**) and decrease its average
distance from the stimulation port (**F**) during the stimulation period.
(**G**) Viral strategy for selective retrograde chemoinhibition of projection-defined plCoA neurons.
(**H**) Schematic for selective retrograde chemoinhibition of projection-defined plCoA neurons.
(**I-L**) Chemoinhibition of NAc-projecting plCoA neurons significantly eliminates the 2PE-evoked
increase in time spent in the odor quadrant (**I**) and decreases in mean port distance (**J**). The
response to TMT is unaffected in time spent in odor quadrant (**K**) or port distance (**L**).
(**M-P**) Chemoinhibition of MeA-projecting plCoA neurons does not affect 2PE-evoked increase
in time spent in the odor quadrant (**M**) or decrease in mean port distance (**N**) significantly
decreases the TMT-evoked reduction in time spent in the odor quadrant (**O**) or increase in mean
port distance (**P**).
Across panels, ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Additional specific
details of statistical tests can be found in Supplemental Table 1.

in the four-quadrant assay following administration of CNO or a vehicle control, as above.

Inhibition of plCoA-NAc projection neurons abolished innate attraction to 2PE without having

any effect on aversion to TMT (**Figure 7I-L**). Conversely, inhibition of plCoA-MeA projection

neurons had no effect on innate 2PE attraction, but significantly decreased the aversion to TMT

(**Figure 7M-P**). Silencing these neurons had no effect in the EPM and OFT assays, indicating

the effects of silencing these neurons are limited to valence or olfaction, and not anxiety or

exploration (**Figure S8M-X**). Thus, plCoA-MeA projection neurons are necessary and sufficient

for innate olfactory aversion, whereas plCoA-NAc projection neurons are necessary and

sufficient for innate olfactory attraction.

DISCUSSION

Topographic Organization of Valence in plCoA

The neural circuits mediating innate valence responses to odor have not been fully
defined. Here, we have advanced our knowledge of the circuitry underlying innate olfactory
behaviors by defining its activity and organization within plCoA and further extending the innate
olfactory pathway from a third order olfactory brain area to limbic structures. We have identified
a novel functional axis for valence with the plCoA that is defined by histologically and
functionally distinct domains along the anteroposterior axis. We have characterized odor
encoding in plCoA, identifying a sparse population code for identity of an odor, but not its innate
valence. We have also determined the composition of molecular cell types in plCoA and
identified spatially biased populations enriched within each domain, which we find sufficient to
drive their respective domain-specific behaviors, though only partially necessary for their
functions in olfaction. Moreover, we identified the outputs of plCoA and quantitatively
characterized the relative strength of each, as well as how it relates to plCoA topography and
domain-specific molecular cell types, demonstrating that neurons projecting to the MeA and
NAc are topographically and molecularly biased. Finally, we demonstrate that neurons
projecting to the to the NAc and MeA are capable of driving approach and avoidance
responses, and loss of function experiments demonstrate that the neurons projecting to the NAc
or MeA are selectively involved in innate olfactory attraction and avoidance, respectively. Thus,
plCoA is composed of dissociable, spatially segregated ensembles for divergent valence,
defined by their downstream projection target.

Spatial organization for features of sensory stimuli is common in sensory cortex, and has
been previously proposed as fundamental to sensory processing.[52] Visual cortex is
topographically organized by retinotopic locations in space, somatosensory cortex contains a
map of the body, auditory cortex has a crude tonotopic organization by frequency, and gustatory
cortex is segregated by taste qualities.[34,35,53–55] However, topographic organization by
perceptual feature has not been observed in the olfactory system, which has been attributed to
the high dimensional nature of olfactory information.[56,57] Our findings identify a topographically
organized divergence motif for valence in plCoA, where activation of aplCoA evokes aversive
behavioral responses, while activation of pplCoA evokes attractive behavioral responses, with a
graded transition between the two domains. This is consistent with prior studies implicating a
spatial organization to inputs from the OB to plCoA, where glomerulus-specific anterograde
tracing from OB shows specific glomeruli send fibers to invariant, densely clustered,
anatomically distinct locations within plCoA, and retrograde tracing from plCoA shows that
upstream glomeruli are spatially biased within OB and closer examination reveals that OB input
can co-vary with plCoA anteroposterior position.[15] Interestingly, spatial segregation of divergent
features is present elsewhere in olfaction as well, although not by perceptual feature. Olfactory
sensory receptors display stereotyped spatial organization even within zones of the olfactory
epithelium, with a corresponding spatially-stereotyped glomerular topography in OB.[16,24,58–60]
Further, these topographic domains in OB have functional relevance, where region-specific OB
manipulations selectively alter different olfactory valence behaviors.[61–63] However, the piriform
cortex generally lacks apparent spatial organization or spatial patterning for odor reponses.[22] To

our knowledge, the topographical organization of valence observed here in plCoA represents
one of the first descriptions of these spatial patterns occurring in central olfactory areas.

Topographic gradients for valence have been observed in other limbic regions, including
the BLA for overall valence, the medial amygdala for innate social behaviors, *Drosophila*
dopaminergic mushroom body neurons for general valence, and the gustatory insular cortex for
taste valence.[19,34,35,64,65] The gustatory insular cortex is of particular interest, given that it also
primarily encodes taste identity and hedonics in a spatially-nonspecific manner.[31,35,66–69] Our
results extend this phenomenon of spatially segregated valence into both olfaction and into a
less well-described amygdala nucleus, and suggest that it could serve as a potential common
motif within the limbic system to organize motivational information, especially for innate
behaviors, which require stereotyped neurocircuitry. We believe a model where aplCoA and
pplCoA are parts of the same region with similar underlying composition, but with a gradual
change in the factor that defines the valence output for a given part of the gradient would best
explain the underlying gradient-like effect on positive and negative valence. We considered two
such organizational principles underlying this topography: molecular cell types and projection
targets. We found that broad divisions of cell types by Slc17a6 and Slc17a7 did not fully define
valence, though more specific subtypes could be responsible. Our results demonstrate that the
projection targets of plCoA neurons are a fundamental feature for imparting valence on this
circuitry.

Molecular Cell Types in plCoA Segregate Topographically and Drive Behaviors of
Divergent Valence

While examining the cell types composing plCoA, we noticed numerous notable, novel
features. First, despite its small area (~1.04 mm^3 and ~170,000 cells), plCoA displays
remarkable diversity, hosting dozens of distinct, robustly separable cell types.[70] The plCoA
appears to have multiple domains positioned at the transition between highly dissimilar brain
tissue types. Interestingly, the high-dimensional structure of molecular variation differs between
glutamatergic and GABAergic neurons. Glutamatergic molecular variation within plCoA is
continuous, with two broadly nested groups generally marked by either *Slc17a6* or *Slc17a7*
along with one or more additional marker gene(s)*,* though it should be noted that expression of
the two broad glutamatergic markers is not necessarily mutually exclusive and a few low
abundance "transition" cell types can express both. This leaves open the possibility that a more
specific cell type could be necessary for aversion, given that this necessity for this behavior did
not perfectly map onto either broad molecular cell type. In contrast, molecular variation in
GABAergic neurons is far more discretized, with ensembles expressing one of a few well-
characterized interneuron markers found throughout the brain, such as *Sst, Vip,* and *Pvalb*,
among others. This is consistent with other studies in neocortex, hippocampus, and subiculum
that find similar patterns of variation, where variation within glutamatergic neurons is more
continuous than in GABAergic neurons.[42,43,71] In this way, we find the continuous gradient-like
structure of valence in plCoA is recapitulated with gradient-like variation in glutamatergic neuron
gene expression in plCoA.

In these molecular datasets, we further observed highly specific differences in cell type
enrichment within aplCoA and pplCoA within glutamatergic neurons, but not GABAergic
neurons or glia. Within plCoA, *Slc17a6+* neurons are enriched in aplCoA and *Slc17a7+* neurons
are enriched in pplCoA, though there is gradient-like intermingling of populations, especially
toward the middle of plCoA, and all glutamatergic neuron types are present, albeit with high
variability along the anteroposterior axis. This molecularly defined order suggests a
programmed organization, rather than stochastically distributed populations within the region,
especially given that its boundaries match the domains previously identified based on behavior
and histology.[72] This phenomenon also broadly matches observations in the neocortex,
hippocampus, and subiculum, where glutamatergic neurons across subdivisions molecularly
diverge to a greater degree than GABAergic neurons or glia, albeit across a correspondingly
greater area than within plCoA, which is generally accompanied by distinct morphological and
electrophysiological properties broadly corresponding to these transcriptomic
differences.[42,43,71,73–76] Spatial segregation of molecular cell types is also observed within deeper
brain regions including BLA, thalamus and habenula, and these molecular differences are also
accompanied by extended phenotypic differences.[19,77–80] Investigation of such properties held in
common and diverging within and between *Slc17a6+* and *Slc17a7+* glutamatergic neuron types
could also serve to further define the local neurocircuitry and information processing dynamics
within plCoA and along its anteroposterior axis.

Interestingly, few if any populations within plCoA are clearly separable from most of the
regions surrounding it (e.g. piriform cortex, basomedial amygdala, and MeA) based on primary
marker gene identity. Instead, plCoA ensembles seem to be defined by the interplay of all three
regions within the same tissue. The predominance of populations resembling different regions
does appear related to this anteroposterior organization, though, where *Slc17a6+* neurons
predominate in both aplCoA and MeA, and more specific marker genes like *Meis2* are
expressed in both regions as well.[81] Conversely, *Slc17a7+* neurons predominate in both pplCoA
and piriform cortex, and the major marker genes like *Satb2* are similarly expressed in both
regions.[82] Such phenomena also cohere with more general characterizations made in whole-
brain molecular taxonomies, which divide plCoA along its axis, grouping aplCoA with MeA and
pplCoA with paleocortex.[83–85] It would be misleading to characterize plCoA populations as mere
extensions of surrounding populations into an adjacent region, though. *Slc17a7+* neurons from
the cortex-amygdala transition zone are also present in the dataset and are continuously
separable from *Slc17a7+* plCoA neurons based on the expression of marker genes like *Fign*.
Instead, plCoA may itself be a transition region, given that such a relationship with its
neighboring regions is very similar to that of the amygdalostriatal transition area, one of the only
transition regions to undergo high-resolution molecular profiling.[86] Given such commonalities
between these two putatively dissimilar regions, molecular characterization of additional
transition regions could potentially uncover similar organizational motifs, especially if compared
with adjacent regions of interest, and allow for a much more in-depth exploration and
characterization of the boundaries and transitions between proximally located, distantly related
brain regions.

These broad molecular groups of glutamatergic cell types themselves do not completely
explain valence in the plCoA. While aplCoA-enriched *Slc17a6+* neurons are sufficient to drive

aversion and pplCoA-enriched *Slc17a7+* neurons are sufficient to drive attraction, as would be
predicted from the valence responses evoked from each anterior-posterior domain enrichments,
it might be expected that these populations would also be necessary for the respective odor-
evoked valence. However, only *Slc17a7+* neurons are necessary for 2PE attraction, whereas
the *Slc17a6+* neurons were not required for aversion to TMT. Given that the plCoA is necessary
for TMT aversion, it is unlikely that such a difference is due to additional redundant function
within other regions for TMT aversion.[3] Instead, although *Slc17a6+* neurons likely contribute to
the valence behavior, other populations within the region not expressing the marker gene could
also contribute to the behavioral response. Thus, although these two broad glutamatergic
groups can drive innate responses of valence, the *Slc17a6+* population doesn't fully represent
the population that control aversive responses, which is supported by other findings presented
here showing that both *Slc17a6+* and *Slc17a7+* neurons project to MeA, while the projection to
NAc is almost entirely composed of *Slc17a7+* neurons. Regarding more specific molecular cell
types, it could be possible that only a subset of *Slc17a7+* neurons are required for 2PE
attraction. Similarly, the neurons required for TMT aversion could be marked by a gene
orthogonal to the observed *Slc17a6/Slc17a7* gradient, and genetic access to olfactory aversion
could potentially be established by investigating these more sparsely expressed marker genes.
In both cases, though, further investigation into these other cell types would enhance our
understanding of both plCoA and innate olfactory valence and allow more precise manipulations
in the future.

Downstream Projection Targets of plCoA Divergently Control Innate Olfactory Valence

856   In this work, we perform the first characterization of plCoA's downstream outputs in the
mouse brain. These outputs are dominated by regions generally involved in valence and
emotion, such as the NAc, BNST, MeA, BLA, and other amygdalar nuclei, or olfactory areas,
such as the pmCoA, OT, and PIR. These outputs are consistent with a role for plCoA in
motivational valence for odor. The plCoA also appears to form numerous intra-regional
connections, where a significant proportion of synapses formed with other neurons are within
the region itself. This raises the possibility that plCoA is not simply a feedforward relay but
performs local recurrent processing as well. Recurrent networks in other sensory systems
expand the dimensionality of encoding schemata and incorporate additional features to
generate mixed, continuously updating representations of relevant information.[87] This raises a
number of interesting questions regarding information processing within plCoA. For instance,
how does odor representation change based on differences in experience and internal state?
Further explorations of information transformation and encoding within plCoA will certainly
enrich our understanding of both this region and olfaction in general.

870   The NAc and MeA are interesting downstream targets given their pre-existing known
relationships to attraction and aversion, respectively. NAc has historically been critical to the
manifestation and processing of reward and motivated behaviors, though this view has been
expanded and made more subtle with a recent focus on action selection in general.[88] On the
other hand, MeA has been linked to defensive and stress-related behaviors at baseline and in
response to aversive stimuli, as well as chemosensation.[89] MeA has also been specifically
linked to olfactory aversion in past studies, as TMT has previously been shown to activate the

MeA, which is necessary for TMT-induced defensive behaviors, though the upstream circuits
and processing were not yet investigated.[48,90,91] These circuits are also notable regarding other
features of plCoA spatiomolecular organization, as the projections to the downstream regions of
interest are the two that diverge to the greatest extent between aplCoA and pplCoA, and
between plCoA$^{Slc17a6+}$ and plCoA$^{Slc17a7+}$ glutamatergic neurons. Given the relationship between
spatiomolecular patterning and the simple wiring and organizational rules used to structure
innate circuits it would be interesting to investigate the plCoA through the lenses of development
and genetic variation. These intersect in recent discussions of genetic bottlenecking, where the
genome encodes general rules for circuit organization and development that nevertheless yield
specific responses to specific stimuli.[92] Such networks have numerous theoretical advantages,
such as reduced information requirements and higher performance at criterion, providing a
conceptual basis for why spatiomolecularly stereotyped circuits yield innate behaviors, and why
these innate behaviors are adaptive in naturalistic settings.[92–94]

Neuronal Activity in plCoA Encodes Odor Identity Via Sparse Population Code

The plCoA receives spatially-ordered inputs and has a cell-type-specific topographical
organization with divergent outputs that mediate responses of opposing valence. This
anatomical organization is consistent with either labeled-line or divergent paths motifs, though
precise connectivity from OB has not yet been described. However, label line coding motif is not
apparent in neural activity, but instead there is a sparse population code for odor identity with no
apparent valence-specific responses, indicating that the plCoA cannot function as a pure
labeled-line relay. Moreover, the absence of valence encoding is distinct from that seen in other
divergent path motifs, suggesting the plCoA represents a different circuit model for valence.

We were surprised to find that despite a robust organization of plCoA cell types and
projections that support behaviors of valence, the neurons do not appear to encode valence at
the single neuron or population level. However, when put in the context of the olfactory system,
these results are less surprising. Sparse, distributed population encoding appears to be a
general feature of all olfactory regions observed thus far, regardless of their specific structure or
computational function within olfaction. In the OE, epithelium-resident OSNs expressing a single
receptor will bind multiple odorants and establish odor identity through a combinatorial code.[21,23]
In the OB, axon terminals from OSNs expressing the same receptor gather into specific
stereotyped glomeruli, where mitral/tufted cells (M/T) then transmit this information to third-order
olfactory regions, where both steps represent the odor identity via a sparse population code in a
spatially-distributed manner.[24,26,32,95] The primary olfactory cortex further represents odor identity
and chemical space similarly through sparse, spatially distributed combinatorial population
activity in a similar manner to its M/T inputs.[17,22,32,96] The OT, a striatal region primarily
composed of *Drd1*- or *Drd2*-expressing medium spiny neurons, also broadly represents odor
identity across both cell types, despite their opposing roles elsewhere in striatum, though a
secondary contribution by hedonic value is currently debated.[25,97–101] Although direct
comparisons of olfactory regions find some differences in features like sparsity, clustering, or
correlation structure, these are generally reported as differences in degree, not in
kind.[17,30,32,95,99,100] Given the remarkable similarity in encoding between these structurally and
functionally distinct regions, it logically follows that an olfactory region with amygdalar structure

would follow similar population encoding principles. Indeed, were plCoA to predominantly
encode stimulus valence in a manner akin to other amygdalar regions, it would reflect a greater
divergence from olfactory coding principles than what we observe. Instead, the olfactory
encoding scheme employed by plCoA reveals a novel framework for valence encoding that is
distinct from that in the extended amygdala.

How does innate valence arise from a population code imposed on stereotyped valence
circuitry? One possibility is that the valence of an odor is defined by the proportion of NAc and
MeA-projecting neurons activated, mixing features of divergent paths and opposing components
motifs. In this model a given odor will activate some balance of NAc and MeA projecting
neurons and the proportional balance determines the valence. If true, one should be able to
predict the valence of the odor by recording from a large sample of these neurons, which is
beyond the scope of this study. Another possibility is that activity could evolve over the time
course of behavior such that a mixture of inputs from local circuitry and long-range circuits could
function like attractor networks to shape output into one of a few convergent states. A third
possibility is that valence may emerge in combination with behavior and internal state. In this
model, the plCoA integrates sensory information with other state variables to shift the activity
towards one output. If either of these latter two models is correct, recording activity over longer
time scales in freely moving animals could provide insight into how activity correlates with
behavior. Alternatively, valence may arise downstream of plCoA, but molecular mechanisms to
support this model are currently undescribed, and such a model would be at odds with our
finding of divergent valence-specific circuitry required for innate valence responses. Innate
behavior is indeed dynamic and more complex than attraction and avoidance. These different
models for valence encoding could serve to increase the flexibility of the system to adapt to
changes in experience or internal state.

A Novel Circuit Motif for Valence

The widespread circuit model for valence encoding centers on divergence, in which
information is routed to different pathways depending on the valence of the stimulus. The plCoA
circuit we have identified is anatomically similar, but unique in its encoding properties. Most
divergent circuits function to generalize stimuli, discarding information about stimulus identity to
simplify to low dimensional valence signals from distinct neuronal populations. In contrast, the
plCoA groups stimuli, using a high dimensional code of odor identity that is routed through
divergent projections to mediate opposing valence responses. In this model, the valence of an
odor could be determined based on the population dynamics and composition within a
distributed, sparse population code that ultimately funnels information through divergent
pathways corresponding to their innate valence. Such a model could serve to increase the
flexibility of the system while retaining the ability to yield stereotyped responses.

ACKNOWLEDGMENTS

We thank the entire Root lab for helpful discussion and support. We thank B. Lim for provision of the AAVDJ-hSyn-Flex-mRuby-T2A-SynEGFP virus. We thank G. Pekkurnaz, N. Spitzer, and M. Pratelli for reagents and facilities support. We also thank C. O'Connor and L. Boggeman at the Salk flow cytometry core, and N. Hah at the Salk next-generation sequencing core. We thank D. Jimenez for assistance with histology and colony management. This publication includes data generated at the UC San Diego IGM Genomics Center utilizing an Illumina NovaSeq 6000 that was purchased with funding from a National Institutes of Health SIG grant (#S10 OD026929). This work was generously supported by the National Defense Science and Engineering Fellowship (J.R.H.), CIHR Postdoctoral Fellowship (F.M.), JPB Foundation, the PIIF, PNDRF, JFDP, New York Stem Cell Foundation, Klingenstein Foundation, McKnight Foundation, Salk Institute, Howard Hughes Medical Institute, Clayton Foundation, Kavli Foundation, Dolby Family Fund (K.M.T.), the Hellman Fellowship (C.M.R.), and the National Institutions of Health, via grants through the NIA (RF1AG061831-01S1, P.A.D.), NIMH (K99MH121563-02, F.M.; R01MH115920-03 and R37MH102441, K.M.T.), NIDDK (DP2DK102256-01S1, K.M.T.), and NCCIH (DP1AT009925-04, K.M.T.), and the NIDCD (R00DC014516-05 and R01DC018313-01A1, C.M.R.).

AUTHOR CONTRIBUTIONS

J.R.H. and C.M.R. conceived the project. J.R.H., C.L.C., D.L., M.B., and A.N.Z. performed surgeries and analyzed histological data. J.R.H., C.L.C., and A.N.Z. performed behavioral experiments, managed the colony, and analyzed behavioral data. J.R.H. and D.L. performed calcium imaging experiments and analyzed calcium imaging data. J.R.H. performed sequencing experiments, analyzed sequencing data, and prepared all figures. H.K.R. and P.A.D. designed and performed spatial transcriptomics experiments and provided data for analysis. F.M., M.E.L., and K.M.T. provided the CellProfiler processing pipeline, as well as intellectual, facilities, and logistic support and guidance. C.M.R. supervised the project. J.R.H., D.L., M.B., H.K.R., and C.M.R. wrote the manuscript. All authors edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Cory M. Root (cmroot@ucsd.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All snRNA-seq and Visium spatial sequencing data generated for this study is deposited on
Gene Expression Omnibus (GEO): GSE270798. All other data reported in this paper will be
shared by the lead contact upon request. The code used to analyze all data and generate all
graphs can be found online at Zenodo: . Any additional information required to reanalyze the
data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All procedures at the University of California, San Diego, and Columbia University were
performed in accordance with Institutional Animal Care and Use Committee protocols in
accordance with NIH guidelines. All mice were provided food and water ad libitum and
maintained on a regular 12-hour reverse light/dark cycle at room temperature, with weight,
health, and immune status monitored daily and verified to be within normal ranges. Mouse
cages were changed regularly based on degree of soiling. All mice were group-housed with
randomly assigned littermates prior to surgery, and single-housed after surgery. All animals
were used in a single experiment each, except for a subset of mice who underwent 4-quad,
EPM, and OFT experiments, who performed each test in no specific order.

Subject Details for Sequencing Experiments

Subject details for single-nucleus sequencing

All mice for snRNA-seq in the study were males on the wild-type C57BL/6J background
(RRID:IMSR_JAX:000664) and received directly from Jackson Laboratories at 6 weeks of age
and acclimated to the colony prior to experiments. Animals were single-housed and placed into
sensory deprivation 24 hours prior to sacrifice to reduce artifactual immediate early gene
expression. Sacrifice was performed at P60 ± 3 days (n = 5-10 mice per pool). Sample size was
determined based on number of expected nuclei per region per mouse: estimates of expected
nuclei were determined empirically, though nuclear recovery was approximately 20% of total
based on cellular density estimates from the Blue Brain Cell Atlas.[70] 20,000 nuclei were targeted
per combination of assay, condition, and region, which was determined using SCOPIT v1.1.4,
allowing potential detection of at least 10 nuclei from 10 rare subpopulations at 0.1% frequency
with 95% probability.[103] A total of 50 mice were used for this purpose.

Subject details for spatial transcriptomics

In the separate study for spatial transcriptomics, APP23 (B6.Cg-Tg(Thy1-APP)3Somm/J;
C57BL/6J background, RRID:IMSR_JAX:030504) non-transgenic (NTG) littermates control mice
were housed in light-tight enclosures.[104] The mice were given ad libitum food and water access.
This study used a total of 17 mice almost equally distributed across sex, of which sections from
11 sagitally bisected the plCoA and were used in downstream analysis. No analysis of sex
differences was performed due to inaccessibility of that information on a per-section basis.

However, no such differences were apparent from per-section gene expression correlations
reported in supplementary information.

Subject Details for Calcium Imaging, Tracing, and Activity Manipulations

Subject details for wild-type experiments

All mice for topographic and projection-defined manipulation and tracing experiments, as well as
calcium imaging, were males on the wild-type C57BL/6J background and received directly from
Jackson laboratories before 12 weeks of age. After surgery, mice incubated for at least 21 days
if injected with virus or at least 7 days if injected with a retrograde tracer (e.g. retrobeads,
cholera toxin B) to allow virus to express and tracers to travel in retrograde, respectively. All
surgeries and downstream experiments were performed on mice at least 8 weeks of age.

Subject details for transgenic experiments

We used VGlut2-IRES-Cre (Slc17a6tm2(cre)Lowl; C57BL/6J background,
RRID:IMSR_JAX:028863; *Slc17a6::Cre*) and VGlut1-IRES2-Cre-D (Slc17a7tm1.1(cre)Hze;
C57BL/6J background, RRID:IMSR_JAX:023527; *Slc17a7::Cre*) strain mice for molecularly
defined optogenetic stimulation experiments and genotype-specific tracing.[44,45] These mice
were bred on-site at UCSD and were genotyped in-house using genomic DNA from ear tissue
amplified with the default primer sets listed by Jackson Laboratories. All mice used for
experiments had a heterozygous genotype for the transgenic construct of interest. After surgery,
mice incubated for at least 21 days to allow viral expression.

METHODS DETAILS

Stereotactic surgery procedures

All surgeries were performed under aseptic conditions using a model 1900 digital small animal
stereotaxic instrument (Kopf Instruments). Mice were initially anesthetized in a sealed box
containing 5% gaseous isoflurane, and then deeply anesthetized using isoflurane (2.5% in
1L/min of O_2) during surgeries (VetFlo, Kent Scientific Corporation). We immobilized and leveled
the head in a stereotaxic apparatus (Kopf Instruments), removed fur from the scalp by shaving,
applied eye lubricant (Optixcare), cleaned the incision site with 70% ethanol and betadine prior
to incision, peeled off connective tissue, and dried the surface of the skull prior to craniotomy
before proceeding with injections and implantations specific to certain experiments. All virus
injections were performed at 2 nL/sec using a pulled glass pipette (Drummond) and a Nanoject
III pressure injector (Drummond). To prevent backflow, the pipette was left in the brain for 15
minutes after injection.

Surgeries for calcium imaging

Surgeries for imaging experiments were performed in a manner similar to that previously
described.[33] The skull was prepared with OptiBond[TM] XTR primer and adhesive (KaVo Kerr)

prior to the craniotomy. After performing a craniotomy 1 mm in diameter centered around the
virus injection site, a 27G blunt needle was used to aspirate ~2.5 mm below the brain surface.
600 ul of AAV9-hSyn-FLEX-jGCaMP8s-WPRE (2.5×10^{13} gc/ml, Addgene) was diluted to ~1 ×
10^{12} gc/ml gc/ml and injected into the left middle plCoA in two 300 ul boli, one in layer 2 and one
400 um dorsal (+0.4 DV) in layer 3. Following the viral injection, a head-plate (Model 4,
Neurotar) was secured to the mouse's skull using light-curing glue (Tetric Evoflow, Ivoclar
Group). At least 30 minutes after viral injection, a 1 mm GRIN lens (NA, ~1.9 pitch, GrinTech)
was sterilized with Peridox-RTU then slowly lowered at a rate of 500 um/min into the craniotomy
until it was 200 um dorsal (+0.2 DV) to the injection coordinate. The lens was adhered to the
surface of the skull using Tetric Evoflow. We then placed a hollow threaded post (AE825ES,
Thorlabs) to act as a housing for the lens and adhered it using Tetric Evoflow. Any part of the
skull that was still visible was covered using dental cement (Lang Dental). Finally, the housing
was covered with a Nylon cap nut (94922A325, McMaster-Carr) screwed onto the thread post to
protect the lens in between imaging. Animals were left on the heating pad until they fully
recovered from anesthesia.

Surgeries for optogenetic stimulation

For optogenetic topographic- or projection-specific stimulation experiments, we injected wild-
type C57BL/6J mice between 2-4 months of age with 100 (if topographic) or 150 (if projection-
specific) nL of either AAV5-hSyn-eYFP (3.3×10^{12} gc/ml, UNC Vector Core) or AAV5-hSyn-
ChR2-mCherry-WPRE-PA (4.1×10^{12} gc/ml, UNC Vector Core). For Cre-dependent molecularly
defined optogenetic stimulation, we injected 200 nl AAV5-EF1A-DIO-hChR2(H134R)-eYFP (5.5
$\times 10^{12}$ gc/ml, UNC Vector Core) or AAV5-EF1A-DIO-eYFP (4.0×10^{12} gc/ml, UNC Vector Core)
All such injections were left unilateral, into either aplCoA or pplCoA for topographic
photostimulation, and into middle plCoA for projection- or genotype-specific photostimulation.
For topographic and genotype-specific photostimulation animals, we implanted the fiber 300 um
(+0.3 DV) directly above the injection site with all other coordinates remaining constant.
Anterior-posterior axis positioning arose from stochastic variation in virus and fiber placement.
For projection-specific photostimulation animals, we implanted the fiber 300 um (+0.3 DV)
directly above either the MeA or NAc, holding all other coordinates for the two regions constant
as described above. For all optogenetic stimulation experiments, we implanted a fiber optic
cannula (2.5 mm ferrule outer diameter, 200 um core, 0.39 numerical aperture; RWD) 300 um
above the targeted stimulation site. These fibers were affixed onto the skull using OptiBond XTR
(Kerr) and stably secured with Tetric Evoflow (Ivoclar Vivadent) coated with cyanoacrylate
(Toagosei). After surgery, we injected all mice with 0.04 mL Buprenorphine SR (Ethiqa XR,
Fidelis) for pain management. All mice were given their own cage immediately after surgery and
returned to the colony once ambulatory.

Surgeries for chemogenetic inhibition

For all plCoA inhibition experiments, we bilaterally injected C57BL/6J mice between 2-4 months
of age with 250 nL of either AAV2-hSyn-DIO-hM4D(Gi)-mCherry (7.1×10^{12} gc/ml, Addgene) or
AAV2-hSyn-mCherry (1.8×10^{13} gc/ml, Addgene) virus. For projection-specific inhibition
experiments, wild-type C57BL6/J mice were used, and AAVretro-hSyn-EBFP-Cre (1.5×10^{12}

gc/ml, Addgene) either 50 µL were injected into MeA or 300 µL were injected into NAc. For

genotype specific inhibition experiments, *Slc17a6::Cre* or *Slc17a7::Cre* mice were used. All

injections were bilateral and targeted to middle plCoA.

Surgeries for fluorescent tracing

For non-topographic anterograde tracing experiments, we bilaterally co-injected of mixed 50 nl

AAVDJ-hSyn-FLEX-mRuby-T2A-SynEGFP (4.0 × 10^{12} gc/ml, Addgene, a gift from Byungkook

Lim) and AAV5-EF1A-mCherry-IRES-Cre-WPRE (1.9 × 10^{12} gc/ml, UNC Vector Core) into

middle plCoA (-1.8 AP, +/-2.9 ML, -5.95 DV).[105,106] For topographic anterograde tracing

experiments, we unilaterally injected 20 nl AAV8-hSyn-hChR2(H134R)-mCherry (2.1 × 10^{13}

gc/ml, Salk GT3 Viral Vector Core) and AAV8-hSyn-hChR2(H134R)-eYFP (3.2 × 10^{13} gc/ml,

Salk GT3 Viral Vector Core) into aplCoA (-1.4 AP, -2.8 ML, -5.95 DV) and pplCoA (-2.1 AP, -3.0

ML, -5.95 DV), counterbalancing region by fluorophore. For retrograde topographic tracing

experiments, we unilaterally injected Red Retrobeads IX (Lumafluor) into either MeA (50 nl, -1.2

AP, -2.0 ML, -5.5 DV) or NAc (+1.1 AP, -1.35 ML, -4.5 DV), at volumes of 50 nl or 300 nl,

respectively. For anterograde collateralization experiments, AAVretro-EF1A-IRES-Cre (1.3 ×

10^{13} gc/ml, Addgene) into either MeA or NAc, and AAVDJ-EF1A-DIO-hChR2(H134R)-eYFP-

WPRE-pA (4.03 × 10^{13} gc/ml, Salk GT3 Viral Vector Core) was injected into middle plCoA. For

genotype-specific anterograde tracing experiments we injected 50 µl AAVDJ-Ef1a-DIO-

ChR2(H134R)-eYFP-WPRE-pA into middle plCoA in Slc17a6::Cre or Slc17a7::Cre mice.

Calcium Imaging

Odor exposure

Odor exposure for imaging experiments was adapted from methods previously described

elsewhere.[25] Odor was delivered to the mouse using a custom-built olfactometer. Compressed

medical air was split into 2 gas-mass flow controllers (Aalborg). One flow controller directed a

constant rate of 1.5 L/min to a hollowed out teflon cylinder. The other flow regulator was

connected to a 3-way solenoid valve (The Lee Co.). Prior to odor delivery, the 3-way valve

directs clean air at 0.5 L/min to the teflon cylinder. During odor delivery, the 3-way valve directs

air to an odor manifold, which consists of an array of 2-way solenoid valves (The Lee Co.), each

connected to a different odor bottle. The kinetics and consistency of odor delivery were

characterized using a miniature Photoionization Detector (mPID) (Aurora Scientific). Depending

on the trial type, the appropriate 2-way valve opens, directing 0.5 L/min of air flow through the

odor bottle containing a kimwipe blotted with 20 ul of odorant, except for 2PE and peanut oil,

which were undetectable to human perception and the mPID after repeated presentation, and

were provided in excess to ensure consistent detectability after numerous exposures.

Prior to imaging, mice were habituated to the head fixation device (Neurotar) and treadmill for at

least 3 days for at least 5 minutes per session beginning at least 10 weeks after surgery. The

treadmill parts were 3D printed using an LCD printer (EPAX) from publicly available designs.[116]

Walking behaviors were measured using a quadrature encoder (Broadcom). A video feed of the

animal's face was also recorded using a camera (Basler) with an 8-50mm zoom lens (Arducam)

at 20 Hz with infrared illumination (Lorex Technology). Animals were exposed to the following
odors for 2 seconds: the appetitive odors 2-phenylethanol (Sigma-Aldrich) and peanut oil
(Spectrum), the neutral odors 1-heptanol (Sigma-Aldrich) and isoamyl acetate (Sigma-Aldrich),
and the aversive odors trimethylthiazoline (BioSRQ) and 4-methylthiazole (Sigma-Aldrich).
Within a single contiguous exposure session, each of the 6 odors were provided 20 times with
12-18 seconds of inter-trial interval. Trials were organized into 6 blocks, each of which consisted
of 20 trials of each of the 6 odors in counterbalanced order without any odors of similar innate
valence adjacent to each other in the trial structure.

2-photon microscope data acquisition

Ca^{2+} imaging data was acquired using an Olympus FV-MPE-RS Multiphoton microscope with
Spectra Physics MaiTai HPDS laser, tuned to 920 nm with 100 fs pulse width at 80 MHz. Each
128 × 128 pixel scan was acquired with a 20x air objective (LCPLN20XIR, Olympus), using a
Galvo-Galvo scanner at 5 Hz. Stimulus delivery and behavioral measurements were controlled
through a custom software written in LabVIEW (National Instruments) and operated through a
DAQ (National Instruments). Each imaging session lasted up to 90 minutes and was
synchronized with the stimulus delivery software through a TTL pulse. Animals were excluded
from analysis if histology showed that either the GRIN lens or the jGCaMP8s virus was
mistargeted or the motion during imaging was too severe for successful motion correction.

Behavioral Assays

Mice had been handled for 5 days prior to experiments and acclimated to the room for an hour
prior to testing. All behavioral experiments were performed during the dark period of the
light/dark cycle at least an hour away from the switch between the two photoperiods. Not all
mice were run in all assays, as elevated plus maze and open field tests were added after a
significant proportion of four-quadrant data was collected at targeted sample sizes and mice had
already been sacrificed.

For all optogenetic experiments, optical fibers (200 mm, 0.39 numerical aperture, Thorlabs)
were epoxied to 2.5 mm stainless steel ferrules (Precision Fibre Products), and polished with a
fiber optic polishing kit (Thorlabs) to achieve a minimum of 80% transmission. After surgical
implantation, the ferrules protruding from the mouse's head were coupled to an ADR-800A 100
mW 473 nm laser (LaserCentury) via custom-made patch cables with a single rotary joint (Doric
Lenses) between the mouse and laser. Laser intensity was set to 5-8 mW at the end of the
patch cable. For inhibition experiments, all mice were injected intraperitoneally 60 minutes prior
to the beginning of the behavioral trial with either sterile PBS vehicle or with clozapine-N-oxide
(CNO) dihydrochloride (Hello Bio) dissolved in sterile PBS for a dosage of 2 mg/kg.

Four quadrant open field assay

The four-quadrant open-field task was performed as previously described.[3] In short, all
behavioral assays took place in a four-quadrant open field chamber. Airflow was pumped into
each quadrant via gas-mass flow controllers 150 mL/min (Cole-Parmer). Airflow exited the

chamber via a 1-inch outlet in the center of the chamber's floor covered by steel mesh, and the
outlet was connected a vacuum line with a gas-mass controller set to 750 ml/min. The chamber
was housed in the dark and illuminated from below by infrared lighting. A Basler A601FM
camera (Edmund Optics) mounted above the chamber recorded videos of behavioral trials at 4
Hz, and custom software written in Labview (National Instruments) tracked the position of the
mouse in real time for each frame. The symmetrical four-quadrant open field chamber was
contained in a lightproof structure (0-10 lux) and illuminated by infrared lights, removing any
potential spatial cues available to the animals with respect to the room or its surroundings. In
optogenetic experiments, an additional 5 cm spacer was added to the chamber flush with the
walls to raise their height for more naturalistic behavior, and an acrylic ceiling with a top with a
circular opening 30.5 cm in diameter was added to prevent escape while allowing the fiber optic
cable to move freely.

In optogenetic experiments, the laser was pulsed with 50 ms bins at 10 Hz, and there was a
steep gradient from 1 to 10 Hz along the perimeter of the quadrant, increasing as proximity to
the corner of the quadrant decreased. Preliminary experiments in topographical stimulation
animals identified no clear behavioral effects from the location of the stimulus quadrant itself
(data not shown), and all other downstream stimulation experiments exclusively used the lower
right quadrant for stimulation to simplify data analysis. The lasers were controlled by TTL
modulation from custom Labview software synchronized to the video capture system.

For inhibition experiments, odor was applied by solenoid valves redirecting airflow through 100
mL glass bottles containing 1 µL of a pure odorant on a small piece of Kimwipes. Odorants used
were either the previously-validated innately aversive 2,5-dihydro-2,4,5-trimethylthiazoline
(BioSRQ) or the innately appetitive 2-phenylethanol (Sigma-Aldrich) on a small piece of
Kimwipe.[3] All odors were presented in the lower-right quadrant, and all trials were spaced out
with at least an hour between runs, during which vacuum was applied to the chamber. Odors
and injection treatments were given in counterbalanced, independent order within experimental
groups.

Open field test

The open field is a square arena illuminated to 100-150 lux by ambient lighting. Mice were
habituated to the room for at least an hour prior to testing, but otherwise had no prior experience
in the arena prior to exposure. Mice were placed in the center of a square arena (27.3 × 27.3 ×
20.3 cm, Med Associates) with four transparent plexiglass walls. Overall locomotion, immobility,
and time spent in corners and center regions of arena during each epoch was analyzed for each
mouse. Immobility was defined as movement under 0.5 cm/s for a period of at least 1 s, while
the center was defined as the middle 13.7 × 13.7 cm square in the center of the arena and the
corners as the corner regions that do not overlap with the center square in either direction (25%
of arena area for each region). For optogenetic experiments, mice were allowed to move freely
throughout the arena for 25 min total, with 5-8 mW 473 nm light stimulation pulsed with 50 ms
bins at 20 Hz, alternately delivered during the 5-10 min and 15-20 min epochs (OFF, ON, OFF,
ON, OFF). For chemogenetic experiments, mice were allowed to freely move through the area
for 10 min total.

Elevated plus maze

The arms of the elevated plus maze were 30.5 × 5.5 cm. The height of the closed arm walls was
15 cm. The maze was elevated 40 cm from the floor and was placed in the center of the
behavior room away from other stimuli. Arms were illuminated to 0-10 lux, with infrared
illumination. Mice were placed in the center of maze at the beginning of each trial. For
optogenetic experiments, mice were allowed to move freely throughout the maze for 15 min
total, with 5-8 mW 473 nm light stimulation pulsed with 50 ms bins at 20 Hz delivered during the
5-10 min epoch (OFF, ON, OFF). For chemogenetic experiments, mice were allowed to freely
move through the area for 10 min total.

Histology

Non-RNAscope section preparation

All sacrifices were performed during the dark period of the light cycle. Animals were
anesthetized prior to sacrifice via combined intraperitoneal injection of 150 mg/kg ketamine
(Zetamine, Vet One) and 15 mg/kg xylazine (AnaSed, AMRI Rensselaer). Except for animals
used in sequencing studies, animals were subject to transcardial perfusion with 10 mL of sterile
phosphate-buffered saline (PBS), followed by 10 mL 4% paraformaldehyde (PFA) solution. The
brain was then extracted from the animal and placed into a 4% (PFA) solution in PBS for at
least 36 hours until it was sectioned on a VT1000S vibratome (Leica). For tissue extracted for
non-RNAscope histology, mice were transcardially perfused with 20 ml phosphate buffered
saline (PBS) followed by 20 ml 4% paraformaldehyde (PFA) in PBS. All brains were extracted
and post-fixed for at least 24 hours in 4% PFA. For tissue extracted for RNAscope, mice under
6 months of age were decapitated once unconscious and their brains were extracted into a
square Peel-A-Way embedding mold (Polysciences) and embedded in OCT (Fisher), and then
snap-frozen a dry-ice/isopentane slurry and stored at -80°C until cryosectioning within a month
of sacrifice.

Tissue was mounted in 5% agarose and sectioned sagittally on a vibratome for retrograde
experiments, or sectioned coronally without mounting for all other non-RNAscope experiments.
These sections were cut at 50 μm and stored in PBS before processing. All connectomic
quantitation was performed on samples using epifluorescence without immunolabeling to avoid
potential bias due to non-stoichiometric antibody binding, while all others were immunolabeled
for visualization of viral targeting accuracy and collection of representative images.
Immunolabeling of eYFP and other GFP-derived fluorophores was performed using goat anti-
GFP primary antibodies (Abcam) and Alexa Fluor 488-conjugated anti-goat secondary
antibodies (Invitrogen), while immunolabeling of mCherry and other DsRed-derived fluorophores
used rabbit anti-DsRed primary antibodies (Takara) and Alexa Fluor 588-conjugated anti-rabbit
antibodies (Invitrogen), all diluted 1:1000 in PBS-T. All non-RNAscope sections were mounted
on Superfrost Plus microscope slides (Fisher) and counterstained with Fluoromount-G
containing DAPI (SouthernBiotech). Sections were stored long-term at 4°C.

RNAscope fluorescence in situ *hybridization*

RNAscope sections were cut at 15 μm on a CM 1950 cryostat (Leica) and mounted on
Superfrost Plus slides and stored at -80˚C until processing via RNAscope within a month of
mounting. RNAscope was performed in an RNA-free environment according to manufacturer
instructions using the Multiplex Fluorescent Reagent Kit v2 (Advanced Cell Diagnostics).[107]
RNAscope was performed using the probes *mm-Slc17a7* in the C2 channel, and *mm-Slc17a6* in
the C3 channel, dyed with Opal 520 and Opal 690 in a counterbalanced manner at 1:15,000
concentration to reduce background fluorescence and allow quantitation of unsaturated, clearly-
distinguishable puncta.[108] Processed RNAscope sections were then mounted with Prolong
Antifade Diamond (ThermoFisher) and stored long-term at 4˚C.

Fluorescence image acquisition

Non-RNAscope Images were acquired at 10X magnification with an VS120 slide scanner
(Olympus), with settings held constant within experiments. Confocal fluorescence images for
RNAscope were acquired on an SP8 (Leica) confocal laser scanning microscope using a
40x/1.30NA oil immersion objective. Serial Z-stack images were acquired using the LASX
software at a thickness of 1 μm per Z stack, with 14-21 planes taken per image. Images were
acquired with identical settings for laser power, detector gain, and amplifier offset for each set of
counterbalanced probe-fluorophore combinations.

Sequencing Data Acquisition

Tissue extraction and cryopreservation for single-cell sequencing

Once unconscious, mice animals were transcardially perfused with ice-cold, carbogen-bubbled
(95% O2, 5% CO2), nuclease-free, 0.22 μm sterile-filtered artificial cerebrospinal fluid (ACSF)
with a composition of 93 mM N-methyl-D-glucamine, 2.5 mM KCl, 1.2 mM NaH2PO4, 30 mM
NaHCO3, 20 mM HEPES, 25 mM glucose, 5 mM sodium ascorbate, 2 mM thiourea, 3 mM
sodium pyruvate, 13.2 mM trehalose, 12 mM N-acetyl-cysteine, 0.5 mM CaCl2, 10 mM MgSO4,
and 93 mM HCl, at pH 7.3-7.4.[42,109] Following transcardial perfusion, brains were immediately
extracted and submerged into ice-cold carbogen-bubbled ACSF, with less than 5 minutes
between the beginning of perfusion and final submersion after extraction. Brains were serially
sectioned in ice-cold, carbogen-bubbled ACSF on a VT1000S vibratome (Leica) with
polytetrafluoroethane-coated razor blades (Ted Pella) at 0.15 mm/sec and 100 Hz, dividing the
whole cerebrum into 400 μm coronal slices. Target regions were microdissected from these
slices under a stereomicroscope using a sterile blunt-end needle (22 gauge for CeA, ASt, and
tail of striatum, 16 gauge for dorsal striatum). All regions were targeted using Paxinos &
Franklin, 5th Edition, as reference.[36] Extracted tissue samples were recovered in ice-cold,
nuclease-free, 0.22 μm sterile-filtered cryoprotective nuclear storage buffer, composed of 0.32
M sucrose, 5 mM CaCl2, 3 mM magnesium acetate, 10 mM Trizma hydrochloride buffer (pH
8.0), 1 mM dithiothreitol, 0.02 U/μl SUPERase•In RNAse Inhibitor (Invitrogen), and 1X cOmplete
Protease Inhibitor Cocktail with EDTA (Roche). Tissue was then snap frozen using a metal
CoolRack M90 (Biocision) pre-chilled to -80˚C and stored at -80˚C until nuclear isolation.
Following extraction of tissue regions of interest, remaining portions of sections were fixed in 4%
paraformaldehyde and 4′,6-diamidino-2-phenylindole (DAPI) was applied to sections at 1 μg/ml.

After fixation and staining, sections were mounted and imaged on an VS120 slide scanner
(Olympus). From these images, dissection accuracy was assessed for each region, and
individual samples were only selected for downstream nuclear isolation if the extracted tissue
fell entirely within the defined target regions.

Nuclear isolation procedures were adapted from multiple methods described previously.[102,110] All
procedures were performed on ice, and all solutions were ice-cold, nuclease-free, and 0.22 μm
sterile-filtered. Cryopreserved tissue pieces were slow thawed by incubation at 4°C for 1 hour
prior to isolation. Tissue pieces were then pooled and resuspended in nuclear isolation medium
composed of 0.25 M sucrose, 25 mM KCl, 5 mM MgCl2, 10 mM Trizma hydrochloride buffer (pH
7.4), 1 mM dithiothreitol, 0.04 U/μl RNasin Plus RNAse Inhibitor (Promega), 1X cOmplete
Protease Inhibitor Cocktail with EDTA (Roche), and 0.1% Triton-X. The pooled tissue pieces in
nuclear isolation medium were transferred to a 2 mL Dounce tissue grinder. Tissue was
homogenized by 5 strokes from the loose pestle and 15 followed by the tight pestle, and the
resulting homogenate was filtered through a 40 μm Flowmi cell strainer (Bel-Art) into a 1.5 ml
Lo-Bind tube (Eppendorf). The homogenate was then centrifuged with a swinging bucket rotor
at 4°C and 1000 x g for 8 minutes. Nuclei were then washed with nuclear flow buffer composed
of DPBS with 1% bovine serum albumin, 1 mM dithiothreitol, and 0.04 U/μl RNasin Plus RNAse
Inhibitor (Promega) and centrifuged at 4°C and 500 x g for 5 minutes, which was subsequently
repeated. Nuclei were finally resuspended in nuclear flow buffer containing 3 μm DRAQ7 (Cell
Signaling Technology) and again filtered through a 40 μm Flowmi cell strainer into a 5 ml round-
bottom polystyrene tube. Each isolation took under 45 minutes to perform, from homogenization
to final suspension.

Fluorescence-activated nuclei sorting (FANS)

FANS was carried out on a FACSAria II SORP (BD Biosciences) using a 70 μm nozzle at 52
PSI sheath pressure. For FANS, debris was first excluded by gating on forward and side scatter
pulse area parameters (FSC-A and SSC-A), followed by exclusion of aggregates (FSC-W and
SSC-W), and finally gating for nuclei based on DRAQ7 fluorescence (APC-Cy7). Nuclei were
successively sorted into 1.5 ml LoBind tubes (Eppendorf) under the purity sort mode. The tube
contained 10X RT master mix without RT Buffer C. 16,000 total nuclei were targeted for
downstream processing, and to account for cytometer errors and subsequent loss of nuclei,
21,000 were sorted into the tube. Nuclei were then immediately processed for snRNA-seq.
FANS conditions were optimized for isolation of debris-free nuclei using the LIVE/DEAD
Viability/Cytotoxicity Kit for Mammalian Cells (Molecular Probes), adding to the final suspension
according to manufacturer instructions and examining on a hemocytometer using an EVOS FL
Cell Imaging System (Thermo Fisher) for enrichment of ethidium homodimer-1-positive nuclei
and the absence of Calcein AM-labeled cellular debris.

Tissue extraction and sample preparation for spatial transcriptomics

Mice were euthanized with CO_2 followed by decapitation, either in the dark or in the light. Brain
hemispheres were collected and placed in OCT and then flash frozen in isopentane in liquid
nitrogen. One hemibrain from each mouse was cryosectioned at -18°C sagittally to a thickness

of 10 mm (~2.8 mm from the midline) using a standard Leica CM1860 cryostat and processed
according to the recommended protocols (Tissue optimization: CG000240 Visium 10X
Genomics; Gene expression: CG000239). The tissue was immediately mounted on a Visium
spatially barcoded slide (10X Genomics). The tissue was covered with OCT and kept at -80
degrees C until it was cryosectioned again starting at the same position to a thickness of 10mm
and mounted onto a Superfrost plus microscope slide (Fisherbrand) for staining. Each section
covered approximately 80% of the 5,000 total spots within their fiducial frame. Slides were
stored at -80˚C until use.

Library Preparation

Library preparation for single nucleus sequencing

Nuclear suspensions were converted into barcoded snRNA-seq libraries using the Chromium
Next GEM Single Cell 3' v3.1 Reagent Kits v3.1 Single Index (10X Genomics). Library
preparation for both assays was performed in accordance with the manufacturer's instructions.
10,000 nuclei were targeted during each snRNA-seq library preparation run. 10X libraries were
first sequenced at low depth on a NextSeq 550 Sequencing System (Illumina) to estimate
quality and number of nuclei for each library, followed by deep sequencing on a NovaSeq 6000
Sequencing System. All runs were performed using 2 x 100-bp paired-end reads, outputting
data in 28/8/91-bp read format for snRNA-seq runs.

Library preparation for Visium spatial transcriptomics

Visium spatial gene expression slides and reagents were used according to the manufacturer
instructions (10X Genomics). Each capture area was 6.5mm x 6.5 mm and contained 5,000
barcoded spots that were 55 μm in diameter (100 μm center to center between spots) provide
an average resolution of about 1 to 10 cells per spot. Optimal permeabilization time was
measured at 24 min. Libraries were prepared according to the Visium protocol (10X Genomics)
and sequenced on a NovaSeq4 (Illumina) at a sequencing depth of 182 million read pairs.
Sequencing was performed with the recommended protocol in a 28/10/10/100-bp read format.
H&E (Hematoxylin, Thermo; Dako bluing buffer, Dako; Eosin Y, Sigma) staining and image
preparation was performed according to the Visium protocol. H&E-stained sections were
imaged using a Nanozoomer slide scanner (Hamamatsu) Spatial gene expression assay was
performed according to the protocol CG000239.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical Analysis

All statistical details can be found in the figure legends. Sample sizes for behavioral studies
were chosen based on past optogenetic studies for each behavior, which had used 6-15
animals per group. Blinding experimenters was not possible for behavioral, imaging, or
sequencing experiments, given familiarity with subjects, but manual quantitation for connectivity
experiments was performed blinded to group with random assignment. All statistical tests were
performed in R (v4.2.3) unless otherwise specified. All statistical tests were performed with two

tails. Group comparisons were made using two-way analysis of variance (ANOVA) followed by
Bonferroni post hoc tests, except where otherwise specified. All behavioral experiments were
performed by multiple experimenters across multiple cohorts each composed of multiple litters,
with littermates distributed across control and treatment groups, with all such cohorts yielding
similar results (data not shown), and topography stimulation experiments were performed
across multiple facilities and institutions. Numbers of mice used for all non-sequencing
experiments are reported within the relevant figures, figure legends, and the text.

Calcium Imaging Data Analysis

Image processing

Data analysis for calcium imaging experiments was adapted from methods previously described
elsewhere.[25] Ca^{2+} imaging data were first motion-corrected using the non-rigid motion correction
algorithm NoRMCorre.[111] Afterwards, neural traces were extracted from the motion-corrected
data using constrained nonnegative matrix factorization (CNMF).[112,113] Spatial components
identified by CNMF were inspected by eye to ensure they were not artifacts. A Gaussian Mixture
Model (GMM) was used to estimate the baseline fluorescence of each neuron. To account for
potential low-frequency drift in the baseline, the GMM was applied along a moving window of
2,500 frames (500 seconds). The fluorescence of each neuron at each time point t was then
normalized to the moving baseline to calculate $\Delta F/F = F_t - F_{baseline} / F_{baseline}$. All subsequent
analysis was performed using custom code written in MATLAB (R2022b).

Hierarchical clustering of pooled averaged responses

1397 $\Delta F/F$ in response to all 6 odors were averaged across trials then Z-scored. The resulting trial-
1398   average values from the following time bins were averaged across time: the first second during
1399   each odor, the last second during each odor, and the first second after each odor. The resulting
1400   18-element vectors were sorted into 6 clusters after agglomerative hierarchical clustering using
1401   Euclidean distance and Ward linkage.

Responsiveness criteria

To determine how many neurons were responsive to a given odor, we compared $\Delta F/F$ at each
frame during the 2 second odor period against a pooled distribution of $\Delta F/F$ values from the 2-
seconds prior to odor onset using a Wilcoxon rank sum test. The resulting p-values were
evaluated with Holm-Bonferroni correction to ensure that familywise error rate (FWER) was
below 0.05. We also counted the number of neurons that were significantly responsive for at
least 4 frames during the odor period to report the total percentage of responsive neurons
during odor.

Single-neuron 6-odor classifiers

To test how reliably a single neuron's fluorescence could discriminate between the 6 odors, we
assessed the performance of multinomial regression (MNR) classifiers trained on a single
neuron's responses to 6 odors. For each neuron and odor pair, we averaged the $\Delta F/F$ during

the last second of the odor exposure for each trial then Z-scored across all trials. The resulting
120-element vector was used to train the MNR classifiers. The 5-fold cross-validated accuracies
are reported and plotted as violin plots.[114] As a control for each neuron, 100 shuffled classifiers
were trained on the data with the odor labels randomly assigned.

Pairwise euclidean distance

To quantify the differences among population-level responses to the 6 odors, we quantified the
pairwise Euclidean distance between the trajectories of odor responses. First, we subtracted the
ΔF/F values during the 2 seconds prior to odor delivery from each frame then averaged these
values across trials for each odor. The pairwise Euclidean distance at each frame was
computed for each odor pair and normalized to the maximum pairwise distance measured in all
odor pairs at any time bin. These calculations were carried out separately for each animal and
then averaged across biological replicates to report the mean and the standard error.

Population classifiers

To assess the discriminability of odor responses in high-dimensional space, we measured the
accuracy of error-correcting output codes (ecoc) classifiers. At each time point relative to odor
delivery, we pooled ΔF/F values from all trials during which either odor was presented. These
values were then normalized and used to train a multinomial ecoc classifier using a Support
Vector Machine (SVM). The accuracy of the classifier was evaluated via 5-fold cross-validation.
To compare the classifier accuracies across different numbers of neurons used for training, we
randomly selected varying numbers of pooled neurons and used the ΔF/F values during the last
second of odor exposure for training.

Behavioral Data Analysis

Behavioral metrics (i.e. performance index, port distance, center distance, open field time, and
total distance) for the four-quadrant preference test, open field test, and elevated plus maze
were calculated on sets of coordinates created by identifying the centroid of the mouse in real
time in video collected from an overhead camera (Basler) at 4 Hz using custom Labview code
and outputting the centroid's coordinates for each frame. The mouse was automatically
identified by taking a background greyscale image of the behavioral assay's environment at the
beginning of each trial and detecting shapes of a minimum size deviating from the background
image by a specific threshold. The centroid was then determined by automated generation of a
bounding box for the mouse in each frame in real time and recording the coordinate of the
centroid of this rectangle.

Four quadrant task data analysis

Mice were tested as previously described.[3] Mice were placed in the chamber for 25 min
experiments and tested no more than once per day. The first 10 min served as a baseline test
for spatial or temporal bias within the chamber during the trial, and no stimulus of any sort was
provided, while the last 15 min were the test of the manipulation. 15 minutes was chosen to
balance time courses of odor responses observed in previous experiments, where appetitive

odors tend to yield initial responses that decay, while aversive odors tend to yield responses
that grow in magnitude over time. To ensure effects did not arise from ceiling or floor effects in
the baseline or from a nonstandard baseline internal state, the mouse had to remain within the
stimulus quadrant during the baseline test between ~20-30% of the time or else the experiment
was terminated, and the mouse was tested again later. The first 2 minutes of data after the
stimulus was introduced were excluded from the analysis to reduce variance and account for
novelty of the stimulus without affecting the overall valence of the behavioral response, and the
last minute of data was excluded to ensure no minor differences in frames captured could affect
analysis. For chemogenetic odor response silencing experiments, animals with vehicle odor
responses below an absolute value of 0.1 were excluded from experiments to avoid false
negatives from attempting to silence a response that was not observed at baseline, which
applied to less than a quarter of overall animals tested across experimental conditions.
Performance index represents the percent difference from chance occupancy in the
manipulation quadrant, calculated as PI = $(P - 25) / 0.25$; where P is the percentage of time the
animal spends in the manipulation quadrant. Mean port distance represents the mean distance
of each point to the deepest point into the manipulation quadrant observed at baseline.

Open field test data analysis

For elevated plus maze analysis, all chemogenetic inhibition trials used the entirely of the 10
min test length as a single period, while optogenetic stimulation trials used the mean of the
three "OFF" periods to compare to the mean of the two "ON" periods. Three metrics of interest
were calculated. Center time was calculated as the proportion of time spent in the middle
square of the open field comprising 50% of its total area. Corner time was calculated as the
proportion of time spent in the corner squares bounded by the walls and the lines bounding the
center region. Time immobile was calculated as the proportion of time when the animal moved
less than 1 cm/s for at least a one-second period. Location of the open field and bounding
regions was kept constant from trial to trial by registering the apparatus to a bounding box with
the same top-down dimensions, and every measured centroid outside of the registered region
resulting from shadows cast and other artifacts was interpolated between the closest points
before and after within the region.

Elevated plus maze data analysis

For elevated plus maze analysis, all chemogenetic inhibition trials used the entirely of the 10
min test length as a single period, while optogenetic stimulation trials used the mean of the two
"OFF" periods to compare to the "ON" period. Three metrics of interest were calculated. Time in
the open arms was calculated as the proportion of time spent in the open arms compared to the
whole period of interest and did not include time in the center between the two arms. Open arm
entries measures number of episodes where the centroid is observed outside of the bounds of
the closed arms or the center region, without any minimum time or distance out onto the open
arms. Finally, distance was simply calculated as the distance traveled during each period of
interest. Location of open and closed arms was kept constant from trial to trial by registering the
apparatus to a cross-shaped bounding box with the same top-down dimensions, and every
measured centroid outside of the registered region due to factors like the mouse leaning over

the edge of the open arms, among others, was interpolated between the closest points before
and after within the region.

Analysis of Single-Nucleus RNA Sequencing Data

1495

Sequence alignment

1496

All samples were processed using Cell Ranger (v5.0.0).[37] All processing was done by using Cell
Ranger's implementation of STAR to align sample sequence reads to their pre-built mm10
vm23/Ens98 reference transcriptome index 2020-A, with predicted and non-validated transcripts
removed. All sequencing reads were aligned to both the exons and the introns present in the
index. Samples were demultiplexed to produce a pair of FASTQ files for each sample. FASTQ
files containing raw read sequence information were aligned to the Cell Ranger index using the
cellranger count command with --chemistry SC3Pv3 and --include-introns flags enabled. Cell
Ranger corrected sequencing errors in cell barcodes to pre-defined sequences in the 10X v3
single-index whitelist within Hamming distance 1. PCR duplicates were removed by selecting
unique combinations of corrected cell barcodes, unique molecular identifiers, gene names, and
location within the transcript. Raw unfiltered count data was read into R (v4.2.1) using the
Seurat package (v4.2.0).[115–118] The final result of the pipeline was a barcode x gene expression
matrix for further analysis downstream.

Quality control

1510

We used the raw, unfiltered matrix output from Cell Ranger as the input to the beginning of the
pipeline. However, to apply a more stringent filter, the emptyDrops dirichlet-multinomial model
from the DropletUtils package (v1.10.2) was applied to each library individually.[119,120] Droplets
with less than 100 total counts were used to construct the ambient RNA profile and an FDR
threshold below 0.001 was used to select putatively occupied droplets. All barcodes with greater
than 1000 UMIs were further assumed non-empty. Most quality filtration choices were heavily
influenced by the recommendations presented in pipeComp.[121] All quality control was performed
on each library individually prior to merging. Minimal quality filtering for each barcode was
performed by setting a floor of 1000 features per barcode for downstream inclusion to ensure
the dataset is entirely composed of high-quality nuclei. Next, to remove highly likely multiplet
barcodes, barcodes were filtered out if their count depth was more than 5 median absolute
deviations above the median count depth. Barcodes were then removed if their proportion of
ribosomal or mitochondrial reads was more than 5 interquartile ranges above the 75th percentile
(median absolute deviations cannot be used, because in many cases the median absolute
deviation is 0). Heterotypic doublets were identified by creating simulated artificial doublets in
scDblFinder (v1.4.1), which uses a DoubletFinder-like model to remove barcodes similar to
simulated doublets, with an assumed doublet rate of 1% per 1000 nuclei in the library.[122,123]
Scater (v1.18.3) was used to produce initial diagnostic tSNE and UMAP plots for visually
checking the influence of each above metric on the structure of the data.[124]

Data processing/transformation

1530

All datasets (initially for all nuclei and again for selected subclusters) were formatted into Seurat
objects (v4.0.0), merged, and then normalized and transformed individually using the
SCTransform (v2) variance stabilizing transform, which performs best according to prior
comparisons in pipeComp.[121,125,126] Following the merge, all genes expressed in 3 or fewer
nuclei of interest were removed from analysis. SCTransform was run returning Pearson
residuals regressing out mitochondrial gene expression, retaining the top 5000 highly variable
features. Dimensionality of the dataset was first reduced using principal component analysis, as
implemented in Seurat's RunPCA function, retaining the top 50 principal components.[127]
Principal components were selected for downstream use by using the lower value of either the
number of principal components where the lowest contributes 5% of standard deviation and all
cumulatively contribute 90% of the standard deviation, or the number of principal components
where the percent change in variation between the consecutive components is lower than 0.1%.
These principal components were used as input to the non-linear tSNE and UMAP
dimensionality reduction methods as implemented by Seurat's RunTSNE and/or RunUMAP
functions with 1000 epochs at 0.5 minimum distance, with otherwise default settings.[128–130]

Clusters were identified via Leiden clustering in latent space using the previously selected
principal components as input.[131] Optimal clustering resolution was identified in a supervised
manner using clustree, finding the highest resolution for each dataset where clustering remains
stable.[132] Cluster annotation was performed in a semi-hierarchical semi-supervised manner,
where known marker genes were first used to separate all nuclei into neuronal and non-
neuronal cell types, and then these cells were re-analyzed and neurons were respectively
separated into glutamatergic and GABAergic neurons, while non-neuronal cells were separated
into astrocytes, microglia, macrophages, oligodendrocytes and their precursors/lineage, mural
cells, endothelia, and vascular leptomeningeal cells. This lower level of cells was then
reanalyzed, and novel cell types were then identified within these more-granular known cell
types. For identification of known cell types, clusters expressing the same marker genes were
manually merged to ensure all cells of a known type were analyzed together, which did not
occur for novel cell type identification. Clusters resulting from specific difference in nuclei quality
instead of true changes in gene expression (i.e. markedly lower mean UMI/features per nucleus,
increased ribosomal/mitochondrial gene expression proportion) were removed prior to final
clustering. Relationships between cell type proportion and plCoA zone were quantitated using
propeller, treating each library as an independent replicate.[133]

Differential expression

Marker genes were identified using Wilcoxon rank-sum test as implemented by the
FindConservedMarkers function in Seurat, using the region as a grouping variable. Genes were
accepted as differentially expressed with a minimum proportion cutoff at 0.1 and minimum fold
change at 1.5-fold (log2-fold change of 0.585), with a p-value cutoff of 0.01 after Bonferroni
correction. To identify genes differentially expressed by region, single-cell values were
converted to pseudo-bulk by batch using the run_de function as implemented in the Libra
package (v1.0.0) using default settings with a minimum proportion cutoff at 0.1, and tested for
differential expression using edgeR's likelihood ratio test.[134] Zone-specific gene expression was
identified by comparing batches from the two isolated zones.

Analysis of Spatial Transcriptomics Data

Sequence and image alignment

All samples were processed using Space Ranger (v1.3.0). All processing was done by using Space Ranger's implementation of STAR to align sample sequence reads to their pre-built mm10 vm23/Ens98 reference transcriptome index 2020-A, with predicted and non-validated transcripts removed, as in snRNA-seq data alignment. Samples were demultiplexed to produce a pair of FASTQ files for each sample. FASTQ files containing raw read sequence information were aligned to the index using the spaceranger count command. Space Ranger corrected sequencing errors in cell barcodes to pre-defined sequences in the single-index whitelist within Hamming distance 1. PCR duplicates were removed by selecting unique combinations of corrected cell barcodes, unique molecular identifiers, gene names, and location within the transcript. Imaging data was processed using automatic fiducial alignment and tissue detection on a brightfield input.

Data processing/transformation

We used the image-filtered matrix output from Space Ranger as the input to the beginning of the pipeline. In a similar manner to snRNA-seq data, all datasets were formatted into Seurat objects (v5.0.0), merged, and then normalized and transformed using the SCTransform (v2) variance stabilizing transform. SCTransform was run returning Pearson residuals regressing out mitochondrial gene expression, retaining the top 5000 highly variable features. Dimensionality of the dataset was first reduced using principal component analysis, as implemented in Seurat's RunPCA function, retaining the top 50 principal components, all of which were used in downstream processing. These principal components were used as input to the non-linear tSNE and UMAP dimensionality reduction methods as implemented by Seurat's RunTSNE and/or RunUMAP functions with 1000 epochs at 0.2 minimum distance, with otherwise default settings. Clusters were identified via Leiden clustering in latent space using all 50 principal components as input. Optimal clustering resolution was identified in a supervised manner using clustree, finding the highest resolution for each dataset where clustering remains stable, choosing a resolution of 0.7. Cluster annotation was performed in a semi-supervised manner, observing where in captured plCoA regions each cluster's spots localized to. For clusters that could not be annotated from spatial location alone (e.g. OLG), marker genes were examined to determine the molecular identity of relevant spots. Spatial data was projected onto neuronal molecular cell types from snRNA-seq data and cell type likelihood was predicted using Seurat's FindTransferAnchors and TransferData functions using snRNA-seq data as a reference and plCoA spatial data as the query, using all 50 PCs. Prediction of a minority of subtypes failed, likely due to low abundance in tissue and/or due to mediolateral spatial differences, alluded to in a separate study, causing the section not to intersect with the part of the tissue containing the relevant neuronal subtypes.[135] Glutamatergic and GABAergic molecular subtype likelihoods were predicted separately to remove noise and increase modeled prediction confidence.

Histological Image Analysis

Registration and localization

Histology for all animals and samples was examined prior to inclusion. Localization within the
coronal plane was determined by registering the coronal slice to the Allen Brain Atlas via the
ABBA plugin, using elastix to sequentially perform affine and spline registration of the DAPI
channel of the slice to the Nissl channel of the atlas.[136,137] The region of interest was then
compared to the Paxinos and Franklin atlas to confirm localization, and find the region's
anteroposterior distance from bregma.[36] This combined method was used because sections
cannot be accurately registered to the Paxinos and Franklin atlas due to low Z-resolution, while
the Allen Brain Atlas lacks information about anteroposterior distance from bregma. Exclusion
based on histology would occur when most of the intervention fell outside of the region of
interest. Due to these differences, individual representative images use the individually
registered Allen Reference Atlas schematics with the comparable Paxinos and Franklin
anteroposterior coordinates noted, while consolidated targeting schematics use the Paxinos and
Franklin atlas for visualization.

Quantification of histological fluorescence

In anterograde tracing experiments, output quantification was performed based on background-
corrected total fluorescence. For all non-collateralization anterograde experiments, fluorescence
intensities were quantified using FIJI (v2.9.0) throughout the whole brain in a series of evenly-
space 50 μm coronal sections, manually segmenting by region with all settings held constant
within experiments.[138] For collateralization experiments, we exclusively examined fluorescence
in the MeA and NAc. We calculated background-corrected total fluorescence using the equation
$F_{total} = ID - (Area \times F_{background})$, where F_{total} is the background-corrected total fluorescence, *ID* is
the integrated density, and $F_{background}$ is the mean background fluorescence measured from four
randomly selected areas per section not receiving input from plCoA. Overall proportion was
calculated by taking the sum of background-corrected fluorescence values across all sections
for a given region and dividing it by the sum of all background-corrected values. For retrograde
experiments, we quantified number of cells using the Cell Counter plugin (v3.0.0) in FIJI. The
sagittal brain slices containing the plCoA were then compared to Paxinos and Franklin, 5th
Edition to count the number of cells found per distance away from bregma from -1.3 to -2.5 mm
in increments of 100 μm.[36] At least two sections per region per animal were analyzed.
Representative images were produced from slide scanner image output, with background
subtraction and uniform brightness and contrast thresholds applied equally to all fluorescent
channels in FIJI to avoid potential distortion of visible fluorescence levels.

Quantification and analysis of RNAscope images

RNAscope images were analyzed as previously described.[86] Images were opened in FIJI and
individual Z-planes encompassing the entire ROI were selected from each image for further
image processing. Background was subtracted from all channels in all images using the
subtract background feature. Masks of each region were drawn based on the mouse brain atlas,
and images were then saved as 8bit TIFFs for further cell and puncta identification in
CellProfiler (v4.2.4).[36,139] Image TIFFs were run through CellProfiler using an optimized version
of the CellProfiler Colocalization pipeline. The pipeline was optimized to identify DAPI labelled
cells (15-45 pixels in diameter) and then subsequently identify mRNA puncta (4-10 pixels in

diameter). DAPI cell detection was further restricted by shrinking DAPI ROIs by 1 pixel. Puncta
overlapping with DAPI-identified cells (using the relate objects module) were considered for
analysis to assess the level of mRNA expression per cell. To determine if cells were expressing
mRNA, a threshold of 5 or more puncta within twice the diameter of nucleus centered over the
nucleus was used.[140] Total number and density of *Slc17a6*+ and *Slc17a7*+ cells in each region
of interest were calculated from CellProfiler .csv outputs using custom R scripts.

References

1. Stowers, L., Cameron, P., and Keller, J.A. (2013). Ominous Odors: olfactory control of instinctive fear and aggression in mice. Curr. Opin. Neurobiol. *23*, 339–345. https://doi.org/10.1016/j.conb.2013.01.007.

2. Stowers, L., and Kuo, T.-H. (2015). Mammalian pheromones; emerging properties and mechanisms of detection. Curr. Opin. Neurobiol. *34*, 103–109. https://doi.org/10.1016/j.conb.2015.02.005.

3. Root, C.M., Denny, C.A., Hen, R., and Axel, R. (2014). The participation of cortical amygdala in innate, odour-driven behaviour. Nature *515*, 269–273. https://doi.org/10.1038/NATURE13897.

4. Li, Q., and Liberles, S.D. (2015). Aversion and Attraction through Olfaction. Curr. Biol. *25*, R120–R129. https://doi.org/10.1016/j.cub.2014.11.044.

5. Hayden, S., Bekaert, M., Crider, T.A., Mariani, S., Murphy, W.J., and Teeling, E.C. (2010). Ecological adaptation determines functional mammalian olfactory subgenomes. Genome Res. *20*, 1–9. https://doi.org/10.1101/gr.099416.109.

6. Saraiva, L.R., Riveros-McKay, F., Mezzavilla, M., Abou-Moussa, E.H., Arayata, C.J., Makhlouf, M., Trimmer, C., Ibarra-Soria, X., Khan, M., Van Gerven, L., et al. (2019). A transcriptomic atlas of mammalian olfactory mucosae reveals an evolutionary influence on food odor detection in humans. Sci. Adv. *5*, eaax0396. https://doi.org/10.1126/sciadv.aax0396.

7. Ibarra-Soria, X., Nakahara, T.S., Lilue, J., Jiang, Y., Trimmer, C., Souza, M.A., Netto, P.H., Ikegami, K., Murphy, N.R., Kusma, M., et al. (2017). Variation in olfactory neuron repertoires is genetically controlled and environmentally modulated. eLife *6*, e21476. https://doi.org/10.7554/eLife.21476.

8. Tye, K.M. (2018). Neural Circuit Motifs in Valence Processing. Neuron *100*, 436–452. https://doi.org/10.1016/j.neuron.2018.10.001.

9. Dewan, A., Pacifico, R., Zhan, R., Rinberg, D., and Bozza, T. (2013). Non-redundant coding of aversive odours in the main olfactory pathway. Nature *497*, 486–489. https://doi.org/10/f4xchh.

10. Li, Q., Korzan, W.J., Ferrero, D.M., Chang, R.B., Roy, D.S., Buchi, M., Lemon, J.K., Kaur, A.W., Stowers, L., Fendt, M., et al. (2013). Synchronous Evolution of an Odor Biosynthesis Pathway and Behavioral Response. Curr. Biol. *23*, 11–20. https://doi.org/10.1016/j.cub.2012.10.047.

11. Semmelhack, J.L., and Wang, J.W. (2009). Select Drosophila glomeruli mediate innate olfactory attraction and aversion. Nature *459*, 218–223. https://doi.org/10.1038/nature07983.

12. Beyeler, A., Namburi, P., Glober, G.F., Simonnet, C., Calhoon, G.G., Conyers, G.F., Luck, R., Wildes, C.P., and Tye, K.M. (2016). Divergent Routing of Positive and Negative

Information from the Amygdala during Memory Retrieval. Neuron *90*, 348–361.
https://doi.org/10.1016/j.neuron.2016.03.004.

13. Chen, Y., Chen, X., Baserdem, B., Zhan, H., Li, Y., Davis, M.B., Kebschull, J.M., Zador,
A.M., Koulakov, A.A., and Albeanu, D.F. (2022). High-throughput sequencing of single
neuron projections reveals spatial organization in the olfactory cortex. Cell *185*, 4117-
4134.e28. https://doi.org/10.1016/j.cell.2022.09.038.

14. Sosulski, D.L., Lissitsyna Bloom, M., Cutforth, T., Axel, R., and Datta, S.R. (2011). Distinct
representations of olfactory information in different cortical centres. Nature *472*, 213–219.
https://doi.org/10/fdwfsk.

15. Miyamichi, K., Amat, F., Moussavi, F., Wang, C., Wickersham, I., Wall, N.R., Taniguchi, H.,
Tasic, B., Huang, Z.J., He, Z., et al. (2011). Cortical representations of olfactory input by
trans-synaptic tracing. Nature *472*, 191–199. https://doi.org/10/bmj88g.

16. Wang, I.-H., Murray, E., Andrews, G., Jiang, H.-C., Park, S.J., Donnard, E., Durán-Laforet,
V., Bear, D.M., Faust, T.E., Garber, M., et al. (2022). Spatial transcriptomic reconstruction of
the mouse olfactory glomerular map suggests principles of odor processing. Nat. Neurosci.
25, 484–492. https://doi.org/10.1038/s41593-022-01030-8.

17. Iurilli, G., and Datta, S.R. (2017). Population Coding in an Innately Relevant Olfactory Area.
Neuron *93*, 1180-1197.e7. https://doi.org/10/f9tpmx.

18. Beyeler, A., Chang, C.-J., Silvestre, M., Lévêque, C., Namburi, P., Wildes, C.P., and Tye,
K.M. (2018). Organization of Valence-Encoding and Projection-Defined Neurons in the
Basolateral Amygdala. Cell Rep. *22*, 905–918. https://doi.org/10/gg6473.

19. Kim, J., Pignatelli, M., Xu, S., Itohara, S., and Tonegawa, S. (2016). Antagonistic negative
and positive neurons of the basolateral amygdala. Nat. Neurosci. *19*, 1636–1646.
https://doi.org/10.1038/nn.4414.

20. Tsuji, T., Tsuji, C., Lozic, M., Ludwig, M., and Leng, G. (2019). Coding of odors in the
anterior olfactory nucleus. Physiol. Rep. *7*, e14284. https://doi.org/10.14814/phy2.14284.

21. Malnic, B., Hirono, J., Sato, T., and Buck, L.B. (1999). Combinatorial Receptor Codes for
Odors. Cell *96*, 713–723. https://doi.org/10.1016/S0092-8674(00)80581-4.

22. Stettler, D.D., and Axel, R. (2009). Representations of Odor in the Piriform Cortex. Neuron
63, 854–864. https://doi.org/10.1016/j.neuron.2009.09.005.

23. Nara, K., Saraiva, L.R., Ye, X., and Buck, L.B. (2011). A Large-Scale Analysis of Odor
Coding in the Olfactory Epithelium. J. Neurosci. *31*, 9179–9191.
https://doi.org/10.1523/JNEUROSCI.1282-11.2011.

24. Soucy, E.R., Albeanu, D.F., Fantana, A.L., Murthy, V.N., and Meister, M. (2009). Precision
and diversity in an odor map on the olfactory bulb. Nat. Neurosci. *12*, 210–220.
https://doi.org/10.1038/nn.2262.

25. Lee, D., Liu, L., and Root, C.M. (2023). Transformation of value signaling in a striatopallidal
circuit. eLife *12*. https://doi.org/10.7554/eLife.90976.

26. Chae, H., Kepple, D.R., Bast, W.G., Murthy, V.N., Koulakov, A.A., and Albeanu, D.F.
(2019). Mosaic representations of odors in the input and output layers of the mouse
olfactory bulb. Nat. Neurosci. *22*, 1306–1317. https://doi.org/10.1038/s41593-019-0442-z.

27. Ma, L., Qiu, Q., Gradwohl, S., Scott, A., Yu, E.Q., Alexander, R., Wiegraebe, W., and Yu,
C.R. (2012). Distributed representation of chemical features and tunotopic organization of
glomeruli in the mouse olfactory bulb. Proc. Natl. Acad. Sci. *109*, 5481–5486.
https://doi.org/10.1073/pnas.1117491109.

28. Roland, B., Deneux, T., Franks, K.M., Bathellier, B., and Fleischmann, A. (2017). Odor
identity coding by distributed ensembles of neurons in the mouse olfactory cortex. eLife *6*,
e26337. https://doi.org/10.7554/eLife.26337.

29. Payton, C.A., Wilson, D.A., and Wesson, D.W. (2012). Parallel odor processing by two
anatomically distinct olfactory bulb target structures. PloS One *7*, e34926.
https://doi.org/10.1371/journal.pone.0034926.

30. Cousens, G.A. (2020). Characterization of odor-evoked neural activity in the olfactory
peduncle. IBRO Rep. *9*, 157–163. https://doi.org/10.1016/j.ibror.2020.07.010.

31. Peng, Y., Gillis-Smith, S., Jin, H., Tränkner, D., Ryba, N.J.P., and Zuker, C.S. (2015). Sweet
and bitter taste in the brain of awake behaving animals. Nature *527*, 512–515.
https://doi.org/10.1038/nature15763.

32. Pashkovski, S.L., Iurilli, G., Brann, D., Chicharro, D., Drummey, K., Franks, K., Panzeri, S.,
and Datta, S.R. (2020). Structure and flexibility in cortical representations of odor space.
Nature *583*, 253–258. https://doi.org/10.1038/s41586-020-2451-1.

33. Wiltschko, A.B., Johnson, M.J., Iurilli, G., Peterson, R.E., Katon, J.M., Pashkovski, S.L.,
Abraira, V.E., Adams, R.P., and Datta, S.R. (2015). Mapping Sub-Second Structure in
Mouse Behavior. Neuron *88*, 1121–1135. https://doi.org/10.1016/j.neuron.2015.11.031.

34. Wang, L., Gillis-Smith, S., Peng, Y., Zhang, J., Chen, X., Salzman, C.D., Ryba, N.J.P., and
Zuker, C.S. (2018). The coding of valence and identity in the mammalian taste system.
Nature *558*, 127–131. https://doi.org/10/gdq4gk.

35. Chen, X., Gabito, M., Peng, Y., Ryba, N.J.P., and Zuker, C.S. (2011). A Gustotopic Map of
Taste Qualities in the Mammalian Brain. Science *333*, 1262–1266. https://doi.org/10/cv4x7s.

36. Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates, Compact - 5th Edition
https://www.elsevier.com/books/paxinos-and-franklins-the-mouse-brain-in-stereotaxic-
coordinates-compact/franklin/978-0-12-816159-3.

37. Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B.,
Wheeler, T.D., McDermott, G.P., Zhu, J., et al. (2017). Massively parallel digital
transcriptional profiling of single cells. Nat. Commun. *8*, 14049. https://doi.org/10/f9mbtp.

38. Tasic, B., Menon, V., Nguyen, T.N., Kim, T.K., Jarsky, T., Yao, Z., Levi, B., Gray, L.T.,
Sorensen, S.A., Dolbeare, T., et al. (2016). Adult mouse cortical cell taxonomy revealed by
single cell transcriptomics. Nat. Neurosci. *19*, 335–346. https://doi.org/10/f778w5.

39. Zeisel, A., Hochgerner, H., Lönnerberg, P., Johnsson, A., Memic, F., van der Zwan, J.,
Häring, M., Braun, E., Borm, L.E., La Manno, G., et al. (2018). Molecular Architecture of the
Mouse Nervous System. Cell *174*, 999-1014.e22. https://doi.org/10/gdxxnt.

40. Marques, S., Zeisel, A., Codeluppi, S., van Bruggen, D., Mendanha Falcão, A., Xiao, L., Li,
H., Häring, M., Hochgerner, H., Romanov, R.A., et al. (2016). Oligodendrocyte
heterogeneity in the mouse juvenile and adult central nervous system. Science *352*, 1326–
1329. https://doi.org/10.1126/science.aaf6463.

41. Yasuda, K., Cline, C., Vogel, P., Onciu, M., Fatima, S., Sorrentino, B.P., Thirumaran, R.K.,
Ekins, S., Urade, Y., Fujimori, K., et al. (2013). Drug Transporters on Arachnoid Barrier
Cells Contribute to the Blood–Cerebrospinal Fluid Barrier. Drug Metab. Dispos. *41*, 923–
931. https://doi.org/10.1124/dmd.112.050344.

42. Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,
Garren, E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct
transcriptomic cell types across neocortical areas. Nature *563*, 72. https://doi.org/10/gfgqmf.

43. Yao, Z., Velthoven, C.T.J. van, Nguyen, T.N., Goldy, J., Sedeno-Cortes, A.E., Baftizadeh,
F., Bertagnolli, D., Casper, T., Chiang, M., Crichton, K., et al. (2021). A taxonomy of
transcriptomic cell types across the isocortex and hippocampal formation. Cell *184*, 3222.
https://doi.org/10.1016/j.cell.2021.04.021.

44. Harris, J.A., Hirokawa, K.E., Sorensen, S.A., Gu, H., Mills, M., Ng, L.L., Bohn, P., Mortrud,
M., Ouellette, B., Kidney, J., et al. (2014). Anatomical characterization of Cre driver mice for
neural circuit mapping and manipulation. Front. Neural Circuits *8*, 76.
https://doi.org/10.3389/fncir.2014.00076.

45. Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., and Lowell, B.B. (2011). Leptin action on
GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. Neuron
71, 142–154. https://doi.org/10.1016/j.neuron.2011.05.028.

46. Krashes, M.J., Koda, S., Ye, C., Rogan, S.C., Adams, A.C., Cusher, D.S., Maratos-Flier, E.,
Roth, B.L., and Lowell, B.B. (2011). Rapid, reversible activation of AgRP neurons drives
feeding behavior in mice. J Clin Invest *121*, 1424–1428. https://doi.org/10/bd3bv9.

47. Ikemoto, S. (2007). Dopamine reward circuitry: two projection systems from the ventral
midbrain to the nucleus accumbens-olfactory tubercle complex. Brain Res. Rev. *56*, 27–78.
https://doi.org/10.1016/j.brainresrev.2007.05.004.

48. Müller, M., and Fendt, M. (2006). Temporary inactivation of the medial and basolateral
amygdala differentially affects TMT-induced fear behavior in rats. Behav. Brain Res. *167*,
57–62. https://doi.org/10.1016/j.bbr.2005.08.016.

49. Lebow, M.A., and Chen, A. (2016). Overshadowed by the amygdala: the bed nucleus of the
stria terminalis emerges as key to psychiatric disorders. Mol. Psychiatry *21*, 450–463.
https://doi.org/10.1038/mp.2016.1.

50. Sedwick, V.M., and Autry, A.E. (2022). Anatomical and molecular features of the
amygdalohippocampal transition area and its role in social and emotional behavior

processes. Neurosci. Biobehav. Rev. *142*, 104893.
https://doi.org/10.1016/j.neubiorev.2022.104893.

51. Pardo-Bellver, C., Vila-Martin, M.E., Martínez-Bellver, S., Villafranca-Faus, M., Teruel-
Sanchis, A., Savarelli-Balsamo, C.A., Drabik, S.M., Martínez-Ricós, J., Cervera-Ferri, A.,
Martínez-García, F., et al. (2022). Neural activity patterns in the chemosensory network
encoding vomeronasal and olfactory information in mice. Front. Neuroanat. *16*.
https://doi.org/10.3389/fnana.2022.988015.

52. Kaas, J.H. (1997). Topographic maps are fundamental to sensory processing. Brain Res.
Bull. *44*, 107–112. https://doi.org/10.1016/s0361-9230(97)00094-4.

53. Garrett, M.E., Nauhaus, I., Marshel, J.H., and Callaway, E.M. (2014). Topography and Areal
Organization of Mouse Visual Cortex. J. Neurosci. *34*, 12587–12600.
https://doi.org/10.1523/JNEUROSCI.1124-14.2014.

54. Kaas, J.H., Nelson, R.J., Sur, M., Lin, C.-S., and Merzenich, M.M. (1979). Multiple
Representations of the Body Within the Primary Somatosensory Cortex of Primates.
Science *204*, 521–523. https://doi.org/10.1126/science.107591.

55. Schreiner, C.E., and Winer, J.A. (2007). Auditory Cortex Mapmaking: Principles,
Projections, and Plasticity. Neuron *56*, 356–365.
https://doi.org/10.1016/j.neuron.2007.10.013.

56. Imai, T., Sakano, H., and Vosshall, L.B. (2010). Topographic Mapping—The Olfactory
System. Cold Spring Harb. Perspect. Biol. *2*, a001776.
https://doi.org/10.1101/cshperspect.a001776.

57. Auffarth, B. (2013). Understanding smell--the olfactory stimulus problem. Neurosci.
Biobehav. Rev. *37*, 1667–1679. https://doi.org/10.1016/j.neubiorev.2013.06.009.

58. Ressler, K.J., Sullivan, S.L., and Buck, L.B. (1993). A zonal organization of odorant receptor
gene expression in the olfactory epithelium. Cell *73*, 597–609. https://doi.org/10/b7gqvk.

59. Zapiec, B., and Mombaerts, P. (2020). The Zonal Organization of Odorant Receptor Gene
Choice in the Main Olfactory Epithelium of the Mouse. Cell Rep. *30*, 4220-4234.e5.
https://doi.org/10.1016/j.celrep.2020.02.110.

60. Pacifico, R., Dewan, A., Cawley, D., Guo, C., and Bozza, T. (2012). An olfactory subsystem
that mediates high-sensitivity detection of volatile amines. Cell Rep. *2*, 76–88.
https://doi.org/10/gfgp62.

61. Kobayakawa, K., Kobayakawa, R., Matsumoto, H., Oka, Y., Imai, T., Ikawa, M., Okabe, M.,
Ikeda, T., Itohara, S., Kikusui, T., et al. (2007). Innate versus learned odour processing in
the mouse olfactory bulb. Nature *450*, 503–508. https://doi.org/10/d899x4.

62. Midroit, M., Chalençon, L., Renier, N., Milton, A., Thevenet, M., Sacquet, J., Breton, M.,
Forest, J., Noury, N., Richard, M., et al. (2021). Neural processing of the reward value of
pleasant odorants. Curr. Biol. CB *31*, 1592-1605.e9.
https://doi.org/10.1016/j.cub.2021.01.066.

63. Kermen, F., Midroit, M., Kuczewski, N., Forest, J., Thévenet, M., Sacquet, J., Benetollo, C.,
Richard, M., Didier, A., and Mandairon, N. (2016). Topographical representation of odor
hedonics in the olfactory bulb. Nat. Neurosci. *19*, 876–878. https://doi.org/10.1038/nn.4317.

64. Choi, G.B., Dong, H., Murphy, A.J., Valenzuela, D.M., Yancopoulos, G.D., Swanson, L.W.,
and Anderson, D.J. (2005). Lhx6 Delineates a Pathway Mediating Innate Reproductive
Behaviors from the Amygdala to the Hypothalamus. Neuron *46*, 647–660.
https://doi.org/10.1016/j.neuron.2005.04.011.

65. Cohn, R., Morantte, I., and Ruta, V. (2015). Coordinated and Compartmentalized
Neuromodulation Shapes Sensory Processing in Drosophila. Cell *163*, 1742–1755.
https://doi.org/10.1016/j.cell.2015.11.019.

66. Levitan, D., Lin, J.-Y., Wachutka, J., Mukherjee, N., Nelson, S.B., and Katz, D.B. (2019).
Single and population coding of taste in the gustatory cortex of awake mice. J.
Neurophysiol. *122*, 1342–1356. https://doi.org/10.1152/jn.00357.2019.

67. Chen, K., Kogan, J.F., and Fontanini, A. (2021). Spatially Distributed Representation of
Taste Quality in the Gustatory Insular Cortex of Behaving Mice. Curr. Biol. *31*, 247-256.e4.
https://doi.org/10.1016/j.cub.2020.10.014.

68. Fletcher, M.L., Ogg, M.C., Lu, L., Ogg, R.J., and Boughter, J.D. (2017). Overlapping
Representation of Primary Tastes in a Defined Region of the Gustatory Cortex. J. Neurosci.
37, 7595–7605. https://doi.org/10/gbsjxs.

69. Lavi, K., Jacobson, G.A., Rosenblum, K., and Lüthi, A. (2018). Encoding of Conditioned
Taste Aversion in Cortico-Amygdala Circuits. Cell Rep. *24*, 278–283.
https://doi.org/10.1016/j.celrep.2018.06.053.

70. Erö, C., Gewaltig, M.-O., Keller, D., and Markram, H. (2018). A Cell Atlas for the Mouse
Brain. Front. Neuroinformatics *12*. https://doi.org/10/ggc9xg.

71. Cembrowski, M.S., Phillips, M.G., DiLisio, S.F., Shields, B.C., Winnubst, J., Chandrashekar,
J., Bas, E., and Spruston, N. (2018). Dissociable Structural and Functional Hippocampal
Outputs via Distinct Subiculum Cell Classes. Cell *173*, 1280-1292.e18.
https://doi.org/10/gdb82r.

72. Cembrowski, M.S., and Spruston, N. (2019). Heterogeneity within classical cell types is the
rule: lessons from hippocampal pyramidal neurons. Nat. Rev. Neurosci. *20*, 193–204.
https://doi.org/10.1038/s41583-019-0125-5.

73. Ding, S.-L., Yao, Z., Hirokawa, K.E., Nguyen, T.N., Graybuck, L.T., Fong, O., Bohn, P., Ngo,
K., Smith, K.A., Koch, C., et al. (2020). Distinct Transcriptomic Cell Types and Neural
Circuits of the Subiculum and Prosubiculum along the Dorsal-Ventral Axis. Cell Rep. *31*,
107648. https://doi.org/10/gg63q5.

74. Cembrowski, M.S., Wang, L., Lemire, A.L., Copeland, M., DiLisio, S.F., Clements, J., and
Spruston, N. (2018). The subiculum is a patchwork of discrete subregions. eLife *7*, e37701.
https://doi.org/10/gg63q4.

75. Scala, F., Kobak, D., Bernabucci, M., Bernaerts, Y., Cadwell, C.R., Castro, J.R., Hartmanis,
L., Jiang, X., Laturnus, S., Miranda, E., et al. (2021). Phenotypic variation of transcriptomic
cell types in mouse motor cortex. Nature *598*, 144–150. https://doi.org/10.1038/s41586-020-
2907-3.

76. Gouwens, N.W., Sorensen, S.A., Baftizadeh, F., Budzillo, A., Lee, B.R., Jarsky, T., Alfiler,
L., Baker, K., Barkan, E., Berry, K., et al. (2020). Integrated Morphoelectric and
Transcriptomic Classification of Cortical GABAergic Cells. Cell *183*, 935-953.e19.
https://doi.org/10.1016/j.cell.2020.09.057.

77. O'Leary, T.P., Sullivan, K.E., Wang, L., Clements, J., Lemire, A.L., and Cembrowski, M.S.
(2020). Extensive and spatially variable within-cell-type heterogeneity across the basolateral
amygdala. eLife *9*, e59003. https://doi.org/10.7554/eLife.59003.

78. Calvigioni, D., Fuzik, J., Le Merre, P., Slashcheva, M., Jung, F., Ortiz, C., Lentini, A.,
Csillag, V., Graziano, M., Nikolakopoulou, I., et al. (2023). Esr1+ hypothalamic-habenula
neurons shape aversive states. Nat. Neurosci. *26*, 1245–1255.
https://doi.org/10.1038/s41593-023-01367-8.

79. Phillips, J.W., Schulmann, A., Hara, E., Winnubst, J., Liu, C., Valakh, V., Wang, L., Shields,
B.C., Korff, W., Chandrashekar, J., et al. (2019). A repeated molecular architecture across
thalamic pathways. Nat. Neurosci. *22*, 1925. https://doi.org/10.1038/s41593-019-0483-3.

80. Mandelbaum, G., Taranda, J., Haynes, T.M., Hochbaum, D.R., Huang, K.W., Hyun, M.,
Venkataraju, K.U., Straub, C., Wang, W., Robertson, K., et al. (2019). Distinct Cortical-
Thalamic-Striatal Circuits through the Parafascicular Nucleus. Neuron *102*, 636-652.e7.
https://doi.org/10.1016/j.neuron.2019.02.035.

81. Hochgerner, H., Singh, S., Tibi, M., Lin, Z., Skarbianskis, N., Admati, I., Ophir, O.,
Reinhardt, N., Netser, S., Wagner, S., et al. (2023). Neuronal types in the mouse amygdala
and their transcriptional response to fear conditioning. Nat. Neurosci. *26*, 2237–2249.
https://doi.org/10.1038/s41593-023-01469-3.

82. Klingler, E. (2017). Development and Organization of the Evolutionarily Conserved Three-
Layered Olfactory Cortex. eNeuro *4*. https://doi.org/10.1523/ENEURO.0193-16.2016.

83. Shi, H., He, Y., Zhou, Y., Huang, J., Maher, K., Wang, B., Tang, Z., Luo, S., Tan, P., Wu,
M., et al. (2023). Spatial atlas of the mouse central nervous system at molecular resolution.
Nature *622*, 552–561. https://doi.org/10.1038/s41586-023-06569-5.

84. Yao, Z., van Velthoven, C.T.J., Kunst, M., Zhang, M., McMillen, D., Lee, C., Jung, W.,
Goldy, J., Abdelhak, A., Aitken, M., et al. (2023). A high-resolution transcriptomic and spatial
atlas of cell types in the whole mouse brain. Nature *624*, 317–332.
https://doi.org/10.1038/s41586-023-06812-z.

85. Zhang, M., Pan, X., Jung, W., Halpern, A.R., Eichhorn, S.W., Lei, Z., Cohen, L., Smith, K.A.,
Tasic, B., Yao, Z., et al. (2023). Molecularly defined and spatially resolved cell atlas of the
whole mouse brain. Nature *624*, 343–354. https://doi.org/10.1038/s41586-023-06808-9.

86. Mills, F., Lee, C.R., Howe, J.R., Li, H., Shao, S., Keisler, M.N., Lemieux, M.E., Taschbach,
F.H., Keyes, L.R., Borio, M., et al. (2022). Amygdalostriatal transition zone neurons encode

sustained valence to direct conditioned behaviors. Preprint at bioRxiv,
https://doi.org/10.1101/2022.10.28.514263 https://doi.org/10.1101/2022.10.28.514263.

87. Singer, W. (2021). Recurrent dynamics in the cerebral cortex: Integration of sensory
evidence with stored knowledge. Proc. Natl. Acad. Sci. *118*, e2101043118.
https://doi.org/10.1073/pnas.2101043118.

88. Floresco, S.B. (2015). The Nucleus Accumbens: An Interface Between Cognition, Emotion,
and Action. Httpsdoiorg101146annurev-Psych-010213-115159.
https://doi.org/10.1146/annurev-psych-010213-115159.

89. Petrulis, A. (2020). Chapter 2 - Structure and function of the medial amygdala. In Handbook
of Behavioral Neuroscience Handbook of Amygdala Structure and Function., J. H. Urban
and J. A. Rosenkranz, eds. (Elsevier), pp. 39–61. https://doi.org/10.1016/B978-0-12-
815134-1.00002-7.

90. Day, H.E.W., Masini, C.V., and Campeau, S. (2004). The pattern of brain c-fos mRNA
induced by a component of fox odor, 2,5-dihydro-2,4,5-trimethylthiazoline (TMT), in rats,
suggests both systemic and processive stress characteristics. Brain Res. *1025*, 139–151.
https://doi.org/10.1016/j.brainres.2004.07.079.

91. Saito, H., Nishizumi, H., Suzuki, S., Matsumoto, H., Ieki, N., Abe, T., Kiyonari, H., Morita,
M., Yokota, H., Hirayama, N., et al. (2017). Immobility responses are induced by
photoactivation of single glomerular species responsive to fox odour TMT. Nat. Commun. *8*,
1–10. https://doi.org/10.1038/ncomms16011.

92. Zador, A.M. (2019). A critique of pure learning and what artificial neural networks can learn
from animal brains. Nat. Commun. *10*, 3770. https://doi.org/10.1038/s41467-019-11786-6.

93. Koulakov, A., Shuvaev, S., Lachi, D., and Zador, A. (2022). Encoding innate ability through
a genomic bottleneck. Preprint at bioRxiv, https://doi.org/10.1101/2021.03.16.435261
https://doi.org/10.1101/2021.03.16.435261.

94. Barabási, D.L., Beynon, T., Katona, Á., and Perez-Nieves, N. (2023). Complex computation
from developmental priors. Nat. Commun. *14*, 2226. https://doi.org/10.1038/s41467-023-
37980-1.

95. Qiu, Q., Wu, Y., Ma, L., and Yu, C.R. (2021). Encoding innately recognized odors via a
generalized population code. Curr. Biol. *31*, 1813-1825.e4.
https://doi.org/10.1016/j.cub.2021.01.094.

96. Illig, K.R., and Haberly, L.B. (2003). Odor-evoked activity is spatially distributed in piriform
cortex. J. Comp. Neurol. *457*, 361–373. https://doi.org/10.1002/cne.10557.

97. Tritsch, N.X., and Sabatini, B.L. (2012). Dopaminergic modulation of synaptic transmission
in cortex and striatum. Neuron *76*, 33–50. https://doi.org/10.1016/j.neuron.2012.09.023.

98. Martiros, N., Kapoor, V., Kim, S.E., and Murthy, V.N. (2022). Distinct representation of cue-
outcome association by D1 and D2 neurons in the ventral striatum's olfactory tubercle. eLife
11, e75463. https://doi.org/10.7554/eLife.75463.

99. Millman, D.J., and Murthy, V.N. (2020). Rapid Learning of Odor–Value Association in the
Olfactory Striatum. J. Neurosci. *40*, 4335–4347. https://doi.org/10.1523/JNEUROSCI.2604-
19.2020.

100.   Gadziola, M.A., Stetzik, L.A., Wright, K.N., Milton, A.J., Arakawa, K., Cortijo, M. del M.,
and Wesson, D.W. (2020). A Neural System that Represents the Association of Odors with
Rewarded Outcomes and Promotes Behavioral Engagement. Cell Rep. *32*.
https://doi.org/10.1016/j.celrep.2020.107919.

101.   Gadziola, M.A., Tylicki, K.A., Christian, D.L., and Wesson, D.W. (2015). The olfactory
tubercle encodes odor valence in behaving mice. J. Neurosci. Off. J. Soc. Neurosci. *35*,
4515–4527. https://doi.org/10.1523/JNEUROSCI.4750-14.2015.

102.   Preissl, S., Fang, R., Huang, H., Zhao, Y., Raviram, R., Gorkin, D.U., Zhang, Y., Sos,
B.C., Afzal, V., Dickel, D.E., et al. (2018). Single-nucleus analysis of accessible chromatin in
developing mouse forebrain reveals cell-type-specific transcriptional regulation. Nat.
Neurosci. *21*, 432–439. https://doi.org/10/gc9gx2.

103.   Davis, A., Gao, R., and Navin, N.E. (2019). SCOPIT: sample size calculations for single-
cell sequencing experiments. BMC Bioinformatics *20*, 566. https://doi.org/10/ggc9ws.

104.   Sturchler-Pierrat, C., Abramowski, D., Duke, M., Wiederhold, K.-H., Mistl, C., Rothacher,
S., Ledermann, B., Bürki, K., Frey, P., Paganetti, P.A., et al. (1997). Two amyloid precursor
protein transgenic mouse models with Alzheimer disease-like pathology. Proc. Natl. Acad.
Sci. *94*, 13287–13292. https://doi.org/10.1073/pnas.94.24.13287.

105.   Oh, S.W., Harris, J.A., Ng, L., Winslow, B., Cain, N., Mihalas, S., Wang, Q., Lau, C.,
Kuan, L., Henry, A.M., et al. (2014). A mesoscale connectome of the mouse brain. Nature
508, 207–214. https://doi.org/10.1038/nature13186.

106.   Knowland, D., Lilascharoen, V., Pacia, C.P., Shin, S., Wang, E.H.-J., and Lim, B.K.
(2017). Distinct Ventral Pallidal Neural Populations Mediate Separate Symptoms of
Depression. Cell *170*, 284-297.e18. https://doi.org/10.1016/j.cell.2017.06.015.

107.   Wang, F., Flanagan, J., Su, N., Wang, L.C., Bui, S., Nielson, A., Wu, X., Vo, H.T., Ma,
X.J., and Luo, Y. (2012). RNAscope: A novel in situ RNA analysis platform for formalin-
fixed, paraffin-embedded tissues. J. Mol. Diagn. *14*, 22–29. https://doi.org/10/dp2pmw.

108.   Pratelli, M., Thaker, A., Li, H., Godavarthi, S., and Spitzer, N.C. (2022). Phencyclidine
and methamphetamine cause cognitive deficits by changing pyramidal neuron transmitter
identity in the prefrontal cortex. Preprint at bioRxiv,
https://doi.org/10.1101/2022.06.16.496480 https://doi.org/10.1101/2022.06.16.496480.

109.   Ting, J.T., Daigle, T.L., Chen, Q., and Feng, G. (2014). Acute Brain Slice Methods for
Adult and Aging Animals: Application of Targeted Patch Clamp Analysis and Optogenetics.
In (Humana Press, New York, NY), pp. 221–242. https://doi.org/10.1007/978-1-4939-1096-
0_14.

110.   Krishnaswami, S.R., Grindberg, R.V., Novotny, M., Venepally, P., Lacar, B., Bhutani, K.,
Linker, S.B., Pham, S., Erwin, J.A., Miller, J.A., et al. (2016). Using single nuclei for RNA-

seq to capture the transcriptome of postmortem neurons. Nat. Protoc. *11*, 499–524.
https://doi.org/10/f8d3qc.

111.   Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for
piecewise rigid motion correction of calcium imaging data. J. Neurosci. Methods *291*, 83–94.
https://doi.org/10.1016/j.jneumeth.2017.07.031.

112.   Pnevmatikakis, E.A., Soudry, D., Gao, Y., Machado, T.A., Merel, J., Pfau, D., Reardon,
T., Mu, Y., Lacefield, C., Yang, W., et al. (2016). Simultaneous Denoising, Deconvolution,
and Demixing of Calcium Imaging Data. Neuron *89*, 285–299.
https://doi.org/10.1016/j.neuron.2015.11.037.

113.   Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B.L., Koay, S.A., Taxidis, J.,
Najafi, F., Gauthier, J.L., Zhou, P., et al. (2019). CaImAn an open source tool for scalable
calcium imaging data analysis. eLife *8*, e38173. https://doi.org/10.7554/eLife.38173.

114.   Bechtold, B., Fletcher, P., seamusholden, and Gorur-Shandilya, S. (2021).
bastibe/Violinplot-Matlab: A Good Starting Point. Version v0.1 (Zenodo).
https://doi.org/10.5281/zenodo.4559847 https://doi.org/10.5281/zenodo.4559847.

115.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species. Nat.
Biotechnol. *36*, 411–420. https://doi.org/10/gc87v6.

116.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,
Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell
data. Cell *184*, 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

117.   Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial
reconstruction of single-cell gene expression data. Nat. Biotechnol. *33*, 495–502.
https://doi.org/10.1038/NBT.3192.

118.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. Cell *177*, 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031.

119.   Griffiths, J.A., Richard, A.C., Bach, K., Lun, A.T.L., and Marioni, J.C. (2018). Detection
and removal of barcode swapping in single-cell RNA-seq data. Nat. Commun. *9*.
https://doi.org/10/gdw373.

120.   Lun, A.T.L., Riesenfeld, S., Andrews, T., Dao, T.P., Gomes, T., Marioni, J.C., and
participants in the 1st Human Cell Atlas Jamboree (2019). EmptyDrops: distinguishing cells
from empty droplets in droplet-based single-cell RNA sequencing data. Genome Biol. *20*,
63. https://doi.org/10/gfxdhf.

121.   Germain, P.-L., Sonrel, A., and Robinson, M.D. (2020). pipeComp, a general framework
for the evaluation of computational pipelines, reveals performant single cell RNA-seq
preprocessing tools. Genome Biol. *21*, 227. https://doi.org/10/ghb3tq.

122.   Germain, P.-L., and Lun, A. (2021). scDblFinder: scDblFinder. Version 1.4.0
        (Bioconductor version: Release (3.12)). https://doi.org/10.18129/B9.bioc.scDblFinder
        https://doi.org/10.18129/B9.bioc.scDblFinder.

123.   McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet
        Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst.
        8, 329-337.e4. https://doi.org/10/gg5m57.

124.   McCarthy, D.J., Campbell, K.R., Lun, A.T.L., Wills, Q.F., and Hofacker, I. (2017). Scater:
        pre-processing, quality control, normalization and visualization of single-cell RNA-seq data
        in R. Bioinformatics *33*, btw777. https://doi.org/10/gfgp3m.

125.   Choudhary, S., and Satija, R. (2022). Comparison and evaluation of statistical error
        models for scRNA-seq. Genome Biol. *23*, 27. https://doi.org/10.1186/s13059-021-02584-9.

126.   Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-
        cell RNA-seq data using regularized negative binomial regression. Genome Biol. *20*, 296.
        https://doi.org/10.1186/s13059-019-1874-1.

127.   Jolliffe, I.T., and Cadima, J. (2016). Principal component analysis: a review and recent
        developments. Philos. Trans. R. Soc. Math. Phys. Eng. Sci. *374*, 20150202.
        https://doi.org/10.1098/rsta.2015.0202.

128.   Becht, E., McInnes, L., Healy, J., Dutertre, C.-A., Kwok, I.W.H., Ng, L.G., Ginhoux, F.,
        and Newell, E.W. (2019). Dimensionality reduction for visualizing single-cell data using
        UMAP. Nat. Biotechnol. *37*, 38–44. https://doi.org/10.1038/nbt.4314.

129.   McInnes, L., Healy, J., and Melville, J. (2018). UMAP: Uniform Manifold Approximation
        and Projection for Dimension Reduction. ArXiv180203426 Cs Stat.

130.   Maaten, L. van der, and Hinton, G. (2008). Visualizing Data using t-SNE. J. Mach.
        Learn. Res. *9*, 2579–2605.

131.   Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden:
        guaranteeing well-connected communities. Sci. Rep. *9*, 5233.
        https://doi.org/10.1038/s41598-019-41695-z.

132.   Zappia, L., and Oshlack, A. (2018). Clustering trees: a visualization for evaluating
        clusterings at multiple resolutions. GigaScience *7*, giy083.
        https://doi.org/10.1093/gigascience/giy083.

133.   Phipson, B., Sim, C.B., Porrello, E.R., Hewitt, A.W., Powell, J., and Oshlack, A. (2022).
        propeller: testing for differences in cell type proportions in single cell data. Bioinformatics *38*,
        4720–4726. https://doi.org/10.1093/bioinformatics/btac582.

134.   Squair, J.W., Gautier, M., Kathe, C., Anderson, M.A., James, N.D., Hutson, T.H.,
        Hudelle, R., Qaiser, T., Matson, K.J.E., Barraud, Q., et al. (2021). Confronting false
        discoveries in single-cell differential expression. Nat. Commun. *12*, 5692.
        https://doi.org/10.1038/s41467-021-25960-2.

2080   135.   Costantini, D. (2020). Neuronal Topography in a Cortical Circuit for Innate Odor Valence.
2081   https://doi.org/10.7916/d8-ryea-m313.

2082   136.   Klein, S., Staring, M., Murphy, K., Viergever, M.A., and Pluim, J.P.W. (2010). elastix: a
2083   toolbox for intensity-based medical image registration. IEEE Trans. Med. Imaging *29*, 196–
2084   205. https://doi.org/10.1109/TMI.2009.2035616.

2085   137.   Shamonin, D.P., Bron, E.E., Lelieveldt, B.P.F., Smits, M., Klein, S., Staring, M., and
2086   Alzheimer's Disease Neuroimaging Initiative (2013). Fast parallel image registration on CPU
2087   and GPU for diagnostic classification of Alzheimer's disease. Front. Neuroinformatics *7*, 50.
2088   https://doi.org/10.3389/fninf.2013.00050.

2089   138.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
2090   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
2091   platform for biological-image analysis. Nat. Methods *9*, 676–682.
2092   https://doi.org/10.1038/nmeth.2019.

2093   139.   Carpenter, A.E., Jones, T.R., Lamprecht, M.R., Clarke, C., Kang, I.H., Friman, O.,
2094   Guertin, D.A., Chang, J.H., Lindquist, R.A., Moffat, J., et al. (2006). CellProfiler: image
2095   analysis software for identifying and quantifying cell phenotypes. Genome Biol. *7*, R100.
2096   https://doi.org/10.1186/gb-2006-7-10-r100.

2097   140.   McCullough, K.M., Morrison, F.G., Hartmann, J., Carlezon, W.A., and Ressler, K.J.
2098   (2018). Quantified Coexpression Analysis of Central Amygdala Subpopulations. eNeuro *5*.
2099   https://doi.org/10.1523/ENEURO.0010-18.2018.

2100

Figure S1. Additional information for imaging experiments. Related to **Figure 2.**

(A) Histologically verified placements for GRIN lenses (bars) in wild type animals infected with
AAV-hSyn-GCaMP8s for calcium imaging experiments. Horizontal lines indicate locations where
lens placements were visible in coronal sections at various positions in the plCoA anterior-
posterior axis. Different color bars represent placements in each of 13 animals. Those scored as
anterior are outlined in red, and those in posterior are outlined in blue.
(B) Histology for anterior plCoA (left) and posterior plCoA (right) lens implant and jGCaMP8s
(green) viral injection sites in representative animals. Scale bar for full size images, 500 μm;
scale bar for magnified images, 200 μm.
(C) Proportion responsive to each odor, pooled across regions.
(D) Heatmap of the normalized average pairwise Euclidean distance of odor response vectors
across biological replicates.

Fig. S2. Targeting of pICoA neurons for optogenetic stimulation. Related to **Figures 2**, **5**,
7, **S3**, and **S8**.

(**A**) Histologically verified placements for optic fiber implants (bars) and viral injection sites
(circles) in wild type animals infected with AAV-hSyn-ChR2-mCherry (blue) or AAV-hSyn-eYFP
(green) in topographic optogenetic cell body stimulation experiments in **Figure 1** and **S2**.
(**B**) Same as (**A**), but for cell-type specific optogenetic cell body stimulation experiments for
Slc17a7::Cre (top) and *Slc17a6::Cre* (bottom) animals infected with AAV-DIO-EF1A-ChR2-
eYFP (blue) or AAV-EF1A-DIO-eYFP (green) in **Figure 5**.
(**C**) Respective placements for fiber implants (bars) and injection sites (pICoA-NAc, circles;
pICoA-MeA, diamonds) in wild-type animals infected with AAV-hSyn-ChR2-mCherry (blue) or
AAV-hSyn-eYFP (green) in projection-specific optogenetic axon terminal stimulation
experiments in **Figure 7** and **S8.**
n denotes number of mice per group batched across 4-quad, elevated plus maze, and open
field test experiments, exceeding *n* values for individual experiments due to behavioral cohort
design (see STAR Methods). Relevant regions are highlighted in grey and outlined: pICoA (red),
NAc (purple, only in **C**), and MeA (pink, only in **C**). All mouse brain sections reproduced from
Paxinos and Franklin, 5th Edition, and numbers below all images denote its anterior-posterior
distance from bregma in this atlas.[36] All scale bars, 500 μm.

Fig. S3. Behavioral effects of topographic plCoA stimulation are limited to valence alone.
Related to **Figure 2**.

(A) Behavioral paradigm for optogenetic stimulation in the open field test.
(B-D) Optogenetic stimulation-induced change in time spent (**B**) and number of entries (**C**) into
the open arms, as well as distance traveled (**D**) in the elevated plus maze is not correlated to
anteroposterior axis position in plCoA.
(E-G) Effects of optogenetic stimulation of aplCoA neurons in the elevated plus maze in 1
minute bins over time (left) and during off and on periods (right).Photostimulation of aplCoA
neurons does not induce a significant change in time spent (**E**) and number of entries (**F**) into
the open arms, as well as distance traveled (**G**) in the elevated plus maze.
(H-J) Effects of optogenetic stimulation of pplCoA neurons in the elevated plus maze in 1minute
bins over time (left) and during off and on periods (right). Photostimulation of pplCoA neurons
does not induce a significant change in time spent (**H**) and number of entries (**I**) into the open
arms, as well as distance traveled (**J**) in the elevated plus maze.
(K) Behavioral paradigm for optogenetic stimulation in the open field test.
(L-N) Optogenetic stimulation-induced change in time spent in the center (**L**) and corners (**M**),
as well as distance traveled (**N**) in the open field test is not correlated to anteroposterior axis
position in plCoA.
(O-Q) Effects of optogenetic stimulation of aplCoA neurons in the open field test in 1 mintue
time bins (left) and during off and on periods (right). Photostimulation of aplCoA neurons does
not induce a significant change time spent in the center (**O**) and corners (**P**), as well as distance
traveled (**Q**) in the open field test.
(R-T) Effects of optogenetic stimulation of pplCoA neurons in the open field test in 1 minute time
bins (left) and during off and on periods (right). Photostimulation of pplCoA neurons does not
induce a significant change time spent in the center (**R**) and corners (**S**), as well as distance
traveled (**T**) in the open field test.
All "ON" and "OFF" comparisons in bar graphs and linear regressions are on a per 5-minute
basis. **(B-D, L-M)** Least-squares linear regression $\pm$ 95% confidence interval. Across panels: ns,
not significant. Additional specific details of statistical tests can be found in Supplemental Table
1.

Fig. S4. Additional information and quality-control for single-nucleus sequencing
experiments. Related to **Figure 3**.

(A) Representative images of tissue microdissection sites from aplCoA and pplCoA following
extraction and DAPI staining (blue). Scale bars, 500 μm.
(B) Location of all tissue sample sites for used for snRNA-seq, color coded by plCoA zone (n =
3 pools per zone, 4-11 sections per pool). Scale bars, 500 μm.
(C) Validation of nuclear enrichment after FANS. Ethidium homodimer-1 (EthD-1, red) labels
nuclei on a hemocytometer after sorting, with an absence of non-nuclear, EthD-1-negative
debris. Scale bar, 100 μm.
(D) Common gating strategy for FANS sorts for snRNA-seq in a representative sample.[102] Far
left, morphology gate on forward and side scatter area excludes likely debris. Middle left,
forward scatter gate excludes nuclear doublets with high forward scatter width. Middle right, side
scatter gate excludes nuclear multiplets with high side scatter width. Far right, stoichiometric
DRAQ7+ fluorescence allows enrichment of single nuclei and exclusion of debris and multiplets.
(E) Absolute number and proportion of snRNA-seq nuclei passing quality control filters from
each replicate in each plCoA zone (n = 27,726 in aplCoA, 19,406 in pplCoA, 3 libraries/batches
each).
(F) Violin plot of UMIs detected per snRNA-seq nucleus for each replicate, filtered at the median
per library + five times the median absolute deviation within each library (median 6081
UMIs/nucleus).
(G) Violin plot of genes detected per nucleus from each replicate, filtered at a minimum of 1000
features per nucleus (median 2547 genes/nucleus).
(H) Percent mitochondrial gene UMIs per snRNA-seq nucleus, filtered at median + five times
the median absolute deviation per library (median 0.02% mitochondrial UMIs/nucleus).
(I) Percent ribosomal gene UMIs per snRNA-seq nucleus, filtered at median + five times the
median absolute deviation per library (median 0.17% ribosomal UMIs/nucleus).
(J) Principal component analysis of pseudobulk snRNA-seq samples created from each batch,
colored based on their combination of zone and batch identity.
(K) Evaluation of transcriptomic homology between batches, where the distance matrix is based
on Spearman correlation between median expression of highly variable features for the whole
dataset, and the dendrogram was created via hierarchical clustering of batches on this
correlation matrix.
(L) UMAP of all snRNA-seq nuclei colored by both target region and batch identity.
(M) Relative proportion of nuclei of each type for all snRNA-seq batches.
Brain diagrams were reproduced from Paxinos and Franklin (2005).[36]

Fig. S5. Additional information and quality control for spatial gene expression. Related to
Figure 4.

(**A-C**) Allen ISH data for marker genes from molecular cell types adjacent to, but not within
plCoA. *Sim1* (**A**) marks cells in the NLOT, *Etv1* (**B**) marks cells in the BLA and posterior
basomedial amygdala, and *Fign* (**C**) marks cells in the cortex-amygdalar transition area, but
none of these mark cells in the plCoA.
(**D**) UMAP of all plCoA GABAergic neurons, colored by domain of origin.
(**E**) Relative proportion of molecular subtype nuclei from each domain within GABAergic
neurons. Dotted line indicates chance level for plCoA GABAergic neuron nuclei.
(**F**) UMAP of all plCoA OPCs, colored by domain of origin.
(**G**) Two-dimensional UMAP of astrocytes, colored by molecular cell type.
(**H**) Heatmap of astrocyte subtype marker genes.
(**I**) UMAP of all plCoA astrocytes, colored by domain of origin.
(**J**) Left, relative proportion of molecular subtype nuclei from each domain within astrocytes.
Dotted line indicates chance level for plCoA astrocyte nuclei. Right, relative abundance of each
astrocyte subtype within plCoA.
(**K**) Two-dimensional UMAP of immune cells, colored by molecular cell type.
(**L**) Heatmap of immune cell subtype marker genes.
(**M**) UMAP of all plCoA immune cells, colored by domain of origin.
(**N**) Left, relative proportion of molecular subtype nuclei from each domain within immune cells.
Dotted line indicates chance level for plCoA immune cell nuclei. Right, relative abundance of
each immune cell type within plCoA.
(**O**) Two-dimensional UMAP of VLMC nuclei, colored by molecular cell type.
(**P**) Heatmap of VLMC subtype marker genes.
(**Q**) UMAP of all plCoA VLMC nuclei, colored by domain of origin.
(**R**) Left, relative proportion of molecular subtype nuclei from each domain within VLMCs. Dotted
line indicates chance level for plCoA VLMC nuclei. Right, relative abundance of each VLMC
subtype within plCoA.
(**S**) Left, representative section on a Visium slide capture area stained with hematoxylin and
eosin. Right, representative section with capture spots overlaid (grey) and plCoA-overlapping
spots highlighted (red).
(**T**) Violin plots of quality metrics for individual Visium sections on a per-spot basis in plCoA-
overlapping spots (N = 21 sections). Upper left, UMIs per spot; upper right, features per spot;
lower left, proportion mitochondrial UMIs per spot; lower right, proportion ribosomal UMIs per
spot.
(**U**) Number of plCoA-overlapping capture spots per section (n = 3,616 total spots).
(**V**) Two-dimensional UMAP of all plCoA-overlapping capture spots, colored by section of origin.
(**W**) Evaluation of transcriptomic homology between sections, where the distance matrix is
based on Spearman correlation between median expression of highly variable features for the
whole dataset, and the dendrogram was created via hierarchical clustering of sections on this
correlation matrix.
Across panels: ** $p < 0.01$; *** $p < 0.001$; ns, not significant. Additional specific details of
statistical tests can be found in Supplemental Table 1.

Fig. S6. Additional information for Cre-dependent molecularly targeted chemogenetic
inhibition experiments. Related to **Figure 5**.

(A) Representative histology for inhibition experiments for AAV-hM4D(Gi)-mCherry-infected
Slc17a6::Cre (left) or *Slc17a7::Cre* (right) animals.
(B) Strategy for chemogenetic inhibition in the open field test and elevated plus maze.
(C-E) Effect of chemogenetic inhibition of plCoA$^{Slc17a6+}$ neurons in the elevated plus maze.
Inhibition does not induce a significant change in time spent (**C**) or number of entries (**D**) into the
open arms, as well as distance traveled (**E**) in the elevated plus maze.
(F-H) Effect of chemogenetic inhibition of plCoA$^{Slc17a6+}$ neurons in the open field test. Inhibition
does not induce a significant change in time spent in the center (**F**) or corners (**G**), as well as
distance traveled (**H**) in the open field test.
(I-K) Effect of chemogenetic inhibition of plCoA$^{Slc17a7+}$ neurons in the elevated plus maze.
Inhibition does not induce a significant change in time spent (**I**) and number of entries (**J**) into
the open arms, as well as distance traveled (**K**) in the elevated plus maze.
(L-N) Effect of chemogenetic inhibition of plCoA$^{Slc17a7+}$ neurons in the open field test. Inhibition
does not induce a significant change in time spent in the center (**L**) or corners (**M**), as well as
distance traveled (**N**) in the open field test.
(**O**) Histologically verified placements for viral injection sites in *Slc17a6::Cre* animals infected
with AAV-DIO-hSyn-mCherry (red) or AAV-DIO-hSyn-hM4D(Gi) (light orange) in molecularly
targeted chemogenetic inhibition experiments.
(**P**) Same as (**O**), but in *Slc17a7::Cre* animals.
n denotes number of mice per group batched across 4-quad, elevated plus maze, and open
field test experiments, exceeding *n* values for individual experiments due to behavioral cohort
design (see STAR Methods). All mouse brain sections reproduced from Paxinos and Franklin,
5[th] Edition, with plCoA highlighted in grey and outlined in purple, and numbers below all images
denote its anterior-posterior distance from bregma in this atlas.[36] All scale bars, 500 µm. Across
panels: ns, not significant. Additional specific details of statistical tests can be found in
Supplemental Table 1.

Fig. S7. Additional information for plCoA anterograde tracing experiments. Related to
Figure 6.

(A) Schematic for general anterograde output mapping strategy, where a virus labeling cell
bodies in plCoA with mRuby and presynaptic terminals in downstream regions with
synaptophysin-bound eYFP.
(B) Histological image of plCoA injection site in a representative animal. Scale bar, 200 μm.
(C) Histologically verified placements for viral injection sites in initial anterograde tracing
experiments.
(D) Schematic for dual-color topographic anterograde tracing: two counterbalanced fluorophores
were injected into aplCoA and pplCoA, and each color was quantified in major projection targets
of plCoA.
(E) Histologically verified placements for viral injection sites in dual-color anterograde tracing
experiments.
(F) Schematic for Cre-dependent anterograde output mapping strategy: a Cre-dependent virus
expressing eYFP was injected into either an *Slc17a6::Cre* or *Slc17a7::Cre* animal to determine
relative output enrichment for either broad cell type.
(G) Histologically verified placements for viral injection sites in Cre-dependent molecular
anterograde tracing experiments.
(H) Schematic of anterograde viral strategy to explore collateralization of plCoA MeA and NAc
projection neurons to other regions.
(I) Histologically verified placements for viral injection sites in collateralization experiments.
Relevant regions are highlighted in grey and outlined: plCoA (red), NAc (purple, only in **I**), and
MeA (pink, only in **I**). All mouse brain sections reproduced from Paxinos and Franklin, 5th
Edition, and numbers below all images denote its anterior-posterior distance from bregma in this
atlas.[36] All scale bars 500 μm unless noted elsewhere.

Fig. S8. Manipulation of plCoA projections to MeA or NAc in either direction does not
change features of behavior unrelated to innate valence. Related to **Figure 7**.

(A-C) Effects of optogenetic stimulation of plCoA terminals in MeA in the elevated plus maze in
time bins of 1 minute (left) and during off and on periods (right). Photostimulation does not
induce a significant change in time spent (**A**) and number of entries (**B**) into the open arms, as
well as distance traveled (**C**) in the elevated plus maze.
(D-F) Effects of optogenetic stimulation of plCoA terminals in NAc in the elevated plus maze in
time bins of 1 minute (left) and during off and on periods (right). Photostimulation does not
induce a significant change in time spent (**H**) and number of entries (**I**) into the open arms, as
well as distance traveled (**J**) in the elevated plus maze.
(G-I) Effects of optogenetic stimulation of plCoA terminals in MeA in the open field test in time
bins of 1 minute (left) and during off and on periods (right). Photostimulation does not induce a
significant change time spent in the center (**G**) and corners (**H**), as well as distance traveled (**I**)
in the open field test.
(J-L) Effects of optogenetic stimulation of plCoA terminals in NAc in the open field test in time
bins of 1 minute (left) and during off and on periods (right). Photostimulation does not induce a
significant change time spent in the center (**G**) and corners (**H**), as well as distance traveled (**I**)
in the open field test.
(**M-O**) Effects of chemogenetic inhibition of plCoA-MeA projection neurons in the elevated plus
maze. Inhibition does not induce a significant change in time spent (**M**) and number of entries
(**N**) into the open arms, as well as distance traveled (**O**) in the elevated plus maze.
(**P-R**) Effects of chemogenetic inhibition of plCoA-MeA projection neurons in the open field test.
Inhibition does not induce a significant change in time spent in the center (**P**) and corners (**Q**),
as well as distance traveled (**R**) in the open field test.
(**S-U**) Effects of chemogenetic inhibition of plCoA-NAc projection neurons in the elevated plus
maze. Inhibition does not induce a significant change in time spent (**S**) and number of entries
(**T**) into the open arms, as well as distance traveled (**U**) in the elevated plus maze.
(**V-X**) Effects of chemogenetic inhibition of plCoA-NAc projection neurons in the open field test.
Inhibition does not induce a significant change in time spent in the center (**V**) and corners (**W**),
as well as distance traveled (**X**) in the open field test.
Across panels: ns, not significant. Additional specific details of statistical tests can be found in
Supplemental Table 1.