Discovery of a Heparan Sulfate Binding Domain in Monkeypox Virus H3 as an Anti-poxviral Drug Target Combing AI and MD Simulations

Bin Zheng[1,#], Meimei Duan[3,#], Yifen Huang[1,#], Shangchen Wang[1,#], Jun Qiu[1], Zhuojian Lu[1], Lichao Liu[1], Guojin Tang[1], Lin Cheng[3]* and Peng Zheng[1,2]*

1. State Key Laboratory of Coordination Chemistry, Chemistry and Biomedicine Innovation Center (ChemBIC), School of Chemistry and Chemical Engineering, Nanjing University, Nanjing, Jiangsu, 210023, China
2. Department of Intensive Care Unit, Nanjing Drum Tower Hospital, Affiliated Hospital of Medical School, Nanjing University, Nanjing, Jiangsu Province, China.
3. Institute for Hepatology, National Clinical Research Center for Infectious Disease, Shenzhen Third People's Hospital, Shenzhen, 518112, China

#These authors contributed equally to this work
*Email: pengz@nju.edu.cn & chenglin252@163.com

Abstract: Viral adhesion to host cells is a critical step in infection for many viruses, including monkeypox virus (MPXV). In MPXV, the H3 protein mediates viral adhesion through its interaction with heparan sulfate (HS), yet the structural details of this interaction have remained elusive. Using AI-based structural prediction tools and molecular dynamics (MD) simulations, we identified a novel, positively charged α-helical domain in H3 that is essential for HS binding. This conserved domain, found across *orthopoxviruses*, was experimentally validated and shown to be critical for viral adhesion, making it an ideal target for antiviral drug development. Targeting this domain, we designed a protein inhibitor, which disrupted the H3-HS interaction, inhibited viral infection *in vitro* and viral replication *in vivo*, offering a promising antiviral candidate. Our findings reveal a novel therapeutic target of MPXV, demonstrating the potential of combination of AI-driven methods and MD simulations to accelerate antiviral drug discovery.

Introduction

The mpox virus (MPXV), a zoonotic pathogen within the *Orthopoxvirus* genus,

has emerged as a global health concern. This genus also includes the Variola virus

(VARV), the causative agent of smallpox, and Vaccinia virus (VACV), which has been

used as a live vaccine against smallpox[1, 2] (Fig. 1A). Historically endemic to Central

and West Africa, MPXV gained international attention during the global outbreaks in

2022 (clade IIb) and 2024 (clade Ib), leading the World Health Organization (WHO)

to declare it a public health emergency of international concern twice within three

years[3-5]. Although MPXV has a lower fatality rate than smallpox (2-10% versus 30%),

the absence of specific antiviral treatments continues to pose significant health risks[6-

11].

Viral adhesion to host cells is a critical initial step in the infection process for

many viruses. Proteins that mediate these interactions have been identified as essential

therapeutic targets, offering promising opportunities for antiviral therapies. In

orthopoxviruses such as monkeypox virus (MPXV), the H3 protein plays a pivotal

role in facilitating viral adhesion by binding to cell-surface heparan sulfate (HS)[12-14], a

glycosaminoglycan essential for viral entry. H3-specific neutralizing monoclonal

antibodies have shown to protective effects for rabbits and immunodeficient mice

against lethal poxvirus infections[15]. And H3 is a key component in recent mpox

vaccines[16, 17]. Despite the central role of H3 in MPXV infection and protection, the

structural details of its specific interaction with HS remain elusive.

Understanding the molecular mechanisms underlying the H3-HS interaction is

critical for advancing antiviral drug development. H3 is highly conserved across

poxviruses (Fig. 1A), including variola and vaccinia viruses, making it a potential

target for broad-spectrum therapies. The recent resurgence of MPXV infections has

underscored the urgent need for specific antiviral treatments. Given the conserved

nature of H3 across all known MPXV lineages, including Clade Ib, which has been

linked to the 2024 outbreak, designing therapies that target H3 could offer a robust

solution to current and future outbreaks.

However, efforts to resolve the complete structure of H3-HS complex, have been

hampered by the dynamic and flexible nature of HS and the tendency of H3 self-

cleavage. Classic structural biology techniques, such as X-ray crystallography, have

been unable to capture the full interaction between H3 and HS due to these

complexities. Therefore, alternative approaches are required to fully elucidate this

critical interaction.

To address these challenges, we employed advanced AI-based protein structural

prediction and generation tools, including AlphaFold2(AF2) and RFdiffusion,

alongside classic molecular dynamics (MD) simulations. These state-of-the-art

computational techniques enabled us to identify key HS-binding sites within H3 and

to characterize the binding mechanisms of the previously uncharacterized α-helical
domain[18-22]. Corroborated by various experimental methods[23-26], the use of AI-driven
tools proved pivotal in overcoming the limitations of traditional structural methods,
allowing us to predict and validate the structure-function relationship of the H3-HS
interaction. A powerful approach for unraveling complex protein-glycan interactions
is demonstrated, offering new pathways for the development of broad-spectrum
antiviral therapies[27, 28].

90
Fig. 1 **Structural Analysis of the H3 Protein in Monkeypox Virus** (A)
Phylogenetic tree depicts the evolutionary relationships of H3 within the *Poxviridae*
family, highlighting MPXV (blue circle), VARV (red circle), and VACV (green
circle). (B) Schematic of MPXV adhesion to the cell surface. Viral particles bind to
cell surface via specific interaction such as between adhesion protein H3 and HS,
followed by membrane fusion mediated by the fusion complex, allowing entry into
the host cell. (C) The amino acid sequence of MPXV H3 shows the newly discovered
helical structure (240-282, highlight in yellow), the Mg(II) (green) binding site and
other potential HS binding motifs (blue underlines). (D) AF2 prediction of MPXV H3
structure. The left and right panels show different orientations of the H3 structure
(rotated 180°). The blue region corresponds to H3(1-239), which has a homologous
crystal structure (VACV H3, PDB code: 5EJ0). The yellow region represents the AF2-
predicted structure of H3(240-282), which remains unresolved in the crystal structure
of the homologous protein. All potential GAGs-binding motifs are highlighted with a
yellow background. (E) MD simulation snapshot of H3 on a DPPC membrane. H3 is

anchored to the membrane through its transmembrane region (residues 283-306, in
gray).
108
Result
Identification of a Novel Helical Domain for HS Binding in H3
The H3 protein of MPXV is a 324-amino-acid transmembrane protein conserved
across all 12 identified *orthopoxviruses*. Previous studies have shown that its
extracellular domain (residues 1-282) binds to cell-surface heparan sulfate (HS)[29, 30],
facilitating viral entry in conjunction with other adhesion proteins and the fusion
protein system (Fig. 1B)[31]. Inhibiting the HS binding to H3, therefore, represents a
promising strategy for developing broad-spectrum antiviral therapies against
orthopoxviruese. While most structure of the extracellular region of Vaccinia virus H3
(residues 1-240; PDB code: 5EJ0) has been solved, the dynamic and heterogeneous
nature of HS as a biopolymer complicates the identification of precise binding sites.
To identify HS binding sites on the H3 protein, we first performed a
comprehensive sequence and structural analysis. As a highly negatively charged
glycosaminoglycans, it is believed that HS interacts with positive charge residues on
proteins by electrostatic interaction. Previous studies have indicated that the positively
charged Mg (II) ion in H3 is critical for HS binding. Therefore, we focused on regions
rich in basic amino acids—lysine (K), arginine (R), and histidine (H). Given that only
the H3 structure from VACV was available[28], we used AlphaFold2 (AF2) to model
the MPXV H3 structure (Fig. S1A), Our analysis revealed three potential HS binding
motifs rich in positively charge residues: Motif 1 (K95, R96, R100), Motif 2 (K141,
K146, K147), and Motif 3 (K161, K163)[32], all located within the homologous
structure of VACV H3 (Fig. 1C, D).
Interestingly, we also identified a cluster of seven additional positively charged
residues (R242, H247, R248, K253, R259, K266, R267) in the protein C-terminal
region. This region had not been previously structurally characterized due to its self-
cleavage during sample preparation for X-ray diffraction studies, but was retained in
all other biochemical and viral assays[28-30]. AF2 predicted these residues to form a
three-helical domain with high confidence (pLDDT score: 82.9 for overall structural;
80.6 for the helical domain, 100 for the maximum score), suggesting it may serve as
an HS binding site as a structured domain (Fig. 1D). We also tested other models,
such as ESMFold (pLDDT:70.6; 50.7) and RoseTTAFold2 (pLDDT: 70.2; 50.7) (Fig.
S1B&C) [33, 34]. They all predict a helical structure. But their results are different in
detail and had lower confidence scores. Thus, we used AF2 for further studies.
Another concern is that the helical domain near the virus surface. Thus, we modeled
the complete structure of H3 in virus using a model DPPC
(dipalmitoylphosphatidylcholine) membrane. It showed a dynamic movement
between the helical domain and the membrane, with a maximum distance exceeding 1
nm (Fig. 1E, Fig. S1D&E, Supplementary Video 1). Thus, enough space be present
for possible HS binding.
Flexible molecular docking and MD simulations identified the exact HS binding
sites in the H3 protein.[35, 36]. We used a 20-repeat HS unit with a specific composition

for detailed analysis ([IdoA2S-GlcNS6S-IdoA-GlcNS(3,6S)]$_5$) (Fig. S2). A 500 ns MD simulation established the equilibrium conformation of H3, and subsequent docking with AutoDock Vina identified four high-scoring conformations for each motif (Fig. 2A, Fig. S3). The docking score calculated by the software reflects the total energy of interactions between molecules. Lower scores (more negative values) indicate more stable binding between the molecules. To address the dynamic nature of HS, we extended our simulations to 3 times additional 1000 ns MD simulations for each conformation. RMSD (Root Mean Square Deviation) analysis demonstrated notably stable binding for HS to both Motif 1 and the newly identified helical domain (Supplementary Video 2), with RMSD values consistently below 4 nm, indicating minimal structural fluctuations during interaction (Fig. 2B).

Moreover, we quantitatively analyzed the binding free energy between H3 and HS in different conformational states using umbrella sampling (US) techniques. A harmonic potential was applied to HS and gradually pulled away from H3, generating a series of reaction coordinates (Fig. 2C, Fig. S4, Supplementary Video 3). The green force-extension curve represents the force during this stretching process. This curve, along with the corresponding PMF (Potential of Mean Force) variation shown in Figure S4, demonstrated the free energy changes throughout the umbrella sampling process, revealing that the helical domain possesses the highest binding affinity with a ΔG of -45 kCal/mol (Fig. 2D).

Previous studies have suggested the critical role of the Mg(II) ion in stabilizing HS binding, where its removal reduces binding efficiency[29]. Indeed, the α-helical domain is very close to the Mg(II)-bound region in H3. Notably, these two parts appear to form a distinct cavity-like binding pocket for HS (Fig. 1E, Fig. 2E). Thus, we conducted a 1000 ns REMD (Replica Exchange Molecular Dynamics) simulation of the H3-HS complex to further understand the binding dynamics. REMD, an advanced sampling technique, helps overcome the kinetic barriers typically encountered in conformational transitions by utilizing a series of temperature-controlled replicas to enhance the exploration of the conformational landscape[37]. For this study, we set up 64 replicas over a temperature range from 310K to 387K, cumulatively simulating for 64 μs. This comprehensive simulation captured the HS molecule progressively binding deeper into a cavity formed by the α-helical domain, eventually interacting with the Mg (II) ion as hypothesized (Fig. 2E, Supplementary Video 4).

The free energy landscape, mapped out from these simulations, shows HS transitioning from a stable interaction with the helical domain across an energy barrier into the Mg (II)-enhanced cavity. This dual interaction—first with the helical domain and then with the Mg(II) site—was consistently observed to stabilize the complex further, evidenced by a significantly lower binding free energy of -67 kCal/mol (Fig. S4)[38]. These results underscore the dynamic nature of the H3-HS interaction and validate our model of sequential binding, which could be critical for designing inhibitors that target these specific interactions.

Given the crucial role of electrostatic interactions in the HS-H3 binding process, we conducted an exhaustive analysis of salt bridge formations across all MD

simulation trajectories. This analysis focused on the formation of salt bridges over
time between the basic amino acids (R, K, H) of H3 and HS, with brighter areas
indicating more frequent formation of salt bridges and robust interaction (Fig. 2F, Fig.
S5). A bar graph accompanied these maps, quantifying the average number of salt
bridges each amino acid formed during the simulations, further illustrating the
intensity and frequency of these interactions. Our findings demonstrated a high
affinity of HS for the helical domain of H3, with basic amino acids in this region
forming more salt bridges compared to other parts of the protein. The surface hotspot
map of H3 (Fig. 2G) visually highlighted these interactions, with varying shades of
blue representing the frequency of salt bridge formation, emphasizing their
distribution and intensity. In the helical domain, a focus view showed that the 1st, 7th,
9th, and 18th sugar units of HS formed salt bridges with R248, R242, R267, and R259
of H3, respectively (Fig. 2H, left), while the sulfate group on the 2nd sugar side chain
of HS bound to Mg(II) (Fig. 2H, right).

Fig. 2 **Identification of HS Binding Motifs in H3 by MD Simulation and Docking**

(A-B) Cartoons show docking results of HS with Motifs 1, 2, 3, and the helical

domain, respectively. Panels (B) show RMSD values from 1μs MD simulations,

color-coded to match the configurations in (A). (C) Schematics illustrates the reaction

coordinate in umbrella sampling, highlighting HS dissociation from H3 with a green

force curve. (D) Histogram shows the binding free energies for HS-H3 interactions in

motifs 1, 2, 3, and the helical domain. (E) A free energy landscape map from a 1000

ns REMD simulation shows HS binding configurations to the helical and Mg(II)

regions. (F) Panel provides salt bridge formation statistics between HS and H3's basic

amino acids, with a bar chart of average formations. (G) A surface plot shows the

frequency of salt bridge formations within H3, with areas of frequent formations in

blue. (H) Detailed views of HS-H3 interactions, with the left image showing salt

bridges and the right image displaying electrostatic interactions with Mg(II). This

panel illustrates the impact on HS binding stability to H3 following the removal of Mg(II) during the simulation. (J) The effects of mutating all basic amino acids in the helical domain on the binding stability of HS are shown. (K) The "palm-binding" model is depicted where HS is secured by the helical "fingers" and interacts with the Mg(II)-bound "palm."

To further prove these findings, we performed two control experiments. First, we removed the Mg(II) ion during MD simulation. According to the results of umbrella sampling, we extended the simulation of the most stable H3-HS binding conformation to 1000 ns, then removed the Mg(II) ion and continued the simulations. The resulting increase in RMSD fluctuations confirmed the stabilizing role of Mg(II) in the binding process (Fig. 2I, Supplementary Video 5). Second, we mutated all seven positively charged amino acids in the helical domain to serine, and name it H3 (uncharged). While AF2 predicted that the helical structure would remain intact (Fig. S6), a significant alteration in HS binding dynamics was observed, supporting the critical role of these charged residues in stabilizing the interaction (Fig. 2J, Supplementary Video 6).

Our interaction model revealed that the α-helical domain of H3 functions like a thumb, guiding HS into a stable binding position, while the Mg(II) region acts like a palm, securing HS in place. This dynamic interaction allows HS to transition from an initial surface-level binding to being deeply anchored within the protein cavity, illustrating a complex yet well-organized binding process (Fig. 2K).

Bioinformatic analysis of the H3 protein across the *Poxviridae* family highlights its evolutionary conservation and its significance in HS binding. Given the crucial role of electrostatic interactions in the H3-HS binding process, we analyzed the charge distribution within the helical domain. To date, H3 proteins have been discovered in 66 of 118 known *Poxviridae* species. Using NCBI's BLAST, we retrieved sequences of these 66 H3 proteins and performed multiple sequence aligment[39]. A heatmap detailing the side-chain charge distribution at a physiological pH of 6.5—mimicking the microenvironment of H3-HS interaction—revealed a significant concentration of positive charges at the helical domain (Fig. 3A). Furthermore, structural predictions using AF2 showed that all these 66 sequences generally share a similar architecture to MPXV H3, particularly in the presence of the α-helical domain (Fig. S7).

Notably, this domain consistently exhibited an accumulation of basic amino acids such as lysine, arginine, and histidine, which are essential for binding the negatively charged HS. These amino acids, including residues identified as 358, 361, 367, 373, 379, 386, and 387 in *Poxviridae* H3 (global position in alignment), corresponding to residues H239, R242, R248, K253, R259, K266, R267 in MPXV H3, show high conservation (Fig. 3B, Fig. S8). Surface charge analysis of H3 proteins from four *orthopoxviruses*, including MPXV, VACV, VARV, and CPXV confirmed the predominance of strong positive charges in their α-helical domains (Fig. 3C), crucial for effective binding to HS. This consistent feature across different viruses underscores the evolutionary importance of the helical domain, suggesting a universal

266 mechanism in orthopoxviral adhesion that could be targeted in antiviral strategy.

Experimental Confirmation of the Structure and Function of α-Helical Domain

To validate the computational predictions, we first conducted experiments using
atomic force microscopy-based single-molecule force spectroscopy (AFM-SMFS).
This technique, widely used for studying protein (un)folding, allowed us to directly
assess the structure and stability of the α-helical domain within the H3 protein[40-42].
We engineered the H3 construct with two GB1 domains, each 18 nm in length upon
unfolding, and one Cohesin module for precise measurement (Fig. 3D)[43-45]. These
proteins are immobilized on the surface by click chemistry and enzymatic ligations
(Fig. S9)[46-48]. During the experiment, the coated AFM tip with GB1 and Dockerin
initiated a Coh-Doc interaction, producing a characteristic sawtooth-like force-
extension curve as it was stretched (Fig. 3D&E). Notably, two peaks corresponding to
the unfolding of the α-helical domain were observed. The first peak showed a contour
length increment (ΔLc) of 13 nm, closely matching the theoretical unfolding length
for the 42 amino acid-long α-helical domain of H3 (residue 240-282, 42aa*0.36
nm/aa-2.6 nm)[49]. The measured unfolding force was 67.4 ± 2.1 pN (mean $\pm$ SEM,
n=139), while the second peak indicated the unfolding of the remaining structure of
H3, with a ΔLc of 84 nm (Fig. 3F). The total unfolding, represented by a cumulative
ΔLc of 97 nm, confirmed the structured and stable nature of the α-helical domain
within H3.

Fig. 3 **Charge Characteristics and Structural Analysis of H3 Protein** (A) The heatmap shows the amino acid charge distribution in the H3 protein across 66 *Poxviridae* viruses, following multiple sequence alignment. Blue indicates areas with more positive charges, and red indicates more negative charges. The accompanying curve shows the average charge of all amino acids in *Poxviridae* H3. (B) Logo plot of the amino acid sequence of the helical region, highlighting the conservation of basic amino acids at specific positions. (C) The surface charge analysis of H3 from the MPXV, VACV, VARV, and cowpox viruses (CPXV) with the helical domain showing a significantly positive charge. (D) Schematic of the single-molecule force spectroscopy unfolding experiment on H3, illustrating the unfolding process of the helical domain (yellow) followed by the main body (blue). (E) Representative curves of H3 unfolding, color-coded to show the helical region (yellow), main body (blue), and full-length (purple) unfolding. (F) Histograms depict the force spectroscopy signals for helical domain, main body, and full-length unfolding, with ΔLc statistics provided. The inset shows a Gaussian fit of unfolding forces.

We further explored the functional role of the α-helical domain in mediating H3's binding to HS using AFM on live cells[50, 51]. The H3 protein was linked to the AFM tip and approached cultured CHO-K1 cells, which express HS, mounted on a Petri dish. Inverted microscopy was used to precisely control the positioning of the AFM tip. Upon contact, the interaction between H3 and HS was initiated, and the binding event was monitored by retracting the tip to generate a force-extension curve. A distinct peak, corresponding to H3-HS interaction, was observed with a dissociation force of 33.7 ± 0.2 pN and a binding probability of 43% (Fig. 4A, B). Repeated trials across different cell surface areas produced a detailed force map, quantifying the distribution of HS unbinding forces and further confirming the critical role of the helical domain in viral adhesion (Fig. 4C, D).

Fig. 4 **Analysis of the Helical Domain's Interaction with HS at Cellular Level** (A)
Schematic of the cell force spectroscopy experiment setup shows three scenarios:
wild-type H3 on an AFM tip interacting with HS on CHO-K1 cells, mutation of all
basic amino acids in the H3 helical region to serine, and cells treated with HS
hydrolase to remove surface HS before testing. (B) Force-extension curves depict
interactions for the wild type, mutant, and control groups, marked with blue and red
asterisks for dissociation events. An inset shows the optical microscope positioning
the AFM probe. (C) Histograms of dissociation signals comparing the wild type,
mutant, and control groups, with an inset detailing the surface distribution of
dissociation forces. (D) Statistical graph showing binding probabilities for different
groups, highlighting significant differences determined by t-tests (p<1e-5). (E) FCM
results illustrate interactions of wild-type (WT) and Uncharged H3 fused with eGFP
with CHO-K1 cells, alongside GFP control (green) and cell-only control (Blank,
grey). (F) Statistical analysis of FCM data, showing significant differences between
groups as determined by t-tests. *****, P<1e-5

Moreover, we performed mutagenesis studies on the α-helical domain by
replacing all positively charged residues with serine. AFM experiments on this variant
H3(uncharged) showed a detectable force peak with a lower unbinding force of $28.8 \pm
0.2$ pN, compared to 33.7 ± 0.3 pN for the wild type—representing a 14% reduction
(ΔForce > 2*SEM, 95% CI)—and a significant decreased HS binding probability
from 43% to 25% (Fig. 4C, D). These results suggest that the basic amino acids in the
helical domain are crucial for HS binding. To confirm the specificity of the
interaction, CHO-K1 cells were treated with heparinase II, an enzyme that degrades

HS. This treatment further reduced the dissociation force to 23.0 pN and the binding probability to 12.6%, confirming that the observed interactions were specific to the HS-H3 complex.

Further validation was carried out using flow cytometry (FCM). We fused eGFP, a fluorescent reporter protein, to the C-terminus of both the H3(WT) and the H3(uncharged). Following incubation of these fusion proteins with cells, FCM analysis showed a significant reduction in mean fluorescence intensity in cells treated with H3(uncharged) compared to H3(WT) (Fig. 4E). Statistical analysis of multiple experiments highlighted a significant difference between the two (Fig. 4F). These findings, along with the results from MD simulations and AFM force spectroscopy, underscore the importance of the positively charged amino acids in the α-helical domain of H3 for efficient HS binding.

The Helical Domain Serves as a Target for H3 Infection Inhibition

Targeting the α-helical domain of H3 as H3 binding sites, we *de novo* designed a series of inhibitors. Using deep-learning model RFdiffusion, we designed a series of inhibitors for the H3-HS interaction and named it AI-PoxBlock (Fig. 5A). Sequence recovery for each series was performed on 1000 backbones using ProteinMPNN, generating ten sequences per backbone. These designs were further validated through complex structure predictions using AF2-multimer, and candidates were selected based on PAE (Predicted Aligned Error) values below 7 and iPTM (interface predicted Template Modelling) scores above 0.9. The predicted structures of these candidates also exhibited RMSD values of less than 2 nm when compared to the RFdiffusion-generated structures (Fig. S10). To assess the binding capabilities of these inhibitors, we conducted 500 ns MD simulations, analyzing RMSD trajectories. Five inhibitors—AI-PoxBlock302, 602, 614, 723, and 761—demonstrated stable RMSD values under 0.4 nm, confirming their potential for stable binding to the helical domain of H3 (Fig. S11, S12, Supplementary Video 7).

Although AI-PoxBlock302 could not be successfully expressed, the remaining four inhibitors underwent further testing in FCM to evaluate their efficacy in inhibiting H3 binding to CHO-K1 cell surfaces. AI-PoxBlock723 stood out, significantly reducing H3 binding at a concentration of 10 μM, while the other inhibitors displayed no tendency to inhibit H3 binding to HS on cell surfaces (Fig. 5B, Fig. S13). Further analysis using biolayer interferometry (BLI) confirmed that AI-PoxBlock723 binds to H3 with an equilibrium dissociation constant (K_D) of 9 μM (Fig. 5C). To ensure specificity, we also assessed the interaction between AI-PoxBlock723 and a truncated version of H3 (residues 1-239), lacking the α-helical domain. BLI results indicated a marked reduction in binding, reinforcing the specificity of AI-PoxBlock723 for the helical domain (Fig. S14). Circular Dichroism spectroscopy also confirmed that AI-PoxBlock723 predominantly consists of α-helices, consistent with its design specifications (Fig. S15).

To evaluate the antiviral activity of AI-PoxBlocks, the inhibitors were serially diluted and incubated with the clade II MPXV isolate SZTH42 before infecting Vero E6 cells. After overnight culture, virus infection foci were determined as previously described.[52] Notably, AI-PoxBlock723 showed an antiviral effect with the half

maximal inhibitory concentration (IC_{50}) of 88.6 μM, whereas other AI-PoxBlocks exhibited no inhibitory efficiency (Fig. 5D). Given the high conservation of the H3 helical domain across the four *orthopoxviruses* pathogenic to humans (Fig. S16), we also assessed AI-PoxBlock activity against VACV infection. Consistently, AI-PoxBlock723 is also effective for inhibiting VACA infection in vitro with a similar IC50 of 86.7 μM (Fig. S17), suggesting the helical domain as a universal antiviral target for *orthopoxvirus*.

To evaluate the efficacy of AI-PoxBlock723 in vivo, groups of BALB/c mice were challenged with 4×10^5 FFU (focus forming unit) MPXV/SZTH42 as we previously reported[53]. Mice were administered intraperitoneally with single dose of AI-PoxBlock723 (10 mg/kg) or PBS immediately after MPXV challenge, and were sacrificed 4 days post-infection (dpi) for the determination of pulmonary MPXV titers. The MPXV infectious particles and genome copies in lungs of the AI-PoxBlock723 treated mice were significantly lower than that in the control mice (Fig. 5E-F, Fig. S18).

These results demonstrate the efficacy of the designed inhibitors targeting the H3 helical domain and validate the crucial role of this domain in viral adhesion, providing a promising foundation for the development of novel antiviral agents.

Fig. 5 **Inhibitor Design Targeting the H3-HS Interaction and Efficacy of AI-PoxBlock723** (A) Diagrams depict protein inhibitors designed to target the H3 helical region, created using RFdiffusion. Sequences capable of folding into the target

scaffold structures were generated using ProteinMPNN, and were validated through
AF2, followed by 500 ns MD simulations for structural stability and interaction
scoring. (B) FCM analysis demonstrates the inhibitory effect of AI-Poxblock723 at
various concentrations (0, 10, 50 and 100 μM, x axis). The control group consisted of
2 μM H3-eGFP without the inhibitor, with the relative fluorescence intensity
normalized to 1. (C) BLI confirms direct interaction between AI-Poxblock723 and
the H3 helical domain. (D) Graphs display the inhibitory effect of the indicated AI-
Poxblocks on MPXV infection of Vero E6 cells, with quantitative analysis of virus-
infected foci provided on the right. (E-F) BALB/c mice were infected intranasally
with 4×10^5 FFU MPXV and treated with single dose of AI-PoxBlock723 (10 mg/kg)
or PBS immediately after challenge ($n = 5$). Infectious MPXV particles and MPXV
viral loads in mice lungs at 4 dpi were determined by focus-forming assay (E) and
qPCR (F).

Discussion:

Our study provides new mechanistic insights into MPXV infection by identifying
the α-helical domain of the H3 protein as a critical mediator of heparan sulfate (HS)
binding. This domain, previously uncharacterized, plays a pivotal role in viral
adhesion to host cells, a key step in MPXV pathogenesis. Using a combination of AI-
based tools and molecular dynamics (MD) simulations, we successfully identified the
specific interaction sites between H3 and HS, offering a novel target for antiviral drug
design[54].

Previous studies have highlighted the importance of viral adhesion proteins in
orthopoxvirus infection, but the specific mechanisms underlying the H3-HS
interaction remained unresolved. Our findings provide the first detailed structural
insight into this interaction, confirming the importance of the electrostatic nature of
H3-HS binding. These results align with prior reports indicating the role of HS in
viral entry, but advance the field by identifying specific residues within the α-helical
domain that are crucial for binding.

One of the strengths of this study lies in the application of advanced AI-based
tools, such as AlphaFold2 and RFdiffusion, alongside MD simulations. These cutting-
edge computational approaches allowed us to overcome the limitations imposed by
the dynamic and flexible nature of HS. In addition, HS serves as a cellular ligand for
various viruses[55-57], underscoring the importance of studying its interactions with viral
adhesion proteins. By predicting the structure-function relationship of the H3-HS
interaction, we were able to pinpoint key binding residues and mechanisms that were
previously inaccessible. This demonstrates the growing potential of AI in structural
biology, particularly for the complex protein-glycan interactions[56, 58, 59].

The identification of the α-helical domain in H3 as a key player in HS binding
offers a promising new target for antiviral drug development[17, 60-67]. Given the
conservation of this domain across *orthopoxviruses*, including variola and vaccinia
viruses, targeting this region could lead to the development of broad-spectrum
antivirals effective against multiple *orthopoxviruses*[68]. Our study highlights the power
of AI-driven drug design, as demonstrated by the promising results of AI-

PoxBlock723, which showed efficacy in inhibiting viral infection. Further optimization and testing of these inhibitors in clinical models will be critical steps toward developing effective antiviral therapies. For example, conjugate the peptides with carrier molecules, such as liposomes, nanoparticles, or dendrimers, which can protect the peptides from immune detection and improve their delivery to target cells.

Looking ahead, the methodologies and findings from this study are likely to catalyze further investigations into viral adhesion mechanisms, not only in *orthopoxviruses* but also across other viral families that utilize similar adhesion strategies. Although challenges remain, we hope our discovery will attract structural biologists to resolve this domain's structure, and even the structure of H3-HS complex, to further enhance drug design efficiency for this target. As AI-driven approaches continue to evolve, they are poised to play a transformative role in advancing our understanding of viral pathogenesis and revolutionizing the future of drug discovery.

Acknowledgments: We appreciate the inspiring comments from previous anonymous reviewers. We acknowledge the funding support from the National Natural Science Foundation of China (22222703, 22477058), the Fundamental Research Funds for the Central Universities (020514380335), the Natural Science Foundation of Jiangsu Province (BK20202004), Shenzhen Medical Research Funds (B2302052). The numerical calculations in this work have been done on the computing facilities in the High-Performance Computing Center (HPCC) of Nanjing University.

Author contributions:
Conceptualization: PZ
Investigation: BZ, MD, YX, JQ, SW, RG, GT, LC
Funding acquisition: PZ, LC
Project administration & Supervision: PZ
Writing – original draft: PZ, BZ, LC
Competing interests: The authors declare no competing interests.
Data and materials availability: All data are available in the main text or the supplementary materials
Supplementary Materials: includes supplementary materials and methods, notes, figures and videos.

References:
1. F.-M. Lum, A. Torres-Ruesta, M. Z. Tay, R. T. P. Lin, D. C. Lye, L. Renia and L. F. P. Ng, Monkeypox: disease epidemiology, host immunity and clinical interventions, *Nat. Rev. Immunol*, 2022, **22**, 597-613.
2. D. The Lancet Infectious, Monkeypox: a neglected old foe, *Lancet Infect. Dis.*, 2022, **22**, 913.
3. A. Otoole, R. A. Neher, N. Ndodo, V. Borges, B. Gannon, J. P. Gomes, N. Groves, D. J. King, D. Maloney, P. Lemey, K. Lewandowski, N. Loman, R. Myers, I. F. Omah, M. A. Suchard, M. Worobey, M. Chand, C. Ihekweazu, D. Ulaeto, I. Adetifa and A. Rambaut, APOBEC3 deaminase editing in mpox

virus as evidence for sustained human transmission since at least 2016, *Science*, 2023, **382**, 595-600.

4. T. Kirby, What happened to the mpox pandemic?, *The Lancet*, 2023, **402**, 949-950.

5. E. H. Vakaniaki, C. Kacita, E. Kinganda-Lusamaki, A. OToole, T. Wawina-Bokalanga, D. Mukadi-Bamuleka, A. Amuri-Aziza, N. Malyamungu-Bubala, F. Mweshi-Kumbana, L. Mutimbwa-Mambo, F. Belesi-Siangoli, Y. Mujula, E. Parker, P.-C. Muswamba-Kayembe, S. S. Nundu, R. S. Lushima, J.-C. Makangara-Cigolo, N. Mulopo-Mukanya, E. Pukuta-Simbu, P. Akil-Bandali, H. Kavunga, O. Abdramane, I. Brosius, E. Bangwen, K. Vercauteren, N. A. Sam-Agudu, E. J. Mills, O. Tshiani-Mbaya, N. A. Hoff, A. W. Rimoin, L. E. Hensley, J. Kindrachuk, C. Baxter, T. de Oliveira, A. Ayouba, M. Peeters, E. Delaporte, S. Ahuka-Mundeke, E. L. Mohr, N. J. Sullivan, J.-J. Muyembe-Tamfum, J. B. Nachega, A. Rambaut, L. Liesenborghs and P. Mbala-Kingebeni, Sustained human outbreak of a new MPXV clade I lineage in eastern Democratic Republic of the Congo, *Nature Medicine*, 2024, DOI: 10.1038/s41591-024-03130-3.

6. P. Li, S. T. Pachis, G. Xu, R. Schraauwen, R. Incitti, A. C. de Vries, M. J. Bruno, M. P. Peppelenbosch, I. Alam, K. Raymond and Q. Pan, Mpox virus infection and drug treatment modelled in human skin organoids, *Nat. Microbiol.*, 2023, **8**, 2067-2079.

7. J. Liu, E. M. Mucker, J. L. Chapman, A. M. Babka, J. M. Gordon, A. V. Bryan, J. L. W. Raymond, T. M. Bell, P. R. Facemire, A. J. Goff, A. Nalca and X. Zeng, Retrospective detection of monkeypox virus in the testes of nonhuman primate survivors, *Nat. Microbiol.*, 2022, **7**, 1980-1986.

8. M. Zahmatyar, A. Fazlollahi, A. Motamedi, M. Zolfi, F. Seyedi, S. A. Nejadghaderi, M. J. M. Sullman, R. Mohammadinasab, A.-A. Kolahi, S. Arshi and S. Safiri, Human monkeypox: history, presentations, transmission, epidemiology, diagnosis, treatment, and prevention, *Front. Med.*, 2023, **10**.

9. L. M. Zaeck, M. M. Lamers, B. E. Verstrepen, T. M. Bestebroer, M. E. van Royen, H. Götz, M. C. Shamier, L. P. M. van Leeuwen, K. S. Schmitz, K. Alblas, S. van Efferen, S. Bogers, S. Scherbeijn, G. F. Rimmelzwaan, E. C. M. van Gorp, M. P. G. Koopmans, B. L. Haagmans, C. H. GeurtsvanKessel and R. D. de Vries, Low levels of monkeypox virus-neutralizing antibodies after MVA-BN vaccination in healthy individuals, *Nat. Med.*, 2023, **29**, 270-278.

10. J. Yu, X. Zhang, J. Liu, L. Xiang, S. Huang, X. Xie, L. Fang, Y. Lin, M. Zhang, L. Wang, J. He, B. Zhang, B. Di, B. Peng, J. Liang, C. Shen, W. Zhao and B. Li, Phylogeny and molecular evolution of the first local monkeypox virus cluster in Guangdong Province, China, *Nat. Commun.*, 2023, **14**, 8241.

11. Q. Peng, Y. Xie, L. Kuai, H. Wang, J. Qi, G. F. Gao and Y. Shi, Structure of monkeypox virus DNA polymerase holoenzyme, *Science*, 2023, **379**, 100-105.

12. B. Moss, Investigating Viruses During the Transformation of Molecular Biology: Part II, *Annu. Rev. Virol.*, 2020, **7**, 15-36.

13. R. D. M. Gray, D. Albrecht, C. Beerli, M. Huttunen, G. H. Cohen, I. J. White,

J. J. Burden, R. Henriques and J. Mercer, Nanoscale polarization of the entry fusion complex of vaccinia virus drives efficient fusion, *Nat. Microbiol.*, 2019, **4**, 1636-1644.

14. F. G. d. Fonseca, E. J. Wolffe, A. Weisberg and B. Moss, Effects of Deletion or Stringent Repression of the H3L Envelope Gene on Vaccinia Virus Replication, *J. Virol.*, 2000, **74**, 7518-7528.

15. L. Crickard, T. Babas, S. Seth, P. Silvera, L. Koriazova and S. Crotty, Protection of Rabbits and Immunodeficient Mice against Lethal Poxvirus Infections by Human Monoclonal Antibodies, *Plos One*, 2012, **7**, e48706.

16. A. Zuiani, C. L. Dulberger, N. S. De Silva, M. Marquette, Y.-J. Lu, G. M. Palowitch, A. Dokic, R. Sanchez-Velazquez, K. Schlatterer, S. Sarkar, S. Kar, B. Chawla, A. Galeev, C. Lindemann, D. A. Rothenberg, H. Diao, A. C. Walls, T. A. Addona, F. Mensa, A. B. Vogel, L. M. Stuart, R. van der Most, J. R. Srouji, O. Tureci, R. B. Gaynor, U. Sahin and A. Poran, A multivalent mRNA monkeypox virus vaccine (BNT166) protects mice and macaques from orthopoxvirus disease, *Cell*, 2024, **187**, 1363-1373.e1312.

17. P. Gulati, J. Chadha, K. Harjai and S. Singh, Targeting envelope proteins of poxviruses to repurpose phytochemicals against monkeypox: An in silico investigation, *Front. Microbiol.*, 2023, **13**.

18. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli and D. Hassabis, Highly accurate protein structure prediction with AlphaFold, *Nature*, 2021, **596**, 583-589.

19. J. C. Phillips, R. Braun, W. Wang, J. Gumbart, E. Tajkhorshid, E. Villa, C. Chipot, R. D. Skeel, L. Kale and K. Schulten, Scalable molecular dynamics with NAMD, *J. Comput. Chem.*, 2005, **26**, 1781-1802.

20. F. Gräter, J. Shen, H. Jiang, M. Gautel and H. Grubmuller, Mechanically Induced Titin Kinase Activation Studied by Force-Probe Molecular Dynamics Simulations, *Biophy. J.*, 2005, **88**, 790-804.

21. E. Kim, S. Lee, A. Jeon, J. M. Choi, H.-S. Lee, S. Hohng and H.-S. Kim, A single-molecule dissection of ligand binding to a protein with intrinsic dynamics, *Nat. Chem. Biol.*, 2013, **9**, 313-318.

22. C. L. Brooks, A. D. MacKerell, C. B. Post and L. Nilsson, Biomolecular dynamics in the 21st century, *Biochim. Biophys. Acta. Gen. Subj.*, 2024, **1868**, 130534.

23. M. Rief, M. Gautel, F. Oesterhelt, J. M. Fernandez and H. E. Gaub, Reversible unfolding of individual titin immunoglobulin domains by AFM, *Science*, 1997, **276**, 1109-1112.

24. C. Sieben, C. Kappel, R. Zhu, A. Wozniak, C. Rankl, P. Hinterdorfer, H. Grubmuller and A. Herrmann, Influenza virus binds its host cell using multiple

dynamic interactions, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 13626-13631.

25. A. Raab, W. Han, D. Badt, S. J. Smith-Gill, S. M. Lindsay, H. Schindler and P. Hinterdorfer, Antibody recognition imaging by force microscopy, *Nat. Biotech.*, 1999, **17**, 901.

26. D. J. Muller, J. Helenius, D. Alsteens and Y. F. Dufrene, Force probing surfaces of living cells to molecular resolution, *Nat. Chem. Biol.*, 2009, **5**, 383-390.

27. J. Dauparas, I. Anishchenko, N. Bennett, H. Bai, R. J. Ragotte, L. F. Milles, B. I. M. Wicky, A. Courbet, R. J. de Haas, N. Bethel, P. J. Y. Leung, T. F. Huddy, S. Pellock, D. Tischer, F. Chan, B. Koepnick, H. Nguyen, A. Kang, B. Sankaran, A. K. Bera, N. P. King and D. Baker, Robust deep learning-based protein sequence design using ProteinMPNN, *Science*, 2022, **378**, 49-56.

28. J. L. Watson, D. Juergens, N. R. Bennett, B. L. Trippe, J. Yim, H. E. Eisenach, W. Ahern, A. J. Borst, R. J. Ragotte, L. F. Milles, B. I. M. Wicky, N. Hanikel, S. J. Pellock, A. Courbet, W. Sheffler, J. Wang, P. Venkatesh, I. Sappington, S. V. Torres, A. Lauko, V. De Bortoli, E. Mathieu, S. Ovchinnikov, R. Barzilay, T. S. Jaakkola, F. DiMaio, M. Baek and D. Baker, De novo design of protein structure and function with RFdiffusion, *Nature*, 2023, **620**, 1089-1100.

29. K. Singh, A. G. Gittis, R. K. Gitti, S. A. Ostazeski, H.-P. Su and D. N. Garboczi, The Vaccinia Virus H3 Envelope Protein, a Major Target of Neutralizing Antibodies, Exhibits a Glycosyltransferase Fold and Binds UDP-Glucose, *J. Virol.*, 2016, **90**, 5020-5030.

30. F. G. d. Fonseca, E. J. Wolffe, A. Weisberg and B. Moss, Characterization of the Vaccinia Virus H3L Envelope Protein: Topology and Posttranslational Membrane Insertion via the C-Terminal Hydrophobic Tail, *J. Virol.*, 2000, **74**, 7508-7517.

31. C.-L. Lin, C.-S. Chung, H. G. Heine and W. Chang, Vaccinia Virus Envelope H3L Protein Binds to Cell Surface Heparan Sulfate and Is Important for Intracellular Mature Virion Morphogenesis and Virus Infection In Vitro and In Vivo, *J. Virol.*, 2000, **74**, 3353-3365.

32. A. Morgan and X. Wang, The Novel Heparin-Binding Motif in Decorin-Binding Protein A from Strain B31 of Borrelia burgdorferi Explains the Higher Binding Affinity, *Biochemistry-Us*, 2013, **52**, 8237-8245.

33. Z. Lin, H. Akin, R. Rao, B. Hie, Z. Zhu, W. Lu, N. Smetanin, R. Verkuil, O. Kabeli, Y. Shmueli, A. dos Santos Costa, M. Fazel-Zarandi, T. Sercu, S. Candido and A. Rives, Evolutionary-scale prediction of atomic-level protein structure with a language model, *Science*, 2023, **379**, 1123-1130.

34. M. Baek, R. McHugh, I. Anishchenko, H. Jiang, D. Baker and F. DiMaio, Accurate prediction of protein-nucleic acid complexes using RoseTTAFoldNA, *Nature Methods*, 2024, **21**, 117-121.

35. J. Eberhardt, D. Santos-Martins, A. F. Tillack and S. Forli, AutoDock Vina 1.2.0: New Docking Methods, Expanded Force Field, and Python Bindings, *J. Chem. Inf. Model.*, 2021, **61**, 3891-3898.

36. J. Lee, X. Cheng, J. M. Swails, M. S. Yeom, P. K. Eastman, J. A. Lemkul, S.

Wei, J. Buckner, J. C. Jeong, Y. Qi, S. Jo, V. S. Pande, D. A. Case, C. L. Brooks, A. D. MacKerell, J. B. Klauda and W. Im, CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field, *J. Chem. Theory Comput.*, 2015, **12**, 405-413.

37. Y. Sugita and Y. Okamoto, Replica-exchange molecular dynamics method for protein folding, *Chem. Phys. Lett.*, 1999, **314**, 141-151.

38. V. Govind Kumar, A. Polasa, S. Agrawal, T. K. S. Kumar and M. Moradi, Binding affinity estimation from restrained umbrella sampling simulations, *Nat. Comput. Sci.*, 2022, **3**, 59-70.

39. S. F. Altschul, T. L. Madden, A. A. Schaffer, J. H. Zhang, Z. Zhang, W. Miller and D. J. Lipman, Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucleic Acids Res.*, 1997, **25**, 3389-3402.

40. H. Yu, M. G. W. Siewny, D. T. Edwards, A. W. Sanders and T. T. Perkins, Hidden dynamics in the unfolding of individual bacteriorhodopsin proteins, *Science*, 2017, **355**, 945-950.

41. H. Dietz and M. Rief, Exploring the energy landscape of GFP by single-molecule mechanical experiments, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 16192-16197.

42. M. Goktas, C. F. Luo, R. M. A. Sullan, A. E. Bergues-Pupo, R. Lipowsky, A. V. Verde and K. G. Blank, Molecular mechanics of coiled coils loaded in the shear geometry, *Chem. Sci.*, 2018, **9**, 4610-4621.

43. Y. Cao, C. Lam, M. Wang and H. Li, Nonmechanical protein can have significant mechanical stability, *Angew. Chem. Int. Ed.*, 2006, **45**, 642-645.

44. S. W. Stahl, M. A. Nash, D. B. Fried, M. Slutzki, Y. Barak, E. A. Bayer and H. E. Gaub, Single-molecule dissection of the high-affinity cohesin-dockerin complex, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 20431-20436.

45. B. Zheng, Y. Xiao, B. Tong, Y. Mao, R. Ge, F. Tian, X. Dong and P. Zheng, S373P Mutation Stabilizes the Receptor-Binding Domain of the Spike Protein in Omicron and Promotes Binding, *JACS Au*, 2023, **3**, 1902-1910.

46. S. Shi, Z. Wang, Y. Deng, F. Tian, Q. Wu and P. Zheng, Combination of Click Chemistry and Enzymatic Ligation for Stable and Efficient Protein Immobilization for Single-Molecule Force Spectroscopy, *CCS Chem.*, 2022, **4**, 598-604.

47. Y. Deng, T. Wu, M. Wang, S. Shi, G. Yuan, X. Li, H. Chong, B. Wu and P. Zheng, Enzymatic biosynthesis and immobilization of polyprotein verified at the single-molecule level, *Nat. Commun.*, 2019, **10**, 2775-2785.

48. Y. Liu, D. Song, S. Li, Z. Guo and P. Zheng, Click Chemistry-Based Force Spectroscopy Revealed Enhanced Binding Dynamics of Phosphorylated HMGB1 to Cisplatin-DNA, *J. Am. Chem. Soc.*, 2024, **146**, 13126-13132.

49. M. Carrion-Vazquez, A. F. Oberhauser, S. B. Fowler, P. E. Marszalek, S. E. Broedel, J. Clarke and J. M. Fernandez, Mechanical and chemical unfolding of a single protein: A comparison, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 3694-3699.

| 669 | 50. | F. Tian, B. Tong, L. Sun, S. Shi, B. Zheng, Z. Wang, X. Dong and P. Zheng, N501Y mutation of spike protein in SARS-CoV-2 strengthens its binding to receptor ACE2, *Elife*, 2021, **10**, e69091. |
| 672 | 51. | D. Alsteens, R. Newton, R. Schubert, D. Martinez-Martin, M. Delguste, B. Roska and D. J. Muller, Nanomechanical mapping of first binding steps of a virus to animal cells, *Nat. Nanotechnol.*, 2016, **12**, 177. |

50.   F. Tian, B. Tong, L. Sun, S. Shi, B. Zheng, Z. Wang, X. Dong and P. Zheng,
N501Y mutation of spike protein in SARS-CoV-2 strengthens its binding to
receptor ACE2, *Elife*, 2021, **10**, e69091.

51.   D. Alsteens, R. Newton, R. Schubert, D. Martinez-Martin, M. Delguste, B.
Roska and D. J. Muller, Nanomechanical mapping of first binding steps of a
virus to animal cells, *Nat. Nanotechnol.*, 2016, **12**, 177.

52.   L. Cheng, L. Yang, M. Wang, Y. Peng, H. Wang, X. Yang, J. Zhao, M. Zhang,
F. Wang and Z. Zhang, Isolation and characterization of mpox virus from the
first mpox case in Shenzhen, China, *Virologica Sinica*, 2024, **39**, 335-337.

53.   L. Cheng, W. Huang, M. Duan, Z. Li, Q. Chen, M. Zhang and Z. Zhang,
Pathogenic BALB/c mice infection model for evaluation of mpox
countermeasures, *Cell Discovery*, 2024, **10**, 105.

54.   S. Monzon, S. Varona, A. Negredo, S. Vidal-Freire, J. A. Patino-Galindo, N.
Ferressini-Gerpe, A. Zaballos, E. Orviz, O. Ayerdi, A. Munoz-Gomez, A.
Delgado-Iribarren, V. Estrada, C. Garcia, F. Molero, P. Sanchez-Mora, M.
Torres, A. Vazquez, J.-C. Galan, I. Torres, M. Causse del Rio, L. Merino-Diaz,
M. Lopez, A. Galar, L. Cardenoso, A. Gutierrez, C. Loras, I. Escribano, M. E.
Alvarez-Arguelles, L. del Rio, M. Simon, M. A. Melendez, J. Camacho, L.
Herrero, P. Jimenez, M. L. Navarro-Rico, I. Jado, E. Giannetti, J. H. Kuhn, M.
Sanchez-Lockhart, N. Di Paola, J. R. Kugelman, S. Guerra, A. Garcia-Sastre,
I. Cuesta, M. P. Sanchez-Seco and G. Palacios, Monkeypox virus genomic
accordion strategies, *Nat. Commun.*, 2024, **15**, 3059.

55.   T. M. Clausen, D. R. Sandoval, C. B. Spliid, J. Pihl, H. R. Perrett, C. D.
Painter, A. Narayanan, S. A. Majowicz, E. M. Kwong, R. N. McVicar, B. E.
Thacker, C. A. Glass, Z. Yang, J. L. Torres, G. J. Golden, P. L. Bartels, R. N.
Porell, A. F. Garretson, L. Laubach, J. Feldman, X. Yin, Y. Pu, B. M. Hauser,
T. M. Caradonna, B. P. Kellman, C. Martino, P. L. S. M. Gordts, S. K. Chanda,
A. G. Schmidt, K. Godula, S. L. Leibel, J. Jose, K. D. Corbett, A. B. Ward, A.
F. Carlin and J. D. Esko, SARS-CoV-2 Infection Depends on Cellular Heparan
Sulfate and ACE2, *Cell*, 2020, **183**, 1043-1057.e1015.

56.   Z. Shriver, I. Capila, G. Venkataraman and R. Sasisekharan, in *Heparin - A
Century of Progress*, eds. R. Lever, B. Mulloy and C. P. Page, Springer Berlin
Heidelberg, Berlin, Heidelberg, 2012, DOI: 10.1007/978-3-642-23056-1_8,
pp. 159-176.

57.   M. Koehler, M. Delguste, C. Sieben, L. Gillet and D. Alsteens, Initial Step of
Virus Entry: Virion Binding to Cell-Surface Glycans, *Annu. Rev. Virol.*, 2020,
7, 143-165.

58.   P.-C. Shih, M.-S. Yang, S.-C. Lin, Y. Ho, J.-C. Hsiao, D.-R. Wang, S. S. F. Yu,
W. Chang and D.-L. M. Tzou, A Turn-like Structure "KKPE" Segment
Mediates the Specific Binding of Viral Protein A27 to Heparin and Heparan
Sulfate on Cell Surfaces*, *J. Bio. Chem.*, 2009, **284**, 36535-36546.

59.   P. E. Marszalek, A. F. Oberhauser, H. Li and J. M. Fernandez, The Force-
Driven Conformations of Heparin Studied with Single Molecule Force

Microscopy, *Biophy. J.*, 2003, **85**, 2696-2704.

60. Z. Q. Zhang, J. Xie, H. Y. Liu, J. Liu and R. J. Linhardt, Quantification of Heparan Sulfate Disaccharides Using Ion-Pairing Reversed-Phase Microflow High-Performance Liquid Chromatography with Electrospray Ionization Trap Mass Spectrometry, *Anal. Chem.*, 2009, **81**, 4349-4355.

61. S. Podduturi, D. Vemula, S. Singothu and V. Bhandari, In-silico investigation of E8 surface protein of the monkeypox virus to identify potential therapeutic agents, *J. Biomol. Struct. Dyn.*, 2023, DOI: 10.1080/07391102.2023.2245041, 1-14.

62. R. C. Bernardi, E. Durner, C. Schoeler, K. H. Malinowska, B. G. Carvalho, E. A. Bayer, Z. Luthey-Schulten, H. E. Gaub and M. A. Nash, Mechanisms of Nanonewton Mechanostability in a Protein Complex Revealed by Molecular Dynamics Simulations and Single-Molecule Force Spectroscopy, *J. Am. Chem. Soc.*, 2019, **141**, 14752-14763.

63. M. Zhao and M. T. Woodside, Mechanical strength of RNA knot in Zika virus protects against cellular defenses, *Nat. Chem. Biol.*, 2021, **17**, 975-981.

64. T. Serdiuk, D. Balasubramaniam, J. Sugihara, S. A. Mari, H. R. Kaback and D. J. Müller, YidC assists the stepwise and stochastic folding of membrane proteins, *Nat. Chem. Biol.*, 2016, **12**, 911-917.

65. P. S. Sen Gupta, S. K. Panda, A. K. Nayak and M. K. Rana, Identification and Investigation of a Cryptic Binding Pocket of the P37 Envelope Protein of Monkeypox Virus by Molecular Dynamics Simulations, *J. Phys. Chem. Lett.*, 2023, **14**, 3230-3235.

66. R. Petrosyan, S. Patra, N. Rezajooei, C. R. Garen and M. T. Woodside, Unfolded and intermediate states of PrP play a key role in the mechanism of action of an antiprion chaperone, *Proc. Natl. Acad. Sci. U. S. A.*, 2021, **118**.

67. G. Chen, H. Wang, L. Bumba, J. Masin, P. Sebo and H. Li, The adenylate cyclase toxin RTX domain follows a series templated folding mechanism with implications for toxin activity, *J. Biol. Chem.*, 2023, **299**.

68. J. Wang, M. Shahed-Ai-Mahmud, A. Chen, K. Li, H. Tan and R. Joyce, An Overview of Antivirals against Monkeypox Virus and Other Orthopoxviruses, *J. Med. Chem.*, 2023, **66**, 4468-4490.