1　Title: **TCTEX1D2 has two distinct functions in sperm flagellum formation in mice;**
2　**cytoplasmic dynein 2 and inner dynein arm.**
3

4　Ryua Harima[1], Kenshiro Hara[1], Kentaro Tanemura[1,2,*]
5
6　[1]Laboratory of Animal Reproduction and Development, Graduate School of Agricultural
7　Science, Tohoku University, 468-1 Aramaki-Aoba, Aoba-ku, Sendai, Miyagi 980-8572,
8　Japan; [2]Laboratory of Reproductive Technology (Repro-SOLEIL), Graduate School of
9　Agricultural Science, Tohoku University, 468-1 Aramaki-Aoba, Aoba-ku, Sendai, Miyagi
10　980-8572, Japan
11
12　*** For correspondence:** kentaro.tanemura.e4@tohoku.ac.jp

13　**Competing interest:** The authors declare that no competing interests exist.
14
15

Abstract
Flagella and cilia are widely conserved motile structures. Mammalian sperm
possess flagella, which confer motility to the sperm and, therefore, are essential
for fertilization. Large protein complexes called dynein, including cytoplasmic
dynein 2 and axonemal dynein, play a role in the formation of cilia and flagella.
The function of each dynein complex subunit in sperm flagellum formation is unclear.
One such subunit is TCTEX1D2. *Tctex1d2* $^{-/-}$ mice generated in this study showed male
infertility because of flagellar dysplasia arising from round spermatids and
disruption of the axonemal structure inside the flagella. In contrast, the motile
cilia of *Tctex1d2* $^{-/-}$ mice were normal. Co-immunoprecipitation studies showed that
TCTEX1D2 interacted with the cytoplasmic dynein 2 component proteins WDR34, WDR60,
and DYNLT1 in the testes and that the localization of these proteins was abnormal
in *Tctex1d2* $^{-/-}$ mice. Furthermore, TCTEX1D2 also interacted with WDR63 and WDR78,
component proteins of the inner dynein arm, which is axonemal dynein. Overall, we
revealed that TCTEX1D2 has two distinct functions in mouse sperm flagellum formation
the assembly of cytoplasmic dynein 2 and organization of inner dynein arm which is
not the case in cilia formation where TCTEX1D2 functions only as cytoplasmic dynein
2.

Introduction

Flagella are widely conserved structures present in a large number of organisms ranging from protists to mammals, and their most distinctive feature is their motility [1,2]. In mammals, the sperm is the only cell with flagella and thereby has motility, which is essential for fertilization [3]. In addition to flagella, mammalian cells also have a similar motile structure called cilia [4]. The flagellar and cilia axonemes have 9+2 microtubule internal structure, consisting of two central microtubules surrounded by nine peripheral microtubules [5-7]. Large protein complexes called dynein, which move along the microtubules, play important roles in the formation of cilia and flagella and maintenance of motility [8-10]. Dynein is composed of different subunits that are classified as heavy, intermediate, light intermediate, and light chain subunits based on their molecular weight [11]. Dynein is divided into cytoplasmic and axonemal dynein, and cytoplasmic dynein is further divided into cytoplasmic dynein 1 and cytoplasmic dynein 2 [12]. Cytoplasmic dynein 2 and axonemal dynein are closely related to cilia and flagella [13-16]. Cytoplasmic dynein 2 primarily plays a role in cilia formation [17,18] and is responsible for retrograde transport as part of the intraflagellar transport (IFT) system within cilia [18-20]. Axonemal dynein is present as part of the axonemal structure, and divided into outer dynein arm and inner dynein arm. Axonemal dynein generates a sliding force that allows the flagella and cilia to acquire motility [15]. The functions of cytoplasmic dynein 2 and axonemal dynein are very different; therefore, most subunits are specific to either cytoplasmic or axonemal dynein [21,22]. However, light chains are an exception to this rule. In *Chlamydomonas* flagella, some light chains are common between cytoplasmic dynein 2 and axonemal dynein [23,24]. This suggests that dynein light chains are involved in many processes about related to flagella, ranging from formation to structural maintenance. In *Chlamydomonas*, dynein light chains contribute to the assembly of the components, structural stability, and motility of the flagella [24].

In mammals, most studies on dynein light chains have focused on somatic cilia, and very few have focused on sperm flagella. Thus, our understanding of the dynein light chains involved in sperm flagellum formation is very poor. Although flagella are widely conserved structures across species, it is not appropriate to apply the findings from other species to mammalian sperm flagella, because there are differences in the formation mechanisms and constituent proteins between species [2,25-27]. The formation mechanism of sperm flagella and cilia is considered to be similar; however, sperm flagella are generally much longer than cilia and have unique structures, such as mitochondrial sheath, outer dense fiber, and fibrous sheath. Furthermore, it has

been reported that the functions of proteins related to axoneme formation differ between sperm flagella and cilia [16], suggesting that there are differences in the mechanism of their formation. Therefore, it is important to use mammalian sperm flagella, to understand the function of dynein light chain in mammalian sperm flagella.

In the present study, we focused on Tctex1 domain-containing 2 (TCTEX1D2, also known as DYNLT2B) a dynein light chain present in the testes. In mammalian somatic cells, TCTEX1D2 is known as one of the cytoplasmic dynein 2 light chains and is involved in the formation of primary cilia (non-motile cilia) [11,28,29]. Cytoplasmic dynein 2 has seven light chains (DYNLL1, DYNLL2, DYNLT1, DYNLT3, DYNLRB1, DYNLRB2, and TCTEX1D2) [11,21]; however, the precise function of each subunit is unclear. Among the light chains, only TCTEX1D2 is specific to cytoplasmic dynein 2 [11,28,29], whereas the others are also found in cytoplasmic dynein 1 [11]. In addition, DYNLL1 and DYNLL2, DYNLT1 and DYNLT3, and DYNLRB1 and DYNLRB2 are homologous to each other, and the homologs appear to have interchangeable functions [30]. Therefore, TCTEX1D2 is unique and irreplaceable among the light chains of cytoplasmic dynein 2 and is thought to play a critical role in cytoplasmic dynein 2. However, there are no reports on the role of TCTEX1D2 in sperm flagellum formation, and it is yet unknown whether TCTEX1D2 is a component of cytoplasmic dynein 2 in germ cells. Interestingly, *Tctex1d2* mRNA is highly expressed in mouse testes according to NCBI gene (https://www.ncbi.nlm.nih.gov/gene/66061) and is mainly expressed in spermatocytes according to single-cell RNA-seq in mouse testes [31]. Therefore, it is hypothesized that *Tctex1d2* might play a role in sperm flagellum formation. In this study, we aimed to elucidate the function of *Tctex1d2* in sperm flagellum formation.

In this study, we created *Tctex1d2*$^{-/-}$ mice to investigate the function of *Tctex1d2*. Male *Tctex1d2*$^{-/-}$ mice were infertile and displayed severe morphological abnormalities in sperm flagella and disorganized distribution of axonemal components. However, there was no effect on the formation of motile cilia; only the formation of sperm flagella was significantly impaired in *Tctex1d2*$^{-/-}$ mice. TCTEX1D2 interacted with cytoplasmic dynein 2 component proteins in the testes, and we observed that the assembly of the cytoplasmic dynein 2 subunit is blocked in *Tctex1d2*$^{-/-}$ mice. TCTEX1D2 also interacted with the subunits of axonemal dynein in the testes. This study revealed that TCTEX1D2 has two distinct functions cytoplasmic dynein 2 and axonemal dynein and plays an important role in sperm flagellum formation in mice.

Results
TCTEX1D2 is highly expressed in the testes and localized in flagella from round
spermatids to elongated spermatids
Tctex1d2 mRNA is highly expressed in mouse testes according to NCBI gene
(https://www.ncbi.nlm.nih.gov/gene/66061). Western blotting (WB) was performed to
investigate TCTEX1D2 protein expression. TCTEX1D2 was found to be highly expressed
in the testes and spleen (fig. 1A). This shows that TCTEX1D2 may have certain functions
in the testes. To determine the localization of TCTEX1D2 in germ cells, we performed
immunohistochemistry (IHC) and immunocytochemistry (ICC) experiments. However, no
antibodies were available for the IHC or ICC analyses. Therefore, we generated
Tctex1d2 – 3×FLAG mice, in which a 3×FLAG was knocked in at the N-terminus of *Tctex1d2*
(fig. S1A) and implemented ICC in germ cells isolated from the testes of these mice.
Spermatogenesis in *Tctex1d2* – 3×FLAG mice was comparable to that in WT mice, and
the sperm flagella of *Tctex1d2* – 3×FLAG mice showed normal morphology (fig. S1B,
C). Thus, 3×FLAG knock-in in *Tctex1d2* had no effect on the function of TCTEX1D2.
We found that TCTEX1D2 was localized along the flagella from round spermatids to
elongated spermatids. (fig. 1B–D). TCTEX1D2 was also localized in the manchette, which
are transiently formed in step 10–14 spermatids (fig. 1C). Furthermore, TCTEX1D2 was
also expressed in the cilia of ventricles, trachea, and oviducts (fig. S2A–C).
125

TCTEX1D2 is a component of cytoplasmic dynein 2
In mammalian somatic cells, TCTEX1D2 is a component of cytoplasmic dynein 2 (fig.
2A). Therefore, we analyzed whether TCTEX1D2 is a component of cytoplasmic dynein
2 in germ cells. Cytoplasmic dynein 2 consists of 11 subunits: heavy chain (DYNC2H1),
intermediate chain (WDR34 and WDR60), light intermediate chain (DYNC2LI1), and light
chain (DYNLRB1, DYNLRB2, DYNLL1, DYNLL2, DYNLT1, DYNLT3, and TCTEX1D2) (fig. 2A).
In mammalian somatic cells, TCTEX1D2 interacts with DYNLT1 and WDR60 [30]. In the testes,
TCTEX1D2 interacted not only with WDR60 and DYNLT1, but also with WDR34 (fig. 2B).
These results indicate that TCTEX1D2 is one of the components of cytoplasmic dynein
2 in germ cells.
136

TCTEX1D2 is an axoneme component and interacts with subunits of the inner dynein arm
In the flagella of *Chlamydomonas*, Tctex2b (a homolog of mouse TCTEX1D2) is present
as a light chain of the inner dynein arm in the axoneme dynein [32]. The inner dynein
arm is important for controlling the size and shape of the bend of the flagella,
features collectively referred to as waveforms [33]. In *Chlamydomonas*, loss of function

of the inner arm dynein subunit causes impaired flagellar motility [34–36]. Thus, it is

believed that in mammals, TCTEX1D2 is present in the sperm flagella as a light chain

of the inner dynein arm. We analyzed the protein expression of TCTEX1D2 in the

spermatozoa of the cauda epididymis. WB analysis of sperm proteins fractionated into

Triton X-100 soluble, SDS soluble, and SDS-insoluble fractions was performed. SLC2A3,

Ac-α-tubulin, and AKAP3 are markers for sperm membrane, axoneme, and other insoluble

fractions, respectively. TCTEX1D2 was present in both the SDS-soluble and

SDS-resistant fractions (fig. 2C). ICC studies also revealed that TCTEX1D2 localized

along the sperm flagella (fig. 2D). These results showed that TCTEX1D2 is an axoneme

component in mice. Further, co-immunoprecipitation studies revealed that TCTEX1D2

interacted with WDR63 and WDR78 (fig. 2E). Both proteins are intermediate chains of

the inner dynein arm, which is a component of the 9+2 microtubule structure. These

results indicate that TCTEX1D2 has some functions in the light chain of the inner

dynein arm in mouse sperm flagella, as well as in that of *Chlamydomonas* flagella.

Tctex1d2 is essential for male fertility

To explore the function of TCTEX1D2 in the mouse testes, we generated *Tctex1d2*

knockout (*Tctex1d2* $^{-/-}$) mice using the CRISPR/Cas9 system (fig. 3A). We confirmed a

4696-bp deletion in the generated knockout mice using genotyping and Sanger sequencing

(fig. 3B, C). Furthermore, the generated knockout mice completely lost TCTEX1D2

protein expression, as shown by results of WB experiments (fig. 3D). Fertility tests

were conducted to assess the fertility of *Tctex1d2* $^{-/-}$ male mice. *Tctex1d2* $^{-/-}$ male mice

were mated with two wild-type (WT) female mice. *Tctex1d2* $^{-/-}$ male mice were found to

be completely infertile (fig. 3E). In contrast, the testis/body weight ratio was

unchanged in *Tctex1d2* $^{-/-}$ mice in comparison with *Tctex1d2* $^{+/+}$ mice (fig. 3F).

Tctex1d2 $^{-/-}$ mice exhibit abnormal morphology of sperm flagella, but not of cilia

To identify the cause of infertility in *Tctex1d2* $^{-/-}$ male mice, we performed

periodic acid-Schiff hematoxylin (PAS-H) staining of the testes and epididymis. In

stage I-III seminiferous tubules, a layered structure of germ cells was observed (fig.

4A, yellow box) and numerous spermatid flagella were present in the tubule lumen of

Tctex1d2 $^{+/+}$ mice (fig. 4A, red box arrow). In contrast, a layered structure of germ

cells was observed (fig. 4B, yellow box), but only a few flagella were present in

the tubule lumen of *Tctex1d2* $^{-/-}$ mice (fig. 4B, red box). In stage Ⅶ seminiferous

tubules, a large number of elongated spermatids were observed in the lumen in *Tctex1d2*

$^{+/+}$ mice (fig. 4C). However, only a few elongated spermatids were observed in the lumen

of *Tctex1d2* $^{-/-}$ mice (fig. 4D), and the number of spermatozoa was significantly

decreased in the cauda epididymis (fig. 4E, F). Furthermore, almost all the sperm

flagella of *Tctex1d2* $^{-/-}$ mice showed markedly abnormal morphology compared with those

of *Tctex1d2* $^{+/+}$ mice (fig. 4G), such as shortened and coiled tails (fig. 4H). In mice,

the epithelial cells of the ventricles, trachea, and oviducts have motile cilia

similar to sperm flagella. We analyzed the morphology of these cilia using Hematoxylin

–Eosin (HE) staining. In contrast to the severe defects in sperm flagella, the cilia

of the ventricles, trachea, and oviducts showed no abnormal morphology in *Tctex1d2*

$^{-/-}$ mice (fig. S3A–C). Therefore, we concluded that only the sperm flagella showed

severely abnormal morphology in *Tctex1d2* $^{-/-}$ mice.

Tctex1d2 $^{-/-}$ mice show abnormal sperm flagella and disruption of axoneme structures

in early round spermatids

The results shown in fig. 1B–D indicate that TCTEX1D2 is expressed from the early

stage of sperm flagellum formation and may have some functions at that stage. To

further understand the role of TCTEX1D2, we analyzed flagellum formation during

spermiogenesis. Flagellum formation in *Tctex1d2* $^{-/-}$ mice was normal until steps 2–3

of the spermatids (fig. 5A). However, the axoneme failed to elongate normally and

showed significant defects from step 4 spermatids onwards in *Tctex1d2* $^{-/-}$ mice (fig.

5A). Using transmission electron microscopy, we examined the internal structure of

the flagella and found that the axonemal components were disorganized in *Tctex1d2*

$^{-/-}$ mice. Doublet microtubules were not formed and many single microtubules were

observed in *Tctex1d2* $^{-/-}$ mice (fig. 5B). Moreover, the 9＋2 microtubule structure, inner

dynein arm (IDA), outer dynein arm (ODA), outer dense fiber (ODF), and fibrous sheath

(FS) were disorganized and scattered in *Tctex1d2* $^{-/-}$ mice (fig. 5B). Thus, the cause

of the failure of flagellar elongation in *Tctex1d2* $^{-/-}$ mice was disruption of the

axonemal structure. These results suggested that TCTEX1D2 plays an important role

in the organization of axonemal components.

TCTEX1D2 functions in the assembly of cytoplasmic dynein 2 subunits

TCTEX1D2 is one of the components of cytoplasmic dynein 2 in germ cells. Therefore,

we investigated the expression patterns of TCTEX1D2 interacted subunit (WDR60, WDR34,

and DYNLT1) in *Tctex1d2* $^{-/-}$ mice using ICC. WDR60 and WDR34 were not expressed in the

flagella both *Tctex1d2* $^{+/+}$ and *Tctex1d2* $^{-/-}$ mice round spermatids (fig. 6A, B). In the

flagella in *Tctex1d2* $^{+/+}$ mice elongated spermatids, WDR60 localized along the flagellar,

especially strongly in principal piece (fig. 6A). WDR34 also localized along the

flagellar, especially strongly in middle piece (fig. 6B). However, expression
patterns of WDR60 and WDR34 were abnormal in *Tctex1d2* $^{-/-}$ mice elongated spermatids,
including loss of signal (fig. 6A, B above) and ectopic signal (fig, 6A, B bellow
arrow head). DYNLT1 uniformly localized along the flagellar in *Tctex1d2* $^{+/+}$ mice round
spermatids to elongated spermatids (fig. 6C). However, in *Tctex1d2* $^{-/-}$ mice, expression
of DYNLT1 decreased in the flagella of round spermatids and loss in elongated
spermatids (fig. 6C). These results suggest that TCTEX1D2 plays a role in the assembly
and structural stability of the cytoplasmic dynein 2 subunits.

During flagellum formation, the components that constitute the flagella are
transported along microtubules. The IFT system is responsible for this transport and
consists mainly of IFT proteins, kinesin 2, cytoplasmic dynein 2, and the BBSome
complex [9,37-39]. As *Tctex1d2* $^{-/-}$ mice showed abnormal cytoplasmic dynein 2 localization,
we analyzed the expression patterns of other IFT systems. IFT88, an IFT protein, and
DYNC2LI1 showed decreased signals and ectopic expression in the flagella of elongated
spermatids in *Tctex1d2* $^{-/-}$ mice (fig, S4A, B). In contrast, IFT140 (an IFT protein),
KIF3A (a component of kinesin 2), and DYNC2H1 were localized normally in the flagella
of elongated spermatids in *Tctex1d2* $^{-/-}$ mice (fig. S4C~E). These results suggest that
the IFT system is partially defective in *Tctex1d2* $^{-/-}$ mice. These results are
represented in the schematic diagram (fig. 6D). In *Tctex1d2* $^{+/+}$ mice elongated
spermatids, the 11 subunits of cytoplasmic dynein 2 form complexes and move on
microtubules (fig. 6D). However, *Tctex1d2* $^{-/-}$ mice elongated spermatids, DYNC2LI1,
WDR60, WDR34 and DYNLT1 among the 11 subunits showed abnormal localization and were
not assembled in the cytoplasmic dynein 2 complex (fig. 6D). Therefore, TCTEX1D2 plays
a role in the assembly of cytoplasmic dynein 2 subunits.

Discussion

In this study, we identified TCTEX1D2 as a dynein light chain important for sperm

flagellum formation in mice. *Tctex1d2*[-/-] mice showed abnormal flagella in round

spermatids and disorganization of axonemal components in the flagella. TCTEX1D2 was

identified as a component of cytoplasmic dynein 2, based on its interaction with WDR34,

WDR60, and DYNLT1, and the expression patterns of these proteins was found to be

aberrant in the elongated spermatids of *Tctex1d2*[-/-] mice. In addition, IFT88 and

DYNC2LI1 showed similar abnormal expression patterns in the elongated spermatids of

Tctex1d2[-/-] mice. These results suggested that the assembly of cytoplasmic dynein 2

was impaired in *Tctex1d2*[-/-] mice, consequently affecting the stability of the IFT

system. Cilia formation is impaired when the functions of the cytoplasmic dynein 2

and IFT proteins are completely lost [40-43]. However, the formation of primary cilia

is normal in TCTEX1D2 knockout somatic cells in mammalian [30], and the formation of

motile cilia was normal in *Tctex1d2*[-/-] mice generated in this study. The expression

patterns of cytoplasmic dynein 2, kinesin 2, and IFT proteins were normal in *Tctex1d2*

[-/-] mice (fig. S5A–C). Therefore, the loss of TCTEX1D2 had no significant effect on

the IFT system, and cytoplasmic dynein 2 and IFT proteins functioned normally in the

sperm flagella and cilia of *Tctex1d2*[-/-] mice. Indeed, as some components of the IFT

system, such as KIF3A, DYNC2H1, and IFT140, are expressed normally in the flagella

of *Tctex1d2*[-/-] mice, it can be concluded that the IFT system works properly during

sperm flagellum and cilia formation in *Tctex1d2*[-/-] mice. Moreover, TCTEX1D2 was

expressed in the flagella of round spermatids, and *Tctex1d2*[-/-] mice showed abnormal

flagella in early round spermatids, whereas WDR34 and WDR60 were not expressed in

the flagella of round spermatids. As flagellar abnormalities in *Tctex1d2*[-/-] mice

occurred before WDR34 and WDR60 expression, TCTEX1D2 has some functions prior to its

function as cytoplasmic dynein 2. Considering these points, the impairment in

cytoplasmic dynein 2 by *Tctex1d2* deficit has only a minor effect on sperm flagellum

formation therefore, it should be other major causes the flagellar disorders seen

in *Tctex1d2*[-/-] mice.

Another explanation is that TCTEX1D2 functions as an axonemal structure. In

Chlamydomonas, Tctex2b (a homolog of mouse TCTEX1D2) is present in the axoneme as

the light chain of the inner dynein arm [32,44]. In the present study, we found that

TCTEX1D2 was localized in the sperm axoneme and interacted with WDR63 and WDR78. Thus,

it was confirmed that TCTEX1D2 is present in the axoneme as a light chain of the inner

dynein arm. Transmission electron microscopy revealed that the inner dynein arm was

aberrant in the axoneme of the sperm of *Tctex1d2*[-/-] mice. These results indicate that

TCTEX1D2 is important for inner dynein arm assembly. In a previous study, it was shown
that *Wdr63* knockout mice exhibited deficiency of sperm flagella and a disruption of
the axoneme structure similar to what was seen in *Tctex1d2* $^{-/-}$ mice [45]. Disorganization
and structural instability of the inner dynein arm causes sperm flagellar dysplasia.
Therefore, flagellar disorders observed in *Tctex1d2* $^{-/-}$ mice are mainly owing to defects
in the inner dynein arm. We intended to perform ICC studies using anti-WDR63 and
anti-WDR78 antibodies in *Tctex1d2* $^{-/-}$ mice. However, we could not confirm whether the
loss of TCTEX1D2 affects the dynamics of WDR63 and WDR78.

Despite *Tctex1d2* $^{-/-}$ mice having a severely phenotype in sperm flagella, there were
no defects in the cilia of the ventricle, trachea, and oviduct. This difference is
likely owing to the fact that TCTEX1D2 expressed in the cilia of the ventricles,
trachea and oviduct functions only as cytoplasmic dynein 2, not as inner dynein arm,
thus weakening the effects of TCTEX1D2 deficit. In a previous study, it was shown
that the axonemal components and proteins necessary for axoneme formation differed
between sperm flagella and cilia [16]. Therefore, the function of TCTEX1D2 as an inner
dynein arm is unique to sperm flagella. On the other hand, sperm flagella were not
completely absent in *Tctex1d2* $^{-/-}$ mice, normally formed up to a certain length in the
steps 2-3 round spermatids. As flagella are characterized by a much greater length
than cilia, flagellum formation may be undergone a specific change to retain TCTEX1D2
inside the axonemes to maintain the length of flagella. A phenotype similar to that
observed in the sperm of *Tctex1d2* $^{-/-}$ mice was observed with CFAP61 deletion. CFAP61,
which is a conserved component of the calmodulin- and radial spoke-associated complex
of cilia and flagella [46,47], when deleted in mice, resulted in motile cilia of normal
length; however, the sperm flagella failed to maintain their length [47]. Furthermore,
some knockout mice with axonemal components show normal formation of motile cilia
but severe defects in sperm flagella [16]. Therefore, despite the sperm flagella being
regarded as having a structure similar to that of motile cilia, unique molecular
mechanisms give rise to the characteristic length and internal structure during its
formation.

In the dynein light chain related to cilia and flagella formation, *Tctex1d2* was
the first gene to be identified as leading to severe defects in sperm flagella when
knocked out in mice. This study showed that dynein light chains, like other chains,
play an essential role in sperm flagellum formation in mammals. During cilia formation
in mammalian somatic cells, it is known that TCTEX1D2 functions as cytoplasmic dynein
2 [11,18]. However, during sperm flagellum formation, TCTEX1D2 functions as an inner
dynein arm in addition to its function as cytoplasmic dynein 2. Thus, dynein light

chains are involved in many processes related to sperm flagella from formation to
structural maintenance in mammals. *Tctex1d2* [-/-] mice showed defective sperm flagella,
whereas the Tctex2b mutant *Chlamydomonas* did not show defective flagella but only
reduced motility [32,44]. This indicates that the function of homologous genes differs
among species, and it is important to analyze each species.

Multiple morphological abnormalities of the sperm flagella (MMAF) is a male
infertility with genetic component, one of the most severe form of sperm defect [48].
MMAF is characterized abnormal flagella, including short, coiled, absent and of
irregular caliber flagellar [49]. A number of genes have been identified to be associated
with MMAF [50], yet only 35-60% of MMAF cases have been identified and need for further
study to discover new MMAF-related genes [51,52]. The phenotype of *Tctex1d2* [-/-] mice is
similar to cases seen in MMAF, suggesting that Tctex1d2 deficiency may cause MMAF
in human. WDR63, which interacted with TCTEX1D2, is an MMAF-related gene and biallelic
variants in WDR63 were identified in infertility men [45]. Therefore, it is possible
that TCTEX1D2 is also a candidate gene causing MMAF. Our study leads to uncover the
new MMAF causative genes and could be used as a genetic diagnostic indicator for male
infertility.

In summary, we identified *Tctex1d2* as a novel gene essential for the formation
of mouse sperm flagella. *Tctex1d2* [-/-] mice exhibited a severely phenotype only in the
sperm flagella, whereas motile cilia had normal structure and function. TCTEX1D2 is
weakly involved in the assembly of cytoplasmic dynein 2 and IFT proteins and is
important for the assembly and structural stability of the inner dynein arm; this
function is unique to sperm flagellum formation, and is not a part of cilia formation.
These findings provide novel insights into the function of the dynein light chain
in sperm flagellum formation and contribute to the elucidation of the differences
between the formation mechanism of sperm flagella and cilia.

Materials and Methods

Animals

All mice were C57BL/6N background. C57BL/6N mice were purchased from SLC and

maintained under a 12-h light/dark cycle. All animal care and experimental procedures

were conducted according to protocols approved by the Tohoku University Institutional

Animal Care and Use Committee (2019noudou-003-02, 2019noukumikae-030-06).

Generation of _Tctex1d2_ $^{-/-}$ mice and _Tctex1d2_ - 3✕FLAG mice

Tctex1d2 knockouts of the C57BL/6N strain were generated using _i_-GONAD as previously

reported [53]. Briefly, crRNA was designed with the following sequences:

5'-GGTTTGAACCACTACGGTGT-3' and 5'- TATTCAAGAACTAAGCCCTC-3'. In addition, crRNA

and tracrRNA (Integrated DNA Technologies) were annealed at 94°C for 2 min, mixed

with Cas9 nuclease, and injected into the oviduct at E0.7 of plug confirmed mice.

Primer sequences for genotyping are given below.

Fw1:5'-ATGCGTGCAAGGATGAGCTA-3', Rv1:5'-AGGATTTGATCCCCAGCGTG-3', and Rv2:

GCTGGACAACCAGGAAGTGT.

The PCR product from the mutant gDNA was sequenced by Sanger sequencing to confirm

successful deletion. To fix the mutation in the strain, the F0 founder was mated with

wild-type (WT) mice, and the resulting F1 heterozygous male and female mice were

crossed to obtain an F2 generation containing homozygous mutants. The established

strain was maintained by sibling breeding and used in the current study.

The method for the generating _Tctex1d2_ - 3✕FLAG mice was the same as described above.

crRNA was designed with the following sequence: 5'- TTGGACCAGACGCAGGGATA-3'. ssODN

was designed with the following sequence:

5'-GAGCCGCAACCGACCACACTTCCTTCTAGAGATCTGTTGGACCAGCCGCTGAGCTATGGACTACAAGGACCACGAC

GGCGATTATAAGGATCACGACATCGACTACAAAGACGACGATGACAAGGCTGTGTCCTTCCGCGGCCTGTCCTTATCGGC

TCACTCCGAGGGCTTGAGTGAGGTCG-3'. crRNA and tracrRNA (Integrated DNA Technologies)

were annealed, mixed with ssODN and Cas9 nuclease, and injected into the oviduct.

Fertility test

A single 12-week-old male mouse (_Tctex1d2_ $^{+/+}$ or _Tctex1d2_ $^{-/-}$) was caged with two

8-week-old female WT mice for 4 weeks. Mating was verified by monitoring the presence

of vaginal plugs. The plugged females were then replaced with fresh mice. Litter size

was scored for the females that had plugs.

Histological analysis

Mice were euthanasia, and the testes, epididymis, brain, trachea, and oviducts were

collected from *Tctex1d2* [+/+] and *Tctex1d2* [-/-] mice. Testes and epididymides were fixed

in Bouin's solution, whereas brain, trachea, and oviducts were fixed in 4 %

paraformaldehyde. The fixed tissues were gradually dehydrated using graded series

of ethanol and xylene at different concentrations and finally embedded in paraffin.

Sections were rehydrated, the testes and epididymis were stained with periodic

acid-Schiff (PAS), and the ventricle, trachea, and oviduct were stained with

hematoxylin-eosin and examined under a microscope (Olympus BX50).

Immunohistochemistry

Immunohistochemistry (IHC) studies were performed as previously described [54]. Tissue

sections were placed in antigen retrieval buffer (pH 6.0; citrate buffer) for 20 min

at 90 ℃. The following primary antibodies were used: rabbit anti-DYKDDDDK tag

antibody (20543-1-AP, Proteintech), mouse anti-acetylated α tubulin antibody

(sc-23950, Santa Cruz), rabbit anti-TCTEX1 antibody (ab202583, Abcam), rabbit

anti-IFT88 antibody (13967-1-AP, Proteintech), and rabbit anti-KIF3A (13930-1-AP,

Proteintech). The samples were examined under a fluorescence microscope (Keyence

BZ-X710).

Immunocytochemistry of testicular germ cells

For staining individual cells, samples were prepared using the following protocol.

Seminiferous tubules were fixed with 2 % paraformaldehyde containing 0.1 % Triton

X-100 for 10 min at room temperature (RT). Pieces of tubules were placed in a drop

of the fixing solution on a glass slide, and the pieces were minced in the drop. The

slides were gently tapped on the coverslips using tweezers and frozen in liquid

nitrogen for 30 s and coverslips were removed. The slides were washed thrice with

PBS for 5 min and immunostained. The slides were treated with blocking buffer (Blocking

One, Nacalai Tesque, 03953-95) for 1 h at RT and incubated with primary antibody

overnight at 4 ℃. The following primary antibodies were used: mouse anti-acetylated

α tubulin antibody (sc-23950, Santa Cruz), mouse anti-FLAG M2 antibody (F1804,

Sigma-Aldrich), rabbit anti-WDR60 antibody (NBP1-90437, Novus Biologicals), rabbit

anti-WDR34 antibody (NBP1-88805, Novus Biologicals), rabbit anti-TCTEX1 antibody

(ab202583, Abcam), rabbit anti-DYNC2H1 antibody (MBS7050711, MyBioSource), rabbit

anti-DYNC2LI1 antibody (PA5-98717, Thermo Fisher Scientific), rabbit anti-IFT88

antibody (13967-1-AP, Proteintech), rabbit anti-KIF3A (13930-1-AP, Proteintech), and

rabbit anti-IFT140 antibody (17460-1-AP, Proteintech). Acrosomes were stained with

Alexa Fluor 568-conjugated Lectin PNA (L32458; Thermo Fisher Scientific). The samples

were examined under a fluorescence microscope (Keyence BZ-X710).

Western blotting

The tissues (brain, trachea, lung, heart, liver. spleen, kidney, testis) were lysed

using RIPA buffer (Nacalai Tesque, 08714-04) containing 1 % phosphatase inhibitor

(Nacalai Tesque, 07575-51), and incubated for 30 min on ice. The lysates were

centrifuged at 15,000 × g for 15 min at 4 ℃ and eluted with sample buffer for 5

min at 100 ℃. The samples were separated by SDS-PAGE, transferred to a PVDF membrane,

and treated with a commercial blocking buffer (Blocking One, Nacalai Tesque, 03953-95)

for 1 h at RT. The membranes were then incubated with primary antibody overnight at

4 ℃. The following primary antibodies were used: rabbit anti-TCTEX1D2 antibody

(PA5-116015, Thermo Fisher Scientific) and mouse anti-β-actin antibody (Santa Cruz

sc-47778). After washing three times with TBS-T for 5 min, the membranes were

incubated with anti-rabbit or anti-mouse HRP-conjugated antibodies (W4011 and W4021,

Promega) for 1 h at RT. Images were captured using an ImageQuant LAS 500 (Cytiva).

Immunoprecipitation

Testes were lysed using RIPA buffer (Nacalai Tesque, 08714-04), and incubated for

30 min on ice. The lysates were centrifuged at 15,000 × g for 15 min at 4 ℃ and

were incubated for overnight at 4 ℃ with anti-FLAG M2 antibody (F1804,

Sigma-Aldrich). Subsequently, the lysates were conjugated with 25 μl Protein G Mag

Sepharose™ (28944008, cytiva) for 2 h at 4 ℃. The immune complexes were washed three

times with TBS-T for 5 min and eluted with sample buffer for 10 min at 100 ℃. Samples

were separated using SDS-PAGE and western blot analysis was subsequently performed.

Membranes were stripped with a stripping buffer (Nacalai Tesque, 05364-55), and used

for other antibody reactions, as required. The following primary antibodies were used

for western blotting: rabbit anti-TCTEX1D2 (PA5-116015, Thermo Fisher Scientific),

rabbit anti-WDR60 (NBP1-90437, Novus Biologicals), rabbit anti-WDR34 (NBP1-88805,

Novus Biologicals), rabbit anti-TCTEX1 (ab202583, Abcam), mouse anti-WDR78

(sc-390633, Santa Cruz Biotechnology), and rabbit anti-WDR63 antibody (NBP2-32639,

Novus).

Images were captured using the ImageQuant LAS 500 (cytiva).

Fractionation of spermatozoa

Cauda epididymis spermatozoa were suspended in 1 % Triton X-100 lysis buffer (50 mM

NaCl, 20 mM Tris-HCl, pH 7.5, protease inhibitor mixture) and incubated for 2 h at

4 ℃. The sample was centrifuged at 15,000 × g for 10 min to separate the

Triton-soluble fraction (supernatant) from the Triton-resistant fraction (pellet).

The pellet was resuspended in 1 % SDS lysis buffer (75 mM NaCl, 24 mM EDTA, pH 6.0,

protease inhibitor mixture) and incubated for 1 h at RT. The sample was centrifuged

at 15,000 × g for 10 min to separate the SDS-soluble fraction (supernatant) and the

SDS-resistant fraction (pellet). The pellet was dissolved in sample buffer for 10

min at 100 ℃. Samples were separated using SDS-PAGE and western blot analysis was

subsequently performed. The following primary antibodies were used: rabbit

anti-DYKDDDDK tag antibody (20543-1-AP, Proteintech), rabbit anti-GLUT3 antibody

(20403-1-AP, Proteintech), mouse anti-acetylated α tubulin antibody (sc-23950,

Santa Cruz), and rabbit anti-AKAP3 antibody (13907-1-A, Proteintech). The samples

were examined under a fluorescence microscope (Keyence BZ-X710).

Immunofluorescence for spermatozoa

Cauda epididymis spermatozoa was incubated in human tubular fluid (HTF) medium for

1 h at 37.5 ℃ with 5 % CO_2, and swim-up spermatozoa was collected and washed with

PBS. The suspension was fixed in 4 % paraformaldehyde for 10 min at RT and air dried

on slide glass. The slides were permeabilized with 0.2 % Triton X-100 for 10 min at

37 ℃ and were washed three times in PBS at 5 min. The slides were treated with blocking

buffer (Blocking One, Nacalai Tesque, 03953-95) for 1 h at RT and incubated with

primary antibody overnight at 4 ℃. The following primary antibodies were used: rabbit

anti-DYKDDDDK tag antibody (20543-1-AP, Proteintech) and mouse anti-acetylated α

tubulin antibody (sc-23950, Santa Cruz). The samples were examined under a

fluorescence microscope (Keyence BZ-X710).

Sperm motility, count, and abnormal sperm ratio analyses

Cauda epididymis sperm was incubated in HTF sodium for 1 h at 37.5 ℃ with 5 % CO_2.

Sperm motility parameters were evaluated using a computer-assisted sperm analysis

(CASA) system (SMAS; DITECT). Sperm count was analyzed using a hemocytometer. Abnormal

sperm ratios were analyzed using phase-contrast microscopy, and 200 sperms from each

sample were observed.

Transmission electron microscopy

Tctex1d2$^{+/+}$ or *Tctex1d2*$^{-/-}$ mice testes were cut into 1-mm-thick sections and were fixed

with 2.5 % glutaraldehyde in 0.1 M phosphate buffer (PB) for 1.5 h at RT. After the

tissues were washed in PB three times, they were additionally fixed with 1 % OsO$_4$ in

0.1 M PB for 1.5 h at 4 ℃. The tissues were dehydrated using a graded ethanol series

for 10 min at RT, which was then replaced with 50 % propylene oxide (PO) in ethanol

followed by 100 % PO for 10 min each at RT. The tissues were permeabilized with a

mixture of PO and epoxy resin at RT (PO: epoxy resin = 2:1 for 2 h, PO: epoxy resin

= 1:1 for 4 h, PO: epoxy resin = 1:2 for 12 h, only epoxy resin for 24 h). The tissues

were embedded in epoxy resin for 12 h at 37 ℃, for 12 h at 45 ℃, and for 12 h at

60 ℃. Ultrathin sectioning, staining, and imaging were performed at an external lab.

Statistical analyses

The unpaired Student's t-test (Microsoft Office Excel; Microsoft Corporation) was

used for comparisons with $Tctex1d2^{+/+}$ mice. Values are represented as the mean $\pm$ SE.

Differences were considered significant at $p < 0.05$, indicated as $p < 0.05$ (*).

 Figure legends

Figure. 1. TCTEX1D2 was high expression in the testis and localized along the flagella

from round spermatids to elongated spermatids and the manchette. (A) Protein

expression analysis of mouse TCTEX1D2 in various organs using western blotting (WB);

TCTEX1D2 was highly expressed in the testes and spleen. (B-D) ICC in WT and *Tctex1d2*

- 3×FLAG germ cells to analyze the localization of TCTEX1D2 (green: anti-FLAG, red:

anti-acetylated-α-tubulin, blue: hoechst33342). TCTEX1D2 localized to flagella from

early round spermatids to elongated spermatids. In addition, TCTEX1D2 localized to

the manchette. (B) Round spermatids in steps 1-3. (C) Elongated spermatids of steps

10-12. (D) Elongated spermatids of steps 15-16. Scale bars are 10 μm.

Figure. 2. TCTEX1D2 interacted with the subunits of cytoplasmic dynein 2 and axonemal

dynein. (A) Schematic diagram of the cytoplasmic dynein 2 complex in somatic cell.

(B) Co-immunoprecipitation in the testes of *Tctex1d2* - 3×FLAG mice. In the testes,

TCTEX1D2 interacted with WDR60, WDR34, and DYNLT1. (C) WB of cauda epididymis sperm

fractionated into Triton X-100 soluble, SDS-soluble, and SDS-insoluble fractions from

WT and *Tctex1d2* - 3×FLAG mice. TCTEX1D2 detected in SDS-soluble and SDS-insoluble

fractions. (D) ICC of cauda epididymis sperm from WT and *Tctex1d2* - 3×FLAG mice

(green: FLAG, red: acetylated-α-tubulin, blue: hoechst33342). TCTEX1D2 localized

along the flagella. Asterisks indicate non-specific signals. Scale bars are 10 μm.

(E) Co-immunoprecipitation with WDR63 and WDR78 in the testes of *Tctex1d2* - 3×FLAG

mice. TCTEX1D2 interacted with WDR63 and WDR78. The used membrane was same as that

in (B).

Figure. 3. TCTEX1D2 was required for male fertility. (A) Schematic representation

of the generation of *Tctex1d2* $^{-/-}$ mice. The gRNA was designed between exons 1 and 2

and between exons 3 and 4. Genotyping primer pairs were Fw1 and Rv1 for *Tctex1d2* $^{+/+}$

mice, Fw1 and Rv2 for *Tctex1d2* $^{-/-}$ mice. (B) Genotyping and (C) Sanger sequencing showed

that *Tctex1d2* $^{-/-}$ mice had a 4696 bp deletion within the target region. (D) WB of tissue

from mouse testes confirmed that TCTEX1D2 protein expression was completely lost in

Tctex1d2 $^{-/-}$ mice. (E) Male *Tctex1d2* $^{+/+}$ and *Tctex1d2* $^{-/-}$ mice were mated with two female

WT mice. We observed 33 plugs in *Tctex1d2* $^{-/-}$ mice; however, *Tctex1d2* $^{-/-}$ mice were unable

to sire any offspring. (F) The ratio of testes weight / body weight. There were no

differences between *Tctex1d2* $^{+/+}$ and *Tctex1d2* $^{-/-}$ mice. Error bars indicate means ±

S.E. Two-tailed Student's t-test, n=3, n.s. : no significant.

Figure. 4. *Tctex1d2* is important for sperm flagellum formation. (A–D) Periodic acid
Schiff hematoxylin (PAS-H) stain of seminiferous tubules of *Tctex1d2* $^{+/+}$ and *Tctex1d2*
$^{-/-}$ mice. In stage I-II seminiferous tubules, *Tctex1d2* $^{-/-}$ mice showed layered structures
of germ cells (B; yellow box), similar to that in *Tctex1d2* $^{+/+}$ mice (A; yellow box).
However, sperm flagella were barely observed in the lumen in *Tctex1d2* $^{-/-}$ mice (B; red
box) compared to *Tctex1d2* $^{+/+}$ mice (A; red box arrow). In the lumen of stage VII
seminiferous tubules, only a few elongated spermatids were observed in *Tctex1d2* $^{-/-}$
mice (D; black box) in contrast to *Tctex1d2* $^{+/+}$ mice (C; black box). Scale bars are
50 µm. (E) Hematoxylin and eosin (H&E) stain of the cauda epididymis. Many spermatozoa
were observed in *Tctex1d2* $^{+/+}$ mice, whereas only very few spermatozoa were observed
in *Tctex1d2* $^{-/-}$ mice. Scale bars are 50 µm. (F) Sperm counts in the cauda epididymis.
Tctex1d2 $^{-/-}$ mice had significantly reduced sperm counts compared to *Tctex1d2* $^{+/+}$ mice.
Error bars are means ± S.E. Two-tailed Student's t-test, n=3, $*p < 0.05$. (G) Cauda
epididymis sperm observed using phase contrast microscopy. Sperm flagella of *Tctex1d2*
$^{-/-}$ mice showed morphological abnormalities. Scale bars are 10 µm. (H) The ratio of
sperm with abnormal flagella in the cauda epididymis. Almost all sperm showed abnormal
flagella in *Tctex1d2* $^{-/-}$ mice. Error bars are means ± S.E. Two-tailed Student's t-test,
n=3, $*p < 0.05$.

Figure. 5. Abnormal assembly of sperm flagella in *Tctex1d2* $^{-/-}$ mice occurred from step
4 round spermatids. (A) Flagella formation in spermiogenesis using ICCs (green:
anti-acetylated-α-tubulin, red: PNA lectin, blue: hoechst33342). In *Tctex1d2* $^{+/+}$ mice,
flagella elongated as the spermatids differentiated. However, *Tctex1d2* $^{-/-}$ mice showed
markedly abnormal flagella after step 4 spermatids. Scale bars are 10 µm. (B)
Observation of the internal structure of the flagella of spermatids using transmission
electron microscopy. In *Tctex1d2* $^{+/+}$ mice, microtubule 9+2 structures and other
structures were neatly aligned in the axonemes. In contrast, in *Tctex1d2* $^{-/-}$ mice, the
internal structure of the axonemes were disorganized. The 9+2 structures, inner dynein
arm (IDA), outer dynein arm (ODA), outer dense fiber (ODF), and fibrous sheath (FS)
were not aligned properly. Scale bars are 250 nm. MS, mitochondrial sheath; DMTs,
doublet microtubules; ODF, outer dense fibers; FS, fibrous sheath; ODA, outer dynein
arm; IDA, inner dynein arm; MTs, microtubule (not doublet).

Figure. 6. TCTEX1D2 played a role in the assembly of cytoplasmic dynein 2 subunits.
(A) Expression analysis of WDR60 in *Tctex1d2* $^{-/-}$ mice using ICC (green: WDR60, red:
acetylated-α-tubulin, blue: hoechst33342). There was no expression of WDR60 in

flagella of round spermatids in *Tctex1d2* $^{+/+}$ mice. In elongated spermatids, abnormal

expression was observed in *Tctex1d2* $^{-/-}$ mice (top: loss of signal, bottom: ectopic

expression). (B) Expression analysis of WDR34 in *Tctex1d2* $^{-/-}$ mice using ICC (green:

WDR34, red: acetylated-α-tubulin, blue: hoechst33342). There was no expression of

WDR34 was in flagella of round spermatids, as seen with WDR60. In elongated spermatids,

abnormal expression was observed in *Tctex1d2* $^{-/-}$ mice (top: loss of signal, bottom:

ectopic expression). (C) Expression analysis of DYNLT1 in *Tctex1d2* $^{-/-}$ mice using ICC

(green: DYNLT1, red: acetylated-α-tubulin, blue: hoechst33342). The signal was

decreased in round spermatids and absent in elongated spermatids in *Tctex1d2* $^{-/-}$ mice.

RST, round spermatid; EST, elongated spermatid. (D) Schematic diagram of the defect

of cytoplasmic dynein 2 subunits in *Tctex1d2* $^{-/-}$ mice. In *Tctex1d2* $^{-/-}$ mice elongated

spermatids, DYNC2LI1, WDR60, WDR34 and DYNLT1 showed abnormal localization and were

not assembled in the cytoplasmic dynein 2 complex.

Supplementary Material

Supplement figure. 1. Generation of *Tctex1d2* - 3×FLAG mice. (A) Schematic

representation of *Tctex1d2* - 3×FLAG mice and the results of Sanger sequencing.

3×FLAG was knocked immediately after the start codon. Sanger sequencing indicated

that the 3 × FLAG nucleotide sequence was knocked-in within the target region. (B)

PAS-H staining of seminiferous tubules of WT and *Tctex1d2* - 3×FLAG mice. Scale bars

are 50 µm. (C) Phase contrast microscopy of cauda epididymis sperm. There were no

differences of sperm morphology between WT and *Tctex1d2* - 3×FLAG mice. Scale bars

are 10 µm.

Supplement figure. 2. TCTEX1D2 was also localized to motile cilia. IHC of WT and

Tctex1d2 - 3×FLAG in tissues with motile cilia to analyze the expression of TCTEX1D2

(green: anti-FLAG, red: anti-acetylated-α-tubulin, blue: hoechst33342). (A)

ventricle, (B) trachea, and (C) oviduct. TCTEX1D2 is expressed in the cilia of the

ventricle, trachea, and oviduct. (*); The signals found in the epithelial cells of

each tissue are non-specific. Scale bars are 10 µm.

Supplement figure. 3. Motile cilia in *Tctex1d2* $^{-/-}$ mice were normal. H&E stain of tissues

with motile cilia. (A) ventricle, (B) trachea, and (C) oviduct. In motile cilia, there

were no differences in morphology between *Tctex1d2* $^{+/+}$ and *Tctex1d2* $^{-/-}$ mice. Scale bars

are 20 µm.

Supplement figure. 4. *Tctex1d2* $^{-/-}$ mice exhibited abnormal protein localization in the

IFT system. Expression analysis of cytoplasmic dynein 2 (DYNC2H1 and DYNC2LI1),

kinesin 2 (KIF3A), and IFT proteins (IFT88, 140) in elongated spermatids of *Tctex1d2*

$^{-/-}$ mice using ICC (red: anti-acetylated-α-tubulin, Blue: hoechst33342). (A) IFT88,

(B) DYNC2LI1, (C) IFT140, (D) DYNC2H1, and (E) KIF3A. IFT88 and DYNC2LI1 showed a

reduced signal and were ectopically expressed in the flagella of *Tctex1d2* $^{-/-}$ mice,

whereas IFT140, DYNC2H1, and KIF3A were expressed normally in *Tctex1d2* $^{-/-}$ mice. Scale

bars are 10 μm.

Supplement figure. 5. The IFT system was normal in the cilia of the ventricles.

Expression analysis of cytoplasmic dynein 2 (DYNLT1) and IFT protein (IFT88) in the

ventricle of *Tctex1d2* $^{-/-}$ mice using IHC (red: anti-acetylated-α-tubulin, blue:

hoechst33342). In *Tctex1d2* $^{-/-}$ mice, the two proteins showed abnormal expression in

the flagella of elongated spermatids, but not in the ventricle. Scale bars are 10

μm.

Funding

This work was supported by the JSPS KAKENHI (19H01142 to K.T., 23K21259 to K.H.) and JST FOREST Program (JPMJFR2018 to K.H.) and JST SPRING (JPMJSP2114 to R.H.).

Author contributions

Ryua Harima, Conceptualization, Methodology, Validation, Formal analysis, Investigation, Resources, Data curation, Writing – original draft preparation, Writing – review & editing, Visualization, Project administration, Funding acquisition; Kenshiro Hara, Resources, Writing – original draft preparation, Writing – review & editing, Funding acquisition; Kentaro Tanemura; Resources, Supervision, Writing – original draft preparation, Writing – review & editing, Funding acquisition.

Acknowledgments

We would like to thank Editage (www.editage.com) for English language editing.

Reference

1. Gibbons, I.R. (1981). Cilia and flagella of eukaryotes. J Cell Biol *91*, 107s-124s.
10.1083/jcb.91.3.107s.

2. Carvalho-Santos, Z., Azimzadeh, J., Pereira-Leal, J.B., and Bettencourt-Dias, M.
(2011). Evolution: Tracing the origins of centrioles, cilia, and flagella. J Cell
Biol *194*, 165-175. 10.1083/jcb.201011152.

3. Lindemann, C.B., and Lesich, K.A. (2016). Functional anatomy of the mammalian sperm
flagellum. Cytoskeleton (Hoboken) *73*, 652-669. 10.1002/cm.21338.

4. Satir, P., and Christensen, S.T. (2007). Overview of structure and function of
mammalian cilia. Annu Rev Physiol *69*, 377-400.
10.1146/annurev.physiol.69.040705.141236.

5. Allen, R.D. (1968). A reinvestigation of cross-sections of cilia. J Cell Biol *37*,
825-831. 10.1083/jcb.37.3.825.

6. Luck, D.J. (1984). Genetic and biochemical dissection of the eucaryotic flagellum.
J Cell Biol *98*, 789-794. 10.1083/jcb.98.3.789.

7. Porter, M.E., and Sale, W.S. (2000). The 9 + 2 axoneme anchors multiple inner arm
dyneins and a network of kinases and phosphatases that control motility. J Cell
Biol *151*, F37-42. 10.1083/jcb.151.5.f37.

8. Pazour, G.J., Wilkerson, C.G., and Witman, G.B. (1998). A dynein light chain is
essential for the retrograde particle movement of intraflagellar transport (IFT).
J Cell Biol *141*, 979-992. 10.1083/jcb.141.4.979.

9. Rosenbaum, J.L., and Witman, G.B. (2002). Intraflagellar transport. Nat Rev Mol
Cell Biol *3*, 813-825. 10.1038/nrm952.

10. Mohri, H., Inaba, K., Ishijima, S., and Baba, S.A. (2012). Tubulin-dynein system
in flagellar and ciliary movement. Proc Jpn Acad Ser B Phys Biol Sci *88*, 397-415.
10.2183/pjab.88.397.

11. Asante, D., Stevenson, N.L., and Stephens, D.J. (2014). Subunit composition of
the human cytoplasmic dynein-2 complex. J Cell Sci *127*, 4774-4787.
10.1242/jcs.159038.

12. Pfister, K.K., Shah, P.R., Hummerich, H., Russ, A., Cotton, J., Annuar, A.A., King,
S.M., and Fisher, E.M. (2006). Genetic analysis of the cytoplasmic dynein subunit
families. PLoS Genet *2*, e1. 10.1371/journal.pgen.0020001.

13. Roberts, A.J., Kon, T., Knight, P.J., Sutoh, K., and Burgess, S.A. (2013).
Functions and mechanics of dynein motor proteins. Nat Rev Mol Cell Biol *14*, 713-726.
10.1038/nrm3667.

14. Pazour, G.J., Dickert, B.L., and Witman, G.B. (1999). The DHC1b (DHC2) isoform

of cytoplasmic dynein is required for flagellar assembly. J Cell Biol *144*, 473-481.

10.1083/jcb.144.3.473.

15.   Summers, K.E., and Gibbons, I.R. (1971). Adenosine triphosphate-induced sliding

of tubules in trypsin-treated flagella of sea-urchin sperm. Proc Natl Acad Sci

U S A *68*, 3092-3096. 10.1073/pnas.68.12.3092.

16.   Inaba, K., and Mizuno, K. (2016). Sperm dysfunction and ciliopathy. Reprod Med

Biol *15*, 77-94. 10.1007/s12522-015-0225-5.

17.   Mikami, A., Tynan, S.H., Hama, T., Luby-Phelps, K., Saito, T., Crandall, J.E.,

Besharse, J.C., and Vallee, R.B. (2002). Molecular structure of cytoplasmic dynein

2 and its distribution in neuronal and ciliated cells. J Cell Sci *115*, 4801-4808.

10.1242/jcs.00168.

18.   Vuolo, L., Stevenson, N.L., Mukhopadhyay, A.G., Roberts, A.J., and Stephens, D.J.

(2020). Cytoplasmic dynein-2 at a glance. J Cell Sci *133*. 10.1242/jcs.240614.

19.   Cole, D.G. (2003). The intraflagellar transport machinery of Chlamydomonas

reinhardtii. Traffic *4*, 435-442. 10.1034/j.1600-0854.2003.t01-1-00103.x.

20.   Signor, D., Wedaman, K.P., Orozco, J.T., Dwyer, N.D., Bargmann, C.I., Rose, L.S.,

and Scholey, J.M. (1999). Role of a class DHC1b dynein in retrograde transport

of IFT motors and IFT raft particles along cilia, but not dendrites, in chemosensory

neurons of living Caenorhabditis elegans. J Cell Biol *147*, 519-530.

10.1083/jcb.147.3.519.

21.   Toropova, K., Zalyte, R., Mukhopadhyay, A.G., Mladenov, M., Carter, A.P., and

Roberts, A.J. (2019). Structure of the dynein-2 complex and its assembly with

intraflagellar transport trains. Nat Struct Mol Biol *26*, 823-829.

10.1038/s41594-019-0286-y.

22.   Reck-Peterson, S.L., Redwine, W.B., Vale, R.D., and Carter, A.P. (2018). The

cytoplasmic dynein transport machinery and its many cargoes. Nat Rev Mol Cell Biol

19, 382-398. 10.1038/s41580-018-0004-3.

23.   Braschi, B., Omran, H., Witman, G.B., Pazour, G.J., Pfister, K.K., Bruford, E.A.,

and King, S.M. (2022). Consensus nomenclature for dyneins and associated assembly

factors. J Cell Biol *221*. 10.1083/jcb.202109014.

24.   Viswanadha, R., Sale, W.S., and Porter, M.E. (2017). Ciliary Motility: Regulation

of Axonemal Dynein Motors. Cold Spring Harb Perspect Biol *9*.

10.1101/cshperspect.a018325.

25.   Konno, A., Shiba, K., Cai, C., and Inaba, K. (2015). Branchial cilia and sperm

flagella recruit distinct axonemal components. PLoS One *10*, e0126005.

10.1371/journal.pone.0126005.

26.   Sarpal, R., Todi, S.V., Sivan-Loukianova, E., Shirolikar, S., Subramanian, N.,
Raff, E.C., Erickson, J.W., Ray, K., and Eberl, D.F. (2003). Drosophila KAP
interacts with the kinesin II motor subunit KLP64D to assemble chordotonal sensory
cilia, but not sperm tails. Curr Biol *13*, 1687-1696. 10.1016/j.cub.2003.09.025.
27.   Han, Y.G., Kwok, B.H., and Kernan, M.J. (2003). Intraflagellar transport is
required in Drosophila to differentiate sensory cilia but not sperm. Curr Biol
13, 1679-1686. 10.1016/j.cub.2003.08.034.
28.   Gholkar, A.A., Senese, S., Lo, Y.C., Capri, J., Deardorff, W.J., Dharmarajan, H.,
Contreras, E., Hodara, E., Whitelegge, J.P., Jackson, P.K., and Torres, J.Z. (2015).
Tctex1d2 associates with short-rib polydactyly syndrome proteins and is required
for ciliogenesis. Cell Cycle *14*, 1116-1125. 10.4161/15384101.2014.985066.
29.   Schmidts, M., Hou, Y., Cortés, C.R., Mans, D.A., Huber, C., Boldt, K., Patel, M.,
van Reeuwijk, J., Plaza, J.M., van Beersum, S.E., et al. (2015). TCTEX1D2 mutations
underlie Jeune asphyxiating thoracic dystrophy with impaired retrograde
intraflagellar transport. Nat Commun *6*, 7074. 10.1038/ncomms8074.
30.   Hamada, Y., Tsurumi, Y., Nozaki, S., Katoh, Y., and Nakayama, K. (2018).
Interaction of WDR60 intermediate chain with TCTEX1D2 light chain of the dynein-2
complex is crucial for ciliary protein trafficking. Mol Biol Cell *29*, 1628-1639.
10.1091/mbc.E18-03-0173.
31.   Green, C.D., Ma, Q., Manske, G.L., Shami, A.N., Zheng, X., Marini, S., Moritz,
L., Sultan, C., Gurczynski, S.J., Moore, B.B., et al. (2018). A Comprehensive
Roadmap of Murine Spermatogenesis Defined by Single-Cell RNA-Seq. Dev Cell *46*,
651-667.e610. 10.1016/j.devcel.2018.07.025.
32.   DiBella, L.M., Smith, E.F., Patel-King, R.S., Wakabayashi, K., and King, S.M.
(2004). A novel Tctex2-related light chain is required for stability of inner
dynein arm I1 and motor function in the Chlamydomonas flagellum. J Biol Chem *279*,
21666-21676. 10.1074/jbc.M313540200.
33.   Brokaw, C.J., and Kamiya, R. (1987). Bending patterns of Chlamydomonas flagella:
IV. Mutants with defects in inner and outer dynein arms indicate differences in
dynein arm function. Cell Motil Cytoskeleton *8*, 68-75. 10.1002/cm.970080110.
34.   Huang, B., Piperno, G., and Luck, D.J. (1979). Paralyzed flagella mutants of
Chlamydomonas reinhardtii. Defective for axonemal doublet microtubule arms. J Biol
Chem *254*, 3091-3099.
35.   Yamamoto, R., Obbineni, J.M., Alford, L.M., Ide, T., Owa, M., Hwang, J., Kon, T.,
Inaba, K., James, N., King, S.M., et al. (2017). Chlamydomonas DYX1C1/PF23 is
essential for axonemal assembly and proper morphology of inner dynein arms. PLoS

Genet *13*, e1006996. 10.1371/journal.pgen.1006996.

36.   Yamamoto, R., Yanagi, S., Nagao, M., Yamasaki, Y., Tanaka, Y., Sale, W.S., Yagi,
T., and Kon, T. (2020). Mutations in PIH proteins MOT48, TWI1 and PF13 define common
and unique steps for preassembly of each, different ciliary dynein. PLoS Genet
16, e1009126. 10.1371/journal.pgen.1009126.

37.   Cole, D.G., Diener, D.R., Himelblau, A.L., Beech, P.L., Fuster, J.C., and Rosenbaum,
J.L. (1998). Chlamydomonas kinesin-II-dependent intraflagellar transport (IFT):
IFT particles contain proteins required for ciliary assembly in Caenorhabditis
elegans sensory neurons. J Cell Biol *141*, 993-1008. 10.1083/jcb.141.4.993.

38.   Stepanek, L., and Pigino, G. (2016). Microtubule doublets are double-track
railways for intraflagellar transport trains. Science *352*, 721-724.
10.1126/science.aaf4594.

39.   Nachury, M.V., Loktev, A.V., Zhang, Q., Westlake, C.J., Peränen, J., Merdes, A.,
Slusarski, D.C., Scheller, R.H., Bazan, J.F., Sheffield, V.C., and Jackson, P.K.
(2007). A core complex of BBS proteins cooperates with the GTPase Rab8 to promote
ciliary membrane biogenesis. Cell *129*, 1201-1213. 10.1016/j.cell.2007.03.053.

40.   Wu, C., Li, J., Peterson, A., Tao, K., and Wang, B. (2017). Loss of dynein-2
intermediate chain Wdr34 results in defects in retrograde ciliary protein
trafficking and Hedgehog signaling in the mouse. Hum Mol Genet *26*, 2386-2397.
10.1093/hmg/ddx127.

41.   Vuolo, L., Stevenson, N.L., Heesom, K.J., and Stephens, D.J. (2018). Dynein-2
intermediate chains play crucial but distinct roles in primary cilia formation
and function. Elife *7*. 10.7554/eLife.39655.

42.   Nakayama, K., and Katoh, Y. (2018). Ciliary protein trafficking mediated by IFT
and BBSome complexes with the aid of kinesin-2 and dynein-2 motors. J Biochem *163*,
155-164. 10.1093/jb/mvx087.

43.   Kessler, K., Wunderlich, I., Uebe, S., Falk, N.S., Gießl, A., Brandstätter, J.H.,
Popp, B., Klinger, P., Ekici, A.B., Sticht, H., et al. (2015). DYNC2LI1 mutations
broaden the clinical spectrum of dynein-2 defects. Sci Rep *5*, 11649.
10.1038/srep11649.

44.   Fu, G., Scarbrough, C., Song, K., Phan, N., Wirschell, M., and Nicastro, D. (2021).
Structural organization of the intermediate and light chain complex of
Chlamydomonas ciliary I1 dynein. Faseb j *35*, e21646. 10.1096/fj.202001857R.

45.   Lu, S., Gu, Y., Wu, Y., Yang, S., Li, C., Meng, L., Yuan, W., Jiang, T., Zhang,
X., Li, Y., et al. (2021). Bi-allelic variants in human WDR63 cause male infertility
via abnormal inner dynein arms assembly. Cell Discov *7*, 110.

781   10.1038/s41421-021-00327-5.

782 46. Urbanska, P., Song, K., Joachimiak, E., Krzemien-Ojak, L., Koprowski, P.,

783   Hennessey, T., Jerka-Dziadosz, M., Fabczak, H., Gaertig, J., Nicastro, D., and

784   Wloga, D. (2015). The CSC proteins FAP61 and FAP251 build the basal substructures

785   of radial spoke 3 in cilia. Mol Biol Cell *26*, 1463-1475. 10.1091/mbc.E14-11-1545.

786 47. Liu, S., Zhang, J., Kherraf, Z.E., Sun, S., Zhang, X., Cazin, C., Coutton, C.,

787   Zouari, R., Zhao, S., Hu, F., et al. (2021). CFAP61 is required for sperm flagellum

788   formation and male fertility in human and mouse. Development *148*.

789   10.1242/dev.199805.

790 48. Coutton, C., Escoffier, J., Martinez, G., Arnoult, C., and Ray, P.F. (2015).

791   Teratozoospermia: spotlight on the main genetic actors in the human. Hum Reprod

792   Update *21*, 455-485. 10.1093/humupd/dmv020.

793 49. Ben Khelifa, M., Coutton, C., Zouari, R., Karaouzène, T., Rendu, J., Bidart, M.,

794   Yassine, S., Pierre, V., Delaroche, J., Hennebicq, S., et al. (2014). Mutations

795   in DNAH1, which encodes an inner arm heavy chain dynein, lead to male infertility

796   from multiple morphological abnormalities of the sperm flagella. Am J Hum Genet

797   *94*, 95-104. 10.1016/j.ajhg.2013.11.017.

798 50. Touré, A., Martinez, G., Kherraf, Z.E., Cazin, C., Beurois, J., Arnoult, C., Ray,

799   P.F., and Coutton, C. (2021). The genetic architecture of morphological

800   abnormalities of the sperm tail. Hum Genet *140*, 21-42. 10.1007/s00439-020-02113-x.

801 51. Liu, C., He, X., Liu, W., Yang, S., Wang, L., Li, W., Wu, H., Tang, S., Ni, X.,

802   Wang, J., et al. (2019). Bi-allelic Mutations in TTC29 Cause Male Subfertility

803   with Asthenoteratospermia in Humans and Mice. Am J Hum Genet *105*, 1168-1181.

804   10.1016/j.ajhg.2019.10.010.

805 52. Lorès, P., Dacheux, D., Kherraf, Z.E., Nsota Mbango, J.F., Coutton, C., Stouvenel,

806   L., Ialy-Radio, C., Amiri-Yekta, A., Whitfield, M., Schmitt, A., et al. (2019).

807   Mutations in TTC29, Encoding an Evolutionarily Conserved Axonemal Protein, Result

808   in Asthenozoospermia and Male Infertility. Am J Hum Genet *105*, 1148-1167.

809   10.1016/j.ajhg.2019.10.007.

810 53. Gurumurthy, C.B., Sato, M., Nakamura, A., Inui, M., Kawano, N., Islam, M.A.,

811   Ogiwara, S., Takabayashi, S., Matsuyama, M., Nakagawa, S., et al. (2019). Creation

812   of CRISPR-based germline-genome-engineered mice without ex vivo handling of

813   zygotes by i-GONAD. Nat Protoc *14*, 2452-2482. 10.1038/s41596-019-0187-x.

814 54. Hiradate, Y., Harima, R., Yanai, R., Hara, K., Nagasawa, K., Osada, M., Kobayashi,

815   T., Matsuyama, M., Kanno, S.I., Yasui, A., and Tanemura, K. (2022). Loss of Axdnd1

816   causes sterility due to impaired spermatid differentiation in mice. Reprod Med

817   Biol *21*, e12452. 10.1002/rmb2.12452.

stage I～II

stage VII

figure. 4

figure. 5

figure. 6

A

WT
Tctex1d2 exon1 ... exon5

start codon
ATG **GACTACAAGGACCACGACGGCGATTATAAGGATCACGACATCGACTACAAAGACGACGATGACAAG** GCTG

3×FLAG

B

stage I～II   stage VII

WT

Tctex1d2-3×FLAG

C

WT   *Tctex1d2*-3×FLAG

<u>**Supplement. 1**</u>

A

ventricle

FLAG

WT

B

trachea

FLAG

WT

C

oviduct

FLAG

WT

FLAG   ac-α-tubulin   hoechst

A ventricle

Tctex1d2 $^{+/+}$ *Tctex1d2* $^{-/-}$

B trachea

Tctex1d2 $^{+/+}$ *Tctex1d2* $^{-/-}$

C oviduct

Tctex1d2 $^{+/+}$ *Tctex1d2* $^{-/-}$

A +/+ -/-

IFT88   ac-α-tubulin   hoechst

B +/+ -/-

DYNC2LI1   ac-α-tubulin   hoechst

C +/+ -/-

IFT140   ac-α-tubulin   hoechst

D +/+ -/-

DYNC2H1   ac-α-tubulin   hoechst

E +/+ -/-

KIF3A   ac-α-tubulin   hoechst

A

DYNLT1

+/+   -/-

B

IFT88

+/+   -/-

C

KIF3A

+/+   -/-

ac-α-tubulin   **hoechst**