Polymerization of Tspan7 into Helical Transmembrane Skeletons for Membrane Tether Stabilization

Dongju Wang[1, 2, 5], Xia Jia[1, 2, 5], Qi Zhang[1, 4], Kui Xu[1, 2, 3], Yi Zheng[1, 2], Qiangfeng Cliff Zhang[1, 2, 3], Nan Liu[1, 2*], Hong-Wei Wang[1, 2*], Li Yu[1, 2*]

[1]Beijing Frontier Research Center for Biological Structure, School of Life Sciences, Tsinghua University, Beijing 100084, China.

[2]The State Key Laboratory of Membrane Biology, Tsinghua University-Peking University Joint Centre for Life Sciences, Tsinghua University, Beijing 100084, China.

[3]Ministry of Education Key Laboratory of Bioinformatics, Center for Synthetic and Systems Biology; Beijing, 100084, China.

[4]Institute of Bio-Architecture and Bio-Interactions (IBABI), Shenzhen Medical Academy of Research and Translation, Shenzhen, China

[5]These authors contributed equally to this work.

*Correspondence: Li Yu, liyulab@tsinghua.edu.cn;

Hong-Wei Wang, hongweiwang@tsinghua.edu.cn or Nan Liu, nanliu@hku.hk

Summary

Membrane tethers, thin tubular extensions of the plasma membrane, form during cell migration, adhesion, and signaling. Various types exist, though their maintenance mechanisms are not fully understood. This study employs live cell imaging, *in vitro* reconstitution, and *in situ* cryo-EM analysis to show that Tspan7, typically a freely diffusing transmembrane protein, forms long cylindrical polymers in right-handed helical formations on retraction fibers. These polymers assemble into spiral tubule structures, acting as a "transmembrane skeleton", enriching Tspan7 on retraction fibers and promoting their formation. Our findings reveal a novel assembly strategy for transmembrane proteins and a mechanism for maintaining membrane tethers.

Keywords

Tetraspanin 7, membrane tether, *in situ* cryo-electron tomography, spiral assembly

Introduction

Membrane tethers, characterized by their dynamic, finger-like extensions from the cell membrane, are pivotal in numerous physiological and pathological contexts[1-3]. These structures, which exhibit diversity in their size, shape, and functionality, underpin critical cellular processes such as motility, tissue morphogenesis, immune function, and cancer metastasis[2,4-7]. Among these, retraction fibers stand out as slender, elongated protrusions that maintain attachment to the substrate during cellular migration, acting as mechanical tethers that facilitate both force transmission and intercellular communication[8-10]. Furthermore, retraction fibers play a key role in the formation of

migrasomes, organelles that emerge at retraction fiber junctions and tips in migrating

cells, highlighting their importance in cellular signaling and material transfer[11-15].

Membrane tethers experience constant and dynamic mechanical stress. In the current

model, the mechanical characteristics of these tethers are mainly determined by the

dynamic of cytoskeleton, alongside their interactions with other cytoskeletal

elements[2,16,17]. It remains unclear if additional structures can also influence the

mechanical properties of membrane tethers.

Tetraspanins are a family of integral membrane proteins characterized by four

transmembrane domains. Tetraspanins are relatively small proteins, typically ranging

from 200 to 350 amino acids in length, corresponding to a molecular weight of

approximately 20 to 50 kDa[18]. They play crucial roles in organizing membrane

microdomains, facilitating the formation of multiprotein complexes, and influencing

cell morphology[19]. Tetraspanins are involved in various cellular processes such as

membrane repair, cell migration, adhesion, signal transduction[20-23] and the formation of

specialized membrane structures like filopodia, microvilli, and dendritic spines[24-27].

They also function as curvature sensors and are essential for the formation and function

of structures like retraction fibers and migrasomes[28-30].

Tetraspanins typically aggregate into tetraspanin-enriched microdomains on

biological membranes, where they coordinate with associated proteins and specific

lipids, such as cholesterol, to form nanoscale assemblies[31,32]. Additionally, certain

tetraspanins, notably Tspan21/UP1a and Tspan20/UP1b, are key components of

urothelial plaques[33]. These plaques consist of two-dimensional crystals with a

hexagonal arrangement of 16-nm protein particles on the urothelium membrane[34],

showcasing a different organizational strategy within the tetraspanin family for

structuring specialized microdomains or organized polymers on cellular membranes.

Results

Tspan7 enrichment on retraction fibers

Previously, we have shown that Tspan7 is required for migrasome formation[12] and that

its overexpression can further promote migrasome biogenesis[29], although the

underlying mechanism remains elusive. Given that retraction fibers are essential for

migrasome formation, acting both as a direct membrane source of migrasomes and as

a linkage between migrasomes and cell body, we were prompt to explore Tspan7's

potential role in retraction fiber formation. By expressing Tspan7-GFP in normal rat

kidney (NRK) cells, we discovered that overexpression of Tspan7 notably boosts the

formation of retraction fibers (Figure 1A), with cells expressing Tspan7-GFP exhibiting

a nearly tenfold increase in fiber length relative to those expressing GFP alone (Figure

1B). Similar results were also observed in other cell lines (Figure S1A-S1D).

Furthermore, we found that Tspan7 enhances the formation of retraction fibers in a

dose-dependent manner. An increase in Tspan7 expression levels correlates with more

retraction fibers emanating from the cells (Figure S1E-S1G).

Notably, Tspan7-GFP also appeared to be enriched on retraction fibers relative to

the cell body. To analyze Tspan7-GFP localization further, we stained NRK cells

expressing Tspan7-GFP with Cellmask-Deepred, which stains the plasma membrane.

By capturing images at an optical plane proximal to the cell bottom, we were able to

visualize both retraction fibers and the plasma membrane on the basal side of the cell

(Figure 1C). Quantitative analysis of the Tspan7-GFP and Cellmask-Deepred signals

on these structures revealed that Tspan7-GFP showed much higher enrichment on

retraction fibers than on the basal membrane (Figure 1D), suggesting its preference for

localizing on tubular structures rather than planar membranes.

The observed preference of Tspan7 for cylindrical retraction fibers suggests its

potential sensitivity to high curvature. To directly test this hypothesis, we generated

Giant Plasma Membrane Vesicles (GPMVs) from Tspan7-GFP expressing cells. We

then employed optical tweezers to extrude a membrane tether from a GPMV. Briefly,

GPMVs were tagged with biotin-PE, and streptavidin-coated polystyrene beads were

manipulated via optical tweezers to make contact with the GPMV, resulting in a strong

adhesion between the streptavidin-coated bead and the biotin-PE-labeled GPMV.

Moving the bead away from the GPMV facilitated the extraction of a membrane tether

(Figure 1E). In this experimental setup, similar to *in vivo* observations, Tspan7-GFP

was found to be highly enriched on the highly curved membrane tether compared to the

GPMV, which exhibits lower curvature (Figure 1F-1I). In contrast, the Rho-PE signal

was confined to the GPMV, with no obvious Rho-PE present on the membrane tether

(Figure 1F, 1H). Furthermore, pannexin1, a four-pass transmembrane protein, was also

found to be exclusively localized on the GPMV (Figure 1G, 1I). Collectively, these

results suggest that Tspan7 has an intrinsic preference for highly curved surfaces.

Figure 1 Tspan7 senses high curvature and promotes the formation of retraction fibers

(A, B) NRK cells transiently transfected with GFP or Tspan7-GFP (T7-GFP) were cultured on fibronectin-coated glass for 16 hours and subsequently stained with WGA-561. Representative confocal images were shown in (A) and the area of retraction fibers per cell was quantified in (B). Data from three independent experiments (colored in yellow, blue or gray respectively) was pooled,

and mean (triangles) $\pm$ SD were plotted. N = 179 and 176 cells were analyzed for GFP or T7-GFP respectively. *P* values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). *P* value<0.05 was considered statistically significant. **** *P*<0.0001.

(C, D) NRK cells stably expressing Tspan7-GFP were cultured on fibronectin-coated glass for 16 hours and subsequently stained with Cellmask-Deepred. Representative confocal images were shown in (C, left). The ratio of T7-GFP/Cellmask was shown in fire color map (C, right) The ratio of fluorescent intensity of Cellmask or T7-GFP on retraction fibers versus cell body was analyzed in (D). Data from three independent experiments (colored in yellow, blue or gray respectively) was pooled, and mean (triangles) $\pm$ SD were plotted. In each experiment, N = 125 cells were analyzed. *P* values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). *P* value<0.05 was considered statistically significant. **** *P*<0.0001.

(E) Schematic illustration of GMPV pulling assay. First, GPMVs labeled with biotin-PE were settled in a glass-bottom confocal chamber and a polystyrene bead coated with streptavidin was trapped by an optical tweezer. Next, the trapped bead was allowed to approach and contact with a GPMV by manipulating the laser trap. A strong interaction between GMPV and bead was established due to the pairing of biotin and streptavidin. Finally, a membrane tether was created by pulling the trapped bead away from GPMV. Z-stack confocal images were acquired to record the signals of fluorescent proteins or lipids on GPMV and its corresponding membrane tether.

(F, G) Representative confocal images of GPMV with a membrane tether. GPMVs were either produced by NRK cells expressing T7-GFP and stained with rhodamine-PE (F) or by NRK cells expressing T7-GFP and Pannexin1-mCherry (G).

(H, I) Statistical analysis of the fluorescent intensity of T7-GFP versus rhodamine-PE (H) or pannexin1-mcherry (I) on GPMV or fiber. Data were pooled from three independent experiments, and mean (triangles) $\pm$ SD were plotted. N = 38 and 36 GPMVs were analyzed for (H) and (I) respectively. *P* values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). *P* value<0.05 was considered statistically significant. **** *P*<0.0001.

Scale bars, 5 μm.

***In vitro* reconstitution of Tspan7-induced membrane tethers resembling retraction fibers**

To investigate Tspan7's role in membrane tether formation and avoid interference from other cell components, we first purified recombinant Tspan7-GFP using a mammalian expression system. Subsequently, we reconstituted Giant Unilamellar Vesicles (GUVs) embedded with Tspan7-GFP *in vitro*. We discovered that when GUVs made contact with a hydrophilic surface, they flattened, forming a pancake-like double-membrane structure that became increasingly thin (Figure 2A). As this "pancake" expanded and thinned out, the upper membrane layer eventually ruptured at its center, causing the ruptured edges to retract to the pancake's periphery. This retraction left behind a network of membrane tethers that closely resemble retraction fibers. As the retraction process continued, vesicles began to form at the tethers' branch points. Over a very short period of 390 seconds, these vesicles intensified in brightness as the tethers seemingly dissolved into them (Figure 2B; Video S1), resembling the formation of migrasomes. Remarkably, in GUVs incorporating Tspan7-GFP, the rupture and retraction process led to an accumulation of Tspan7-GFP along the tethers, effectively stabilizing them and preventing their transformation into vesicles over a relatively long time (Figure 2C, 2D; Video S2). These observations confirmed that Tspan7 can detect high curvature and stabilize highly curved membrane tethers (Figure 2E).

Figure 2 Tspan7 senses and stabilizes tubular structures in an *in vitro* reconstitution system

(A) Schematic illustration of GUV-based *in vitro* reconstitution system. Firstly, liposomes w/wo T7-GFP were generated. GUVs w/wo T7-GFP were prepared from corresponding liposomes using electroformation method. Next, GUVs were transferred to plasma-treated hydrophilic glass surface and settled for 1.5 hour to allow attachment. Due to the strong adhesion force, the area of adhesion on attached GUVs automatically expanded, resulting in the avulsion of GUV. Finally, retraction

fiber and migrasome-like structures emerged from the spreading GUVs

(B) Snapshots from a timelapse video showing the spreading of a GUV without Tspan7-GFP. See

also Video S1.

(C) Snapshots from a timelapse video showing the spreading of a GUV with embedded Tspan7-

GFP.See also Video S2.

(D) Representative confocal images showing the morphology of reconstitution systems after setting

at room temperature overnight.

(E) Schematic illustration of the observed morphological transition of GUVs in the reconstitution

system.

Scale bars,10 μm, 5 μm in inserts.

Immobility of Tspan7-GFP on retraction fibers and reconstituted RF-like

membrane tethers

We next investigated whether other members of the tetraspanin family also exhibit

enrichment at retraction fibers. Through phylogenetic analysis[35,36], tetraspanins can be

divided into multiple subgroups (Figure 3A). We overexpressed five tetraspanins fused

with GFP in NRK cells, including Tspan7-GFP, its closest homologue Tspan6-GFP and

three representative members from other clusters including Tspan33-GFP, Tspan4-GFP,

and Tspan27-GFP. We found that all tested tetraspanins could localize to the retraction

fibers, although they showed varying preferences for retraction fibers over the flat basal

plasma membranes (Figure 3B). Among these tetraspanins, Tspan7 displayed the

strongest preference for retraction fibers, followed by Tspan6, which belongs to the

same subgroup as Tspan7. In contrast, other tetraspanins showed relatively minor

differences between their location at retraction fibers and the flat cell membranes

(Figure 3B, 3C).

Interestingly, we found that the Tspan7-GFP fluorescence on retraction fibers did

not recover after photobleaching (Figure 3D, 3E), suggesting that once Tspan7 is

concentrated on retraction fibers, it becomes immobilized. As the closest homologue to

Tspan7, Tspan6 exhibited partial recovery after photobleaching. In contrast,

tetraspanins belonging to other subgroups showed a rapid and nearly complete signal

recovery after photobleaching (Figure 3D, 3E). This distinctive behaviour of

Tspan7,which includes its unique immobility on retraction fibers, is intriguing for a

small membrane protein and deviates from what is typically expected. Subsequently,

we assessed the mobility of Tspan7-GFP on *in vitro* reconstituted membrane tethers.

Our findings revealed that Tspan7-GFP remained immobile (Figure 3F, 3G), similar to

in vivo observations. Given that the *in vitro* system consists solely of Tspan7-GFP and

lipids, this result suggests that the immobility of Tspan7 is an intrinsic property, not a

consequence of interactions with other proteins.

Figure 3 Immobility of Tspan7-GFP on retraction fibers and reconstituted RF-like membrane tethers

(A) Phylogenic tree of tetraspanin family generated by MEGA11. First, the protein sequences of 33 human tetraspanins were aligned using MUSCLE algorithm. Then, a phylogenic tree was constructed using Maximum Likelihood method. Bootstrap method with 1000 replications was used to test phylogeny. Frequencies with values higher than 70% were labeled at each branch. Five tetraspanins were highlighted in red: TSPAN7; its closest homologue TSPAN6 and representative

tetraspanins from three different clusters TSPAN33, TSPAN4 and TSPAN27.

(B-E) NRK cells were transiently transfected with Tspan7-GFP (T7-GFP), Tspan6-GFP (T6-GFP),

Tspan27-GFP (T27-GFP), Tspan4-GFP (T4-GFP) or Tspan33-GFP (T33-GFP) and cultured on

fibronectin-coated glass overnight. The fluorescent intensity of GFP was shown in fire color map in

(B) and the ratio of GFP intensity on retraction fiber (RF) versus cell was analyzed in (C). Data

from three independent experiments (colored in yellow, blue or gray respectively) was pooled, and

mean (triangles) $\pm$ SD were plotted. N = 90 cells were analyzed for each group. P values were

calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). P value<0.05 was

considered statistically significant. **** P<0.0001.

(D, E) FRAP assays were performed on transfected cells in order to analyze the mobility of different

tetraspanins on retraction fibers. Representative time-lapse images were shown in (D). The right

panels represent enlarged area from the left panels. The dashed circles represent areas of bleaching.

A 5-minute movie was acquired after bleaching. Representative images before or after bleaching

were shown. The fluorescent intensity in bleached area were tracked and plotted in (E). For each

curve, data from 15 bleaching areas were normalized and plotted as mean + SEM.

(F, G) FRAP assays were performed on $in\ vitro$ reconstituted fibers as described in Figure 2. Time

lapse images before and after bleaching were shown in (F). Dashed circles represent areas of

bleaching. The GFP intensity in bleached area were tracked and plotted in (G). Data from 9

bleaching areas were normalized and plotted as mean + SEM.

Scale bars, 5 μm in (B); 2 μm in (D) and (F).

Cryo-EM revealed the highly ordered spiral assembly of Tspan7 on retraction fibers

233 As a small transmembrane protein (28 kDa), Tspan7 is expected to diffuse

234 relatively freely on the membrane and recover quickly after photobleaching, similar to

235 other tetraspanins. But our observation is against this notion. One possible explanation

236 for its differing behavior is that Tspan7 forms higher-order oligomers on retraction

fibers. We then sought to explore the structural basis of Tspan7's immobility on retraction fibers. For this purpose, we performed *in-situ* cryo-EM analysis on NRK cells overexpressing either Tspan7-GFP or Tspan4-GFP, the latter serving as a control due to its high mobility on retraction fibers (Figure 3D, 3E). Briefly, NRK cells were seeded and cultured overnight on EM grids. We captured the fluorescent images of the cells to identify retraction fibers with strong green fluorescence signals. The EM grids were then rapidly frozen for cryo-EM imaging. Starting with a low-magnification cryo-EM micrograph, we correlated it with the fluorescent cell images to pinpoint the target retraction fibers. We then switched to high-magnification mode to capture detailed structural information of the retraction fibers (Figure 4A).

We firstly collected the cryo-EM micrographs of retraction fibers and performed single particle averaging analysis to sort relatively homogenous structures and enhance the signal-to-noise ratio of them (Figure 4B). We found that certain portion of retraction fibers overexpressing Tspan7-GFP exhibited a more uniform diameter of approximately 25 nm. Notably, we observed a regular row of dots, each about 5 nm in size, along the cytoplasmic side of these group of Tspan7-GFP retraction fibers (Figure 4B, bottom), a feature absent in those overexpressing Tspan4-GFP (Figure 4B, top). We then collected the tilt series of the retraction fibers in cryo-EM and reconstructed their cryo-eletron tomograms. Analysis of the tomograms confirmed the presence of the same regular row of dots along the inner side of Tspan7-GFP retraction fibers, a feature not observed in Tspan4-GFP retraction fibers (Figure 4C, 4D), consistent with the single-particle analysis.

We hypothesized that the observed dots corresponded to the GFP tags (27 kDa)

fused to the C-terminus, that is, the cytoplasmic side of the Tspan7 protein, as their size

matched well with that of GFP proteins. To test this, we replaced the GFP tag at the

cytoplasmic C-terminal of Tspan7 with a much smaller HA tag consisting of only nine

amino acids. This substitution caused the regular dot pattern on the cytoplasmic side to

disappear (Figure 4E, right). These results collectively confirm that the dot pattern is

linked to the GFP protein, serving as an indicator for the positioning and arrangement

of Tspan7, which displayed a highly ordered and densely packed configuration on the

retraction fibers.

Interestingly, cryo-EM tomography analysis revealed a spiral arrangement in the

Tspan7-GFP retraction fibers (Figure 4E, middle; Figure S2C, S2D; Video S4). In

contrast, the Tspan4-GFP retraction fibers, which are highly mobile, did not show spiral

structures (Figure 4E, left; Figure S2A, S2B; Video S3). This finding, along with

previous observations, indicates that Tspan7 concentrates and forms a highly ordered

spiral structure on retraction fibers. The similar spiral arrangement was also observed

on retraction fibers from cells overexpressing Tspan7-HA (Figure 4E, right; Video S5),

though these fibers were narrower compared to those expressing Tspan7-GFP,

suggesting that the diameter of Tspan7-containing fibers may be influenced by the tag

size. This result indicates that the formation of the spiral structure is an intrinsic

property of Tspan7, independent of its labeling constructs. Segmentation of the Tspan7-

GFP spiral structure revealed a right-handed six-start spiral with a diameter of 25 nm

(Figure 4F; Figure S3), exhibiting varying helical parameters.

Figure 4 Cryo-EM analysis of Tspan7 arrangement on retraction fibers

(A) Diagram of cryo-EM sample preparation and data collection. We first seed cells on golden EM

grids coated with graphene film (Step 1) and use a confocal fluorescent light microscope (fLM) to

take images of the cells (Step 2). After that, the cells cultured on EM grids would be blotted from

their backside and quickly frozen into liquid ethane to prepare the cryo-specimen. The atlas of EM grid was captured and cross-correlated with the fluorescent images in step 2, to locate the target retraction fibers (step 4). Tilt series or single-particle datasets were then acquired at the desired regions (step 5).

(B) Two-dimensional (2D) average results of particles picked from the cryo-EM micrographs of Tspan4-GFP (top) and Tspan7-GFP (bottom) retraction fibers.

(C) Representative central slices extracted from the tomogram reconstructions of Tspan4-GFP (top) and Tspan7-GFP (bottom) retraction fibers. The right panels showed the central slices with the membrane labeled in yellow and the dots in cyan.

(D) The intensity of the 9 line cross-sections across the Tspan4-GFP (top) and Tspan7-GFP (bottom) retraction fibers, plotted as a function of distance. The dark blue line was the averaged trace for all plots. The light blue line was the averaged trace after adding or subtracting the standard deviation. The distances separating the peak maxima were indicated for membrane densities (yellow line) and luminal dot densities (cyan line). a.u., arbitrary units.

(E) Representative bottom (B), middle (M), and top (T) slices of Tspan4-GFP (left), Tspan7-GFP (middle) or Tspan7-HA (right) retraction fiber tomograms. The yellow lines outlined the cell membrane in different slices, and the cyan markers indicated the regular dot density attached to the cytoplasmic side of the cell membrane. See also Video S3, S4, and S5.

(F) On the left, a typical density of a Tspan7-GFP retraction fiber segmented from the tomogram reconstruction. On the right, the density was identified as a right-handed six-start helix, with the six individual strands indicated by the six yellow lines.

Scale bars, 30 nm.

The formation of the Tspan7 spiral structure is negatively regulated by filamentous actin

We further examined the distribution and assembly of Tspan7 on retraction fibers at varying distances from the cell bodies, from proximal to distal (Figure 5A). In the

proximal region, where membrane curvature starts to increase, densely distributed GFP

dots were absent along the cytoplasmic side of the plasma membrane, and no spiral

arrangement was observed (Figure 5B, upper). Conversely, in the distal region, densely

distributed GFP dots along with a spiral arrangement were visible (Figure 5B, middle

and bottom), suggesting a preferential distribution of Tspan7 on the distal region. This

observation is consistent with our earlier findings that Tspan7 favors highly curved

membrane regions (Figure 1).

Filamentous actin (F-actin) is frequently observed in the proximal region of

retraction fibers[11]. To investigate the possible interplay between Tspan7 and F-actin

within these regions, cells expressing Tspan7-GFP were stained by SiR-actin, a

fluorescent probe for F-actin. We found that F-actin is predominantly located in the

proximal region of the retraction fibers, with occasional presence in isolated short

sections of the distal region (Figure 5C). Colocalization analysis of Tspan7-GFP and

SiR-actin signals on retraction fibers showed that these two components have a negative

correlation (Figure 5D). The repulsion between the distribution of F-actin and Tspan7-

GFP signals in the proximal region is further depicted by time-lapse analysis,

characterized by an increase in F-actin fluorescence coinciding with a decrease in

Tspan7-GFP fluorescence within the same area (Figure 5E; Video S6). In regions with

strong F-actin fluorescence, Tspan7-GFP fluorescence is often undetectable. When both

F-actin and Tspan7-GFP are present, their signals are generally weak. Photobleaching

experiments in areas with both signals suggest partial recovery of Tspan7-GFP

fluorescence, indicating some degree of diffusion of Tspan7-GFP in these regions

(Figure 5F, 5G). Collectively, these observations imply that the presence of F-actin in

retraction fibers may hinder Tspan7 recruitment and limit the formation of Tspan7 spiral

structures.

Next, we treated Tspan7-GFP expressing cells with Latrunculin A (LatA) to disrupt

F-actin assembly. LatA treatment did not affect the immobile state of Tspan7-GFP

(Figure 5H, 5I). Cryo-EM tomography analysis revealed a tightly packed Tspan7

configuration and spiral structure in LatA-treated retraction fibers (Figure 5J; Figure

S2E, S2F; Video S7). We then calculated the percentage of spiral formation in LatA-

treated Tspan7 retraction fibers and non-treated Tspan7 retraction fibers. This was

determined by the ratio of the number of retraction fibers exhibiting an obvious spiral

structure to the total number of retraction fibers. We found that LatA treatment

significantly increased the formation of Tspan7 spirals, from 24% to 88%, compared to

the untreated samples (Figure 5K). Furthermore, we also quantified the tightness of the

spiral structure using an angular velocity metric, defined as the ratio of the twist angle

to the rise of the spiral. This angular velocity metric also changed significantly, shifting

from 0.99 to 1.29 °/nm after LatA treatment (Figure 5L). This indicates that the Tspan7

spiral adopted a tighter form when the F-actin fiber was absent (Figure 5M). Together,

these data confirm that the presence of F-actin hinders the formation of Tspan7-GFP

spirals.

Figure 5 The interplay between Tspan7 and F-actin

(A) On the left, a representative low-magnification cryo-EM micrograph shows retraction fibers extruding from the cell body. On the right, the cell boundary is traced using white lines and the retraction fibers are indicated with green lines.

(B) The left panel displays the middle (M) and bottom (B) slices of the tomograms, corresponding to the three areas labeled as I-III in (A). On the right, F-actin is labeled in magenta, the cell membrane is labeled in yellow, and GFP is labeled in cyan.

(C) Representative confocal images of NRK cells stably expressing Tspan7-GFP, which had been stained with SiR-actin prior to imaging. In the segmentation image, cell body, total retraction fibers and SiR-actin-positive areas on retraction fibers were colored in deep blue, cyan and red respectively.

(D) The Pearson correlation coefficient of Tspan7-GFP and SiR-actin on retraction fibers was calculated based on the segmented images in (C). N=40 images were analyzed. Values between 0 and -0.3 indicate weak negative correlations.

(E) Snapshots from a timelapse video of a migrating cell. See also Video S6.

(F) Representative confocal images of NRK cells in order to analyze the mobility of Tepans7 on retraction fibers. The right panels represent enlarged area from the left panels (top: ROI 1, bottom: ROI 2). The dashed circles represent areas of bleaching. A 5-minute movie was acquired after bleaching. Representative images before or after bleaching were shown. ROI: Region Of Interest.

(G) The fluorescent intensity in bleached areas in (F) were tracked and plotted. For each curve, data from 15 bleaching areas were normalized and plotted as mean + SEM.

(H, I) FRAP assays were performed on NRK cells stably expressing tspan7-GFP. Cells were pretreated with 2 μM latrunculin A (LatA) or control solvent ethanol (NT). Confocal images of cells before FRAP assay were shown in (H), contour of the cells were highlighted in white line. The right panels represent enlarged area from the left panels. The dashed circles represent areas of bleaching. A 5-minute movie was acquired after bleaching. Representative images before or after bleaching were shown. The fluorescent intensities in bleached area were tracked and plotted in (I). For each curve, data from 15 bleaching areas were normalized and plotted as mean + SEM.

(J) The middle (M) and bottom (B) slices of the tomogram of tspan7-GFP retraction fibers after LatA treatment. See also Video S7.

(K) Statistical analysis on the formation of Tspan7 spiral after LatA treatment. N=75 or 51 tomograms of Tspan7-GFP retraction fibers that were either nontreated (NT) or treated with LatA were analyzed, respectively. Retraction fibers were categorized as either "non/weak spiral", indicating no spiral or a very slight spiral that was difficult to identify, or "strong spiral", indicating

a very obvious spiral structure on the retraction fibers.

(L) Plotting of the angular velocity of the Tspan7 spirals to assess their tightness. We analyzed N=52

randomly selected spiral structures (each around 30nm in length) from Tspan7-GFP retraction fibers

that were either nontreated (NT) or treated with LatA. The means (dark red line) ± SD were plotted.

P values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). A *P*

value <0.05 was considered statistically significant. ** *P*<0.01

(M) Model showcasing the assembly state of Tspan7 spirals on retraction fibers with or without the

existence of F-actin.

Scale bars, 1 μm in (A); 30 nm in (B); 5 μm in (C); 2 μm in (F) and (H).

Structural analysis of Tspan7 spiral on retraction fiber

Tspan7-GFP retraction fibers exhibited a more uniform diameter and a higher

density of GFP "dot" compared to Tspan7-HA retraction fibers (Figure 4E). As a result,

we chose Tspan7-GFP retraction fibers for subsequent cryo-EM subtomogram

averaging (STA) analysis to elucidate more detail information of the Tspan7 spiral

structure and identify critical interaction interfaces within the spiral.

The STA reconstruction of Tspan7-GFP retraction fibers revealed the general spiral

shape (Figure S3C). A higher resolution reconstruction was hindered by the

heterogeneity of the Tspan7-GFP spiral assembly in the retraction fibers (Figure 6B).

The resulting spiral density revealed a right-handed six-start helix with a twist of 3.14°

and a rise of 6.5 nm, with the GFP densities located in the center of the cylindrical

retraction fiber.

To pinpoint the potential key residues on Tspan7 that contribute to the formation

of spiral structure on retraction fibers, we utilized AlphaFold-Multimer[37,38] to predict

multiple hypothetical models of Tspan7 dimers (Figure 6A; Figure S4, S5). By

analyzing the interaction interfaces, we clustered these Tspan7 dimers into 5 distinct

classes and then selected a representative dimer model from each class (Figure S5) to

dock into the STA density. After correlating the models with the cryo-EM density map

of Tspan7 spiral, two of the dimer models, dimer 1 and dimer 2, exhibited the best

matches to the Tspan7 spiral structure. Following density-guided adjustment, we built

a putative model of how Tspan7 assembles into a spiral along the retraction fiber, using

the interaction interfaces observed in dimer 1 and dimer 2 (Figure 6B). In this model,

six strands of Tspan7 form a right-handed spiral, where the two interaction interfaces

between adjacent Tspan7 monomers are oriented in a head-to-head and back-to-back

arrangement, facilitating the organization of Tspan7 protomers within each strand.

To verify the above-built model of Tspan7 in the spiral, we performed a

comprehensive mutation analysis of the interaction interfaces to identify critical

residues in Tspan7 that contribute to its spiral arrangement and immobility (Figure 6C,

S7A). Our analysis pinpointed several key sites responsible for Tspan7's immobility,

including E115, aa190-199 (TVAATKVNQK) in the large extracellular loop, and aa53-

57 (ENSTN) in the small extracellular loop (Figure 6C). In cells expressing Tspan7-

GFP with mutations E115A, Δ53-57, or Δ190-199, immobility was partially lost. In

contrast, cells expressing the triple mutant Tspan7-GFP-E115A, Δ53-57, Δ190-199 (T7

3M) showed almost complete loss of immobility, suggesting that T7 3M cannot form

spirals. To further validate our findings, we performed in situ cryo-tomography analysis

on retraction fibers overexpressing T7 3M and found that the spiral structure was indeed

absent (Figure 6D; Video S8). Moreover, sequence alignment revealed that these

identified interaction sites are fully conserved in mouse and human Tspan7 (Figure

S6B). Among these sites, E115 is conserved in Tspan6 but not in Tspan4, aa53-57 and

aa190-199 are not conserved in Tspan6 or Tspan4 (Figure S6A). These results indicate

that the identified interaction interfaces are crucial for Tspan7's oligomerization and

spiral assembly on retraction fibers.

Finally, we compared the enrichment of Tspan7 WT-GFP and Tspan7 3M-GFP on

retraction fibers. Compared to Tspan7 WT-GFP, Tspan7 3M-GFP showed significantly

reduced enrichment on retraction fibers (Figure 6E, 6F). Moreover, in cells expressing

Tspan7 3M-GFP, the retraction fibers were significantly shorter than those in cells

expressing Tspan7 WT-GFP (Figure 6G, S7B). This suggests that spiral formation is

crucial for the enrichment of Tspan7 on retraction fibers and for maintaining their length.

Figure 6 Construction and verification of the molecular model of Tspan7 spiral

(A) The process of generating Tspan7 oligomerization models by combining the EM density map

and the AlphaFold-Multimer structural prediction.

(B) Subtomogram average of the Tspan7-GFP retraction fiber and the fitting of the helical model to

the map. The potential interaction surfaces involved in the oligomerization of Tspan7 (aa190-199,

aa53-57, and E115) were also highlighted.

(C) NRK cells were transiently transfected with wild-type Tspan7-GFP (T7 WT), wild-type Tspan4-

GFP (T4 WT), single-site mutant Tspan7-GFP (T7 E115A, T7 Δ53-57, T7 Δ190-199) and multiple-

sites mutant Tspan7-GFP (T7 3M with triple mutations E115A, Δ53-57 and Δ190-199). FRAP assays

were performed on transfected cells in order to analyze the mobility of WT or mutant tetraspanins on retraction fibers (See also Figure S7A.) For each curve, data from 15 bleaching areas were normalized and plotted as mean + SEM.

(D) Representative tomogram slices of the mutated Tspan7-GFP retraction fibers (T7 3M with triple mutations E115A, Δ53-57 and Δ190-199). See also Video S8.

(E, F) NRK cells were transiently transfected with wild-type Tspan7-GFP (T7 WT) or mutant Tspan7-GFP (T7 3M). The fluorescent intensity of GFP was shown in fire color map in (E) and the ratios of GFP intensity on retraction fiber (RF) versus cell were analyzed in (F). Data from three independent experiments (colored in yellow, blue or gray respectively) was pooled, and mean (triangles) ± SD were plotted. N = 94 and 92 cells were analyzed for WT and mutant respectively. *P* values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). *P* value<0.05 was considered statistically significant. **** *P*<0.0001.

(G) NRK cells transiently transfected with GFP, wild-type Tspan7-GFP (T7 WT) or mutant Tspan7-GFP (T7 3M) were cultured on fibronectin-coated glass for 16 hours and subsequently stained with WGA-561 (See also Figure S7B). The area of retraction fibers per cell was quantified. Data from three independent experiments (colored in yellow, blue or gray respectively) was pooled, and mean (triangles) ± SD were plotted. N = 167, 174 and 181 cells were analyzed for GFP, T7 WT and T7 3M respectively. *P* values were calculated using a two-tailed unpaired nonparametric test (Mann-Whitney test). *P* value<0.05 was considered statistically significant. **** *P*<0.0001.

(H) Working model of the transmembrane skeleton Tspan7. Tspan7 monomers sense high membrane curvature, forming polymers via homologous interaction on membrane tethers. When the cytoskeleton retracts, the Tspan7 polymers shape and stabilize the membrane tether through spiral assembly.

Scale bars, 30 nm in (D); 5 μm in (E).

Discussion

In this study, we have identified that Tspan7 is highly concentrated on retraction fibers and reconstituted membrane tethers. Our results revealed that Tspan7's selective

association with retraction fibers stems from its capacity to sense and concentrate on the highly curved membrane tethers. This accumulation is significantly amplified when Tspan7, typically a freely diffusing small transmembrane protein, organizes into long cylindrical polymers, adopting right-handed helical formations. We demonstrated that on retraction fiber, Tspan7 assembles into a tubular six-start spiral structure. Further, our findings indicate that these Tspan7 tubules markedly enhance the length of retraction fibers, and substantially improve their stability. Given the structural parallels between Tspan7 tubules and microtubules — both composed of polymerized basic protein units forming linear protofilaments and both providing structural support — we suggest that Tspan7 tubules act as a form of "transmembrane skeleton". This transmembrane skeleton, akin to the cytoskeleton, not only offers structural support but may also fulfill various other roles.

Membrane tethers, such as filopodia, retraction fibers, tunneling nanotubes, and microvilli, are prevalent features across various specialized cells and play critical roles in physiological processes. A shared characteristic of these protrusions is their exposure to constant mechanical stress, highlighting the potential benefit of a transmembrane skeleton that can stabilize these structures. Whether Tspan7 tubules can form and stabilize different types of membrane tether, or whether other proteins can create a transmembrane skeleton using similar organizational principles, remains to be determined.

Consistent with our previous finding, we found that Tspan7 overexpression not only elongates retraction fibers but also promotes migrasome formation. Earlier studies

suggested that the assembly of tetraspanin-enriched macrodomains triggers migrasome

formation. We observed that Tspan4, a non-polymerizing tetraspanin, integrates into

nanometer-scale tetraspanin microdomains and preferentially assembles in migrasomes

over retraction fibers[29]. This study, however, shows that Tspan7 has a contrasting

distribution, being more concentrated on retraction fibers. Considering the extensive

coverage of the Tspan7 transmembrane skeleton on retraction fibers, we propose that

Tspan7 facilitates migrasome formation by spatially restricting the movement of

tetraspanin-enriched microdomains, effectively channeling them towards migrasomes

and supporting their formation. Further research is required to test this hypothesis.

The absence of a functional Tspan7 antibody prevents us from directly testing the

formation of endogenous Tspan7 tubules. However, our data suggest that the assembly

of Tspan7-GFP tubules is not merely due to GFP's propensity for dimerization, as

replacing the GFP tag with an HA tag does not affect Tspan7's ability to form spiral

tubules, and mutating the potential interaction sites on Tspan7 abolishes the spiral

formation. Based on these findings, we think that Tspan7 tubules are likely a genuine

structure.

Except for a few cases, investigating cellular structures ranging from 10 nm to 100

nm has previously been challenging. Traditional structural biology tools are difficult to

implement *in vivo*, and optical imaging lacks the necessary resolution. Recently,

advancements in cryo-electron tomography (cryo-ET) have shown potential to change

this scenario. However, cryo-ET faces several challenges, including inherently low

signal-to-noise ratio and a limited field of view, which make searching for ordered

cellular structures without specific targets nearly impossible. In this study, we employed live imaging, FRAP (Fluorescence Recovery After Photobleaching) analysis, and *in vitro* reconstitution to investigate the properties of targeted proteins. We then used the insights gained (in this case, the immobility of a protein) to guide cryo-ET investigations, leading to the discovery of the Tspan7 transmembrane skeleton. We believe this integrated approach could be useful for discovering other ordered cellular structures.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (grant nos. 32030023, 92354306 and 32330025 to L.Y., 32130054 to H.-W.W., 32200981 to N.L.), Tsinghua-Toyota Joint Research Fund (grant no. 20233930058), Tsinghua University Dushi Program（grant no. 20241080003）and Beijing Municipal Science and Technology Commission, Administrative Commission of Zhongguancun Science Park (grant no. Z221100003422012) to L.Y.. We thank the State Key Laboratory of Membrane Biology and SLSTU-Nikon Biological Imaging Center for facility support in microscopic imaging and data processing. We thank Cryo-EM and High-Performance Computation platforms of Tsinghua University Branch of the National Protein Science Facility, and Shuimu BioSciences Ltd. (Beijing) for the support in cryo-EM data collection. We thank Mr. Xihang Chen (Molecular-Machines&Industries GmbH) for technical support in the optical tweezer system.

548 Sincere gratitude is expressed towards Dr. Sai Li and Xiaopeng Li for their kind

549 assistance on cryo-ET data processing.

Author contributions

L.Y. and H.-W.W. supervised the project and conceived the experiments. D.J.W.

conducted cell experiments, imaging, mutation analysis and statistics. D.J.W. and Y.Z.

conducted the *in vitro* reconstitution assays. D.J.W., X.J., Q.Z., N.L.and H-W.W.

conducted the cryo-EM analysis. D.J.W., X.J., N.L., and K.X. performed the model

building. L.Y., H-W.W., D.J.W., and N.L. wrote the manuscript with input from all

authors.

Declarification of interests

560 L.Y. is the scientific founder of Migrasome Therapeutics Ltd. All other authors declare

561 no competing interests related to this work.

Materials and Methods

Reagents and antibodies

Fibronectin was purchased from Invitrogen (PHE0023) or Sigma (F0895). WGA-TMR

(Wheat Germ Agglutinin tetramethylrhodamine conjugate, W849), CellMask Deep Red

(C10046) and Lipofectamin-3000 (L3000015) were purchased from Invitrogen. SiR-

Actin Kit (CY-SC001) was purchased from Cytoskeleton. Latrunculin A (10010630)

was purchased from Cayman. PEI MW40000 was purchased from Yeasen.

Lipids including DOPC, DOPE, POPS, cholesterol, 16:0 Biotinyl Cap PE and 18:1 Liss

Rhod PE were purchased from Avanti. Streptavidin (85878) was purchased from Sigma.

Polybead® Polystyrene Sampler Kit (19822-1) was purchased from Polysciences.

Anti-GFP (11814460001) was purchased from Roche. Anti-GAPDH (60004-1-Ig) was

purchased from Proteintech.

Plasmid construction

For transient transfection and imaging, the coding sequence of TSPAN7 and other

tetraspanins were amplified from rat cDNA through polymerase chain reaction (PCR)

and subsequently cloned into a pEGFP-N1 vector (linearized with XhoI and BamHI)

through homologous recombination. TSPAN7 constructs with point mutations were

generated using a standard site-directed mutagenesis protocol[39]. TSPAN7 constructs

with deletions were generated using a standard overlap extension PCR protocol[40].

For protein purification, the coding sequence of TSPAN7 was amplified from human

cDNA and subsequently cloned into a modified pCAG vector. Briefly, an affinity

cassette containing two repeats of Strep-Tag II followed by one FLAG tag was fused to

the N-terminal of TSPAN7 and an EGFP tag was fused to the C-terminal of TSPAN7

using the linker GGSGGS.

Cell culture and transfection

Adherent cells, including NRK, MGC-803 and MEF cells were cultured at 37 °C and

5% CO2 in DMEM high glucose medium supplemented with 10% fetal bovine serum,

1% penicillin–streptomycin and 1% Glutamax. 293F cells were cultured in suspension

at 37 °C, 5% CO2 and 120 rpm. in SMM293-TII supplemented with 1% penicillin–

streptomycin. Electroporation (Amaxa nucleofector, program X-001) was used for the

transfection of NRK cells. Lipofectamine-3000 transfection kit was used for the

transfection of MGC-803 and MEF. PEI was used for the transfection of 293F cells.

595　Retraction fiber reconstitution assay via tether pulling

NRK cells at 90% confluence were washed with GPMV buffer (10 mM HEPES, 2 mM

CaCl$_2$, 150 mM NaCl, pH 7.4) and incubated with GPMV working buffer (GPMV

buffer supplemented with 2 mM DTT and 25 mM formaldehyde before use) for 1 h at

37 °C. Supernatant containing GPMV was collected and incubated with 10 μg/ml

biotin-PE (and 0.25 μg/ml Rhodamine-PE in some cases) for 10 min at room

temperature. Polystyrene beads with 2 μm diameter were incubated with 100 μg/ml

streptavidin for 1 h. GPMVs were added to plasma treated glass surface and settled for

1 h before the addition of polystyrene beads. Optical tweezer experiments were

performed on a Nikon HD25 confocal microscope equipped with MMI CellManipulator.

A polystyrene bead was captured by a laser trap and was allowed to contact with a

GPMV. By pulling the bead at the opposite direction, a membrane tether was generated.

Z-stack images covering GPMV and membrane tether were collected using A1 confocal

microscope.

609　Protein expression and purification

293FT cells were grown to a density of 0.8-1.2 x 10^6 cell/ml and then transfected with

2 mg plasmid / L of cell culture using PEI (DNA : PEI = 1: 2.5). 3 days after transfection,

cells were harvested by centrifugation at 550 × g. The pellet was washed once with PBS

and resuspended in Tris buffer (100 mM Tris, 150 mM NaCl, pH 8.0) supplemented

with protease inhibitor. Cells were lysed by grinding 100 times using a glass Dounce

homogenizer on ice. The lysate was centrifuged at 30,000 × g for 1 hr and the pellet

was resuspended with 10 ml Tris buffer supplemented with protease inhibitor. To

extract Tspan7 protein from membrane, 2.5 ml of 10 % (w/v) Fos-Choline-12 was

added. The mixture was rotated at 4 °C for 2 hrs and then centrifuged at 12,000 × g for

20 min. Supernatant containing extracted protein was loaded to a column containing 1

ml pre-balanced strep-tactin beads and incubated for 1 hr at 4 °C on a rotator. The beads

were washed with 10 ml wash buffer (100 mM Tris, 150 mM NaCl, 0.1 % (w/v) Fos-

Choline-12, pH 8.0). Protein was eluted with 8 ml elution buffer (100 mM Tris, 150

mM NaCl, 0.1 % (w/v) Fos-Choline-12, 10 mM desthiobiotin, pH 8.0) and concentrated

using a 30 kDa MWCO centrifugal filter.

Liposome preparation

DOPC, DOPE, POPS and cholesterol were mixed at a ratio of 4:2.5:2.5:1, making up 3

mg lipids in total. 0.025 % rhodamine-PE was added as a fluorescent indicator. The

lipid mixture (prepared in a capped glass vial) was gently dried under a stream of

nitrogen and set for 1hr at 37 °C to ensure complete solvent evaporation. The resulting

lipid film was hydrated by adding 300 μl HEPES buffer (20 mM HEPES, 150 mM

NaCl, pH 7.4) and shaking vigorously at 37 °C for 1 hr. The mixture containing

liposomes was transferred into a centrifuge tube and freeze-thawed for at least 10 times

by transferring between liquid nitrogen and 42°C water bath. To homogenize liposomes,

a polycarbonate membrane with 1 μm pore size was inserted into an extruder and

liposomes were passed through for 21 times.

Proteoliposome preparation

Liposomes were kept at 4 °C on a rotator and protected from light during the whole

process. First, liposomes were incubated with 0.1 % Fos-Choline-12 for 30 min. Then,

purified Tspan7-GFP protein was added at a protein/lipid ratio of 0.01 (for protein-free

liposomes, equal amount of wash buffer (100 mM Tris, 150 mM NaCl, 0.1 % (w/v)

Fos-Choline-12, pH 8.0) was added). The mixture was incubated for 1 hr. To remove

the detergent, 10 – 12 mg of bio-beads / 100 μl liposome were added every hour for 3

times. After the third addition of bio-beads, the mixture was stirred overnight. A fourth

portion of bio-beads were added on the next day, followed by 1 hr incubation. The

supernatant was transferred into a clean centrifuge tube and store in aliquots at -80°C.

GUV preparation and reconstitution

GUVs were prepared via the electroformation method using Vesicle Prep-Pro (Nanion).

Before use, ITO (indium tin oxide)-covered glass slides were gently cleaned by liquid

soap, triple-washed using distilled water/ethanol and dried under a stream of nitrogen.

A rubber ring was stuck onto the ITO-covered side of the slide, creating a confined

chamber. 20 μl of proteoliposome was deposited onto the confined glass surface and

air-dried. The chamber was gently filled with 280 μl of 300 mM sorbitol and

electroformation was then carried out at 0.24 V, 9.9 Hz for 90 min at 37°C. The solution

containing GUVs was collected and stored at 4 °C before use.

For *in vitro* reconstitution, GUVs were added to a glass-bottom confocal chamber treated with plasma cleanser. The chamber was sealed by parafilm and settled for at least 1.5 h before imaging by a confocal microscope.

Imaging and staining

Cells were seeded on 10 μg/ml fibronectin coated confocal chamber (Cellvis) and transferred to a Nikon A1 HD25 confocal microscope equipped with a live cell imaging system. Images were acquired using a 100× objective at 1024 × 1024 pixels. For long-term time-lapse experiments, cells were allowed to migrate for about 6 hrs before imaging. Images were acquired at 2 min interval for indicated time periods. For confocal snapshot images and FRAP assays, cells were allowed to migrate for about 16 hrs before imaging. An additional staining step was performed before imaging in some experiments. Specifically, cells were incubated with 1 μg/ml WGA-TMR for 10min, 1.25 μg/ml CellMask Deep Red for 10 min or 1 μM SiR-actin for 1h before imaging.

FRAP assay and data processing

For each cell analyzed, 7 ROIs (Region Of Interest) were defined: 3 stimulation ROIs (1.6 μm diameter), 3 reference ROIs (1.6 μm diameter) and 1 background ROI. The experiment contains three phases: first, two frames were captured before stimulation; next, stimulation ROIs were bleached with 100% 488 nm laser for 1.5 s; finally, a 5 min time-lapse movie was recorded after bleaching. Data processing was performed using the NIS-Elements software. First, alignment was performed to correct XY motions. Then, the mean fluorescent signals of 7 ROIs over time were measured. Next, the fluorescent signals of stimulation ROIs were subjected to background correction and

reference correction. Finally, all datasets were linearly normalized to the [0, 100] range.

The largest value of each dataset was defined as 100% and 0 was defined as 0%.

Cryo-EM specimen preparation

Quantifoil R1.2/1.3 Au-grids (200 mesh) were coated with homemade graphene film[41]

and glow-discharged for 15 seconds (Harrick Plasma, low-power mode). Afterwards,

the EM grids were placed in a 35-mm Ibidi μ-Dish under ultraviolet illumination for 10

minutes. Next, human plasma fibronectin (FC010, Millipore, USA) solution (10 μg/ml

in PBS buffer) was dropped onto the EM grids and incubated for 40 minutes at 37 °C.

The fibronectin solution was then removed, and 1.2×10^5 NRK cells were seeded on

the EM grids and cultured overnight at 37°C and 5% CO_2. After a 14-16 hour culture,

we imaged the EM grids using a confocal light microscope with a 100× magnification

lens to obtain a grid atlas. The EM grids with cultured cells were then gently picked up

with tweezers, and 3 μL of nanogold particle solution (10 nm in diameter; Aurion, the

Netherlands) was added onto each grid. We then plunge-froze the grids using a Leica

EM GP. Grids were blotted from the back using the Leica EM GP at 37°C and 90%

humidity with a blotting time of 8-10 seconds. Finally, the grids were transferred into

liquid nitrogen for storage, ready for loading into the cryo transmission electron

microscope.

Single-particle cryo-EM data collection and analysis

The single-particle cryo-EM datasets for Tspan4-GFP and Tspan7-GFP retraction

fibers were collected using a Thermo Fisher Scientific Titan Krios (300 kV) TEM,

equipped with a K3 camera (Gatan, Inc.) and an energy filter. We collected 99

micrographs for Tspan4 and 103 micrographs for Tspan7 samples at a magnification of

18000×, with a defocus range of -1.5 to -2.0 μm. Each micrograph was dose-

fractionated into 32 frames with a cumulative dose of 50 e⁻/Å² and a pixel size of 1.56

Å. The micrographs were motion-corrected using MotionCor2[42] and then imported into

Relion[43] for data processing. We used CTFFIND4 to estimate the CTF parameters[44].

We manually picked 1502 particles for the Tspan4 samples and 1570 particles for the

Tspan7 samples for the subsequent two-dimensional classification, using identical

parameters.

Cryo-electron tomography data collection and reconstruction

Cryo-electron tomography (cryo-ET) tilt series of retraction fibers were collected on a

Titan Krios microscope operated at 300 kV (Thermo Fisher Scientific) equipped with

a GIF Quantum post-column energy filter (Gatan) and a K3 direct detector camera

(Gatan). Data were recorded in dose-fractionation mode using SerialEM software[45].

Tilt-series were collected using a dose symmetric scheme[46] in nanoprobe mode, with a

calibrated pixel size of 1.36 Å, a defocus range of -4.5 to -6.5 μm, a tilt increment of

3°, and an imaging dose of 3 e⁻/Å² for all tilts. The tilt range was from +60° to -60°

starting from 0°, resulting in a total dose of ∼123 e⁻/Å². We collected 12 tilt series for

the Tspan4-GFP retraction fibers, 77 tilt series for the Tspan7-GFP retraction fibers,

and 82 tilt series for the LatA-treated Tspan7-GFP retraction fibers, respectively.

The tilt series were imported to Warp software[47] for motion correction as well as CTF

estimation. The corrected tilt series were firstly aligned in IMOD[48] using a combination

of patch-tracking and gold fiducial markers for tracking, and then the final alignment

was performed using the linear interpolation option. Aligned images were binned by a

factor of 4 to a pixel size of 5.44 Å for sub-tomogram averaging.

For subtomogram averaging of Tspan7-GFP retraction fiber, the retraction fibers were

first traced in Dynamo[49] package. In the Dynamo model generation interface, we

manually selected the extremal points of straight segments along each retraction fiber

to define the central line ''backbone''. Bent regions of the retraction fibers were

discarded. Subsequently, we used the Dynamo filament model with torsion type for

generating subtomograms. The parameters *subunits_dz* and *subunits_dphi* were set to

12 pixels and 30°, repectively, to account for the helical rise of the retraction fibers and

to reduce the effect of the missing wedge in subtomogram averaging[50]. Subtomograms

(4 times binned, with a box size of 128^3 pixels) were cropped using Dynamo with the

model-defined positions, and initial Euler angles were generated relative to filament

axes with extra in-plane rotations derived from torsion angles.

Reference-free subtomogram averaging/refinement was performed using Dynamo.

Initial reference was generated by averaging subtomograms (4 times binned) from 3

retraction fibers without alignment. A cylindrical mask was generated in Dynamo

according to the initial average, and used as alignment mask. Other masks were left as

default. The resulting average of the 3 retraction fibers was used to align all 2208

subtomograms extracted from 29 retraction fibers.

Model building of Tspan7 spiral structure

We used the MMSeq2[51] to generate Multiple Sequence Alignments (MSAs) for the Tspan7 dimer. Following this, we conducted network predictions using AlphaFold-Multimer (version 2.3), employing the default 5-model parameter settings [38] and utilizing 17 distinct random seeds. This process ultimately yielded 85 predicted models of the Tspan7 dimer, which could be clustered into 5 classes based on their interaction interfaces.

To clearly illustrate the interaction modes of these dimer models, we selected one representative structure from each of the five classes of the Tspan7 dimer (dimers 1-5). Upon aligning one monomer of each dimer using the *matchmarker* function in ChimeraX, we found that dimer 1 and dimer 2 were arranged in a line, which is consistent with the feature of the TEM density map of Tspan7 forming a long chain. We then placed all these 5 dimers into the partitioned helical asymmetric unit density, aligning the "V" shape opening of Tspan7 with the orientation of the membrane curvature (Figure S4, S5). The correlation calculation was then performed in ChimeraX.

We used the interaction interfaces from dimer1 and dimer2 to build the Tspan7 spiral model, placing them into the long chains of Tspan7 spiral density map. Consequently, we maintained the interaction interfaces unchanged while manually adjusting the angle between the monomers to modify the distance between their transmembrane regions, thereby reducing the apparent clash. The spiral model was then generated using the spiral parameters (rise 65 Å, twist 3.14, 6-start), which were inferred from the

subtomogram averaging density map.

Supplemental information

Document S1. Figure S1-S7

Video S1. *In vitro* reconstitution using protein-free GUV, related to Figure 2B

Video S2. *In vitro* reconstitution using Tspan7-GFP-embedded GUV, related to Figure 2C

Video S3. Tomogram of retraction fiber from NRK cells expressing Tspan4-GFP, related to Figure 4E (left).

Video S4. Tomogram of retraction fiber from NRK cells expressing Tspan7-GFP, related to Figure 4E (middle).

Video S5. Tomogram of retraction fiber from NRK cells expressing Tspan7-HA, related to Figure 4E (right).

Video S6. The dynamics of SiR-actin and Tspan7-GFP on retraction fibers, related to Figure 5E

Video S7. Tomogram of retraction fiber from NRK cells expressing Tspan7-GFP and treated with 2 μM LatA, related to Figure 5J.

Video S8. Tomogram of retraction fiber from NRK cells expressing Tspan7 3M-GFP, related to Figure 6D.

References

1. Mill, P., Christensen, S.T., and Pedersen, L.B. (2023). Primary cilia as dynamic and diverse signalling hubs in development and disease. Nature Reviews Genetics *24*, 421-441. 10.1038/s41576-023-00587-9.

2. Mattila, P.K., and Lappalainen, P. (2008). Filopodia: molecular architecture and cellular functions. Nature Reviews Molecular Cell Biology *9*, 446-454. 10.1038/nrm2406.

3. Buszczak, M., Inaba, M., and Yamashita, Y.M. (2016). Signaling by Cellular Protrusions: Keeping the Conversation Private. Trends in Cell Biology *26*, 526-534. 10.1016/j.tcb.2016.03.003.

4. Jacquemet, G., Hamidi, H., and Ivaska, J. (2015). Filopodia in cell adhesion, 3D migration and cancer cell invasion. Current Opinion in Cell Biology *36*, 23-31. https://doi.org/10.1016/j.ceb.2015.06.007.

5. Wang, C., Xie, B., Yin, S., Cao, J., Huang, J., Jin, L., Du, G., Zhai, X., Zhang, R., Li, S., et al. (2023). Induction of filopodia formation by α-Actinin-2 via RelA with a feedforward activation loop promoting overt bone marrow metastasis of gastric cancer. Journal of Translational Medicine *21*, 399. 10.1186/s12967-023-04156-w.

6. Bischoff, M.C., Lieb, S., Renkawitz-Pohl, R., and Bogdan, S. (2021). Filopodia-based contact stimulation of cell migration drives tissue morphogenesis. Nature Communications *12*, 791. 10.1038/s41467-020-20362-2.

7. Zhang, C., Brunt, L., Ono, Y., Rogers, S., and Scholpp, S. (2024). Cytoneme-mediated transport of active Wnt5b–Ror2 complexes in zebrafish. Nature *625*, 126-133. 10.1038/s41586-023-06850-7.

8. Taylor, A.C., and Robbins, E. (1963). Observations on microextensions from the surface of isolated vertebrate cells. Developmental Biology *7*, 660-673. https://doi.org/10.1016/0012-1606(63)90150-7.

9. Lee, S.Y., Choi, S.H., Lee, M.S., Kurmashev, A., Lee, H.N., Ko, Y.G., Lee, K., Jeong, S., Seong, J., Kang, J.H., and Kim, H. (2021). Retraction fibers produced by fibronectin-integrin α5β1 interaction promote motility of brain tumor cells. Faseb j *35*, e21906. 10.1096/fj.202100452RR.

10. Porter, K.R., Claude, A., and Fullam, E.F. (1945). A STUDY OF TISSUE CULTURE CELLS BY ELECTRON MICROSCOPY : METHODS AND PRELIMINARY OBSERVATIONS. J Exp Med *81*, 233-246. 10.1084/jem.81.3.233.

11. Ma, L., Li, Y., Peng, J., Wu, D., Zhao, X., Cui, Y., Chen, L., Yan, X., Du, Y., and Yu, L. (2015). Discovery of the migrasome, an organelle mediating release of cytoplasmic contents during cell migration. Cell Res *25*, 24-38. 10.1038/cr.2014.135.

12. Jiang, D., Jiang, Z., Lu, D., Wang, X., Liang, H., Zhang, J., Meng, Y., Li, Y., Wu, D., Huang, Y., et al. (2019). Migrasomes provide regional cues for organ morphogenesis during zebrafish gastrulation. Nat Cell Biol *21*, 966-977. 10.1038/s41556-019-0358-6.

823   13.   Zhu, M., Zou, Q., Huang, R., Li, Y., Xing, X., Fang, J., Ma, L., Li, L., Yang, X.,
824   and Yu, L. (2021). Lateral transfer of mRNA and protein by migrasomes
825   modifies the recipient cells. Cell Res *31*, 237-240. 10.1038/s41422-020-00415-
826   3.

827   14.   Jiao, H., Jiang, D., Hu, X., Du, W., Ji, L., Yang, Y., Li, X., Sho, T., Wang, X., Li,
828   Y., et al. (2021). Mitocytosis, a migrasome-mediated mitochondrial quality-
829   control process. Cell *184*, 2896-2910 e2813. 10.1016/j.cell.2021.04.027.

830   15.   Zhang, C., Li, T., Yin, S., Gao, M., He, H., Li, Y., Jiang, D., Shi, M., Wang, J.,
831   and Yu, L. (2022). Monocytes deposit migrasomes to promote embryonic
832   angiogenesis. Nat Cell Biol *24*, 1726-1738. 10.1038/s41556-022-01026-3.

833   16.   Kim, J., Jo, H., Hong, H., Kim, M.H., Kim, J.M., Lee, J.-K., Heo, W.D., and
834   Kim, J. (2015). Actin remodelling factors control ciliogenesis by regulating
835   YAP/TAZ activity and vesicle trafficking. Nature Communications *6*, 6781.
836   10.1038/ncomms7781.

837   17.   Routledge, D., Rogers, S., Ono, Y., Brunt, L., Meniel, V., Tornillo, G., Ashktorab,
838   H., Phesse, T.J., and Scholpp, S. (2022). The scaffolding protein flot2 promotes
839   cytoneme-based transport of wnt3 in gastric cancer. eLife *11*, e77376.
840   10.7554/eLife.77376.

841   18.   Boucheix, C., and Rubinstein, E. (2001). Tetraspanins. Cellular and Molecular
842   Life Sciences CMLS *58*, 1189-1205. 10.1007/PL00000933.

843   19.   Hemler, M.E. (2005). Tetraspanin functions and associated microdomains.
844   Nature Reviews Molecular Cell Biology *6*, 801-811. 10.1038/nrm1736.

845   20.   Huang, Y., Zhang, X., Wang, H.-W., and Yu, L. (2022). Assembly of
846   Tetraspanin-enriched macrodomains contains membrane damage to facilitate
847   repair. Nature Cell Biology *24*, 825-832. 10.1038/s41556-022-00920-0.

848   21.   Zijlstra, A., Lewis, J., DeGryse, B., Stuhlmann, H., and Quigley, J.P. (2008).
849   The Inhibition of Tumor Cell Intravasation and Subsequent Metastasis via
850   Regulation of In Vivo Tumor Cell Motility by the Tetraspanin CD151. Cancer
851   Cell *13*, 221-234. 10.1016/j.ccr.2008.01.031.

852   22.   Lammerding, J., Kazarov, A.R., Huang, H., Lee, R.T., and Hemler, M.E. (2003).
853   Tetraspanin CD151 regulates α6β1 integrin adhesion strengthening.
854   Proceedings of the National Academy of Sciences *100*, 7616-7621.
855   10.1073/pnas.1337546100.

856   23.   Rubinstein, E., Le Naour, F., Lagaudrière-Gesbert, C., Billard, M., Conjeaud,
857   H., and Boucheix, C. (1996). CD9, CD63, CD81, and CD82 are components of
858   a surface tetraspan network connected to HLA-DR and VLA integrins.
859   European   Journal   of   Immunology   *26*,   2657-2665.
860   https://doi.org/10.1002/eji.1830261117.

861   24.   Menager, M.M., and Littman, D.R. (2016). Actin Dynamics Regulates Dendritic
862   Cell-Mediated Transfer of HIV-1 to T Cells. Cell *164*, 695-709.
863   10.1016/j.cell.2015.12.036.

864   25.   Runge, K.E., Evans, J.E., He, Z.-Y., Gupta, S., McDonald, K.L., Stahlberg, H.,
865   Primakoff, P., and Myles, D.G. (2007). Oocyte CD9 is enriched on the
866   microvillar membrane and required for normal microvillar shape and

distribution. Developmental Biology *304*, 317-325. https://doi.org/10.1016/j.ydbio.2006.12.041.

26. Bassani, S., Cingolani, L.A., Valnegri, P., Folci, A., Zapata, J., Gianfelice, A., Sala, C., Goda, Y., and Passafaro, M. (2012). The X-linked intellectual disability protein TSPAN7 regulates excitatory synapse development and AMPAR trafficking. Neuron *73*, 1143-1158. 10.1016/j.neuron.2012.01.021.

27. Blumenthal, A., Giebel, J., Ummanni, R., Schlüter, R., Endlich, K., and Endlich, N. (2012). Morphology and migration of podocytes are affected by CD151 levels. American Journal of Physiology-Renal Physiology *302*, F1265-F1277. 10.1152/ajprenal.00468.2011.

28. Dharan, R., Goren, S., Cheppali, S.K., Shendrik, P., Brand, G., Vaknin, A., Yu, L., Kozlov, M.M., and Sorkin, R. (2022). Transmembrane proteins tetraspanin 4 and CD9 sense membrane curvature. Proceedings of the National Academy of Sciences *119*, e2208993119. 10.1073/pnas.2208993119.

29. Huang, Y., Zucker, B., Zhang, S., Elias, S., Zhu, Y., Chen, H., Ding, T., Li, Y., Sun, Y., Lou, J., et al. (2019). Migrasome formation is mediated by assembly of micron-scale tetraspanin macrodomains. Nat Cell Biol *21*, 991-1002. 10.1038/s41556-019-0367-5.

30. Dharan, R., Huang, Y., Cheppali, S.K., Goren, S., Shendrik, P., Wang, W., Qiao, J., Kozlov, M.M., Yu, L., and Sorkin, R. (2023). Tetraspanin 4 stabilizes membrane swellings and facilitates their maturation into migrasomes. Nat Commun *14*, 1037. 10.1038/s41467-023-36596-9.

31. Yáñez-Mó, M., Barreiro, O., Gordon-Alonso, M., Sala-Valdés, M., and Sánchez-Madrid, F. (2009). Tetraspanin-enriched microdomains: a functional unit in cell plasma membranes. Trends in Cell Biology *19*, 434-446. https://doi.org/10.1016/j.tcb.2009.06.004.

32. Charrin, S., Manié, S., Thiele, C., Billard, M., Gerlier, D., Boucheix, C., and Rubinstein, E. (2003). A physical and functional link between cholesterol and tetraspanins. European Journal of Immunology *33*, 2479-2489. https://doi.org/10.1002/eji.200323884.

33. Wu, X.-R., Kong, X.-P., Pellicer, A., Kreibich, G., and Sun, T.-T. (2009). Uroplakins in urothelial biology, function, and disease. Kidney International *75*, 1153-1165. https://doi.org/10.1038/ki.2009.73.

34. Oostergetel, G.T., Keegstra, W., and Brisson, A. (2001). Structure of the major membrane protein complex from urinary bladder epithelial cells by cryo-electron crystallography1 1Edited by W. Baumeister. J Mol Biol *314*, 245-252. https://doi.org/10.1006/jmbi.2001.5128.

35. Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular biology and evolution *38*, 3022-3027. 10.1093/molbev/msab120.

36. Jones, D.T., Taylor, W.R., and Thornton, J.M. (1992). The rapid generation of mutation data matrices from protein sequences. Computer applications in the biosciences : CABIOS *8*, 275-282. 10.1093/bioinformatics/8.3.275.

37. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

911 Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
912 accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.
913 10.1038/s41586-021-03819-2.

38. Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek,
915 A., Bates, R., Blackwell, S., Yim, J., et al. (2021). Protein complex prediction
916 with AlphaFold-Multimer. bioRxiv, 2021.2010.2004.463034.
917 10.1101/2021.10.04.463034.

39. Zheng, L., Baumann U Fau - Reymond, J.-L., and Reymond, J.L. An efficient
919 one-step site-directed and site-saturation mutagenesis protocol.

40. Nelson, M.D., and Fitch, D.H.A. (2011). Overlap Extension PCR: An Efficient
921 Method for Transgene Construction. In Molecular Methods for Evolutionary
922 Genetics, V. Orgogozo, and M.V. Rockman, eds. (Humana Press), pp. 459-470.
923 10.1007/978-1-61779-228-1_27.

41. Zheng, L., Liu, N., Gao, X., Zhu, W., Liu, K., Wu, C., Yan, R., Zhang, J., Gao,
925 X., Yao, Y., et al. (2023). Uniform thin ice on ultraflat graphene for high-
926 resolution cryo-EM. Nature Methods *20*, 123-130. 10.1038/s41592-022-01693-
927 y.

42. Zheng, S.Q., Palovcak, E., Armache, J.-P., Verba, K.A., Cheng, Y., and Agard,
929 D.A. (2017). MotionCor2: anisotropic correction of beam-induced motion for
930 improved cryo-electron microscopy. Nature Methods *14*, 331-332.
931 10.1038/nmeth.4193.

43. Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E.,
933 and Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM
934 structure determination in RELION-3. Elife *7*. 10.7554/eLife.42166.

44. Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus
936 estimation from electron micrographs. J Struct Biol *192*, 216-221.
937 10.1016/j.jsb.2015.08.008.

45. Mastronarde, D.N. (2005). Automated electron microscope tomography using
939 robust prediction of specimen movements. Journal of Structural Biology *152*,
940 36-51. https://doi.org/10.1016/j.jsb.2005.07.007.

46. Hagen, W.J.H., Wan, W., and Briggs, J.A.G. (2017). Implementation of a cryo-
942 electron tomography tilt-scheme optimized for high resolution subtomogram
943 averaging. Journal of Structural Biology *197*, 191-198.
944 https://doi.org/10.1016/j.jsb.2016.06.007.

47. Tegunov, D., and Cramer, P. (2019). Real-time cryo-electron microscopy data
946 preprocessing with Warp. Nature Methods *16*, 1146-1152. 10.1038/s41592-019-
947 0580-y.

48. Kremer, J.R., Mastronarde, D.N., and McIntosh, J.R. (1996). Computer
949 Visualization of Three-Dimensional Image Data Using IMOD. Journal of
950 Structural Biology *116*, 71-76. https://doi.org/10.1006/jsbi.1996.0013.

49. Castaño-Díez, D., Kudryashev, M., Arheit, M., and Stahlberg, H. (2012).
952 Dynamo: A flexible, user-friendly development tool for subtomogram
953 averaging of cryo-EM data in high-performance computing environments.
954 Journal of Structural Biology *178*, 139-151.

955   https://doi.org/10.1016/j.jsb.2011.12.017.

956   50.   Wan, W., and Briggs, J.A.G. (2016). Chapter Thirteen - Cryo-Electron Tomography and Subtomogram Averaging. In Methods in Enzymology, R.A. Crowther, ed. (Academic Press), pp. 329-367. https://doi.org/10.1016/bs.mie.2016.04.014.

960   51.   Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nature Biotechnology *35*, 1026-1028. 10.1038/nbt.3988.