Melanization interacts with soil mineral and microbial properties to determine fungal carbon and

nitrogen persistence in soils

Katilyn V. Beidler[1,2*], Elizabeth Huenupi [1], Lang C. DeLancey[3], François Maillard[4], Bowen Zhang[5], Per

Persson[4,5], Peter G. Kennedy[2] & Richard Phillips[1]

1. Department of Biology, Indiana University, Bloomington, IN, 47405 , United States of America

2. Department of Plant and Microbial Biology, University of Minnesota, St. Paul, MN, 55108,

United States of America

3. Ecology, Evolution, and Behavior Graduate Program, University of Minnesota, St. Paul, MN,

55108, United States of America

4. Department of Biology, Microbial Ecology Group, Lund University, Sölvegatan 35, SE-22362,

Lund, Sweden

5. Centre for Environmental and Climate Science, Faculty of Science, Lund University, Ecology

Building, Sölvegatan 37, SE-22362, Lund, Sweden

*Corresponding Author

Katilyn V. Beidler

beidl006@umn.edu

1030 Manvel Street,

Saint Paul MN 55114

843-830-4679

Abstract

Despite the importance of mineral-associated organic matter (MAOM) in long-term soil carbon (C) and nitrogen (N) persistence, and the significant contribution of fungal necromass to this pool, the factors controlling the formation of fungal-derived MAOM remain unclear. This study investigated how fungal necromass chemistry, specifically melanin, interacts with soil mineral properties and microbial communities to influence MAOM formation and persistence. We cultured the fungus *Hyaloscypha bicolor* to produce ^{13}C- and $^1\square$N-labeled necromass with varying melanin content (high or low) and incubated it in both live and sterile soils collected from six Indiana forests that differed in their clay and iron oxide (FeOx) content. After 38 days, we found that seven times more fungal-derived N was incorporated into MAOM than fungal-derived C, with fungal N comprising 20% of the MAOM-N pool. Low melanin necromass formed more MAOM-C than high melanin necromass, although site-level differences in overall MAOM formation were substantial. Soil clay and FeOx content were strong predictors of MAOM formation, explaining ~60% and ~68% of the variation in MAOM-C and MAOM-N, respectively. However, microbial communities significantly influenced MAOM formation, with MAOM-C formation enhanced and MAOM-N formation reduced in sterile soils. Furthermore, the relative abundance of fungal saprotrophs was negatively correlated, and bacterial richness was positively correlated with MAOM formation, and these relationships were influenced by necromass melanin content. This study reveals that microbial communities and soil properties interactively mediate the incorporation of fungal necromass C and N into MAOM, with microbes differentially influencing C and N incorporation, and these processes being further modulated by necromass melanization.

Keywords: Fungal necromass, Melanin, Mineral-associated soil organic matter, Iron oxide

    1. Introduction

To reach sparse soil resources, it is estimated that fungi produce an average of one kilometer of hyphae

per cubic centimeter of soil (See et al., 2022). High hyphal densities combined with rapid turnover make

fungi major, if not the main microbial contributors to soil organic carbon (SOC) and nitrogen (SON;

Godbold et al., 2006; Wang et al., 2021). An estimated 30-50% of SOC and 60-80% of SON is microbial

in origin (Simpson et al., 2007; Hu et al., 2020; Wang et al., 2021; Warren, 2021), with dead fungal

hyphae (hereafter fungal necromass) forming up to two-thirds of all microbial necromass (Liang et al.,

2019; Wang et al., 2021). While the accumulation of organic soil C and N occurs gradually over years to

centuries, microbial compounds can be stabilized within the soil matrix much more rapidly (days to

months) and this rapid stabilization is facilitated by the small size and charged nature of microbial

compounds, which readily bind to the surfaces of soil clays and metal oxides (Lützow et al., 2006; Kleber

et al., 2007; Cotrufo et al., 2013; Lehmann and Kleber, 2015; Cotrufo and Lavallee, 2022). These

mineral-organic associations may persist over hundreds of years, as they are biologically, chemically, and

physically protected from further decay

(Sollins et al., 1996; Kögel-Knabner, 2002; Kleber et al., 2015). Given the extensive presence of fungal

hyphae in soils, their close proximity to soil minerals, and the unique chemical composition of fungal

cells

(Kögel-Knabner, 2002; Godbold et al., 2006; See et al., 2022), it is important to identify the dominant

ecological factors controlling fungal-derived SOM formation and persistence.

The soil mineral matrix or physicochemical context in which fungal hyphae live and die strongly

influences the accumulation and persistence of SOM (Lützow et al., 2006; Frey, 2019). The composition

of this matrix, including the abundance of clay minerals and metal oxides, plays a crucial role in

protecting organic matter from further decomposition (Sollins et al., 1996; Kleber et al., 2007). In similar

climates, soil clay content correlates positively with SOC (Matus, 2021). This relationship is reflected in

the frequent use of percent clay to estimate SOM content in the field and as a proxy variable in terrestrial

C cycling models (Sulman et al., 2014; Bailey et al., 2018). Together with clay-sized particles, amorphous

metal oxides provide the greatest surface area on which organic matter (OM) can bind (Lützow et al.,

2006). Iron (hydro)oxides (FeOx) have high densities of reactive hydroxyl (-OH) sites and thus increased

capacity for ligand exchange or -OH exchange between carboxyl or phenolic functional groups in OM

and -OH functionalities on mineral surfaces (Harsh et al., 2018; Wu et al., 2023). Ligand exchange results

in stronger bonds compared to OM-cation-mineral interactions (i.e., polyvalent cation bridges) and thus

has increased capacity to stabilize OM

(Lützow et al., 2006; Mikutta et al., 2007). However, the charge and reactivity of mineral constituents

with OM are ultimately dependent upon soil pH, which is considered a "master soil variable" for soil

chemistry and microbial activity (Deng and Dixon, 2018; Rasmussen et al., 2018; Wang and Kuzyakov,

2024). Furthermore, soil pH and mineralogy interact to have indirect effects on SOM dynamics by

shaping microbial decomposer communities through the availability of oxygen, water, and nutrients in

soil pores

(Fierer and Schimel, 2002; Kallenbach et al., 2016).

Fungal necromass in forest soil supports a distinct microbial community whose composition and activity

are influenced by necromass initial chemistry (Maillard et al., 2020; Cantoran et al., 2023; Maillard et al.,

2023b). These microbial decomposers, in turn, can shape the chemical makeup of the resulting SOM

(Domeignoz-Horta et al., 2021). In early decomposition, fungal necromass is dominated by saprotrophic

fungi (i.e., fast growing moulds and yeasts) and copiotrophic bacteria (Brabcová et al., 2016; Beidler et

al., 2020; Kennedy and Maillard, 2023) that can capitalize on an abundance of labile, C-, and N-rich

compounds (See et al., 2021). Fast-growing microbial taxa may have lower carbon use efficiency (CUE),

meaning they respire more C than they convert into biomass (Manzoni et al., 2018), which may limit the

persistence of necromass-derived C (Štursová et al., 2012; Adingo et al., 2021). Maillard et al. (2023a)

found that relative to other edaphic factors (e.g., soil pH, moisture and temperature), initial soil bacterial

richness and fungal community composition were the strongest predictors of fungal necromass decay. It

has also been shown that increased melanin content slows the uptake of fungal necromass C and N by

both bacterial and fungal communities during decomposition (Maillard et al., 2023a). Melanins are high

molecular weight, hydrophobic polyphenols that often complex with proteins and carbohydrates (Bell and

Wheeler, 1986; Butler and Day, 1998; Rillig et al., 2007), potentially limiting C and N release from

fungal necromass. However, melanin is also known to be rich in carboxyl and phenolic hydroxyl groups

(Fogarty and Tobin, 1996; Harki et al., 1997; Fu et al., 2022), which interact readily with charged

surfaces on clay minerals. Additionally, melanin contributes to metal binding and sorption in fungal cell

walls (Fomina and Gadd 2003), suggesting that melanized hyphae may have a high sorption affinity for

metal oxides in soils (Butler and Day, 1998). Given that melanized hyphae can constitute over 50% of

total hyphal mass in some forests (Van Der Wal et al., 2009; Clemmensen et al., 2013; Siletti et al.,

2017), it's important to understand how melanization influences the formation of mineral-associated

fungal C and N.

The molecular composition of fungal cells reflects the varied ecology of individual fungal species and

results in a complex mixture of detrital proteins, polysaccharides, and phenols post-senescence (See et al.,

2021). The two-pool nature of fungal necromass decay suggests the existence of a larger labile or 'fast'

decaying fraction (~80% of initial mass) and a smaller recalcitrant or 'slow' decaying fraction (~20% of

initial mass; Fernandez et al., 2019; Ryan et al., 2020; See et al., 2021). Labile substrates form dissolved

organic matter (DOM) that can be assimilated by microbes and converted into residues and biomass (i.e.,

the microbial synthesis pathway of SOM formation), which may then become "entombed" via mineral-

organic interactions (Cotrufo et al., 2015; Liang et al., 2017). Alternatively, DOM may bind directly to

soil minerals (i.e., the direct sorption pathway of SOM formation; Kalbitz et al., 2005; Cotrufo et al.,

2015), with the strength of mineral bonding being higher for polar compounds (e.g., carboxyl and

phenolic acids) compared with compounds that bind through weaker hydrogen bonds or van der Waals

forces (e.g., simple sugars; (Jagadamma et al., 2012; Sokol and Bradford, 2019). In general, peptidic

compounds have a strong affinity for a wide array of mineral and organic surfaces in soils and make up

the largest store of soil organic N (Rillig et al., 2007; Norén et al., 2008; Theng, 2024). Previous studies

have used isotopically-enriched fungal necromass to demonstrate that fungal biomass can be a significant

source of SOM in forests (Schweigert et al., 2015), that environmental factors and microbial communities

influence necromass recycling efficiency (Buckeridge et al., 2022), and that minerals play a key role in

necromass C and N stabilization (Wang et al., 2024). Building on this previous work, our study

investigates how melanization, in soils differing in initial mineral and microbial properties, might

promote fungal C and N persistence, either together or independently.

To gain insights into how both soil mineral and microbial properties interact to determine fungal C and N

persistence, we performed two experiments utilizing necromass of high and low melanin phenotypes of

Hyaloscypha bicolor (previously *Meliniomyces bicolor*). *H. bicolor* is a widely distributed ascomycete

fungus capable of forming both ectomycorrhizal and ericoid mycorrhizal associations with plants (Grelet

et al., 2009; Fehrer et al., 2019). Its ability to produce varying levels of melanin influences both its cell

wall chemistry and decay rates (Fernandez et al., 2019; Ryan et al., 2020), making it a valuable organism

for studying the impact of necromass chemistry on microbial decomposition processes independent of

species differences. In the first experiment, we decayed dual enriched (^{13}C and ^{15}N) low and high melanin

necromass types in either live or sterilized soils differing in initial microbial and physicochemical

properties. Specifically, we utilized six temperate forest soils representing a gradient of increasing iron-

oxide (FeOx) content collected from various sites in Indiana, USA (Table 1). We predicted that higher

FeOx and clay content would promote the retention of fungal-derived C and N in mineral-associated

organic matter (MAOM). Beyond initial soil mineral properties, we were interested in the potential for

initial microbial community properties to predict the transformation of fungal necromass into MAOM.

We hypothesized that initial soil bacterial richness and relative abundances of fungal saprotrophs would

influence necromass-derived MAOM formation and chemistry, but likely to a lesser extent than soil

mineral properties. We predicted that MAOM formation would be higher in sterile than live soils due to

direct adsorption of soluble necromass compounds in the absence of microbial decomposers. Further, we

hypothesized that the chemical classes available for adsorption onto iron oxide minerals would differ

between necromass types. To test this hypothesis, we performed a second experiment to identify

compounds from low and high melanin that adsorbed directly onto synthesized goethite, an iron oxide

mineral ubiquitous in soils, using simultaneous infrared and potentiometric titration (SIPT). Collectively,

these two experiments allowed us to assess the relative amounts of fungal necromass-derived C and N as

well as the necromass chemical constituents that are converted to MAOM, which may indicate

persistence in soil. Given global efforts to increase soil C (Minasny et al., 2017), we require a more

mechanistic understanding of how microbial necromass is converted into MAOM and the factors that

predict its persistence.

Table 1. Site locations and initial physicochemical properties of six different soil types chosen to

represent contrasting soil iron oxide (FeOx), clay, carbon (C), and nitrogen (N) content, C to N ratio and

pH.

Site ID	Soil Taxonomic Class	FeOx (%)	Clay (%)	C (%)	N (%)	C:N	pH
C153	Fine-silty, mixed, active, acid, mesic Fluventic Endoaquepts	1.28	28.9	9.0	0.23	39	7.06
C55	Fine-silty, mixed, active, mesic Oxyaquic Fragiudalfs	0.52	14.9	1.8	0.15	12	3.98
C169	Fine, illitic, mesic Aeric Epiaqualfs	0.42	19.3	4.6	0.28	16	6.34
C71	Loamy-skeletal, mixed, active, mesic Typic Dystrudepts	0.25	11.9	1.6	0.14	11	4.46
C13	Loamy-skeletal, mixed, active, mesic Typic Hapludults	0.18	12.2	1.3	0.10	13	3.94
C61	Fine, mixed, active, mesic Typic Hapludalfs	0.08	12.4	0.67	0.04	17	4.15

174   <u>2 Methods</u>

2.1 Fungal Necromass Preparation

2.1.1 Necromass Production

We generated dual ^{13}C and ^{15}N enriched fungal necromass ex-situ from *Hyaloscypha bicolor*. Depending on culture oxygenation, *H. bicolor* will produce both high and low melanin phenotypes and therefore has been utilized in a number of field and laboratory incubation studies to test the effect of melanization on necromass decay while keeping necromass species constant (Fernandez and Kennedy, 2018; Ryan et al., 2020; Maillard et al., 2023b; Novak et al., 2024). Mycelial plugs (3 mm diameter) taken from modified Melin-Norkans (MMN) agar plates were transferred into either 125- or 250-ml Erlenmeyer flasks containing 100 ml of liquid MMN media (pH = 5). Different sized Erlenmeyer flasks were used to vary liquid levels without changing media volume, as it has been demonstrated that increasing the submersion level of *H. bicolor* reduces cell wall melanization without altering fungal mass or growth rates (Fernandez and Kennedy, 2018). The sole carbon source in the media was D-Glucose-**^{13}C**$\square$, (99 atom % ^{13}C; Cambridge Isotope Laboratories) together with D-Glucose-^{12}C$_6$, for a targeted enrichment of ~1 atom % or ~37 ‰ (see Table S1 for initial necromass isotopic enrichment values). As an N source, we added ^{15}N ammonium chloride (99 atom % ^{15}N; Cambridge Isotope Laboratories) in combination with ^{14}N ammonium chloride for a targeted enrichment of ~2 atom % or ~700 ‰. We aimed for higher enrichment of N because of its disproportionally smaller mass relative to the C contained in necromass (Table S1) and enrichment targets were set according to the highest isotopic standards available through the Purdue Stable Isotope (PSI) Facility, West Lafayette, IN, USA.

Flasks were shaken at 80 rpm in the dark for 30 days or until colonies stopped growing. During harvest, fungal biomass was rinsed with distilled water (di-H$_2$O) over 0.5 mm polyester mesh and any remaining agar from the mycelial plugs was carefully removed with forceps. The rinsed biomass was then dried at 25ºC for 48 hours, which was shown to be sufficient for killing the fungus, while limiting chemical

transformation. To homogenize the resulting necromass, dried fungal biomass was lightly ground using a mortar and pestle and passed through a 2 mm and 0.125 mm sieve stack; the cumulative mass retained on the 0.125 mm sieve was considered fungal necromass. Fungal necromass was subsampled for initial chemistry measurements and stored in a desiccator until the start of the incubation.

2.1.2 Initial Necromass Chemistry

Necromass subsamples were ground to a fine powder using a GenoGrinder (SPEX® SamplePrep; 1600 rpm for 1 minute) and analyzed for total C and N, as well as $\delta^{13}C$ and $\delta^{15}N$ (three technical replicates per necromass type) using an elemental analyzer coupled to a gas-isotope ratio mass spectrometer (PDZ Europa Elemental Analyzer coupled to a Sercon 20-22 IRMS) at the PSI facility. To approximate the proportion of initial C quality fractions (e.g., cell-soluble versus acid unhydrolyzable fractions) we followed forage fiber analysis procedures as described by ANKOM Technology (Macedon, NY, USA) and used in previous fungal necromass studies (See et al., 2021; Maillard et al., 2023b). This method separates necromass into fractions based on their solubility in neutral detergent and resistance to acid hydrolysis. We weighed necromass samples before and after sequential extractions to calculate the proportion in each fraction. First, to determine the mass of the cell-soluble fraction, which includes simple carbohydrates, lipids, and soluble proteins, we quantified the portion of necromass lost after constant agitation in a neutral detergent solution at 100ºC for two hours. We then dried and weighed the remaining necromass, attributing the mass difference to the cell-soluble fraction. Next, to determine the mass of the acid unhydrolyzable fraction, which consists of aromatic compounds including melanin, residual necromass was then extracted with a strong acid (72% H_2SO_4) at room temperature for three hours with repeated agitation (every 30 minutes; (Maillard et al., 2023b). We then dried and weighed the remaining residue, attributing the mass difference to the acid unhydrolyzable fraction.

To identify potential differences in chemical bonds and functional groups in a non-destructive manner, we used Diffuse Reflectance Infrared Fourier transform spectroscopy (DRIFTS). Briefly, initial necromass

samples (one per necromass type) were mixed with KBr (1% sample and 99% KBr) and analyzed on a Bruker Vertex 80 v FTIR spectrometer (Bruker Vertex 80 v, Ettlingen, Germany) in a temperature-controlled room at 21ºC. Sixty-four scans were averaged across the 4000–400 cm^{-1} range at a resolution of 4 cm^{-1}. Background subtraction was performed using a pure KBr spectrum, and a baseline correction was applied to eliminate baseline distortions. We carried out both background subtraction and baseline correction in OPUS software (Bruker, Ettlingen, Germany) and normalized peak heights by calculating z-scores. Peaks were identified based on literature references for fungi (Niemeyer et al., 1992; Cocozza et al., 2003; Maillard et al., 2023c).

2.2 Soil Microcosm Preparation

2.2.1 Soil collection

Soils were taken from hardwood forest plots at sites established by the Indiana Department of Natural Resources (IDNR) as part of the Continuous Forest Inventory (CFI) program (See Table 1 and the accompanying map). We selected soils from six CFI plot sites (diameter = 14.6 m) sampled in the summer of 2020 (Late May-Late July). Soil samples were collected 7.62 m from the plot center at 90° and 270° azimuth and a depth of 10–20 cm (5 cm core diameter). The two soil cores from a given plot were then homogenized over a 2 mm sieve and air dried. See Table S2 for specific site soil series and overstory information.

2.2.2 Initial soil physicochemical measurements

To measure soil pH, a subsample of air-dried soil (5.0 g) was shaken at 180 rpm in 40 mL of 0.01M CaCl$_2$ for 30 minutes and vortexed prior to taking readings with a benchtop SymPHony Benchtop pH Meter (VWR International, Radnor, PA, USA). The percentage of clay within soils was measured using a standard hydrometer procedure (Gavlak et al., 2005). We dispersed approximately 40 g of air-dried soil in a 5% (w/v) sodium hexametaphosphate solution and placed on a reciprocal shaker (120 rpm) for 16 hours.

This suspension was transferred to a 1000 mL graduated cylinder and brought to volume with deionized

water (di-H_2O). A hydrometer was used to measure the specific gravity of the suspension at 40 seconds

and 6 hours to determine the proportions of sand, silt, and clay, respectively. To estimate noncrystalline

and poorly crystalline Fe-oxides (FeOx) content in initial soils, we measured oxalate-extractable Fe

content (Ross and Wang, 1993; Craig et al., 2022). In short, 40 mL of 0.2 M $(NH_4)_2C_2O_4$ (pH = 3) was

added to 0.2 g (dwt equivalent) of soil. After shaking in the dark for 4 h, we gravity filtered the samples

(Whatman No. 40 filter paper) and stored the extracts at 4°C until analysis (within 4 d). Iron in the extract

was determined using atomic-adsorption spectrometry; Fe was atomized using an Aanalyst 800 air-

acetylene flame (Perkin Elmer, Waltham, MA, USA).

2.2.3 Initial soil microbial community measurements

Air-dried soils were subsampled to identify fungal and bacterial communities in initial soils (three

technical replicates per site or 18 samples total), we applied a two-step high throughput sequencing (HTS)

DNA metabarcoding approach. Total genomic DNA was extracted from soils using a Qiagen DNEasy

PowerSoil Kit (Qiagen, Benelux BV) according to the manufacturer's instructions. We first PCR

amplified extracted DNA from each sample, targeting the ITS2 region of the fungal rRNA operon (5.8S-

Fun and ITS4-Fun primer pair; (Taylor et al., 2016) and the V4 region of the bacterial 16S rRNA gene

(515F-806R primer pair; (Caporaso et al., 2012), respectively. In a second, shorter PCR, we added unique

Golay barcodes Golay barcodes and sequencing adaptors were added to samples with amplicons. Further

details on PCR reagents and conditions are available in (Maillard et al., 2023a). We quantified amplified

products using a Qubit dsDNA HS Fluorometer (Life Technologies, Carlsbad, CA, USA) and cleaned

using a Charm Just-a-Plate Purification and Normalization Kit (Charm Biotech, San Diego, CA, USA).

We included negative controls, consisting of a lysis tube without soil during DNA extraction and PCR

blanks (molecular grade water used in place of DNA template), as well as positive controls consisting of

fungal synmock (Palmer et al., 2018) or bacterial zymock (Zymo Research, Irvine, CA,USA)

communities. Each of the samples was pooled into a single library and sequenced at the University of

Minnesota Genomics Center using the 2 x 300 base pair (bp) paired-end MiSeq Illumina platform.

2.3 Fungal Necromass Incubation

2.3.1 Live soil conditions

Incubations were carried out in 50 ml centrifuge tubes (Chang Bioscience, Inc., Fremont, CA, USA) with

modified caps, which included a 0.2 µm vent filter to allow for air exchange and central needle (14 gauge

1 ½'' needle) for sterile water additions (needle was capped when not in use; See Fig. S1). We added 5.0

g (dwt equivalent) of soil to each microcosm and one week prior to the start of the incubation soils were

brought up to 65% water holding capacity (WHC). Three soil microcosms were prepared for each soil-

necromass combination & soil only controls (54 microcosms total). To begin the incubation, 70 mg of

necromass was added to microcosms and soils were mixed using a dissecting needle. To maintain

consistency among treatments we mixed soil only controls in a similar manner as microcosms receiving

necromass additions. Throughout the duration of the 38-day incubation microcosms were kept in the dark

at 20ºC and soils were held at 65% WHC via weekly additions of sterile di-H_2O in a laminar flow hood.

To stop incubations, soils were dried at 60ºC for 4 days.

2.3.2 Sterile soil conditions

To assess MAOM formation in the absence of microbial decomposition, we sterilized soils for a subset of

the sites (C153, C169, and C61). Briefly, dry soils and necromass were autoclaved separately (60-minutes

at 121ºC and 15 psi in the absence of water). One week prior to the start of the incubation, sterile

deionized water was added to the autoclaved soils under aseptic conditions to bring them to 65% water

holding capacity (WHC). Then, 70 mg of autoclaved necromass was mixed into each soil using a flame

sterilized dissecting probe. Note, there was not enough initial autoclaved necromass remaining to measure

initial C quality fractions. Three soil microcosms were prepared for each soil-necromass combination as

well as three soil only controls (27 microcosms total). Sterile soil microcosms were maintained and

sampled in a similar manner as live soil microcosms.

2.4 Necromass ^{13}C & ^{15}N Tracing

2.4.1 Necromass-derived MAOM measurements

We isolated mineral-associated organic matter (MAOM) using a size fractionation procedure

(Cambardella and Elliott, 1992), modified to minimize the inclusion of dissolved organic matter (DOM)

leached from undecomposed necromass (Craig et al., 2022). First, we added 30 mL of deionized water to

the samples, inverted them to mix, and then removed any floating necromass fragments. Soil solutions

were then centrifuged at 2500 rpm for 10 minutes, and the resulting supernatant containing DOM was

discarded. The soil pellet was resuspended in 5% (w/v) sodium hexametaphosphate solution and shaken

at 120 rpm for 24 hours to disperse soil aggregates. Following dispersion, soils were wet-sieved through a

53-μm mesh sieve. The soil fraction that passed through the sieve, consisting of clay and silt-sized

particles, was considered the MAOM fraction. This MAOM fraction was dried at 80°C and ground for

%C, δ^{13}C, %N, and δ^{15}N analysis. To account for any residual sodium hexametaphosphate, subsample

weights for chemical analysis were adjusted based on the initial mass of sodium hexametaphosphate

added. Similar to initial necromass samples, total C and N content, as well as δ^{13}C and δ^{15}N values, were

determined on ground MAOM samples using an elemental analyzer coupled to a gas-isotope ratio mass

spectrometer (PDZ Europa Elemental Analyzer coupled to a Sercon 20-22 IRMS) at the PSI facility.

2.4.2 Isotopic Mixing Model

We used a two-end member isotopic mixing model to determine the amount of fungal necromass-derived

C and N in the MAOM fraction (Comeau et al., 2013; Buckeridge et al., 2022). This model assumes that

the measured isotopic signature (δ^{13}C or δ^{15}N) of the MAOM fraction is a mixture of two sources: (1) the

original unlabeled SOM and (2) the added ^{13}C- and ^{15}N-labeled fungal necromass. The fraction of

necromass-derived C or N (f_{necro}) in the MAOM, expressed as a proportion of the total C or N in the

MAOM fraction, was calculated using the following equation:

$$f_{necro} = \frac{\delta_{sample} - \delta_{control}}{\delta_{necro} - \delta_{control}}$$

Where δ_{sample} is the measured δ¹³C or δ¹⁵N value for the MAOM fraction in the necromass-amended

soil, $\delta_{Control}$ is the average δ¹³C or δ¹⁵N value for the MAOM fraction in the soil-only control samples,

and δ_{necro} is the average initial δ¹³C or δ¹⁵N value for the added fungal necromass. This equation

calculates the proportion of C or N in the MAOM fraction that is derived from the added fungal

necromass. To calculate the amount of necromass-derived C or N, the calculated fraction (f_{necro}) was

multiplied by the total mass of C or N in the MAOM fraction at the end of the 38-day incubation. Data are

presented as the amount of necromass-derived MAOM-C or -N formed relative to the initial necromass C

or N (% added necromass C or N).

2.4.3 MAOM chemistry

We used DRIFTS to investigate the biochemical composition of MAOM generated at the end of the 38-

day incubation. MAOM samples were processed in a similar manner as initial necromass samples and

annotated using previous literature on the biochemical composition of SOM (Ellerbrock and Kaiser, 2005;

Artz et al., 2008; Pärnpuu et al., 2022). Multiple annotations have been proposed for some peaks since

MAOM is composed of both minerals and organic compounds, and some regions are characteristic of

both (Table S3;

Stuart, 2004).

2.5 Simultaneous Infrared and Potentiometric Titration

2.5.1 Goethite synthesis

To assess the capacity for necromass-soluble compounds to adsorb onto soil minerals abiotically, we

utilized goethite as a model mineral. Naturally occurring in forest soils (Krumina et al., 2017), goethite

was synthesized following the method described by Villacís García et al. (2015). A 2.5 M NaOH solution

was slowly added to a 0.5 M $Fe(NO_3)_3 \cdot 9H_2O$ solution under stirring and N_2 bubbling until the pH reached

approximately 12. The product was aged at 60ºC for 7 days. The suspension was then dialyzed with a

dialysis bag (12-14 kDa) in MiliQ water in a cold room at 4°C, replacing the MiliQ water every 12 hours

until its electrical conductivity was less than 2.0 μS cm^{-1}. The goethite product was transferred to a

polyethylene bottle, purged with N_2 gas to remove carbonate species, and stored under 4°C. The

concentration of the goethite suspension was determined to be 10.99 g L^{-1}. The morphology of goethite

particles was assessed for synthesis success by dropping suspensions onto C-coated copper grids, air-

drying, and observing with a TEM (JEOL-1400 PLUS) operating at 100 keV. The morphology of

synthesized goethite (Fig. S2) is typical for this iron oxide. Using goethite minerals without pre-existing

soil organic matter allowed us to isolate the effects of labile necromass compounds on MAOM chemistry

under abiotic conditions.

2.5.2 Adsorption of necromass compounds on goethite

The adsorption of fungal necromass compounds onto goethite was assessed by infrared spectroscopy

using the Simultaneous Infrared and Potentiometric Titration (SIPT) method, as described by Villacís

García et al. (2015) and Tian et al. (2020). For the adsorption analysis, goethite over-layers were prepared

on an attenuated total reflectance (ATR) ZnSe crystal by evaporating 0.7 mL of a goethite suspension.

The ATR crystal was then coupled to a titration vessel and placed inside an evacuated FTIR (Bruker

Vertex 80 v, Ettlingen, Germany) in a temperature-controlled room at 21°C. A 0.01 M NaCl solution was

added to the titration vessel and gently stirred under N_2 gas to mitigate the effects of atmospheric CO_2.

The pH was adjusted to 5.0 and maintained constant with an automated, computer-controlled burette

system (Methrom 907 Titrando and Tiamo 2.4 software). After equilibrating the over-layer and medium

for approximately 24 hours, a background spectrum of 1024 scans was collected. Subsequently, either

low or high melanin bags, consisting of 80 mg of dry necromass in 40 µm polyester mesh bags (Sintab,

Malmö, Sweden), were immersed in the vessel under constant stirring. IR collection time was set at 256

scans per spectrum. IR spectra were recorded between 700–4000 cm^{-1} at a resolution of 4 cm^{-1} over one

day, with one spectrum recorded every 3.56 minutes, amounting to roughly 400 spectra per necromass

type. Retained spectra ranged from 920 to 3050 cm^{-1}. Peaks in the final spectra (number 400) for both

necromass types were annotated based on literature on the biochemical composition of fungal biomass

and necromass (Dzurendova et al., 2020; Langseter et al., 2021; Maillard et al., 2023c). Necromass

remaining in the mesh bags after incubation accounted for 69.7% and 68.3% of the initial mass for low

and high melanin necromass types, respectively, suggesting similar leachate sizes.

2.6 Data Processing and Statistical Analyses

All statistical analyses were performed in R version 4.3.1 (2023-06-16). Response variables that did not

meet assumptions of normality were transformed or analyzed via non-parametric tests indicated below.

Means are presented in text and tables plus or minus one standard deviation (SD).

2.6.1 Initial Necromass Data

Analysis of variance (ANOVA) was used to test for differences in initial necromass δ^{13}C between

necromass types (high vs low melanin), sterilization treatment (autoclaved vs. non-autoclaved) and their

interaction. Initial necromass % C, % N, and δ^{15}N did not meet the assumptions of ANOVA (data were

non-normal and had unequal variances), as a result, we collapsed necromass type and sterilization

treatment into one factor and performed Welch's Analysis of variance ANOVA with Games–Howell post

hoc tests. Because sterilization treatment did not have an effect on initial δ^{13}C or δ^{15}N (Table S1),

autoclaved and non-autoclaved necromass isotopic values were combined and mean values were used for

high and low melanin necromass initials.

2.6.2 Microbial Community Data

Bioinformatic processing of HTS data was performed using the amptk pipeline (Palmer et al., 2018;

v1.5.4). Briefly, we removed primers and then trimmed sequences to 300 bp. Paired reads were denoised

using DADA2 algorithm (Callahan et al., 2016), using the default parameters (minLen = 50,

maxN = 0, truncQ = 2, maxEE = 2), and then clustered into OTUs at 97% similarity. To assign

taxonomy, we used a hybrid algorithm integrating results from a USEARCH global alignment against the

UNITE (v8, Nilsson et al., 2019) and RDP (training set 19, Wang and Cole, 2024) databases, including

both UTAX and SINTAX classifiers. Data are deposited in the NCBI Short Read Archive (Bioproject #

for bacteria PRJNA1094420; for fungi PRJNA1094416).

We processed data to remove any non-target OTUs and spurious sequences, with OTUs not classified as

Fungi or Bacteria excluded from the analysis. We also discarded bacterial OTUs identified as chloroplast

and lacking genus-level classification were discarded. To account for potential contamination, sequence

reads present in the PCR negative controls were subtracted from corresponding OTU counts in the soil

samples. We used mock communities to establish a threshold for spurious OTU detection. Based on this

analysis, we implemented a read count threshold of four; OTUs with fewer than five reads were removed

from the dataset. Further, we only included samples that had read totals in the 90% data quantile (more

than 2159 and 15347 reads for bacteria and fungi respectively) in the final analyses. From the original 18

samples, we removed two samples due to low sequence totals for bacteria (one replicate from site C55 &

one from C169) and two samples for fungi (one replicate from C153 & one from C71). Finally, to

account for differences in sequence reads among the remaining 16 samples, we rarefied the bacterial and

fungal OTU table to 2202 and 15507 reads per sample, respectively. There were 35232 bacterial reads

and 248112 fungal reads total in the quality-controlled dataset. Fungal OTUs were classified by trophic

mode using FUNGuild (Nguyen et al., 2016). When there were multiple fungal guilds assigned to a single

OTU, we determined the most commonly occurring lifestyle using the FungalTraits database (Põlme et

al., 2020). Of the 1384 fungal OTUs identified, 856 (62%) were assigned a primary lifestyle.

To further characterize the soil bacterial communities, we calculated a weighted mean predicted 16S

rRNA gene copy number (16S GCN) for each soil sample. The number of copies of the 16S rRNA gene

in a bacterial genome is a useful indicator of bacterial life history strategies, with higher copy numbers

generally associated with a more copiotrophic lifestyle (Stoddard et al., 2015). We used the rrnDB

database (https://rrndb.umms.med.umich.edu/) to predict 16S GCN for bacterial OTUs identified in our

study (Lee et al., 2009). Of the 603 bacterial OTUs identified, 342 (57%) were present in the rrnDB

database. While the rrnDB may not fully capture the diversity of soil bacterial communities, it provides a

valuable tool for estimating 16S GCN. Based on empirical data, this offers a more robust approach than

relying solely on phylum-level classifications (Jia et al., 2023). For each soil sample, we calculated a

weighted mean 16S GCN by multiplying the predicted 16S GCN of each OTU by its relative abundance

in the sample and summing these values (Nemergut et al., 2016; West et al., 2023). A higher weighted

mean 16S GCN suggests a community dominated by copiotrophic bacteria, while a lower value indicates

a more oligotrophic community.

We used type III ANOVAs ('Anova' function in the Car package) were used to test for site differences in

mean richness (number of unique OTUs) and diversity (Shannon diversity) in bacterial and fungal

communities, as well as the weighted mean 16S GCN and fungal saprotroph abundances. We investigated

differences in bacterial and fungal community composition across sites by calculating a Bray-Curtis

dissimilarity matrix from the rarefied sequence read abundances using the 'metaMDS()' function in the

Vegan package (Oksanen et al., 2001). To visualize the soil physiochemical variables covarying with

microbial community composition across sites, we fit vectors to non-metric multidimensional scaling

(NMDS) ordination plots using the 'envfit' function in the Vegan package. We used permutational

multivariate analyses of variance (PERMANOVA) to assess site effects on microbial community

composition.

2.6.3 MAOM Formation and DRIFTS Data

One sample from site C71 amended with high melanin necromass was removed from the dataset because

both fungal-derived MAOM-C (2.83%) and MAOM-N (19.6%) formation was more than two standard

deviations below mean values (5.74 and 41.6% for MAOM-C and MAOM-N respectively). We used

ANOVA with type III sum of squares to test for significant effects of necromass type and site, as well as

their interaction on MAOM formation (MAOM-C and N data were log transformed prior to analysis for

sterile microcosm soils). When interactions were not significant only main effects were included in final

models. We carried out post-hoc tests using Sidak multiple comparison in the Emmeans package (Lenth,

2017). Correlations between initial soil properties and response variables were determined by calculating

Pearson's correlation coefficients.

To assess the relative contributions of initial necromass melanization, soil physiochemical and microbial

properties on fungal necromass contributions to mineral-associated organic matter (MAOM) formation,

we employed hierarchical partitioning analysis. Hierarchical partitioning is a variance partitioning

method that quantifies the independent and joint contributions of multiple predictor variables to the total

variance explained in a response variable (Chevan and Sutherland, 1991). This analysis addresses

multicollinearity by systematically examining all possible combinations of predictor variables and

calculating the goodness-of-fit (e.g., R-squared) for each combination (Mac Nally, 2002). Using the

rdacca.hp package (Lai et al., 2021), we partitioned the variance in MAOM formation among three

predictor sets: (1) initial necromass type (high vs. low melanin), (2) soil physiochemical properties (pH,

% C, % N, % clay, and % FeOx), and (3) microbial properties (fungal richness, fungal Shannon diversity,

fungal saprotroph relative abundance, bacterial richness, bacterial diversity, and weighted mean predicted

16S GCN).

2.6.4 DRIFT and SIPT Spectral Data

We identified IR peak heights using the quick peaks function with a constant baseline in OriginLab

(Northampton, MA, USA). We included peaks that fell within a range of $\pm$ 10cm^{-1} of annotated peaks

(Table S3). We used principal component analysis (PCA) biplots of DRIFT spectra to visualize separation

of MAOM samples based on site or necromass treatment. Principle component loadings were plotted

against wavenumber to identify peaks that correspond with the first two PCA axes. To explore changes in

SIPT peak height over time, we fit asymptotic regression models for annotated peaks, for both high and

low melanin necromass-goethite suspensions using the 'nls' and 'SSasymp' functions in the Stats

package. The model below describes peak height (y) as a function of hourly time (x), where $\emptyset_1$ is the

horizontal asymptote on the right of the curve or the maximum value of y, $\emptyset_2$ is a numeric parameter

representing the response when the input is zero and $\emptyset_3$ is the natural logarithm of the rate constant (lrc;

Drever et al., 2023).

$$y = \emptyset_1 + (\emptyset_2 - \emptyset_1)exp[-exp(\emptyset_3)x]$$

Initial Soil Microbial Properties						
Site ID	Bacterial OTU Richness	Fungal OTU Richness	Bacterial Shannon Diversity	Fungal Shannon Diversity	Weighted Mean Predicted 16S rRNA GCN	Fungal Saprotroph Relative Abundance
C153	190 (33)[a]	113 (25)[b]	4.45 (0.19)[a]	4.03 (0.03)[ab]	2.03 (0.08)[d]	0.370 (0.04)[b]
C55	146 (61)[a]	245 (16)[a]	3.80 (0.25)[b]	4.05 (0.05)[ab]	4.50 (0.28)[a]	0.302 (0.07)[b]
C169	125 (40)[a]	262 (24)[a]	3.83 (0.27)[ab]	3.64 (0.17)[bc]	2.94 (0.22)[c]	0.263 (0.05)[b]
C71	121 (37)[a]	258 (25)[a]	3.54 (0.22)[b]	4.38 (0.03)[a]	3.80 (0.27)[b]	0.612 (0.03)[a]
C13	112 (15)[a]	132 (25)[b]	3.71 (0.10)[b]	2.86 (0.29)[d]	2.50 (0.11)[cd]	0.580 (0.06)[a]
C61	144 (16)[a]	162 (13)[b]	4.00 (0.10)[ab]	2.30 (0.16)[cd]	2.40 (0.09)[cd]	0.378 (0.03)[b]

Figure 1. A. Initial soil microbial properties including alpha diversity metrics (OTU richness and Shannon diversity), as well as a weighted mean predicted 16S rRNA gene copy number (16S rRNA GCN) and fungal saprotroph relative abundances for initial soils (mean, SD, n = 3*). Letters denote differences in site means as determined by ANOVA. B. NMDS ordinations for B. Bacterial and C. Fungal communities with soil property vectors. *Due to some initial soil samples not passing quality control (see methods), n=2 for sites C55 and C169 for bacterial metrics & n=2 for sites C71 and C153 for fungal metrics.

3. Results

3.1 Initial physicochemical and microbial properties of microcosm soils

While the sites were primarily selected to vary in soil oxalate-extractable Fe, FeOx was significantly correlated with clay content (r = 0.95 , p < 0.01) and total C (r = 0.94, p < 0.01). Initial microbial community composition varied more across sites than OTU richness and Shannon diversity (Figure 1A). PERMANOVA results demonstrated significant differences in the soil bacterial (R^2 = 0.90, F = 17.2, p =

0.01) and fungal (R^2 = 0.95, F = 35.9, p = 0.01) communities by site. In NMDS ordination biplots,

microbial communities from sites with high soil pH, C, N, clay and FeOx grouped together, with bacteria

exhibiting stronger correlations with soil properties (Fig. 1B) relative to fungi (Fig. 1C). Bacterial OTU

richness (r = 0.57, p = 0.03) and Shannon diversity (r = 0.67, p < 0.01) were positively correlated with

soil FeOx content.

3.2 Fungal necromass-derived MAOM formation and chemistry in live soil microcosms

After 38 d of incubation, seven times more initial fungal necromass N (42.2 ± 5.2%, CV = 0.12) was

incorporated into MAOM than necromass C (5.82 ± 0.96%, CV = 0.16). Site identity explained 49.2%

and 37.6% of the variation in necromass-derived MAOM-C (F = 6.62, p < 0.001) and MAOM-N (F =

3.79, p = 0.01) formation. The accumulation of fungal-derived MAOM-C ranged from 4.72 ± 0.47% of

initial necromass C for soils from the Jackson County site (C71) to 6.92 ± 1.1% of initial necromass C

for soils from the Sullivan County site (C153) which had the highest initial clay and FeOx content.

Fungal-derived MAOM-N also differed among sites (Table S4), ranging from 40 ± 4% for soils from

the Harrison County site (C61) to 48 ± 6% of initial necromass N for soils from Greene County site

(C55). Though the greatest differences in MAOM formation were among sites, necromass type

significantly affected MAOM-C formation (Fig. 2A, F = 6.21, p = 0.02) and had a moderately significant

effect on MAOM-N formation (Fig. 2B, F = 3.37, p = 0.08). On average, low-melanin necromass

contributed 9% more to MAOM-C (HM: 5.6 ± 0.8%, LM: 6.1 ± 1.0%) and 6% more to MAOM-N (HM:

41 ± 6%, LM: 43 ± 5%) than high-melanin necromass. Interaction terms between site and necromass type

were not significant for both MAOM-C (F = 1.3; p = 0.3) and MAOM-N (F = 0.17; p = 0.9) formation.

Figure 2. Box and whisker plots of fungal necromass-derived mineral-associated organic matter
(MAOM) A. carbon (C) and B. nitrogen (N) formation in live soil microcosms after a 38-day incubation.

Correlations between initial soil properties and MAOM formation differed between necromass types (Fig.
3). For soils amended with high-melanin necromass (Fig. 3A), necromass-derived MAOM-C was
negatively correlated with initial fungal saprotroph abundance ($r = -0.60$, $p = 0.03$), and MAOM-N
formation was positively correlated with the weighted mean predicted 16S GCN ($r = 0.63$, $p = 0.02$). For
soils amended with low-melanin necromass (Fig. 3B), MAOM-C formation was positively correlated
with soil FeOx ($r = 0.74$, $p < 0.01$), clay content ($r = 0.71$, $p < 0.01$), total C ($r = 0.63$, $p = 0.02$), bacterial
OTU richness ($r = 0.61$, $p = 0.02$), and Shannon diversity ($r = 0.74$, $p < 0.01$). In contrast, in these soils,
fungal saprotroph relative abundance was negatively correlated with both MAOM-C ($r = -0.57$, $p = 0.04$)
and MAOM-N formation ($r = -0.62$, $p = 0.02$).

Figure 3. Pearson's correlation matrix and coefficients (Corr) for relationships among initial soil properties and fungal necromass-derived MAOM-C (MAOM_Fun_C) and MAOM-N (MAOM_Fun_N) formation for soils amended with A. high melanin and B. low melanin necromass. Significant Pearson correlation coefficients are indicated in squares and non-significant correlations (P > 0.05) are represented by white squares. Soil physiochemical properties have "Soil_" before them and include initial soil % carbon (C), % nitrogen (N), % clay, % Iron Oxide (FeOx) and pH. Microbial properties have "Micro_" before them and include fungal OTU richness (FunRich), fungal Shannon diversity (FunDiv), fungal saprotroph relative abundance (FunSapRelab), bacterial OTU richness (BacRich), bacterial Shannon diversity (BacDiv), and weighted mean predicted 16S gene copy number (Bac16SGCN) .

Because soil variables were highly collinear (Fig. 3), we used hierarchical partitioning to determine the unique contribution of each initial predictor set (soil physiochemical, microbial, and necromass type) to MAOM formation, as well as their combined effects (Table 2). Overall, initial predictors explained more of the total variation in fungal-derived MAOM-N (adj. $R^2 = 0.42$) than in MAOM-C formation (adj. $R^2 = 0.35$). Soil physiochemical properties were the strongest predictors of fungal-derived MAOM formation,

explaining 59.9% and 68.4% of the variation in MAOM-C and MAOM-N formation, respectively (Table

2). While soil microbial properties also contributed substantially to both MAOM-C (32.5%) and MAOM-

N (32.8%) formation, the unique contribution of microbial predictors was less pronounced due to their

correlation with soil physiochemical properties (Fig 3). Interestingly, necromass type exhibited a minor

negative relative importance for MAOM-N formation (–1.12%), meaning that necromass type alone did

not directly enhance MAOM-N formation. The joint effects of soil and microbial predictor sets were

considerable, indicating complex interactions driving MAOM formation.

Table 2. Results of hierarchical partitioning analysis including total average shared effects with other

predictors and individual importance (Individual effect divided by total adjusted R^2) of different initial

soil predictor sets[*]

Response Variable	Predictor Set	Unique	Average Shared	Individual Importance	Relative Importance (% of R^2)
Fungal Derived MAOM-C ($R^2 = 0.35$)	Necromass Type	0.0383	-0.0122	0.0261	7.50
	Soil Physiochemical	0.0186	0.1898	0.2084	59.9
	Soil Microbial	-0.0782	0.1913	0.1131	32.5
Fungal Derived MAOM-N ($R^2 = 0.42$)	Necromass Type	0.0004	-0.0051	-0.0047	-1.12
	Soil Physiochemical	0.0467	0.2413	0.2880	68.4
	Soil Microbial	-0.1067	0.2448	0.1381	32.8

*Initial soil physiochemical properties include initial soil % carbon (C), % nitrogen (N), %

clay, % Iron Oxide (FeOx) and pH. Initial soil microbial properties include alpha diversity

metrics (OTU richness and Shannon diversity), as well as a weighted mean predicted 16S

rRNA gene copy number and fungal saprotroph relative abundances for initial soils.

The chemical composition of MAOM from soils at the end of the 38-day incubation was characterized by

strong mineralogy-associated bands at 3695 – 3621 cm^{-1} assigned to O-H bonds, indicative of either clay

or aluminum minerals, as well as Si-O bonds in the 695-474 cm^{-1} region, associated with silicate minerals

(Fig. 4A & Table S3). The mean intensity of bands at 1284 cm^{-1} and from 940-779 cm^{-1} varied notably

among sites, attributed to phenolic and aromatic compounds with C-O and C-H functionalities. While

first two principal components explained ~70% of the variation in MAOM chemistry, the PCA plot did

not show strong separation of spectra among necromass amended and soil only controls (Fig. 4B). The

clearest separation, however, occurred between soils originating from the Sullivan County site (C153) and

other samples along the first principal component axis (PC1). Site explained a much more significant

amount of spectral variation (R^2 = 0.60, F = 14.4, p < 0.001) than necromass type (R^2 = 0.02, F = 1.41, p

= 0.20), supporting the PCA visualization. Peaks at 3621–3695, 1186–1360, and 779–940 cm^{-1}, indicative

of oxygenated functionalities associated with clay and silicate minerals, were negatively correlated with

the loading of PC1 (Fig. 4C). These findings support the unique mineralogy of the Sullivan County site

(C153), which formed the greatest amount of necromass-derived MAOM-C (Table S4).

Figure 4. A. Diffuse Reflectance Infrared Fourier transform spectroscopy (DRIFTS) spectra with

annotated peaks for mineral-associated organic matter (MAOM) grouped by site. B. Principal Component

Analysis (PCA) biplot of DRIFT spectra for MAOM from live soil microcosms. C. PCA loadings for the

first two principal component (PC) axes.

3.3 Fungal necromass-derived MAOM formation & chemistry in sterile microcosm soils

Sterilization significantly affected both MAOM-C (F = 44.1, p < 0.001) and MAOM-N (F = 32.0, p <

0.001) formation. More necromass-derived MAOM-C (Fig. 5A) and less MAOM-N (Fig. 5B)

accumulated in sterile soils than in live soils for both high- and low-melanin necromass types.

Specifically, for high- and low-melanin necromass, MAOM-C formation was 50% and 40% higher,

respectively, in sterile soils, while MAOM-N formation was 30% lower. Moreover, for soils amended

with low melanin necromass, fungal-derived MAOM-C was higher in sterile soils despite the autoclaved

necromass having lower initial % C (Tables S1). In sterile soils MAOM formation was similar between

high and low melanin necromass types regardless of site (Table S5). Similar to live soil microcosms,

DRIFT spectra of MAOM from sterile microcosms did not differ between necromass amended and soil

only controls (PERMANOVA: R^2 = 0.01, F = 0.59, p = 0.70; Fig S4). Additionally, spectra did not differ

between sterile and live soil only control microcosms (PERMANOVA: R^2 = 0.01, F = 0.14, p = 0.9, Fig.

S3).

Figure 5: Box and whisker plots of fungal necromass-derived mineral-associated organic matter (MAOM) A. carbon and B. nitrogen formation in live and sterile (autoclaved soils) after a 38-day incubation. Dark gray boxes and points represent high melanin (HM) necromass and light gray boxes and points represent low melanin (LM) necromass in either live (filled boxes and points) or sterile (open boxes and points) soil conditions.

3.4 The adsorption of fungal necromass compounds onto synthesized goethite

With the exception of higher $\delta^{13}C$ enrichment for low melanin necromass, the initial C values did not differ between the two necromass types (Table S1). The cell-soluble and unhydrolyzable fractions were also statistically similar between high and low melanin necromass. However, DRIFT spectra of initial necromass types did show a higher aromatic peak (C=C) at 1400 cm^{-1} for high melanin necromass (Fig. S4). We also observed a number of clear spectral differences in the labile compounds that adsorbed onto goethite minerals between necromass types (Fig 6). Specifically, IR spectra showed much higher intensity of amide bond peaks (N-H & C=O bonds between 1650 cm^{-1} and 1540 cm^{-1}) for low melanin necromass as well as greater intensity of the 1108 cm^{-1} aromatic bond peak (Fig. 6A). Conversely, IR spectra for high melanin necromass had higher intensities of aliphatic (C-H) and polysaccharide (C-O) bond peaks located at absorbances of 2925 and 1108 cm^{-1}, respectively (Fig 6B). The IR signal intensities also varied over time, with peak heights changing rapidly at first and then stabilizing or reaching an asymptote within

635   24 hours (Fig. 6B). We found that compounds from high melanin necromass tended to converge to their

636   asymptote (maximum peak height) at faster rates, especially for the aforementioned aliphatic (2925 cm^{-1})

637   and polysaccharide (1108 cm^{-1}) associated bonds (Fig S5).

Figure 6: A. Simultaneous infrared and potentiometric titration (SIPT) infrared spectrum (spectra number

400) for high and low melanin necromass suspensions after 24 hours. B. Changes in peak intensity over

time for aliphatic (C-H), amide (N-H), aromatic (C-O) and polysaccharide (C-O) derived functionalities

(numbers represent peak IDs- see Fig. S5) from SIPT spectra of high (black line) and low (gray line)

necromass compounds adsorbed onto synthesized goethite minerals under abiotic conditions.

4. Discussion

Long-term soil C and N persistence is thought to be largely determined by associations with soil minerals,

with a diversity of microbial inputs and minerals offering numerous means for binding OM (Cotrufo et

al., 2013; Kallenbach et al., 2016; Liang et al., 2017; Sokol and Bradford, 2019; Lehmann et al., 2020).

Here we assessed how melanization interacts with different soil physicochemical and microbial properties

to determine fungal necromass C and N stabilization via mineral associations. We hypothesized that

MAOM formation and chemistry would differ from high and low melanin fungal necromass inputs, but

that soil iron oxide (FeOx) content would be a strong overall predictor of necromass-derived MAOM-C

and MAOM-N accumulation regardless of melanization. Indeed, soil physicochemical properties,

particularly those related to clay and FeOx content, were the strongest predictors of MAOM formation,

explaining ~60% and ~68% of the variation in MAOM-C and MAOM-N, respectively (Table 2). We

found that high and low melanin necromass differed in amide and polysaccharide associated peaks for

soluble necromass compounds that adsorbed onto synthesized FeOx minerals. However, total MAOM-C

and MAOM-N formation were similar between necromass types in sterile soils (Fig. 5). This suggests that

initial substrate chemistry does not directly predict MAOM-C and MAOM-N accumulation across diverse

soil mineralogies. Instead, it appears that microbial degradation of fungal necromass, which differs

between high and low melanin necromass types, more strongly impacts MAOM-C and MAOM-N

formation. This was evident in the live soil microcosms, where bacterial richness and initial FeOx content

were positively related to MAOM-C formation, but only for soils amended with low melanin necromass

(Fig. 3). Further, we found that initial microbial community properties, including fungal saprotroph

abundance and bacterial richness and life history strategies (as indicated by 16S rRNA gene copy

number), were significant predictors of MAOM formation, contributing ~32% to the explained variation

in both MAOM-C and -N (Table 2). Taken together, these findings point to the potential for different

stabilization pathways for C- and N-containing fungal necromass compounds mediated by necromass

melanization, soil properties, and microbial communities, as discussed below.

4.1 Interactive Effects of Soil Properties and Microbial Communities on MAOM Formation

The persistence of microbial necromass in soil may be promoted abiotically by associating directly with

mineral surfaces (direct sorption pathway) or biotically via microbial assimilation (microbial synthesis

pathway); in reality, both pathways occur simultaneously and their relative importance for fungal

necromass- C and -N stabilization is unclear (Buckeridge et al., 2022). On average, we show that 8% of

total MAOM-C and 20% of MAOM-N originated from fungal necromass decay over a 38-day period.

Greater retention of necromass-derived N than C may be due to differences in bond affinities for N-

containing compounds and/or greater incorporation of fungal-derived N in living microbial biomass

(Omoike and Chorover, 2006; Kleber et al., 2007; Wang et al., 2024), which may also decouple MAOM-

C and MAOM-N formation. Nitrogenous compounds bind readily to mineral surfaces (Kleber et al.,

2007) and microbes may use them with high efficiency, resulting in N-rich microbial products that then

subsequently bind to soil minerals (Cotrufo et al., 2013; Kallenbach et al., 2016).

We found that the intensity of N-H bonds from necromass-derived amide compounds increased on

goethite minerals within 24 hours, in support of the direct sorption pathway. However, 30% less initial

necromass-N was converted into MAOM-N in sterile soils, indicating that microbial synthesis may be the

more important pathway for necromass-derived MAOM-N formation (Wieder et al., 2014; Wang et al.,

2020). These findings agree with those of Keiluweit et al. (2012), who tracked ^{13}C and ^{15}N isotopically

fungal cell wall residues onto FeOx surfaces during a three-week incubation experiment. That study

revealed that amide-N from fungal chitin was consumed by hyphal-associated bacteria, assimilated into

newly synthesized bacterial proteins, and adsorbed onto FeOx surfaces. They also found greater

enrichment of fungal-derived ^{15}N relative to ^{13}C on FeOx minerals, which may be explained by greater

microbial assimilation of fungal-N and greater mineralization of fungal-C (Keiluweit et al., 2012; Wang

et al., 2024).

Our SIPT analyses also showed evidence for direct sorption of C-containing bonds from carboxylic acids,

a functionality known to be important in stabilizing microbial necromass via interactions with FeOx

minerals (Lützow et al., 2006; Kleber et al., 2015). We also found that less necromass C became MAOM-

C in live soils, suggesting that microbial synthesis did not promote the retention of necromass C to the

same extent as necromass N. Buckeridge et al. (2020) also found higher retention of fungal necromass

[13]C in sterile compared to live soils in a short-term (three day) study in grassland soil microcosms. Given

the fast-decaying nature of fungal necromass (Brabcová et al., 2016; Fernandez et al., 2016), it is possible

that the majority of necromass C was lost via respiration (Buckeridge et al., 2020). We did not, however,

track necromass-derived $^{13}CO_2$ loss or incorporation into other soil pools (microbial biomass, dissolved

organic matter, or particulate organic matter), making it challenging to differentiate between abiotic and

biotic processes in stabilizing fungal necromass-C. Additionally, the steam sterilization of soils may have

altered the solubility or leaching of the pre-existing SOM prior to necromass additions (Egli et al., 2006),

although we did not find spectral differences in MAOM chemistry after 38 days.

Overall, these results suggest that the microbial synthesis pathway may be more crucial for N

stabilization, while microbial processing can limit C stabilization by promoting its loss through

respiration or transformation into other soil organic matter pools. This is further supported by the

observed correlations between initial soil predictors and MAOM formation. For instance, fungal

saprotroph relative abundance was negatively correlated with MAOM-C formation, indicating that higher

abundances of these fungi might lead to reduced MAOM formation, potentially due to their rapid

consumption of labile necromass compounds. Sites with highest fungal saprotroph abundance (C13 and

C71) featured increased relative abundances of fast-growing genera such as *Penicillium*, *Ascolobus* and

Umbelopsis, which were likely targeting necromass polysaccharides (Baldrian et al., 2011; Štursová et al.,

2012). Fast growth in fungi is thought to be associated with higher metabolic costs (Arendt, 1997;

Veresoglou et al., 2018), which may limit microbial transfer of necromass C to MAOM.

Correlations between initial microbial communities and MAOM formation may be explained by the

presence of a core fungal necrobiome. Cantoran et al., (2023) found that a consistent set of bacterial and

fungal taxa, including fungal saprotrophs and bacterial copiotrophs, is associated with decaying

necromass across various studies and biomes. We acknowledge that initial microbial community

composition, as determined by high throughput sequencing, does not necessarily reflect microbial

community composition or activity over the course of our study. That said, the presence of this core

necrobiome might contribute to similar outcomes in MAOM formation across different soil environments.

Additionally, Maillard et al. (2023a) demonstrated that initial microbial community composition was the

strongest predictor of necromass decay across a diverse set of forest plots, suggesting assessments of the

initial microbial community can be ecologically informative. Future research investigating the temporal

dynamics of microbial communities associated with necromass decomposition would provide valuable

insights into the specific roles of these core taxa and their functional contributions to MAOM formation.

4.2 Indirect Role of Necromass Melanization in MAOM Formation

Among fungal species, initial substrate chemistry has been shown to be an important control on

necromass decay (Fernandez and Koide, 2012, 2014), with increased melanin content consistently

slowing necromass decomposition across a range of soil conditions (Fernandez and Kennedy, 2018;

Fernandez et al., 2019; Maillard et al., 2020; See et al., 2021). Consistent with this, we found that high

melanin necromass contributed less C to MAOM when compared with low melanin necromass in our live

soil microcosms, despite similar sized cell-soluble and acid-unhydrolyzable fractions in initial necromass.

This similarity in starting C quality fractions might be attributed to our culturing method, where fungi

were grown in C-starved media, potentially limiting the production of labile compounds and masking

differences in specific chemical components between the necromass types. However, it's important to

recognize that the composition of those fractions, rather than just their size, likely differed between the

necromass types. For example, the cell-soluble fraction of fungal necromass typically includes simple

sugars, amino acids, lipids, and soluble proteins (Ekblad et al., 2013; See et al., 2021), but it can also

contain a variety of aromatic compounds not associated with melanin, such as those involved in signaling,

defense, or pigment production (Avalos and Limón, 2021). This difference in composition is supported by

our SIPT experiment, which revealed distinct adsorption patterns for high and low melanin necromass on

goethite. For instance, we observed higher intensity of amide and aromatic bond peaks for low melanin

necromass and greater intensity of aliphatic and polysaccharide bond peaks for high melanin necromass.

These differences in labile chemistry likely contributed to the observed indirect effects of melanization on

MAOM formation. By "indirect effects," we refer to the influence of melanization on factors beyond

direct sorption to minerals. Melanin has been shown to limit microbial uptake of necromass C and N

during both early (7-14 days) and later (35-77 days) stages of decay (Maillard et al., 2023b), likely also

preventing microbial assimilation in this study. Additionally, Novak et al. (2024) found that bacterial

growth was higher on low melanin necromass. We observed a positive relationship between initial soil

bacterial richness and MAOM-C formation, but only for soils amended with low melanin necromass. An

increased number of bacterial species might enhance resource partitioning and niche complementarity,

leading to more efficient resource utilization and potentially higher carbon use efficiency (Bell et al.,

2005; Domeignoz-Horta et al., 2021). Interestingly, we found that 16S rRNA copy number was positively

related to MAOM-N formation but only for high melanin necromass. Melanized necromass can have

significant quantities of N in protein-melanin complexes (Bull, 1970), and copiotrophic bacteria, with

their diverse metabolic capabilities, may be able to access and efficiently target this cell wall-bound N

(Maillard et al., 2023a). While numerous studies have demonstrated the stability of melanin in soils (e.g.,

Fernandez et al., 2013; Siletti et al., 2017), our results suggest that melanized necromass might be

primarily incorporated into the SOM pool as particulate organic matter (POM). This could be due to

several factors, including the hydrophobicity of melanin, which can limit the access of microbes and their

enzymes to necromass compounds and may also contribute to the formation of stable aggregates within

SOM (Fernandez and Kennedy, 2018; Netherway and Bahram, 2024). Additionally, melanin has been

shown to form complexes with N-rich cell wall components like chitin, further reducing decomposer

access to labile sources of N (Bull, 1970; Fernandez and Koide, 2014; Ryan et al., 2020) and reducing the

availability of necromass components for direct sorption to minerals (Ryan et al., 2020). Although our

study did not include the effects of detritivore activity on necromass fragmentation, recent findings by

Pérez⬚Pazos et al. (2024) suggest this may not be a major limitation. Their work, which focused on

microbial decomposers, showed that particle size did not significantly influence *H. bicolor* necromass

decomposition rates. However, it's important to consider that grinding may still have influenced other

aspects of necromass-soil interactions in our study, such as microbial access or adsorption to mineral

surfaces. Future studies comparing ground and intact necromass, including the role of detritivores, could

further explore these potential effects.

Although we found differences in MAOM-C formation between necromass types, the chemical analyses

of MAOM generated after 38 days did not differ between soils amended with high and low melanin

necromass. Our ability to detect differences in MAOM chemistry in necromass amended soils may have

been restricted by the short time span of the study and relatively small starting amounts of necromass.

Further, we emphasize that it is important to note that MAOM is rich in minerals, which, like organic

compounds, absorb infrared light and thus DRIFTs peaks could be indicative of either organic or mineral

compounds or a combination of both. Notwithstanding difficulties in differentiating peaks between

organic and mineral functionalities, the strongest spectral differences in ordination space were between

the Sullivan County site (C153) and the other five sites. In addition to being a poorly drained riparian soil,

differences in the dominant tree species at the Sullivan County site may have contributed to spectral

variation in MAOM chemistry. Though microbes contribute to chemically diverse MAOM (Golchin et

al., 1996; Kallenbach et al., 2016), the direct incorporation of plant compounds into MAOM (Angst et al.,

2021) is increasingly recognized as an important pathway of SOM stabilization and may contribute

significantly more to SOM formation under certain conditions (Whalen et al., 2022). We also

acknowledge that a major limitation of our microcosm design was the absence of living plants, thus

preventing live plant inputs as well as inputs and activities of microbial groups directly associated with

plant roots (e.g. ectomycorrhizal fungi, Fernandez and Kennedy, 2018). Future studies are needed to track

fungal necromass contributions to MAOM formation and chemistry under field conditions with diverse

plant communities and more complete soil food webs.

4.3 Conclusions and Future Directions

On the whole, our study showed that microbial decomposition decreased the retention of fungal

necromass-derived-C while increasing the retention of necromass-derived-N (Fig. 5), highlighting the

importance of microbial transformations in the loss of fungal C and preservation of fungal N. Further, we

demonstrate the direct potential of fungal cell wall melaninization to influence soluble necromass

chemistry (Fig. 6) and limit mineral-associated C accumulation (Fig. 2). These findings align with

previous studies that report slower decay and microbial assimilation of melanized fungal necromass under

field conditions.

While our study provides valuable insights into the factors controlling fungal necromass contribution to

MAOM in temperate forest soils, it is important to consider how these findings might apply to other

ecological settings and soil types. For instance, the relative importance of soil mineralogy and microbial

communities in driving MAOM formation might differ in ecosystems with contrasting soil properties,

such as grasslands or tropical forests. In grasslands, where soils are often characterized by lower clay and

FeOx content (Jobbágy and Jackson, 2000; Rasmussen et al., 2018), microbial processing might play a

more dominant role in necromass stabilization. Conversely, in tropical forests with highly weathered soils

and abundant FeOx (Kleber et al., 2007), mineral sorption might be the primary mechanism for

necromass persistence. Furthermore, the influence of necromass chemistry, including melanization, might

vary depending on the dominant fungal species and their associated decomposer communities (Klink et

al., 2022). Further, in ecosystems with high fungal diversity, such as tropical forests (Tedersoo et al.,

2014), the range of necromass chemistries might be broader, leading to more diverse interactions with soil

minerals and microbes. Finally, environmental factors such as climate and nutrient availability could

further modulate the effects of necromass chemistry and microbial communities on MAOM formation

(Heckman et al., 2023).

By demonstrating the interactive effects of soil mineralogy, microbial communities, and necromass

chemistry, we highlight the need for integrated approaches that consider multiple factors when predicting

and managing soil C dynamics. Future research should focus on scaling up these findings to different

ecosystems and soil types through comparative studies or meta-analyses, allowing us to develop more

comprehensive models of soil organic matter formation and persistence. We acknowledge that using a

single species may limit the generalizability of our findings to other fungal taxa with potentially different

chemical compositions. Future studies incorporating diverse fungal species are needed to confirm the

broader applicability of our results. Additionally, future experiments should aim to incorporate greater

realism by utilizing necromass from diverse fungal species, including those produced and decayed in situ,

and by conducting experiments under field conditions with intact soil food webs. This will allow for a

more comprehensive understanding of the factors controlling fungal necromass contribution to MAOM

across different ecosystems and soil types. This knowledge will be crucial for developing effective

strategies to enhance soil health and promote carbon sequestration in the face of global environmental

change.

<u>Acknowledgments</u>

We thank Joy Gallion the former CFI director at DNR for her help in obtaining the soils used in this study

and Brien Beidler for help in fabricating vented microcosm tops. We are grateful for the contributions of

the three anonymous reviewers whose comments and careful edits significantly improved this manuscript.

This work was funded by a National Science Foundation DEB MacroSysBIO & NEON-Enabled Science

grant (Award# 2106096) and the Department of Energy, Office of Biological and Environmental

Research, Genomic Science Program as part of the *Microbes Persist* Soil Microbiome SFA (Award #

SCW1632).

<u>References</u>

Adingo, S., Yu, J.-R., Xuelu, L., Li, X., Jing, S., Xiaong, Z., 2021. Variation of soil microbial carbon use

efficiency (CUE) and its Influence mechanism in the context of global environmental change: a

review. PeerJ 9, e12131. doi:10.7717/peerj.12131

Angst, G., Mueller, K.E., Nierop, K.G.J., Simpson, M.J., 2021. Plant- or microbial-derived? A review on

the molecular composition of stabilized soil organic matter. Soil Biology and Biochemistry 156,

108189. doi:10.1016/j.soilbio.2021.108189

Arendt, J.D., 1997. Adaptive Intrinsic Growth Rates: An Integration Across Taxa. The Quarterly Review

of Biology 72, 149–177. doi:10.1086/419764

Artz, R.R.E., Chapman, S.J., Jean Robertson, A.H., Potts, J.M., Laggoun-Défarge, F., Gogo, S., Comont,

L., Disnar, J.-R., Francez, A.-J., 2008. FTIR spectroscopy can be used as a screening tool for

organic matter quality in regenerating cutover peatlands. Soil Biology and Biochemistry 40, 515–

527. doi:10.1016/j.soilbio.2007.09.019

Avalos, J., Limón, M.C., 2021. Fungal Secondary Metabolism. Encyclopedia 2, 1–13.

doi:10.3390/encyclopedia2010001

Bailey, V.L., Bond◻Lamberty, B., DeAngelis, K., Grandy, A.S., Hawkes, C.V., Heckman, K., Lajtha, K.,

Phillips, R.P., Sulman, B.N., Todd◻Brown, K.E.O., Wallenstein, M.D., 2018. Soil carbon

cycling proxies: Understanding their critical role in predicting climate change feedbacks. Global

Change Biology 24, 895–905. doi:10.1111/gcb.13926

Baldrian, P., Voříšková, J., Dobiášová, P., Merhautová, V., Lisá, L., Valášková, V., 2011. Production of

extracellular enzymes and degradation of biopolymers by saprotrophic microfungi from the upper

layers of forest soil. Plant and Soil 338, 111–125. doi:10.1007/s11104-010-0324-3

Beidler, K.V., Phillips, R.P., Andrews, E., Maillard, F., Mushinski, R.M., Kennedy, P.G., 2020. Substrate

quality drives fungal necromass decay and decomposer community structure under contrasting

vegetation types. Journal of Ecology 108, 1845–1859. doi:10.1111/1365-2745.13385

Bell, A.A., Wheeler, M.H., 1986. Biosynthesis and Functions of Fungal Melanins. Annual Review of

Phytopathology 24, 411–451. doi:10.1146/annurev.py.24.090186.002211

Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L., Lilley, A.K., 2005. The contribution of species

richness and composition to bacterial services. Nature 436, 1157–1160. doi:10.1038/nature03891

Brabcová, V., Nováková, M., Davidová, A., Baldrian, P., 2016. Dead fungal mycelium in forest soil

represents a decomposition hotspot and a habitat for a specific microbial community. New

Phytologist 210, 1369–1381. doi:10.1111/nph.13849

Buckeridge, K.M., La Rosa, A.F., Mason, K.E., Whitaker, J., McNamara, N.P., Grant, H.K., Ostle, N.J.,

2020. Sticky dead microbes: Rapid abiotic retention of microbial necromass in soil. Soil Biology

and Biochemistry 149, 107929. doi:10.1016/j.soilbio.2020.107929

Buckeridge, K.M., Mason, K.E., Ostle, N., McNamara, N.P., Grant, H.K., Whitaker, J., 2022. Microbial

necromass carbon and nitrogen persistence are decoupled in agricultural grassland soils.

Communications Earth & Environment 3, 114. doi:10.1038/s43247-022-00439-0

Bull, A.T., 1970. Chemical Composition of Wild-type and Mutant Aspergillus nidulans Cell Walls. The

Nature of Polysaccharide and Melanin Constituents. Journal of General Microbiology 63, 75–94.

doi:10.1099/00221287-63-1-75

Butler, M.J., Day, A.W., 1998. Fungal melanins: a review. Canadian Journal of Microbiology 44, 1115–

1136. doi:10.1139/w98-119

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2:

High-resolution sample inference from Illumina amplicon data. Nature Methods 13, 581–583.

doi:10.1038/nmeth.3869

Cambardella, C.A., Elliott, E.T., 1992. Particulate Soil Organic Matter Changes across a Grassland

Cultivation Sequence. Soil Science Society of America Journal 56, 777–783.

doi:10.2136/sssaj1992.03615995005600030017x

Cantoran, A., Maillard, F., Baldrian, P., Kennedy, P.G., 2023. Defining a core microbial necrobiome
associated with decomposing fungal necromass. FEMS Microbiology Ecology 99, fiad098.
doi:10.1093/femsec/fiad098

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M.,
Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R., 2012. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. The
ISME Journal 6, 1621–1624. doi:10.1038/ismej.2012.8

Chevan, A., Sutherland, M., 1991. Hierarchical Partitioning. The American Statistician 45, 90–96.
doi:10.1080/00031305.1991.10475776

Clemmensen, K.E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J.,
Finlay, R.D., Wardle, D.A., Lindahl, B.D., 2013. Roots and Associated Fungi Drive Long-Term
Carbon Sequestration in Boreal Forest. Science 339, 1615–1618. doi:10.1126/science.1231923

Cocozza, C., D'Orazio, V., Miano, T.M., Shotyk, W., 2003. Characterization of solid and aqueous phases
of a peat bog profile using molecular fluorescence spectroscopy, ESR and FT-IR, and comparison
with physical properties. Organic Geochemistry 34, 49–60. doi:10.1016/S0146-6380(02)00208-5

Comeau, L.-P., Lemke, R.L., Knight, J.D., Bedard-Haughn, A., 2013. Carbon input from 13C-labeled
crops in four soil organic matter fractions. Biology and Fertility of Soils 49, 1179–1188.
doi:10.1007/s00374-013-0816-4

Cotrufo, M.F., Lavallee, J.M., 2022. Soil organic matter formation, persistence, and functioning: A
synthesis of current understanding to inform its conservation and regeneration, in: Advances in
Agronomy. Elsevier, pp. 1–66. doi:10.1016/bs.agron.2021.11.002

Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H., Parton, W.J., 2015.
Formation of soil organic matter via biochemical and physical pathways of litter mass loss.
Nature Geoscience 8, 776–779. doi:10.1038/ngeo2520

Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2013. The M icrobial E fficiency⬚ M
atrix S tabilization (MEMS) framework integrates plant litter decomposition with soil organic

926  matter stabilization: do labile plant inputs form stable soil organic matter? Global Change

927   Biology 19, 988–995. doi:10.1111/gcb.12113

928 Craig, M.E., Geyer, K.M., Beidler, K.V., Brzostek, E.R., Frey, S.D., Stuart Grandy, A., Liang, C.,

929   Phillips, R.P., 2022. Fast-decaying plant litter enhances soil carbon in temperate forests but not

930   through microbial physiological traits. Nature Communications 13, 1229. doi:10.1038/s41467-

931   022-28715-9

932 Deng, Y., Dixon, J.B., 2018. Soil Organic Matter and Organic-Mineral Interactions, in: Dixon, J.B.,

933   Schulze, D.G. (Eds.), SSSA Book Series. Soil Science Society of America, Madison, WI, USA,

934   pp. 69–107. doi:10.2136/sssabookser7.c3

935 Domeignoz-Horta, L.A., Shinfuku, M., Junier, P., Poirier, S., Verrecchia, E., Sebag, D., DeAngelis, K.M.,

936   2021. Direct evidence for the role of microbial community composition in the formation of soil

937   organic matter composition and persistence. ISME Communications 1, 64. doi:10.1038/s43705-

938   021-00071-7

939 Drever, M.C., Douglas, T.J., Domalik, A.D., Balke, E., Scott, D.C., Johnson, M.S., Gurd, D.B., 2023.

940   Water diversion structures and river discharge dictate salinity dynamics on tidal flats of the Fraser

941   River delta, British Columbia. Estuarine, Coastal and Shelf Science 291, 108436.

942   doi:10.1016/j.ecss.2023.108436

943 Dzurendova, S., Zimmermann, B., Kohler, A., Tafintseva, V., Slany, O., Certik, M., Shapaval, V., 2020.

944   Microcultivation and FTIR spectroscopy-based screening revealed a nutrient-induced co-

945   production of high-value metabolites in oleaginous Mucoromycota fungi. PLOS ONE 15,

946   e0234870. doi:10.1371/journal.pone.0234870

947 Egli, M., Mirabella, A., Kägi, B., Tomasone, R., Colorio, G., 2006. Influence of steam sterilisation on soil

948   chemical characteristics, trace metals and clay mineralogy. Geoderma 131, 123–142.

949   doi:10.1016/j.geoderma.2005.03.017

950 Ekblad, A., Wallander, H., Godbold, D.L., Cruz, C., Johnson, D., Baldrian, P., Björk, R.G., Epron, D.,

951   Kieliszewska-Rokicka, B., Kjøller, R., Kraigher, H., Matzner, E., Neumann, J., Plassard, C.,

2013. The production and turnover of extramatrical mycelium of ectomycorrhizal fungi in forest

soils: role in carbon cycling. Plant and Soil 366, 1–27. doi:10.1007/s11104-013-1630-3

Ellerbrock, R.H., Kaiser, M., 2005. Stability and composition of different soluble soil organic matter

fractions–evidence from δ13C and FTIR signatures. Geoderma 128, 28–37.

doi:10.1016/j.geoderma.2004.12.025

Fehrer, J., Réblová, M., Bambasová, V., Vohník, M., 2019. The root-symbiotic *Rhizoscyphus ericae*

aggregate and *Hyaloscypha* (*Leotiomycetes*) are congeneric: Phylogenetic and experimental

evidence. Studies in Mycology 92, 195–225. doi:10.1016/j.simyco.2018.10.004

Fernandez, C.W., Heckman, K., Kolka, R., Kennedy, P.G., 2019. Melanin mitigates the accelerated decay

of mycorrhizal necromass with peatland warming. Ecology Letters 22, 498–505.

doi:10.1111/ele.13209

Fernandez, C.W., Kennedy, P.G., 2018. Melanization of mycorrhizal fungal necromass structures

microbial decomposer communities. Journal of Ecology 106, 468–479. doi:10.1111/1365-

2745.12920

Fernandez, C.W., Koide, R.T., 2014. Initial melanin and nitrogen concentrations control the

decomposition of ectomycorrhizal fungal litter. Soil Biology and Biochemistry 77, 150–157.

doi:10.1016/j.soilbio.2014.06.026

Fernandez, C.W., Koide, R.T., 2012. The role of chitin in the decomposition of ectomycorrhizal fungal

litter. Ecology 93, 24–28. doi:10.1890/11-1346.1

Fernandez, C.W., Langley, J.A., Chapman, S., McCormack, M.L., Koide, R.T., 2016. The decomposition

of ectomycorrhizal fungal necromass. Soil Biology and Biochemistry 93, 38–49.

doi:10.1016/j.soilbio.2015.10.017

Fernandez, C.W., McCormack, M.L., Hill, J.M., Pritchard, S.G., Koide, R.T., 2013. On the persistence of

Cenococcum geophilum ectomycorrhizas and its implications for forest carbon and nutrient

cycles. Soil Biology and Biochemistry 65, 141–143. doi:10.1016/j.soilbio.2013.05.022

Fierer, N., Schimel, J.P., 2002. Effects of drying–rewetting frequency on soil carbon and nitrogen

transformations. Soil Biology and Biochemistry 34, 777–787. doi:10.1016/S0038-

0717(02)00007-X

Fogarty, R.V., Tobin, J.M., 1996. Fungal melanins and their interactions with metals. Enzyme and

Microbial Technology 19, 311–317. doi:10.1016/0141-0229(96)00002-6

Frey, S.D., 2019. Mycorrhizal Fungi as Mediators of Soil Organic Matter Dynamics. Annual Review of

Ecology, Evolution, and Systematics 50, 237–259. doi:10.1146/annurev-ecolsys-110617-062331

Fu, X., Xie, M., Lu, M., Shi, L., Shi, T., Yu, M., 2022. Characterization of the physicochemical

properties, antioxidant activity, and antiproliferative activity of natural melanin from S. reiliana.

Scientific Reports 12, 2110. doi:10.1038/s41598-022-05676-z

Godbold, D.L., Hoosbeek, M.R., Lukac, M., Cotrufo, M.F., Janssens, I.A., Ceulemans, R., Polle, A.,

Velthorst, E.J., Scarascia-Mugnozza, G., De Angelis, P., Miglietta, F., Peressotti, A., 2006.

Mycorrhizal Hyphal Turnover as a Dominant Process for Carbon Input into Soil Organic Matter.

Plant and Soil 281, 15–24. doi:10.1007/s11104-005-3701-6

Golchin, A., Clarke, P., Oades, J.M., 1996. The heterogeneous nature of microbial products as shown by

solid-state13C CP/MAS NMR spectroscopy. Biogeochemistry 34, 71–97.

doi:10.1007/BF02180974

Grelet, G., Johnson, D., Paterson, E., Anderson, I.C., Alexander, I.J., 2009. Reciprocal carbon and

nitrogen transfer between an ericaceous dwarf shrub and fungi isolated from *Piceirhiza*

bicolorata ectomycorrhizas. New Phytologist 182, 359–366. doi:10.1111/j.1469-

8137.2009.02813.x

Harki, E., Talou, T., Dargent, R., 1997. Purification, characterisation and analysis of melanin extracted

from Tuber melanosporum Vitt. Food Chemistry 58, 69–73. doi:10.1016/S0308-8146(96)00215-

4

Harsh, J., Chorover, J., Nizeyimana, E., 2018. Allophane and Imogolite, in: Dixon, J.B., Schulze, D.G.

(Eds.), SSSA Book Series. Soil Science Society of America, Madison, WI, USA, pp. 291–322.

doi:10.2136/sssabookser7.c9

Heckman, K.A., Possinger, A.R., Badgley, B.D., Bowman, M.M., Gallo, A.C., Hatten, J.A., Nave, L.E.,

SanClements, M.D., Swanston, C.W., Weiglein, T.L., Wieder, W.R., Strahm, B.D., 2023.

Moisture-driven divergence in mineral-associated soil carbon persistence. Proceedings of the

National Academy of Sciences 120, e2210044120. doi:10.1073/pnas.2210044120

Hu, Y., Zheng, Q., Noll, L., Zhang, S., Wanek, W., 2020. Direct measurement of the in situ

decomposition of microbial-derived soil organic matter. Soil Biology and Biochemistry 141,

107660. doi:10.1016/j.soilbio.2019.107660

Jagadamma, S., Mayes, M.A., Phillips, J.R., 2012. Selective Sorption of Dissolved Organic Carbon

Compounds by Temperate Soils. PLoS ONE 7, e50434. doi:10.1371/journal.pone.0050434

Jia, W., Zheng, T., Zhao, Y., Deng, F., Yang, Y., Liang, C., He, H., Zhang, X., 2023. Nitrogen application

influences the effect of bacteria on the belowground allocation of photosynthesized carbon under

elevated CO2. Soil Biology and Biochemistry 180, 109021. doi:10.1016/j.soilbio.2023.109021

Jobbágy, E.G., Jackson, R.B., 2000. The Vertical Distribution of Soil Organic Carbon and Its Relation to

Climate and Vegetation. Ecological Applications 10, 423–436. doi:10.1890/1051-

0761(2000)010[0423:TVDOSO]2.0.CO;2

Kalbitz, K., Schwesig, D., Rethemeyer, J., Matzner, E., 2005. Stabilization of dissolved organic matter by

sorption to the mineral soil. Soil Biology and Biochemistry 37, 1319–1331.

doi:10.1016/j.soilbio.2004.11.028

Kallenbach, C.M., Frey, S.D., Grandy, A.S., 2016. Direct evidence for microbial-derived soil organic

matter formation and its ecophysiological controls. Nature Communications 7, 13630.

doi:10.1038/ncomms13630

Keiluweit, M., Bougoure, J.J., Zeglin, L.H., Myrold, D.D., Weber, P.K., Pett-Ridge, J., Kleber, M., Nico,

P.S., 2012. Nano-scale investigation of the association of microbial nitrogen residues with iron

(hydr)oxides in a forest soil O-horizon. Geochimica et Cosmochimica Acta 95, 213–226. doi:10.1016/j.gca.2012.07.001

Kennedy, P.G., Maillard, F., 2023. Knowns and unknowns of the soil fungal necrobiome. Trends in Microbiology 31, 173–180. doi:10.1016/j.tim.2022.08.011

Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., Nico, P.S., 2015. Mineral–Organic Associations: Formation, Properties, and Relevance in Soil Environments, in: Advances in Agronomy. Elsevier, pp. 1–140. doi:10.1016/bs.agron.2014.10.005

Kleber, M., Sollins, P., Sutton, R., 2007. A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces. Biogeochemistry 85, 9–24. doi:10.1007/s10533-007-9103-5

Klink, S., Keller, A.B., Wild, A.J., Baumert, V.L., Gube, M., Lehndorff, E., Meyer, N., Mueller, C.W., Phillips, R.P., Pausch, J., 2022. Stable isotopes reveal that fungal residues contribute more to mineral-associated organic matter pools than plant residues. Soil Biology and Biochemistry 168, 108634. doi:10.1016/j.soilbio.2022.108634

Kögel-Knabner, I., 2002. The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. Soil Biology and Biochemistry 34, 139–162. doi:10.1016/S0038-0717(01)00158-4

Krumina, L., Lyngsie, G., Tunlid, A., Persson, P., 2017. Oxidation of a Dimethoxyhydroquinone by Ferrihydrite and Goethite Nanoparticles: Iron Reduction versus Surface Catalysis. Environmental Science & Technology 51, 9053–9061. doi:10.1021/acs.est.7b02292

Lai, J., Zou, Y., Zhang, J., Peres-Neto, P., 2021. rdacca.hp: an R package for generalizing hierarchical and variation partitioning in multiple regression and canonical analysis. doi:10.1101/2021.03.09.434308

Langseter, A.M., Dzurendova, S., Shapaval, V., Kohler, A., Ekeberg, D., Zimmermann, B., 2021. Evaluation and optimisation of direct transesterification methods for the assessment of lipid

accumulation in oleaginous filamentous fungi. Microbial Cell Factories 20, 59. doi:10.1186/s12934-021-01542-1

Lee, Z.M.-P., Bussema, C., Schmidt, T.M., 2009. rrnDB: documenting the number of rRNA and tRNA genes in bacteria and archaea. Nucleic Acids Research 37, D489–D493. doi:10.1093/nar/gkn689

Lehmann, J., Hansel, C.M., Kaiser, C., Kleber, M., Maher, K., Manzoni, S., Nunan, N., Reichstein, M., Schimel, J.P., Torn, M.S., Wieder, W.R., Kögel-Knabner, I., 2020. Persistence of soil organic carbon caused by functional complexity. Nature Geoscience 13, 529–534. doi:10.1038/s41561-020-0612-3

Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. Nature 528, 60–68. doi:10.1038/nature16069

Lenth, R.V., 2017. emmeans: Estimated Marginal Means, aka Least-Squares Means. doi:10.32614/CRAN.package.emmeans

Liang, C., Amelung, W., Lehmann, J., Kästner, M., 2019. Quantitative assessment of microbial necromass contribution to soil organic matter. Global Change Biology 25, 3578–3590. doi:10.1111/gcb.14781

Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial control over soil carbon storage. Nature Microbiology 2, 17105. doi:10.1038/nmicrobiol.2017.105

Lützow, M.V., Kögel￿Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., Flessa, H., 2006. Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions – a review. European Journal of Soil Science 57, 426–445. doi:10.1111/j.1365-2389.2006.00809.x

Mac Nally, R., 2002. Multiple regression and inference in ecology and conservation biology: further comments on identifying important predictor variables. Biodiversity and Conservation 11, 1397–1401. doi:10.1023/A:1016250716679

Maillard, F., Beatty, B., Park, M., Adamczyk, S., Adamczyk, B., See, C.R., Cavender-Bares, J., Hobbie, S.E., Kennedy, P.G., 2023a. Microbial community attributes supersede plant and soil parameters

in predicting fungal necromass decomposition rates in a 12-tree species common garden

experiment. Soil Biology and Biochemistry 184, 109124. doi:10.1016/j.soilbio.2023.109124

Maillard, F., Michaud, T.J., See, C.R., DeLancey, L.C., Blazewicz, S.J., Kimbrel, J.A., Pett-Ridge, J.,

Kennedy, P.G., 2023b. Melanization slows the rapid movement of fungal necromass carbon and

nitrogen into both bacterial and fungal decomposer communities and soils. mSystems e00390-23.

doi:10.1128/msystems.00390-23

Maillard, F., Pflender, S., Heckman, K.A., Chalot, M., Kennedy, P.G., 2023c. Fungal necromass presents

a high potential for Mercury immobilization in soil. Chemosphere 311, 136994.

doi:10.1016/j.chemosphere.2022.136994

Maillard, F., Schilling, J., Andrews, E., Schreiner, K.M., Kennedy, P., 2020. Functional convergence in

the decomposition of fungal necromass in soil and wood. FEMS Microbiology Ecology 96,

fiz209. doi:10.1093/femsec/fiz209

Manzoni, S., Čapek, P., Porada, P., Thurner, M., Winterdahl, M., Beer, C., Brüchert, V., Frouz, J.,

Herrmann, A.M., Lindahl, B.D., Lyon, S.W., Šantrůčková, H., Vico, G., Way, D., 2018. Reviews

and syntheses: Carbon use efficiency from organisms to ecosystems – definitions, theories, and

empirical evidence. Biogeosciences 15, 5929–5949. doi:10.5194/bg-15-5929-2018

Matus, F.J., 2021. Fine silt and clay content is the main factor defining maximal C and N accumulations

in soils: a meta-analysis. Scientific Reports 11, 6438. doi:10.1038/s41598-021-84821-6

Mikutta, R., Mikutta, C., Kalbitz, K., Scheel, T., Kaiser, K., Jahn, R., 2007. Biodegradation of forest floor

organic matter bound to minerals via different binding mechanisms. Geochimica et

Cosmochimica Acta 71, 2569–2590. doi:10.1016/j.gca.2007.03.002

Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V.,

Chen, Z.-S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley, C.B., Hong, S.Y., Mandal,

B., Marchant, B.P., Martin, M., McConkey, B.G., Mulder, V.L., O'Rourke, S., Richer-de-Forges,

A.C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V.,

Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., Van Wesemael, B., Winowiecki, L.,

2017. Soil carbon 4 per mille. Geoderma 292, 59–86. doi:10.1016/j.geoderma.2017.01.002

Nemergut, D.R., Knelman, J.E., Ferrenberg, S., Bilinski, T., Melbourne, B., Jiang, L., Violle, C., Darcy,

J.L., Prest, T., Schmidt, S.K., Townsend, A.R., 2016. Decreases in average bacterial community

rRNA operon copy number during succession. The ISME Journal 10, 1147–1156.

doi:10.1038/ismej.2015.191

Netherway, T., Bahram, M., 2024. Melanized root-associated fungi: key players in plant–soil systems.

Trends in Microbiology S0966842X24001690. doi:10.1016/j.tim.2024.06.006

Nguyen, N.H., Song, Z., Bates, S.T., Branco, S., Tedersoo, L., Menke, J., Schilling, J.S., Kennedy, P.G.,

2016. FUNGuild: An open annotation tool for parsing fungal community datasets by ecological

guild. Fungal Ecology 20, 241–248. doi:10.1016/j.funeco.2015.06.006

Niemeyer, J., Chen, Y., Bollag, J. M., 1992. Characterization of Humic Acids, Composts, and Peat by

Diffuse Reflectance Fourier Transform Infrared Spectroscopy. Soil Science Society of America

Journal 56, 135–140. doi:10.2136/sssaj1992.03615995005600010021x

Norén, K., Loring, J.S., Persson, P., 2008. Adsorption of alpha amino acids at the water/goethite interface.

Journal of Colloid and Interface Science 319, 416–428. doi:10.1016/j.jcis.2007.11.046

Novak, J.K., Kennedy, P.G., Gardner, J.G., 2024. Transcriptomic analyses of bacterial growth on fungal

necromass reveal different microbial community niches during degradation. Applied and

Environmental Microbiology e01062-24. doi:10.1128/aem.01062-24

Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B.,

Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B.,

Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista, H.B.A.,

FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.O., Lahti, L., McGlinn, D.,

Ouellette, M.-H., Ribeiro Cunha, E., Smith, T., Stier, A., Ter Braak, C.J.F., Weedon, J., 2001.

vegan: Community Ecology Package. doi:10.32614/CRAN.package.vegan

Omoike, A., Chorover, J., 2006. Adsorption to goethite of extracellular polymeric substances from

1129   Bacillus subtilis. Geochimica et Cosmochimica Acta 70, 827–838. doi:10.1016/j.gca.2005.10.012

1130   Palmer, J.M., Jusino, M.A., Banik, M.T., Lindner, D.L., 2018. Non-biological synthetic spike-in controls

1131   and the AMPtk software pipeline improve mycobiome data. PeerJ 6, e4925.

1132   doi:10.7717/peerj.4925

1133   Pärnpuu, S., Astover, A., Tõnutare, T., Penu, P., Kauer, K., 2022. Soil organic matter qualification with

1134   FTIR spectroscopy under different soil types in Estonia. Geoderma Regional 28, e00483.

1135   doi:10.1016/j.geodrs.2022.e00483

1136   Põlme, S., Abarenkov, K., Henrik Nilsson, R., Lindahl, B.D., Clemmensen, K.E., Kauserud, H., Nguyen,

1137   N., Kjøller, R., Bates, S.T., Baldrian, P., Frøslev, T.G., Adojaan, K., Vizzini, A., Suija, A.,

1138   Pfister, D., Baral, H.-O., Järv, H., Madrid, H., Nordén, J., Liu, J.-K., Pawlowska, J., Põldmaa, K.,

1139   Pärtel, K., Runnel, K., Hansen, K., Larsson, K.-H., Hyde, K.D., Sandoval-Denis, M., Smith,

1140   M.E., Toome-Heller, M., Wijayawardene, N.N., Menolli, N., Reynolds, N.K., Drenkhan, R.,

1141   Maharachchikumbura, S.S.N., Gibertoni, T.B., Læssøe, T., Davis, W., Tokarev, Y., Corrales, A.,

1142   Soares, A.M., Agan, A., Machado, A.R., Argüelles-Moyao, A., Detheridge, A., De Meiras-

1143   Ottoni, A., Verbeken, A., Dutta, A.K., Cui, B.-K., Pradeep, C.K., Marín, C., Stanton, D., Gohar,

1144   D., Wanasinghe, D.N., Otsing, E., Aslani, F., Griffith, G.W., Lumbsch, T.H., Grossart, H.-P.,

1145   Masigol, H., Timling, I., Hiiesalu, I., Oja, J., Kupagme, J.Y., Geml, J., Alvarez-Manjarrez, J.,

1146   Ilves, K., Loit, K., Adamson, K., Nara, K., Küngas, K., Rojas-Jimenez, K., Bitenieks, K., Irinyi,

1147   L., Nagy, L.G., Soonvald, L., Zhou, L.-W., Wagner, L., Aime, M.C., Öpik, M., Mujica, M.I.,

1148   Metsoja, M., Ryberg, M., Vasar, M., Murata, M., Nelsen, M.P., Cleary, M., Samarakoon, M.C.,

1149   Doilom, M., Bahram, M., Hagh-Doust, N., Dulya, O., Johnston, P., Kohout, P., Chen, Q., Tian,

1150   Q., Nandi, R., Amiri, R., Perera, R.H., Dos Santos Chikowski, R., Mendes-Alvarenga, R.L.,

1151   Garibay-Orijel, R., Gielen, R., Phookamsak, R., Jayawardena, R.S., Rahimlou, S., Karunarathna,

1152   S.C., Tibpromma, S., Brown, S.P., Sepp, S.-K., Mundra, S., Luo, Z.-H., Bose, T., Vahter, T.,

1153   Netherway, T., Yang, T., May, T., Varga, T., Li, W., Coimbra, V.R.M., De Oliveira, V.R.T., De

Lima, V.X., Mikryukov, V.S., Lu, Y., Matsuda, Y., Miyamoto, Y., Kõljalg, U., Tedersoo, L., 2020. FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. Fungal Diversity 105, 1–16. doi:10.1007/s13225-020-00466-2

Rasmussen, C., Heckman, K., Wieder, W.R., Keiluweit, M., Lawrence, C.R., Berhe, A.A., Blankinship, J.C., Crow, S.E., Druhan, J.L., Hicks Pries, C.E., Marin-Spiotta, E., Plante, A.F., Schädel, C., Schimel, J.P., Sierra, C.A., Thompson, A., Wagai, R., 2018. Beyond clay: towards an improved set of variables for predicting soil organic matter content. Biogeochemistry 137, 297–306. doi:10.1007/s10533-018-0424-3

Rillig, M.C., Caldwell, B.A., Wösten, H.A.B., Sollins, P., 2007. Role of proteins in soil carbon and nitrogen storage: controls on persistence. Biogeochemistry 85, 25–44. doi:10.1007/s10533-007-9102-6

Ross, G.L., Wang, C., 1993. Extractable Al, Fe, Mn and Si, in: Soil Sampling and Methods of Analysis. Lewis Publication, Boca Raton, pp. 239–246.

Ryan, M.E., Schreiner, K.M., Swenson, J.T., Gagne, J., Kennedy, P.G., 2020. Rapid changes in the chemical composition of degrading ectomycorrhizal fungal necromass. Fungal Ecology 45, 100922. doi:10.1016/j.funeco.2020.100922

Schweigert, M., Herrmann, S., Miltner, A., Fester, T., Kästner, M., 2015. Fate of ectomycorrhizal fungal biomass in a soil bioreactor system and its contribution to soil organic matter formation. Soil Biology and Biochemistry 88, 120–127. doi:10.1016/j.soilbio.2015.05.012

See, C.R., Fernandez, C.W., Conley, A.M., DeLancey, L.C., Heckman, K.A., Kennedy, P.G., Hobbie, S.E., 2021. Distinct carbon fractions drive a generalisable two pool model of fungal necromass decomposition. Functional Ecology 35, 796–806. doi:10.1111/1365-2435.13728

See, C.R., Keller, A.B., Hobbie, S.E., Kennedy, P.G., Weber, P.K., Pett Ridge, J., 2022. Hyphae move matter and microbes to mineral microsites: Integrating the hyphosphere into conceptual models of soil organic matter stabilization. Global Change Biology 28, 2527–2540. doi:10.1111/gcb.16073

Siletti, C.E., Zeiner, C.A., Bhatnagar, J.M., 2017. Distributions of fungal melanin across species and

soils. Soil Biology and Biochemistry 113, 285–293. doi:10.1016/j.soilbio.2017.05.030

Simpson, A.J., Simpson, M.J., Smith, E., Kelleher, B.P., 2007. Microbially Derived Inputs to Soil

Organic Matter: Are Current Estimates Too Low? Environmental Science & Technology 41,

8070–8076. doi:10.1021/es071217x

Sokol, N.W., Bradford, M.A., 2019. Microbial formation of stable soil carbon is more efficient from

belowground than aboveground input. Nature Geoscience 12, 46–53. doi:10.1038/s41561-018-

0258-6

Sollins, P., Homann, P., Caldwell, B.A., 1996. Stabilization and destabilization of soil organic matter:

mechanisms and controls. Geoderma 74, 65–105. doi:10.1016/S0016-7061(96)00036-5

Stoddard, S.F., Smith, B.J., Hein, R., Roller, B.R.K., Schmidt, T.M., 2015. rrnDB: improved tools for

interpreting rRNA gene abundance in bacteria and archaea and a new foundation for future

development. Nucleic Acids Research 43, D593–D598. doi:10.1093/nar/gku1201

Stuart, B.H., 2004. Infrared Spectroscopy: Fundamentals and Applications, 1st ed, Analytical Techniques

in the Sciences. Wiley. doi:10.1002/0470011149

Štursová, M., Žifčáková, L., Leigh, M.B., Burgess, R., Baldrian, P., 2012. Cellulose utilization in forest

litter and soil: identification of bacterial and fungal decomposers. FEMS Microbiology Ecology

80, 735–746. doi:10.1111/j.1574-6941.2012.01343.x

Sulman, B.N., Phillips, R.P., Oishi, A.C., Shevliakova, E., Pacala, S.W., 2014. Microbe-driven turnover

offsets mineral-mediated storage of soil carbon under elevated CO_2. Nature Climate Change 4,

1099–1102. doi:10.1038/nclimate2436

Taylor, D.L., Walters, W.A., Lennon, N.J., Bochicchio, J., Krohn, A., Caporaso, J.G., Pennanen, T.,

2016. Accurate Estimation of Fungal Diversity and Abundance through Improved Lineage-

Specific Primers Optimized for Illumina Amplicon Sequencing. Applied and Environmental

Microbiology 82, 7217–7226. doi:10.1128/AEM.02576-16

Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., Ruiz, L.V., Vasco-Palacios, A.M., Thu, P.Q., Suija, A., Smith, M.E., Sharp, C., Saluveer, E., Saitta, A., Rosas, M., Riit, T., Ratkowsky, D., Pritsch, K., Põldmaa, K., Piepenbring, M., Phosri, C., Peterson, M., Parts, K., Pärtel, K., Otsing, E., Nouhra, E., Njouonkou, A.L., Nilsson, R.H., Morgado, L.N., Mayor, J., May, T.W., Majuakim, L., Lodge, D.J., Lee, S.S., Larsson, K.-H., Kohout, P., Hosaka, K., Hiiesalu, I., Henkel, T.W., Harend, H., Guo, L., Greslebin, A., Grelet, G., Geml, J., Gates, G., Dunstan, W., Dunk, C., Drenkhan, R., Dearnaley, J., De Kesel, A., Dang, T., Chen, X., Buegger, F., Brearley, F.Q., Bonito, G., Anslan, S., Abell, S., Abarenkov, K., 2014. Global diversity and geography of soil fungi. Science 346, 1256688. doi:10.1126/science.1256688

Theng, B.K.G., 2024. The Chemistry of Clay-Organic Reactions, 2nd ed. CRC Press, Boca Raton. doi:10.1201/9781003080244

Tian, Z., Wang, T., Tunlid, A., Persson, P., 2020. Proteolysis of Iron Oxide-Associated Bovine Serum Albumin. Environmental Science & Technology 54, 5121–5130. doi:10.1021/acs.est.0c00860

Van Der Wal, A., Bloem, J., Mulder, C., De Boer, W., 2009. Relative abundance and activity of melanized hyphae in different soil ecosystems. Soil Biology and Biochemistry 41, 417–419. doi:10.1016/j.soilbio.2008.10.031

Veresoglou, S.D., Rillig, M.C., Johnson, D., 2018. Responsiveness of plants to mycorrhiza regulates coexistence. Journal of Ecology 106, 1864–1875. doi:10.1111/1365-2745.13026

Villacís García, M., Ugalde Arzate, M., Vaca Escobar, K., Villalobos, M., Zanella, R., Martínez Villegas, N., 2015. Laboratory synthesis of goethite and ferrihydrite of controlled particle sizes. Boletín de La Sociedad Geológica Mexicana 67, 433–446. doi:10.18268/BSGM2015v67n3a7

Wang, B., An, S., Liang, C., Liu, Y., Kuzyakov, Y., 2021. Microbial necromass as the source of soil organic carbon in global ecosystems. Soil Biology and Biochemistry 162, 108422. doi:10.1016/j.soilbio.2021.108422

Wang, C., Kuzyakov, Y., 2024. Soil organic matter priming: The pH effects. Global Change Biology 30, e17349. doi:10.1111/gcb.17349

Wang, C., Wang, X., Pei, G., Xia, Z., Peng, B., Sun, L., Wang, J., Gao, D., Chen, S., Liu, D., Dai, W.,

Jiang, P., Fang, Y., Liang, C., Wu, N., Bai, E., 2020. Stabilization of microbial residues in soil

organic matter after two years of decomposition. Soil Biology and Biochemistry 141, 107687.

doi:10.1016/j.soilbio.2019.107687

Wang, Q., Cole, J.R., 2024. Updated RDP taxonomy and RDP Classifier for more accurate taxonomic

classification. Microbiology Resource Announcements 13, e01063-23. doi:10.1128/mra.01063-23

Wang, Xu, Wang, C., Fan, X., Sun, L., Sang, C., Wang, Xugao, Jiang, P., Fang, Y., Bai, E., 2024.

Mineral composition controls the stabilization of microbially derived carbon and nitrogen in soils:

Insights from an isotope tracing model. Global Change Biology 30, e17156.

doi:10.1111/gcb.17156

Warren, C., 2021. What are the products of enzymatic cleavage of organic N? Soil Biology and

Biochemistry 154, 108152. doi:10.1016/j.soilbio.2021.108152

West, J.R., Lauer, J.G., Whitman, T., 2023. Tillage homogenizes soil bacterial communities in

microaggregate fractions by facilitating dispersal. Soil Biology and Biochemistry 186, 109181.

doi:10.1016/j.soilbio.2023.109181

Whalen, E.D., Grandy, A.S., Sokol, N.W., Keiluweit, M., Ernakovich, J., Smith, R.G., Frey, S.D., 2022.

Clarifying the evidence for microbial⎕ and plant⎕derived soil organic matter, and the path

toward a more quantitative understanding. Global Change Biology 28, 7167–7185.

doi:10.1111/gcb.16413

Wieder, W.R., Grandy, A.S., Kallenbach, C.M., Bonan, G.B., 2014. Integrating microbial physiology and

physio-chemical principles in soils with the MIcrobial-MIneral Carbon Stabilization (MIMICS)

model. Biogeosciences 11, 3899–3917. doi:10.5194/bg-11-3899-2014

Wu, S., Konhauser, K.O., Chen, B., Huang, L., 2023. "Reactive Mineral Sink" drives soil organic matter

dynamics and stabilization. Npj Materials Sustainability 1, 3. doi:10.1038/s44296-023-00003-7

1254

1255

1256

1257

1258

1259

1260

1261   Supplemental Information

1262

1263   **Table S1**. Initial necromass chemistry (mean, SD, n = 3) for high and low melanin *Hyaloscypha bicolor*

1264   necromass types before (non-sterile) and after autoclaving (sterile).

Necromass Type	High Melanin		Low Melanin	
Initial	Non-Sterile	Sterile[*]	Non-Sterile	Sterile[*]
% C	43.3 (1.3)[a]	48.2 (9.9)[a]	51.4 (0.33)[b]	48.3 (0.90)[c]
% N	3.28 (0.18)[a]	3.94 (0.61)[a]	3.93 (0.06)[a]	3.62 (0.11)[a]
‰ δ ^{13}C	31.4 (1.2)[a]	31.7 (1.3)[a]	35.5 (0.90)[b]	36.39 (0.36)[b]
‰ δ ^{15}N	680.4 (21)[a]	849.6 (140)[a]	731.8 (23)[a]	728.8 (8.1)[a]
% Soluble fraction	76.2 (1.2)[a]	---------------	77.1 (7.5)[a]	---------------
% Acid unhydrolyzable fraction	18.1 (3.4)[a]	---------------	16.7 (4.1)[a]	---------------

1265

1266   *Sterile necromass was added to sterile soil microcosms in 38-day necromass incubations. There was not

1267   enough initial autoclaved necromass available to measure initial C quality fractions.

1268

1269

1270   **Table S2.** Site locations and overstory tree species for Indiana Department of Natural Resources (IDNR)

1271   Continuous Forest Inventory (CFI) plots selected for soil collection.

Latitude	Longitude	Site ID	County	Plot ID	Sampling Date	Soil Series	Tree Species
38.979306	-87.255861	C153	Sullivan	1619	6/5/20	Stendal silt loam	*Pinus virginiana* & *Robinia pseudoacacia*
38.988417	-87.230556	C55	Greene	1601	6/1/20	Ava silt loam	*Acer rubrum, Asimina triloba, Cornus florida, Morus rubra, Prunus serotina,* & *Quercus ellipsoidalis*
40.812436	-85.714392	C169	Wabash	13	5/21/20	Blount silt loam	*Celtis occidentalis, Morus alba, Pinus resinosa, Prunus serotina, Quercus muehlenbergii,* & *Ulmus americana*
38.801075	-86.056	C71	Jackson	1859	5/22/20	Brownstown channery silt loam	*Acer saccharum, Paulownia tomentosa,* & *Liriodendron tulipifera*
39.238972	-86.314778	C13	Brown	856	5/26/20	Berks-Trevlac-Wellston complex	*Acer rubrum, Acer saccharum, Fagus grandifolia, Sassafras albidum,* & *Quercus velutina*
38.170672	-86.252908	C61	Harrison	3667	7/24/20	Caneyville-Rock outcrop complex	*Acer saccharum, Nyssa sylvatica,* & *Quercus alba*

Table S3. Mineral-associated organic matter peak annotations for diffuse reflectance infrared Fourier transform spectroscopy (DRIFTS).

Wavenumber (cm^{-1})	Peak ID Number	Functional Group	Potential Compound	Alternative Functional Group	Alternative Compound
3695	1	O-H	Clay, minerals	Al-OH	Aluminum, minerals
3650	2	O-H	Clay, minerals	Al-OH	Aluminum, minerals
3621	3	O-H	Clay, minerals	Al-OH	Aluminum, minerals
1683	4	C=O	Carboxylic acids	C=O	Proteins
1609	5	C=C	Aromatic	N-H	Proteins
1521	6	C=C	Aromatic		
1493	7	C=C	Aromatic	N-H	Proteins
1384	8	C-H	Aromatic	C-H	Phenolic
1284	9	C-O	Phenolic		
1159	10	C-O	Polysaccharides	Si-O	Silicate, minerals
1104	11	C-O	Polysaccharides	Si-O	Silicate, minerals
1033	12	C-O	Polysaccharides	Si-O	Silicate, minerals
1013	13	C-O	Polysaccharides	Si-O	Silicate, minerals
940	14	C-H	Aromatic	C-H	Alkenes
916	15	C-H	Aromatic	C-H	Alkenes
798	16	C-H	Aromatic	C-H	Alkenes
779	17	C-H	Aromatic	C-H	Alkenes
695	18	Si-O	Silicate, minerals		
647	19	Si-O	Silicate, minerals		
528	20	Si-O	Silicate, minerals		
474	21	Si-O	Silicate, minerals		

Table S4. Site means and standard deviations for fungal necromass-derived mineral-associated organic matter (MAOM) carbon (C) and nitrogen (N) formation in live soil microcosms. Letters denote differences in site means as determined by ANOVA.

Site ID	Fungal Derived MAOM-C (% initial necromass C)	Fungal Derived MAOM-N (% initial necromass N)
C153	6.92 (1.1)[a]	40.1 (4.2)[b]
C55	6.12 (0.72)[ab]	48.0 (5.9)[a]
C169	5.82 (0.48)[abc]	44.5 (3.5)[ab]
C71	4.72 (0.47)[c]	40.1 (5.6)[ab]
C13	5.36 (0.45)[bc]	40.6 (2.6)[ab]
C61	5.83 (0.94)[abc]	39.6 (4.3)[b]

Table S5. Necromass-derived mineral-associated organic matter (MAOM) carbon (C) and nitrogen (N) formation (mean, SD, n = 3) in sterile soil microcosms. Letters denote differences in site means as determined by ANOVA.

Site ID	Sterilization Treatment	Necromass Type	Fungal Derived MAOM-C (% initial necromass C)	Fungal Derived MAOM-N (% initial necromass N)
C153	Sterile	High Melanin	11.7 (2.8)[a]	30.2 (7.5)[a]
C153	Sterile	Low Melanin	11.5 (3.4)[a]	30.2 (7.6)[a]
C169	Sterile	High Melanin	7.92 (0.86)[b]	28.5 (1.9)[a]
C169	Sterile	Low Melanin	6.87 (1.1)[b]	39.1 (4.1)[a]
C61	Sterile	High Melanin	9.71 (3.1)[ab]	29.3 (9.5)[a]
C61	Sterile	Low Melanin	10.9 (0.23)[ab]	26.9 (0.24)[a]

1292

1293 **Figure S1**. A. Soil microcosm design with modified vented caps and B. sterile soil microcosm receiving a

1294 sterile water addition.

1295

1296

1297

1298

1299 **Figure S2**. Morphology of goethite under Transmission Electron Microscopy.

1300

1301

1302

1303

1304

Figure S3. A. Diffuse Reflectance Infrared Fourier transform spectroscopy (DRIFTS) spectra with annotated peaks for MAOM samples grouped by necromass treatment. Principal Component Analysis (PCA) biplots of DRIFT spectra for mineral-associated organic matter (MAOM) samples from microcosms colored by B. necromass treatment and C. sterilization treatment.

1309

1310

1311 **Figure S4.** A. Diffuse Reflectance Infrared Fourier transform spectroscopy (DRIFTS) spectra and B. table

1312 of annotated peaks for initial *H. bicolor* necromass types (high and low melanin). Photos in top left corner

1313 are of initial necromass.

1314

1315

1316

1317

1318

Figure S5. A. Peak annotations for SIPT spectra B. Rate constants and C. Asymptotic peak heights

generated from asymptotic regression models. White rows represent Peak IDs for which we were unable

to fit an asymptotic regression model. *Peak 5 is mainly attributed to C=O stretching of Amide I, and

partially attributed to C-N stretching and N-H bending. While peak 6 is mainly attributed to N-H bending

and partially to C-N stretching. Peak 9 at 1248 cm^{-1} and the shoulder 1230 cm^{-1} might also attributed to

PO2- asymmetric stretching (Stuart, 2004).