1

Towards a More Objective and High-throughput Spheroid Invasion Assay Quantification Method

Rozanne W. Mungai[1], Roger J. Hartman II[2], Grace E. Jolin[1], Kevin W. Piskorowski[1] and Kristen L. Billiar[1*]

[1]Department of Biomedical Engineering, Worcester Polytechnic Institute, Worcester, MA, USA 01605

[2]No affiliations, independent researcher

11
12

13
14
*Correspondence to:
      Kristen L. Billiar
      Biomedical Engineering Department
      Worcester Polytechnic Institute
      100 Institute Road
      Worcester, MA 01609
      kbilliar@wpi.edu
      Phone: +1 (508) 8315000 x5384

ABSTRACT

Multicellular spheroids embedded in 3D hydrogels are prominent *in vitro* models for 3D cell

invasion. Yet, quantification methods for spheroid cell invasion that are high-throughput, objective and

accessible are still lacking. Variations in spheroid sizes and the shapes of the cells within render it

difficult to objectively assess invasion extent. The goal of this work is to develop a high-throughput

quantification method of cell invasion into 3D matrices that minimizes sensitivity to initial spheroid size

and cell spreading and provides precise integrative directionally-dependent metrics of invasion. By

analyzing images of fluorescent cell nuclei, invasion metrics are automatically calculated at the pixel

level. The initial spheroid boundary is segmented and automated calculations of the nuclear pixel

distances from the initial boundary are used to compute common invasion metrics (i.e., the change in

invasion area, mean distance) for the same spheroid at a later timepoint. We also introduce the area

moment of inertia as an integrative metric of cell invasion that considers the invasion area as well as the

pixel distances from the initial spheroid boundary. Further, we show that principal component analysis

can be used to quantify the directional influence of a stimuli to invasion (e.g., due to a chemotactic

gradient or contact guidance). To demonstrate the power of the analysis for cell types with different

invasive potentials and the utility of this method for a variety of biological applications, the method is

used to analyze the invasiveness of five different cell types. In all, implementation of this high-throughput

quantification method results in consistent and objective analysis of 3D multicellular spheroid invasion.

We provide the analysis code in both MATLAB and Python languages as well as a GUI for ease of use

for researchers with a range of computer programming skills and for applications in a variety of biological

research areas such as wound healing and cancer metastasis.

INTRODUCTION

Cell invasion and migration are driving factors for various biological events such as development [1,2], wound healing [3-6], tumor metastasis [7-9] and host cell infiltration of implanted scaffolds [10-13]. The effects of soluble and immobilized chemical gradients on cell migration on 2D surfaces and infiltration into 3D matrices and tissues is well established [9,14]; however, it is also becoming clear that mechanical cues play a large role in cell invasion and migration [7,15]. Cells move in response to physical cues such as stiffness gradients [9,15,16], contact guidance [9,15,17,18] and mechanical restraint [15,19-29].

In vitro models consisting of multicellular spheroids embedded in ECM are commonly used for investigating the mechanisms of cell invasion into extracellular matrices (ECM) in a variety of diseases [30-36]. Low contrast phase imaging is often used to capture images of cell invasion as it accommodates live-imaging. However, phase images of spheroid invasion are difficult to analyze with common automated software. Thus, this method often requires manual tracing of cell boundaries in programs such as ImageJ/FIJI to quantify the extent of invasion via measuring the distances and areas of migrating cells [8,30,37-43]. This manual tracing is time consuming and introduces subjectivity and possible technical error. Alternatively, high-contrast fluorescent images are amenable to automated quantification using common analysis software which allows for more objective high-throughput analysis. However, fluorescent images are less accommodating for live imaging as they require transfection of fluorescent proteins [44] or adding live cell stains which do not give a strong signal-to-noise ratio [45]. To overcome this limitation, clear high contrast fluorescence images can be captured by fixing and staining samples at the experiment endpoint using common accessible dyes. After imaging, the cell invasion can be quantified (such as with migration tracks [8,46]) using image analysis software or custom programs, and invasion metrics can be calculated.

Quantification metrics for cell invasion into 3D matrices commonly involve the overall invasion area

and/or the distance travelled by cells at a given timepoint [30,34,37,38,41,43,44,47-49]. The invasion area

is used to represent the number of invaded cells [30,34,37,38,41,43,44,49-51], and the distance travelled

is often reported as either the mean or maximum distance travelled by cells [8,30,41,49,52]. Yet, these

metrics are sensitive to changes in cell size and shape that occur as cells migrate through the matrix

[53,54]. In the case of multicellular spheroid *in vitro* models, when invasion area and distance travelled

are calculated without taking into account the initial spheroid boundary, these metrics are also sensitive to

differences in size and shape of the spheroids. A metric that takes into account both the invasion area and

distance travelled is especially useful, such as the migration index published by Liu *et al.* [30], as it

informs both how many and how far cells are invading. The types of quantification metrics that can be

calculated are limited by the methods used to capture and quantify images, especially if a common or

microscope-based software is used. Therefore, there is a need for an easy-to-implement method and

corresponding computational analysis tool that is precise and objective, offers flexibility in quantification,

and is simple to use for a researcher unfamiliar with programming.

Here we present a high-throughput automated quantification method of cell invasion into 3D

matrices using a complementary pixel-based method to segmentation methods. Multicellular spheroids

embedded in collagen hydrogels are live-stained for cell nuclei, which are easy to image and are more

resistant to shape changes during invasion than the cell body, and the distribution of nuclear pixels is

quantified. Live fluorescent imaging allows for precise segmentation of the initial spheroid boundary,

thus minimizing the effects of spheroid initial size and shape. Automated calculations of the invasion area

as well as distances and angles of nuclear pixels from the boundary are calculated—metrics that are

relatively insensitive to initial spheroid size. We introduce the area moment of inertia, which takes into

account both the area and distances of nuclear pixels of invading cells, as an integrative metric for the

overall extent of invasion. In combination with principal component analysis, the area moment of inertia

metric is also used to quantify invasion directionality in response to anisotropic mechanical constraint.

Five different cell types are included to demonstrate the applicability of the analysis for cell types with different invasive potentials, and we provide an open-source graphical interface and the detailed code in both MATLAB and Python languages for ease of adoption by researchers with a range of computer programming skills.

MATERIALS AND METHODS

Ethics

No experiments were performed on humans or animals for this study; therefore, ethics approval is not required. Freshly isolated cells used in this study were harvested from animal (pig) tissue while all other cells, including human cells, were obtained as a gift.

Cell types and culture conditions

Cell spheroids of various cell types were embedded in collagen hydrogels and allowed to invade the matrix for two days. The cell types were chosen to span mesenchymal and epithelial cell type classifications as they are known to be motile and immotile respectively. Valvular interstitial cells (VIC), dermal fibroblasts and smooth muscle cells (SMC) are the mesenchymal cell types studied to highlight differences in invasive potential within the group and were compared to epithelial RPE-1 cells. PC9 cells were also included as we are interested in how a cancerous epithelial cell type (having undergone the endothelial-to-mesenchymal transformation) compares to non-cancerous mesenchymal cell types.

Porcine aortic VICs were isolated from fresh pig hearts obtained from a local abattoir (Blood Farm, Groton, MA), and the VICs were isolated within three hours as per published protocols [20] and used in experiments between passages 2-8. Neonatal human dermal fibroblasts (NHF), originally harvested from de-identified donated male foreskin tissue, were obtained as a gift from Dr. George Pins (Worcester Polytechnic Institute) and used in experiments between passages 6-12. Immortalized WKY

3M-22 male rat aortic SMCs were obtained as a gift from Dr. Marsha Rolle (Worcester Polytechnic

Institute). The immortalized female human retinal pigment epithelial cell line (RPE-1) was obtained as a

gift from Dr. Amity Manning (Worcester Polytechnic Institute). The same culture medium base

formulation and conditions were used for all of the above cell types: DMEM (Gibco) supplemented with

10% v/v fetal bovine serum and 1% v/v antibiotic-antimycotic (Gibco). For the SMCs, 1% v/v MEM non-

essential amino acids (Gibco) were added to maintain a growth phenotype. The PC9 human lung

adenocarcinoma cell line was also obtained as a gift from Dr. Amity Manning and were cultured in

RPMI-1640 (GenClone, Genessee Scientific) supplemented with 10% v/v fetal bovine serum and 1% v/v

antibiotic-antimycotic (Gibco). The identity of the PC9 cancer cells have been previously confirmed by

STR analysis [55]. The cell cultures were maintained at 37°C in a humidified 5% CO_2-containing

incubator.

Generation of multicellular spheroids

3D spheroids were generated for each cell type as in our previous work [56]. Briefly, the cells were

trypsinized, resuspended, and stained with 5μg/mL Hoechst 33342 (Invitrogen) for 10 minutes at 37°C for

visualization of cell nuclei. The stain was then diluted, and the cells were resuspended at a cell

concentration of one million cells per 200 μL. Non-adherent agarose microwells were generated by

pipetting a pre-warmed 2% w/v agarose solution (MilliporeSigma), made in DMEM, into negative molds

with 500 μm diameter wells (24-96 or 12-256; Microtissues®), and the agarose was allowed to solidify.

The stained cell suspension was seeded into the prepared agarose microwells, and the cells were allowed

to settle into spheroids for one day (Supplementary Fig. 1) before embedding them into collagen

hydrogels.

Embedding of multicellular spheroids into collagen gels

The multicellular spheroids were harvested from the microwells and suspended in media. Cooled rat tail collagen type I (Advanced Biomatrix, RatCol®) was then mixed with neutralization solution following manufacturer's recommendations to make collagen hydrogels. The spheroid suspension was mixed into the collagen solution for ~ 20 spheroids/mL in a final collagen concentration of 2 mg/mL. Next, the solution was poured into well plates and allowed to gel for 30 minutes at 37°C. The well plates used were either a tissue culture plastic 6-well plate for maximizing gel volume, or a flexible silicone well plate with small side posts (CellScale MechanoCulture FX™) to restrict gel compaction (4x4 array of 8 mm x 8 mm square wells).

Fresh medium was added over the samples, and the samples were imaged (Day 0) as described below. After imaging, the samples were returned to 37°C in a humidified 10% CO_2-containing incubator and allowed to culture for two days (Supplementary Fig. 2). The timepoint of two days was chosen to allow sufficient time for cells to invade into the matrix while minimizing migration past the camera field of view.

Staining and Imaging

The extent of cell invasion and migration was assessed by capturing images of the cell nuclei as 10µm-spaced z-stack slices through the thickness of the spheroids using a Keyence BZ-X810 fluorescence microscope (6.1µm DOF, High Resolution setting: 6dB gain, Binning off). The z-slices demonstrated consistent staining intensity through the depth of the spheroids (Supplementary Fig. 3). On Day 0, the spheroids, pre-stained with Hoechst during spheroid formation, were imaged live immediately after gelation of the collagen, and the spheroid locations were saved for use again during Day 2 imaging. On Day 2, the samples were fixed with 4% paraformaldehyde, permeabilized using 0.25% Triton-X 100 and stained again with Hoechst, at a 1:1000 dilution to obtain a strong fluorescent signal in the nuclei. They were further stained for F-actin visualization using Alexa Fluor® 488 phalloidin (Life Technologies) at a 1:100 dilution. The images of each z-stack were processed using the Keyence BZ-X800 Analyzer to

create maximum z-projection images (Fig. 1A, Supplementary Fig. 4), increase the image contrast and

convert them to grayscale.

Because the spheroids are mixed into the collagen hydrogel solution, it was common to observe

spheroids that were close to each other or out of the vertical field-of-view. Spheroids that were out of the

field of view or too close together (less than 400 µm apart or with migration paths of different spheroids

that could cross) were not imaged (Supplementary Fig. 5).

Quantification of cell invasion and migration into matrix

Image quantification was carried out using a custom MATLAB program (MathWorks®) using the image

processing package. The grayscale images were contrast enhanced, background subtracted if needed, and

binarized with a global threshold strategy. A circular mask was then applied to the images to prevent

biases towards the long axis and corners of the frame by excluding those pixels. For each set of Day 0 and

Day 2 binarized images, the Day 0 spheroid edge was located and automatically segmented to determine

the boundary of cell invasion (Fig. 1B). The centroids of the Day 0 and Day 2 spheroids were identified

and used to overlap the Day 0 boundary and the Day 2 binarized image. From there, the locations of all

the pixels of the Day 2 image that lay outside of the boundary were found (i.e., "the outer pixels") (Fig.

1C), and the distances of each of these pixels from both the boundary and the spheroid centroid were

calculated as well as their angles from the centroid, measured clockwise from the x-axis (as in a polar

coordinate system) (Fig. 1D).

A.	B.	C.	D.
	Trace Day 0 boundary	Find Day 2 outer pixels	Calculate distances and angles

Figure 1: Z-stack image processing and quantification. Images were processed to create maximum z-

projection images from each z-stack as demonstrated by a representative Day 2 image (A). The Day 0

boundary was segmented (B) and the pixels of the Day 2 image located past the boundary were identified

(C). The distances and angles of pixels outside of the boundary were calculated with reference to the

spheroid boundary and center. For clarity, only a portion of the distance lines are shown (D). Scale bar:

200 µm. Panel A: Representative spheroid is imaged at 20X magnification and off-set to highlight

invading cells. Panels B-D: 10X magnification.

The change in spheroid area *(ΔA)* i.e., the stained pixel area outside of the initial spheroid boundary, is

calculated by summing the number of pixels in the image and subtracting the Day 2 area to the Day 0

area, equation (1).

12
$$\Delta A = Area_{Day\ 2} - Area_{Day\ 0}$$

13   *(1)*

15   By locating the individual pixels in the images, rather than cell or nuclei centroids, pixel-level distance

16   calculations can be performed which avoids the difficulty of distinguishing clustered cells. The distance

17   *(d)* is calculated from the differences of the x and y distances from the boundary (x_b, y_b) and the outer

18   pixels (x_p, y_p) along a radial line emanating from the centroid of the Day 0 image to the outer pixel,

19   equation (2) (Fig. 1D):

20
$$d = \sqrt{(x_p - x_b)^2 + (y_p - y_b)^2}$$

21   *(2)*

22   The angles of invasion are obtained by calculating the arctangent of the x and y distances to the spheroid

23   centroid, (x_0, y_0), equation (3).

24
$$\theta = \tan^{-1}\frac{y_p - y_0}{x_p - x_0}$$

25   *(3)*

1   The invasion area moment of inertia *(I)* was evaluated as an integrative measure of the area change and

2   the distance from the spheroid boundary. The radial moment, I_r, was calculated by summing over the

3   squared radial distances (*r*) of each outer pixel (*i*) to the boundary, where *dA* is the area of each pixel,

4   equation (5).

5
$$I_r = \sum_{i=0}^{n} r^2 \, dA$$

6   (5)

Principal component analysis

In addition to being able to quantify the directional migration along the camera x- and y-axes, the

invasion area moment of inertia can be calculated along any arbitrary axes (x', y') using coordinate

transformation. Most notably, the directions of maximum and minimum invasion and extent of invasion

in each direction can be expediently calculated using principal component analysis (PCA). Coordinate

transformations were performed in MATLAB to reorient the distance values along the principal

components (directions of maximum and minimum invasion). The extent of invasion is then determined

by calculating the directional moments of inertia (*I$_x$'* and *I$_y$'*) along the principal axes, by summing the

transformed directional distances, equation (6), and calculating the mean distance in the principal

directions.

17
$$I_x' = \sum_{i=0}^{n} y'^{\,2} \, dA \;\; ; \;\; I_y' = \sum_{i=0}^{n} x'^{\,2} \, dA$$

18   (6)
19
20

Code availability

Our custom MATLAB image analysis program is shared on GitHub to facilitate use by other researchers.

<https://github.com/rmungai/SpheroidInvasionAnalysis>

To expand user accessibility, we have published a Python version of the program as well as a

downloadable GUI for users without computer programming backgrounds.

<https://github.com/rogerh2/SpheroidInvasionAnalysis>

Detailed descriptions of the scripts are also included on the GitHub pages.

Statistical analysis

An outlier analysis was performed on the dataset using MATLAB by detecting and removing values more

than three scaled median absolute deviations (MAD) from the median. The normality of the data set was

assessed using MATLAB via the D'Agostino & Pearson omnibus normality test employing a significance

level $\alpha = 0.05$. After determination of normality, significant differences among groups were analyzed

using the sjstats library in R using one-way analysis of variance (ANOVA) followed by Tukey's Honest

Significant Difference (HSD) post hoc test. Violin plots of the area change, mean distance from the

boundary, and area moment of inertia per spheroid were generated using the ggplot library in R and were

utilized to show the entire data distribution of all spheroids. Numbers of spheroids and biological

replicates are provided in the figure captions.

RESULTS

Image quantification of spheroid invasion

Images of the embedded spheroids demonstrate that all cell types invaded into the collagen hydrogel with

the extent of invasion varying substantially among cell types (Fig. 2, Supplementary Fig. 4).

Qualitatively, the RPE-1 cells demonstrated the least invasion while the SMCs demonstrated the most

invasion. The PC9 cells demonstrated the second-most invasion with the NHFs and VICs following,

respectively.

4
5

Figure 2: Invasion of cells from spheroids into collagen hydrogels. Representative images of invasion on
Day 0 compared to Day 2, VIC (A), NHF (B), SMC (C), RPE-1 (D) and PC9 (E). Binarized maximum
projection images of cell nuclei stained with Hoechst fluorescent dye at 10X magnification. Scale bar:
200 μm.

The binarized images were analyzed using our custom MATLAB program to obtain polar plots depicting

spheroid invasion past the boundary and calculate quantitative metrics for each cell type to demonstrate

the nuances of each invasion metric (Fig. 3). The area change is not necessarily correlated with distance

as demonstrated by comparing representative NHF and SMC polar plots (Fig. 3A&B). The SMCs do not

invade as far as the NHFs, resulting in a lower distance, but they do have a higher area change which

results in an area moment of inertia with a nearly equal value (~0.001 mm^4) for both cell types since the

moment incorporates both the distance of invasion and area change.

Figure 3: Visualization of variation of invasion behavior across cell types. Polar plots demonstrate invasion behavior of individual spheroids at Day 2. The distance (µm) versus angle is plotted for each pixel past the Day 0 spheroid boundary for VIC (A), NHF (B), SMC (C), RPE-1 (D) and PC9 (E). The boundary is represented as the plot center (0,0). Invasion metrics for representative spheroids, area change, ΔA, mean distance, $\bar{D}$, radial moment, I_r, as defined in the text, are provided below each plot.

Directional invasion assessment

We aimed to demonstrate that the area moment of inertia can be used to assess invasion directionality

resulting from a guidance cue, particularly in cases where directional invasion does not exclusively occur

along the x- and y- axes. We performed principal coordinate analysis (PCA) to a representative spheroid

exhibiting directional invasion due to its proximity to a stiff silicone post within the gel (Fig. 4A, right).

The post introduces constraint in the hydrogel which causes directional invasion towards it via contact

guidance [46,57-59]. A binarized image of the spheroid demonstrating directional invasion was corrected

to mask the post from the image (Fig. 4C, right) and quantified as described above. PCA resulted in the

determination of the principal angles of maximum and minimum cell invasion (Fig. 4D, right). A

coordinate transformation was then performed to calculate the extent of invasion along the principal

angles via the mean distance (0.104 mm and 0.071 mm) and the directional area moments of inertia

(0.227e-3 mm^4 and 0.103e-3 mm^4). The fold change of mean distances and directional moments were

calculated (by taking the ratio of the maximum to minimum directions) and found to be 1.5X and 2.2X

greater in the direction of the post, respectively, thus indicating directional invasion. To validate the

directional analysis method, a control spheroid (which was not located next to a post) was also quantified

for comparison (Fig 4, left). The mean distances along the principal angles for the control spheroid were

found to be 0.0984 and 0.0942 mm, and the directional moments were found to be 0.369e-3 mm^4 and

0.329e-3 mm^4 (Fig 4D, left) resulting in fold change values of 1.1X for both metrics. In comparison, the

mean distances and directional moments fold change values are 1.4X and 2X greater for the spheroid with

the post (Fig 4, right). Taken together, these results demonstrate that this method effectively detects and

quantifies invasion directionality.

Figure 4: Quantification program also assesses invasion directionality. Directional invasion was
demonstrated by imaging a collagen gel-embedded VIC spheroid growing near a structural post (indicated
by arrow) of the culture well for four days. Phase, Hoechst-stained and binarized spheroid images on Day
2 (no post control) and Day 4 (spheroid with post). Scale bar: 200 µm (A-C). PCA of quantified data
plotted as a polar plot centered at the mean. Principal angles are indicated (blue line: maximum invasion
at 125° and -12°; black line: minimum invasion). Polar plot depicts the distance from the boundary (mm)
versus angle for each pixel. Mean distances and directional moments along the principal angles were
calculated to quantify invasion directionality and are provided below the polar plot (D).

Statistical comparison of invasion between groups

Comparing the five cell types together, the change in area from Day 2 compared to Day 0 was found to be

the highest for the SMCs and all other cell types had a significantly reduced area change in comparison

indicating SMCs as the most invasive (Fig. 5A). The PC9 cells had the second-highest area change

followed by the NHF, VIC, and RPE-1 cells respectively. The area change of the PC9 cells was

significantly higher than that of the RPE-1 demonstrating that, as expected, the cancerous epithelial cell

type is much more invasive than the wild-type epithelial cell type.

For the mean invasion distance, SMCs and NHFs were found to have the longest distance

followed by VIC, PC9 and RPE-1 cells (Fig. 5B). All cells had a significantly higher mean distance than

the RPE-1 cells, demonstrating that the epithelial cell type was, as expected, the least motile compared to

the more migratory mesenchymal-type and cancer cells. SMCs did not demonstrate a significantly longer

distance than other cells apart from RPE-1. This difference in the SMC results for area change and

distance demonstrates that a high invasion area change does not necessarily indicate a longer invasion

distance.

For the radial area moment of inertia of invasion (I_r), the SMCs were found to have the highest

value which was significantly higher than all the other cell types, as with area change (Fig. 5C). Since the

area moment of inertia metric incorporates both invasion area and distance into the calculation, the

moment calculation demonstrates that SMCs are the most invasive cell type overall, a finding that may

have been overlooked if invasion distance was used alone.

Figure 5: Quantitative metrics of invasion. Quantified area change (A), mean distance (B) and radial area
moment of inertia (I_r) (C) for VIC, NHF, SMC, PC9 and RPE-1 cells. Biological replicates are VIC=3,
NHF & SMC=2, and RPE-1 & PC9=1. Number of spheroids are indicated by dots on each plot and are:
VIC n=9, 10, 8 (A-C respectively), NHF n=12, 10, 13 (A-C respectively), SMC n=11, 9, 10 (A-C
respectively), RPE n=8 (all), PC9 n=5 (all). "x" indicates the mean value. Significance at $p<0.05$
determined by one-way ANOVA with Tukey's HSD post hoc test (exact p-values provided in
Supplementary Table 1). * and ** indicate significant difference to SMC (A) and (C) and RPE-1 (B)
respectively while __#__ indicates significant difference between the specified groups (A).

DISCUSSION

In this work, we develop an *in vitro* imaging and analysis method to objectively assess the extent of cell invasion from multicellular spheroids into a surrounding 3D hydrogel. Individual spheroids are imaged at initial and final time points allowing for accurate determination of cell invasion over the culture period and overcoming limitations of methods that only analyze the final state of invasion [34,38,41]. The projected distribution of nuclear pixels past the Day 0 boundary is automatically identified to calculate common invasion metrics, e.g., the increase in area and mean distance from the boundary, as well as the area moment of inertia, our new integrative metric which quantifies how the pixels of the nuclei of invading cells are distributed in the image. These metrics are calculated with reference to the initial boundary of the spheroid at Day 0 which is automatically segmented. We introduce coordinate transformation to identify the directions and extent of most and least invasion for cases where the invasion may not be radially symmetric due to directional mechanical or chemical signaling gradients. We demonstrate the utility of our method using five different cell types of varying invasive potentials and show that the metrics quantitatively capture the differences in invasiveness among the cell types.

By automatically segmenting the initial spheroid to generate the boundary of invasion, we lessen the effect that spheroid size differences can have on the invasion metrics. While this approach is not an all-encompassing solution to the challenge of generating consistently sized spheroids, as demonstrated by the variability in our invasion metric results (Figure 5), it aids in reducing the effect of spheroid size on invasion metrics as previously described [47]. This approach is especially useful for cell types that do not reliably form uniform spheroids (Supplementary Fig. 6) [30,37,60,61]. A limitation of this method is that it assumes that invasion occurs from cells located at the spheroid boundary, thus it does not account for the additional distance traveled by cells invading from deeper within the spheroid. However, this method provides a key advantage over the majority of invasion quantification methods that assume all spheroids are the same size [30,38,41,43,49]. Automated boundary segmentation makes spheroid invasion analysis

more flexible for non-uniform spheroid sizes and shapes and, by extension, makes it easier to quantify

spheroids of various cell types.

To our knowledge, we are the first to introduce a pixel-based method for invasion analysis. Using

pixels is less subjective than segmenting cells because it can be difficult to automatically distinguish

individual cells when they are clustered (Supplementary Fig. 7), and manual tracing of cell boundaries is

user-dependent and time consuming and cannot be completed if the cell boundaries are not evident

[8,30,37-43,62]. Automated spheroid invasion quantification methods such as TASI segment the spheroid

mass as one object and, as a result, are limited to assessing the distance of invasion from the spheroid

centroid or core boundary to the edge of a dense invasion front [47,48]. This limitation is also present in

machine learning approaches [63] as well as in methods that quantify morphological features for

spheroids not embedded in ECM, such as AnaSP and INSIDIA [62,64]. Our method has the key

advantage of accounting for all pixels past the boundary—a distinction especially important for cell types

(such as primary cells) which separate into individual clumps and do not form a dense collective invasion

front, as we have demonstrated in Figure 3 and 5.

To determine the sensitivity of our pixel-based approach to thresholding, we analyzed the dissociated

single nuclei invading from a representative spheroid at low, medium and high global threshold values

and compared the results to a standard object-based method (Supplementary Fig. 8). As the threshold is

decreased, an increasing number of pixels surpass the threshold value leading to larger area, distance and

moment values for both the pixel-based and object-based methods at comparable degrees. We expected

that the number of single nuclei identified in the object-based method would be less sensitive to threshold

than the pixel-based metrics because changing the threshold value would simply change the sizes of

binarized nuclei. However, we were surprised to find a non-monotonic relationship between the number

of single nuclei and the threshold. The number of nuclei increased when the threshold was decreased from

the high to medium values as expected, but the number of nuclei decreased from the medium and low

threshold values due to nuclei either coalescing into the spheroid bulk or merging together into a single

object too large to meet the size criteria for a single nucleus. Therefore, our pixel-based approach is not

only comparable to an object-based approach for calculating invasion metrics but also potentially

advantageous as it does not strictly report the "number" of cells or nuclei which, as we have

demonstrated, is highly sensitive to thresholding.

Maximum projection z-stack images of fluorescently stained nuclei were utilized since the nuclei are

high contrast and easy to binarize and can be easily stained for live imaging using inexpensive and readily

available stains, (e.g., Hoechst) and are common for 3D spheroid invasion analysis [30,41,44,48,65]. If

analysis of 3D invasion along the z-direction is needed, these methods can potentially be applied to

individual z-stack images [66]. Further, quantifying cell nuclei rather than the cell body results in more

accurate invasion area measurements because the cell nucleus, unlike the cell body, commonly resists

large changes in shape during cell migration and invasion [53] due to being 5-10 times stiffer than the

surrounding cytoskeleton [54]. This effect of cell shape on perceived cell size in 2D images is an issue

overlooked by studies that assume invading cells are all the same size [8,30,34,37,43,44,49]. Thus,

utilizing individual pixels of fluorescently stained nuclei is a powerful approach to quantifying spheroid

cell invasion.

We introduce the application of the classical area moment of inertia parameter, which reflects

how the points of an area are distributed with regard to an arbitrary axis, and apply it towards spheroid

invasion as an integrative "invasiveness" metric encompassing both the projected distance and area of

invasion. Thus, the area moment of inertia is a more informative metric than the area change or the mean

distance alone. While the area change metric is useful for representing the change in cell number over the

culture period [30,37,38,43,44], it may be skewed by variations in cell proliferation rates among cell

types. The mean distance does not have this limitation, but it does not provide insight into the cell number

[41]. Liu *et al.* use a similar metric combining cell area and distance of invasion (not squared as for the

area moment of inertia) [30]; however, their method requires manual image tracing of cells from a

uniform spherical bead whereas our metric is implemented more automatically and can be applied to

common, arbitrarily shaped spheroids. Using specific cases, we show that NHFs and SMCs from

individual spheroids can have nearly identical invasiveness as measured by radial moment (~1.3e-3 mm^4)

yet substantially different area change and distance values (c.f., panels A and B in Fig. 3). Further, the

invasion of VICs can look qualitatively similar to that of NHFs and have a similar area of invasion, yet

the radial moment is almost 30% smaller (c.f., panels A and 3 in Fig. 3). These results highlight the

limitations of the area change and distance metrics when used alone and demonstrate the importance of

the area moment of inertia metric which incorporates both area and distance to quantify cell invasion.

Quantifying invasion directionality for cell populations is important for determining the effects of

spatial gradients of chemical and mechanical factors. Most previous studies either track single cell

directional migration [8,38,39,46,67] or quantify invasion of a population of cells along one direction

[46,47]. We directly calculate the extent of invasion along x- and y-image axes using the area moment of

inertia (I_x and I_y) as well as estimate it using the mean distances in the x- and y-directions. For simpler

comparison, the extent of anisotropic invasion between cell types can be calculated by the fold change of

the directional moments and distances. Ibrahim *et al.* similarly calculate the mean distance in the

maximum invasion direction (by fitting it to an ellipse) [47]; however, their approach limits analysis to

only the contiguous portion of the spheroid and omits the isolated cells whereas our approach includes all

invaded cells. For greater generalizability, we also introduce the use of coordinate transformation to

account for cases when invasion does not occur strictly along the axes of the image. We showed that

calculating the mean distances and directional moments along the axes of maximum and minimum

invasion using PCA is a straight-forward way to characterize the extent of directionality of invasion. As a

demonstration, we applied the method to a spheroid cultured near a stiff silicone post which increases cell

invasion along its direction via contact guidance and found the invasion (as measured by area moment of

inertia) to be 2.2-fold higher in the direction of the post. This method can be used to assess the directional

invasion response to guidance cues applied in any direction (e.g., mechanical constraints and chemical

gradients); to our knowledge, we are the first to apply PCA to cell invasion analysis.

To expand user accessibility to our computer program, we have published an open-source graphical user interface for researchers without training in computer programming. In addition, the source code is also published in both Python (which is open-source) and MATLAB to allow for custom adaptation as desired and to cater to the researcher's programming language preference. Taken together, these options broaden the ease of use of this spheroid invasion quantification method to biomedical researchers with a broad array of backgrounds.

CONCLUSIONS

Here we develop a high-throughput objective quantification method of cell invasion from multicellular spheroids into a 3D extracellular matrix and demonstrate its utility with different metrics of invasion applied to cell types of various invasive potentials. This pixel-based method can be applied to a variety of spheroid invasion studies, such as exploring the roles of ECM components and guidance cues (e.g., chemokines and dynamic stretch) in cell invasion. The use of a nuclear stain minimizes the effect of cell size variations, while live-staining the Day 0 spheroid allows for objective spheroid segmentation. Calculating the distances of all pixels past the Day 0 boundary, which is unique to each spheroid, minimizes the effects of spheroid size and shape on the calculated metrics of invasion. Thus, this automated custom image analysis tool is complementary to traditional segmentation methods and is a considerable addition to the field. We also introduce the area moment of inertia as an integrative metric of cell invasion. The program also utilizes coordinate transformation to allow researchers to assess invasion directionality in response to a guidance cue. This innovative approach to 3D cell invasion analysis has the potential to advance research in fields including wound healing, cancer metastasis, and the repopulation of decellularized tissue engineered scaffolds.

1   **DATA AVAILABILITY**

2   The datasets used and/or analyzed during the current study are available from the corresponding author on

3   reasonable request and are also available at the Harvard Dataverse using the following link:

4   <https://doi.org/10.7910/DVN/VQH0BK >

REFERENCES

Butcher, J. T. & Markwald, R. R. Valvulogenesis: the moving target. *Philos Trans R Soc Lond B Biol Sci* **362**, 1489-1503, doi:10.1098/rstb.2007.2130 (2007).

Ahuja, N., Ostwald, P., Bark, D. & Garrity, D. Biomechanical Cues Direct Valvulogenesis. *J Cardiovasc Dev Dis* **7**, doi:10.3390/jcdd7020018 (2020).

Elsdale, T. & Bard, J. Collagen substrata for studies on cell behavior. *J Cell Biol* **54**, 626-637, doi:10.1083/jcb.54.3.626 (1972).

Singer, A. J. & Clark, R. A. Cutaneous wound healing. *N Engl J Med* **341**, 738-746, doi:10.1056/nejm199909023411006 (1999).

Grinnell, F., Rocha, L. B., Iucu, C., Rhee, S. & Jiang, H. Nested collagen matrices: a new model to study migration of human fibroblast populations in three dimensions. *Exp Cell Res* **312**, 86-94, doi:10.1016/j.yexcr.2005.10.001 (2006).

Guillamat-Prats, R. The Role of MSC in Wound Healing, Scarring and Regeneration. *Cells* **10**, doi:10.3390/cells10071729 (2021).

Hayward, M.-K., Muncie, J. M. & Weaver, V. M. Tissue mechanics in stem cell fate, development, and cancer. *Developmental cell* **56**, 1833-1847 (2021).

Heussner, R. K., Zhang, H., Qian, G., Baker, M. J. & Provenzano, P. P. Differential contractility regulates cancer stem cell migration. *Biophys J* **122**, 1198-1210, doi:10.1016/j.bpj.2023.02.008 (2023).

SenGupta, S., Parent, C. A. & Bear, J. E. The principles of directed cell migration. *Nat Rev Mol Cell Biol* **22**, 529-547, doi:10.1038/s41580-021-00366-6 (2021).

VeDepo, M. C., Detamore, M. S., Hopkins, R. A. & Converse, G. L. Recellularization of decellularized heart valves: Progress toward the tissue-engineered heart valve. *J Tissue Eng* **8**, 2041731417726327, doi:10.1177/2041731417726327 (2017).

Syedain, Z. H., Maciver, R. & Tranquillo, R. T. Vascular grafts and valves that animate, made from decellularized biologically-engineered tissue tubes. *J Cardiovasc Surg (Torino)* **61**, 577-585, doi:10.23736/s0021-9509.20.11615-x (2020).

Syedain, Z. H. *et al.* Pediatric tri-tube valved conduits made from fibroblast-produced extracellular matrix evaluated over 52 weeks in growing lambs. *Sci Transl Med* **13**, doi:10.1126/scitranslmed.abb7225 (2021).

Bouten, C. V. C., Smits, A. & Baaijens, F. P. T. Can We Grow Valves Inside the Heart? Perspective on Material-based In Situ Heart Valve Tissue Engineering. *Front Cardiovasc Med* **5**, 54, doi:10.3389/fcvm.2018.00054 (2018).

Shellard, A. & Mayor, R. All Roads Lead to Directional Cell Migration. *Trends Cell Biol* **30**, 852-868, doi:10.1016/j.tcb.2020.08.002 (2020).

Zonderland, J. & Moroni, L. Steering cell behavior through mechanobiology in 3D: A regenerative medicine perspective. *Biomaterials* **268**, 120572 (2021).

Deng, B. *et al.* Biological role of matrix stiffness in tumor growth and treatment. *J Transl Med* **20**, 540, doi:10.1186/s12967-022-03768-y (2022).

Roby, T., Olsen, S. & Nagatomi, J. Effect of sustained tension on bladder smooth muscle cells in three-dimensional culture. *Ann Biomed Eng* **36**, 1744-1751, doi:10.1007/s10439-008-9545-5 (2008).

Tondon, A. & Kaunas, R. The direction of stretch-induced cell and stress fiber orientation depends on collagen matrix stress. *PLoS One* **9**, e89592, doi:10.1371/journal.pone.0089592 (2014).

Throm Quinlan, A. M., Sierad, L. N., Capulli, A. K., Firstenberg, L. E. & Billiar, K. L. Combining dynamic stretch and tunable stiffness to probe cell mechanobiology in vitro. *PLoS One* **6**, e23272, doi:10.1371/journal.pone.0023272 (2011).

Quinlan, A. M. & Billiar, K. L. Investigating the role of substrate stiffness in the persistence of valvular interstitial cell activation. *J Biomed Mater Res A* **100**, 2474-2482, doi:10.1002/jbm.a.34162 (2012).

Cirka, H. *et al.* Active Traction Force Response to Long-Term Cyclic Stretch Is Dependent on Cell Pre-stress. *Biophys J* **110**, 1845-1857, doi:10.1016/j.bpj.2016.02.036 (2016).

Jebeli, M. *et al.* Multicellular aligned bands disrupt global collective cell behavior. *Acta Biomater* **163**, 117-130, doi:10.1016/j.actbio.2022.10.041 (2023).

Desai, L. P., Chapman, K. E. & Waters, C. M. Mechanical stretch decreases migration of alveolar epithelial cells through mechanisms involving Rac1 and Tiam1. *Am J Physiol Lung Cell Mol Physiol* **295**, L958-965, doi:10.1152/ajplung.90218.2008 (2008).

Boulter, E., Tissot, F. S., Dilly, J., Pisano, S. & Féral, C. C. Cyclic uniaxial mechanical stretching of cells using a LEGO(®) parts-based mechanical stretcher system. *J Cell Sci* **133**, doi:10.1242/jcs.234666 (2020).

Klingberg, F. *et al.* Prestress in the extracellular matrix sensitizes latent TGF-β1 for activation. *The Journal of cell biology* **207**, 283-297, doi:10.1083/jcb.201402006 (2014).

Zhang, B. *et al.* Cyclic mechanical stretching promotes migration but inhibits invasion of rat bone marrow stromal cells. *Stem Cell Research* **14**, 155-164, doi:https://doi.org/10.1016/j.scr.2015.01.001 (2015).

Song, J. *et al.* Physiological cyclic stretch up-regulates angiotensin-converting enzyme 2 expression to reduce proliferation and migration of vascular smooth muscle cells. *Biosci Rep* **40**, doi:10.1042/bsr20192012 (2020).

Balachandran, K., Sucosky, P., Jo, H. & Yoganathan, A. P. Elevated cyclic stretch alters matrix remodeling in aortic valve cusps: implications for degenerative aortic valve disease. *American Journal of Physiology-Heart and Circulatory Physiology* **296**, H756-H764 (2009).

Balachandran, K. *et al.* Cyclic strain induces dual-mode endothelial-mesenchymal transformation of the cardiac valve. *Proceedings of the National Academy of Sciences* **108**, 19943-19948 (2011).

Liu, H., Lu, T., Kremers, G.-J., Seynhaeve, A. L. B. & ten Hagen, T. L. M. A microcarrier-based spheroid 3D invasion assay to monitor dynamic cell movement in extracellular matrix. *Biological Procedures Online* **22**, 3, doi:10.1186/s12575-019-0114-0 (2020).

Bakker, E. R. M. *et al.* Wnt5a promotes human colon cancer cell migration and invasion but does not augment intestinal tumorigenesis in Apc 1638N mice. *Carcinogenesis* **34**, 2629-2638, doi:10.1093/carcin/bgt215 (2013).

Janmaat, V. T. *et al.* HOXA9 mediates and marks premalignant compartment size expansion in colonic adenomas. *Carcinogenesis* **40**, 1514-1524, doi:10.1093/carcin/bgz038 (2019).

Dietrich, F. & Lelkes, P. I. Fine-tuning of a three-dimensional microcarrier-based angiogenesis assay for the analysis of endothelial-mesenchymal cell co-cultures in fibrin and collagen gels. *Angiogenesis* **9**, 111-125 (2006).

Arevalos, C. A. *et al.* Valve Interstitial Cells Act in a Pericyte Manner Promoting Angiogensis and Invasion by Valve Endothelial Cells. *Ann Biomed Eng* **44**, 2707-2723, doi:10.1007/s10439-016-1567-9 (2016).

Kniazeva, E. & Putnam, A. J. Endothelial cell traction and ECM density influence both capillary morphogenesis and maintenance in 3-D. *American Journal of Physiology-Cell Physiology* **297**, C179-C187 (2009).

Juliar, B. A., Keating, M. T., Kong, Y. P., Botvinick, E. L. & Putnam, A. J. Sprouting angiogenesis induces significant mechanical heterogeneities and ECM stiffening across length scales in fibrin hydrogels. *Biomaterials* **162**, 99-108 (2018).

Cisneros Castillo, L. R., Oancea, A. D., Stüllein, C. & Régnier-Vigouroux, A. Evaluation of Consistency in Spheroid Invasion Assays. *Sci Rep* **6**, 28375, doi:10.1038/srep28375 (2016).

Elosegui-Artola, A. *et al.* Matrix viscoelasticity controls spatiotemporal tissue organization. *Nature Materials*, doi:10.1038/s41563-022-01400-4 (2022).

Ray, A., Slama, Z. M., Morford, R. K., Madden, S. A. & Provenzano, P. P. Enhanced Directional Migration of Cancer Stem Cells in 3D Aligned Collagen Matrices. *Biophys J* **112**, 1023-1036, doi:10.1016/j.bpj.2017.01.007 (2017).

Solbu, A. A. *et al.* Assessing cell migration in hydrogels: An overview of relevant materials and methods. *Materials Today Bio* **18**, 100537, doi:https://doi.org/10.1016/j.mtbio.2022.100537 (2023).

Ducker, M., Millar, V., Ebner, D. & Szele, F. G. A Semi-automated and Scalable 3D Spheroid Assay to Study Neuroblast Migration. *Stem Cell Reports* **15**, 789-802, doi:https://doi.org/10.1016/j.stemcr.2020.07.012 (2020).

Dolega, M. *et al.* Mechanical behavior of multi-cellular spheroids under osmotic compression. *Journal of the Mechanics and Physics of Solids* **147**, 104205 (2021).

Nandi, S. & Brown, A. C. Characterizing Cell Migration Within Three-dimensional In Vitro Wound Environments. *J Vis Exp*, doi:10.3791/56099 (2017).

Baskaran, J. P. *et al.* Cell shape, and not 2D migration, predicts extracellular matrix-driven 3D cell invasion in breast cancer. *APL Bioeng* **4**, 026105, doi:10.1063/1.5143779 (2020).

Schneckenburger, H. & Richter, V. in *Photonics.* 275 (MDPI).

Rouillard, A. D. & Holmes, J. W. Mechanical boundary conditions bias fibroblast invasion in a collagen-fibrin wound model. *Biophys J* **106**, 932-943, doi:10.1016/j.bpj.2013.12.002 (2014).

Ibrahim, H. *et al.* A Biomimetic high throughput model of cancer cell spheroid dissemination onto aligned fibrillar collagen. *SLAS technology* **27**, 267-275 (2022).

Hou, Y., Konen, J., Brat, D. J., Marcus, A. I. & Cooper, L. A. TASI: A software tool for spatial-temporal quantification of tumor spheroid dynamics. *Scientific reports* **8**, 7248 (2018).

Mina, S. G., Huang, P., Murray, B. T. & Mahler, G. J. The role of shear stress and altered tissue properties on endothelial to mesenchymal transformation and tumor-endothelial cell interaction. *Biomicrofluidics* **11**, 044104, doi:10.1063/1.4991738 (2017).

Menon, S. & Beningo, K. A. Cancer cell invasion is enhanced by applied mechanical stimulation. *PLoS One* **6**, e17277, doi:10.1371/journal.pone.0017277 (2011).

O'Leary, R. & Wood, E. J. A novel in vitro dermal wound-healing model incorporating a response to mechanical wounding and repopulation of a fibrin provisional matrix. *In Vitro Cell Dev Biol Anim* **39**, 204-207, doi:10.1290/1543-706x(2003)039<0204:Anivdw>2.0.Co;2 (2003).

Petroll, W. M., Lakshman, N. & Ma, L. Experimental models for investigating intra-stromal migration of corneal keratocytes, fibroblasts and myofibroblasts. *J Funct Biomater* **3**, 183-198, doi:10.3390/jfb3010183 (2012).

McGregor, A. L., Hsia, C.-R. & Lammerding, J. Squish and squeeze—the nucleus as a physical barrier during migration in confined environments. *Current Opinion in Cell Biology* **40**, 32-40, doi:https://doi.org/10.1016/j.ceb.2016.01.011 (2016).

Friedl, P., Wolf, K. & Lammerding, J. Nuclear mechanics during cell migration. *Curr Opin Cell Biol* **23**, 55-64, doi:10.1016/j.ceb.2010.10.015 (2011).

Rassias, D. J. & Weathers, P. J. Dried leaf Artemisia annua efficacy against non-small cell lung cancer. *Phytomedicine* **52**, 247-253, doi:10.1016/j.phymed.2018.09.167 (2019).

Coutts, C. W. *et al.* The Role of Apoptosis and Oxidative Stress in a Cell Spheroid Model of Calcific Aortic Valve Disease. *Cells* **13**, 45 (2024).

Miron-Mendoza, M., Seemann, J. & Grinnell, F. The differential regulation of cell motile activity through matrix stiffness and porosity in three dimensional collagen matrices. *Biomaterials* **31**, 6425-6435, doi:10.1016/j.biomaterials.2010.04.064 (2010).

Matsumoto, T. *et al.* Three-dimensional cell and tissue patterning in a strained fibrin gel system. *PloS one* **2**, e1211-e1211, doi:10.1371/journal.pone.0001211 (2007).

Pelham, R. J., Jr. & Wang, Y. l. Cell locomotion and focal adhesions are regulated by substrate flexibility. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 13661-13665, doi:10.1073/pnas.94.25.13661 (1997).

Vinci, M. *et al.* Advances in establishment and analysis of three-dimensional tumor spheroid-based functional assays for target validation and drug evaluation. *BMC biology* **10**, 1-21 (2012).

Dolznig, H. *et al.* Modeling colon adenocarcinomas in vitro: A 3D co-culture system induces cancer-relevant pathways upon tumor cell and stromal fibroblast interaction. *The American journal of pathology* **179**, 487-501 (2011).

Piccinini, F. AnaSP: A software suite for automatic image analysis of multicellular spheroids. *Computer Methods and Programs in Biomedicine* **119**, 43-52, doi:https://doi.org/10.1016/j.cmpb.2015.02.006 (2015).

Ngo, T. K. N. *et al.* A deep learning-based pipeline for analyzing the influences of interfacial mechanochemical microenvironments on spheroid invasion using differential interference contrast microscopic images. *Materials Today Bio* **23**, 100820 (2023).

Perini, G. *et al.* INSIDIA 2.0 high-throughput analysis of 3D cancer models: multiparametric quantification of graphene quantum dots photothermal therapy for glioblastoma and pancreatic cancer. *International Journal of Molecular Sciences* **23**, 3217 (2022).

Huang, Y. L. *et al.* Tumor spheroids under perfusion within a 3D microfluidic platform reveal critical roles of cell-cell adhesion in tumor invasion. *Scientific Reports* **10**, 9648, doi:10.1038/s41598-020-66528-2 (2020).

Weiss, F. *et al.* 3D spheroid culture to examine adaptive therapy response in invading tumor cells. *In Vitro Model* **1**, 463-471, doi:10.1007/s44164-022-00040-x (2022).

Pang, Y., Wang, X., Lee, D. & Greisler, H. P. Dynamic quantitative visualization of single cell alignment and migration and matrix remodeling in 3-D collagen hydrogels under mechanical force. *Biomaterials* **32**, 3776-3783, doi:10.1016/j.biomaterials.2011.02.003 (2011).

ACKNOWLEDGEMENTS

The assistance provided by Jamie Baines in conducting experiments is gratefully acknowledged.

This work was supported by the National Science Foundation (NSF) [CMMI 1761432], the National Institutes of Health (NIH) [1R15HL167235-01] and the American Heart Association (AHA) [20AIREA35120448]. The funders had no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

AUTHOR INFORMATION

Affiliations

Department of Biomedical Engineering, Worcester Polytechnic Institute, Worcester, MA, USA 01605.

Rozanne W. Mungai, Grace E. Jolin, Kevin W. Piskorowski and Kristen L. Billiar

No affiliations. Independent researcher, Worcester, MA, USA 01605.

Roger J. Hartman II

Contributions

K.L.B. and R.W.M. conceived the experiment which was performed by R.W.M., G.E.J., and K.W.P.. Results analysis was performed by R.W.M. who also quantified the data using MATLAB. R.J.H.II translated the quantification methods to Python and designed the GUI. R.W.M. wrote the manuscript draft which was edited by K.L.B. All authors approved the final version of the manuscript.

ETHICS DECLARATIONS

Competing interests

1   The authors declare no competing financial interests.

 FIGURE LEGENDS

Figure 1: Z-stack image processing and quantification. Images were processed to create maximum z-

projection images from each z-stack as demonstrated by a representative Day 2 image (A). The Day 0

boundary was segmented (B) and the pixels of the Day 2 image located past the boundary were identified

(C). The distances and angles of the pixels outside of the boundary were calculated with reference to the

spheroid boundary and center. For clarity, only a portion of the distance lines are shown (D). Scale bar:

200µm. Panel A: Representative spheroid is imaged at 20X magnification and off-set to highlight

invading cells. Panels B-D: 10X magnification.

Figure 2: Invasion of cells from spheroids into collagen hydrogels. Representative images of invasion

on Day 0 compared to Day 2, VIC (A), NHF (B), SMC (C), RPE-1 (D) and PC9 (E). Binarized maximum

projection images of cell nuclei stained with Hoechst fluorescent dye at 10X magnification. Scale bar:

200 µm

Figure 3: Visualization of variation of invasion behavior across cell types. Polar plots demonstrate

invasion behavior of individual spheroids at Day 2. The distance (µm) versus angle is plotted for each

pixel past the Day 0 spheroid boundary for VIC (A), NHF (B), SMC (C), RPE-1 (D) and PC9 (E). The

boundary is represented as the plot center (0,0). Invasion metrics for representative spheroids, area

change, ΔA, mean distance, $\overline{D}$, radial moment, I_r, as defined in the text, are provided below each plot.

Figure 4: Quantification program also assesses invasion directionality. Directional invasion was

demonstrated by imaging a collagen gel-embedded VIC spheroid growing near a structural post (indicated

by arrow) of the culture well for four days. Phase, Hoechst-stained and binarized spheroid images on Day

2 (no post control) and Day 4 (spheroid with post). Scale bar: 200 µm (A-C). PCA of quantified data

plotted as a polar plot centered at the mean. Principal angles are indicated (blue line: maximum invasion

at 125° and -12°; black line: minimum invasion). Polar plot depicts the distance from the boundary (mm)

versus angle for each pixel. Mean distances and directional moments along the principal angles were

calculated to quantify invasion directionality and are provided below the polar plot (D).

Figure 5: Quantitative metrics of invasion. Quantified area change (A), mean distance (B) and radial

area moment of inertia (I_r) (C) for VIC, NHF, SMC, PC9 and RPE-1 cells. Biological replicates are

VIC=3, NHF & SMC=2, and RPE-1 & PC9=1. Number of spheroids are indicated by dots on each plot

and are: VIC n=9, 10, 8 (A-C respectively), NHF n=12, 10, 13 (A-C respectively), SMC n=11, 9, 10 (A-C

respectively), RPE n=8 (all), PC9 n=5 (all). "x" indicates the mean value. Significance at p<0.05

determined by one-way ANOVA with Tukey's HSD post hoc test (exact p-values provided in

Supplementary Table 1). * and ** indicate significant difference to SMC (A) and (C) and RPE-1 (B)

respectively while __#__ indicates significant difference between the specified groups (A).