Metabolic mode estimated by breathing reflects long-term motor memory

Takuji Hayashi[1*], Nobuyasu Nakano[2,3,4], Sohei Washino[3], Akihiko Murai[3].

1. Graduate School of Education, The University of Tokyo, Tokyo 113-0033,, Japan

2. Center for Information and Neural Networks, National Institute of Information and Communications

Technology, Osaka 565-0871, Japan

3. Human Augmentation Research Center, National Institute of Advanced Industrial Science and Technology,

Chiba 277-0882, Japan

4. Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

*Corresponding author.

Keywords

11 Respiratory exchange ratio, Sensorimotor learning, Reaching movement, Long-term memory, Glucose

12 intervention

Correspondence

14 Takuji Hayashi

15 The University of Tokyo

16 Graduate School of Education

17 Division of Physical and Health Education

18 Email: thayashi@p.u-tokyo.ac.jp

Acknowledgements

20 We thank Ryota Furuno, On Kobayashi, Yuta Nozawa, Kotaro Beppu, and Asako Munakata for their help in

21 organizing and conducting the experiments. We thank Daichi Nozaki and Maurice Smith for helpful

22 discussions. This work was supported by KAKENHI grants from JSPS (JP23H03296) and JST/PRESTO

23 (JPMJPR23S8) to TH. This work was also partly supported by JST Moonshot R&D Grant Number

24 JPMJMS2239.

Author contributions

26 T.H., N.N., S.W., and A.M. designed the experiments. T.H. and N.N. performed the experiments. T.H.

27 analyzed the data. T.H., N.N., S.W., and A.M. wrote the paper.

Competing interests

29 The authors declare no competing interests.

Abstract

Respiration is a crucial metabolic process that converts macronutrients, carbohydrates and fats, and oxygen (O_2) into energy and carbon dioxide (CO_2) to support motor actions. In addition to the energy demands of movements, the brain is a significant energy consumer, accounting for approximately 20% of the body's total energy expenditure and relying primarily on carbohydrates for neural activity and plasticity. However, it is not known whether O_2-CO_2 gas composition in breathing can serve as an indicator of neural activity and plasticity as they can for movement intensity. In the human reaching movement tasks, we evaluated time-constants of sensorimotor learning while recording O_2-CO_2 gas exchange. We computed the respiratory exchange ratio (RER), indicating which carbohydrate or fat is used preferentially, and found that the RER was unaffected by the execution and learning of reaching movements and that it was stable within individuals but varied across individuals. Interestingly, using computational modeling to identify short and long-time constants of sensorimotor learning, individual RER levels correlated with the magnitude of long-term, but not short-term memory. Furthermore, to experimentally manipulate the individual RER, we provided 200 kcal of glucose immediately before the task. Surprisingly, this simple intervention dramatically increased 24-hour retention by 21%. Together, the RER served as a remarkable proxy for long-term motor memory, and glucose intake shifted the physiological idling state for sensorimotor learning.

INTRODUCTION

Respiration is a crucial process in metabolism responsible for the generation of adenosine triphosphate (ATP), an energy currency that supports all motor actions (Fig. 1a). Metabolism converts macronutrients, primarily carbohydrates and fats, along with oxygen (O_2), into ATP and carbon dioxide (CO_2). These metabolic processes exhibit varying rates and efficiencies. Quick, high-intensity motor actions require rapid energy production from carbohydrates, whereas prolonged, low-intensity motor actions require significant energy production from fat. Several studies have shown that the intensity of movement, measured by force or muscle activity, can be approximated with gas composition in breathing and is energetically optimized by motor experience (Abram et al., 2022; Huang et al., 2012; Selinger et al., 2015; Shadmehr et al., 2016). Therefore, breathing is recognized as a reliable measure for estimating the energy expenditure associated with movement execution.

In addition to the energy demands for muscle contraction, the brain is a significant energy consumer, utilizing 20% of the total energy expenditure despite comprising only 2% of body weight (Raichle and Gusnard, 2002). In particular, cortical energy is crucial for both neural activity (Oakes et al., 2004; Sokoloff, 1977) and plasticity (Lee et al., 2013; zur Nedden et al., 2011), which must play pivotal roles for sensorimotor learning. Sensorimotor learning not only establishes new patterns of neural activity (Li et al., 2001; Paz et al., 2003; Sun et al., 2022) but also induces novel cortical connections through long-term potentiation-like mechanisms (Kida et al., 2016; Landi et al., 2011; Rioult-Pedotti et al., 2000, 1998). Notably, these processes operate on different time scales: trial-by-trial emergence of new neural activity (Perich et al., 2018; Veuthey et al., 2020) contrasts with the slower, structural consolidation of connections, such as dendritic spine growth within an hour at the earliest (Harms et al., 2008; Xu et al., 2009). This temporal dichotomy may mirror a prevalent computational framework in sensorimotor learning, in which fast and slow processes interact to update motor memory (Smith et al., 2006). The fast process adapts quickly but decays rapidly, whereas the slow process learns gradually and supports longer-term retention. Computational models incorporating these dynamics successfully capture behavioral features of both short- and long-term memory formation (Joiner and Smith, 2008). Given the energetic demands of both fast and slow learning processes, we hypothesized that breathing metrics—particularly those reflecting cerebral metabolism—might provide a window into the temporal characteristics of sensorimotor learning.

To test this hypothesis, we conducted human behavioral experiments involving reaching movement tasks while recording O_2-CO_2 gas exchange. We focused on the respiratory exchange ratio (RER, Fig. 1b), which is the volume of CO_2 produced divided by the volume of O_2 consumed (McArdle et al., 2010), as it indicates the metabolic mode. Carbohydrate metabolism converts O_2 to CO_2 in a 1:1 ratio (RER = 1.0), while fat metabolism converts O_2 to CO_2 in a 1:0.7 ratio (RER = 0.7). Thus, the RER theoretically varies between 0.7-1.0, depending on the macronutrient usage. Since the brain utilizes only carbohydrates for the energy resource, its RER in the cortex is 0.97-0.99 (Himwich and Nahum, 1932), implying that a higher RER could signal increased cerebral energy expenditure. We observed that the RER was not influenced by the execution and learning of reaching movements; rather, it was stable within individuals but varied across individuals. Interestingly, using a computational decomposition of motor memory into fast and slow components, we

discovered that individual RER levels correlated with the magnitude of slow motor memory, but not fast

memory. Furthermore, we provided 200 kcal of glucose immediately before the experiment to test the

hypothesis that manipulation of the RER can modulate long-term motor memory. This simple intervention

increased the RER on day 1, and surprisingly, enhanced motor memory retention by 21% on day 2. Together,

these findings suggest that the RER can serve as a physiological proxy for long-term motor memory, and that

modulating metabolic state with glucose intake before learning can shift the energetic baseline, thereby

enhancing memory consolidation.

RESULTS

Sensorimotor learning of dynamic perturbation in Experiment 1

Experiment 1 aimed to distinguish the short- and long-time constants of sensorimotor learning while

monitoring O2-CO2 gas exchange and muscle activity in 18 human participants (Fig. 1c, 18 males aged 21-52

years). Participants engaged in planar reaching movements controlling a visual cursor on a display, where a

visuomotor rotation (VMR) shifted the visual cursor away from hand movements angularly (Fig. 1d). To

dissect the time constant of sensorimotor learning (Fig. 1e), we introduced a dynamic VMR perturbation (Fig.

1f and g), comprising 10°-amplitude sinusoidal sequences of VMR with different time constants ([2, 4, 8, 16,

and 32] cycles over 900 trials) (Miyamoto et al., 2020). The two-state model posits that, in sensorimotor

learning, two motor memories are updated concurrently with different time constants (Smith et al., 2006). This

unique idea, other from more recent frameworks (Heald et al., 2021), is leveraged to reliably estimate time

features of motor memory, where slow motor memory predicts next-day retention at 99% of variances in the

group-level analysis and at 50% of variances in the individual level (Joiner and Smith, 2008). Accordingly, we

predicted that fast motor memory would adapt to VMR changes more quickly (16 and 32 cycles), while the

slow motor memory would adapt to VMR changes more gradually (2 and 4 cycles). The dynamic VMR

allowed for an adequate assessment of learning components with different time constants.

Overall movement directions largely followed changes in the dynamic VMR (Fig. 2a). Fourier

transform analysis revealed large amplitudes only at trained frequencies ([2, 4, 8, 16, and 32] Hz), indicating

appropriate adaption to the dynamic VMR (Fig. 2b). Despite not reaching the perturbation levels of 10°, the

mean amplitudes at the trained frequencies (6.97° ± 0.29°, mean ± SEM) significantly surpassed those at the

untrained frequencies between 2 and 32 Hz (0.56° ± 0.10°) (Fig. 2c, $t(17) = 19.99$ and $p = 3.0 \times 10^{-13}$). These

results indicate appropriate participant adjustment to the dynamic VMR.

Effects of reaching movements on RER

The RER is known to covary with the intensity of movement execution (Goedecke et al., 2000); therefore,

changes in muscle activity could potentially confound the relationship between the RER and sensorimotor

learning. Thus, we investigated how sensorimotor learning altered muscle activity and whether these changes

affected the RER. Electromyography (EMG) data from representative muscles including the pectoralis major,

posterior deltoid, biceps brachii, and triceps brachii were recorded to observe movement execution intensity

effects. Muscle activity normalized by the trial average was high at the experiment's outset but quickly

decreased, stabilizing at approximately one-third of the experiment's duration (Fig. 2d). This pattern,

recognized as the relaxation of co-contraction into a novel environment (Franklin *et al.*, 2003), contrasted with

the stability observed in the mean RER across participants, which did not fluctuate across trials (Fig. 2e),

suggesting a deviation from typical sensorimotor learning patterns (Fig. 2a).

We examined whether muscle activity could account for RER variance across trials. Performing a

linear regression with trial lags, we found that only 3.6% of the variance could be explained without lags, with

minimal improvement observed across trials, even when accounting for trial lags (Fig. 2f). This was in

comparison with the permuted data from randomized trial blocks without significance noted for all trial lags

after false discovery rate (FDR) correction ($p > 0.05$). A slight enhancement was detected with 36-46 trial lags

($p < 0.05$, without FDR correction). However, these trials elapsed approximately 190-243 seconds (trial

duration was 5.3 ± 0.1 seconds), which is 1.5-2.0 times longer considering the time constant of the metabolic

loop compared with the previous report showing a 120-second lag (Short and Sedlock, 1997).

To further explore the impact of movement execution, we compared the average RER during the

reaching movement tasks and breaks (Fig. 2g). Despite high consistency within participants ($r = +0.91$ and p

$= 2.0 \times 10^{-7}$), the absence of a statistically significant difference between task and break RERs ($t(17) = 0.29$

and $p = 0.78$) indicated minimal effect of movement execution. Additionally, RERs during tasks exhibited no

correlation with mean peak velocity (Fig. 2h, $r = -0.10$ and $p = 0.69$) or reaction time (Fig. 2i, $r = -0.26$ and $p =$

0.30). These results suggest minimal influence of movement execution on the RER owing to the insufficient

intensity of reaching movements.

Effects of sensorimotor learning on RER

Next, we investigated whether there was any indication of sensorimotor learning reflected in the RER (Fig.

2e). However, we found that the RER remained constant during sensorimotor learning when comparing the

RER between the early and late training blocks (Fig. 2j, $t(17) = 1.91$ and $p = 0.07$). Nonetheless, these values

were highly consistent within participants ($r = +0.65$ and $p = 0.0033$).

Given that we used the dynamic VMR as the sum of the sinusoidal function, the amount of VMR to be

learned in the early training block was equal to that in the late training block. Consequently, if the RER were to

fluctuate with the amount of motor memory, it should remain stable during sensorimotor learning. In such a

scenario, the peaks in the frequency domain of the RER are the same as those in the dynamic VMR.

Alternatively, the RER might fluctuate with changes in motor memory, particularly following the derivatives of

the sum of the sinusoidal functions, which would manifest as cosinusoidal function at the same frequencies.

Thus, we would expect to observe peaks at [2, 4, 8, 16, and 32] Hz if sensorimotor learning affected the RER.

Figure 2k illustrates the amplitudes of the frequency analysis, revealing no discernible peaks. However, we

found a slight increase in the mean amplitudes at the trained frequencies compared to those at the untrained

frequencies (Fig. 2l, $t(17) = 4.01$ and $p = 9.1 \times 10^{-4}$).

The significant difference in the RER at the trained and untrained frequencies may not necessarily

signal sensorimotor learning for the following three reasons. First, the absolute difference (0.0017) was

smaller than the standard deviation of the within-participant RER fluctuation (SD: 0.044 ± 0.003). Second, the

RER may not behave as a random process as white noise, it may exhibit characteristics of pink noise (or 1/f

noise), where the power is inversely proportional to the frequency. Such patterns of pink noise are widely

found in human physiology (Pritchard, 1992; Zarahn et al., 1997) and cognition (Gilden et al., 1995), and the

RER's tendency to display larger amplitudes at low frequencies (Fig. 2k) could imply a pattern of pink noise. If

the RER is 1/f scaled, the mean amplitudes at the trained frequencies will be 2.41 times larger than those at

the untrained frequencies. Third, the Fourier analysis could not identify polarity, meaning that we cannot

ascertain whether the RER increased or decreased with sensorimotor learning. To further investigate whether

the RER changed in tandem with sensorimotor learning, we analyzed RER in Experiment 2 (see below).

165 Individual variability in RER reflects slow motor memory

We designed the dynamic VMR to rigorously estimate both fast and slow components of sensorimotor

learning. Individual data were fit into a two-state model. The overall motor output followed this behavior (Fig.

3a, $R^2 = 0.82 \pm 0.04$). As predicted, we observed peaks at the trained frequencies for both fast and slow

components, as well as overall (Fig. 3b). The mean amplitude at the trained frequencies was significantly

higher for the overall data (Fig. 3c, $t(17) = 24.17$ and $p = 1.3 \times 10^{-14}$). However, neither the experimental data

nor the overall data correlated with the RER during the task (Fig. 3d, $r = 0.04$ and $p = 0.88$ for experimental

data, $r = 0.16$ and $p = 0.53$ for overall), suggesting that the RER does not reflect overall sensorimotor learning.

Figure 3e illustrates that the peak amplitudes for the fast components gradually increased with

frequency (slope > 0, $t(17) = 3.39$ and $p = 0.0035$), while those for the slow components gradually decreased

(slope < 0, $t(17) = 8.07$ and $p = 3.3 \times 10^{-7}$). A statistically significant difference between fast and slow

components was found at the highest frequency of 32 Hz ($t(17) = 7.45$ and $p = 9.4 \times 10^{-7}$), with a marginal

difference at the lowest frequency of 2 Hz ($t(17) = 1.32$ and $p = 0.21$). The mean amplitudes at the trained

frequency were higher than those at the untrained frequency (Fig. 3f, $t(17) = 10.98$ and $p = 3.8 \times 10^{-9}$ for the

fast component, $t(17) = 6.71$ and $p = 3.7 \times 10^{-6}$ for the slow component). Interestingly, a statistically significant

negative correlation was found for the fast component (Fig. 3g, $r = -0.58$ and $p = 0.011$), but a positive

correlation was found for the slow component (Fig. 3h, $r = 0.67$ and $p = 0.0024$) with the RER during the task.

Note that only the slow component and the RER before the experiment were significantly correlated ($r = -0.34$

and $p = 0.17$ for the fast component, $r = 0.47$ and $p = 0.048$ for the slow component), suggesting that RER

can predict the slow motor memory to be learned. However, this correlation was weaker than during the task

(Fig. 3h), possibly owing to the time at baseline being slightly too far from the training phase.

Fast and slow motor memories were negatively correlated ($r = -0.71$ and $p = 8.6 \times 10^{-4}$) because an

increase in one memory decreased the other, suggesting that the opposite correlation shown in Figure 3g and

h would not satisfy the assumption that the RER reflects both of them. Therefore, we further examined

whether the RER reflected the fitting parameters for the fast and slow, namely the retention and learning

factors (Fig. 3i and j). Interestingly, we found a negative correlation with the retention factor ($r = -0.53$ and $p = $

0.025) and a positive correlation with the learning factor ($r = 0.63$ and $p = 0.0053$) of the slow component.

However, no significant correlations were found for the fast component ($r = -0.35$ and $p = 0.15$ for retention

factor, $r = -0.14$ and $p = 0.59$ for learning factor). These findings suggested that participants with a higher

RER demonstrated faster learning but also exhibited greater forgetting in slow motor memory. Given that both

changes in hand motion away from and toward the 0° target direction necessitate the updating of slow motor
memory in the dynamic VMR, the results suggest that the slow component of sensorimotor learning is more
flexible. The RER does not alter the execution and learning of reaching movements; instead, it reflects the
slow component of sensorimotor learning, characterized by a motor memory with a relatively longer time
constant.

Long-term retention with glucose intervention: Experiment 2

We found that the RER reflected the slow component of sensorimotor learning in Experiment 1. The slow
component serves as an estimate, offering indirect means to measure long-term memory. To directly
investigate whether the RER reflects long-term memory, we conducted a test for 24-hour retention, wherein
the slow components played a significant role (Fig. 4a) (Joiner and Smith, 2008). Furthermore, while the RER
remained constant in Experiment 1, several methods have been proposed to alter it, including circadian
rhythm (Zitting et al., 2018), dietary factors, and genetic predispositions (Toubro et al., 1998). Since glucose
intake is a simple but powerful way to increase the RER (Tellez et al., 2013), we tested whether glucose
intervention resulted in improved long-term motor memory (Fig. 4a).

We recruited an additional 24 human participants (14 males and 10 females aged 19-39 years) for
Experiment 2. This was a two-day experiment in which they adapted to a constant VMR on day 1 and tested
retention on day 2 while monitoring the RER (Fig. 4b). The participants were divided into two groups: water
and glucose (WG, n = 16) and water alone (Wa, n = 8) (Fig. 4c). On day 1, the WG group consumed a gel-
type drink containing 50 g glucose (200 kcal) and water, whereas the Wa group consumed only water. Before
and after the intervention, resting-state gas exchange was recorded for 20 min to observe the rapid effects of
glucose intervention on the RER. After a short baseline of 20 trials, they learned a constant VMR for 220 trials
while the gas exchange was consistently recorded. On day 2, the WG and Wa groups returned to the
laboratory, and we examined 20 min of resting-state gas exchange, followed by the retention test without
visual feedback (Fig. 4b), which can measure long-term memory but not learning, while the gas exchange was
recorded.

Increase in RER following glucose intake

We found that the RER increased only after glucose intake. The RER before the intervention was not
significantly different between the WG and Wa groups (Fig. 4d, $t(22) = 0.85$ and $p = 0.41$). The RER
immediately after the intervention did not increase but rather slightly decreased from the pre- to post-
intervention phases ($t(15) = 2.23$ and $p = 0.041$ for WG, $t(7) = 1.24$ and $p = 0.25$ for Wa) and was not
significantly different between these groups ($t(22) = 0.42$ and $p = 0.68$). Notably, however, we found a
significant increase in the RER from the pre-intervention to task phases only in the WG group ($t(15) = 5.94$
and $p = 2.7 \times 10^{-5}$ for WG, $t(7) = 1.15$ and $p = 0.29$ for Wa), with a significant difference noted between these
groups ($t(22) = 2.77$ and $p = 0.011$). The increase in the RER did not persist for the next day, and no
significant differences were found between these groups in either the pre-retention ($t(22) = 0.43$, $p = 0.67$) or
retention phases ($t(22) = 0.07$ and $p = 0.95$). This result suggests that the RER can be successfully controlled
by glucose intervention. Notably, there were no changes in the RER during the training phase in the Wa

group, providing important evidence that sensorimotor learning does not affect the RER.

Association between individual RER responses to glucose intake and retention

We observed basic trends in the VMR learning curves. After short baseline periods, the hand movement
directions gradually shifted toward ideal learning (Fig. 4e). The resulting learning levels were 24.2° ± 0.6° for
the WG group and 21.8° ± 2.6° for the Wa group, which were not significantly different (t(22) = 1.25 and p =
0.23). The learning amounts at the retention period on day 2 were obviously reduced from the end of the
training period, but were still greater than the baseline at 5.5° ± 0.7° for the WG group but 3.9° ± 0.8° for the
Wa group. Unfortunately, the retention amounts were not significantly different between groups (t(22) = 1.36
and p = 0.19). Because the RER in the training phase was so variable (Fig. 4d), a comparison at the group
level would not reveal the effects of glucose intake on retention.

The results of Experiment 1 suggested that RER during the task reflected slow motor memory (Fig.
3h). To provide a consistent analysis, we performed a correlation of retention on day 2 with RER at three
representative phases: in the pre-intervention, during the task on day 1, and during the task on day 2. We
found no significant correlation between the retention and the RER in the pre-intervention phase (r = 0.16 and
p = 0.45) or during the task on day 2 (r = 0.31 and p = 0.14), but a significant correlation was found with the
RER during the task on day 1 (r = 0.55 and p = 0.0051) (Fig. 4g). The results indicated that the original RER
that the participants inherently had in the pre-intervention phase did not predict retention, but the intervention
RER that increased with glucose intake in the training phase predicted retention, suggesting that the control of
RER via glucose intake can increase retention at the intra-individual level.

How much can retention be improved by a 200 kcal glucose intake? Using the regression parameters,
slope and offset, from the RER during the task on day 1 to retention on day 2 (Fig. 2g), we predicted the
retention levels if the participants inherently had the original RER during training. We also projected the RER
during the task on day 1 onto the regression line and calculated the increase in RER within individuals (See
Methods section). Surprisingly, the glucose intake improved up to 21.4% ± 4.1% in the WG group (t(15) =
5.25 and p = 9.8×10^{-5}), but the water alone did not improve at −5.1% ± 6.2% in the Wa group (t(7) = 0.83
and p = 0.43) (t(22) = 3.67 and p = 0.013 between the two). The results demonstrate that a simple glucose
intervention improves long-term motor memory acquisition.

DISCUSSION

Respiration plays a crucial role in energy production and supports neural activity and plasticity (Lee et al.,
2013; Oakes et al., 2004; Sokoloff, 1977; zur Nedden et al., 2011), which are essential for sensorimotor
learning. Breathing patterns provide detailed information about whole-body metabolism including movement
intensity (Abram et al., 2022; Huang et al., 2012; Selinger et al., 2015; Shadmehr et al., 2016); however, it is
largely unknown whether they are also a prominent index of brain metabolism, that is, whether breathing
patterns predict sensorimotor learning. According to the two-state sensorimotor learning model (Smith et al.,
2006), the faster component reflects short-term memory, whereas the slower component reflects long-term
memory. We investigated whether the metabolic mode estimated by the RER reflects short-term or long-term

memory. The results showed that the RER was not influenced by the execution or learning of reaching
movements (Fig. 2). Instead, the RER predicted the amount of estimated slow rather than fast motor memory
(Fig. 3). We also investigated whether metabolic interventions could improve long-term memory, that is, next-
day retention. To accomplish this, we provided 200 kcal of glucose, which immediately shifted the participants
towards more carbohydrate utilization. Surprisingly, this intervention not only increased the RER but also
improved next-day retention by up to 21% (Fig. 4). Together, with two separate cohorts of human participants,
these findings showed that the metabolic mode estimated by the RER can serve as an indicator of long-term
motor memory, particularly at the intra-individual level, implying a potential avenue for enhancing human
motor performance.

Methodological consideration

The metabolic mode estimated by the RER varied widely from 0.74 to 0.97 among individuals even at
baseline, which is very similar range to the previous findings (e.g., from 0.718 to 0.927 in (Goedecke et al.,
2000)). Previous studies have shown that during low-intensity activities such as unloaded or lightly loaded
pedaling—activities likely to exert more load than the reaching movements in our study—RER does not
significantly deviate from baseline levels (Wasserman et al., 1973). This is consistent with our findings (Fig.
2e). We also found no explanation of muscle activity on RER (Fig. 2f) as well as unchanged RER between at-
task vs at-break (Fig. 2g). Furthermore, the RER is highly stable even during sensorimotor learning (Figs. 2e
and 4d gray) and no correlation against sensorimotor learning amplitudes (Fig. 3d). The results suggest that
the execution and learning of reaching movement do not affect RER accordingly.

We believe that a comparison between the presence and absence of glucose intake provides a direct
examination of how the RER intervention affects long-term motor memory. However, the baseline RER
variability was high (SD across participants: 0.049) compared to the increased RER (0.046 ± 0.008 from pre-
intervention to task phases) with only 200 kcal of glucose intake, implying that the glucose intervention on
next-day retention (Fig. 4f) would be buried under large individual differences at baseline. The long-term
motor memory would be enhanced by maximizing the physiological effects of glucose intake, such as the
glucose dose or glucose digesting time. Notably, a literature review showed enhancement of memory with
glucose intake is limited even in the other memory types, such as verbal word memory (Smith et al., 2011).
Our results underscore a new type of memory retention improvement at the intra-individual level, and it needs
to turn the tide for investigation of physiological basis for sensorimotor learning.

Glucose intake may contribute to other cognitive improvements during learning. For example, glucose
intake may induce a positive mood and reduce hunger, perhaps leading to an upregulation of motivation and
attention. These cognitive improvements would affect the fast component, reflecting the use of an explicit
strategy (McDougle et al., 2015), rather than the slow component. Thus, if glucose intake elicited only
cognitive improvements, the slow component would not change. Our findings that the glucose intake affects
slow, long-term motor memory suggest that cognitive improvements from the glucose intake are not the main
driver of its enhancement of long-term motor memory.

The cortical metabolism and time-scale dynamics in sensorimotor Learning

Long-term motor memory is shaped by multiple factors, and our computational approach serves to simplify the
intricate interplay of biological factors of sensorimotor learning, such as protein synthesis and gene
expression (Kandel et al., 2014). For instance, neural plasticity is regulated by protein synthesis (Luft et al.,
2004). The metabolic demands associated with sensorimotor learning may further influence neuroplasticity
through mechanisms involving ATP availability and mitochondrial function. However, the specific connections
between metabolic mode and processes of motor memory retention remain unclear and require further
investigation.

Another possible explanation is cortical-area-dependent time constants (Kim et al., 2015). The
parietal areas and cerebellum, which are considered as key regions in the neural basis of sensorimotor
learning, have been suggested to exhibit neural activity changes at different time scales—fast for the parietal
areas and slow for the cerebellum. A study by Vaishnavi and colleagues demonstrated that these areas also
differ in their metabolic profiles, particularly in terms of aerobic glycolysis (Vaishnavi et al., 2010). Specifically,
aerobic glycolysis is markedly elevated in the medial and lateral parietal cortices, while the cerebellum
exhibits levels well below the brain average. These findings suggest a potential coupling between regional
metabolic signatures and time-dependent learning processes, suggesting that metabolic measures may serve
as physiological markers of engagement in distinct brain systems during sensorimotor learning. Future work
should aim to integrate metabolic modeling with neural network dynamics to develop a more comprehensive
picture of motor memory formation.

323 The RER would indicate physiological idling state for sensorimotor learning

The brain accounts for 20% of total energy expenditure (Raichle and Gusnard, 2002). This proportion
underscores that RER may indeed provide valuable insights into the neural states associated with learning
and memory. However, it is important to note that the RER is an indicator of the net metabolic mode of the
entire body. Although the contribution of brain metabolism is significant, it is certainly not a large contribution.
One could argue that the effect of cortical metabolism on RER is limited. How does RER reflect long-term
motor memory?

It should be noted that the glucose intake did not immediately increase the RER in the post-
intervention phase, but rather slowly increased it during the subsequent training phase (Fig. 4d). This
temporal pattern suggests a gradual physiological response that unfolds over time, that is, insulin secretion.
Insulin secretion is not an instantaneous response but occurs gradually in response to increasing levels of
glucose in the bloodstream. Insulin secretion is usually accompanied by an increase in RER (Dickson et al.,
1924), and presumably regulates glucose levels in the body and neuronal and glial cells (Milstein and Ferris,
2021; Uemura and Greenlee, 2006) for future energy production. Thus, this increase in cerebral glucose
levels would meet the cortical energy needs, especially for learning and memory.

Interestingly, insulin modulates memory formation, as evidenced by studies on Alzheimer's disease, a
model of impaired declarative memory function. Alzheimer's disease is accompanied by disrupted glucose
regulation and utilization (Kyrtata et al., 2021); however, surprisingly, insulin injection improves memory
function in Alzheimer's patients (Craft et al., 1996; Reger et al., 2008). Consistent results have been found in

healthy subjects, indicating that the intranasal administration of insulin upregulates memory function (Benedict
et al., 2007, 2004). Furthermore, insulin resistance in healthy obese individuals impairs their ability to learn;
however, the administration of liraglutide, which temporarily stimulates insulin release, can restore this ability
(Hanssen et al., 2023). Collectively, these studies suggest that insulin secretion and its receptors regulate
memory. In line with these studies, our results suggest that the RER, which may be linked to insulin release,
would reflect a physiological idling state for sensorimotor learning.

METHODS

Participants

In total, 18 participants (18 males, aged between 21 and 52 years) in Experiment 1 and 27 participants (16
males and 11 females, aged between 19 and 39 years) in Experiment 2 were recruited; however, we removed
data from three participants in Experiment 2 owing to their data quality (See Data Analysis). All participants
were right-handed, reported no neurophysiological disorders, and provided written informed consent prior to
the experiments in accordance with the Declaration of Helsinki. The Committee on Ergonomic Experiments of
the National Institute of Advanced Industrial Science and Technology (No. HF2022-1183) approved the
protocol of Experiment 1 and the research ethics committee of the University of Tokyo (No. 23-204) approved
the protocol for Experiments 1 and 2, respectively.

Reaching movement task

Participants performed rapid 20-cm point-to-point arm reaching movements with their right hand while sitting
on a chair positioned at the center of the task space. We used an exoskeleton robotic manipulandum in
Experiment 1 (KINARM Exoskeleton Lab, Bkin Technologies, Canada), which can prevent unwanted fatigue
and maintain a constant posture to reliably record muscle activity with electromyography (EMG), while we
used a digitizing tablet and pen encased in a light grip in Experiment 2 (Wacom Intuos Pro L, Wacom, Japan)
to reduce muscle activity that might affect the breathing data. The sampling rates were 1000 Hz in Experiment
1 and 200 Hz in Experiment 2, and the refresh rates of the visual screen were both 60 Hz.

At the beginning of each trial, the participants were asked to hold a white circle (1.0 cm in diameter)
representing their hand position on a red start circle (1.5 cm). One second later, the red target circle (1.5 cm)
was displayed 20 cm in front of the start circle. One second later, the target color changed to green, which
was a cue to initiate a forward-reaching movement. Upon reaching the green target, the target color changed
to blue and the start circle turned green, which was a cue to initiate a backward-reaching movement. Outback
movements were completed, and the next trial started after an inter-trial interval of 1 s. A preliminary
experiment showed that the outback movements took approximately 2 s. A trial duration of approximately 5 s
was chosen because each trial would involve at least one breath, which occurred approximately once every 4
s. After the participants familiarized themselves with the task, we introduced VMR, which involved angularly
shifting the cursor position.

To dissociate the fast and slow components of sensorimotor learning in Experiment 1, we designed

the sum of sinusoidal functions(Miyamoto et al., 2020). The frequencies of the functions were [2, 4, 8, 16, 32]

Hz over 900 trials; that is, the perturbation sequence combined gradual components (2 and 4 Hz) and sudden

components (16 and 32 Hz). The amplitude of the function was 10°. The maximum amplitudes of the dynamic

perturbation were ±30.9° and maximum changes in a trial were ±4.3°. We counterbalanced the signs of the

amplitudes of the 18 participants. A total of 900 training trials were divided into 12 blocks, with an

approximately one-minute break between blocks.

In Experiment 2, to test the 24-h retention with the glucose intervention, participants adapted to a

constant VMR perturbation on day 1 and were tested for motor memory without visual feedback on day 2. The

constant perturbation was 30°, which was counterbalanced across the 24 participants. There were 220 trials

after 20 null trials on day 1 and 40 non-visual trials on day 2. The one-minute break was given every 10 trials

to remove short-term memory and encourage long-term memory (Hadjiosif et al., 2023).

Gas exchange

We recorded the O2-CO2 gas exchange using a portable metabolic measurement system in the breath-by-

breath mode (COSMED K5, COSMED Srl, Italy). In Experiment 1, we started recording the RER from the

familiarization phase to the end of the experiment to examine how the baseline RER in the pre-experiment, as

well as training the RER in the experiment, can predict the fast and slow components of sensorimotor learning

and how the RER alters during sensorimotor learning. In Experiment 2, to investigate how glucose intake

modulates the RER, we recorded before and after the intervention phase (pre- and post-intervention phases)

on day 1. Both phases took 20 min, as determined by the data analysis in Experiment 1, and the data

reliability, calculated by the mean squared error against the mean of the whole data, was saturated at 20 min.

The gas exchange was recorded continuously during the training phase. On day 2, we recorded the pre-

experiment baseline gas exchange for 20 min and then continuously recorded it during the retention phase.

Before the experiment, the K5 system was calibrated according to the manufacturer's instructions.

Electromyography

In Experiment 1, muscle activity was recorded using a wireless surface electromyography system (WavePlus

EMG system, Cometa, Italy). The wireless system had a very low and consistent delay - 14 ms reported by

the product specifications - which had little to no impact on the results in reaching movements that took 1980

± 87 ms. Data were recorded from four muscles: the pectoralis major, posterior deltoid, biceps brachii, and

triceps brachii, which govern large portions of shoulder extension/flexion and elbow extension/flexion during

horizontal reaching movements. An electrode size was 41 × 15.5 × 11.3 mm and the inter-electrode distance

was 20 mm (Pico EMG, Cometa, Italy). The electrode sites were carefully determined by visually checking the

data quality during the ballistic reaching movements before the experiment. The EMG and kinematic data

were recorded using an A/D signal converter (PCI-6229, National Instruments, USA) at a sampling rate of

1000 Hz. The EMG signals were amplified (×1000) and band-pass filtered from 10 to 500 Hz via the

hardware.

Glucose administration

In Experiment 2, we examined the effects of glucose intake on RER and 24-h retention. We asked the

participants in both the WG and Wa groups not to consume any food or drink with sweeteners for at least 1 h

before the experiment, to maximize the effects of the intervention. The WG group consumed water and gel-

type drinks (Glucorescue, Arkray, Japan), while the Wa group consumed only water during the intervention

phase; therefore, they had to take off their masks to record gas exchange. The main ingredient of the gel-type

drinks was 50 g of glucose, corresponding to 200 kcal. Because the WG group took 387 ± 40 seconds, the

Wa group also waited for approximately 360 seconds, to match the periods of the intervention phase.

Data analysis

We smoothed the handle position using a fourth-order Butterworth filter with a cutoff frequency of 10 Hz.

Subsequently, we found the midpoints of the forward movement, that is, 10 cm away from the start position.

This is almost identical to the position at peak velocity, which represents the feedforward predictive

component of movement. We computed the movement directions at the midpoints relative to the target

direction, because VMR is an angular perturbation. We performed outlier removal in each trial of each

experiment if the data were outside the median ± 3 IQRs across participants, which was nearly 0.05% under

the hypothetical standard distribution. We removed one participant from Experiment 2 because the data were

very distant from the others, especially in the retention phase (38 out of 40 retention trials were outliers).

We used a two-state model to represent short- and long-term memory in sensorimotor learning(Smith

et al., 2006). The model considered that the total motor outputs (x) were determined by the sum of two distinct

memories ($x = x_f + x_s$) and that the memories are parallelly updated with the error ($e = x + p$, p is VMR) as

$x_f(n + 1) = \alpha_f x_f(n) + \beta_f e(n)$ and $x_s(n + 1) = \alpha_s x_s(n) + \beta_s e(n)$, where f and s indicate labels of fast and

slow components of sensorimotor learning, n indicates the trial number, and α and β are retention and

learning factors. The retention and learning factors are thought to be different between the components: $0 <$

$\alpha_f < \alpha_s < 1$ and $0 < \beta_s < \beta_f$. We fitted the model to individual data with expectation maximization

methods(Albert and Shadmehr, 2018), which allowed us to robustly estimate the parameters of the model

from the noisy data; thus, we did not have to obtain a number of samples. A previous report showed that

almost 20-95% fewer participants were needed from the power analysis compared to the least-squares

algorithm (Albert and Shadmehr, 2018). In Experiment 1, we recruited a number of participants comparable to

those in previous studies (Albert and Shadmehr, 2018). Basically, we utilized the original code that is available

at http://shadmehrlab.org/tools, although we fixed a few initial parameters for the parameter searching:

narrower ranges of initial fast and slow motor memories from ±30° to ±5°. Narrower ranges that emphasized

zero learning amounts at baseline were more reflective of the current situation.

The O2-CO2 gas exchange ratio was used to compute the RER offline for each breath. The RER was

recorded breath-by-breath and it was obtained at discrete times over a few seconds. For the RER during

tasks, we performed linear interpolation between the RER to match the kinematic data with a sampling rate of

1000 Hz and used the RER at the midpoint of movement as a representative index in the trial. The range of

RER was theoretically defined as 0.7-1.0, the outsides of which were regarded as outliers. In particular, we

removed two participants in Experiment 2 because about 35% of trials were determined as outliers, which is a

high ratio compared with the others (7.5 ± 1.0%). For the steady-state RER, such as the pre-experiment

phase in Experiment 1 and the pre- and post-intervention phases in Experiment 2, we computed the mean of

the RER from the individual breathing data after outlier removal in each breathing data.

EMG data were subtracted from the mean and rectified for each trial. Here, we used muscle activity to

estimate the intensity of movement execution, as it affects the RER(Goedecke et al., 2000). Therefore,

instead of EMG activity during the middle movement, we computed a trial-wide summation of the EMG of

each muscle. We then performed self-normalization with the mean across trials for each muscle. This allowed

us to examine how muscle activity changes with sensorimotor learning, independent of the different EMG

responses of the muscles.

To examine whether the intensity of the reaching movements influenced the subsequent RER in

Experiment 1, we computed a linear regression from muscle activity to RER with a lag within blocks. There

were four muscle activities [900 × 4] and RER [900 × 1] over 900 trials. Seventy-five trials were performed in

each of 12 blocks; therefore, we cut the blocks for both activities [75 × 12 × 4] and RER [75 × 12 × 1], and slid

them every single trial within blocks, as in the preceding trials [1, …, 75-i] and the RER with the i-th lag [i-75].

From the regression results, we computed the coefficient of determination (R^2), which indicates the

percentage of variance explained. For comparison, we randomly permuted the blocks 5000 times, performed

the same procedures as described above, and obtained the mean R^2 across the iterations. We performed this

for each participant and then tested the differences in R^2 between the experimental and permuted data with

each of the 75 lags. To compensate for the 75 repetitive tests, we corrected the p-values using the FDR.

For the estimates of intra-individual enhancements in the 24-h retention in Experiment 2, we

computed the changes from baseline RER to intervention RER over the regression line ($y = ax + b$),

indicating how much glucose intake increased next-day retention along with the RER changes. Using the

slopes (a) and offset (b) of the regression line, we first predicted the retention levels from the baseline RER

($y_{pre} = ax_{pre} + b$), which indicates the retention of the original RER. We then projected the RER during the

task on day 1 onto the regression line ($y_{task} = ax_{task} + b$) and calculated the extent to which the RER

increased within individuals as follows: ($y_{task} - y_{pre})/y_{pre} \times 100$ (%). Thus, this value estimates the intra-

individual changes in retention due to glucose intake. For comparison, we performed the same procedure in

the Wa group.

We performed one-/paired-sample or two-sample two-tailed t-tests to compare the mean variables

and performed Pearson's correlation when testing the covering variables. The significant p-value was set at

0.05.

Data and code availability

482   We prepare an online repository on Zenodo (https://zenodo.org/) that contains the data and codes in the

483   current study. The uploads will be completed when the paper is published.

Figure 1

Figure 2

Figure 3

Figure 4

FIGURE CAPTIONS

Figure 1. Respiratory exchange ratio and its potential role for sensorimotor learning.

a. The metabolic cycle converts oxygen (O2) and macronutrients, carbohydrates, and fats, into carbon dioxide (CO2) and adenosine triphosphate (ATP), the energy currency that supports neural activity, plasticity, and muscle activity. Breathing, which involves O2-CO2 exchange, plays an important role in this process, but it is unknown whether breathing can predict learning and memory, which are influenced by neural activity and plasticity. b. The respiratory exchange ratio (RER), calculated as the ratio of CO2 production (VCO2) to O2 consumption (VO2), reflects the metabolic preference for using carbohydrates or fats as fuel. In the cortex, where only carbohydrates are used, the RER approaches 1.0. c. We designed Experiment 1 to test whether the RER, calculated by O2-CO2 exchange, represents sensorimotor learning of reaching movements while recording muscle activity. d. Visuomotor rotation (VMR), which angularly shifts the position of the visual cursor from the hand position, was introduced to elicit sensorimotor learning. This example shows counterclockwise VMR; the leftward visual error shifts the reaching movement to the right. e. A theory posits that two motor memories with different time constants, fast and slow, are stored in sensorimotor learning. The short- or long-time constants of sensorimotor learning, which presumably correspond to neural activity and plasticity. f. To accurately estimate motor memories with varying time constants, a dynamic VMR comprising five sinusoidal functions with different frequencies ranging from 2 to 32 Hz. It is hypothesized that rapidly changing VMR induces fast motor memory formation, while gradually changing VMR elicits the development of slow motor memory. g. The sum of the sinusoidal function is used in the reaching movement task.

Figure 2. Few effects of execution and learning of reaching movements on RER.

a. Movement directions changed with the VMR. Error bars show the standard errors of their means. b. Significant peaks at the trained frequencies were found with the Fourier transformation for a series of movement directions. c. The average of the peaks was significantly larger than that at the untrained frequencies. d,e. Self-normalized muscle activity (d) was high at the beginning of the experiment and decreased rapidly over the first 300 trials. In contrast, RERs (e) were very stable over the course of the experiment. f,g. Regression analysis with trial lags revealed that self-normalized muscle activity cannot explain the changes in RER (f, permutation test, $p > 0.05$ for all lags with FDR correction). Furthermore, there was no change between task and break (g). h,i. No relationships with possible factors, peak velocity (h) and reaction time (i), were observed. j. No differences between RER in the first and last block; Sensorimotor learning did not modulate RER. k,l. No obvious peaks at the trained frequency were found with the Fourier transformation for a series of RER (k), although the mean amplitude was statistically higher at the trained frequency (l).

Figure 3. Individual RER reflects slow components of sensorimotor learning

a-c. The two-state model captured the sensorimotor learning processes (a). Obvious peaks of the overall, fast, and slow components at the trained frequency were revealed by the Fourier transformation (b), and the mean amplitudes at the trained frequency were higher than those at the untrained frequency (c), analogous to the

experimental data shown in Figure 2c. d. The experimental and fitted data were not correlated with the RER

during the task. e,f. The amplitudes of the fast and slow components were increased and decreased,

respectively, with the trained frequency (e). The mean amplitudes at the trained frequency were higher than

those at the untrained frequency (f). g,h. The mean amplitudes of the fast (g) and slow (h) components were

negatively and positively correlated with the RER during the task, respectively. i,j. For comparison with the

model parameters, the RER during the task was correlated only with the slow retention factor (i) and the slow

learning factor (j).

Figure 4. RER predicts glucose-intake enhancement of long-term motor memory

a. Based on the results of Experiment 1, which showed that the RER reflects slow motor memory, we tested

the hypothesis that increasing the RER with glucose intake would improve 24-hour retention. b. Experiment 2

was designed to examine how much of the 30° constant VMR learned on day 1 was retained on day 2 during

reaching movements without visual feedback. c. On day 1, the RER was recorded for 20 min before and after

an intervention in which one group received 200 kcal glucose (WG) while the other group did not (Wa). Both

then performed the same training task. On day 2, the RER was recorded for 20 minutes, followed by the

retention task. d. No difference in the original RER was found in the pre-intervention phase. Furthermore, the

change in the RER was not found in the post-intervention phase but in the training phase. The effects did not

persist on day 2; no differences were found in the pre-retention and retention phases. e,f. We observed the

typical learning curve for VMR training on day 1 (left panel in e). On day 2, the amount of learning largely

dropped, but we can still see significant retention (right panel in e). Notably, no differences were found

between the groups (f). g. The correlation analysis showed a significant positive relationship between

retention and intervention RER, but no relationships against baseline RER or RER on retention. h. Changes in

the RER from the pre-intervention phase to the training phase estimated a 21.4% ± 4.1% increase in retention

with the glucose intervention.

References

Abram SJ, Poggensee KL, Sánchez N, Simha SN, Finley JM, Collins SH, Donelan JM. 2022. General
variability leads to specific adaptation toward optimal movement policies. *Curr Biol* **32**:2222-2232.e5.

Albert ST, Shadmehr R. 2018. Estimating properties of the fast and slow adaptive processes during
sensorimotor adaptation. *J Neurophysiol* **119**:1367–1393.

Benedict C, Hallschmid M, Hatke A, Schultes B, Fehm HL, Born J, Kern W. 2004. Intranasal insulin improves
memory in humans. *Psychoneuroendocrinology* **29**:1326–1334.

Benedict C, Hallschmid M, Schultes B, Born J, Kern W. 2007. Intranasal insulin to improve memory function in
humans. *Neuroendocrinology* **86**:136–142.

Craft S, Newcomer J, Kanne S, Dagogo-Jack S, Cryer P, Sheline Y, Luby J, Dagogo-Jack A, Alderson A.
1996. Memory improvement following induced hyperinsulinemia in Alzheimer's disease. *Neurobiol*
Aging **17**:123–130.

Dickson BR, Eadie GS, MacLeod JJR, Pember FR. 1924. The effect of insulin on the respiratory exchange of
normal animals. *Q J Exp Physiol* **14**:123–149.

Franklin DW, Osu R, Burdet E, Kawato M, Milner TE. 2003. Adaptation to stable and unstable dynamics

achieved by combined impedance control and inverse dynamics model. *J Neurophysiol* **90**:3270–
3282.

Gilden DL, Thornton T, Mallon MW. 1995. 1/f noise in human cognition. *Science* **267**:1837–1839.

Goedecke JH, St Clair Gibson A, Grobler L, Collins M, Noakes TD, Lambert EV. 2000. Determinants of the
variability in respiratory exchange ratio at rest and during exercise in trained athletes. *Am J Physiol*
Endocrinol Metab **279**:E1325-34.

Hadjiosif AM, Morehead JR, Smith MA. 2023. A double dissociation between savings and long-term memory
in motor learning. *PLoS Biol* **21**:e3001799.

Hanssen R, Rigoux L, Kuzmanovic B, Iglesias S, Kretschmer AC, Schlamann M, Albus K, Edwin Thanarajah
S, Sitnikow T, Melzer C, Cornely OA, Brüning JC, Tittgemeyer M. 2023. Liraglutide restores impaired
associative learning in individuals with obesity. *Nat Metab* **5**:1352–1363.

Harms KJ, Rioult-Pedotti MS, Carter DR, Dunaevsky A. 2008. Transient spine expansion and learning-
induced plasticity in layer 1 primary motor cortex. *J Neurosci* **28**:5686–5690.

Heald JB, Lengyel M, Wolpert DM. 2021. Contextual inference underlies the learning of sensorimotor
repertoires. *Nature* **600**:489–493.

Himwich HE, Nahum LH. 1932. The respiratory quotient of the brain. *Am J Physiol* **101**:446–453.

Huang HJ, Kram R, Ahmed AA. 2012. Reduction of metabolic cost during motor learning of arm reaching
dynamics. *J Neurosci* **32**:2182–2190.

Joiner WM, Smith MA. 2008. Long-term retention explained by a model of short-term learning in the adaptive
control of reaching. *J Neurophysiol* **100**:2948–2955.

Kandel ER, Dudai Y, Mayford MR. 2014. The molecular and systems biology of memory. *Cell* **157**:163–186.

Kida H, Tsuda Y, Ito N, Yamamoto Y, Owada Y, Kamiya Y, Mitsushima D. 2016. Motor training promotes both
synaptic and intrinsic plasticity of layer II/III pyramidal neurons in the primary motor cortex. *Cereb*
Cortex **26**:3494–3507.

Kim S, Ogawa K, Lv J, Schweighofer N, Imamizu H. 2015. Neural substrates related to motor memory with
multiple timescales in sensorimotor adaptation. *PLoS Biol* **13**:e1002312.

Kyrtata N, Emsley HCA, Sparasci O, Parkes LM, Dickie BR. 2021. A systematic review of glucose transport
alterations in Alzheimer's disease. *Front Neurosci* **15**:626636.

Landi SM, Baguear F, Della-Maggiore V. 2011. One week of motor adaptation induces structural changes in
primary motor cortex that predict long-term memory one year later. *J Neurosci* **31**:11808–11813.

Lee HU, Yamazaki Y, Tanaka KF, Furuya K, Sokabe M, Hida H, Takao K, Miyakawa T, Fujii S, Ikenaka K.
2013. Increased astrocytic ATP release results in enhanced excitability of the hippocampus. *Glia*
61:210–224.

Li CS, Padoa-Schioppa C, Bizzi E. 2001. Neuronal correlates of motor performance and motor learning in the
primary motor cortex of monkeys adapting to an external force field. *Neuron* **30**:593–607.

Luft AR, Buitrago MM, Ringer T, Dichgans J, Schulz JB. 2004. Motor skill learning depends on protein
synthesis in motor cortex after training. *J Neurosci* **24**:6515–6520.

McArdle WD, Katch FI, Katch VL. 2010. Essentials of exercise physiology, 4th ed. Philadelphia, PA: Lippincott
Williams and Wilkins.

McDougle SD, Bond KM, Taylor JA. 2015. Explicit and Implicit Processes Constitute the Fast and Slow
Processes of Sensorimotor Learning. *J Neurosci* **35**:9568–9579.

Milstein JL, Ferris HA. 2021. The brain as an insulin-sensitive metabolic organ. *Mol Metab* **52**:101234.

Miyamoto YR, Wang S, Smith MA. 2020. Implicit adaptation compensates for erratic explicit strategy in human
motor learning. *Nat Neurosci* **23**:443–455.

Oakes TR, Pizzagalli DA, Hendrick AM, Horras KA, Larson CL, Abercrombie HC, Schaefer SM, Koger JV,
Davidson RJ. 2004. Functional coupling of simultaneous electrical and metabolic activity in the human
brain. *Hum Brain Mapp* **21**:257–270.

Paz R, Boraud T, Natan C, Bergman H, Vaadia E. 2003. Preparatory activity in motor cortex reflects learning
of local visuomotor skills. *Nat Neurosci* **6**:882–890.

Perich MG, Gallego JA, Miller LE. 2018. A neural population mechanism for rapid learning. *Neuron* **100**:964-
976.e7.

Pritchard WS. 1992. The brain in fractal time: 1/f-like power spectrum scaling of the human
electroencephalogram. *Int J Neurosci* **66**:119–129.

Raichle ME, Gusnard DA. 2002. Appraising the brain's energy budget. *Proc Natl Acad Sci U S A* **99**:10237–
10239.

Reger MA, Watson GS, Green PS, Wilkinson CW, Baker LD, Cholerton B, Fishel MA, Plymate SR, Breitner
JCS, DeGroodt W, Mehta P, Craft S. 2008. Intranasal insulin improves cognition and modulates beta-
amyloid in early AD. *Neurology* **70**:440–448.

Rioult-Pedotti MS, Friedman D, Donoghue JP. 2000. Learning-induced LTP in neocortex. *Science* **290**:533–
536.

Rioult-Pedotti MS, Friedman D, Hess G, Donoghue JP. 1998. Strengthening of horizontal cortical connections
following skill learning. *Nat Neurosci* **1**:230–234.

Selinger JC, O'Connor SM, Wong JD, Donelan JM. 2015. Humans can continuously optimize energetic cost
during walking. *Curr Biol* **25**:2452–2456.

Shadmehr R, Huang HJ, Ahmed AA. 2016. A representation of effort in decision-making and motor control.
Curr Biol **26**:1929–1934.

Short KR, Sedlock DA. 1997. Excess postexercise oxygen consumption and recovery rate in trained and
untrained subjects. *J Appl Physiol* **83**:153–159.

Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting adaptive processes with different timescales
underlie short-term motor learning. *PLoS Biol* **4**:e179.

Smith MA, Riby LM, van Eekelen JAM, Foster JK. 2011. Glucose enhancement of human memory: a
comprehensive research review of the glucose memory facilitation effect. *Neurosci Biobehav Rev*
35:770–783.

Sokoloff L. 1977. Relation between physiological function and energy metabolism in the central nervous
system. *J Neurochem* **29**:13–26.

Sun X, O'Shea DJ, Golub MD, Trautmann EM, Vyas S, Ryu SI, Shenoy KV. 2022. Cortical preparatory activity
indexes learned motor memories. *Nature* **602**:274–279.

Tellez LA, Ren X, Han W, Medina S, Ferreira JG, Yeckel CW, de Araujo IE. 2013. Glucose utilization rates
regulate intake levels of artificial sweeteners: Glucose utilization and artificial sweeteners. *J Physiol*
591:5727–5744.

Toubro S, Sørensen TI, Hindsberger C, Christensen NJ, Astrup A. 1998. Twenty-four-hour respiratory
quotient: the role of diet and familial resemblance. *J Clin Endocrinol Metab* **83**:2758–2764.

Uemura E, Greenlee HW. 2006. Insulin regulates neuronal glucose uptake by promoting translocation of
glucose transporter GLUT3. *Exp Neurol* **198**:48–53.

Vaishnavi SN, Vlassenko AG, Rundle MM, Snyder AZ, Mintun MA, Raichle ME. 2010. Regional aerobic
glycolysis in the human brain. *Proc Natl Acad Sci U S A* **107**:17757–17762.

Veuthey TL, Derosier K, Kondapavulur S, Ganguly K. 2020. Single-trial cross-area neural population
dynamics during long-term skill learning. *Nat Commun* **11**:4057.

Wasserman K, Whipp BJ, Koyl SN, Beaver WL. 1973. Anaerobic threshold and respiratory gas exchange
during exercise. *J Appl Physiol* **35**:236–243.

Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, Tennant K, Jones T, Zuo Y. 2009. Rapid formation and selective
stabilization of synapses for enduring motor memories. *Nature* **462**:915–919.

Zarahn E, Aguirre GK, D'Esposito M. 1997. Empirical analyses of BOLD fMRI statistics. I. Spatially
unsmoothed data collected under null-hypothesis conditions. *Neuroimage* **5**:179–197.

Zitting K-M, Vujovic N, Yuan RK, Isherwood CM, Medina JE, Wang W, Buxton OM, Williams JS, Czeisler CA,
Duffy JF. 2018. Human resting energy expenditure varies with circadian phase. *Curr Biol* **28**:3685-
3690.e3.

zur Nedden S, Hawley S, Pentland N, Hardie DG, Doney AS, Frenguelli BG. 2011. Intracellular ATP
influences synaptic plasticity in area CA1 of rat hippocampus via metabolism to adenosine and
activity-dependent activation of adenosine A1 receptors. *J Neurosci* **31**:6221–6234.