Ecologically and economically sustainable level of timber harvesting in boreal forests – defining the safe operating space for forest use

Mönkkönen, Mikko [1,2]

Blattert, Clemens [1,2,3]

Cours, Jérémy [1,2]

Duflot, Rémi [1,2]

Elo, Merja [1,2]

Eyvindson, Kyle [4,5]

Kouki, Jari [6]

Triviño, María [1,2]

Burgas, Daniel [1,2 *]

[1] Department of Biological and Environmental Science, University of Jyvaskyla, FI-40014 Jyvaskyla, Finland

[2] School of Resource Wisdom, University of Jyvaskyla, FI-40014 Jyvaskyla, Finland

[3] Forest Resources and Management, Swiss Federal Research Institute WSL, Zürcherstrasse 111, CH-8903, Birmensdorf, Switzerland

[4] Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, NMBU, P.O. Box 5003, NO-1433, Ås, Norway

[5] Natural Resource Institute Finland (LUKE), Latokartanonkaari 9, FI-00790, Helsinki, Finland

[6] School of Forest Sciences, University of Eastern Finland, P. O. Box 111, FI-80101, Joensuu, Finland

* Corresponding author: daniel.d.burgas@jyu.fi

Abstract

Planetary-level analyses indicate that we are exceeding the ecological limits. However, we need approaches to implement global sustainability frameworks at the regional scale. This implies using natural resources at levels that allow for their regeneration. We present a framework to define the limits beyond which ecosystems are threatened to collapse, to answer how much we can extract from ecosystems, and to manage natural resources for both human and ecosystem wellbeing. We exemplify this approach with the heath forest habitat types in Finland, representing most of the

national timber production forests. We use the IUCN Red List of Habitats to set favourable reference values for volume of deadwood, proportion of old growth forest cover and proportion of deciduous trees. Through forest growth simulation and management optimization, we found that the proportion of old growth forest is the most challenging criteria to be reached, only by 2100. This would require not only a larger use of extensive forest management practices than hitherto, but also to drastically reduce the maximum economic sustainable harvest level from the current 96% to 60%. By combining threat assessments with ecosystem modelling and management planning, this approach can support regional decision makers to make informed decisions to stay within safe limits of the use of natural resources.

Main

Humans have already exceeded six of the nine Earth System boundaries beyond which humanity can operate safely (i.e., planetary boundaries, *sensu*[1-3]). This overshoot of, for example, climate and biodiversity loss (or biosphere integrity, *sensu*[1]), poses a significant risk for ecosystems and their functions and, ultimately, also for human societies[4,5]. As human appropriation of natural resources and habitat encroachment are the main factors deteriorating the Earth system[6], we should set boundaries for human activities, yet, defining how this could be done in practice remains a significant challenge. As species always require functional ecosystems and habitats therein, a useful approach to prevent biodiversity loss is to reduce the threat for habitats. IUCN[7] defines that "an ecosystem is collapsed when it is virtually certain that its defining biotic or abiotic features are lost, and the characteristic native biota is no longer sustained". To sustain a favourable conservation status of habitats and avoid ecosystem collapse, we must bring human influence to a safe operating space, that is limited by the ecological ceiling and the social foundation—i.e., the unavoidable impacts of human activities to fulfil human needs. (Fig. 1a;[8])

Many environmental policy initiatives (e.g., Global Biodiversity Framework) address the challenge by asking "*how much conservation is enough?*" to maintain biodiversity and strive to set the minimum coverage of protected areas (e.g.,[9,10]). In addition, we need to tackle the root causes of biodiversity loss rather than only remedy their symptoms. Biodiversity is lost because we consume natural resources faster than nature can regenerate them[11]. Even when not exceeding the regeneration rate, excessive human appropriation of natural resources is a threat to biodiversity and ecosystem integrity (e.g., functional), as remaining resources (e.g., carbon, nutrients, water, space) may become insufficient. Thus, we should shift the focus and address "*how much resource use is too much?*", and quantify an ecological ceiling that prevents ecosystem collapse. Defining a

maximum biodiversity-safe level of natural resource use would provide a clear, practical, and accountable policy target that relates directly to biodiversity loss due to human activities.

Fig. 1 | Schematic overview of the framework developed to explore ecologically and economically sustainable timber harvest levels applied to Finland. a) Our study combines the frameworks of Planetary Boundaries[1] and doughnut economics[8]. We evaluate what would be the maximum level of timber extraction, i.e., timber flow and net present value (social/economic foundation) that would still allow critical forest structural features to develop and remain at necessary levels (ecological ceiling). **b**) Case study context in South and Central Finland. **c**) Current level of the three criteria defining the conservation status of heath forest habitats and, thus, used to define the ecological ceiling (see criteria values in Table 1).

Taking the boreal forest ecosystems and forestry sector as an example, we explored how to define an ecological ceiling for the safe operating space. The boreal forest is one of the largest biomes on Earth, contains approximately 45% of the world's stock of growing timber and accounts for 25-33% of wood-biomass products within the global export market[12,13]. About two-thirds of the biome's area is actively managed and harvested for timber production, with changes in structure and tree species composition, and impacts on wildlife and ecosystem functioning[14–16]. Particularly in Fennoscandia, protected areas cover only less than 5% of productive forest land and forests are considered primarily a source of biomass[17]. Therefore, habitat and resource availability for species associated with processes and structures characteristic for pristine forests has severely declined. Consequently, forest management is the most prominent threat for biodiversity[18–20]. It is evident that the ecological ceiling of forest ecosystems and their functional integrity *sensu* Rockström *et al.*[21] have largely been exceeded in Fennoscandia.

Fennoscandian boreal forests are perhaps the most thoroughly studied terrestrial ecosystems of the biosphere. From a global perspective, national economies of the Nordic countries are affluent, stable, and predictable. Therefore, Fennoscandia is an ideal test arena to find out if sustainable forestry, in ecological, economic, and social terms, is an achievable goal. Implementation of the safe operating space consequently implies securing forest habitats while maintaining timber production for the forestry sector. Indeed, the latter represent a significant economic sector, providing raw material, energy, and job opportunities to society (i.e., social foundation, Fig. 1a).

Here, we introduce a new approach to address the question of "*how much is too much?*" and test it by defining an ecological ceiling for human forest use, that is, a maximum timber harvest level below which forest ecosystems are no longer threatened. We apply this approach to boreal heath forests in Finland using the recent Red-List assessment of forest habitat types (Fig. 1b;[22]). Specifically, we ask (i) how fast the forest habitat types can be removed from the IUCN Red List, (ii) how forests should be managed to achieve this goal, and finally, (iii) what is the maximum annual harvest to stay in the safe space. To answer these questions, we use the IUCN Red-List criteria[22], which define criteria for endangerment of forest habitat types, and identify the level of those criteria that should be reached to remove the forest habitats from the risk of collapse, i.e., Favourable Reference Value (FRV, Fig. 1c). Then, we use forest growth simulations and a multi-objective optimization framework[23] to define the time, the management combination, and the maximum harvesting level needed to reach the FRVs (Fig. 2).

Fig. 2 | Workflow used for the analyses and main answers we expect with the outputs of this framework. FRVs stands for Favourable Reference Values.

Results

How much time will it take to reach favourable reference values for all forest habitat types?

The time needed to render the habitat types not threatened ranges from 40 to 80 years (2060-2100), depending on the habitat type (Fig. 3). In sub-xeric forests (representing 20% of forest area in Southern Finland;[24]) it may take till 2100 before all criteria can reach their FRV. Herb-rich forests (23% of forest area) can reach FRV already by 2060, and mesic and xeric forests (representing 50 % and 3 % of forest area, respectively) by 2080. The main limiting criteria affecting the time needed to achieve a favourable status was the proportion of old-growth forest area followed by the volume of deadwood (Fig. 3, see Table 1 in the Methods section). The required proportion of deciduous trees from total timber volume is swiftly achievable across the forest types. To project the development of forests, we used the intermediate climate change Representation Concentration Pathway (RCP) 4.5, which approximates our current climate pathway[25]. Sensitivity analyses with the "business as usual" high-emission scenario RCP8.5 did not show significant changes in the timing or criteria limiting the achievement of favourable status (Supplementary Material Fig. S1).

Fig. 3 | Possibility to achieve and exceed the favourable reference value (FRV) in South and Central Finland for the four main habitat types in heath forest (herb-rich, mesic, sub-xeric and xeric). The horizontal axis represents the target year when the FRVs were strived to be reached. Values for alternative criteria are expressed as percentages from the FRV (see Table 1), with red colour indicating that the target has not yet been achieved, white that the target has been achieved and green that the target has been surpassed.

How forests should be managed and what is the maximum harvest level to stay within a safe operating space?

The faster we strive to render forest habitat types not threatened, the more we need to rely on setting aside forests from timber harvesting (Fig. 4a). Targeting 2100, the earliest year when all forest habitat types can reach favourable reference values, we would need to set aside (i.e., no management) more than on third of forests; currently less than 6% of forest land in southern Finland is protected[26]. The optimal combination of management categories also included 40% of intensive management where the focus is more on timber production, and 25% of extensive management also targeting biodiversity conservation and other ecosystem services than timber (Fig. 4a, and Supplementary Material Fig. S2 for management sub-categories). There was variation among the forest habitat types in the optimal combination of management categories. In general, on the more productive habitat types (herb-rich and mesic forests) set aside was a more prominent option, while on less productive habitat types (sub-xeric & xeric forests) extensive management gained more prevalence (see Supplementary Material Fig. S3 for the proportion of management showed by habitat type categories).

The sooner we want to reach FRVs, the more we should give up on the maximum economically sustainable annual harvesting potential (Fig. 4b). If we set 2100 as the target year, when all forest habitat types can reach a favourable status, the ecological ceiling is a harvest level of 58% of the maximum economically sustainable annual harvesting potential. This is considerably less than the realized recent harvest level in southern Finland (Fig. 4b).

Fig. 4 | (a) Proportion of management categories and (b) maximum annual timber harvest level that would maximise the favourable reference values in all four forest habitat types. The horizontal axis represents the target-year when the FRVs were strived to be reached (but might not be reached, see Fig. 3). In (b), the vertical axis shows the harvest level relative to the maximum economically sustainable annual harvesting potential (i.e., no ecological constraints). The dashed horizontal line represents the average harvest level between the years 2016-2021 in southern Finland[27].

Discussion

Our economies are not external to nature but embedded within it[11]. Building on this premise, we operationalized the planetary boundaries and doughnut economy frameworks at large spatial scale and defined a maximum biodiversity-safe level of harvesting timber from a boreal region. Maximum harvest level was estimated so that it does not exceed the capacity of the forest ecosystems to supply biodiversity-critical habitat characteristics. Results indicate that harvesting should be capped well below current levels, but the exact level depends on how fast we want to enter safe operating space. Setting a realistic target year of 2100 requires utilizing at most 60% of the annual harvesting potential. Curbing forest harvesting to this level creates a challenge for the

Fennoscandian forest industry, which currently is largely based on producing large quantities of pulp and sawn timber for exporting. The challenge can be tackled by developing regenerative business models[28], where the emphasis is higher value-added wood products that provide livelihoods for the entire value-chain relying on wood biomass. More profoundly, resolving this challenge will require moving away from a life frame where nature is considered a mere provider of resources, to partnerships with nature that focus on nature's life-supporting processes[29].

Simultaneously with the ecological ceiling, we evaluated the social foundation by the maximum annual timber flow and monetary benefits (net present value) that are possible under the ecological ceiling. This follows the current societal values, which emphasize instrumental values and where forest resources are seen as a provider of job opportunities, economic value, and raw materials to fulfil human needs[30]. In affluent postindustrial societies, such as in Fennoscandia, we see an increasing endorsement of relational and intrinsic values such as seeing nature as a part of one's meaningful life and much more than means to an end. Such values are typically tied to local context, and thus, implementing the idea of staying within the safe operating space requires local deliberation and planning—e.g., on how to sustain livelihoods with less resource extraction in a just manner.

Staying within a safe operating space requires a constant flow of forest biomass, which, according to our results, is best achieved with a diverse combination of management regimes to ensure a diversity of habitats[31,32] and the flow of timber and non-timber ecosystem services from forests[33]. Our results suggest that high shares of both protected areas supporting biodiversity conservation, and intensively managed areas—maximizing timber production and economic benefits—are needed, complemented by extensive management. The utility of such functional zoning outcome is evident[34,35]. This requires a big change in current management practices, where intensive clear-cut forestry dominates, by increasing adoption of alternatives such as continuous cover forestry (see Supplementary Material Fig. S2).

Among the biodiversity indicators, achieving the favourable reference value for old-growth forests is the largest challenge. More than a century of intensive forest management has shifted the forest composition towards young forests[36] and recovering a high proportion of old-growth forest will require a long time. This result highlights the need to swiftly start actions for biodiversity as recovery processes and formation time of most critical microhabitats in forest ecosystems are slow. Only by acting now we can reduce any further, potentially irreversible biodiversity losses[37]. The

primary focus should be on safeguarding the remaining old-growth forests, in accordance with the EU biodiversity strategy for 2030[9]. However, old-growth forests are not only characterized by tree age but also by their structural features that are important for biodiversity and human well-being[38,39]. Hence, there exists the possibility of actively restoring deadwood and tree-related microhabitats to speed up the recovery process[40].

Deadwood is a critical resource for species in boreal forests[41] and a good indicator of forest biodiversity[42]. In boreal Fennoscandia, 20–25% of the forest-dwelling species are dependent on deadwood habitat and they constitute 60% of red-listed species[43]. The FRVs for deadwood volume can be reached relatively quickly in all forest types, suggesting the potential for relatively rapid recovery of deadwood-dependent biodiversity. However, production of large deadwood pieces, sheltering peculiar communities[44], should need extra-time. This effectively requires more set asides[36], which would eventually result in more old-growth forests, as well.

Further enhancements of this approach could entail adding ecological constraints into simulations as well as considering social factors. For example, our results remained qualitatively similar under different climate change scenarios, but this could change by including natural disturbances, which have increased and will continue to increase in frequency, intensity and spatial extent due to climate change[45,46]. Disturbances may have contrasting effects on FRVs: they contribute in creating deadwood and certain tree-related microhabitats if systematic salvage logging of damaged trees is avoided[36,47,48], while severe disturbances can also reduce the area of old-growth forests[49,50]. Ultimately, cumulative and interactive effects of human and atypical natural disturbances may lead to an ecosystem shift into a compromised functional state[45,51]. Because reaching a favourable conservation status for heath forest habitats in Finland will take several decades, the chance of facing extreme disturbance events is very high, further delaying the achievement of targeted proportion of old-growth forest. Hence, the more ambitious we are initially, the more likely we are to halt the biodiversity crisis.

In our study we consider the ecological and economic limits of the forest ecosystems to identify a safe operating space. Future studies should go one step further and quantify both safe and just Earth system boundaries[21]. Thus, a just transformation of the forest sector requires acknowledging the multiple needs and views of forest stakeholders and understanding the dynamics of the entire forest social-ecological system transition.

Methods

We used a large forest dataset representing Middle and Southern boreal regions in Finland (Fig. 1). We focused on the four main fertility site types of heath forest: herb-rich, mesic, sub-xeric, and xeric heath forest. These four site types represent together over 96% of the productive forest in Southern Finland (NFI12). Forested peatlands were excluded because a large proportion of them have been drained for forestry purposes. Consequently, their conservation status is not directly related to harvest level. We also excluded the northern boreal zone, which contains 28% of heath forests in Finland, because a large proportion of heath forests there are either barren heaths (very low timber production) and/or currently protected. Less than 5% of harvested timber volume in Finland originates from forests in the North boreal zone, and thus, excluding these forests has a small effect on required harvesting levels at the national scale. Moreover, most threatened forest-dwelling species in Finland have southern distributions[20,37].

Favourable reference values for defining an ecological ceiling.

The recent assessment of threatened habitat types in Finland[52] followed IUCN criteria, which consider changes in the quantity of the habitat type, changes in abiotic and biotic quality, and rarity. The assessment shows that three-quarters of the Finnish forest habitat types are threatened[22]; herb-rich and mesic forest habitats are classified as vulnerable (VU), i.e., at a high risk of collapse, and sub-xeric and xeric forest habitats are endangered (EN), i.e., at a very high risk of collapse.

Most of the heath forest habitat types were assessed as threatened according to changes in their abiotic and biotic quality (IUCN criteria D). However, the decline in area (IUCN criteria A) was the most important reason for old heath forests to be threatened[22]. We adopted three quality indicators, which translate the IUCN criteria into measurable characteristics, from Kouki *et al.* (2018): deadwood volume, the proportion of old forest cover, and the proportion of deciduous trees relative to the total tree volume. These indicators are considered critical features also for red-listed species in Finnish boreal forests[20]. Following the logic of the greatest risk of loss and following the IUCN guidelines, Kouki *et al.*[53] used the condition of forests in 1750 as a reference, except for the proportion of deciduous trees for which the situation in the 1960s was the reference (the value in 1750 is not available). The time point 1750 represents the earliest onset of industrial-scale exploitation of ecosystems[54] and is justified in the IUCN Red List of Ecosystems[7]. According to the IUCN criteria D, a habitat type (fertility site type in our study) is considered threatened if the indicator value has decreased more than 50% (30% for the proportion of deciduous trees) from its reference value in 1750 (1960s for the proportion of deciduous trees). We followed this logic to

address what is the required improvement in the quality indicator values that will render the heath forest habitat types not being at risk of collapse (i.e., not threatened). That is, for a currently threatened habitat type to become not threatened, the value of an indicator must improve to a level that is at least 50% of the reference value in 1750 (area of old forest and volume of deadwood), or 70% of the reference value in the 1960s (proportion of deciduous trees) and remain at that level or above ever since. We consider this level of an indicator as a favourable reference value (FRV, Table 1).

Table 1. Reference values for the quality indicators for the four fertility site types in South and Middle boreal regions in Finland. Values for forests in natural state (no human impact) were calculated multiplying the values for each age class in Kouki *et al.*[53] by the proportion of land covered by each age class in natural state forests[55]. Values for the 1750 were taken from Kouki *et al.*[53]. Values for the 1960s were taken from the 5th national forest inventory in Finland (NFI5) and current values were obtained from the input data for the simulation (see section Data and forest simulation). The logic to derive favourable reference values (FRV) is explained in the text.

Indicator	Fertility site type	Natural state	Ref. 1750	1960s (NFI5)	FRV	Current values
Deadwood volume (m³/ha)	Herb-rich	140	56	3.3	28	4.9
	Mesic	110	38	3.7	19	4.7
	Sub-xeric	95	29	3.7	14.5	4.4
	Xeric	70	20	3.2	10	4.3
Proportion of old forest (% cover)*	Herb-rich		11	0.6	5.5	0.4
	Mesic	50	25	4.4	12.5	0.1
	Sub-xeric		35	1.4	17.5	0.3
	Xeric		50	0.6	25	0.5
Proportion of deciduous trees (% volume)	Herb-rich			28.5	20	36
	Mesic	na	na	16.9	11.8	24
	Sub-xeric			8.1	5.7	12
	Xeric			3.3	2.3	9

*Threshold ages for old forest: herb-rich 120 years, mesic and sub-xeric 140 years, xeric forests 160 years

We used IUCN criteria to estimate favourable reference values (FRVs). Another option would have been to derive the FRVs from literature on species distributions and population persistence as

functions of critical structural features. Although there is much variation in species response to variation in critical features[56], literature suggests that when resource availability decreases by 80□90% from the natural state, native biota become depauperated due to local extinctions[57–59]. This approach would result in similar FRVs to what we had (Table 1). Our FRVs for deadwood volume also largely correspond to critical ecological threshold values for deadwood-dependent organisms in boreal forests[60,61].

Data and forest simulation

To obtain a representative sample of the initial forest state for forest growth simulations, we used two data sources. We obtained private forest owners' detailed stand-level information from 2016 from the public database www.metsaan.fi of the Finnish Forest Centre. This data was complemented by the Multi-source National Forest Inventory from 2015, to cover areas not privately owned[62]. Both data sets were sampled along the regional and temporal systematic clusters following the design of the 11th Finnish National Forest Inventory (see [23]). We selected only data points within the Middle and Southern boreal regions.

We performed the forest simulations using the open-source forest simulator SIMO[63]. SIMO simulates individual tree growth, mortality, and regeneration for even-aged[64] and uneven-aged boreal forests[65]. We simulated forest dynamics and management under two climate trajectories; representative concentration pathways RCP4.5 and RCP8.5. Climate variables driving stand growth and soil dynamics (mean and amplitude of temperature, CO_2 concentration, precipitation) were based on[66], and the climate data of the Canadian Earth system model CanESM[67]. The impacts of climate on tree growth are introduced into the calculation of volume growth, and further allocated between diameter and height growth based on the models of Matala *et al.*[68].

For each forest plot, we simulated 29 management regimes in five-year periods over 100 years, with the exact number of management alternatives applied depending on the initial conditions of each individual stand (i.e., dominant stand height, basal area, site type, and stand age). The regimes provided a diverse set of management alternatives for each stand, from which the optimization afterward selected an ideal management based on the management effects on optimization objectives (see optimization section below). We grouped the regimes into three classes based on the harvest level: Intensive, Extensive, and Set aside (Table 2).

Table 2. Summary table of the simulated management regimes for Finland and their allocation to the six management classes. The classification to intensive and extensive regimes follows Blattert *et al.*[35]. BAU stands for business-as-usual rotation forestry; CCF for continuous cover forestry.

Management class	Abbreviation	Description
Intensive managements	*BAU*	Rotation forestry, no thinnigs prior clearfelling
	BAU F	= *BAU*, with fertilization
	BAU -5	= *BAU*, with 5 years shorter rotation age
	BAU -5 F	= *BAU -5*, with fertilization
	BAU + 5	= *BAU*, with 5 year extended rotation age
	BAU w thin	Rotation forestry, with thinnings prior clearfelling
	BAU w thin B	= *BAU w thin*, with increased broadleaf planting
	BAU w thin F	= *BAU w thin*, with fertilization
	BAU w thin -5	= *BAU w thin*, with 5 years shorter rotation age
	BAU w thin -5 F	= *BAU w thin -5*, with fertilization
	BAU w thin +5	= *BAU w thin*, with 5-year extended rotation age
	BAU w thin +5 B	= *BAU w thin +5*, with increased broadleaf planting
	BAU w thin GTR	= *BAU w thin*, with 30 retention trees or 30m^2 per ha left
	BAU w thin GTR B	= *BAU w thin GTR*, with increased broadleaf planting
	BAU w GTR	= *BAU*, with 30 retention trees or 30m^2 per ha left
	CCF 1	Thinning from above, basal area threshold -3 m^2/ha
	CCF 2	Thinning from above, basal area threshold +/-0 m^2/ha
Extensive managements	*CCF 3*	Thinning from above, basal area threshold + 3 m^2/ha
	CCF 4	Thinning from above, basal area threshold + 6 m^2/ha
	BAU w/o thin	Rotation forestry, no thinnings prior to or after clearfelling
	BAU w/o thin -20	= *BAU w/o thin*, with 20 years shorter rotation age
	BAU w/o thin -20 F	= *BAU w/o thin -20*, with fertilization
	BAU + 15	= *BAU*, with 15 years extended rotation age
	BAU w thin +15	= *BAU w thin*, with 15 years extended rotation age
	BAU w thin +15 B	= *BAU w thin +15*, with increased broadleaf planting
	BAU +30	= *BAU*, with 30 years extended rotation age
	BAU w thin +30	= *BAU w thin*, with 30 years extended rotation age
	BAU w thin +30 B	= *BAU w thin +30*, with increased broadleaf planting
Set aside (SA)	*SA*	No management, only growth and mortality

Optimization

To quantify the potential ability of the forest to reach the specific FRVs at a specific time period, we applied a systematic optimization framework. This framework is based on the MultiOptForest framework[69] which relies on the application of an achievement scalarizing function (ASF, [70]) and the epsilon constraint method[71]. This allowed evaluating the possibility to reach the specific targets, and the economic timber availability possible once the maximum potential (or targeted) habitat indicators are reached. The general optimization problem can be formulated as:

(1)

$$\min_{x} \max_{i \in I} \left(f_i(x) - z_i^{ref}\right) \Big/ \left(z_i^{ideal} - z_i^{nadir}\right) + \rho \sum_{i \in I} \left(f_i(x) \Big/ \left(z_i^{ideal} - z_i^{nadir}\right)\right)$$

subject to:

(2)
$$f_i(x) \leq \varepsilon_i, \forall i \in I$$

(3)
$$f_{NPV}(x) = \sum_{p \in P} \sum_{j \in J} \sum_{k \in K_j} \frac{\left(x_{jk} z_{jkp}^i\right)}{(1+r)^{(2.5+5(p-1))}} + \sum_{j \in J} \sum_{k \in K_j} \frac{\left(x_{jk} z_{jk}^e\right)}{(1+r)^{(5(\#P))}}$$

(4)
$$f_{EF}(x) \leq \min \left(\sum_{j \in J} \sum_{k \in K_j} \left(x_{jk} z_{jkp}^h\right)\right) \forall p \in P$$

(5)
$$f_{b,s}(x) \leq \sum_{j \in J_s} \sum_{k \in K_j} \left(x_{jk} z_{jkp}^b\right) - t_b \forall p \in P \setminus Y_{\geq target\ year}, s \in S, b \in B$$

(6)
$$\sum_{k \in K_j} x_{jk} = 1, \forall j \in J$$

where I is the set of objectives; $f_i(x)$ is the obtained value for the specific objective; $z_i^{ref}, z_i^{ideal}, z_i^{nadir}$ are the target, ideal and nadir values for the specific objective; ρ is an arbitrary small constant; ε_i is the epsilon constraint target for the specific objective; P is the set of periods under consideration; J is the set of forest plots; K_j is the management options available for stand j; z_{jkp} is the value provided for each stand, management action at the specific period, with superscripts i, e, h and b respectively referring to net income, end stand value, timber harvest quantity and, biodiversity indicators; x_{jk} is the decision to conduct management action k to stand j.

The optimization problem starts with equation (1), where the objective function is a multi-objective optimization, formulated as an ASF. The epsilon constraint is formulated in equation (2), which requires the objective to be greater than the specified value (ε_i). Equation (3) evaluates the net present value of the problem, with equation (4) calculates the minimum timber harvest value across all periods. Equation (5) calculates the biodiversity indicators, striving to meet the target value at a specific period, and exceeding it for all other periods.

To assess when the indicators are possible to be attained, we performed a re-iteration of the optimization striving to achieve the habitat targets at the beginning of each decade. We first ran the optimization without any epsilon constraints (ε_i) and set the targets (z_i^{ref}) for the ASFs to the specific habitat indicator value desired. We then ran the optimization setting the obtained values for the habitat indicators as epsilon constraints (requiring the specific indicator value at the specific year), and re-ran the optimization three times: once were we strived to maximize NPV, to maximize the minimum periodic timber harvested across all time periods exceeding the targeted year, to maximize both the NPV and minimum periodic timber harvested across all time periods exceeding the targeted year.

References

1. Steffen, W. *et al.* Planetary boundaries: Guiding human development on a changing planet. *Science* **347**, 1259855 (2015).

2. Rockström, J. *et al.* A safe operating space for humanity. *Nature* **461**, 472–475 (2009).

3. Richardson, K. *et al.* Earth beyond six of nine planetary boundaries. *Science Advances* **9**, eadh2458 (2023).

4. IPBES. *IPBES Global Assessment on Biodiversity and Ecosystem Services. IPBES Global Assessment on Biodiversity and Ecosystem Services* (2019).

5. Millennium Ecosystem Assessment (Program). *Ecosystems and human well-being: synthesis.* (Island Press, 2005).

6. Erb, K.-H. *et al.* Analyzing the global human appropriation of net primary production — processes, trajectories, implications. An introduction. *Ecological Economics* **69**, 250–259 (2009).

7. IUCN. *An Introduction to the IUCN Red List of Ecosystems: The Categories and Criteria for Assessing Risks to Ecosystems.* (IUCN International Union for Conservation of Nature, 2016). doi:10.2305/IUCN.CH.2016.RLE.2.en.

8. Raworth, K. *Doughnut Economics: Seven Ways to Think Like a 21st-Century Economist.* (Chelsea Green Publishing, 2017).

9. European Commission. Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. EU Biodiversity Strategy for 2030. *COM/2020/380 final* (2020).

10. Convention on Biological Diversity. Kunming–Montreal Global Biodiversity Framework. (2022).

11. Dasgupta, P. *The economics of biodiversity: the Dasgupta review.* (Hm Treasury, 2021).

12. Burton, P. J. *et al.* Sustainability of boreal forests and forestry in a changing environment. *Forests and society: responding to global drivers of change* 247–282 (2010).

13. Vanhanen, H., Jonsson, R., Gerasimov, Y., Krankina, O. N. & Messier, C. Making Boreal Forests Work for People and Nature. 15 (2012).

14. Östlund, L., Zackrisson, O. & Axelsson, a-L. The history and transformation of a Scandinavian boreal forest landscape since the 19th century. *Canadian Journal of Forest Research* **27**, 1198–1206 (1997).

15. Bradshaw, C. J. A., Warkentin, I. G. & Sodhi, N. S. Urgent preservation of boreal carbon stocks and biodiversity. *Trends in Ecology & Evolution* **24**, 541–548 (2009).

16. Kuuluvainen, T., Tahvonen, O. & Aakala, T. Even-aged and uneven-aged forest management in boreal Fennoscandia: a review. *Ambio* **41**, 720–37 (2012).

17. Mönkkönen, M. *et al.* Solving Conflicts among Conservation, Economic, and Social Objectives in Boreal Production Forest Landscapes: Fennoscandian Perspectives. in *Ecosystem Services from Forest Landscapes* 169–219 (Springer, 2018).

18. ArtDatabanken 2015. *Rödlistade arter i Sverige 2015 [Red-listed species in Sweden 2015].* (ArtDatabanken 2015, 2015).

19. S. Henriksen & O. Hilmo. *Norsk Rødliste for Arter 2015 [Norwegian Red List for Species 2015].* (Artsdatabanken, Norge, 2015).

20. *Suomen lajien uhanalaisuus: Punainen kirja 2019 [The 2019 Red List of Finnish Species]*. (Ympäristöministeriö & Suomen ympäristökeskus, 2019).

21. Rockström, J. *et al.* Safe and just Earth system boundaries. *Nature* **619**, 102–111 (2023).

22. Kouki J *et al.* Forests. in *Threatened habitat types in Finland 2018. Red List of habitats. Part I: Results and basis for assessment* 113–124 (Finnish Environment Institute and Ministry of Environment, 2019).

23. Blattert, C. *et al.* Climate targets in European timber-producing countries conflict with goals on forest ecosystem services and biodiversity. *Commun Earth Environ* **4**, 1–12 (2023).

24. Korhonen, K. *et al.* Forests of Finland 2014–2018 and their development 1921–2018. *Silva Fennica* **55**, 10662 (2021).

25. Lee, H. *et al.* Climate change 2023: synthesis report. Contribution of working groups I, II and III to the sixth assessment report of the intergovernmental panel on climate change. (2023).

26. Forest protection. *Natural Resources Institute Finland* https://www.luke.fi/en/statistics/forest-protection.

27. Commercial fellings. *Natural Resources Institute Finland* https://www.luke.fi/en/statistics/commercial-fellings.

28. Konietzko, J., Das, A. & Bocken, N. Towards regenerative business models: A necessary shift? *Sustainable Production and Consumption* **38**, 372–388 (2023).

29. Pascual, U. *et al.* Diverse values of nature for sustainability. *Nature* **620**, 813–823 (2023).

30. Carmenta, R. *et al.* Exploring the relationship between plural values of nature, human well-being, and conservation and development intervention: Why it matters and how to do it? *People and Nature* **5**, 1720–1738 (2023).

31. Augustynczik, A. L. D. *et al.* Diversification of forest management regimes secures tree microhabitats and bird abundance under climate change. *Science of The Total Environment* **650**, 2717–2730 (2019).

32. Duflot, R., Fahrig, L. & Mönkkönen, M. Management diversity begets biodiversity in production forest landscapes. *Biological Conservation* **268**, 109514 (2022).

33. Eyvindson, K. *et al.* High boreal forest multifunctionality requires continuous cover forestry as a dominant management. *Land Use Policy* **100**, 104918 (2021).

34. Himes, A., Betts, M., Messier, C. & Seymour, R. Perspectives: Thirty years of triad forestry, a critical clarification of theory and recommendations for implementation and testing. *Forest Ecology and Management* **510**, 120103 (2022).

35. Blattert, C. *et al.* Enhancing multifunctionality in European boreal forests: The potential role of Triad landscape functional zoning. *Journal of Environmental Management* **348**, 119250 (2023).

36. Mönkkönen, M. *et al.* More wood but less biodiversity in forests in Finland: a historical evaluation. *Memo Soc Fauna Flora Fenn* **98**, 1–11 (2022).

37. Hanski, I. Extinction debt and species credit in boreal forests: modelling the consequences of different approaches to biodiversity conservation. *Annales Zoologici Fennici* **37**, 271–280 (2000).

38. DellaSala, D. A. *et al.* Mature and old-growth forests contribute to large-scale conservation targets in the conterminous United States. *Frontiers in Forests and Global Change* **5**, (2022).

39. Martin, M., Shorohova, E. & Fenton, N. J. Embracing the Complexity and the Richness of Boreal Old-Growth Forests: A Further Step Toward Their Ecosystem Management. in *Boreal Forests in the Face of Climate Change* (eds. Girona, M. M., Morin, H., Gauthier, S. & Bergeron, Y.) vol. 74 191–218 (Springer International Publishing, 2023).

40. Elo, M., Halme, P., Toivanen, T. & Kotiaho, J. S. Species richness of polypores can be increased by supplementing dead wood resource into a boreal forest landscape. *Journal of Applied Ecology* **56**, 1267–1277 (2019).

41. Stokland, J. N., Siitonen, J. & Jonsson, B. G. *Biodiversity in Dead Wood*. (Cambridge University Press, 2012). doi:10.1017/CBO9781139025843.

42. Lassauce, A., Paillet, Y., Jactel, H. & Bouget, C. Deadwood as a surrogate for forest biodiversity: Meta-analysis of correlations between deadwood volume and species richness of saproxylic organisms. *Ecological Indicators* **11**, 1027–1039 (2011).

43. Tikkanen, O.-P., Martikainen, P., Hyvärinen, E., Junninen, K. & Kouki, J. Red-listed boreal forest species of Finland: associations with forest structure, tree species, and decaying wood. *Annales Zoologici Fennici* **43**, 373–383 (2006).

44. Juutilainen, K., Mönkkönen, M., Kotiranta, H. & Halme, P. The effects of forest management on wood-inhabiting fungi occupying dead wood of different diameter fractions. *Forest Ecology and Management* **313**, 283–291 (2014).

45. Turner, M. G. & Seidl, R. Novel Disturbance Regimes and Ecological Responses. *Annual Review of Ecology, Evolution, and Systematics* **54**, 63–83 (2023).

46. Triviño, M. *et al.* Enhancing Resilience of Boreal Forests Through Management Under Global Change: a Review. *Curr Landscape Ecol Rep* **8**, 103–118 (2023).

47. Thorn, S. *et al.* Impacts of salvage logging on biodiversity: A meta-analysis. *Journal of Applied Ecology* **55**, 279–289 (2018).

48. Bouget, C. *et al.* Trait-Based Response of Deadwood and Tree-Related Microhabitats to Decline in Temperate Lowland and Montane Forests. *Ecosystems* (2023) doi:10.1007/s10021-023-00875-9.

49. McDowell, N. G. *et al.* Pervasive shifts in forest dynamics in a changing world. *Science* **368**, eaaz9463 (2020).

50. Turner, M. G. *et al.* The magnitude, direction, and tempo of forest change in Greater Yellowstone in a warmer world with more fire. *Ecological Monographs* **92**, e01485 (2022).

51. Lindenmayer, D. B., Hobbs, R. J., Likens, G. E., Krebs, C. J. & Banks, S. C. Newly discovered landscape traps produce regime shifts in wet forests. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 15887–15891 (2011).

52. *Red List of Habitats – Results and Basis for Assessment*. (Finnish Environment Institute and Ministry of the Environment).

53. Kouki J *et al.* Metsät [Forests]. in *Suomen luontotyyppien uhanalaisuus 2018. Luontotyyppien punainen kirja – Osa 1: Tulokset ja arvioinnin perusteet [Threatened habitats in Finland 2018 - Red Book of Habitats - Part 1: Results and assessment criteria]* vol. 2018 171–201 (Finnish Environment Institute and Ministry of Environment, 2018).

54. Keith, D. A. *et al.* Scientific Foundations for an IUCN Red List of Ecosystems. *PLoS ONE* **8**, e62111 (2013).

55. Berglund, H. & Kuuluvainen, T. Representative boreal forest habitats in northern Europe, and a revised model for ecosystem management and biodiversity conservation. *Ambio* 1–15 (2021) doi:10.1007/s13280-020-01444-3.

56. Mönkkönen, M. & Reunanen, P. On Critical Thresholds in Landscape Connectivity: A Management Perspective. *Oikos* **84**, 302–305 (1999).

57. Hanski, I. Habitat loss, the dynamics of biodiversity, and a perspective on conservation. *Ambio* **40**, 248–55 (2011).

58. Hanski, I. Habitat fragmentation and species richness. *Journal of Biogeography* **42**, 989–993 (2015).

59. Rybicki, J. & Hanski, I. Species–area relationships and extinctions caused by habitat loss and fragmentation. *Ecology Letters* **16**, 27–38 (2013).

60. Müller, J. & Bütler, R. A review of habitat thresholds for dead wood: a baseline for management recommendations in European forests. *European Journal of Forest Research* **129**, 981–992 (2010).

61. Junninen, K. & Komonen, A. Conservation ecology of boreal polypores: A review. *Biological Conservation* **144**, 11–20 (2011).

62. Mäkisara, K., Katila, M. & Peräsaari, J. *The Multi-Source National Forest Inventory of Finland – methods and results 2015*. (Luonnonvarakeskus, 2019).

63. Rasinmäki, J., Mäkinen, A. & Kalliovirta, J. SIMO: An adaptable simulation framework for multiscale forest resource data. *Computers and Electronics in Agriculture* **66**, 76–84 (2009).

64. Hynynen, J. *et al. Models for predicting stand development in MELA System.* (Metsäntutkimuslaitos, 2002).

65. Pukkala, T., Lähde, E. & Laiho, O. Species interactions in the dynamics of even- and uneven-aged boreal forests. *Journal of Sustainable Forestry* **32**, 371–403 (2013).

66. Lehtonen, I., Venäläinen, A., Kämäräinen, M., Peltola, H. & Gregow, H. Risk of large-scale fires in boreal forests of Finland under changing climate. *Nat. Hazards Earth Syst. Sci.* **16**, 239–253 (2016).

67. Von Salzen, K. *et al.* The Canadian Fourth Generation Atmospheric Global Climate Model (CanAM4). Part I: Representation of Physical Processes. *Atmosphere-Ocean* **51**, 104–125 (2013).

68. Matala, J., Ojansuu, R., Peltola, H., Raitio, H. & Kellomäki, S. Modelling the response of tree growth to temperature and CO_2 elevation as related to the fertility and current temperature sum of a site. *Ecological Modelling* **199**, 39–52 (2006).

69. Eyvindson, K. *et al.* MultiOptForest: An interactive multi-objective optimization tool for forest planning and scenario analysis. *Open Research Europe* **3**, 103 (2023).

70. Wierzbicki, A. P. On the completeness and constructiveness of parametric characterizations to vector optimization problems. *OR Spektrum* **8**, 73–87 (1986).

71. Haimes, Y. Y., Lasdon, L. S. & Wismer, D. A. On a Bicriterion Formulation of the Problems of Integrated System Identification and System Optimization. *IEEE Trans. Syst., Man, Cybern.* **1**, 296–297 (1971).

Acknowledgements

This work was partly funded by the Strategic Research Council at the Academy of Finland (Mönkkönen 358473). We would also like to thank Kaisa Junninen for the help with threat assessments of forest habitats.

Author contributions

M.M., C.B., R.F., M.E., K.E., M.T. and D.B. conceived and designed the study, performed the experiments, and analysed the data. C.B., K.E. and J.K. contributed materials/analysis tools. All authors contributed to the writing of the manuscript.