1 **Reward integration in prefrontal-cortical and ventral-hippocampal nucleus accumbens**
2 **inputs cooperatively modulates engagement**
3

4 *Eshaan S. Iyer[1], Peter Vitaro[2], Serena Wu[1], Jessie Muir[4,5], Yiu Chung Tse[2], Vedrana*
5 *Cvetkovska[2], Rosemary C. Bagot[2,3]*
6
1. Integrated Program in Neuroscience, McGill University, Montréal, Canada
2. Department of Psychology, McGill University, Montréal, QC, Canada
3. Ludmer Centre for Neuroinformatics and Mental Health, Montréal, QC, Canada
4. Center for Neuroscience, University of California, Davis; Davis, 95618, USA.
5. Department of Neurology, School of Medicine, University of California, Davis; Sacramento,
95817, USA.

Abstract
15

The NAc, a highly integrative brain region controlling motivated behavior, is thought to receive
distinct information from various glutamatergic inputs yet strong evidence of functional
specialization of inputs is lacking. While circuit neuroscience commonly seeks specific functions
for specific circuits, redundancy can be highly adaptive and is a critical motif in circuit
organization. Using dual-site fiber photometry in an operant reward task, we simultaneously
recorded from two NAc glutamatergic afferents to assess circuit specialization. We identify a
common neural motif that integrates reward history in medial prefrontal cortex (mPFC) and
ventral hippocampus (vHip) inputs to NAc. Then, by systematically degrading task complexity,
dissociating reward from choice and action, we identify key circuit-specificity in the behavioral
conditions that recruit encoding. While mPFC-NAc invariantly encodes reward, vHip-NAc
encoding is uniquely anchored to loss. Ultimately, using optogenetic stimulation we demonstrate
that both inputs co-operatively modulate task engagement. We illustrate how similar encoding,
with differential gating by behavioral state, supports state-sensitive tuning of reward-motivated
behavior.

INTRODUCTION
Redundancy is a defining property of nervous system organization (Marder, 2011; Marder &
Goaillard, 2006; Mizusaki & O'Donnell, 2021) yet there has been limited consideration of the
role of redundancy in neural circuit mechanisms of motivated behavior. Redundancy in neural
circuits may confer various advantages, including increasing robustness of cognition and
behavior to perturbation, enhancing encoding accuracy, and facilitating coherent integration of
multiple inputs, suggesting it should be a frequently observed motif (Ghanbari et al., 2023;
Hiratani & Fukai, 2018; Nguyen et al., 2019). While the literature abounds with examples of
apparent circuit-specific cognitive and behavioral functions, the potential for redundancy is
rarely examined. To better understand these opposing motifs in nucleus accumbens (NAc)
circuits we leveraged dual circuit recordings and computational modeling to rigorously test the
specificity and redundancy of information processing in a fully controlled, within animal
comparison of two NAc glutamatergic inputs.
44

The NAc integrates glutamatergic inputs with dopaminergic input from the ventral tegmental
area, with multiple glutamatergic inputs converging at the level of individual medium spiny
neurons in the NAc medial shell (Britt et al., 2012; Carter et al., 2007; Christoffel et al., 2021;
Floresco, 2015; French & Totterdell, 2002; Lind et al., 2023; Muir et al., 2024; O'Donnell &
Grace, 1995). Prominent theoretical perspectives hold that these inputs send qualitatively
distinct information which the NAc then integrates to orchestrate motivated behavior (Floresco,
2015; Grace et al., 2007; Lind et al., 2023; Parker et al., 2022). For example, the mPFC
contributes information about rewarding events and executive control while the vHip contributes
emotional context and behavioral inhibition (Bagot et al., 2015; Barker et al., 2019; Hamel et al.,
2022; Lindenbach et al., 2022; Muir et al., 2020; Otis et al., 2017; Parker et al., 2022; Spellman
et al., 2021; Wenzel et al., 2023; Yoshida et al., 2020). Despite predictions of distinct encoding
and behavioral function for mPFC-NAc and vHip-NAc, strong evidence of functional
specialization is lacking. To date, most studies have examined a single input and the few
studies that examined one or more inputs in the same task compared across animals leaving
open the possibility that inter-individual variation in behavior and other variables influence neural
encoding (Britt et al., 2012; Chen et al., 2023; Reed et al., 2018).
61
To systematically interrogate functional redundancy versus specialization we simultaneously
probed neural encoding using dual-site *in vivo* fiber photometry to record activity in two
glutamatergic circuits during reward-guided choice in a two-armed bandit task. The mPFC-NAc
is widely appreciated to mediate reward processing and, given that vHip-NAc inputs converge
with mPFC-NAc, we asked if the vHip-NAc might also contribute to this function (Otis et al.,
2017; Parker et al., 2022; Spellman et al., 2021). Using trial-by-trial modeling of neural activity,
we identify a novel mechanism for integrating outcome information across trials that is common
to both circuits. Analyzing the redundancy across signals revealed an additional dimension of
uniqueness to vHip-NAc encoding. By sequentially degrading task complexity we show that,
despite sharing a common mechanism for outcome integration, each circuit is recruited in
distinct behavioral states, with the vHip-NAc preferentially encoding reward after loss.
Optogenetically manipulating circuit-specific activity revealed that, once recruited, both inputs
cumulatively mediate dynamic behavioral engagement. Our findings reveal co-operative circuit
organization in NAc wherein redundant encoding in two inputs is gated by circuit-specific
mechanisms for state-sensitive tuning of reward-motivated behavior.
77
78 **Results**
79
80 *mPFC-NAc and vHip-NAc similarly encode outcomes in a probabilistically rewarded*
81 *environment*
82
To assess redundancy versus specificity in outcome encoding in two distinct circuits under
matched conditions and trial histories, we injected retrograding AAV-GCaMP7f in NAc medial
shell and implanted optic fibers in mPFC and vHip to record Ca^{2+}-associated fluorescence while
mice engaged in reward-guided choice (Fig. 1E). We trained mice in a two-lever probabilistic
reward learning task (i.e. a two-armed bandit task) in which lever pressing probabilistically earns
a chocolate milk reward (Fig. 1A). Following each lever press, one of two different auditory cues

signaled trial outcome (rewarded, unrewarded) and start of the inter-trial interval (ITI). To
maintain a dynamic environment with robustly encountered rewarded and unrewarded
outcomes, levers were probabilistically rewarded on 80% or 20% of presses with probabilities
switched after five consecutive responses on the high probability lever. Female (n=10) and male
(n=12) mice experienced similarly high numbers of unrewarded and rewarded trials and low
numbers of omission trials (Fig. 1B-D). Examining behavior across sessions shows decreasing
staying probability after unrewarded outcomes and increasing rewards earned, indicating
animals use information about outcomes to guide behavior (Fig. S1).

Trial based tasks are ideal for probing neural encoding, generating large numbers of trials.
However, standard analysis approaches either analyze individual trials, failing to account for the
within animal nested data structure and inappropriately inflating effects, or average all trials
within animals, thereby underestimating effects. Choice tasks are additionally challenging with
the number of instances of each trial type varying across animals. To preserve the power of
trial-by-trial data while accounting for the nested structure and unbalanced observations we
used a linear mixed model approach (Yu et al., 2022). To examine how outcome is encoded in
each projection, we modeled normalized Ca^{2+}-associated fluorescence change as a function of
trial outcome, while controlling for inter-individual variability.

Reward strongly suppressed mPFC-NAc and vHip-NAc activity in female and male mice. In
mPFC-NAc, a peak following the lever press and outcome delivery is followed by gradually
emerging reward-associated suppression across the ITI (Fig 1H,J,P,Q). In vHip-NAc, an initial
peak is followed by suppression after the lever press and outcome delivery, with suppression
sustained following reward or activity gradually increasing following unrewarded outcomes (Fig.
1 L,N,R,S). We focused analysis on the end of the ITI (8-10 sec after lever press) when trial
outcome has been integrated prior to next trial start. By ITI end, reward robustly suppressed
mPFC-NAc activity in female and male mice (Fig. 1I,K). Reward also robustly suppressed vHip-
NAc activity in female and male mice (Fig. 1M,O). This indicates that outcome encoding
emerges across the ITI with reward suppressing mPFC-NAc and vHip-NAc activity. To explore
modulation by other task factors, we examined neural encoding time-locked to licking and
decision-relevant behaviors. We did not observe clear neural encoding of licking (Fig. S2A,D),
the identity of the chosen lever (Fig. S2B,E), or the decision to stay or shift (Fig. S2C,F)
suggesting that outcome is the primary source of modulation in mPFC-NAc and vHip-NAc in this
task.

Having observed that mPFC-NAc and vHip-NAc are similarly modulated by reward, we then
examined if one circuit leads the other. Calculating the time lag for the maximum cross-
correlation, we found that the regression coefficients of the mPFC-NAc and vHip-NAc cross-
correlation did not significantly differ from zero in rewarded or unrewarded trials in either sex.
This shows that neither circuit drives outcome encoding in the other (Fig. S3A, B). However, we
observe that suppression emerges earlier in vHip-NAc than mPFC-NAc, suggesting that while
overall encoding might be similar, the underlying dynamics likely vary between the pathways
(Fig. S3C).

mPFC-NAc and vHip-NAc similarly integrate reward history

We find that mPFC-NAc and vHip-NAc similarly encode outcomes. Observing that reward-
induced suppression lasted throughout the ITI, we speculated that this enduring modulation
might integrate reward information across successive trials and that this integration might be
more prominent in mPFC-NAc than vHip-NAc, given prior evidence of enduring representation
in mPFC (Parker et al., 2022; Spellman et al., 2021; Sul et al., 2010). To test this, we sorted
trials by both prior and current outcome, identifying trial sequences that were rewarded then
rewarded (R→R), rewarded then unrewarded (R→U), unrewarded then rewarded (U→R), and
unrewarded then unrewarded (U→U). We then compared neural activity across the ITI on the
most recent trial to determine how prior outcome modulates outcome encoding on the current
trial. Analyzing males and females separately revealed similar modulation (Fig. S4) and we
therefore report sex-combined analyses. Both previous and current outcome modulate mPFC-
NAc activity (Fig. 2A). Following a given trial (t -1), reward suppresses mPFC-NAc activity (Fig.
2B) effectively resetting the baseline for the next trial. Reward on the subsequent trial (t0)
similarly suppresses mPFC-NAc by ITI end, regardless of prior outcome. However, when mice
are unrewarded on the subsequent trial (t0), suppression of mPFC-NAc by prior reward is
maintained through ITI end (Fig. 2C). This suggests that a single reward maximally and
enduringly suppresses mPFC-NAc activity and that, in the absence of subsequent reward, this
suppression slowly dissipates.

We then examined if vHip-NAc similarly integrates outcomes (Fig. 2D). Following a given trial (t-
1), reward suppresses vHip-NAc activity (Fig. 2E). As with mPFC-NAc, this resets the baseline
for the next trial (t0), wherein reward suppresses vHip-NAc regardless of prior outcome.
However, when the subsequent trial (t0) is unrewarded, suppression of vHip-NAc activity by
prior reward is maintained through ITI end (Fig. 2F). Together, this shows that mPFC-NAc and
vHip-NAc similarly integrate outcomes across trials. In both circuits, reward maximally
suppresses neural activity and activity gradually increases following subsequent unrewarded
outcomes, such that, by ITI end, the relative degree of suppression represents an integrated
reward outcome history.

mPFC-NAc and vHip-NAc are differentially sensitive to loss

Analyzing neural encoding of reward and outcome integration revealed that mPFC-NAc and
vHip-NAc similarly encode reward suggesting they may provide redundant information to the
NAc. To test redundancy between mPFC-NAc and vHip-NAc we calculated the conditional
entropy of mPFC-NAc given vHip-NAc (□(mPFC-NAc|vHip-NAc)) and vHip-NAc given mPFC-
NAc (□(vHip-NAc|mPFC-NAc)). In this way, we assessed the information contributed by each
circuit beyond that contributed by the other at ITI end, when outcome is fully integrated (Fig.
3A). We contrasted entropy between rewarded and unrewarded outcomes as a function of prior
outcome. Relative to unrewarded outcomes, the entropy of mPFC-NAc given vHip-NAc was
reduced by rewarded outcomes, indicating that vHip-NAc and mPFC-NAc signals are more
redundant after reward than non-reward (Fig. 3B). In contrast, following previous unreward, but
not previous reward, current reward increased the entropy of vHip-NAc given mPFC-NAc (Fig.

3C), indicating that, under these conditions, mPFC-NAc explains less of the vHip-NAc signal.
This shows that, after reward, vHip-NAc and mPFC-NAc encoding converges, becoming more
redundant, but when reward is made more surprising by immediately following loss, vHip-NAc
carries additional information. That is, despite global redundancy in reward encoding motifs, we
identify a dimension of circuit specificity and a potential unique role for vHip-NAc in encoding
reward following loss.
183
If this is true, across outcome histories, vHip-NAc encoding should be most apparent when
reward follows an unrewarded outcome, whereas, following consecutive rewards, vHip-NAc
should become insensitive to outcome as rewards become less surprising. In contrast, mPFC-
NAc encoding is predicted to be relatively invariant across outcome histories. To test this, we
examined current outcome encoding at ITI end while considering prior outcomes up to three
trials back. Consistent with our prediction, mPFC-NAc encoded current outcome regardless of
prior outcome history (Fig. 3D; Supplementary Table 1) while vHip-NAc failed to encode current
outcome after two or more consecutive rewards (Fig. 3E; Supplementary Table 1). This
supports our hypothesis that vHip-NAc encoding anchors to loss such that it is specifically
engaged by surprising rewards whereas mPFC-NAc invariantly encodes outcome.
194
195   *Degrading task requirements reveals circuit-specific roles in reward integration*
196
Analyzing informational redundancy and encoding across varying outcome histories suggested
that, while mPFC-NAc and vHip-NAc encode and integrate reward via a common mechanism,
each may nevertheless serve distinct functions in reward processing. To isolate the specific
conditions under which each circuit integrates outcomes we recorded neural activity while
degrading task requirements to sequentially eliminate choice and action. We first eliminated
choice, extending only a single lever while maintaining the requirement to press to elicit an
outcome. To hold outcome experience constant, the specific sequence of reward and unreward
was yoked to each animal's prior performance on the two-lever task (Fig. 4A). In the absence of
choice, mPFC-NAc continued to encode previous and current outcome (Fig. 4B). On trial t0, by
ITI end, current and prior outcomes were encoded, as in the two-lever task (Fig. 4C, Fig. 2C).
Examining vHip-NAc in the one-lever task also revealed largely similar outcome-mediated
modulation (Fig. 4D, Fig. 2D). At ITI end, prior and current outcomes were integrated, similar to
the two-lever task (Fig. 4E, Fig. 2F). Despite conserved information encoding in both circuits,
the shape of the vHip-NAc signal was more visibly altered than the mPFC-NAc. In particular, the
vHip-NAc signal in the one-lever task appeared noisier and blunted with the expected peak
following lever press largely absent, potentially suggesting heightened sensitivity to task
structure. Overall, we find that both mPFC-NAc and vHip-NAc maintain similar graded
representations of reward history that are largely independent of choice requirements.
215
Removing lever choice minimally impacted reward integration. We then asked if neural
integration of outcome history is entirely independent of response requirements by removing
both levers in a choice-free response-free task. Trials continued to be signaled by cue-lights, but
without lever extension and outcomes were passively delivered yoked to each animal's
individual performance on the full two-lever task (Fig. 4F). Eliminating the response requirement

markedly and distinctly altered reward integration in both circuits. In mPFC-NAc (Fig. 4G),
encoding of prior outcome was erased and only the current outcome encoded (Fig. 4H). This
differs from both the two-lever and one-lever tasks wherein mPFC-NAc encoded a graded
representation of reward history and suggests that mPFC-NAc integrates reward history only in
instrumental settings where a response elicits outcomes. However, even when rewards are
passively encountered (i.e. when no lever press is required), mPFC-NAc continues to encode
reward but with a shortened time constant, such that only the most recent outcome is retained.
228
vHip-NAc representation of reward history was also degraded, yet in a distinct manner (Fig. 4I).
Current outcomes were encoded only when the previous trial, t-1, was unrewarded, suggesting
a dependence on loss (Fig. 4J). This shift in encoding translates into vHip-NAc effectively
overlooking isolated instances of non-reward, likely reflecting an extended time constant.
Critically, this cannot be explained by changes in task engagement given that mPFC-NAc
continued to represent reward in these same animals (Fig. 4G,H) and licking bouts were
similarly maintained across task variants (Fig. S5). Rather, this reveals that task demands
differently shape neural encoding of reward in mPFC-NAc and vHip-NAc. When reward is
passively encountered, independent of a required response, mPFC-NAc maintains a simplified
reward representation across a shortened temporal window, limiting integration across trials. In
contrast, vHip-NAc anchors encoding to loss with an extended time constant, to preferentially
represent surprising rewards. This suggests that while the fundamental function of mPFC-NAc
in rewarding contexts is to encode outcomes, the fundamental function of vHip-NAc is to use
loss to tune outcome encoding.
243
244 *mPFC-NAc and vHip-NAc modulate task engagement*
245
Examining neural representation of outcomes identified both mechanistic redundancy and
functional specificity in mPFC-NAc and vHip-NAc encoding. We then asked how this neural
processing might integrate to modulate behavior. While in general encoding was similar in both
circuits, reducing the requirement for engagement by making reward non-contingent revealed
functional specialization. We hypothesized that outcome-associated neural activity in mPFC-
NAc and vHip-NAc modulates task engagement. To test this, we examined if neural activity at
ITI end predicted latency to lever press on the subsequent trial, a metric operationalizing
engagement (Bari et al., 2019; Beierholm et al., 2013; Cox et al., 2023; Hamid et al., 2016; Niv
et al., 2007). A linear mixed effects model revealed modest yet significant relationships between
latency to lever press and mPFC-NAc, vHip-NAc, and the interaction of mPFC-NAc and vHip-
NAc activity (Fig. 5A, Supplementary Table 2; Fig. S6). This suggests that increased activity
during outcome integration in either circuit increases latency to lever press, indicating reduced
behavioral engagement.
259
From the association between neural activity and latency, we hypothesized that reward
suppresses activity in mPFC-NAc and vHip-NAc to support behavioral engagement, defining a
mechanism whereby recent reward history modulates engagement in reward-motivated
behavior. We predicted that acutely increasing activity in either mPFC-NAc or vHip-NAc would
suppress engagement. To test this, we injected retrograding AAV-ChR2 into NAc and implanted

fibers above mPFC and vHip to deliver blue light stimulation during the ITI on a subset of trials
in the two-armed bandit task (Fig. 5B-E; Fig. S7). To test if mPFC-NAc and vHip-NAc uniquely
or redundantly control behavior we stimulated each circuit alone or both simultaneously.
Stimulating either circuit alone had no effect, whereas stimulating both simultaneously increased
latency to lever press, but did not alter choice behavior (Fig. 5F; Fig. S8A). This could indicate
either a threshold for sufficient cumulative glutamatergic drive or a requirement for synergistic
interaction between inputs. To differentiate these possibilities, we repeated the experiment with
stronger stimulation. Strong stimulation of either circuit alone increased latency to lever press,
again with no effect on choice (Fig. 5G, Fig. S8B). This shows that total glutamatergic input
modulates engagement, independent of input identity. mPFC-NAc stimulation yielded a slightly
weaker effect than vHip-NAc, consistent with previous findings that mPFC projections to NAc
medial shell are sparser than those from vHip (Britt et al., 2012). Stimulation during lever
presentation did not yield any changes in latency or choice behavior, supporting the importance
of neural integration of outcome during the ITI period, prior to action initiation (Fig. S9). Together
our results demonstrate that mPFC-NAc and vHip-NAc dynamically track outcome information
to modulate behavioral engagement according to recent history of reward. While each circuit is
specialized to execute this function under distinct behavioral states, once engaged, they
redundantly modulate behavior pointing to complementary roles in control of reward seeking.
283
284   **DISCUSSION**
285
We examined redundancy and specificity in the function of two distinct glutamatergic inputs to
the NAc. Using dual-site fiber photometry to probe trial-by-trial outcome encoding
simultaneously in two circuits in the same animal during reward-guided choice, we find that
mPFC-NAc and vHip-NAc similarly integrate reward via suppression of neural activity. By then
systematically manipulating the conditions in which outcomes are encountered, we revealed
that each circuit executes this common function under distinct behavioral states. While the
mPFC-NAc invariantly encodes outcome, vHip-NAc uses information about loss to tune
outcome encoding, effectively amplifying surprising reward. By comparing independent or
synchronous circuit-specific optogenetic stimulation we show that, once engaged, these circuits
cooperatively execute a shared function, i.e. modulating task engagement. Taken together, we
identify a redundant mechanism for outcome integration with circuit-specific gating. This
supports convergence of multiple inputs in tuning behavioral engagement to recent history of
reward.
299
Our finding that both mPFC-NAc and vHip-NAc integrate information about outcomes of reward-
motivated actions is consistent with the well-established role of mPFC in reward processing.
Critically, we demonstrate that this function is not specific or limited to the mPFC-NAc.
Globally, the mPFC encodes information about previous actions and outcomes (Sul et al., 2010)
and mPFC projections to the NAc bridge information about current actions and outcomes
across trials (Parker et al., 2022; Spellman et al., 2021). Our findings suggest these functions
are not unique to mPFC-NAc and are shared by vHip-NAc. However, we identify novel state
dependent specialization in how reward integration is engaged in each circuit. We show that the
mPFC-NAc fundamentally functions as a reward ledger, with reward suppressing neural activity

no matter the behavioral state. In contrast, we find that vHip-NAc is tuned to preferentially
encode outcome information after loss.
311
Differential encoding between mPFC-NAc and vHip-NAc emerged upon degrading task
requirements, a manipulation that minimizes cognitive and behavioral demands effectively
reducing the behavioral utility of representing integrated reward history. Under these
circumstances, the base functionality of each circuit is revealed: mPFC-NAc encoding is
anchored to reward whereas vHip-NAc is anchored to loss. Layered on top of this base
functionality, representation of reward history scales with task complexity in support of
behavioral demands. When reward is passively encountered with limited utility for action-
outcome associations, mPFC-NAc encoding is limited to the most recent outcome. In more
complex environments wherein actions elicit reward and action-outcome associations have high
utility, the mPFC-NAc encoding window extends to integrate reward history. In simpler task
structures that no longer require active engagement with a lever to earn rewards, the time-
constant of vHip-NAc encoding shifts such that an activity no longer increases when a single
unrewarded outcome follows reward. As a result, the vHip-NAc effectively comes to encode
consecutive loss against other outcomes. Together, this suggests a role for vHip-NAc in
providing information about the state of reward statistics in the environment, modulating
behavior as a function of loss, and revealing a novel role for this circuit as a parallel and distinct
stream of outcome integration.
329
The NAc has long been implicated in reward processing yet the precise neural circuit
mechanisms are still being resolved. In the NAc medial shell, reward predominantly suppresses
neural activity (Chen et al., 2023). This suppression likely maintains reward seeking as
stimulation of either D1 or D2 medium spiny neurons bidirectionally controls reward seeking
behavior (Lafferty et al., 2020). Here we show that reward suppresses both mPFC-NAc and
vHip-NAc, two major excitatory inputs to NAc medial shell. Reward-associated suppression of
these inputs would lead to reduced NAc activity. As such, our findings are consistent with
reports that optogenetic stimulation of diverse glutamatergic inputs inhibits motivated behavior
and the idea that glutamatergic input to NAc medial shell functions as a brake on motivated
behavior (Lafferty et al., 2020; Millan et al., 2017; Reed et al., 2018; Yoshida et al., 2019, 2021).
We show that outcome integration in mPFC-NAc and vHip-NAc initiates parallel, temporally
integrated, neural signaling that may engage this 'brake' to align ongoing behavior with recent
reward history and so tune behavioral engagement to prevailing environmental conditions.
343
Employing a redundant mechanism in mPFC-NAc and vHip-NAc may serve several functions. A
common mechanism makes for simple integration of multiple inputs and ensures the robustness
of the fundamental function of reward-guided engagement against insults. Further, modulating
redundant encoding with state-dependent circuit-specific sensitivity may increase the granularity
and range of encoding to ultimately amplify the behavioral impact of surprising rewards. We
demonstrate that high levels of reward suppress activity in both mPFC-NAc and vHip-NAc to
favor continued engagement. In contrast, strong activation of either input suppresses
engagement, but, when weakly activated, synchronous recruitment of both circuits is required.
Functionally, this may translate into a mechanism whereby moderate, balanced activity

predominantly modulates task engagement while allowing for strong activation of either circuit to
exert more direct behavioral control.
355
Preferential outcome encoding in vHip-NAc after loss may serve to strengthen engagement in
variably rewarding environments, driving increased engagement when reward is infrequently
encountered. The sensitivity of vHip-NAc to continuous loss, may also serve to gauge reward
statistics of the environment, continually increasing with each consecutive loss to trigger task
disengagement when activity reaches some threshold. Qualitatively, we see suggestion of this
in the shape of the signal after loss: mPFC-NAc tends to plateau while vHip-NAc continues to
increase. Ultimately, dysregulated outcome-encoding in either mPFC-NAc or vHip-NAc could
alter behavioral sensitivity to reward and loss. Relative to mPFC-NAc, The vHip-NAc is poised
to exert an outsized effect on behavioral engagement both in the strength of its input to NAc
medial shell (Britt et al., 2012) and in its role in signaling loss. For example, hyperactivity of
vHip-NAc may erroneously signal sustained loss, causing premature disengagement. Given our
finding that engagement is modulated by the cumulative glutamatergic input to NAc, a
sufficiently strong vHip-NAc signal could effectively jam any reward signal from mPFC-NAc,
compounding insensitivity to reward that manifests as anhedonia. Indeed, disruption of the
balance between NAc inputs and increased vHip-NAc drive is observed following chronic stress
(Bagot et al., 2015; Muir et al., 2020; Pignatelli et al., 2021; Williams et al., 2020), as well as
chronic alcohol (Griffin et al., 2023; Kircher et al., 2019) and cocaine intake (Barrientos et al.,
2018; Cahill et al., 2016; Pascoli et al., 2014; Zinsmaier et al., 2022), manipulations associated
with aberrant reward processing.
375
Here we examined the simultaneous encoding in two key neural circuits for motivated behavior.
By considering outcome encoding within the context of recent outcome history and behavioral
demands we identified a common neural mechanism of sustained temporal integration of
reward outcomes and reveal how the external environment differentially shapes internal
representations within two neural circuits. We also revealed critical circuit specificity: while
mPFC-NAc consistently tracks outcomes, vHip-NAc preferentially encodes outcome information
after loss. By illustrating the interplay of redundancy and specificity in circuit control of
motivated behavior we demonstrate the need to contextualize events within varied behavioral
states to fully understand neural encoding. Overall, our findings point to the importance of
balanced suppression of NAc glutamatergic inputs during outcome integration to maintain
reward-modulated behavioral engagement.
387
388
METHODS
Animals

Mice were maintained on a 12-h light-dark cycle (lights on at 7:00AM) at 22-25ºC, group-housed
with 3-4 same-sex cage-mates with *ad libitum* access to food and water. All experimental
manipulations occurred during the light cycle, in accordance with guidelines of McGill
University's Comparative Medicine and Animal Resources Center and approved by the McGill
Animal Care Committee. 7-week-old male and female C57BL/6J mice were obtained from

Jackson Laboratories and habituated to the colony room one week prior to start of
manipulations. Mice were food restricted to 85% of their free-feeding body weight during
experimentation.

Surgeries

Stereotaxic surgery was performed under ketamine (100 mg/kg)/xylazine (10 mg/kg)
anesthesia. To achieve projection-specific GCaMP7f expression in glutamatergic NAc-projecting
cells, 0.3µl pGP-AAVrg-syn-jGCaMP7f-WPRE virus (1.85× 10^{13}GC/ml; Addgene) was infused
into the NAc (A/P: +1.3, M/L: +/-0.60, D/V: -4.9) at a rate of 0.1µl per min, before raising the
needle to D/V: -4.7 and infusing a further 0.4µl virus, and allowed to diffuse for 10 min before
withdrawing the needle. pGP-AAV-syn-jGCaMP7f-WPRE was a gift from Douglas Kim & GENIE
Project (Addgene plasmid # 104488 ; http://n2t.net/addgene:104488 ; RRID:Addgene_104488)
(Dana et al., 2019). Chronically implantable optic fibers (Neurophotometrics) with 200µm core
and 0.37 NA threaded through ceramic ferrules were implanted above the ventral subiculum of
the vHip (A/P: -3.40, M/L: +/-3.00, D/V: -4.75) and infralimbic mPFC (A/P: 1.90, M/L: +/-0.3, D/V:
-2.80). Recordings began minimum 4 weeks after surgery to allow sufficient time for stable and
robust retrograde virus expression. To achieve projection-specific ChR2 expression in
glutamatergic NAc-projecting cells, 0.3µl pGP-AAVrg-hSyn-hChR2(H134R)-EYFP virus (7×
10^{12}GC/ml; Addgene) or a fluorophore only control, pGP-AAVrg-hSyn-mCherry (7× 10^{12}GC/ml;
Addgene) was infused into the NAc (A/P: +1.3, M/L: +/-0.60, D/V: -4.9) at a rate of 0.1µl per
min, before raising the needle to D/V: -4.7 and infusing a further 0.4µl virus, and allowed to
diffuse for 10 min before withdrawing the needle. pAAV-hSyn-hChR2(H134R)-EYFP was a gift
from Karl Deisseroth (Addgene plasmid # 26973; http://n2t.net/addgene:26973 ;
RRID:Addgene_26973). pAAV-hSyn-mCherry was a gift from Karl Deisseroth (Addgene plasmid
114472; http://n2t.net/addgene:114472 ; RRID:Addgene_114472). Chronically implantable
optic fibers (Neurophotometrics) with 200µm core and 0.22 NA threaded through ceramic
ferrules were implanted above the ventral subiculum of the vHip (A/P: -3.40, M/L: +/-3.00, D/V: -
4.75) and infralimbic mPFC (A/P: 1.90, M/L: +/-0.3, D/V: -2.80). Optogenetic manipulations
began minimum 4 weeks after surgery to allow sufficient time for stable and robust retrograde
virus expression.

Histology

After completion of all behavioral testing, mice were deeply anesthetized with ketamine/xylazine
and transcardially perfused with phosphate buffered saline (PBS) and paraformaldehyde (4%).
Brains were removed and post-fixed in paraformaldeyhde for 24h and stored in PBS until
sectioning on a vibratome (50 µm). Sections were mounted with Vectashield with DAPI (Vector
Laboratories) and examined under a fluorescent microscope (Leica DM6000 B) to confirm viral
expression and fiber placement. A confocal microscope (Zeiss LSM800) was used to obtain
fluorescent images. Images were acquired as tiles with a 20x air objective (NA 0.8) using Zeiss
Zen Blue imaging software. Images were collected in the McGill University Advanced
BioImaging Facility (ABIF), *RRID:SCR_017697*. Mistargeted animals were excluded from
analysis.

Apparatus

Behavioral experiments were performed in standard Med Associates operant boxes (15.24 x
13.34 x 12.7 cm) enclosed in sound attenuating chambers outfitted with a programmable audio
generator, two retractable levers and cue lights either side of a food port for delivering a liquid
chocolate milk reward (30µl, Nesquick) diluted with water in a 2:1 ratio. Boxes were controlled
and data collected by a computer running MED-PC software (Med-Associates).

Lever Press Training

Training was completed in three stages, with all training sessions lasting 30 minutes. In the first
stage, animals were presented with two levers, both of which delivered a chocolate milk reward
with a 100% probability. To signal the start of the trial, both levers extended and the cue lights
above the levers turned on, animals then had 60 seconds to make a response on either lever. A
press on either lever resulted in lever retraction, immediate delivery of a 30 µL chocolate milk
reward, and the start of a 3 second auditory cue (2kHz pure tone or white noise). Following
either a lever press or 60 seconds with no press (i.e. an omission), a 10 second intertrial interval
(ITI) was triggered. After one session with over 25 responses, animals progressed to the second
stage. In this stage animals again were presented with two levers but reward was now delivered
with a 50% probability on both levers. To signal the start of the trial, both levers extended and
the cue lights above the levers turned on, animals then had 60 seconds to make a response. A
lever press resulted in lever retraction and immediate delivery of the outcome, either a 30 µL
chocolate milk reward and a 3 second auditory cue (2kHz pure tone or white noise,
counterbalanced across animals) or just a 3 second auditory cue (white noise or 2kHz pure
tone). Following either a lever press or omission, a 10 second intertrial interval (ITI) was
triggered. Following two consecutive sessions with over 40 responses, animals progressed to
the third stage. This stage was the same as stage two except that animals now had only 10
seconds to make a response before an omission was registered. Following two consecutive
sessions with over 100 responses animals achieved criterion to progress to the two-armed
bandit task.

Two-armed bandit Task

The two-armed bandit task was performed over of 6 days with each session lasting one hour. In
this task, animals were presented with two levers with one lever rewarded on 80% of trials, and
the other lever rewarded on 20% of trials. To signal the start of the trial, both levers extended
and the cue lights above the levers turned on, animals then had 10 seconds to make a
response on either lever or an omission was registered. A lever press resulted in lever retraction
and immediate delivery of the outcome, either a 30 µL chocolate milk reward and a 3 second
auditory cue (2kHz pure tone or white noise, counterbalanced across animals) or simply a
different 3 second auditory cue (white noise or 2kHz pure tone) signaling non-reward. Following
either a lever press or an omission, a 10 second intertrial interval (ITI) was triggered. To
maintain a dynamic learning environment and high rates of rewarded and unrewarded
outcomes, probability of reward was switched between levers after five consecutive responses
on the high probability lever. Four males and four females remained on the two-armed bandit
task during the task degradation (data not shown).

 One-Lever Forced Choice Task

 The one lever forced choice task was performed over 3 days with each session lasting one
 hour. In this task, animals were presented with a single lever (counterbalanced across animals).
 Pressing this lever resulted in probabilistic reward on a predetermined schedule. The outcome
 schedule was matched to each animal's individual performance in the final three days of the
 two-armed bandit task, such that the first session in the one-lever task was yoked to the reward
 schedule experienced by the animal on day four in the two-armed bandit task, the second to day
 five, and the third to day six. To signal the start of the trial, the lever extended and the cue light
 above the lever turned on. Animals then had 10 seconds to make a response. A lever press
 resulted in lever retraction and immediate delivery of the outcome, either a 30 µL chocolate milk
 reward and a 3 second auditory cue (2kHz pure tone or white noise) or simply a different 3
 second auditory cue (white noise or 2kHz pure tone). Following either a lever press or an
 omission, a 10 second intertrial interval (ITI) was triggered.

 No Lever Response Free Task

 The no lever response free task was performed over the course of 3 days with each session
 lasting one hour. In this task, animals were able to retrieve non-contingently delivered rewards
 under a similar trial structure to both the two-armed bandit task and the one-lever forced choice
 task but with no levers available. To signal the start of the trial, cue lights above both levers
 turned on and remained illuminated for a period of time matched to each animal's response time
 in the last three days of the two-armed bandit task. After cue lights turned off, outcomes were
 delivered, either a 30 µL chocolate milk reward and a 3 second auditory cue (2kHz pure tone or
 white noise) or simply a different 3 second auditory cue (white noise or 2kHz pure tone). As in
 the one-lever task, the outcome schedule was matched to each animal's performance in the
 final three days of the two-armed bandit task now also matching the latency to receive the
 outcome to the trial-by-trial latency to lever press on the two-armed bandit task with a 10 second
 intertrial interval (ITI).

 Frame Independent Projected Fiber Photometry

 To measure calcium-associated changes in fluorescence in real time, recordings were made
 from vHip-NAc and mPFC-NAc-projecting cells during the two-armed bandit task, the one-lever
 forced choice task, and the no lever response free task. Samples were collected at a frequency
 of 20 Hz using Neurophotometrics hardware through Bonsai and FlyCap software. Recordings
 were coupled to the start of behavioral analysis by interfacing Bonsai with MED-PC using a
 custom DAQ box (Neurophotometrics).

 Photometry data extraction and normalization

 Photometry data were extracted and analyzed using custom-written scripts in Python. To
 normalize the data, the control channel (415nm) was fitted to the raw (470nm). The fitted control
 was then subtracted from the raw trace. The resultant trace was divided by the fitted control
 giving the ΔF/F and converted to a Z-score. This calculation was performed over the entirety of
 the session to preserve dynamic fluctuation in population activity that persists beyond individual
 trials to allow comparison across trials. For heatmaps Z-scores were baseline subtracted from

average activity in the two seconds prior to lever press to accommodate moving baselines. For
analyses of reward history, Z-scores were baseline subtracted from average activity in the two
seconds prior to lever press on trial t-1 to account for shifted baselines in trial t0.

Optogenetics in Two-armed Bandit Task

Following lever press training, animals started the two-armed bandit task with optogenetic
manipulations of mPFC-NAc and vHip-NAc activity for the duration of the ITI. Each day animals
received either mPFC-NAc, vHip-NAc, or simultaneous mPFC-NAc and vHip-NAc stimulation on
a subset of trials over the course of 9 days such that they received a total of 3 days of
stimulation per condition for each stimulation protocol tested (5 Hz, 10 ms, 1-2 mW; 8 Hz, 10
ms, 2-3 mW). Order of stimulation days was fully counterbalanced within and between mice to
avoid any order effects. Stimulation was delivered by 450 nm lasers controlled by a laser driver
(Doric) running Doric studios software and triggered via a TTL (Med-Associates) at ITI start on a
random subset of trials (30%) and terminated immediately prior to lever extension.

Ex vivo current-clamp electrophysiology

Brain slice preparation

Mice were deeply anesthetized with isofluorane. Transcardial perfusion was performed with 25-
30 ml of ice-chilled carbogenated NNMDG artificial cerebrospinal fluid (aCSF: containing in mM:
92 NMDG, 2.5 KCl, 1.25 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-
ascorbate, 3 Na-pyruvate, 0.5 $CaCl_2 \cdot 4H_2O$ and 10 $MgSO_4 \cdot 7H_2O$; titrated to pH 7.3–7.4 with
concentrated hydrochloric acid). Brain slices (200 µm) were prepared in ice-chilled
carbogenated NMDG aCSF by a vibratome (Lecia VT 1200S). All brain slices were recovery in
32–34 °C carbogenated NMDG aCSF for 10 min and then were transferred into room-
temperature carbogenated HEPES holding aCSF (containing in mM: 92 NaCl, 2.5 KCl, 1.25
NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-ascorbate, 3 Na-pyruvate, 2
$CaCl_2 \cdot 4H_2O$ and 2 $MgSO_4 \cdot 7H_2O$; titrated to pH to 7.3–7.4 with NaOH) for at least 1 hour before
current-clamp recording.

Electrophysiology recordings

Current-clamp recordings were performed in room-temperature carbogenated aCSF (containing
in mM: mM: 128 NaCl, 3 KCl, 1.25 NaH_2PO_4, 2 $MgCl_2$, 2 $CaCl_2$, 24 $NaHCO_3$ and 10 glucose; pH
7.2). The patch pipette solution was composed of (in mM) 115 K-gluconate, 20 KCl, 1.5 $MgCl_2$,
10 Phosphocreatine-Tris, 2 Mg-ATP, 0.54 Na-GTP and 10 HEPES. Blue light (wavelength: 470
nm) from a LED system (DC4100, Thorlabs) was used for optogenetic stimulation to evoke
action potentials. The optogenetic stimulation protocol consisted of trains of 5 Hz (1-2 mW) or 8
Hz (2-3 mW) 10ms light pulses for 5 s. All signals were amplified and digitized by Multiclamp
700B (Molecular Device) and Digidata 1550B (Molecular Device) respectively. Series and
access resistance were monitored during the experiments and signals were bessel filtered at 2
kHz.

Data Analysis & Statistics

Linear Mixed Effects Regression

Linear Mixed Effects Regression Models are a powerful approach to probe variance attributable
to variables of interest (e.g. trial outcome) while simultaneously controlling for random effects
(e.g. session ID) (Fetcho et al., 2023; Kato et al., 2022; Yu et al., 2022). This is useful for
modeling instances where there is nonindependence in the structure of data e.g. multiple trials
recorded within multiple animals. Models were fit using the full interaction of the factors of
interest (trial outcome, previous trial outcome, sex) and using animal ID and session ID as
random effects using the *lme4* package in R (Bates et al., 2014). Where the dependent variable
was latency, a Gamma link function was used to approximate the non-gaussian distribution. The
fitted models were used to calculate estimated marginal means using the *emmeans* package in
R (Lenth et al., 2021). The effect of variables of interest were then examined by comparing
estimated marginal means. Given the large number of samples generated using this approach
(all trials x all animals), comparisons of estimated marginal means were conducted using a Z-
test and Sidak's method to adjust for multiple comparisons.

Cross-Correlation Time Delay Analysis

Time delay analysis was performed by first calculating the cross-correlation between mPFC-
NAc and vHip-NAc during the ITI across a maximum lag of ± 5 seconds using the CCF function
in R. The argument of the maximum (i.e. the time offset of peak correlation) of the resulting
cross-correlation function was used to estimate the delay between mPFC-NAc and vHip-NAc on
a trial-by-trial basis (Abboud & Sadeh, 1984). Linear mixed effects models were then fit to
assess if the delay was non-zero (i.e. non-synchronous) using the following models to test for
effects of sex [Time Delay~Sex-1+(1|ID)+(1|Day)] and for the interaction between sex and
reward [Time Delay~Rewards:Sex-1+(1|ID)+(1|Day)]. The resulting regression coefficients from
each model were examined to determine if the time delay was non-zero in any group (i.e.
regression coefficient significantly different from zero).
580
581   Conditional Entropy Analysis
582
Conditional entropy is an information measure used to estimate the amount of additional
information needed to explain one signal given full knowledge of a second signal. This can be
interpreted as the unique information contributed by a second signal beyond that contributed by
a first with smaller conditional entropy suggesting less unique information carried by the second
signal. Conditional entropy was calculated on the first two seconds and the last two seconds of
the ITI using the PyInform package in Python to calculate the entropy (□) of the mPFC circuit
given the vHip-NAc circuit, □(mPFC-NAc|vHip-NAc), and the entropy of the vHip-NAc circuit
given the mPFC-NAc circuit, □(vHip-NAc|mPFC-NAc) (Cover & Thomas, 1991; Moore et al.,
2018).
592
593   Code Availability
594
595   Code used to perform analyses for all figures available at
596   https://github.com/eshaaniyer/mPFCvHip-NAc_RewardIntegration
597
ACKNOWLEDGMENTS

We would like to thank Dr. Becket Ebitz, Dr. Mihaela Iordanova and Heike Schuler for their
helpful comments and feedback throughout this project. Renderings of mice were created with
BioRender.com. Research was supported by grants to RCB from NSERC and the Ludmer
Centre for Neuroinformatics and Mental Health.

Author contributions

Conceptualization, E.S.I and R.C.B.; Methodology, E.S.I and R.C.B.; Investigation, E.S.I, P.V.,
S.W., J.M., Y.C.T., V.C.; Writing – Original Draft, E.S.I and R.C.B.; Writing – Review & Editing,
E.S.I. and R.C.B.; Funding Acquisition, R.C.B.; Resources, R.C.B.; Supervision, R.C.B.

Declaration of interests

The authors declare no competing interests.

Figures

Figure 1. mPFC-NAc and vHip-NAc similarly encode reward in a probabilistically
rewarded environment. (A) Schematic of two-armed bandit task. Mice lever press in a two
lever task in which one lever is rewarded with chocolate milk on 80% of trials, and the other on
20%. Following a lever press, levers retract, and auditory cues signal outcome and start of a 10
sec inter-trial interval (ITI). Contingencies switch after five consecutive responses on the high
probability lever. Female (n=10) and male (n=12) mice robustly engage with the task,
experiencing similar numbers of (B) unrewarded (C) rewarded and (D) omission trials. (E)
Retrograding jGCaMP7f is injected into the nucleus accumbens (NAc) and optic fibers
implanted in medial prefrontal cortex (mPFC) and ventral hippocampus (vHip) to simultaneously
probe neural activity indicated by Ca^{2+}-associated fluorescence changes in (F) mPFC neurons
projecting to NAc (mPFC-NAc) and (G) vHip neurons projecting to NAc (vHip-NAc) as mice
encounter reward and non-reward. Estimated mean mPFC-NAc activity across all rewarded and
unrewarded trials in (H) female (n=10) and (I) male (n=12) mice. Analysis focused on 8-10 sec
after lever press (ITI end). At ITI end, mPFC-NAc activity is suppressed by rewarded outcomes
in female (J; Z=21.348, p<0.0001) and male (K; Z=19.625, p<0.0001) mice. Estimated mean
vHip-NAc activity across all rewarded and unrewarded trials in (L) female and (M) male mice. At
ITI end, vHip-NAc activity is suppressed by rewarded outcomes in female (N; Z=8.161;
p<0.0001) and male (O; Z=8.924; p<0.0001) mice. Heatmap of mPFC-NAc activity to (P)
rewarded outcomes and (Q) unrewarded outcomes in a representative animal across one
session. Heatmap of vHip-NAc activity to (R) rewarded outcomes and (S) unrewarded outcomes
in a representative animal across one session. Error bars represent SEM around the estimated
mean. ****p<0.0001

Figure 2. mPFC-NAc and vHip-NAc similarly integrate reward history. (A) Estimated mean
mPFC-NAc activity across pairs of consecutive trials (t-1→t0) showing rewarded+rewarded
(R→R), rewarded+unrewarded (R→U), unrewarded+rewarded (U→R), and

unrewarded+unrewarded (U→U) trial pairs in female (n=10) and male (n=12) mice. Analysis
focused on 8-10 sec after lever press (ITI end). (B) On trial t-1, mPFC-NAc activity is
significantly suppressed by reward (U→U vs R→U: Z=28.99496, p<0.0001; U→R vs R→R:
Z=25.6767, p<0.0001). (C) On the subsequent trial, t0, mPFC-NAc activity is significantly
suppressed by current reward (U→U vs U→R: Z=-28.5098, p<0.0001; R→U vs R→R: Z=-
19.8981, p<0.0001). When trial t0 is unrewarded, mPFC-NAc activity remains significantly
suppressed by reward experienced on the previous trial, t-1, (U→U vs R→U: Z= 9.19653,
p<0.0001). (D) Estimated mean vHip-NAc activity across pairs of consecutive trials (t-1→t0)
showing rewarded+rewarded (R→R), rewarded+unrewarded (R→U), unrewarded+rewarded
(U→R), and unrewarded+unrewarded (U→U) trial pairs. (E) On trial t-1, vHip-NAc activity is
significantly suppressed by reward (U→U vs R→U: Z=14.9372, p<0.0001; U→R vs R→R:
Z=11.6962, p<0.0001). (F) On the subsequent trial, t0, vHip-NAc activity is significantly
suppressed by current reward (U→U vs U→R: Z=-17.4993, p<0.0001; U→U vs R→R: Z=-
7.1005, p<0.0001). When trial t0 is unrewarded, vHip-NAc activity remains suppressed by
reward experienced on the previous trial, t-1, (U→U vs R→U: Z= 11.4112, p<0.0001).
Individual-animal averages are indicated by circles for males and triangles for females. Error
bars represent SEM around the estimated mean. ****p<0.0001
660
Figure 3. mPFC-NAc and vHip-NAc are differentially sensitive to loss. (A) Venn diagram
representing the relationship between the mutual information and conditional entropy that exists
between observed mPFC-NAc and vHip-NAc signals. Conditional entropy is a measure of the
additional unique information contributed by a second signal given fully knowledge of a first
signal. (B) Conditional entropy in mPFC-NAc is reduced on rewarded relative to unrewarded
trials regardless of previous outcome (U→U vs U→R: Z=2.8644, p<0.0001; R→U vs R→R:
Z=3.5185, p<0.0001) indicating that less unique information is carried in mPFC-NAc after
reward. (C) Conditional entropy in vHip-NAc is increased on rewarded relative to unrewarded
trials only when the prior outcome was unrewarded (U→U vs U→R: Z=-3.7566, p=0.0003)
indicating that more unique information is carried in vHip-NAc when reward follows nonreward.
Comparison of activity at ITI end on currently rewarded or unrewarded trials considering prior
outcome history up to three trials back shows that (D) mPFC-NAc activity is suppressed on
every currently rewarded trial indicating that mPFC-NAc consistently encodes current outcome
via relative suppression regardless of outcome history. In contrast, (E) vHip-NAc activity is
suppressed on currently rewarded trials except when current reward is preceded by two
(Z=1.2310, p=0.8606) or three (Z=0.8398, p=0.9834) prior consecutive rewards indicating that
vHip-NAc ceases to encode current outcome via relative suppression after consistent reward.
See supplementary table 29 for all comparisons. Error bars represent SEM around the
estimated mean. *p<0.05, ***p<0.001, ****p<0.0001
680
Figure 4. Degrading task requirements reveals circuit specialization in integrating reward
history. (A) Schematic of one-lever forced choice task in which lever presses are rewarded on
a schedule yoked to each animal's individual performance in the final three days of the two-
armed bandit task. Following a lever press, levers retract, and auditory cues signal outcome and
start of a 10 sec inter-trial interval (ITI). (B) Estimated mean mPFC-NAc activity across pairs of
consecutive trials (t-1→t0) showing rewarded+rewarded (R→R), rewarded+unrewarded

(R→U), unrewarded+rewarded (U→R), and unrewarded+unrewarded (U→U) trial pairs (male
n= 8, female n=6). Analysis focused on 8-10 sec after lever press (ITI end). (C) On trial t0,
mPFC-NAc activity is suppressed by reward (U→U vs U→R: Z=18.8757, p<0.0001; R→U vs
R→R: Z=12.0687, p<0.0001). When trial t0 is unrewarded, mPFC-NAc activity remains
suppressed by reward experienced on the previous trial, t-1, (U→U vs R→U: Z= 6.3467,
p<0.0001). (D) Estimated mean vHip-NAc activity across pairs of consecutive trials (t-1→t0)
showing R→R, R→U, U→R, and U→U trial pairs (male n= 8, female n=6). (E) On trial t0, vHip-
NAc activity is suppressed by reward (U→U vs U→R: Z=8.5245, p<0.0001; R→U vs R→R:
Z=4.0519, p=0.0001). When trial t0 is unrewarded, mPFC-NAc activity remains suppressed by
reward experienced on the previous trial, t-1, (U→U vs R→U: Z= 6.2425, p<0.0001). (F)
Schematic of no lever response free task. Mice are allowed to collect rewards delivered on a
schedule yoked to each animal's individual trial statistics (latency and outcome) of the two-
armed bandit task. Trial structure is signaled by cue-light illumination and after a predetermined
delay auditory cues signal outcome and start of a 10 sec ITI. (G) Estimated mean mPFC-NAc
activity across pairs of consecutive trials (t-1→t0)   showing R→R, R→U, U→R, and U→U trial
pairs (male n= 8, female n=6). (H) On trial t0, mPFC-NAc activity is suppressed by reward
(U→U vs U→R: Z=8.2136, p<0.0001; R→U vs R→R: Z=7.4647, p<0.0001). (I) Estimated mean
vHip-NAc activity across pairs of consecutive trials (t-1→t0) showing R→R, R→U, U→R, and
U→U trial pairs (male n= 8, female n=6). (J) On trial t0, vHip-NAc activity is suppressed by
reward only if trial t-1 was unrewarded (U→U vs U→R: Z=3.7413, p=0.0003). When trial t0 is
unrewarded, vHip-NAc activity remains suppressed by reward experienced on the previous trial,
t-1, (U→U vs R→U: Z= 3.8661, p=0.0002). Individual-animal averages are indicated by circles
for males and triangles for females. Error bars represent SEM around the estimated mean.
***p<0.001, ****p<0.0001
711
Figure 5. mPFC-NAc and vHip-NAc modulate task engagement. (A) Heatmap of estimated
latency to respond on the subsequent trial given mPFC-NAc and vHip-NAc activity at ITI end
shows that increased activity associates with longer latency. (B) Optogenetic stimulation in the
two-armed bandit task is delivered for the duration of the ITI to either mPFC-NAc, vHip-NAc, or
simultaneously to both circuits. (C) AAVrg-ChR2-mCherry or AAVrg-mCherry is injected into the
NAc and optic fibers implanted in mPFC and vHip to stimulate (D) mPFC-NAc neurons and (E)
vHip-NAc neurons. (F) Simultaneous 5Hz stimulation of mPFC-NAc and vHip-NAc, but neither
circuit individually, increased latency to respond in ChR2 animals (male n=6, female n=7)
compared to mCherry controls (male n=6, female n=6; Z=-18.6984, p<0.0001). (G) 8 Hz
stimulation of mPFC-NAc (Z=-12.6970, p=0.01354), vHip-NAc (Z=-23.8073, p<0.0001), and
simultaneous stimulation of both mPFC-NAc and vHip-NAc (Z=-24.1357, p<0.0001) all
increased latency in ChR2 animals (male n= 5, female n=6) compared to mCherry controls
(male n= 6, female n=6). Individual-animal averages are indicated by circles for males and
triangles for females. Error bars represent SEM around the estimated mean.
p<0.01,**p<0.0001
727

728
729 **REFERENCES**
730

Abboud, S., & Sadeh, D. (1984). The use of cross-correlation function for the alignment of ECG waveforms and rejection of extrasystoles. *Computers and Biomedical Research*, *17*(3), 258–266. https://doi.org/10.1016/S0010-4809(84)80017-8

Bagot, R. C., Parise, E. M., Peña, C. J., Zhang, H.-X., Maze, I., Chaudhury, D., Persaud, B., Cachope, R., Bolaños-Guzmán, C. A., Cheer, J. F., Deisseroth, K., Han, M.-H., & Nestler, E. J. (2015). Ventral hippocampal afferents to the nucleus accumbens regulate susceptibility to depression. *Nature Communications*, *6*, 7062. https://doi.org/10.1038/ncomms8062

Bari, B. A., Grossman, C. D., Lubin, E. E., Rajagopalan, A. E., Cressy, J. I., & Cohen, J. Y. (2019). Stable representations of decision variables for flexible behavior. *Neuron*, *103*(5), 922-933.e7. https://doi.org/10.1016/j.neuron.2019.06.001

Barker, J. M., Bryant, K. G., & Chandler, L. J. (2019). Inactivation of ventral hippocampus projections promotes sensitivity to changes in contingency. *Learning & Memory*, *26*(1), 1–8.

Barrientos, C., Knowland, D., Wu, M. M. J., Lilascharoen, V., Huang, K. W., Malenka, R. C., & Lim, B. K. (2018). Cocaine-Induced Structural Plasticity in Input Regions to Distinct Cell Types in Nucleus Accumbens. *Biological Psychiatry*, *84*(12), 893–904. https://doi.org/10.1016/j.biopsych.2018.04.019

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv Preprint arXiv:1406.5823*.

Beierholm, U., Guitart-Masip, M., Economides, M., Chowdhury, R., Düzel, E., Dolan, R., & Dayan, P. (2013). Dopamine Modulates Reward-Related Vigor. *Neuropsychopharmacology*, *38*(8), 1495–1503. https://doi.org/10.1038/npp.2013.48

Britt, J. P., Benaliouad, F., McDevitt, R. A., Stuber, G. D., Wise, R. A., & Bonci, A. (2012). Synaptic and Behavioral Profile of Multiple Glutamatergic Inputs to the Nucleus Accumbens. *Neuron*, *76*(4), 790–803. https://doi.org/10.1016/j.neuron.2012.09.040

Cahill, M. E., Bagot, R. C., Gancarz, A. M., Walker, D. M., Sun, H., Wang, Z.-J., Heller, E. A., Feng, J., Kennedy, P. J., Koo, J. W., Cates, H. M., Neve, R. L., Shen, L., Dietz, D. M., & Nestler, E. J. (2016). Bidirectional Synaptic Structural Plasticity after Chronic Cocaine Administration Occurs through Rap1 Small GTPase Signaling. *Neuron*, *89*(3), 566–582. https://doi.org/10.1016/j.neuron.2016.01.031

Carter, A. G., Soler-Llavina, G. J., & Sabatini, B. L. (2007). Timing and Location of Synaptic Inputs Determine Modes of Subthreshold Integration in Striatal Medium Spiny Neurons. *The Journal of Neuroscience*, *27*(33), 8967–8977. https://doi.org/10.1523/JNEUROSCI.2798-07.2007

Chen, G., Lai, S., Bao, G., Ke, J., Meng, X., Lu, S., Wu, X., Xu, H., Wu, F., Xu, Y., Xu, F., Bi, G.-Q., Peng, G., Zhou, K., & Zhu, Y. (2023). Distinct reward processing by subregions of the nucleus accumbens. *Cell Reports*, *42*(2). https://doi.org/10.1016/j.celrep.2023.112069

Christoffel, D. J., Walsh, J. J., Hoerbelt, P., Heifets, B. D., Llorach, P., Lopez, R. C., Ramakrishnan, C., Deisseroth, K., & Malenka, R. C. (2021). Selective filtering of excitatory inputs to nucleus accumbens by dopamine and serotonin. *Proceedings of the National Academy of Sciences*, *118*(24), e2106648118. https://doi.org/10.1073/pnas.2106648118

Cover, T. M., & Thomas, J. A. (1991). Entropy, relative entropy and mutual information. *Elements of Information Theory*, *2*(1), 12–13.

Cox, J., Minerva, A. R., Fleming, W. T., Zimmerman, C. A., Hayes, C., Zorowitz, S., Bandi, A., Ornelas, S., McMannon, B., Parker, N. F., & Witten, I. B. (2023). A neural substrate of sex-dependent modulation of motivation. *Nature Neuroscience*, *26*(2), Article 2. https://doi.org/10.1038/s41593-022-01229-9

Dana, H., Sun, Y., Mohar, B., Hulse, B. K., Kerlin, A. M., Hasseman, J. P., Tsegaye, G., Tsang, A., Wong, A., Patel, R., Macklin, J. J., Chen, Y., Konnerth, A., Jayaraman, V., Looger, L. L., Schreiter, E. R., Svoboda, K., & Kim, D. S. (2019). High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nature Methods*, *16*(7), 649–657. https://doi.org/10.1038/s41592-019-0435-6

Fetcho, R. N., Hall, B. S., Estrin, D. J., Walsh, A. P., Schuette, P. J., Kaminsky, J., Singh, A., Roshgodal, J., Bavley, C. C., Nadkarni, V., Antigua, S., Huynh, T. N., Grosenick, L., Carthy, C., Komer, L., Adhikari, A., Lee, F. S., Rajadhyaksha, A. M., & Liston, C. (2023). Regulation of social interaction in mice by a frontostriatal circuit modulated by established hierarchical relationships. *Nature Communications*, *14*, 2487. https://doi.org/10.1038/s41467-023-37460-6

Floresco, S. B. (2015). The Nucleus Accumbens: An Interface Between Cognition, Emotion, and Action. *Annual Review of Psychology*, *66*(1), 25–52. https://doi.org/10.1146/annurev-psych-010213-115159

French, S. J., & Totterdell, S. (2002). Hippocampal and prefrontal cortical inputs monosynaptically converge with individual projection neurons of the nucleus accumbens. *Journal of Comparative Neurology*, *446*(2), 151–165.

Ghanbari, M., Li, G., Hsu, L., & Yap, P. (2023). Accumulation of network redundancy marks the early stage of Alzheimer's disease. *Human Brain Mapping*, *44*(8), 2993–3006. https://doi.org/10.1002/hbm.26257

Grace, A. A., Floresco, S. B., Goto, Y., & Lodge, D. J. (2007). Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends in Neurosciences*, *30*(5), 220–227. https://doi.org/10.1016/j.tins.2007.03.003

Griffin, W. C., Lopez, M. F., Woodward, J. J., & Becker, H. C. (2023). Alcohol dependence and the ventral hippocampal influence on alcohol drinking in male mice. *Alcohol*, *106*, 44–54. https://doi.org/10.1016/j.alcohol.2022.10.004

Hamel, L., Cavdaroglu, B., Yeates, D., Nguyen, D., Riaz, S., Patterson, D., Khan, N., Kirolos, N., Roper, K., Ha, Q. A., & Ito, R. (2022). Cortico-Striatal Control over Adaptive Goal-Directed Responding Elicited by Cues Signaling Sucrose Reward or Punishment. *Journal of Neuroscience*, *42*(18), 3811–3822. https://doi.org/10.1523/JNEUROSCI.2175-21.2022

Hamid, A. A., Pettibone, J. R., Mabrouk, O. S., Hetrick, V. L., Schmidt, R., Vander Weele, C. M., Kennedy, R. T., Aragona, B. J., & Berke, J. D. (2016). Mesolimbic Dopamine Signals the Value of Work. *Nature Neuroscience*, *19*(1), 117–126. https://doi.org/10.1038/nn.4173

Hiratani, N., & Fukai, T. (2018). Redundancy in synaptic connections enables neurons to learn optimally. *Proceedings of the National Academy of Sciences*, *115*(29), E6871–E6879. https://doi.org/10.1073/pnas.1803274115

Kato, T., Mitsukura, Y., Yoshida, K., Mimura, M., Takata, N., & Tanaka, K. F. (2022). Oscillatory Population-Level Activity of Dorsal Raphe Serotonergic Neurons Is Inscribed in Sleep Structure. *The Journal of Neuroscience*, *42*(38), 7244–7255. https://doi.org/10.1523/JNEUROSCI.2288-21.2022

Fischer, D. M., Aziz, H., Mangieri, R. A., & Morrisett, R. A. (2019). Ethanol Experience Enhances Glutamatergic Ventral Hippocampal Inputs To D1 Receptor-Expressing Medium Spiny Neurons In The Nucleus Accumbens Shell. *The Journal of Neuroscience*, 3051–18. https://doi.org/10.1523/JNEUROSCI.3051-18.2019

Lafferty, C. K., Yang, A. K., Mendoza, J. A., & Britt, J. P. (2020). Nucleus Accumbens Cell Type- and Input-Specific Suppression of Unproductive Reward Seeking. *Cell Reports*, *30*(11), 3729-3742.e3. https://doi.org/10.1016/j.celrep.2020.02.095

Lenth, R. V., Buerkner, P., Herve, M., Love, J., Riebl, H., & Singmann, H. (2021). *Emmeans: Estimated marginal means, aka least-squares means. CRAN*.

LeGates, E. B., Sweis, B. M., Asp, A. J., Esguerra, M., Silvis, K. A., David Redish, A., & Thomas, M. J. (2023). A quadruple dissociation of reward-related behaviour in mice across excitatory inputs to the nucleus accumbens shell. *Communications Biology*, *6*(1), Article 1. https://doi.org/10.1038/s42003-023-04429-6

Loewenbach, D., Vacca, G., Ahn, S., Seamans, J. K., & Phillips, A. G. (2022). Optogenetic modulation of glutamatergic afferents from the ventral subiculum to the nucleus accumbens: Effects on dopamine function, response vigor and locomotor activity. *Behavioural Brain Research*, *434*, 114028. https://doi.org/10.1016/j.bbr.2022.114028

Marder, E. (2011). Variability, compensation, and modulation in neurons and circuits. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(Suppl 3), 15542–15548. https://doi.org/10.1073/pnas.1010674108

Marder, E., & Goaillard, J.-M. (2006). Variability, compensation and homeostasis in neuron and network function. *Nature Reviews Neuroscience*, *7*(7), 563–574. https://doi.org/10.1038/nrn1949

Millan, E. Z., Kim, H. A., & Janak, P. H. (2017). Optogenetic activation of amygdala projections to nucleus accumbens can arrest conditioned and unconditioned alcohol consummatory behavior. *Neuroscience*, *360*, 106–117. https://doi.org/10.1016/j.neuroscience.2017.07.044

Mishra, B. E. P., & O'Donnell, C. (2021). Neural circuit function redundancy in brain disorders. *Current Opinion in Neurobiology*, *70*, 74–80. https://doi.org/10.1016/j.conb.2021.07.008

Moore, D. G., Valentini, G., Walker, S. I., & Levin, M. (2018). Inform: Efficient information-theoretic analysis of collective behaviors. *Frontiers in Robotics and AI*, *5*, 60.

Nair, J., Iyer, E. S., Tse, Y.-C., Sorensen, J., Wu, S., Eid, R. S., Cvetkovska, V., Wassef, K., Gostlin, S., Vitaro, P., Spencer, N. J., & Bagot, R. C. (2024). Sex-biased neural encoding of threat discrimination in nucleus accumbens afferents drives suppression of reward behavior. *Nature Neuroscience*, *27*(10), 1966–1976. https://doi.org/10.1038/s41593-024-01748-7

Nair, J., Tse, Y. C., Iyer, E. S., Biris, J., Cvetkovska, V., Lopez, J., & Bagot, R. C. (2020). Ventral Hippocampal Afferents to Nucleus Accumbens Encode Both Latent Vulnerability and Stress-Induced Susceptibility. *Biological Psychiatry*, *88*(11), 843–854. https://doi.org/10.1016/j.biopsych.2020.05.021

Nguyen, A. T., Xu, J., Luu, D. K., Zhao, Q., & Yang, Z. (2019). Advancing System Performance with Redundancy: From Biological to Artificial Designs. *Neural Computation*, *31*(3), 555–573. https://doi.org/10.1162/neco_a_01166

Niv, Y., Daw, N. D., Joel, D., & Dayan, P. (2007). Tonic dopamine: Opportunity costs and the control of response vigor. *Psychopharmacology*, *191*(3), 507–520. https://doi.org/10.1007/s00213-006-0502-4

O'Donnell, P., & Grace, A. (1995). Synaptic interactions among excitatory afferents to nucleus
accumbens neurons: Hippocampal gating of prefrontal cortical input. *The Journal of*
Neuroscience, *15*(5), 3622–3639. https://doi.org/10.1523/JNEUROSCI.15-05-03622.1995

Otis, J. M., Namboodiri, V. M. K., Matan, A. M., Voets, E. S., Mohorn, E. P., Kosyk, O., McHenry, J. A.,
Robinson, J. E., Resendez, S. L., Rossi, M. A., & Stuber, G. D. (2017). Prefrontal cortex output
circuits guide reward seeking through divergent cue encoding. *Nature*, *543*(7643), Article 7643.
https://doi.org/10.1038/nature21376

Parker, N. F., Baidya, A., Cox, J., Haetzel, L. M., Zhukovskaya, A., Murugan, M., Engelhard, B.,
Goldman, M. S., & Witten, I. B. (2022). Choice-selective sequences dominate in cortical relative
to thalamic inputs to NAc to support reinforcement learning. *Cell Reports*, *39*(7).
https://doi.org/10.1016/j.celrep.2022.110756

Pascoli, V., Terrier, J., Espallergues, J., Valjent, E., O'Connor, E. C., & Lüscher, C. (2014). Contrasting
forms of cocaine-evoked plasticity control components of relapse. *Nature*, *509*(7501), Article
7501. https://doi.org/10.1038/nature13257

Pignatelli, M., Tejeda, H. A., Barker, D. J., Bontempi, L., Wu, J., Lopez, A., Palma Ribeiro, S.,
Lucantonio, F., Parise, E. M., Torres-Berrio, A., Alvarez-Bagnarol, Y., Marino, R. A. M., Cai, Z.-
L., Xue, M., Morales, M., Tamminga, C. A., Nestler, E. J., & Bonci, A. (2021). Cooperative
synaptic and intrinsic plasticity in a disynaptic limbic circuit drive stress-induced anhedonia and
passive coping in mice. *Molecular Psychiatry*, *26*(6), Article 6. https://doi.org/10.1038/s41380-
020-0686-8

Reed, S. J., Lafferty, C. K., Mendoza, J. A., Yang, A. K., Davidson, T. J., Grosenick, L., Deisseroth, K.,
& Britt, J. P. (2018). Coordinated Reductions in Excitatory Input to the Nucleus Accumbens
Underlie Food Consumption. *Neuron*, *99*(6), 1260-1273.e4.
https://doi.org/10.1016/j.neuron.2018.07.051

Spellman, T., Svei, M., Kaminsky, J., Manzano-Nieves, G., & Liston, C. (2021). Prefrontal deep
projection neurons enable cognitive flexibility via persistent feedback monitoring. *Cell*, *184*(10),
2750-2766.e17. https://doi.org/10.1016/j.cell.2021.03.047

Sul, J. H., Kim, H., Huh, N., Lee, D., & Jung, M. W. (2010). Distinct Roles of Rodent Orbitofrontal and
Medial Prefrontal Cortex in Decision Making. *Neuron*, *66*(3), 449–460.
https://doi.org/10.1016/j.neuron.2010.03.033

Wenzel, J. M., Zlebnik, N. E., Patton, M. H., Smethells, J. R., Ayvazian, V. M., Dantrassy, H. M., Zhang,
L.-Y., Mathur, B. N., & Cheer, J. F. (2023). Selective chemogenetic inactivation of
corticoaccumbal projections disrupts trait choice impulsivity. *Neuropsychopharmacology*,
48(12), Article 12. https://doi.org/10.1038/s41386-023-01604-5

Williams, E. S., Manning, C. E., Eagle, A. L., Swift-Gallant, A., Duque-Wilckens, N., Chinnusamy, S.,
Moeser, A., Jordan, C., Leininger, G., & Robison, A. J. (2020). Androgen-Dependent
Excitability of Mouse Ventral Hippocampal Afferents to Nucleus Accumbens Underlies Sex-
Specific Susceptibility to Stress. *Biological Psychiatry*, *87*(6), 492–501.
https://doi.org/10.1016/j.biopsych.2019.08.006

Yoshida, K., Drew, M. R., Kono, A., Mimura, M., Takata, N., & Tanaka, K. F. (2021). Chronic social
defeat stress impairs goal-directed behavior through dysregulation of ventral hippocampal
activity in male mice. *Neuropsychopharmacology*, *46*(9), 1606–1616.
https://doi.org/10.1038/s41386-021-00990-y

Yoshida, K., Drew, M. R., Mimura, M., & Tanaka, K. F. (2019). Serotonin-mediated inhibition of ventral hippocampus is required for sustained goal-directed behavior. *Nature Neuroscience*, *22*(5), Article 5. https://doi.org/10.1038/s41593-019-0376-5

Yoshida, K., Tsutsui-Kimura, I., Kono, A., Yamanaka, A., Kobayashi, K., Watanabe, M., Mimura, M., & Tanaka, K. F. (2020). Opposing Ventral Striatal Medium Spiny Neuron Activities Shaped by Striatal Parvalbumin-Expressing Interneurons during Goal-Directed Behaviors. *Cell Reports*, *31*(13), 107829. https://doi.org/10.1016/j.celrep.2020.107829

Yu, Z., Guindani, M., Grieco, S. F., Chen, L., Holmes, T. C., & Xu, X. (2022). Beyond t test and ANOVA: Applications of mixed-effects models for more rigorous statistical analysis in neuroscience research. *Neuron*, *110*(1), 21–35. https://doi.org/10.1016/j.neuron.2021.10.030

Zinsmaier, A. K., Dong, Y., & Huang, Y. H. (2022). Cocaine-induced projection-specific and cell type-specific adaptations in the nucleus accumbens. *Molecular Psychiatry*, *27*(1), Article 1. https://doi.org/10.1038/s41380-021-01112-2

A

One Lever Forced Choice Task

Rewarded
Unrewarded

rewarded

3s

unrewarded

3s

7s

Reward Distribution
Yoked to PRL Performance

F

No Lever Response Free Task

Rewarded
Unrewarded

rewarded

3s

unrewarded

3s

7s

Reward Distribution
Yoked to PRL Performance

mPFC-NAc

B

Lever Press
Outcome

T -1

ITI End
Lever Extends

Lever Press
Outcome

T 0

ITI End
Lever Extends

0.1
zScore

1s

G

Cue Light Off
Outcome

T -1

ITI End
Cue Light Off

Cue Light Off
Outcome

T 0

ITI End
Cue Light Off

0.05
zScore

1s

Rewarded
Rewarded

Rewarded
Unrewarded

Unrewarded
Rewarded

Unrewarded
Unrewarded

C

mPFC (zSc0re)

0.4
0.2
0.0
-0.2

****

****

****

T -1   Rewarded   Rewarded   Unrewarded   Unrewarded
T 0   Rewarded   Unrewarded Rewarded   Unrewarded

H

mPFC (zScore)

0.1
0.0
-0.1
-0.2

****

****

T -1   Rewarded   Rewarded   Unrewarded   Unrewarded
T 0   Rewarded   Unrewarded Rewarded   Unrewarded

vHip-NAc

D

Lever Press
Outcome

T -1

ITI End
Lever Extends

Lever Press
Outcome

T 0

ITI End
Lever Extends

0.1
zScore

1s

I

Cue Light Off
Outcome

Trial 1

ITI End
Cue Light Off

Cue Light Off
Outcome

T 0

ITI End
Cue Light Off

0.05
zScore

1s

Rewarded
Rewarded

Rewarded
Unrewarded

Unrewarded
Rewarded

Unrewarded
Unrewarded

E

vHip (zScore)

0.2
0.1
0.0
-0.1
-0.2
-0.3

***

****

****

T -1   Rewarded   Rewarded   Unrewarded   Unrewarded
T 0   Rewarded   Unrewarded Rewarded   Unrewarded

J

vHip (zScore)

0.1
0.0
-0.1
-0.2

***

***

T -1   Rewarded   Rewarded   Unrewarded   Unrewarded
T 0   Rewarded   Unrewarded Rewarded   Unrewarded

A — Latency on Next Trial (Sec); vHip-NAc Activity (Z-Score) vs mPFC-NAc Activity (Z-Score)

B — Reward Probability 80% / 20%; rewarded / unrewarded; 3s; 7s; Reward Probability 80% / 20%

C — AAVrg-ChR2/AAVrg-mCherry; mPFC; NAc; vHip

D — ChR2 / DAPI

E — ChR2 / DAPI

F — 5 Hz Stimulation; Percent Change in Latency; mCherry / ChR2; No Stim, mPFC, vHip, mPFC+vHip; ****

G — 8 Hz Stimulation; Percent Change in Latency; No Stim, mPFC, vHip, mPFC+vHip; **, ****, ****