1

LasR regulates protease IV expression at suboptimal growth temperatures in

Pseudomonas aeruginosa

Running title: LasR regulates protease IV expression

Rachel E. Done[a,b], Joshua K. Robertson[c], Samantha M. Prezioso[a,b*], and Joanna B. Goldberg[b,d#]

[a]Microbiology and Molecular Genetics Program, Graduate Division of Biological and Biomedical Sciences, Laney Graduate School, Emory University, Atlanta, Georgia, USA

[b]Department of Pediatrics, Division of Pulmonary, Asthma, Cystic Fibrosis, and Sleep, Emory University School of Medicine, Atlanta, Georgia, USA

[c]Department of Biology, Emory University, Atlanta, Georgia, USA

[d]Emory+Children's Center for Cystic Fibrosis and Airway Disease Research, Emory University School of Medicine, Atlanta, Georgia, USA

*Present address: Alltrna, Boston, Massachusetts, USA

Abstract word count: 238 words

Importance word count: 134 words

Manuscript text word count: 4,941 words

Address correspondence to Joanna B. Goldberg, joanna.goldberg@emory.edu.

ABSTRACT

The opportunistic pathogen *Pseudomonas aeruginosa* causes debilitating lung infections in people with cystic fibrosis, as well as eye, burn, and wound infections in otherwise immunocompetent individuals. Many of *P. aeruginosa*'s virulence factors are regulated by environmental changes associated with human infection, such as a change in temperature from ambient to human body temperature. One such virulence factor is protease IV (PIV). Interestingly, *piv* expression is higher at ambient temperatures (22-28°C) compared to human body temperature (37°C). We found that *piv* expression was thermoregulated at stationary phase, but not exponential phase, and that *piv* is thermoregulated at the level of transcription. Protein levels of known transcriptional regulators of *piv*, the quorum sensing regulator LasR and the gene-silencing histone nucleoid silencing proteins MvaT/MvaU, were not thermoregulated. Using a transcriptional reporter for *piv*, we show that LasR activates *piv* expression at stationary phase at 25°C but not 37°C, while MvaT/MvaU are not required for *piv* thermoregulation. We also identified a *las* box in the *piv* promoter, which is important for *piv* thermoregulation. We propose that LasR directly regulates *piv* at stationary phase at 25°C but has a negligible impact at 37°C. Here, we show that *piv* is uniquely regulated by LasR in a temperature-dependent manner. Our findings suggest that the LasRI quorum sensing regulon of *P. aeruginosa* may not be fully characterized and that growth at non-standard laboratory conditions such as lower temperatures could reveal previously unrecognized quorum sensing regulated genes.

IMPORTANCE

Pseudomonas aeruginosa is a versatile opportunistic pathogen capable of causing many different types of infections that are often difficult to treat, such as lung infections in people with cystic fibrosis. Temperature regulates the expression of many virulence factors that contribute to *P. aeruginosa*'s ability to cause infection, yet our mechanistic understanding of virulence factor

thermoregulation is poor. In this study, we show that the virulence factor protease IV is

thermoregulated at the level of transcription via temperature-dependent upregulation by the

quorum sensing regulator, LasR. Mechanistic studies of virulence factor thermoregulation will

expand our understanding of how *P. aeruginosa* experiences different environments, including

the mammalian host. Our work also highlights the importance of growth conditions in studying

gene regulation, as it elucidates the regulation of protease IV by LasR, which was previously not

well understood.

INTRODUCTION

Pseudomonas aeruginosa is a highly adaptable bacterial pathogen that causes lung, burn, wound, and corneal infections. To survive in the diverse niches of the human body, *P. aeruginosa* adapts to its environment and regulates the expression of many virulence factors that assist in nutrient acquisition and immune evasion. A major environmental cue that regulates virulence in many pathogenic bacteria, including *P. aeruginosa*, is the change in temperature from ambient or room temperature to that encountered in or on the human body (1, 2). Human body temperature is generally accepted to be 37°C but can be lower at body sites exposed to the environment, such as the skin and eyes.

Many virulence factors in *P. aeruginosa* are regulated by temperature and mechanisms of thermoregulation have been studied at the post-transcriptional (2) and post-translational levels (3–5). Thermoregulation at the level of transcription has not been as extensively studied, despite the finding that expression of 6.4% of the genome is regulated by temperature, with many virulence factors upregulated at 37°C compared to ambient temperatures (6, 7). Interestingly, expression of the gene encoding the virulence factor protease IV (PIV) is greatly upregulated at ambient temperatures compared to 37°C in both of the laboratory strains PAO1 and PA14 (6, 7). PIV is a serine protease that degrades a broad range of immune molecules, including the iron sequestering lactoferrin and transferrin, complement C3 found in blood, as well as alveolar surfactant proteins and IL-22 that provide protection in the lungs (8–11). PIV has been extensively characterized as a key virulence factor for *P. aeruginosa* eye infections and strains lacking PIV are rendered avirulent in corneal infections (12–15). Corneal infections can be difficult to treat with severe patient outcomes, as seen in a recent carbapenem-resistant *P. aeruginosa* outbreak caused by contaminated artificial tears that resulted in multiple deaths, instances of moderate to complete loss of vision, and enucleation due to the infection (16).

PIV is initially translated as a ~48 kD pre-proenzyme that contains an N-terminal signal

sequence, an inactivating propeptide, and the protease domain (17). The signal sequence is

cleaved during secretion and the remaining propeptide renders the 45 kD proenzyme

catalytically inactive; release of the propeptide is required to yield the 26 kD, enzymatically

active ("mature"), protease. Cleavage of the propeptide has been reported to occur

autocatalytically and by the *P. aeruginosa* elastase LasB in bacterial supernatants (9, 18, 19).

Transcriptional regulation of *piv* is also complex and not fully understood. *piv* expression

is directly upregulated by the sigma factor PvdS under iron starvation conditions, leading to the

gene's alternative name *prpL* (P̲vdS-r̲egulated endop̲rotease, l̲ysyl class) (20). *piv* expression is

directly repressed by the histone nucleoid structuring (H-NS) family members MvaT and MvaU,

which act together to negatively regulate the same global regulon (21, 22). *piv* is also regulated

by the LasRI quorum sensing system (23–25). The LasRI quorum sensing system is a

mechanism for *P. aeruginosa* to coordinate expression of certain genes at high cell densities

through the audoinducer synthase LasI synthesizing the freely diffusible homoserine lactone

(HSL) autoinducer 3-oxo-C12-HSL (3O-C12-HSL), which complexes with transcription factor

LasR to activate its function as a positive regulator of hundreds of genes (23–26). LasR has not

been found to directly bind to the *piv* promoter and the mechanism(s) by which it regulates *piv* is

not clear (27). Additionally, how *piv* expression is regulated by temperature is not understood

and is important for our understanding of how *P. aeruginosa* adapts to body sites which can

experience lower temperatures due to exposure to the environment.

We found that *piv* thermoregulation occurs at the level of transcription due to significantly

higher promoter activity at room temperature (25°C) compared to human body temperature

(37°C). Higher *piv* expression at ambient temperatures was reflected in a higher level of mature

PIV protein, suggesting that temperature regulation could affect the amount of this virulence

factor produced by *P. aeruginosa* in infection sites experiencing different temperatures. Our

study outlines the role of LasR as an activator of *piv* expression at 25°C but not 37°C and

provides evidence for direct LasR regulation of *piv*. This suggests that LasRI quorum sensing

may regulate certain genes disparately under conditions different from those standardly used in

the laboratory.

RESULTS

Thermoregulation of *piv* depends on growth phase.

A previous transcriptomic study from our group showed by microarray analysis that *piv* was

more highly expressed at 22°C compared to 37°C in *P. aeruginosa* PAO1 (6). Similar results

were found in an RNA-seq analysis of *P. aeruginosa* PA14 grown at 28°C compared to 37°C (7).

However, both transcriptomes were determined for cells at stationary phase and only examined

two temperatures: a "low" or ambient temperature and human body temperature, 37°C. We

wanted to examine *piv* expression at additional temperatures to characterize if *piv*

thermoregulation is binary with a 'high' and 'low' expression state or continuous with expression

changing gradually as temperature changes. We also wanted to determine if growth phase

affected *piv* thermoregulation. Wild-type *P. aeruginosa* PAO1 was grown in parallel at 25°C,

30°C, 37°C, and 42°C and RNA extracted from cultures at each temperature, first at exponential

phase and then from the same cultures at early stationary phase (Fig. 1A). *piv* expression at

each temperature was determined relative to 37°C (a standard laboratory condition) at each

growth phase using RT-qPCR. Interestingly, we found that *piv* thermoregulation depended on

growth phase. At exponential phase, *piv* expression was not regulated by temperature (Fig. 1B).

However, at stationary phase, *piv* expression at 25°C and 30°C was ~16-fold and ~10-fold

higher, respectively, than at 37°C and ~6-fold lower at 42°C than 37°C. These results support

piv thermoregulation as continuous within the range of tested temperatures, with highest

expression observed at 25°C and expression decreasing gradually as temperature increases to

42°C.

We then determined whether *piv* transcript levels were reflected in levels of PIV protein

in cells grown at different temperatures. To test this, we used PAO1 PIV VSV-G, in which the

native *piv* locus was tagged at the C-terminus with the vesicular stomatitis virus G (VSV-G)

epitope. The C-terminal placement of the VSV-G tag would allow for detecting the 48 kD pre-

proenzyme, the 45 kD proenzyme, and the 26 kD mature enzymatic forms of PIV, as processing

of the full-length gene product involves proteolysis from the N-terminus (9, 17). PAO1 PIV VSV-

G was grown at 25°C, 30°C, 37°C, and 42°C to early stationary phase and cell lysates were

subjected to quantitative western blotting with VSV-G antibodies (αVSV-G) as well as with RNA

polymerase α subunit antibodies (αRpoA) as a temperature independent control (Fig 2A,B). We

found that the abundance of the 26 kD mature PIV VSV-G domain was significantly higher at

25°C and 30°C than at 37°C. Mature PIV VSV-G was hardly detected at 42°C. An unprocessed

form of PIV could be detected at approximately 50 kD at 25°C and 30°C but not at 37°C or

42°C; however, we were not able to distinguish between the 48 kD pre-proenzyme and 45 kD

proenzyme forms of PIV. The abundance of mature PIV VSV-G at each temperature agrees well

with *piv* expression data, suggesting there is likely not temperature-dependent post-

transcriptional modification of *piv*. Together, these results show that *piv* thermoregulation

depends on growth phase and thermoregulation of *piv* gene expression is reflected in the

abundance of PIV protein.

Transcriptional regulators of *piv* are not thermoregulated.

Previous studies have shown that LasRI quorum sensing positively regulates *piv* (23–25) and

the H-NS family members MvaT and MvaU directly repress *piv* expression (21, 22). One

possible mechanism for the thermoregulation of *piv* expression could be thermoregulation of

transcriptional regulators of *piv*. We asked if the quorum sensing regulator LasR was more

abundant at 25°C than 37°C, and/or if the repressors MvaT and MvaU were more abundant at

37°C than 25°C, as a possible mechanism for *piv* thermoregulation. We first examined gene

expression of *lasR*, *mvaT*, and *mvaU* in PAO1 grown at 25°C and 37°C at both exponential and

stationary phases (Fig. 3A). The expression of each transcriptional regulator was not

significantly regulated by temperature, even at stationary phase, when *piv* is most

thermoregulated. This suggested that LasR, MvaT, and MvaU are not themselves

thermoregulated, which is consistent with other studies that did not identify these genes as

thermoregulated (6, 7). To confirm that the protein abundance of LasR, MvaT, and MvaU was

not regulated by temperature, we assessed the level of each protein in cells grown at 25°C and

37°C to stationary phase by quantitative western blotting. Blotting in PAO1 with antibodies

raised to LasR in rabbits (25) confirmed that the level of LasR protein was not significantly

different at 25°C compared 37°C (Fig. 3B). Blotting with αVSV-G antibodies in the PAO1 MvaT

VSV-G (Fig. 3C) and PAO1 MvaU VSV-G strains (Fig. 3D) with C-terminal VSV-G epitope

tagged *mvaT* and *mvaU* at the native loci also showed that levels of MvaT VSV-G and MvaU

VSV-G were also not significantly different at 25°C compared to 37°C. These results indicate

that transcriptional regulators LasR, MvaT, and MvaU are not themselves thermoregulated.

***piv* expression is regulated by temperature at the level of transcription.**

Another possible mechanism for the thermoregulation of *piv* expression could be temperature

dependent *piv* promoter activity. To test this, we made a transcriptional reporter by fusing the

200 bp upstream of the *piv* coding sequence to an unstable green fluorescent protein variant,

gfp(ASV) (28) that has been previously used for transcriptional reporters (29). PAO1 carrying

the P$_{piv}$-*gfp*(ASV) reporter plasmid (pRD91) was grown at 25°C and 37°C and assayed for

fluorescence at exponential phase (Fig. 4A) and then again at stationary phase (Fig. 4B). PAO1

carrying a no-promoter *gfp*(ASV) reporter plasmid (pRD87) was used to assess background

fluorescence, which was overall low at both temperatures and growth phases (Fig. S1). At both

exponential and stationary phases, *piv* promoter activity in PAO1 was higher at 25°C than 37°C,

with promoter activity at 37°C not above the background (Fig. 4A,B). Promoter activity at 25°C

was also higher at stationary phase than exponential phase. The finding that promoter activity

was thermoregulated and affected by growth phase suggests that the growth phase dependent

thermoregulation of *piv* observed in Fig. 1 is due to differences in *piv* promoter activity.

181   Although the abundance of transcriptional regulators LasR, MvaT, and MvaU was not

thermoregulated, we wondered if their regulation of *piv* was affected by temperature since the

piv promoter is sufficient to drive thermoregulation. To test if LasR was required for *piv*

thermoregulation, we assayed a Δ*lasR* mutant carrying the P$_{piv}$-*gfp*(ASV) reporter for

fluorescence at 25°C and 37°C as described; there was no promoter activity above background

levels at either 25°C or 37°C, at both exponential and stationary phases (Fig. 4A, B). Expressing

LasR in single copy from the ectopic Tn*7* site complemented the Δ*lasR* mutation, restoring wild-

type *piv* promoter thermoregulation. This shows that LasR is required for *piv* promoter activity at

both 25°C and 37°C. To further confirm that *piv* was regulated by LasRI quorum sensing, we

deleted the gene encoding the LasI autoinducer synthase and assayed the Δ*lasI* mutant for

activity from the *piv* transcriptional reporter (Fig. 4A,B). Promoter activity in the Δ*lasI* mutant

phenocopied that of the Δ*lasR* mutant, supporting *piv* being regulated by LasRI quorum sensing.

193   In the Δ*mvaT*Δ*mvaU* mutant, promoter activity at exponential phase was overall higher

at both temperatures than in PAO1 and was slightly thermoregulated; however, at stationary

phase promoter activity was highly thermoregulated (Fig. 4). Expressing either MvaT or MvaU in

single copy from the ectopic Tn*7* site complemented the Δ*mvaT*Δ*mvaU* mutation. Since

promoter activity is thermoregulated at stationary phase in the Δ*mvaT*Δ*mvaU* mutant, this result

indicates that MvaT and MvaU are not required for *piv* thermoregulation. To test if LasR was

responsible for thermoregulation in the Δ*mvaT*Δ*mvaU* mutant and to assess the role of LasR in

piv thermoregulation in the absence of strong repressors MvaT and MvaU, we deleted *lasR*

from the Δ*mvaT*Δ*mvaU* strain. At exponential phase, promoter activity in Δ*mvaT*Δ*mvaU*Δ*lasR*

was overall derepressed and slightly thermoregulated, phenocopying Δ*mvaT*Δ*mvaU* (Fig. 4A,B).

However, at stationary phase, promoter activity in Δ*mvaT*Δ*mvaU*Δ*lasR* was significantly less

thermoregulated due to a decrease in *piv* promoter activity at 25°C with no change at 37°C.

Thus, deleting *lasR* from Δ*mvaT*Δ*mvaU* revealed that LasR only increases promoter activity at

stationary phase at 25°C and that LasR does not seem to regulate *piv* at 37°C.

Collectively, these results reveal that *piv* thermoregulation is occurring due to

temperature dependent promoter activity which requires the transcriptional regulator LasR but

not MvaT nor MvaU.

LasR directly regulates *piv*.

To determine how *piv* expression was being thermoregulated by LasR, we first analyzed the *piv*

promoter region more closely. We had found that 200 nucleotides upstream of the *piv* coding

region (Fig. 5A) was sufficient for thermoregulation of its promoter activity. The *piv* upstream

region was recently identified as containing a σ^{70}-dependent promoter (22). Using SAPPHIRE

(30), a predictor for σ^{70} promoters in *Pseudomonas* spp., we identified the putative -35 and -10

elements for the sigma factor RpoD (Fig. 5A, underlined regions). Putative transcription start

sites (TSSs) for *piv* were mapped using 5' rapid amplification of cDNA ends (5' RACE) in PAO1

(Fig. 5A, highlighted bases). We found that transcription began at positions 151C or 152G of the

promoter region sequence. The predicted -35 and -10 elements are appropriately positioned

upstream of our TSSs such that RpoD could be directing transcription of *piv* beginning at 151C

or 152G of the promoter.

We then analyzed the region upstream of the TSSs and predicted σ^{70} -35 and -10

elements and identified a 16 bp sequence (Fig. 5A, boxed region) that resembles a canonical

las box, particularly the one identified for non-cooperative LasR binding (27). We hypothesized

that LasR could be directly regulating *piv* by binding to this region. To test this, we introduced

point mutations to the putative *las* box in the P_{piv}-*gfp*(ASV) reporter plasmid (pRD91) at highly

conserved positions that we predict would either disrupt (Fig. 5B, pink base changes) or

improve (Fig. 5B, green base changes) potential LasR binding. The predicted deleterious

mutations were among ones that have been previously shown to abolish LasR binding to and

greatly reduce activity of the *pqsR* promoter (31, 32). We then assayed the effects of these

mutations on *piv* promoter activity at 25°C and 37°C in PAO1 (Fig. 5C). The predicted

deleterious 1C→A and 14T→A mutations of the putative *las* box reduced *piv* promoter activity

slightly at exponential phase and greatly at stationary phase at 25°C only. Since these

mutations greatly reduced the effect of growth phase on *piv* promoter activity only at 25°C,

these results are consistent with the role of LasR as a quorum sensing regulator that is more

active at stationary phase. The 16T→G mutation, predicted to improve LasR binding, increased

promoter activity at both 25°C and 37°C; however, promoter activity was still highly

thermoregulated. These results are consistent with our prediction for the location of a *las* box in

the *piv* promoter.

DISCUSSION

Here, we found that *piv* thermoregulation occurs at the level of transcriptional regulation and

requires the quorum sensing regulator LasR but not the H-NS family members MvaT/MvaU. By

leveraging a Δ*mvaT*Δ*mvaU* mutant strain in which *piv* expression was depressed, we show that

LasR acts as a positive regulator of *piv* at 25°C with little to no effect on *piv* expression at 37°C.

This is consistent with our findings that *piv* expression is significantly thermoregulated at

stationary phase but not at exponential phase, as LasR abundance and activity is higher at

stationary phase (25). We also identify a putative *las* box in the *piv* promoter and provide

evidence that LasR directly regulates *piv*.

The role of *piv* and its thermoregulation in *P. aeruginosa* virulence

Thermoregulation of *piv* may underpin the importance of this protease for infections of surface

exposed body sites, such as the eyes or skin. Corneal infections by *P. aeruginosa* can lead to

permanent loss of vision and death as seen in a recent outbreak caused by contamination of

artificial tear drops (16). PIV causes epithelial erosion in the cornea and contributes to bacterial

virulence, and *P. aeruginosa* lacking PIV is practically avirulent in corneal models of infection

(12–15). PIV also severely impaired wound healing and contributed to elevated levels of

inflammation in a wound model of infection in mice (33, 34). At temperatures lower than 37°C,

which could be experienced by the bacteria in surface exposed infections, many virulence

factors such as LasB and type III secretion are downregulated, while PIV is upregulated (6, 7).

The unique thermoregulation of PIV may in part contribute to the severity of corneal infections

and to the success of *P. aeruginosa* as an ocular pathogen, as well as the ability of *P.*

aeruginosa to cause chronic infections of wounds.

Thermoregulation of *piv* has impacted its study.

Many studies of PIV processing, secretion, and post-secretion activation have utilized *in vitro*

experiments or expressed PIV in heterologous hosts, which has led to some inconsistencies in

the model for PIV post-secretion activation. One study in PAO1 showed that the propeptide

could only be degraded by protease LasB in *in vitro* assays (although the ability of PIV to

degrade its own propeptide was not tested) and proposed that the PIV proenzyme is secreted

into supernatant and then processed into the mature form by LasB (18). However, earlier

studies found that PIV purified from the supernatant of PA103-29, a derivative of PA103 that

lacks LasB, was enzymatically active, as was PIV when expressed in *Pseudomonas putida* and

Escherichia coli (14, 17, 35). In both heterologous hosts, mature PIV was found in cell

supernatants, indicating that LasB is not required for the removal of the propeptide from the

proenzyme and suggesting that this maturation step may occur autocatalytically. Mutants of PIV

that lacked catalytic activity could not be processed from the proenzyme into the mature

protease, further supporting an autocatalytic cleavage mechanism of the propeptide (9). We

found significant amounts of mature PIV inside cells of PAO1 at 25°C and 30°C (Fig. 2), as well

as low levels of precursor forms, which supports a path for the maturation of PIV inside cells

and not just in the supernatant. As LasB is inactive in the cytoplasm of *P. aeruginosa*, we

believe our results show that PIV can mature autocatalytically in PAO1.

Prior studies on the secretion and protease activity of PIV have been hampered due to

the low amount of PIV produced from the native chromosomal locus (8, 17). We postulate that

the low level of PIV observed by others was due to thermoregulation of *piv* expression, as the

studies were all conducted at 37°C. Consistent with this, we detected abundant PIV in both

mature and precursor(s) forms at 25°C and 30°C but very little mature and no precursor PIV at

37°C or higher (Fig. 2). Our findings demonstrate that PIV produced from its native locus can be

easily detected in cells grown at 25°C and propose this as a viable alternative to yield high *piv*

expression and/or high PIV protein levels for further studies of PIV secretion and maturation.

A model for thermoregulation of *piv* by LasR

We propose a model for *piv* thermoregulation in which LasR activates *piv* transcription at

stationary phase by interacting with a newly identified *las* box in the *piv* promoter at 25°C but not

at all, or very lowly, at 37°C and higher (Fig. 6). In wild-type cells growing at 25°C, LasR

increases *piv* promoter activity while MvaT and MvaU appear to have no regulatory effect. Given

that the loss of *lasR* results in no detectable promoter activity at 25°C and that further deletion of

mvaT and *mvaU* from Δ*lasR* restores promoter activity (Fig. 4), we suspect that MvaT/MvaU do

not repress at 25°C due to LasR interfering with MvaT/MvaU interacting with the *piv* promoter. In

a wild-type cell growing at 37°C, *piv* promoter activity is barely detectable and LasR appears to

have no regulatory effect on *piv* (Fig. 4). MvaT/MvaU repress *piv* at 37°C, which we suggest is

due to the lack of LasR interference at the promoter. And while PvdS is a known regulator of *piv*

in low iron conditions, since our experiments were all conducted in iron replete LB media in

which PvdS is inactive, *piv* thermoregulation must not require PvdS.

It remains unclear how temperature is modulating LasR regulation of *piv*. In agreement with

prior studies (36), we determined that LasR expression and protein levels are not temperature

dependent (Fig. 3) and thus the amount of LasR is not driving higher *piv* expression at lower

temperatures. Additionally, although translation of LasI is slightly higher at 37°C than 30°C due

to an RNA thermometer, this has been shown to only result in a small increase of 3O-C12-HSL

levels that had no effect on expression of the LasR-regulated *lasB* or the amount of the gene

product elastase (36). Given this and that *piv* expression is higher at ambient temperatures of

25°C-30°C than 37°C (the opposite of how LasI is thermoregulated), we do not believe that the

general activity of the LasRI quorum system itself is thermoregulated nor responsible for

thermoregulating *piv*. A second quorum sensing system in *P. aeruginosa*, the RhlRI system, is

thermoregulated due to an RNA thermometer that results in higher RhlR levels at 37°C. The

thermoregulation of the transcriptional activator RhlR does result in higher levels of RhlR-

regulated virulence factors, like rhamnolipids, at 37°C (36); however, *piv* is not regulated by the

RhlRI quorum sensing system, and furthermore is upregulated at temperatures lower than 37°C,

indicating that RhlRI is not responsible for *piv* thermoregulation (25).

We postulate that thermoregulation of *piv* by LasR involves an interaction between LasR

and specifically the *piv* promoter that is temperature sensitive. Lower temperatures such as

25°C may favor LasR interacting with the *piv* promoter at the identified *las* box to mediate *piv*

thermoregulation. A previous study that identified promoters directly bound by LasR in PAO1

failed to identify the *piv* promoter as one, despite identifying several known and novel LasR

targets (27). In retrospect, this was likely because the experiment in this previous study was

conducted at 37°C (27), a temperature at which our findings show that LasR does not regulate

piv. We find a mutation designed to 'improve' the *las* box sequence within the *piv* promoter

increases promoter activity at 37°C to detectable, albeit still low, levels (Fig. 5). This further

supports that LasR directly regulates *piv*, but only under certain temperature conditions, which

could explain why prior experiments conducted at 37°C failed to identify *piv* as a directly LasR-

regulated gene. To the best of our knowledge, this is the only gene known to be directly

regulated by LasR in a temperature-dependent manner.

It is interesting that mutating the *las* box sequence toward consensus at highly conserved

positions did not result in equal promoter activity at 25°C and 37°C (Fig 5). Thermoregulation of

piv by LasR must not be solely due to the sequence of the *las* box. It is notable that the position

of the *las* box is farther upstream of the *piv* TSSs we identified compared to the position of the

las box in some other LasR bound promoters (7). The *piv* promoter is also unusually AT rich at

~59% compared to the *P. aeruginosa* PAO1 genomic average of ~33%. Genomic DNA that is

high in AT content is more flexible and has been documented to adopt temperature sensitive

secondary structures as "DNA thermometers" that can mediate temperature-dependent

transcription factor binding and subsequent gene expression (1). The position of the putative *las*

box within the *piv* promoter could be accessible to LasR at 25°C, but not 37°C, due to

temperature-dependent structural changes of the promoter itself.

As previously discussed, LasR regulated genes have been extensively studied as part of the

quorum sensing regulon in *P. aeruginosa* (23–25, 37). However, *piv* had not been identified as

directly regulated by LasR prior to this study, likely due to low or nonexistent direct LasR

regulation of *piv* under the standard laboratory growth temperature of 37°C. This raises the

possibility that LasR activates transcription of other unknown genes only at specific conditions,

included but not limited to non-optimal growth temperatures. Studying gene expression under

non-standard laboratory conditions that mimic various other infection environments may expand

our understanding of virulence factor regulation and quorum sensing in the diverse

environments that *P. aeruginosa* can survive in.

ACKNOWLEDGEMENTS

We thank Drs. Simon Dove and Michael Gebhardt for providing key Δ*mvaT*Δ*mvaU* and parent

PAO1 strains, as well as discussions regarding MvaT/MvaU and technical advice on quantitative

immunoblotting. The αLasR antibodies were kindly gifted by Drs. Ajai Dandekar and Pete

Greenberg. We also thank the Goldbeg Lab for discussion and feedback on the manuscript.

This work was supported by NIH grant EY028320 (to JBG). RED was supported by a NIH NRSA

Pre-Doctoral Fellowship (F31 AI172335); RED and SMP were supported by the NIH funded

Antimicrobial Resistance and Therapeutic Discovery Training Program at Emory University

(AI106699). JKR was supported by the Emory Summer Undergraduate Research Experience

(SURE) Program.

MATERIALS AND METHODS

Bacterial strains and plasmids

Bacterial strains, plasmids, and oligonucleotides are listed in the Supplemental Material, Tables

S1-S3. Standard molecular biology practices were used to construct plasmids and all

constructed plasmids were confirmed by sequencing. Mutant strains were confirmed by PCR

and sequencing. Full plasmid and strain construction details are available in the Supplemental

Material.

Culture conditions

Overnight cultures of *P. aeruginosa* were routinely grown in 3 mL of lysogeny broth (LB) at 37°C

in a rolling drum. As appropriate, *E. coli* growth media was supplemented with gentamicin (15

μg/mL), carbenicillin (100 μg/mL), or tetracycline (10 μg/mL) and *P. aeruginosa* media with

gentamicin (60 μg/mL), carbenicillin (300 μg/mL), or tetracycline (100 μg/mL), unless otherwise

noted.

RNA extraction and DNase treatment

To measure transcript levels at different growth temperatures, biological triplicates of PAO1

grown overnight at 37°C were subcultured to an initial OD_{600} of 0.05 in 25 mL LB and incubated

shaking at 200 rpm at the indicated temperature. At exponential phase (OD_{600} ~0.5), 10^9 cells

were centrifuged and resuspended in 1 mL Tri-Reagent (Millipore Sigma) and RNA extracted

according to the manufacturer's instructions, using chloroform. The cultures were allowed to

continue growing until early stationary phase (OD_{600} ~2.0) and RNA extracted again as

described. 5 μg of extracted RNA was DNase treated with TURBO DNase (ThermoFisher)

according to the manufacturer's instructions for rigorous DNase treatment.

Real time quantitative PCR (RT-qPCR)

One-step RT-qPCR was performed on DNase treated RNA using Power SYBR™ Green RNA-

to-CT™ 1-Step Kit (Applied Biosystems) on a LightCycler 96 (Roche) using LightCycler

software v1.1.0.1320. Reactions were conducted in technical triplicate per each biological

sample and the C_q values were averaged for gene expression analysis using the temperature

insensitive *omlA* (6) as an internal control gene. The relative expression of a gene at a given

temperature $X°C$ compared to 37°C was calculated and plotted as $\log_2(2^{-\Delta\Delta Ct})$, where $\Delta\Delta Ct = (C_q$

gene at $X°C$ - C_q *omlA* at $X°C$) – (C_q gene at 37°C – C_q *omlA* at 37°C).

Immunoblotting

Overnight biological triplicates were subcultured to an initial OD_{600} of 0.05 in 25 mL LB and

incubated shaking at 200 rpm at the indicated temperature. At early stationary (OD_{600} ~2.0)

phase, 0.5 mL was collected and resuspended in 475 μL Laemlli buffer with 25 μL β-

mercaptoethanol and boiled for 15 minutes. 10 μL of prepared cell lysate sample was

electrophoresed, transferred to a low-fluorescence PVDF membrane (Bio-Rad), and blocked for

1 hour in Intercept Blocking Buffer (Li-Cor). Primary antibodies against the VSV-G epitope tag

(Sigma, batch # 0000143676) were used at 1:5,000 overnight at 4°C and primary antibodies

against LasR (25) used at 1:1,000 overnight at 4°C with fluorophore conjugated secondary

antibodies against rabbits used at 1:10,000 (Li-Cor, Lot No. D20803-09). Primary antibodies

against the α-subunit of RNA polymerase were used at 1:5,000 (BioLegend, Lot No. B376827)

with fluorophore conjugated antibodies against mice (Li-Cor, Lot No. D20601-01) used at

1:10,000. Membranes were imaged on a BioRad ChemiDoc Imager. Primary antibodies against

LasR were validated by immunoblotting cell lysate from PAO1 and PAO1 $\Delta lasR$ (Fig. S2).

Densitometry analysis was conducted on raw image files using ImageJ.

Transcriptional reporter assays

Biological triplicates of strains with a transcriptional reporter plasmid were grown overnight at

37$°$C and subcultured to an initial OD_{600} of 0.1 in 25 mL LB supplemented with gentamicin 30

μg/mL and incubated shaking at 200 rpm at each 25$°$C and 37$°$C. At exponential phase (OD_{600}

~0.5), 10^9 cells were centrifuged at 10,000 rpm for 2 minutes, supernatant removed, and the

pellet resuspended in 1 mL PBS. 200 μL of each sample was then added to a black sided 96

well plate in technical triplicate and the fluorescence measured (excitation 485 nm, emission

515 nm) on a Synergy H1 plate reader (BioTek). At stationary phase (OD_{600} ~2.0), 10^9 cells were

sampled from the same cultures and fluorescence measured as described. The RFU of 200 μL

PBS on each plate was subtracted from technical replicates, which were then averaged per

biological replicate. Background was determined by averaging all values from a transcriptional

reporter assay of PAO1 carrying pRD87, a no-promoter control of the reporter plasmid.

5' rapid amplification of cDNA ends (5' RACE)

PAO1 was grown at 25$°$C in LB to late log phase (OD_{600} ~0.8), RNA extracted using MasterPure

RNA Purification Kit (Epicentre), and treated with TURBO DNase as described. 5' RACE was

conducted using the 5' RACE System for Rapid Amplification of cDNA Ends (ThermoFisher)

according to manufacturer's instructions with modifications: ThermoScript Reverse

Transcriptase (Invitrogen) was used in place of the kit's reverse transcriptase with betaine

added to 0.5 M for the reverse transcriptase step.

Statistical analyses

All statistical tests were conducted as described using Prism version 10.

REFRENCES

1. Shapiro RS, Cowen LE. 2012. Thermal Control of Microbial Development and Virulence:

Molecular Mechanisms of Microbial Temperature Sensing. mBio 3.

2. Grosso-Becera MV, Servín-González L, Soberón-Chávez G. 2015. RNA structures are

involved in the thermoregulation of bacterial virulence-associated traits. Trends Microbiol

23:509–518.

3. Almblad H, Randall TE, Liu F, Leblanc K, Groves RA, Kittichotirat W, Winsor GL, Fournier

N, Au E, Groizeleau J, Rich JD, Lou Y, Granton E, Jennings LK, Singletary LA, Winstone

TML, Good NM, Bumgarner RE, Hynes MF, Singh M, Stietz MS, Brinkman FSL, Kumar A,

Brassinga AKC, Parsek MR, Tseng BS, Lewis IA, Yipp BG, MacCallum JL, Harrison JJ.

2021. Bacterial cyclic diguanylate signaling networks sense temperature. Nat Commun

12:1986.

4. Barbier M, Oliver A, Rao J, Hanna SL, Goldberg JB, Albertí S. 2008. Novel

Phosphorylcholine-Containing Protein of Pseudomonas aeruginosa Chronic Infection

Isolates Interacts with Airway Epithelial Cells. J Infect Dis 197:465–473.

5. Owings JP, Kuiper EG, Prezioso SM, Meisner J, Varga JJ, Zelinskaya N, Dammer EB,

Duong DM, Seyfried NT, Albertí S, Conn GL, Goldberg JB. 2016. Pseudomonas

aeruginosa EftM Is a Thermoregulated Methyltransferase. J Biol Chem 291:3280–3290.

6. Barbier M, Damron FH, Bielecki P, Suárez-Diez M, Puchałka J, Albertí S, Santos VM dos,

Goldberg JB. 2014. From the Environment to the Host: Re-Wiring of the Transcriptome of

Pseudomonas aeruginosa from 22°C to 37°C. PLOS ONE 9:e89941.

7. Wurtzel O, Yoder-Himes DR, Han K, Dandekar AA, Edelheit S, Greenberg EP, Sorek R,

Lory S. 2012. The Single-Nucleotide Resolution Transcriptome of Pseudomonas

aeruginosa Grown in Body Temperature. PLOS Pathog 8:e1002945.

8. Engel LS, Hill JM, Caballero AR, Green LC, O'Callaghan RJ. 1998. Protease IV, a unique

extracellular protease and virulence factor from Pseudomonas aeruginosa. J Biol Chem

273:16792–16797.

9. Traidej M, Marquart ME, Caballero AR, Thibodeaux BA, O'Callaghan RJ. 2003.

Identification of the active site residues of Pseudomonas aeruginosa protease IV.

Importance of enzyme activity in autoprocessing and activation. J Biol Chem 278:2549–

2553.

10. Malloy JL, Veldhuizen RAW, Thibodeaux BA, O'Callaghan RJ, Wright JR. 2005.

Pseudomonas aeruginosa protease IV degrades surfactant proteins and inhibits surfactant

host defense and biophysical functions. Am J Physiol Lung Cell Mol Physiol 288:L409-418.

11. Guillon A, Brea D, Morello E, Tang A, Jouan Y, Ramphal R, Korkmaz B, Perez-Cruz M,

Trottein F, O'Callaghan RJ, Gosset P, Si-Tahar M. 2016. Pseudomonas aeruginosa

proteolytically alters the interleukin 22-dependent lung mucosal defense. Virulence 8:810–

820.

12. Engel LS, Hobden JA, Moreau JM, Callegan MC, Hill JM, O'Callaghan RJ. 1997.

Pseudomonas deficient in protease IV has significantly reduced corneal virulence. Invest

Ophthalmol Vis Sci 38:1535–1542.

13. Engel LS, Hill JM, Moreau JM, Green LC, Hobden JA, O'Callaghan RJ. 1998.

Pseudomonas aeruginosa protease IV produces corneal damage and contributes to

bacterial virulence. Invest Ophthalmol Vis Sci 39:662–665.

14.   O'Callaghan RJ, Engel LS, Hobden JA, Callegan MC, Green LC, Hill JM. 1996.

Pseudomonas keratitis. The role of an uncharacterized exoprotein, protease IV, in corneal

virulence. Invest Ophthalmol Vis Sci 37:534–543.

15.   Willcox MDP. 2007. Pseudomonas aeruginosa Infection and Inflammation During Contact

Lens Wear: A Review. Optom Vis Sci 84:273–278.

16.   Grossman MK, Rankin DA, Maloney M, Stanton RA, Gable P, Stevens VA, Ewing T,

Saunders K, Kogut S, Nazarian E, Bhaurla S, Mephors J, Mongillo J, Stonehocker S,

Prignano J, Valencia N, Charles A, McNamara K, Fritsch WA, Ruelle S, Plucinski CA, Sosa

LE, Ostrowsky B, Ham DC, Walters MS, for the Multistate Pseudomonas Outbreak

Investigation Group. 2024. Extensively Drug-Resistant Pseudomonas aeruginosa

Outbreak Associated With Artificial Tears. Clin Infect Dis ciae052.

17.   Traidej M, Caballero AR, Marquart ME, Thibodeaux BA, O'Callaghan RJ. 2003. Molecular

analysis of Pseudomonas aeruginosa protease IV expressed in Pseudomonas putida.

Invest Ophthalmol Vis Sci 44:190–196.

18.   Oh J, Li X-H, Kim S-K, Lee J-H. 2017. Post-secretional activation of Protease IV by

quorum sensing in Pseudomonas aeruginosa. 1. Sci Rep 7:4416.

19.   O'Callaghan R, Caballero A, Tang A, Bierdeman M. 2019. Pseudomonas aeruginosa

Keratitis: Protease IV and PASP as Corneal Virulence Mediators. Microorganisms 7.

20.   Wilderman PJ, Vasil AI, Johnson Z, Wilson MJ, Cunliffe HE, Lamont IL, Vasil ML. 2001.

Characterization of an Endoprotease (PrpL) Encoded by a PvdS-Regulated Gene in

Pseudomonas aeruginosa. Infect Immun 69:5385–5394.

21. Castang S, McManus HR, Turner KH, Dove SL. 2008. H-NS family members function

coordinately in an opportunistic pathogen. Proc Natl Acad Sci U S A 105:18947–18952.

22. Lippa AM, Gebhardt MJ, Dove SL. 2021. H-NS-like proteins in Pseudomonas aeruginosa

coordinately silence intragenic transcription. Mol Microbiol 115:1138–1151.

23. Wagner VE, Bushnell D, Passador L, Brooks AI, Iglewski BH. 2003. Microarray Analysis of

Pseudomonas aeruginosa Quorum-Sensing Regulons: Effects of Growth Phase and

Environment. J Bacteriol 185:2080–2095.

24. Schuster M, Lostroh CP, Ogi T, Greenberg EP. 2003. Identification, timing, and signal

specificity of Pseudomonas aeruginosa quorum-controlled genes: a transcriptome

analysis. J Bacteriol 185:2066–2079.

25. Schuster M, Greenberg EP. 2007. Early activation of quorum sensing in Pseudomonas

aeruginosa reveals the architecture of a complex regulon. BMC Genomics 8:287.

26. Papenfort K, Bassler B. 2016. Quorum-Sensing Signal-Response Systems in Gram-

Negative Bacteria. Nat Rev Microbiol 14:576–588.

27. Gilbert KB, Kim TH, Gupta R, Greenberg EP, Schuster M. 2009. Global position analysis of

the Pseudomonas aeruginosa quorum-sensing transcription factor LasR. Mol Microbiol

73:1072–1085.

28. Andersen JB, Sternberg C, Poulsen LK, Bjørn SP, Givskov M, Molin S. 1998. New

Unstable Variants of Green Fluorescent Protein for Studies of Transient Gene Expression

in Bacteria. Appl Environ Microbiol 64:2240–2246.

29. Rybtke MT, Borlee BR, Murakami K, Irie Y, Hentzer M, Nielsen TE, Givskov M, Parsek MR,

Tolker-Nielsen T. 2012. Fluorescence-Based Reporter for Gauging Cyclic Di-GMP Levels

in Pseudomonas aeruginosa. Appl Environ Microbiol 78:5060–5069.

30. Coppens L, Lavigne R. 2020. SAPPHIRE: a neural network based classifier for σ70

promoter prediction in Pseudomonas. BMC Bioinformatics 21:415.

31. Farrow JM, Pesci EC. 2017. Distal and proximal promoters co-regulate pqsR expression in

Pseudomonas aeruginosa. Mol Microbiol 104:78–91.

32. Farrow JM, Hudson LL, Wells G, Coleman JP, Pesci EC. 2015. CysB Negatively Affects

the Transcription of pqsR and Pseudomonas Quinolone Signal Production in

Pseudomonas aeruginosa. J Bacteriol 197:1988–2002.

33. Prasad ASB, Shruptha P, Prabhu V, Srujan C, Nayak UY, Anuradha CKR, Ramachandra L,

Keerthana P, Joshi MB, Murali TS, Satyamoorthy K. 2020. Pseudomonas aeruginosa

virulence proteins pseudolysin and protease IV impede cutaneous wound healing. Lab

Investig J Tech Methods Pathol 100:1532–1550.

34. Kim T-H, Li X-H, Lee J-H. Alleviation of Pseudomonas aeruginosa Infection by Propeptide-

Mediated Inhibition of Protease IV. Microbiol Spectr 9:e00782-21.

35. Caballero AR, Moreau JM, Engel LS, Marquart ME, Hill JM, O'Callaghan RJ. 2001.

Pseudomonas aeruginosa protease IV enzyme assays and comparison to other

Pseudomonas proteases. Anal Biochem 290:330–337.

36. Grosso-Becerra MV, Croda-García G, Merino E, Servín-González L, Mojica-Espinosa R,

Soberón-Chávez G. 2014. Regulation of Pseudomonas aeruginosa virulence factors by

two novel RNA thermometers. Proc Natl Acad Sci 111:15562–15567.

37.   Whiteley M, Lee KM, Greenberg EP. 1999. Identification of genes controlled by quorum

536   sensing in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A 96:13904–13909.

537   38.   Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic

538   assembly of DNA molecules up to several hundred kilobases. Nat Methods 6:343–345.

539   39.   Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE, Tseng BS, Lin C, Irie Y, Storek

540   KM, Yang JJ, Siehnel RJ, Howell PL, Singh PK, Tolker-Nielsen T, Parsek MR, Schweizer

541   HP, Harrison JJ. 2015. Precision-engineering the Pseudomonas aeruginosa genome with

542   two-step allelic exchange. Nat Protoc 10:1820–1841.

543   40.   Chen W, Zhang Y, Zhang Y, Pi Y, Gu T, Song L, Wang Y, Ji Q. 2018. CRISPR/Cas9-based

544   Genome Editing in Pseudomonas aeruginosa and Cytidine Deaminase-Mediated Base

545   Editing in Pseudomonas Species. iScience 6:222–231.

546   41.   Engler C, Gruetzner R, Kandzia R, Marillonnet S. 2009. Golden Gate Shuffling: A One-Pot

547   DNA Shuffling Method Based on Type IIs Restriction Enzymes. PLOS ONE 4:e5553.

548   42.   Choi K-H, Schweizer HP. 2006. mini-Tn 7 insertion in bacteria with single att Tn 7 sites:

549   example Pseudomonas aeruginosa. Nat Protoc 1:153–161.

A

B

Figure 1. Temperature regulation of *piv* depends on growth phase. A) PAO1 was grown at 25°C, 30°C, 37°C, and 42°C. RNA was extracted from an equal number of cells first at exponential phase and then from the same cultures at early stationary phase, at points during bacterial growth indicated by arrows on the y-axis. The average growth of three biological replicates is shown with error bars representing standard deviation. B) Expression of *piv* at each temperature was calculated relative to 37°C using *omlA* as a temperature insensitive, internal control gene. RT-qPCR was conducted in technical triplicate and the mean of three biological replicates is shown with error bars representing standard deviation. Statistical significance was determined by two-way ANOVA with Dunnett's multiple comparisons: * $p < 0.0332$, ** $p < 0.0021$, *** $p < 0.0002$, **** $p < 0.0001$, ns – not significant. Values between the dotted lines were not considered biologically significant.

Figure 2. Temperature regulation of *piv* expression is reflected by levels of PIV protein. A) Cell lysates of PAO1 PIV VSV-G grown to early stationary phase at 25°C, 30°C, 37°C, and 42°C were probed with antibodies against the VSV-G tag (αVSV-G) and the RNA polymerase α subunit (αRpoA). A representative image of three biological replicates is shown. B) The amount of mature PIV VSV-G protein at each temperature was determined relative to the amount of RpoA using ImageJ. The mean of three biological replicates is shown with standard deviation. Statistical significance was determined by one-way ANOVA with Dunnett's multiple comparisons: * $p < 0.0332$, ** $p < 0.0021$, *** $p < 0.0002$, **** $p < 0.0001$, ns – not significant.

A

B

C

D

Figure 3. Transcriptional regulators of *piv* are not thermoregulated. A) PAO1 was grown at 25°C and 37°C and RNA extracted at exponential and stationary phases as previously described. Expression of genes was determined by RT-qPCR using *omlA* as an internal control gene and calculated for 25°C relative to 37°C. RT-qPCR was conducted in technical triplicate. The mean of three biological replicates analyzed is shown with error bars representing standard deviation. Values between the dotted lines were not considered biologically significant. B,C,D) Western blots (left) and subsequent quantification (right) of indicated transcription factors in cells grown at 25°C and 37°C to stationary phase. B) PAO1 was probed with αLasR antibodies. C) PAO1 MvaT VSV-G was probed with αVSV-G antibodies. D) PAO1 MvaU VSV-G was probed with αVSV-G antibodies. All samples were probed with αRpoA as a loading control. Biological triplicates for each protein are shown. For densitometry analysis, the amount of LasR (B), MvaT VSV-G (C), and MvaU VSV-G (D) protein was determined relative to the amount of RpoA in the same sample using ImageJ. The mean of three biological replicates is shown with standard deviation. Statistical significance was determined by two-tailed t test, ns – not significant.

Figure 4. Temperature alters activity of the *piv* promoter through transcriptional regulator LasR, but not MvaT/MvaU. The indicated strains carrying the P*piv*-*gfp*(ASV) transcriptional reporter (pRD91) were grown at 25°C and 37°C. An equal number of cells were sampled for fluorescence (RFU) measurements (excitation 485 nm, emission 515 nm) first at exponential phase (A) and then at early stationary phase (B). The dotted line indicates background fluorescence of PAO1 carrying a no-promoter *gfp*(ASV) reporter plasmid (pRD87). Three biological replicates were performed, with RFU measurements conducted in technical triplicate per each. The mean of three biological replicates is shown with error bars representing standard deviation. Statistical significance was determined by two-way ANOVA with Šídák multiple comparisons: * $p < 0.0332$, ** $p < 0.0021$, *** $p < 0.0002$, **** $p < 0.0001$, ns – not significant. Statistical comparisons shown are for a given strain grown at 25°C versus 37°C.

A

1
TGGTAGAGAGAGCAATCCAACATCAATGGCAAGTGAAGGA

41
AATAGCTATCTATTCTGCTATCCACAAACGCTTCTTCATA
las box

81
TTTAATAGCTAGCAGAAACGAATATAAAAAATAT<u>TTGAAC</u>
σ70 −35

121
GACTCCGTCACACCTGCA<u>TATCTTT</u>CGCCA**CG**CGAACGAT
σ70 −10

161
TGAGGAAGTTGCCCTCCCAAAAAACTGAAGGAGTCGATTC

B

consensus *las* box: CTAnnnnnnnnnnTAG

piv las box: CTATCTATTCTGCTAT

1C→A: **A**TATCTATTCTGCTAT

14T→A: CTATCTATTCTGC**A**AT

16T→G: CTATCTATTCTGCTA**G**

C

Figure 5. Mutations to a putative *las* box in the *piv* promoter region alter promoter activity.
A) Sequence of the 200 nucleotides upstream of the *piv* start codon that are sufficient for temperature and growth phase regulation. Putative σ70 -35 and -10 elements identified with

SAPPHIRE (30) are underlined. Nearby transcription start sites (TSSs) were identified by 5' RACE are highlighted. A DNA motif with similarity to canonical *las* boxes (27) was identified upstream of transcription initiation regions and is boxed. B) Mutations made to the *las* box within the *piv* promoter of the P$_{piv}$-*gfp*(ASV) transcriptional reporter plasmid (pRD91). Mutations are predicted to disrupt (in pink) or improve (in green) LasR binding based on comparison to the consensus sequence. C) PAO1 carrying *las* box mutants of the P$_{piv}$-*gfp*(ASV) transcriptional reporter shown in B) were grown at 25°C and 37°C and fluorescence measurements taken at exponential (left) and stationary (right) phases as previously described. The mean of three biological replicates analyzed with standard deviation is shown. Statistical significance was determined two-way ANOVA with Šídák multiple comparisons: * $p < 0.0332$, ** $p < 0.0021$, *** $p < 0.0002$, **** $p < 0.0001$.

 Figure 6. The proposed model for the transcriptional thermoregulation of *piv* by the quorum sensing regulator LasR. At 25°C (blue thermometer), LasR drives high expression of *piv* at stationary phase by directly interacting with the *piv* promoter while MvaT/MvaU do not repress expression. At 37°C (red thermometer), LasR no longer directly regulates *piv*, which allows MvaT/MvaU to repress *piv* expression.