<u>Crispant analysis in zebrafish as a tool for rapid functional screening of</u>
<u>disease-causing genes for bone fragility.</u>

Sophie Debaenst[1]; Tamara Jarayseh[1]; Hanna de Saffel[1]; Jan Willem Bek[1]; Matthieu Boone[2],
Ivan Josipovic[2], Pierre Kibleur[2], Ronald Young Kwon[3,4], Paul J. Coucke[1]; Andy Willaert[1].

1. Center for Medical Genetics Ghent, Department of Biomolecular Medicine, Ghent
University, Ghent, Belgium
2. Center for X-ray Tomography, Department of Physics and Astronomy, Ghent University,
Gent, Belgium
3. Department of Orthopaedics and Sports Medicine, University of Washington, Seattle, United
States
4. Institute for Stem Cell and Regenerative Medicine, University of Washington, Seattle, United
States

Abstract

Heritable Fragile Bone Disorders (FBDs) encompass a spectrum of conditions, from

widespread multifactorial to rare monogenic diseases, all characterized by an elevated risk of

fractures. The process of validating causative genes and elucidating their pathogenic

mechanisms remains a daunting and resource-intensive task. In this study, we evaluated the

feasibility of a semi-high throughput zebrafish screening platform for rapid validation and *in*

vivo functional testing and validation of candidate disease-causing genes for a wide range of

heritable FBDs. Six genes associated with severe recessive forms of Osteogenesis Imperfecta

(OI) and four genes associated with BMD, a key osteoporosis indicator, identified through

genome-wide association studies (GWAS) were selected. The crispant screening approach,

based on CRISPR/Cas9 technology, was used to phenotype directly in F0 mosaic founder

zebrafish. Next-Generation Sequencing (NGS) analysis revealed a mean indel efficiency of

88% across ten different crispants, indicating a high proportion of knock-out alleles and thus

resembling stable knock-out models. We applied multiple techniques to evaluate skeletal

characteristics at 7, 14 and 90 days post-fertilization (dpf), including microscopy for osteoblast

reporter visualization and mineralization by Alizarin Red S staining, and microCT for

quantitative skeletal analysis. While larval crispants exhibited variable differences in

osteoblast-positive and mineralized surface areas, adult-stage crispants displayed more

pronounced and consistent skeletal phenotypes. Notably, all crispants developed malformed

neural and haemal arches, with a majority presenting vertebral fractures and fusions, and some

showing significant alterations in vertebral bone volume and density. In addition, *aldh7a1* and

mbtps2 crispants experienced increased mortality due to severe skeletal deformities. RT-qPCR

analysis of osteoblast differentiation and bone formation markers at larval stages indicated

differential expression of osteogenic markers *bglap* and *col1a1a* in a substantial portion of the

crispants, hinting at their utility as biomarkers for FBD crispant screening. In summary, our

findings demonstrate that crispant screening in zebrafish offers a viable and efficient strategy

for the functional assessment of FBD genes. We advocate for a comprehensive approach that

integrates various techniques and evaluates distinct skeletal and molecular profiles across

different developmental and adult stages. This methodology has the potential to provide new

insights into the role of these genes in skeletal biology.

Key words: Osteogenesis Imperfecta, osteoporosis, candidate genes, zebrafish model, crispant,

skeletal disorders

Introduction

Heritable fragile bone disorders (FBD), which are generally associated with increased fracture risk, range from common multifactorial conditions to rare monogenic diseases (Charoenngam *et al.*, 2023). They present within a wide phenotypic spectrum where monogenic disorders are generally more severe and often include skeletal deformities and extra-skeletal abnormalities. Multifactorial FBD, on the other hand, usually present with milder symptoms that are often restricted to bone fragility. Osteoporosis is a common skeletal disease characterized by low bone mass, low bone mineral density (BMD) and increased fracture risk (Kanis *et al.*, 1994). It affects about 200 million people around the world (Pouresmaeili *et al.*, 2018). In Europe, 5.6% of the population aged over 50 is diagnosed with osteoporosis, making it a highly prevalent disease, especially in the elderly. The risk of osteoporosis is influenced by both environmental factors, such as physical activity, diet preferences, medication use, coexisting diseases, and genetic factors (Ralston and Uitterlinden, 2010). Low BMD and fractures are highly heritable risk factors, with heritability values ranging from 50% to 80% (Brent Richards *et al.*, 2012). Genome-wide association studies (GWAS) in humans have unraveled hundreds of susceptible loci associated with BMD and other bone traits related to risk of osteoporotic fracture (Zhu *et al.*, 2021). GWAS results have great potential for the development of drug targets, the identification of risk factors, and risk prediction for osteoporosis. However, the identification of causal genes at these loci or the mechanisms by which GWAS loci alter bone physiology has faced challenges due to the scarcity of *in vivo* validation tools with sufficient throughput (Sabik and Farber, 2017).

In monogenic FBD, on the other hand, gene identification is generally more straightforward. One example of such a disease is osteogenesis imperfecta (OI), a genetically heterogeneous connective tissue disorder that is mainly characterized by bone fragility, skeletal deformities, and other extra skeletal manifestations (Forlino and Marini, 2016). Although in the majority of cases, it is caused by autosomal dominant mutations in the collagen type 1 genes, mutations in an increasing number of non-collagen genes causing severe recessive forms of OI, are being identified. These genes encode proteins that are involved in different aspects of bone formation, such as osteoblast differentiation (OSTERIX, WNT1, OASIS, S2P and SPARC), bone mineralization (PEDF and BRIL) and collagen processing (HSP47, FKBP65, LH2, KDELR2, CRTAP, P3H1, CyPB and TRIC-B) (Forlino and Marini, 2016)(Jovanovic *et al.*, 2022). Recent sequencing technologies have improved genetic diagnosis, but the investigation of the pathogenesis of the different forms of OI is still a laborious, costly, and time-consuming

task. Hence, it is essential to develop efficient and cost-effective methods for *in vivo* functional
testing to validate the involvement and mechanistic role of candidate and newly identified
causal genes in the pathogenesis of FBD.

Zebrafish are a valuable model organism for studying human diseases, as they share a
high degree of genetic similarity with humans. More than 70% of human protein-coding genes
and 82% of human disease-related genes have zebrafish orthologs, including most of the genes
involved in skeletogenesis (Howe *et al.*, 2013). Additionally, zebrafish share many of their
skeletal features and ossification mechanisms with mammals (Dietrich *et al.*, 2021) and
numerous zebrafish models for skeletal diseases have been established and investigated
(Gistelinck *et al.*, 2018)(Daponte *et al.*, 2023)(Kague and Karasik, 2022). These models have
some advantages over rodent models, such as longer survival, likely due to the reduced impact
of gravity on their fragile skeletons in the aquatic environment, and the availability of transgenic
lines marking specific bone cell types combined with the transparency of early life stages
allowing for *in vivo* observation of the dynamics of skeletal development (Dietrich *et al.*, 2021).

Genome-editing with CRISPR/Cas9 is a fast and easy method to create knockout alleles
and stable zebrafish mutant lines (Uribe-Salazar *et al.*, 2022). However, obtaining zebrafish
lines that are homozygous for the mutant alleles requires two generations of adult animals (>6
months), which limits the efficiency and affordability of genetic screens (Kroll *et al.*, 2021). To
overcome these drawbacks, phenotyping the first-generation (F0) mosaic founder zebrafish,
also known as crispants, is an emerging approach. Adult crispants have a shorter generation
time (~3 months) than stable lines (6 to 9 months), which allows screening more genes in less
time and at lower costs and resources. We previously demonstrated the feasibility of this
approach by comparing the phenotypes and molecular profiles of a stable knockout line and a
crispant for the osteoporosis gene *lrp5*, and found them to be highly similar (Bek *et al.*, 2021).
In a study by Watson et al., 2020, comparison of crispants to homozygous germline mutants for
bmp1a and *plod2*, two causal OI genes, revealed phenotypic convergence, suggesting that
crispants faithfully recapitulate the biology of germline skeletal disease models. Therefore,
while zebrafish crispant screens are promising as an economic, fast, and biologically relevant
genetic screening approach in vertebrate model systems, a rigorous evaluation of their potential
to screen for FBD-associated genes is needed.

In this study, we aimed to conduct a pilot-study to evaluate the feasibility of zebrafish
crispant screening as an approach for rapid *in vivo* functional screening and validation of a set
of causal genes for a broad range of FBD. The selected set includes on one hand genes
associated with traits related to osteoporosis which is a relatively mild late-onset FBD and on

the other hand genes causing recessive forms of OI associated with more severe skeletal

phenotypes. With this study, we want to provide an experimental platform for medium-sized

zebrafish crispant screening of FBD genes.

Results

Generation of zebrafish crispants for osteoporosis and OI genes

To identify and prioritize potential causal genes for osteoporosis, we used the GWAS catalog (http://www.ebi.ac.uk/gwas/). We selected the variants associated with bone density and ranked them by their GWAS evidence, i.e., the number of independent publications that reported significant associations. We then identified the genes that harbored these variants and chose the top four candidates with single zebrafish orthologs: *ALDH7A1*, *ESR1*, *DAAM2* and *SOST*. Furthermore, we selected six genes associated with recessive forms of OI associated with severe skeletal phenotypes, and involved in different aspects of bone formation, including osteoblast differentiation (*CREB3L1, MBTPS2 and SPARC),* bone mineralization (*SERPINF1* and *IFITM*) and collagen transport (*SEC24D*).

For each of the 10 selected genes and "scrambled" sequence, Alt-R™ gRNAs (IDT) were designed through a thorough procedure, via the Benchling platform. For each gene, the gRNA with the highest out-of-frame (OOF) efficiency, as predicted by the InDelphi-mESC prediction tool (Supplementary Figure 1), was selected for co-injection with Cas9 into one-cell stage embryos. Based on previous studies, we assumed that high indel (insertion/deletion) and OOF efficiencies result in a sufficient reduction or inactivation of the corresponding protein in crispants to induce a skeletal phenotype (Naert *et al.*, 2020). DNA from a pool of 1 dpf larvae (n=10) for the different crispants and corresponding sibling controls, was extracted, and subjected to NGS analysis. Sequencing data was analyzed using the Crispresso2 tool (Clement *et al.*, 2019) to determine the fraction of reads containing indels, the OOF rate and the number of in-frame deletions consisting of more than 6 base pairs in these pools (Table 1).

Across the different crispants the minimal indel efficiency was 71%, while the OOF rate ranged between 49 and 73%. We subsequently determined the proportion of reads containing in-frame deletions of at least 6 base pairs, resulting in the removal of 2 or more amino acids (Table 1; Supplementary Figure 1). Such deletions have a relatively high chance of impairing protein function (Emond *et al.*, 2020)(Savino *et al.*, 2022)(Miton and Tokuriki, 2023). The accumulated percentage of reads with indels that likely cause loss-of-function of the respective protein (OOF % + % in-frame indel ≥ 6bp), fall within the range of 60% to 92%, although also reads with short in-frame indels (<6bp) can affect protein function. Finally, we investigated the occurrence of off-target effects in these samples by analyzing NGS data from the three most probable off-target sites per crispant, as determined by the CFD score provided by the CRISPOR tool (Concordet and Haeussler, 2018). In 20 out of the 30 examined off-target sites,

201   no off-target mutations were identified (Supplementary Table 1). For the remaining off-target
202   sites, the off-target rate ranged from 0,11% to 2,82%. This included basepair substitutions
203   which are most likely sequencing errors, or true off-target indel variants, present in such low
204   number of genomes that it is negligible in terms of biological impact in the crispants.

Assessment of the early skeleton in crispants reveals differences in osteoblast-positive and mineralized surface areas.

Crispants and sibling controls were raised until 7 or 14 dpf. At this point the presence
of immature osteoblasts in the head region was visualized using the transgenic *osx:Kaede*
background. Images from 10 larvae per crispant were captured from the ventral perspective at
7dpf and from both ventral and lateral perspectives at 14dpf, as this provides a more
comprehensive view of the operculum's surface area that can be clearly visualized at that time.
The osteoblast-positive areas in both the total head and the operculum were then quantified to
gain insight into the formation of new skeletal tissue during early development (Figure 1a;
Supplementary Figure 2, 4).

At 7 and 14 dpf, *esr1* crispants exhibited a significant increase of the osteoblasts-
positive surface area within the head ($p \leq 0,05$; $p \leq 0.001$) (Figure 1b, 1c). Additionally, *ifitm5*
crispants at 14 dpf also displayed a significant increase ($p \leq 0.01$) in osteoblasts-positive surface
area within the head. Finally, the increased osteoblasts-positive surface area was also observed
in the operculum of both crispants at 14 dpf (*esr1* $p \leq 0.001$; *ifitm5* $p \leq 0.01$) and in the
operculum of *serpinf1* ($p \leq 0.01$) and *sparc* ($p \leq 0.05$) crispants.

Subsequently, Alizarin Red S (ARS) staining was conducted on the same 7 and 14 dpf
crispant zebrafish larvae in order to evaluate the degree of mineralization in the early skeletal
structures. Ventral views were captured at 7 dpf, while both ventral and lateral views were
obtained at 14 dpf. These lateral images at 14 dpf served a dual purpose: they facilitated
measurement of the mineralized surface area of the operculum and allowed for quantification
of the number of mineralized vertebrae. We quantified the mineralized surface areas of the total
head, the notochord tip, and the operculum, as well as determined the number of vertebrae
(Figure 1a; Supplementary Figure 3,5).

At 7dpf, the total head exhibited a significant reduction in mineralized surface in
aldh7a1 ($p \leq 0.05$) and *sec24d* crispants ($p \leq 0.0001$), while the notochord tip displayed a
significant increase in mineralized surface area in *creb3l1* ($p \leq 0.05$), *ifitm5* ($p \leq 0.05$), and
serpinf1 ($p \leq 0.05$) crispants (Figure 1b). At 14 dpf, the significantly reduced mineralized
surface area of the total head persisted in *aldh7a1* crispants ($p \leq 0.05$), with a similar trend

observed in *sec24d* crispants, albeit not reaching statistical significance. Furthermore, the increased mineralization of the notochord tip remained significant in *ifitm5* crispants ($p \leq 0.05$), while it normalized in *creb3l1* and *sec24d* crispants. A novel increase emerged in *esr1* crispants ($p \leq 0.01$). Finally, at 14 dpf, *sec24d* crispants exhibited a significant decrease ($p \leq 0.01$) in the mineralized surface area of the operculum and *serpinf1* crispants showed a notable increase in the number of mineralized vertebrae ($p \leq 0.01$) (Figure 1c).

RT-qPCR expression analysis reveals differential expression of osteogenic markers *bglap* and *col1a1a* in a significant subset of crispants.

To explore the expression level of different osteogenic markers throughout the set of crispants, RT-qPCR analysis was performed on whole larvae at 7 and 14 dpf. At 7dpf, no significant differences were seen in the expression of *runx2*, an early pre-osteoblast marker, between the different crispants and corresponding controls. *Osterix,* also known as *sp7*, a *runx2* downstream gene which plays a critical role for the differentiation of pre-osteoblasts into mature osteoblasts*,* was only upregulated in crispants for *sparc* ($p \leq 0{,}001$). Osteocalcin, also known as *bglap,* which is mainly expressed in mature osteoblasts, was downregulated in almost all crispants, except in crispants for *sost* and *sparc*. Finally, *col1a1a,* encoding the α1 chain of type 1 collagen, and marking extracellular matrix synthesis by both pre- and mature osteoblasts (Valenti *et al.*, 2020), was upregulated in crispants for *aldh7a1* ($p \leq 0{,}001$), *daam2* ($p \leq 0{,}05$), *esr1* ($p \leq 0{,}05$) and *sec24d* ($p \leq 0{,}05$) (Figure 2a).

At 14dpf, the expression of *runx2* remained unchanged in all crispants compared to control. The expression of *sp7* was significantly increased in crispants for *daam2* ($p \leq 0{,}05$) and *sec24d* ($p \leq 0{,}05$), decreased in *esr1* crispants, while expression levels were normalized in *sparc* crispants relative to 7 dpf ($p \leq 0{,}05$). The expression of *bglap* remained significantly downregulated in crispants for *aldh7a1* ($p \leq 0{,}001$), *daam2* ($p \leq 0{,}01$), *esr1* ($p \leq 0{,}01$) and *creb3l1* ($p \leq 0{,}05$) while it reverted to control levels in *ifitm5*, *sec24d* and *serpinf1* crispants. An upregulation in the expression of *col1a1a* was seen in all the crispants, except in crispants for sparc and for *aldh7a1* and *esr1* where expression levels normalized compared to 7dpf (Figure 2b).

Assessment of the adult skeleton in FBD crispants reveals frequent morphological abnormalities and quantitative differences in vertebral bone parameters.

Crispants were further grown into adulthood (90dpf) and subjected to whole-mount
ARS staining for mineralized bone, in order to identify skeletal abnormalities. Crispants for
both *aldh7a1* and *mbtps2* displayed early lethality. Survival analysis indicated that *aldh7a1*
crispants experienced increased lethality from 7dpf onwards, with no survivors beyond 20dpf,
while *mbtps2* crispants exhibited increased lethality from 17dpf onwards, with only a very small
number of fish surviving into adulthood (90dpf) (Figure 3). ARS staining of the skeleton of two
surviving adult *mbtps2* crispant fish revealed severe cranial malformations, including
underdevelopment of the premaxilla, dentary, and articular. However, the vertebral column
seemed to be only mildly affected.

Among the eight crispants that successfully matured into adulthood, none exhibited
significant differences in standard length and head size (n=5 fish per crispant) (Figure 4a).
However, *esr1* crispants were observed to have notably larger eye diameters (Figure 4a,c). All
these crispants demonstrat__ .arious abnormalities in the caudal part of the vertebral column
such as fusions, compressions, fractures, or arch malformations, except for *daam2* crispants
where no obvious abnormalities were detected (Figure 4b,c; Supplementary Figure 6). In
creb3l1 crispants the vertebral phenotype was relatively mild and primarily restricted to arch
malformations.

Next, the adult skeleton of the different crispants and sibling controls was visualized
by microCT scanning and the data was quantified and further analyzed using FishCuT software
(Hur *et al.*, 2017). For each of the vertebral centra, the length, TMD, volume, and thickness
were determined and tested for statistical differences between groups using a regression-based
statistical test (Supplementary Figure 7). To quantify the extent of differences between crispants
and their controls, the means of these parameters in the crispants' vertebral column, were
normalized to the standard deviation of the controls, resulting in a z-score [z-score = (mean
crispant – mean control) / SD control] (Gistelinck et al., 2018). A significant increase in centrum
volume was observed in *daam2* crispants compared to the controls ($p < 0.05$, $z = 2.86$) (Figure
4d). Centrum TMD was significantly higher, and to a similar extent, in crispants for *daam2* (p
< 0.05; $z = 1.54$) , *esr1* ($p < 0.05$; $z = 1.62$), and *sost* ($p < 0.05$; $z = 1.41$), three genes likely
involved in osteoporosis pathogenesis. Centrum thickness was increased in crispants for *esr1*
($p < 0.05$; $z = 4.26$), *ifitm5* ($p < 0.01$; $z = 7.13$), and *sec24d* ($p < 0.05$; $z = 5.04$) with the most
pronounced increase observed in *ifitm5* crispants. Centrum length was increased only in *daam2*
crispants ($p < 0.05$; $z = 4.22$).

Discussion

The generation of a diverse array of null alleles through CRISPR/Cas9-induced F0 mosaic (crispant) zebrafish offers specific advantages compared to stable knockout zebrafish lines, in particular for rapid screening of phenotypes (Naert *et al.*, 2020). In this study we tested the efficacy of a semi-high throughput zebrafish crispant analysis for rapid validation of candidate disease causing genes for skeletal disorders across a broad phenotypic spectrum. On one hand we included genes for non-classical recessive OI, which is an early-onset monogenic skeletal syndrome with severe skeletal deformities. On the other hand, we screened a set of top candidate genes for osteoporosis, which is a rather late-onset multifactorial disease with little to no deformities. An overview of the selected genes with observed mutant phenotypes in human, mice and zebrafish is provided in Supplementary Table 4. In this study we propose a crispant screening platform that combines various phenotypic and molecular assays at three key developmental and adult stages (7, 14 and 90 dpf). To obtain a general overview of the results for the selected crispants, all results were summarized in Table 2.

At two distinct larval stages, 7 dpf and 14 dpf, diverse measurements of the early skeleton were conducted, including an assessment of both the osteoblast-positive and the mineralized surface areas located in the head region. The osteoblast-positive area represents the quantity of osteoblasts within a given structure and is a measure of the formation of bone matrix. Although a general tendency towards increased areas could be noted across different crispants at 14dpf, only crispants for *esr1* showed a consistent increase at both 7 and 14 dpf. Interestingly, Brett et al., 2021 extensively investigated *esr1* mutations in the context of hormone receptor-positive breast cancer but did not evaluate the presence of osteoblasts. However, the depletion of estrogen in an osteoblastic cell line *in vitro* results in the stimulation of osteoblast differentiation (Schiavi *et al.*, 2021).

The extent of mineralization of the head skeletal elements was found to be mostly not different compared to controls, although in the crispants where significant differences were detected at one of the studied time points, the mineralization of the head skeleton elements was mostly lower and of the notochord tip higher. Only crispants for *sec24d*, a gene critical for transporting extracellular matrix (ECM) proteins, exhibited reduced mineralization in different head structures both at 7dpf and 14dpf. These findings aligned with the observed malformations in *bulldog* zebrafish, where disrupted *sec24d* function leads to head abnormalities and growth delays in early larval stages (Sarmah *et al.*, 2010).

Despite considerable variability in the measurements we conducted, a subtle trend

emerged in the expression data for osteogenic markers during early stages. Specifically, the

expression of *bglap*, a late osteoblast differentiation marker appeared to be consistently

downregulated in near all crispants, except for crispants for *sost*, *mbtps2* and *sparc* at 7 dpf.

However, this downregulation became less pronounced at 14 dpf, affecting only crispants for

aldh7a1, daam2, esr1 and *creb3l1*. Interestingly, most crispants for genes involved in the

pathogenesis of osteoporosis exhibited this downregulation, except for *sost*, where the

expression of *bglap* remained unaffected. In contrast, the expression of *col1a1a* showed an

opposing trend. Initially, a modest upregulation of this marker was observed at 7 dpf, limited

to just four crispants. However, this effect intensified over time, with nearly all crispants

displaying upregulation at 14 dpf, except for *aldh7a1*, *esr1*, and *sparc*. The remarkable

upregulation of *col1a1* expression could potentially arise from an abundance of immature

osteoblasts within the cranial skeleton, a phenomenon we frequently observed across the set of

crispants (Amarasekara *et al.*, 2021). These osteoblasts play an active role in the synthesis of

the organic bone matrix, including the production of collagen type I fibers. Interestingly, the

generally diminished expression of the late osteoblastic marker *bglap* corroborates the presence

of less mature osteoblasts (Gavaia *et al.*, 2006). As mainly mature osteoblasts engage in the

deposition of calcium and phosphate ions, essential for the formation of hydroxyapatite crystals,

bone mineralization can be expect to be reduced in the crispants, which results in increased

formation of non-mineralized osteoid deposits. Indeed, we observed that despite often increased

bone osteoblast-positive surfaces, mineralization-positive areas in the head skeleton were

frequently found to be reduced (Salhotra *et al.*, 2020). The differential expression profiles of

the osteogenic markers *bglap* and *col1a1a* highlight their potential as biomarkers for osteoblast

differentiation and bone formation in FBD crispant screens.

In early stages of development, we observed that crispants, although presenting with a

high indel frequency, showed a relatively high variability for the phenotypic parameters we

tested (mineralization and osteoblast surface areas), which often prevented reaching statistical

significance. Phenotypic variability in these zebrafish larvae can be attributed to several factors,

including crispant mosaicism, allele heterogeneity, environmental factors, differences in

genomic background, and slightly variable imaging positioning. In our study, for each crispant

and corresponding controls, all experiments at different ages were conducted on zebrafish from

the same clutch, allowing analysis of a developmental time series in animals with a relative

genetically homogeneous background. Although increasing the number of biological replicates

could potentially enhance statistical power, previous studies have not confirmed this (Winter *et*

al., 2022). Enhancing out-of frame (OOF) rates by simultaneously targeting a single gene with multiple crRNAs could potentially increase the penetrance of phenotypic abnormalities observed in early crispants. By directing crRNAs to multiple loci within one gene, the likelihood of introducing frameshift mutations increases significantly. This is suggested to lead to a higher rate of biallelic knockout and reduce the chances of retaining the functional protein (Kroll et al., 2021). However, this also elevates the likelihood of off-target effects, increases double-strand breaks and toxicity, while simultaneously increasing costs (Wu et al., 2018). Furthermore, also in-frame indels affecting two or more amino acids, detected in a substantial proportion of NGS reads in our study, are likely to have a detrimental effect on protein function (Emond et al., 2020)(Savino et al., 2022)(Miton and Tokuriki, 2023). Therefore, and because we used highly efficient two-part guide RNAs (Alt-R™ CRISPR-Cas9 system), indicated by the high total indel efficiencies we obtained, we opted for single crRNA injections.

In adult crispants, the skeletal phenotype was generally more penetrant. All crispants showed malformed arches, a majority displayed vertebral fractures and fusions and some crispants exhibited distinct quantitative variations in vertebral body measurements. This confirmed the role of the selected genes in skeletal development and homeostasis and their involvement in skeletal disease and established the crispant approach as a valid approach for rapidly providing in vivo gene function data to support candidate gene identification.

Osteoporosis is typically linked with symptoms that appear late in life. However, our results challenged this notion, by showing that in crispants for osteoporosis genes, skeletal abnormalities can manifest earlier and are not solely reflected in changes to bone mineral density (BMD). For instance, we found that aldh7a1 and esr1 crispants displayed significant skeletal anomalies already at larval stages, with aldh7a1 crispants showing early lethality. At adult stages, esr1 and sost crispants demonstrated various morphological abnormalities of the vertebral column such as fusions, compressions, and arch malformations. The more pronounced and earlier phenotypes in these zebrafish crispants are most likely attributed to the quasi knock-out state of the studied genes, while more common less impactful variants in the same genes result in typical late-onset osteoporosis (Laine et al., 2013) . This phenomenon is also observed in knock-out mouse models for these genes (Melville et al., 2014)(Coughlin et al., 2019).

Interestingly, all adult crispants for osteoporosis genes demonstrated a substantial increase in centrum Tissue Mineral Density (TMD). TMD is quantitative measure of the actual mineral volume within bone tissue, computed as bone mineral content over the bone volume (mgHA/cm3) and obtained through mCT or pQCT. In human patients, reduced bone mineral density (aBMD), a measure for the bone mineral content in a projected bone area and assessed

using dual energy X-ray absorptiometry (DXA), is frequently linked to osteoporosis. However,

also TMD, or vBMD, can be evaluated in the trabecular and cortical bone. Knock-out mice

lacking Sost showed an increase in TMD within the cortical bone, which is comparable to the

bone present in the zebrafish vertebral column (Li *et al.*, 2008). As for the other genes, for

daam2, mutant mice with a hypomorphic *Daam2* allele presented with reduced bone strength

and increased cortical bone porosity (Morris *et al.*, 2019). *ERα* knockout mice presented with

a decrease in cortical bone mineral density and increase in trabecular bone mineral density

(Melville *et al.*, 2014). Further studies are needed to better understand how high TMD values

observed in these crispants relate to phenotypic changes observed in these mouse models.

Finally, it is worthwhile to note that, osteoporosis typically correlates with low vBMD values.

While neither the causal variant nor the biological direction of its effects on the target gene are

known for BMD-associated loci harboring these genes, the increased TMD following gene

knockdown in our models may help to assign the directionality of their effects, and puts forth

the possibility that variants at these loci alter the expression of genes that converge on a

common biological process that in zebrafish, when disrupted results in the high TMD.

Conversely, crispants for genes involved in recessive forms of OI, an early-onset severe

skeletal disorder, can intuitively be expected to exhibit pronounced skeletal abnormalities

already in larval stages. Interestingly, during these early stages, the majority of the crispants do

not exhibit consistent skeletal abnormalities. Although this could be partially explained by the

above-discussed variability of the phenotypic outcomes we measured, also human OI subjects

with these severe conditions, often experience disease onset during infancy only, with

progressive worsening of the skeletal phenotype (Selina *et al.*, 2023).

The phenomenon of genetic compensation could play a role in explaining why early

phenotypes manifest differently in human patients compared to zebrafish. Genetic

compensation ensures an organism to maintain its genetic robustness, even when genetic

variation, possibly induced by mutations, occurs. This compensation mechanism can be

triggered by the upregulation of compensating genes, often mediated through the regulation of

nonsense-mediated decay (NMD) in combination with the epigenetic machinery in zebrafish

(El-Brolosy Mohamed A. and Stainier, 2017) (Jakutis and Stainier, 2021). Crispants, in contrast

to morphants, also undergo NMD although in a mosaic manner. In larval crispant,

compensatory mechanisms may effectively mask the impact of gene loss. However, in adult

crispants, the cumulative effect of this loss becomes more pronounced, leading to a later onset

phenotype. Furthermore, specific tissues or cell types within the organism may activate

compensatory pathways more efficiently, thereby influencing the overall manifestation of the

phenotype. When a gene is disrupted in humans, the consequences can be more severe due to

the complexity of human development and biology. Such disruptions may result in

developmental abnormalities, leading to an early onset of phenotypic traits (Rouf *et al.*, 2023).

To conclude, our study proposes a multifaceted approach to investigate zebrafish crispants

associated with heritable FBDs, employing various techniques, and assessing different skeletal

phenotypes and molecular profiles at distinct developmental stages. While this work represents

a pioneering effort in establishing a screening platform for skeletal diseases, it offers

opportunities for future improvement and refinement. For instance, analyzing different skeletal

structures and cell types within various transgenic lines, increasing sample size, and

incorporating a broader range of techniques, such as phenotypic scoring based on the type of

malformations seen in the vertebral column or alcian blue staining for cartilage tissue

evaluation, could enhance the comprehensiveness of the screening process and the skeletal

phenotype. Nonetheless, our initial set-up demonstrates the feasibility of this approach to

identify genes potentially implicated in the pathogenesis of skeletal disorders, as all studied

crispants exhibit skeletal abnormalities in adult stages. These findings support the prospect of

crispant screening as a promising tool for rapid functional assessment of candidate genes for

skeletal disorders, providing valuable insights into the roles of these genes in skeletal biology.

Materials and methods

Zebrafish maintenance and breeding

The zebrafish were incubated at 28°C until 5 days post-fertilization (dpf). Then, they were reared in a rotifer/zebrafish polyculture system from 5dpf to 9dpf following the method of Lawrence et al., 2016. After 10dpf, they were transferred to 3.5 liter tanks in the Core Zebrafish Facility Ghent (ZFG) and housed in semi-closed recirculating systems (Tecniplast). The water parameters were maintained at 27°C, pH 7.5, conductivity 550 µS, and a 14/10 light/dark cycle. The zebrafish were fed with Zebrafeed (Sparos) and Gemma Micro dry food (Inve) in the morning and Micro Artemia (Ocean Nutrition) in the afternoon. The *osx*:Kaede transgenic line was used for all experiments. Breeding and embryo collection were done according to standard protocols (Westerfield, 2000). All animal studies complied with EU Directive 2010/63/EU for animals, permit no. ECD 23/27 (Ghent University). Pain, distress, and discomfort were minimized as much as possible.

Generation of crispants through CRISPR/Cas9 genome editing

Guide RNA design

For the generation of crispants we largely followed the methods described in Bek et al. (2021). In summary, two-part guide RNAs (gRNAs) were used which are functional gRNA duplexes made of a universal 67 mer trans-activating CRISPR-RNA (tracrRNA) and a target-specific CRISPR-RNA (crRNA) (Alt-R™ CRISPR-Cas9 System, IDT). A non-specific crRNA (GCAGGCAAAGAATCCCTGCC), referred to as the 'scrambled' sequence, was used to generate the controls (Wu *et al.*, 2018).

The crRNAs corresponding to the genes of interest were designed with the web-based tool Benchling (https://benchling.com), using GRCz11 as a reference genome. With this tool, 20 bp crRNAs target sequences, that are followed by a 'NGG' PAM sequence, were listed. Further selection was based on a few specific criteria to minimize the risk of producing (partially) active truncated protein products and potential off-target effects. Specifically, crRNA target sequences located in exon 1, in the last exon, within the last 55bp of the second-to-last exon or in exons consisting of (n) x 3 bp were excluded. Additionally, crRNAs with the highest possible scores for both on-target (Doench *et al.*, 2016) and off-target (Hsu *et al.*, 2013)were selected.

For the selected crRNA sequences, the Indelphi tool
(https://indelphi.giffordlab.mit.edu) employing the mESC training set, was used to predict the
DNA repair outcomes (mixture of indels) resulting from non-homologous end-joining (NHEJ)
and microhomology-mediated end-joining (MMEJ) induced by CRISPR/Cas9 cleavage. The
crRNAs that were predicted to result in high percentages of out-of-frame indels and a limited
number of possible genotypes (high precision) were prioritized.

Eventually, 1 crRNA per gene was ordered from IDT (https://www.idtdna.com):
aldh7a1 (AATTGTTCGACAGATTGGAG), *daam2* (GATCTATTGCAGTAAGAAGA), *esr1*
(GCTCACGACAGAAACACAGC), *sost* (CTGCTCACTCCCGCACCCAG), *creb3l1*
(ACACAGTTACTCTCTCAGCG), *ifitm5* (AGCGCAGGAATGCTCAGACA), *mbtps2*
(TTCCATATCAAGTGGCACAC), *sec24d* (GCCTATGGATCTCCAACACA), *serpinf1*
(CAGGTTGTAGCCAAAATCAG), *sparc* (AGAGGAGCCAGCTGTTGAAG).

To predict the potential off-target effects of the selected crRNAs, Crispor
(http://crispor.tefor.net) was used. The cutting frequently determination (CFD) off-target score
was used to predict the probability of off-target cleavage. The three targets with the highest
CFD scores were selected and PCR-amplified (Supplementary Table 1). Next-generation
sequencing using the Miseq platform was used to assess the frequency of the off-target effects
(Concordet and Haeussler, 2018).

Injection

To prepare gRNA duplexes for each assay, we mixed 200µM crRNA and 200µM
tracrRNA, heated the mixture at 95°C for 5 minutes, and then cooled it to room temperature.
Next, we micro-injected one-cell-stage Tg(osx:Kaede) zebrafish embryos. The injection mix
consisted of a 1.4 nL drop containing RNase-free water, KCl (2000 mM) and
ribonucleoproteins (RNPs), consisting of 800 pg of both gRNA duplex and Cas9 (1000 ng/µl)
(ToolGen, Seoul, South Korea; wild-type nuclease with nuclear localization signal) protein. For
each crispant and each corresponding control, a total of 200 eggs, originating from the same
clutch, were injected.

Next-generation sequencing analysis

We extracted DNA from 10 pooled embryos at one day after fertilization by adding 100
µl NaOH and incubating the samples at 95°C for 20 minutes. We then added 10µL Tris-HCl.

We performed PCR amplification with the primers listed in Supplementary Table 2, using either

'FORD58' or 'FORD60' assays depending on the primer. The thermocycling conditions for the

'FORD58' assay were: 95°C for 3 min, followed by 40 cycles of 95°C for 15s, 58°C for 10s,

and 72°C for 15s, and a final extension at 72°C for 10 min. The thermocycling conditions for

the 'FORD60' assay were: 95°C for 3 min, followed by 40 cycles of 95°C for 15s, 60°C for

10s, and 72°C for 15s, and a final extension at 72°C for 10 min. We prepared the PCR amplicons

for sequencing using the Nextera XT DNA Library Preparation Kit (Illumina, San Diego, CA,

USA) and sequenced them on a MiSeq instrument (Illumina) using 2 X 250 bp cycles (Bek *et*

al., 2021). We analyzed the NGS data with CRISPResso2 and determined the mean fraction of

reads containing in-frame and out-of-frame indels over different samples, by exporting the

allele frequency table, counting the number of reads with both in- and out-of-frame indels or

out-of-frame indels only and dividing these numbers by the total number of mapped reads

(Fiume *et al.*, 2019). For both the indel and out-of-frame efficiency, the reads with sequencing

errors and natural variants were excluded.

RNA extraction and RT-qPCR

The RNA extraction and qPCR protocol described by Bek et al., 2021 was followed,

with minor modifications; for each crispant, RNA was extracted from 4 or 5 pools of 10

embryos at 7 dpf and 14 dpf using Trizol (Life Technologies Europe) and purified using the

RNeasy Mini Kit (Qiagen) with on-column DNase I treatment (Qiagen). RNA concentrations

were measured with the Little Lunatic (Unchained Labs) and cDNA was synthesized using the

iScript cDNA synthesis kit (Bio-rad). Primers for skeletal marker genes were designed using

NCBI Primer-BLAST (Supplementary Table 3) and primers for the housekeeping genes (*elfa*

and *bactin2*) were obtained from literature (McCurley & Callard, 2008).

The qPCR reaction mixture was prepared with 2.5 µL of SsoAdvanced Universal SYBR

Green Supermix (Bio-rad), 5 ng of cDNA, and 250 nM of each primer. The qPCR reaction was

run on a LightCycler 480 instrument (Roche) in a white 384-well plate with the following

thermocycling conditions: initial denaturation step at 95 °C for 2 min, followed by 44 cycles of

denaturation at 95 °C for 5 sec, annealing at 60 °C for 30 sec, and extension at 72 °C for 1 sec.

Subsequently, a melting curve analysis was performed with a denaturation step at 95 °C for 5

sec, followed by annealing at 60 °C for 1 min and gradual heating to 95 °C at a ramp-rate of

0.11 °C/s. Finally, the product was cooled to 37 °C for 3 min (Vanhauwaert *et al.*, 2014). The

data was analyzed using the qBase+ software (Biogazelle).

Osteoblast imaging in transgenic *osx:*Kaede larvae.

Ten zebrafish larvae (at 7 dpf and 14 dpf) with the transgenic *osx:Kaede* background were anesthetized using 1x Tricaine. Imaging was performed using the Leica microscope M165FC and mounted camera (DFC450C) operated by LAS V4.3 software (Leica Microsystems, Wetzlar, Germany). Images of 7 dpf larvae were taken from the ventral side. Images of 14 dpf larvae were taken from both ventral and lateral perspectives to provide a more comprehensive view of the operculum's surface area.

Fiji software (version 2.14.0), powered by ImageJ, was employed for all measurements (Schindelin *et al.*, 2012). Osteoblast-positive area of both the total head and operculum was quantified using thresholding, followed by measurement of the positive area. All measurements were performed in triplicate on blinded pictures. The presence of osteoblasts in the operculum was measured by hand using the polygon selection tool. Unpaired T-test was applied in GraphPad Prism (v.9.4.0), with p-values considered significant when ≤0.05.

Alizarin red staining and imaging of zebrafish larvae

The ten zebrafish larvae, that were previously used for osteoblast imaging at 7 dpf and 14 dpf, were euthanized using 25x Tricaine. They were transferred to a cell strainer and washed with dH2O several times to remove the Tricaine. The strainers were placed in a 6-well plate and incubated in 4% PFA overnight at 4°C. The plates were sealed with parafilm to prevent evaporation. The larvae were then washed with dH2O several times to remove the PFA and bleached for 30 minutes at room temperature using a solution containing 3% H_2O_2 and 2% KOH to remove pigmentation. Next, they were stained overnight at 4°C using Alizarin Red S (0.005% in 1% KOH/2% Triton). The samples were washed with dH2O several times and transferred to increasing concentrations of glycerol (30-70-100% for 5 minutes each) for storage (4°C) and imaging. Zebrafish larvae stained with Alizarin red were then imaged using a Leica M165FC microscope and DFC450C camera with the LAS V4.3 software (Leica Microsystems). The ventral side of the larvae was captured at 7 dpf. Both ventral and lateral perspectives were captured at 14 dpf to provide a more comprehensive view of the operculum's surface area and the number of mineralized vertebrae.

Fiji software (version 2.14.0), powered by ImageJ, was employed for all measurements (Schindelin *et al.*, 2012). The red stain in the head elements was assessed using thresholding, followed by quantification of the positive area. These measurements were conducted in

592 triplicate on blinded pictures. The stained notochord tip and operculum was measured by hand

593 using the polygon selection tool. The statistical analysis employed the Unpaired T-test in

594 GraphPad Prism (v.9.4.0), with p-values considered significant when ≤0,05.

Micro-Computed Tomography (MicroCT) analysis in adult zebrafish

Adult zebrafish (3 months post fertilization) were euthanized using a 25x tricaine

solution (4000 mg tricaine powder (Sigma), 979 ml distilled water, and 1M Tris, adjusted to

pH 7). Brightfield images of the zebrafish were taken using a Leica M165FC microscope and

DFC450C camera with the LAS V4.3 software (Leica Microsystems). The standard lengths of

the zebrafish were measured using the Fiji (ImageJ) software (ImageJ 1.52i) (Schindelin *et al.*,

2012).

The fish were fixed for approximately one week in a fixative solution (4%

paraformaldehyde, 5% triton X-100, and 1% potassium hydroxide). The solution was changed

when it turned yellow. A maximum of five fish per 100 ml of fixative solution was used. After

fixation, the fish were transferred to a 70% ethanol solution for storage until scanning.

Before scanning, three fish were transferred to a 15 ml Eppendorf tube and immobilized

using paper towel soaked in 70% ethanol. The Eppendorf tubes were scanned at the custom-

built HECTOR scanner of the Centre for X-ray Tomography (UGCT, www.ugct.ugent.be). A

hardware filter of 0.5 mm Al was used to eliminate beam hardening artefacts, with the tube

voltage at 90 kVp and target power of 10 W. During a 360° rotation, 1201 radiographic images

were acquired at 1 s acquisition time per image. The raw data was reconstructed using the

Octopus Reconstruction software tool, resulting in a 3D volume of approximately

2800x1000x1000 voxels. Using geometrical magnification, a voxel size of 20^3 µm³ was

obtained (Masschaele *et al.*, 2013).

The 3 fish in one tube were virtually separated from the scan, aligned and exported as 3

individual single tiff volume using VGStudioMAX (VolumeGraphics, Heidelberg, Germany).

The generated volumes were analyzed using the FishCut software in Matlab described in Hur

et al., 2017. The resulting plots were generated using R statistics and were color-coded to

highlight significant differences. Significant differences were represented with a lighter color

scheme.

Alizarin red staining and imaging of adult zebrafish

The fish used for microCT were reused for Alizarin Red staining for which the protocol

has been described in Bek et al., 2021. In short, the fish were transferred to cassettes and the

concentration of EtOH was gradually reduced from 70% to distilled water (70%-50%-30%-

dH2O each 30 min). Then, the samples were enhanced, immersed into the bone-staining

medium, stained with bone-staining solution and destained after removal of the scales with

destaining solution. Finally, the samples were gradually (30-70-100%) transferred to 100%

glycerol for storage (4°C) and imaging.

Adult zebrafish were imaged using a Leica M165FC microscope and DFC450C camera

with the LAS V4.3 software (Leica Microsystems). The DSR DsRed Dichroic Mirror filter

was used to visualize the (red) fluorescent signal. Images were obtained from a lateral view to

get a clear picture of the whole fish. Higher magnification was used to get detailed images of

the vertebral column and the head. Images of the head were used to measure the head size, using

the Fiji (ImageJ) software (Schindelin *et al.*, 2012).

References

Amarasekara, D.S., Kim, S. and Rho, J. (2021), "Regulation of osteoblast differentiation by

cytokine networks", *International Journal of Molecular Sciences*, MDPI AG, 2 March,

doi: 10.3390/ijms22062851.

Bek, J.W., Shochat, C., De Clercq, A., De Saffel, H., Boel, A., Metz, J., Rodenburg, F., *et al.*

(2021), "Lrp5 Mutant and Crispant Zebrafish Faithfully Model Human Osteoporosis,

Establishing the Zebrafish as a Platform for CRISPR-Based Functional Screening of

Osteoporosis Candidate Genes", *Journal of Bone and Mineral Research*, John Wiley and

Sons Inc, Vol. 36 No. 9, pp. 1749–1764, doi: 10.1002/jbmr.4327.

Brent Richards, J., Zheng, H.F. and Spector, T.D. (2012), "Genetics of osteoporosis from

genome-wide association studies: Advances and challenges", *Nature Reviews Genetics*,

August, doi: 10.1038/nrg3228.

Brett, J.O., Spring, L.M., Bardia, A. and Wander, S.A. (2021), "ESR1 mutation as an

emerging clinical biomarker in metastatic hormone receptor-positive breast cancer",

Breast Cancer Research, BioMed Central Ltd, 1 December, doi: 10.1186/s13058-021-

01462-3.

Charoenngam, N., Rittiphairoj, T., Ponvilawan, B., Jaroenlapnopparat, A., Waitayangkoon,

P., Suppakitjanusant, P., Prasitsumrit, V., *et al.* (2023), "Bone Fragility in Hereditary

Connective Tissue Disorders: A Systematic Review and Meta-Analysis", *Endocrine*

Practice, Vol. 29 No. 7, pp. 589–600, doi: https://doi.org/10.1016/j.eprac.2023.02.003.

Clement, K., Rees, H., Canver, M.C., Gehrke, J.M., Farouni, R., Hsu, J.Y., Cole, M.A., *et al.*

(2019), "CRISPResso2 provides accurate and rapid genome editing sequence analysis",

Nature Biotechnology, Nature Publishing Group, 1 March, doi: 10.1038/s41587-019-

0032-3.

Concordet, J.P. and Haeussler, M. (2018), "CRISPOR: Intuitive guide selection for

CRISPR/Cas9 genome editing experiments and screens", *Nucleic Acids Research*,

Oxford University Press, Vol. 46 No. W1, pp. W242–W245, doi: 10.1093/nar/gky354.

Coughlin, C.R., Swanson, M.A., Spector, E., Meeks, N.J.L., Kronquist, K.E., Aslamy, M.,

Wempe, M.F., *et al.* (2019), "The genotypic spectrum of ALDH7A1 mutations resulting

in pyridoxine dependent epilepsy: A common epileptic encephalopathy", *Journal of*

Inherited Metabolic Disease, John Wiley and Sons Inc., Vol. 42 No. 2, pp. 353–361, doi:

10.1002/jimd.12045.

Daponte, V., Tonelli, F., Masiero, C., Syx, D., Exbrayat-Héritier, C., Biggiogera, M.,
Willaert, A., *et al.* (2023), "Cell differentiation and matrix organization are differentially
affected during bone formation in osteogenesis imperfecta zebrafish models with
different genetic defects impacting collagen type I structure", *Matrix Biology*, Elsevier
B.V., Vol. 121, pp. 105–126, doi: 10.1016/j.matbio.2023.06.003.

Dietrich, K., Fiedler, I.A.K., Kurzyukova, A., López-Delgado, A.C., McGowan, L.M.,
Geurtzen, K., Hammond, C.L., *et al.* (2021), "Skeletal Biology and Disease Modeling in
Zebrafish", *Journal of Bone and Mineral Research*, John Wiley and Sons Inc, 1 March,
doi: 10.1002/jbmr.4256.

Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I.,
et al. (2016), "Optimized sgRNA design to maximize activity and minimize off-target
effects of CRISPR-Cas9", *Nature Biotechnology*, Nature Publishing Group, Vol. 34 No.
2, pp. 184–191, doi: 10.1038/nbt.3437.

El-Brolosy Mohamed A. AND Stainier, D.Y.R. (2017), "Genetic compensation: A
phenomenon in search of mechanisms", *PLOS Genetics*, Public Library of Science, Vol.
13 No. 7, pp. 1–17, doi: 10.1371/journal.pgen.1006780.

Emond, S., Petek, M., Kay, E.J., Heames, B., Devenish, S.R.A., Tokuriki, N. and Hollfelder,
F. (2020), "Accessing unexplored regions of sequence space in directed enzyme
evolution via insertion/deletion mutagenesis", *Nature Communications*, Nature
Research, Vol. 11 No. 1, doi: 10.1038/s41467-020-17061-3.

Fiume, M., Cupak, M., Keenan, S., Rambla, J., de la Torre, S., Dyke, S.O.M., Brookes, A.J.,
et al. (2019), "Federated discovery and sharing of genomic data using Beacons", *Nature
Biotechnology*, Nature Publishing Group, 1 March, doi: 10.1038/s41587-019-0046-x.

Forlino, A. and Marini, J.C. (2016), "Osteogenesis imperfecta", *The Lancet*, Lancet
Publishing Group, 16 April, doi: 10.1016/S0140-6736(15)00728-X.

Gavaia, P.J., Simes, D.C., Ortiz-Delgado, J.B., Viegas, C.S.B., Pinto, J.P., Kelsh, R.N.,
Sarasquete, M.C., *et al.* (2006), "Osteocalcin and matrix Gla protein in zebrafish (Danio
rerio) and Senegal sole (Solea senegalensis): Comparative gene and protein expression
during larval development through adulthood", *Gene Expression Patterns*, Vol. 6 No. 6,
pp. 637–652, doi: 10.1016/j.modgep.2005.11.010.

Gistelinck, C., Kwon, R.Y., Malfait, F., Symoens, S., Harris, M.P., Henke, K., Hawkins,
M.B., *et al.* (2018), "Zebrafish type I collagen mutants faithfully recapitulate human type
I collagenopathies", *Proceedings of the National Academy of Sciences of the United*

States of America, National Academy of Sciences, Vol. 115 No. 34, pp. E8037–E8046, doi: 10.1073/pnas.1722200115.

Howe, K., Clark, M.D., Torroja, C.F., Torrance, J., Berthelot, C., Muffato, M., Collins, J.E., *et al.* (2013), "The zebrafish reference genome sequence and its relationship to the human genome", *Nature*, Nature Publishing Group, Vol. 496 No. 7446, pp. 498–503, doi: 10.1038/nature12111.

Hsu, P.D., Scott, D.A., Weinstein, J.A., Ran, F.A., Konermann, S., Agarwala, V., Li, Y., *et al.* (2013), "DNA targeting specificity of RNA-guided Cas9 nucleases", *Nature Biotechnology*, Vol. 31 No. 9, pp. 827–832, doi: 10.1038/nbt.2647.

Hur, M., Gistelinck, C.A., Huber, P., Lee, J., Thompson, M.H., Monstad-Rios, A.T., Watson, C.J., *et al.* (2017), "MicroCT-based phenomics in the zebrafish skeleton reveals virtues of deep phenotyping in a distributed organ system", doi: 10.7554/eLife.26014.001.

Jovanovic, M., Guterman-Ram, G. and Marini, J.C. (2022), "Osteogenesis Imperfecta: Mechanisms and Signaling Pathways Connecting Classical and Rare OI Types", *Endocrine Reviews*, Endocrine Society, 1 February, doi: 10.1210/endrev/bnab017.

Kague, E. and Karasik, D. (2022), "Functional Validation of Osteoporosis Genetic Findings Using Small Fish Models", *Genes*, MDPI, 1 February, doi: 10.3390/genes13020279.

Kanis, J.A., Melton, L.J., Christiansen, C., Johnston, C.C. and Khaltaev, N. (1994), "The diagnosis of osteoporosis", *Journal of Bone and Mineral Research*, Vol. 9 No. 8, pp. 1137–1141, doi: 10.1002/jbmr.5650090802.

Kroll, F., Powell, G.T., Ghosh, M., Gestri, G., Antinucci, P., Hearn, T.J., Tunbak, H., *et al.* (2021), "A simple and effective F0 knockout method for rapid screening of behaviour and other complex phenotypes", *ELife*, Vol. 10, doi: 10.7554/eLife.59683.

Laine, C.M., Joeng, K.S., Campeau, P.M., Kiviranta, R., Tarkkonen, K., Grover, M., Lu, J.T., *et al.* (2013), "WNT1 Mutations in Early-Onset Osteoporosis and Osteogenesis Imperfecta ", *New England Journal of Medicine*, New England Journal of Medicine (NEJM/MMS), Vol. 368 No. 19, pp. 1809–1816, doi: 10.1056/nejmoa1215458.

Lawrence, C., Best, J., Cockington, J., Henry, E.C., Hurley, S., James, A., Lapointe, C., *et al.* (2016), "The complete and updated 'rotifer polyculture method' for rearing first feeding zebrafish", *Journal of Visualized Experiments*, Journal of Visualized Experiments, Vol. 2016 No. 107, doi: 10.3791/53629.

Li, X., Ominsky, M.S., Niu, Q.T., Sun, N., Daugherty, B., D'Agostin, D., Kurahara, C., *et al.* (2008), "Targeted deletion of the sclerostin gene in mice results in increased bone

formation and bone strength", *Journal of Bone and Mineral Research*, Vol. 23 No. 6, pp.
860–869, doi: 10.1359/jbmr.080216.

Masschaele, B., Dierick, M., Loo, D. Van, Boone, M.N., Brabant, L., Pauwels, E., Cnudde,
V., *et al.* (2013), "HECTOR: A 240kV micro-CT setup optimized for research", *Journal*
of Physics: Conference Series, Vol. 463, Institute of Physics Publishing, doi:
10.1088/1742-6596/463/1/012012.

McCurley, A.T. and Callard, G. V. (2008), "Characterization of housekeeping genes in
zebrafish: Male-female differences and effects of tissue type, developmental stage and
chemical treatment", *BMC Molecular Biology*, Vol. 9, doi: 10.1186/1471-2199-9-102.

Melville, K.M., Kelly, N.H., Khan, S.A., Schimenti, J.C., Ross, F.P., Main, R.P. and Van Der
Meulen, M.C.H. (2014), "Female mice lacking estrogen receptor-alpha in osteoblasts
have compromised bone mass and strength", *Journal of Bone and Mineral Research*,
Vol. 29 No. 2, pp. 370–379, doi: 10.1002/jbmr.2082.

Miton, C.M. and Tokuriki, N. (2023), "Insertions and Deletions (Indels): A Missing Piece of
the Protein Engineering Jigsaw", *Biochemistry*, American Chemical Society, 17 January,
doi: 10.1021/acs.biochem.2c00188.

Morris, J.A., Kemp, J.P., Youlten, S.E., Laurent, L., Logan, J.G., Chai, R.C., Vulpescu, N.A.,
et al. (2019), "An atlas of genetic influences on osteoporosis in humans and mice",
Nature Genetics, Nature Publishing Group, Vol. 51 No. 2, pp. 258–266, doi:
10.1038/s41588-018-0302-x.

Naert, T., Tulkens, D., Edwards, N.A., Carron, M., Shaidani, N.I., Wlizla, M., Boel, A., *et al.*
(2020), "Maximizing CRISPR/Cas9 phenotype penetrance applying predictive modeling
of editing outcomes in Xenopus and zebrafish embryos", *Scientific Reports*, Nature
Research, Vol. 10 No. 1, doi: 10.1038/s41598-020-71412-0.

Pouresmaeili, F., Kamalidehghan, B., Kamarehei, M. and Goh, Y.M. (2018), "A
comprehensive overview on osteoporosis and its risk factors", *Therapeutics and Clinical*
Risk Management, Dove Medical Press Ltd., doi: 10.2147/TCRM.S138000.

Ralston, S.H. and Uitterlinden, A.G. (2010), "Genetics of osteoporosis", *Endocrine Reviews*,
October, doi: 10.1210/er.2009-0044.

Rouf, M.A., Wen, L., Mahendra, Y., Wang, J., Zhang, K., Liang, S., Wang, Y., *et al.* (2023),
"The recent advances and future perspectives of genetic compensation studies in the
zebrafish model", *Genes and Diseases*, KeAi Communications Co., 1 March, doi:
10.1016/j.gendis.2021.12.003.

Sabik, O.L. and Farber, C.R. (2017), "Using GWAS to identify novel therapeutic targets for
osteoporosis", *Translational Research*, Mosby Inc., 1 March, doi:
10.1016/j.trsl.2016.10.009.

Salhotra, A., Shah, H.N., Levi, B. and Longaker, M.T. (2020), "Mechanisms of bone
development and repair", *Nature Reviews Molecular Cell Biology*, Nature Research, 1
November, doi: 10.1038/s41580-020-00279-w.

Sarmah, S., Barrallo-Gimeno, A., Melville, D.B., Topczewski, J., Solnica-Krezel, L. and
Knapik, E.W. (2010), "Sec24D-dependent transport of extracellular matrix proteins is
required for zebrafish skeletal morphogenesis", *PLoS ONE*, Vol. 5 No. 4, doi:
10.1371/journal.pone.0010367.

Savino, S., Desmet, T. and Franceus, J. (2022), "Insertions and deletions in protein evolution
and engineering", *Biotechnology Advances*, Elsevier Inc., 1 November, doi:
10.1016/j.biotechadv.2022.108010.

Schiavi, J., Fodera, D.M., Brennan, M.A. and McNamara, L.M. (2021), "Estrogen depletion
alters osteogenic differentiation and matrix production by osteoblasts in vitro",
Experimental Cell Research, Elsevier Inc., Vol. 408 No. 1, doi:
10.1016/j.yexcr.2021.112814.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., *et al.* (2012), "Fiji: An open-source platform for biological-image
analysis", *Nature Methods*, July, doi: 10.1038/nmeth.2019.

Selina, A., Kandagaddala, M., Kumar, V., Abraham, S.S.C., Danda, S. and Madhuri, V.
(2023), "SERPINF1 gene variants causing late-onset progressive deforming osteogenesis
imperfecta – A study of 18 patients from India", *Bone Reports*, Elsevier Inc., Vol. 18,
doi: 10.1016/j.bonr.2023.101690.

Uribe-Salazar, J.M., Kaya, G., Sekar, A., Weyenberg, K.C., Ingamells, C. and Dennis, M.Y.
(2022), "Evaluation of CRISPR gene-editing tools in zebrafish", *BMC Genomics*,
BioMed Central Ltd, Vol. 23 No. 1, doi: 10.1186/s12864-021-08238-1.

Valenti, M.T., Marchetto, G., Mottes, M. and Dalle Carbonare, L. (2020), "Zebrafish: A
Suitable Tool for the Study of Cell Signaling in Bone", *Cells*, NLM (Medline), 17
August, doi: 10.3390/cells9081911.

Vanhauwaert, S., Van Peer, G., Rihani, A., Janssens, E., Rondou, P., Lefever, S., De Paepe,
A., *et al.* (2014), "Expressed repeat elements improve RT-qPCR normalization across a
wide range of zebrafish gene expression studies", *PLoS ONE*, Public Library of Science,
Vol. 9 No. 10, doi: 10.1371/journal.pone.0109091.

Watson, C.J., Monstad-Rios, A.T., Bhimani, R.M., Gistelinck, C., Willaert, A., Coucke, P.,

Hsu, Y.H., *et al.* (2020), "Phenomics-Based Quantification of CRISPR-Induced

Mosaicism in Zebrafish", *Cell Systems*, Cell Press, Vol. 10 No. 3, pp. 275-286.e5, doi:

10.1016/j.cels.2020.02.007.

Westerfield, M. (2000), *The Zebrafish Book: A Guide for the Laboratory Use of Zebrafish*

(Danio Rerio), University of Oregon Press.

Wu, R.S., Lam, I.I., Clay, H., Duong, D.N., Deo, R.C. and Coughlin, S.R. (2018), "A Rapid

Method for Directed Gene Knockout for Screening in G0 Zebrafish", *Developmental*

Cell, Cell Press, Vol. 46 No. 1, pp. 112-125.e4, doi: 10.1016/j.devcel.2018.06.003.

Zhu, X., Bai, W. and Zheng, H. (2021), "Twelve years of GWAS discoveries for osteoporosis

and related traits: advances, challenges and applications", *Bone Research*, Springer

Nature, 1 December, doi: 10.1038/s41413-021-00143-3.

Figures

Figure 1: Measurements of osteoblast-positive surface area and mineralized surface area of the head skeleton in 7 and 14 dpf crispants. (a) Schematic overview of the ventral and lateral perspective of the head of zebrafish larvae at 7 and 14 dpf. The surface area of the total head is circled with a dashed line. Arrows point to the notochord tip (n), the operculum (o) and the mineralized vertebrae. The eyes (e) and parasphenoid (p) are also shown. (b) Quantification of the osteoblast-positive surface area of the total head and the mineralized surface area of the total head and notochord tip at 7 dpf in comparison with their respective controls. (c) Quantification of the osteoblast-positive surface area of the total head and the mineralized surface area of the total head, operculum and notochord tip and number of vertebrae at 14 dpf in comparison with their respective controls. For easier visualization, obtained results were normalized to the respective controls (normalization = individual value crispant (or control) / mean control) (n=10). Statistical significance is evaluated using the unpaired T-test on non-normalized data and significant differences were visualized using an asterix (* = p ≤0,05; ** = p ≤0,01; *** = p ≤0,001; **** = p ≤0,0001). Error bars show the standard deviation of non-normalized data.

868 *Figure 2: RT-qPCR expression analysis of runx2, sp7, bglap and col1a1a in crispants at 7 (a) and 14 dpf (b) and*

869 *their respective controls, normalized according to the controls (normalization = individual values crispant (or*

870 *control) / mean control). Statistical significance is evaluated using the unpaired T-test on non-normalized data*

871 *and significant differences were visualized using an asterix (* = p ≤0,05; ** = p ≤0,01; *** = p ≤0,001; *****

872 *= p ≤0,0001). Error bars show the standard deviation of non-normalized data.*

885 *Figure 3: ARS images and survival analysis of mbtps2 crispants. (a) ARS images of control fish and crispants for*

886 *mbtps2, showing severe craniofacial abnormalities (arrows). (b) Survival curve of controls and crispants for*

887 *mbtps2, showing a reduction in survival starting from 17 dpf. (c) Survival curve of controls and crispants for*

888 *aldh7a1, showing a reduction in survival starting from 7 dpf.*

Figure 4: Skeletal phenotyping of adult crispants and their respective controls (a) Pictures of ARS-stained

vertebral column of a control and three crispants (from left to right: esr1, ifitm5 and creb3l1), showing fractures

(arrow), fusions and compressions (squared) and malformations in the arches (arrowhead). (b) Measurements of

the standard length, the head size and the eye diameter of the crispants compared to their control. The standard

length was measured from the snout tip to the tail base. The head size was measured from the snout tip to the supraoccipital bone and the eye diameter was measured from the lateral ethmoid to the hyomandibula. The data was normalized for easier visualization (normalization = individual value crispants (or controls) / mean control) (n=5)). (c) Quantification of the number of fractures, fusions or compressions and malformations in the arches of crispants and the controls. For this quantification, twelve vertebrae were selected per fish and a total of 5 crispants per assay was evaluated. Statistical analysis was evaluated using the unpaired T-test on non-normalized data and significant differences were visualized using an asterix (* = p ≤0,05; ** = p ≤0,01; *** = p ≤0,001; **** = p ≤ 0,0001). Error bars showed the standard deviation of non-normalized data. (d) Quantification of skeletal parameters by quantitative micro-computed tomography (µCT) analysis. In the graphical representation, the different crispants were listed. Statistically significant differences from the control values for a given crispant were depicted as red bars. Significance levels were determined through the global test analysis (Supplementary Figure 7).

Tables

	aldh7a1	daam2	esr1	sost	creb3l1	ifitm5	mbtps2	sec24d	serpinf1	sparc
% reads with indel	87%	96%	92%	80%	71%	90%	94%	85%	92%	92%
% reads with out-of-frame indel	66%	57%	64%	64%	49%	73%	73%	63%	55%	72%
% reads with in frame indel ≥ 6-bp	16%	24%	15%	14%	11%	15%	19%	12%	25%	16%
% reads likely affecting protein function	82%	81%	79%	78%	60%	88%	92%	75%	80%	88%

Table 1: InDelphi-mESC prediction, indel, out-of-frame indel and in-frame rates for the different crispants, determined by NGS from pools of 1dpf embryos. The percentage of reads likely to affect the protein function is calculated based on the percentage of reads with out-of-frame indels combined with the percentage of reads with in-frame deletions of equal to or more than 6 basepairs.

	Osteoblast-positive surface area total head 7 dpf	Osteoblast-positive surface area total head 14 dpf	Osteoblast-positive surface area operculum 14 dpf	Mineralized surface area total head 7 dpf	Mineralized surface area total head 14 dpf	Mineralized surface area notochord tip 7 dpf	Mineralized surface area notochord tip 14 dpf	Mineralized surface area operculum 14 dpf	Number of mineralized vertebrae	runx2 expression 7 dpf	runx2 expression 14 dpf	sp7 expression 7 dpf	sp7 expression 14 dpf	bglap expression 7 dpf	bglap expression 14 dpf	col1a1a expression 7 dpf	col1a1a expression 14 dpf	Standard length 90 dpf	Head size 90 dpf	Diameter eye 90 dpf	Centrum Volume 90 dpf	Centrum TMD 90 dpf	Centrum Thickness 90 dpf	Centrum Length 90 dpf
aldh7a1	ns	ns	ns	(-)*	(-)*	ns	ns	ns	ns	ns	ns	ns	ns	(-)*	(-)***	(+)***	ns	■	■	■	■	■	■	■
daam2	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	(+)*	(-)*	(-)**	(+)*	(+)****	ns	ns	ns	(+)*	(+)*	ns	(+)*
esr1	(+)*	(+)***	(+)***	ns	ns	ns	(+)**	ns	ns	ns	ns	ns	(-)*	(-)*	(-)**	(+)*	ns	ns	ns	ns	ns	(+)*	(+)*	ns
sost	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	(+)***	ns	ns	ns	ns	ns	(+)*	ns	ns
creb3ll	ns	ns	ns	ns	ns	(+)*	ns	ns	ns	ns	ns	ns	(-)*	(-)**	ns	(+)***	ns	ns	ns	ns	ns	ns	ns	ns
ifitm5	ns	(+)**	(+)**	ns	ns	(+)*	(+)*	ns	ns	ns	ns	ns	(-)*	ns	ns	(+)*	ns	ns	ns	ns	ns	ns	(+)*	ns
mbtps2	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	(+)**	ns	■	■	■	■	■	■	■
sec24d	ns	ns	ns	(-)****	ns	(+)*	ns	(-)**	(-)**	ns	ns	ns	(+)*	(-)**	ns	(+)*	(+)****	ns	ns	ns	ns	(+)*	ns	ns
serpinf1	ns	ns	(+)**	ns	ns	ns	ns	ns	ns	ns	ns	ns	(-)*	ns	ns	(+)*	ns	ns	ns	ns	ns	ns	ns	ns
sparc	ns	ns	(+)*	ns	ns	ns	ns	ns	ns	ns	(+)***	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns

Table 2: Overview of the different measurements in crispants at 7, 14 and 90 dpf. (-): downregulation or less, (+) upregulation or more, ns: not significant, asterix defines statistical significance (* = p ≤0,05; ** = p ≤0,01; *** = p ≤0,001; **** = p ≤0,0001).