The effect of long-range linkage disequilibrium on allele-frequency dynamics under stabilizing selection

Sherif Negm[1]   Carl Veller[2,*]

Abstract

Stabilizing selection on a polygenic trait reduces the trait's genetic variance by (i) generating correlations (linkage disequilibria) between opposite-effect alleles throughout the genome and (ii) selecting against rare alleles at polymorphic loci that affect the trait, eroding heterozygosity at these loci. Here, we characterize the impact of the linkage disequilibria, which stabilizing selection generates on a rapid timescale, on the subsequent allele-frequency dynamics at individual loci, which proceed on a slower timescale. We obtain expressions for the expected per-generation change in minor-allele frequency at individual loci, as functions of the effect sizes at these loci, the strength of selection on the trait, its variance and heritability, and the linkage relations among loci. Using whole-genome simulations, we show that our expressions predict allele-frequency dynamics under stabilizing selection more accurately than the expressions that have previously been used for this purpose. Our results have implications for understanding the genetic architecture of complex traits.

1 Introduction

To understand the genetic architecture of polygenic traits, we need to connect population genetic models with data such as from genome-wide association studies (GWASs). For non-disease traits, a particularly plausible model is stabilizing selection, which penalizes deviations from an optimal trait value. Theoretical argument and empirical evidence indicate that many complex traits are under stabilizing selection (Sella and Barton 2019). For example, in humans, Sanjak et al. (2018) used lifetime reproductive success as a proxy for fitness and estimated significant nonlinear selection gradients consistent with stabilizing selection for a large fraction of the traits they analyzed. Additionally, Simons et al. (2018, 2022) have demonstrated that, for many human traits, the joint distribution of allele frequencies and effect sizes (as inferred from GWAS) is particularly consistent with stabilizing selection.

The macroscopic consequence of stabilizing selection on a complex trait is to reduce the trait's genetic variance over time. This is achieved in two ways. First, by selecting for compensating combinations of trait-increasing and trait-decreasing alleles, stabilizing selection rapidly generates negative correlations—linkage disequilibria (LD)—between alleles with the same directional effect on the trait (Bulmer 1971; this has come to be known as the 'Bulmer effect'). Second, stabilizing selection generates weak selection against the rarer allele at each polymorphic locus affecting a complex trait, slowly eroding heterozygosity at these loci (Robertson 1956).

This second variance-reducing effect—which mimics underdominant selection—can be understood intuitively in the following way (see also Section 3.1 below). Consider a focal polymorphic locus affecting

---

[1]Department of Human Genetics, University of Chicago
[2]Department of Ecology & Evolution, University of Chicago
 *cveller@uchicago.edu

the trait, and suppose that the common allele at the locus is trait-increasing, so that the mean effect of the locus is to increase the trait. For the population's average trait value to be at the optimum, the mean effect of the rest of the genome must compensate for the mean effect at the focal locus, and must therefore be trait-decreasing. But this leaves the rare allele at the focal locus—which is also trait-decreasing—maladapted to the rest of the genome: on average, it resides in individuals with trait values further from the optimum than the common allele at the locus.

The speed of the resulting allele-frequency dynamics depends, as the intuition above suggests, on the mean phenotypes experienced by the trait-increasing and trait-decreasing alleles at the focal locus. These mean phenotypes are determined not only by the individual effects of the alleles themselves in concert with the mean effect of the rest of the genome, but also by the effects of alleles with which the alleles at the focal locus are associated via LD. Because of the Bulmer effect, alleles at the focal locus will tend to be associated with opposite-effect alleles elsewhere in the genome, partially masking the individual effects of the alleles at the focal locus. This brings the mean phenotypes experienced by the alleles at the focal locus closer to the optimum, weakening selection between them and therefore slowing down their frequency dynamics.

Here, we quantify the slowdown in allele-frequency dynamics at individual loci caused by the Bulmer effect. Exploiting a separation of the timescales over which stabilizing selection generates LD (rapidly) and changes allele frequencies (slowly; Fig. S1), we obtain simple expressions for the expected change in an allele's frequency across a single generation under stabilizing selection, as a function of its individual effect, the strength of stabilizing selection on the trait, the trait's variance and heritability, and the linkage relations among loci in the genome. As we show, the expressions that we derive predict allele-frequency change in simulations more accurately than other expressions that have commonly been used.

There is a deep theoretical literature on the population genetics of stabilizing selection, and some of the conclusions that we reach echo results from earlier work (see especially Lande 1975, 1977; Turelli and Barton 1990, 1994). For example, assuming that the distribution of allelic effects on a trait under stabilizing selection is Gaussian, Lande (1975) and Turelli and Barton (1990) find that that the linkage relations among loci do not affect the trait's additive genetic variance in equilibrium. This can be understood as reflecting a balance between two consequences of the stronger negative LD that accumulates under stabilizing selection with tighter linkage. First, that the stronger negative LD, as one component of the trait's genetic variance, directly reduces the genetic variance by a greater amount. Second, that the stronger negative LD slows down the allele-frequency dynamics at individual loci more severely, allowing the rarer allele at each locus to be maintained at a higher mutation selection balance in expectation, thus increasing the other component of the trait's additive genetic variance: the variance contributed by individual loci (the 'genic variance').

Previous treatments have usually considered the impact of the Bulmer effect on aggregate quantities such as the genetic and phenotypic variance of the trait under selection. However, for many applications, it is important to characterize the per-locus allele-frequency dynamics themselves. For example, the joint distribution of allele frequencies and effect sizes is an important summary of a trait's genetic architecture, and, under stabilizing selection, is determined by the allele-frequency dynamics at individual loci (Simons et al. 2018, 2022). More generally, given the abundance of GWAS data emerging for human traits, it is important to develop theoretical frameworks for understanding them in a principled way.

2 Model

We consider an additive polygenic trait under stabilizing selection. Genetic variation in the trait is underlain by L polymorphic loci, $l = 1, 2, \ldots, L$. At each locus l, there is a trait-increasing allele, with frequency p_l and haploid effect $+\alpha_l/2$, and a trait-decreasing allele, with frequency $1 - p_l$ and haploid effect $-\alpha_l/2$ (so that the difference in the expected phenotypes of a homozygote for the trait-increasing allele and a homozygote for the trait-decreasing allele is, all else equal, $2\alpha_l$). Since we are interested in characterizing allele-frequency dynamics at these loci, we ignore mutation at alternative causal loci (although we later discuss the implications of our results for the rate of turnover of the set of polymorphic loci underlying genetic variation in a trait).

An individual's trait value Y is given by

$$Y = G + E,$$

where

$$G = \sum_{l=1}^{L} \alpha_l g_l$$

is the individual's additive genetic value for the trait, with their genotype g_l coded as -1, 0, or 1 if the individual carries 0, 1, or 2 trait-increasing alleles at locus l respectively. E is an environmental disturbance that we assume to have mean zero and to be independent of G.

We denote the phenotypic variance by $V_P = \text{Var}(Y)$, the additive genetic variance by $V_G = \text{Var}(G)$, and the environmental variance by $V_E = \text{Var}(E)$. The trait's heritability is $h^2 = V_G/V_P$, and its genic variance—the additive genetic variance ignoring the contribution of linkage disequilibrium (LD) among loci—is $V_g = \sum_{l=1}^{L} 2p_l(1 - p_l)\alpha_l^2$.

The trait is under stabilizing selection around an optimal value of 0: the (relative) fitness of an individual with trait value Y is specified by the Gaussian fitness function

$$\phi(Y) = e^{-\frac{Y^2}{2V_S}}, \tag{1}$$

where V_S modulates the strength of stabilizing selection on the trait (with smaller values of V_S corresponding to stronger selection). It will sometimes be useful to approximate this Gaussian fitness function by a Taylor expansion around the optimal value of zero,

$$\phi(Y) \approx 1 - \frac{Y^2}{2V_S}. \tag{2}$$

We assume that mating is random within the population and that the trait is not affected by any forms of selection other than the stabilizing selection specified above.

Under these assumptions, upon the onset of selection, the mean trait value in the population rapidly converges to the optimal value of zero (e.g., Lande 1976; Hayward and Sella 2022). We therefore assume that the population starts with a mean trait value equal to zero—with the allele frequencies p_l such that this is the case—and study the subsequent allele-frequency dynamics at causal loci.

3 Results

Since we are interested in the allele-frequency dynamics at individual loci, it will be useful to consider the marginal fitnesses of the two alleles at a given locus, that is, the average fitness of the individual carrying

a randomly chosen copy of the one allele versus the other. From these marginal fitnesses, we can calculate an 'effective selection coefficient' s_{eff} for one of the alleles—say, the minor allele. The expected change in frequency of this allele across a single generation is then $p(1-p)s_{\text{eff}}$, where p is its frequency in the earlier generation (Wright 1931).

We begin with a simple calculation of these marginal fitnesses under the fitness model described above, ignoring phenotypic variation from the environment and from loci other than the focal locus, and also ignoring LD between the focal locus and other loci. We will then add in these two ingredients in turn.

3.1   A simple one-locus calculation

Consider a focal locus segregating for a trait-increasing allele A, at frequency p and with effect size $+\alpha/2$, and a trait-decreasing allele a, at frequency $1-p$ and with effect size $-\alpha/2$ (we have omitted the locus subscript since we are considering only a single locus here).

A haploid instance of this locus contains allele A with probability p and allele a with probability $1-p$. The average genetic value at this haploid instance of the locus is therefore

$$p \times (+\alpha/2) + (1-p) \times (-\alpha/2) = (2p-1)\alpha/2.$$

Under stabilizing selection around an optimal trait value of zero, the average genome-wide genetic value must be zero. Therefore, considering the average genetic value of the focal haploid locus above, the genetic value of the rest of the genome (including the homologous instance of the locus) must be $-(2p-1)\alpha/2 = (1-2p)\alpha/2$.

Now suppose that the allele at the focal haploid locus is A. If we ignore any correlation between the allelic state at the locus and the genetic value of the rest of the genome—i.e., if we ignore LD—the average genetic value of the individual carrying this A allele is

$$\underbrace{+\alpha/2}_{\substack{\text{contribution of} \\ \text{the allele itself}}} + \underbrace{(1-2p)\alpha/2}_{\substack{\text{average contribution} \\ \text{of rest of genome}}} = (1-p)\alpha. \tag{3}$$

If the allele at the focal haploid locus is instead a, the average genetic value of the individual is

$$\underbrace{-\alpha/2}_{\substack{\text{contribution of} \\ \text{the allele itself}}} + \underbrace{(1-2p)\alpha/2}_{\substack{\text{average contribution} \\ \text{of rest of genome}}} = -p\alpha. \tag{4}$$

Thus, the average phenotypes inhabited by A and a alleles differ in sign. If $p < 1/2$, that is, if A is the minor allele at the locus, the average phenotype inhabited by an A allele will be further from zero than the average phenotype inhabited by an a allele (and vice versa if $p > 1/2$). This explains why the minor allele is selected against under stabilizing selection: for the average value of the phenotype to be at its optimum, the rest of the genome must adapt predominantly to the more common allele at the locus, leaving the rarer allele maladapted to the rest of the genome.[1]

____

[1]As an interesting aside, we may use the same method to calculate the average phenotypes experienced by the three possible diploid genotypes at the focal locus, and therefore to rank the fitnesses of the three genotypes. The mean phenotypic effect of the diploid locus is twice the mean haploid effect, i.e., $(2p-1)\alpha$, and so mean phenotypic effect of all other loci is $(1-2p)\alpha$. Therefore, again ignoring systematic signed LD among causal loci, the mean phenotype of the aa genotype is $-\alpha + (1-2p)\alpha = -2p\alpha$, that of the Aa genotype is $0 + (1-2p)\alpha = (1-2p)\alpha$, and that of the AA genotype is $+\alpha + (1-2p)\alpha = 2(1-p)\alpha$. Assume that A is the minor allele at the locus, i.e., that $p < 1/2$. Then, if $p < 1/4$,

If we further ignore the phenotypic variation contributed by the environment and from the rest of the genome, then we can substitute the average phenotypic values above into our fitness function to calculate the marginal fitnesses of A and a. Employing the quadratic approximation in Eq. (2), the mean fitness of the bearer of a randomly chosen A allele is

$$w_A = 1 - \frac{(1-p)^2\alpha^2}{2V_S},$$

while the mean fitness of the bearer of a randomly chosen a allele is

$$w_a = 1 - \frac{p^2\alpha^2}{2V_S}.$$

Since the mean fitness in the population $\bar{w}$ is close to 1, the effective selection coefficient on A is

$$s_A^{\text{eff}} = \frac{w_A - w_a}{\bar{w}} \approx w_A - w_a = \frac{\left[p^2 - (1-p)^2\right]\alpha^2}{2V_S^2} = \left(p - \frac{1}{2}\right)\frac{\alpha^2}{V_S},$$

which is positive if $p > 1/2$ (when A is the common allele at the locus) and negative if $p < 1/2$ (when A is the rare allele at the locus).

The expected change in frequency of A across a single generation is then

$$\mathbb{E}\left[\Delta p\right] = p(1-p)s_A^{\text{eff}} = p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha^2}{V_S}, \tag{5}$$

which is a commonly used formula for allele-frequency change under stabilizing selection (e.g., Lande 1976; Simons et al. 2018).

However, this formula overpredicts the rate of allele-frequency change in simulations (Fig. 1). This is for two reasons. First, phenotypic variation from the environment and from other loci obscures the signal that selection sees of each allele's average phenotype. Second, the alleles come into positive LD with opposite-effect alleles elsewhere in the genome, bringing the average phenotypes of their bearers closer to zero and thus weakening selection at the locus. We consider these two effects in turn.

3.2   Background phenotypic variance

We assume that the trait value is normally distributed with mean zero and variance V_P. Still ignoring LD for now, the distribution of trait values experienced by A alleles is then normally distributed with mean $(1-p)\alpha$ (from Eq. 3) and variance $\sim V_P$ (because, for a highly polygenic trait, the genetic variance at the individual haploid locus is small compared to the overall phenotypic variance). Similarly, the distribution of trait values experienced by a alleles is normally distributed with mean $-p\alpha$ (from Eq. 4) and variance $\sim V_P$.

In general, if the trait distribution among the bearers of a given allele is normal with mean μ and variance V_P, then the average fitness of the bearer of a randomly chosen copy of the allele is

$$w(\bar{y} = \mu) = e^{-\frac{\mu^2}{2(V_S + V_P)}}, \tag{6}$$

$|-2p\alpha| < |(1-2p)\alpha| < |2(1-p)\alpha|$, i.e., the fitness ranking of the genotypes is $w(aa) > w(Aa) > w(AA)$. If, instead, $1/4 < p < 1/2$, then $|(1-2p)\alpha| < |-2p\alpha| < |2(1-p)\alpha|$, i.e., the fitness ranking is $w(Aa) > w(aa) > w(AA)$. In neither case is the heterozygous genotype the least fit, and in fact it is the fittest genotype when the minor-allele frequency is greater than 1/4 (Robertson 1956). Therefore, although the frequency dynamics at the locus (Eq. 5) resemble those expected under classical underdominance at the locus, the fitness values of the genotypes at the locus do not.

as shown in Lande (1976), Simons et al. (2018), and Appendix A1. The effect of accounting for background variance in the trait is simply to dilute the strength of selection at the focal locus by a factor $1 + V_P/V_S$, which is greater if the phenotypic variance is large relative to the width of the selection function.

Therefore, in our case, the mean relative fitness of the bearer of a randomly chosen A allele is

$$w_A = w(\bar{y} = (1-p)\alpha) = e^{-\frac{(1-p)^2\alpha^2}{2(V_S+V_P)}} \approx 1 - \frac{(1-p)^2\alpha^2}{2(V_S + V_P)}, \tag{7}$$

while the mean relative fitness of the bearer of a randomly chosen a allele is

$$w_a = w(\bar{y} = -p\alpha) = e^{-\frac{p^2\alpha^2}{2(V_S+V_P)}} \approx 1 - \frac{p^2\alpha^2}{2(V_S + V_P)}, \tag{8}$$

where the approximations hold because $\alpha^2 \ll V_S + V_P$ for a polygenic trait. Since the overall mean fitness is close to 1, the effective selection coefficient on A is

$$s_A^{\text{eff}} = \frac{w_A - w_a}{\bar{w}} \approx w_A - w_a = \frac{p^2\alpha^2}{2(V_S + V_P)} - \frac{(1-p)^2\alpha^2}{2(V_S + V_P)} = \left(p - \frac{1}{2}\right)\frac{\alpha^2}{V_S + V_P}, \tag{9}$$

and so the expected change in frequency of A across a single generation is

$$\mathbb{E}\left[\Delta p\right] = p(1-p)s_A^{\text{eff}} = p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha^2}{V_S + V_P}. \tag{10}$$

This prediction of allele-frequency change is smaller than the prediction of Eq. (5) by a factor $V_S/(V_S+V_P)$, but, while more accurate than Eq. (5), Eq. (10) still overpredicts the rate of allele-frequency change in

Figure 1: Simulated versus predicted average minor-allele frequencies across the polymorphic loci affecting a trait under stabilizing selection, in the case of full symmetry across loci. In the simulations, there are $L = 1{,}000$ polymorphic loci, all unlinked and with equal effect sizes ($\alpha = 1$) and starting minor-allele frequencies ($\min\{p_l, 1 - p_l\} = 0.2$). The loci are initially in linkage equilibrium, and the strength of selection V_S is chosen such that $V_S/V_G = 5$ initially. Further simulation details can be found in the Methods. Eq. (25) (dashed blue line), which takes into account background phenotypic variance and the Bulmer effect, is seen to be a better prediction of the observed trajectory of average minor-allele frequency (solid black line) than either Eq. (5) (dashed pink line), which ignores both background phenotypic variance and the Bulmer effect, and Eq. (10) (dashed yellow line), which ignores the Bulmer effect. The 'simulation' trajectory is averaged over 1,000 replicate trials.

simulations (Fig. 1). As we show next, this is because the mean phenotypes of bearers of A and a are in fact closer to zero than predicted by Eqs. (3) and (4), because A and a have each come into LD with opposite-effect alleles elsewhere in the genome.

3.3   The Bulmer effect

As we have discussed, stabilizing selection reduces a trait's genetic variance in two ways. The first is that, by favoring compensating combinations of trait-increasing and trait-decreasing alleles, stabilizing selection generates negative LD between alleles with the same directional effect on the trait. This leads to a rapid decline in the trait's genetic variance, to a quasi-equilibrium value that reflects a balance between the generation of LD by selection and its destruction by recombination (Fig. S1). The second is that stabilizing selection induces selection against rare alleles affecting the trait, eroding heterozygosity on average at their loci. For a highly polymorphic trait, selection against rare alleles is weak, and so their average frequency decline is slow.

In the limit of high polygenicity and a large population size, allele frequencies do not change at all under selection, and so the reduction in a trait's genetic variance due to stabilizing selection is entirely due to LD. In this limit, and under some other simplifying assumptions including that all loci are unlinked, Bulmer (1971) calculated the equilibrium reduction in the genetic variance of a trait under stabilizing selection, as a function of the trait's initial genetic variance in the absence of LD (its genic variance, V_g), the strength of stabilizing selection on the trait (V_S) and the contribution of the environment to trait variance (V_E). Bulmer (1974) extended this calculation to allow for variable linkage among loci.

An allele's LD with causal alleles elsewhere in the genome will affect the mean trait value experienced by the allele, and therefore also its frequency dynamics under selection. To calculate how the LD generated by stabilizing selection affects allele-frequency dynamics at individual loci, we first determine how LD in general affects the mean trait value experienced by an allele. Thereafter, we calculate the total degree of LD generated by stabilizing selection in equilibrium, and the expected apportionment of this overall amount of LD to specific pairs of loci. Exploiting the difference in the timescales over which LD is generated and rare alleles decline in frequency under stabilizing selection (Fig. S1), we then substitute the equilibrium degree of LD into our calculation for how LD affects the mean trait value experienced by an allele to find the expected effect of LD on allele-frequency dynamics under stabilizing selection. The agreement between our analytical predictions and our simulations validates this separation-of-timescales approximation for polygenic traits.

3.3.1   The effect of LD on the mean trait value experienced by an allele

Recall that there are L loci underlying genetic variation in the trait, and that the trait-increasing allele at locus l is at frequency p_l and increases the value of the trait by $\alpha_l/2$, while the trait-decreasing allele is at frequency $1 - p_l$ and decreases the trait value by $\alpha_l/2$. We denote the coefficient of LD between the trait-increasing alleles at loci l and l' by $D_{ll'}$. Since the population mean value of the trait is, by assumption, at its optimum, 0,

$$\sum_{l=1}^{L}\left[p_l \times \left(+\frac{\alpha_l}{2}\right) + (1 - p_l) \times \left(-\frac{\alpha_l}{2}\right)\right] = 0 \quad \Rightarrow \quad \sum_{l=1}^{L}(2p_l - 1)\alpha_l = 0. \tag{11}$$

Define the random variable I_l to take the value 1 if the allele at locus l on a randomly chosen haploid/gametic genome is trait-increasing, and 0 if the allele is instead trait-decreasing. From Bayes'

theorem, these indicator variables are related to the coefficients of LD via

$$\text{Prob}(I_{l'} = 1 \mid I_l = 1) = p_{l'} + D_{ll'}/p_l. \tag{12}$$

Consider a copy of the trait-increasing allele at l. If it is maternally inherited, then, under random mating, the mean genetic value of the paternally inherited genome is 0, and so the mean trait value experienced by the allele is the mean genetic value of the maternally inherited genome. Similarly, if it is paternally inherited, the mean trait value it experiences is the mean genetic value of the paternally inherited genome. Therefore,

$$\mathbb{E}[Y \mid I_l = 1] = \underbrace{\left(+\frac{\alpha_l}{2}\right)}_{\substack{\text{contribution of} \\ \text{trait-incr. allele} \\ \text{at focal locus } l}} + \underbrace{\sum_{l' \neq l} \left[\text{Prob}(I_{l'} = 1 \mid I_l = 1)\left(+\frac{\alpha_{l'}}{2}\right) + [1 - \text{Prob}(I_{l'} = 1 \mid I_l = 1)]\left(-\frac{\alpha_{l'}}{2}\right) \right]}_{\substack{\text{average contribution of rest of (haploid) genome,} \\ \text{conditional on having trait-increasing allele at } l}}$$

$$= \frac{1}{2}\left(\alpha_l + \sum_{l' \neq l} \left[2\text{Prob}(I_{l'} = 1 \mid I_l = 1) - 1 \right]\alpha_{l'} \right)$$

$$= \frac{1}{2}\left(\alpha_l + \sum_{l' \neq l} \left[2p_{l'} - 1 + \frac{2D_{ll'}}{p_l} \right]\alpha_{l'} \right)$$

$$= \frac{1}{2}\left(2(1 - p_l)\alpha_l + \sum_{l'=1}^{L} \left[2p_{l'} - 1 \right]\alpha_{l'} + \sum_{l' \neq l} \frac{2D_{ll'}\alpha_{l'}}{p_l} \right)$$

$$= (1 - p_l)\alpha_l + \sum_{l' \neq l} \frac{D_{ll'}\alpha_{l'}}{p_l}$$

$$= (1 - p_l)\left(\alpha_l + \frac{1}{p_l(1 - p_l)} \sum_{l' \neq l} D_{ll'}\alpha_{l'} \right). \tag{13}$$

Sensibly, positive LD with trait-increasing alleles elsewhere in the genome ($D_{ll'} > 0$) will tend to increase the trait value experienced by the trait-increasing allele at l, while negative LD with other trait-increasing alleles ($D_{ll'} < 0$) will tend to decrease the trait value it experiences. With no LD ($D_{ll'} = 0$), $\mathbb{E}[Y \mid I_l] = (1 - p_l)\alpha_l$, recovering Eq. (3) above.

By a similar calculation, the average trait value experienced by a randomly chosen trait-decreasing allele at l is

$$\mathbb{E}[Y \mid I_l = 0] = -p_l\left(\alpha_l + \frac{1}{p_l(1 - p_l)} \sum_{l' \neq l} D_{ll'}\alpha_{l'} \right). \tag{14}$$

3.3.2 Symmetric loci

We now calculate the expected coefficients of LD between a focal locus and all of the other loci that affect the trait under stabilizing selection. We begin with the simplest possible case, where all loci are unlinked and have the same minor-allele frequency p and effect size α. In this case, the mean phenotype

experienced by a randomly chosen trait-increasing allele at locus l is, from Eq. (13),

$$\mathbb{E}[Y \mid I_l = 1] = (1-p)\alpha\left(1 + \frac{1}{p(1-p)}\sum_{l' \neq l} D_{ll'}\right), \tag{15}$$

while the mean phenotype experienced by a randomly chosen trait-decreasing allele at l is, from Eq. (14),

$$\mathbb{E}[Y \mid I_l = 0] = -p\alpha\left(1 + \frac{1}{p(1-p)}\sum_{l' \neq l} D_{ll'}\right); \tag{16}$$

that is, LD between the focal locus and other causal loci in the genome causes the mean phenotypes experienced by both alleles at the focal locus to be multiplied by a factor $1 + \frac{1}{p(1-p)}\sum_{l' \neq l} D_{ll'}$, as if the effect size at the locus were not α but instead $\alpha\left(1 + \frac{1}{p(1-p)}\sum_{l' \neq l} D_{ll'}\right)$.

Under the same conditions of symmetry across loci, and in the limit of high polygenicity, the equilibrium reduction d in the trait's genetic variance due to the Bulmer effect satisfies

$$\frac{d}{V_g} = \frac{3 + X - \sqrt{1 + 6X + X^2}}{4}, \tag{17}$$

where $X = (V_S + V_E)/V_g$ (Bulmer 1971; Veller and Coop 2024, S1 Text Section S3.4). (Note that, for neatness, we define d to be the *reduction* in the genetic variance, so that it is a positive number; this is in contrast to Bulmer's d, which is the net change in the genetic variance and therefore negative.) When $V_S \gg V_g$, $d/V_g \approx V_g/(V_S + V_E)$ (Turelli 1988).

While Eq. (17) gives the total reduction in the trait's genetic variance explicitly in terms of the model parameters V_S, V_E, and V_g (which don't themselves depend on d), in practice we will usually not have access to measurements of a trait's genic variance V_g (since we cannot characterize all of the loci that contribute variance to the trait). Fortunately, the equilibrium value of d/V_g can also be expressed implicitly in terms of the heritability of the trait, h^2, and its variance, V_P (which both depend on d), as well as V_S:

$$\frac{d}{V_g} = \frac{h^2}{1 + h^2 + V_S/V_P}, \tag{18}$$

where h^2 and V_P are at their quasi-equilibrium values (see Veller and Coop 2024, S1 Text, Section 3.4.3). Eq. (18) obviates the need to estimate V_g.

The reduction d is entirely due to LD among the polymorphic loci that underlie genetic variation in the trait:

$$-d = 2\alpha^2 \sum_{l=1}^{L} \sum_{\substack{l'=1 \\ l' \neq l}}^{L} D_{ll'} \quad \Rightarrow \quad \sum_{l=1}^{L} \sum_{\substack{l'=1 \\ l' \neq l}}^{L} D_{ll'} = -\frac{d}{2\alpha^2}. \tag{19}$$

Under the assumption of symmetry across loci in their effects and minor-allele frequencies, this total sum of LD is apportioned equally, in expectation, across the $L(L-1)/2$ pairs of loci: for each pair of loci l and l',

$$\mathbb{E}[D_{ll'}] = -\frac{d}{2\alpha^2 L(L-1)}. \tag{20}$$

(Note that the summations in Eq. (19) count each pair of loci twice.) So, for a given causal locus l, the sum of its LD coefficients with the other $L - 1$ causal loci is, in expectation,

$$\mathbb{E}\left[\sum_{\substack{l'=1 \\ l' \neq l}}^{L} D_{ll'}\right] = -\frac{d}{2\alpha^2 L}. \tag{21}$$

Substituting Eq. (21) into Eqs. (15) and (16), we find that the mean trait value of a randomly chosen trait-increasing allele at locus l is

$$\mathbb{E}[Y \mid I_l = 1] = (1-p)\alpha \left(1 - \frac{d}{2p(1-p)\alpha^2 L}\right)$$
$$= (1-p)\alpha \left(1 - \frac{d}{V_g}\right) \tag{22}$$

and, similarly, that the mean trait value of a randomly chosen trait-decreasing allele at locus l is

$$\mathbb{E}[Y \mid I_l = 0] = -p\alpha \left(1 - \frac{d}{V_g}\right). \tag{23}$$

Comparing Eqs. (22) and (23) with Eqs. (3) and (4), we see that the mean phenotypes of the two alleles at the locus are as they would be in the simple scenario considered before with no LD, but with the effect sizes of the alleles attenuated by a factor $1 - d/V_g$ (with d/V_g specified by Eq. 17). That is, the LD generated by stabilizing selection can be incorporated into the formula for allele-frequency change at the locus that ignores LD, Eq. (10), by defining an 'effective' effect size,

$$\alpha_{\text{eff}} = \alpha \left(1 - \frac{d}{V_g}\right). \tag{24}$$

The expected change in frequency across a single generation of the trait-increasing allele at the focal locus is then

$$\mathbb{E}[\Delta p] = p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha_{\text{eff}}^2}{V_S + V_P} = p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha^2\left(1 - \frac{d}{V_g}\right)^2}{V_S + V_P}. \tag{25}$$

(Note that the phenotypic variances experienced by the trait-increasing and trait-decreasing alleles at l are still equal, and close to V_P owing to the trait's high polygenicity.)

In simulations of stabilizing selection under the conditions of symmetry across loci assumed here, Eq. (25) is seen to be a better predictor of the allele-frequency dynamics at causal loci than either Eq. (5) or Eq. (10) (Fig. 1).

3.3.3 Incorporating linkage

The calculations above assume full symmetry across loci, and, in particular, that every locus is on a separate chromosome. This allowed us to apportion the total amount of LD generated by stabilizing selection evenly, in expectation, among the locus pairs.

We now consider the case of variable linkage relations among loci, with some locus pairs lying on the same chromosome and some lying on different chromosomes. Let the recombination fraction between loci l and l' be $r_{ll'}$. For now, we maintain the assumption of constant effect sizes and minor-allele frequencies

across loci. Variable linkage relations among loci affect the total amount of LD generated by stabilizing selection: In the limit of high polygenicity, the equilibrium reduction d in the trait's genetic variance due to the Bulmer effect satisfies

$$\frac{d}{V_g} = \frac{1 - \bar{r}_h \left(\sqrt{1 + 2 \left(1 + \frac{1}{\bar{r}_h}\right) X + X^2} - (1 + X) \right)}{1 + 2\bar{r}_h},$$ (26)

where $X = (V_S + V_E)/V_g$ as before, and $\bar{r}_h = L(L-1)/\left(\sum_l \sum_{l' \neq l} \frac{1}{r_{ll'}}\right)$ is the harmonic mean recombination rate across loci (Bulmer 1974; Veller and Coop 2024, S1 Text Section S3.4). $\bar{r}_h$ can be estimated from various kinds of data, including linkage maps, cytological data, and sequence data, similarly to the arithmetic mean recombination rate $\bar{r} = \sum_l \sum_{l' \neq l} r_{ll'}/L(L-1)$ (Veller et al. 2019). When $V_S \gg V_g$, $d/V_g \approx \frac{1}{2\bar{r}_h} \cdot \frac{V_g}{V_S + V_E}$.

As in the case of no linkage, in practice we do not need to estimate X, and in particular V_g, to estimate the equilibrium value of d/V_g—this quantity can instead be expressed implicitly in terms of V_P and h^2 (which depend on d), as well as V_S and $\bar{r}_h$:

$$\frac{d}{V_g} = \frac{h^2}{2\bar{r}_h(1 + V_S/V_P) + h^2},$$ (27)

where, again, h^2 and V_P are taken to be at their quasi-equilibrium values (see Veller and Coop 2024, S1 Text, Section 3.4.3).

As before, the reduction d is entirely due to LD among loci, with $\sum_l \sum_{l' \neq l} D_{ll'} = -d/(2\alpha^2)$. However, now, this overall LD is not apportioned equally among locus pairs in expectation. Instead, the expected LD between l and l' is proportional to the inverse of the recombination fraction between the loci, $1/r_{ll'}$ (Bulmer 1974), so that

$$\mathbb{E}[D_{ll'}] = -\frac{d}{2\alpha^2 L(L-1)} \cdot \frac{\bar{r}_h}{r_{ll'}}.$$ (28)

For a given locus l, let $\bar{r}_h^{(l)} = (L-1)/\sum_{l' \neq l} (1/r_{ll'})$ be the harmonic mean recombination fraction between l and the $L-1$ other causal loci. Then, summing Eq. (28) across these $L-1$ other loci, we find

$$\mathbb{E}\left[\sum_{\substack{l'=1 \\ l' \neq l}}^{L} D_{ll'}\right] = -\frac{d\bar{r}_h}{2\alpha^2 L(L-1)} \sum_{l' \neq l} \frac{1}{r_{ll'}} = -\frac{d}{2\alpha^2 L} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}}.$$ (29)

Because of our assumption of equal effect sizes and minor-allele frequencies across loci, Eqs. (15) and (16) still specify the mean trait values experienced by a randomly selected trait-increasing and trait-decreasing allele at l respectively. Substituting Eq. (29) into (15) and (16), we find

$$\mathbb{E}[Y \mid I_l = 1] = (1-p)\alpha \left(1 - \frac{d}{2p(1-p)\alpha^2 L} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right)$$

$$= (1-p)\alpha \left(1 - \frac{d}{V_g} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right)$$ (30)

and, similarly,

$$\mathbb{E}[Y \mid I_l = 0] = -p\alpha \left(1 - \frac{d}{V_g} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right).$$ (31)

Therefore, again, comparing the equations above with those for the no-LD case (Eqs. 3 and 4), we can think of the impact of the LD generated by stabilizing selection on the mean phenotypes experienced by the alleles at a particular locus in terms of an attenuation of the 'effective' effect size at the locus, with this 'effective' effect size dependent on the recombination relations of the locus to other causal loci, according to

$$\alpha_{\text{eff}}^{(l)} = \alpha \left(1 - \frac{d}{V_g} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}} \right). \tag{32}$$

Intuitively, alleles at loci that have tighter recombination relations with other loci $(\bar{r}_h/\bar{r}_h^{(l)} > 1)$ will develop stronger average LD with the opposite-effect alleles at these other loci, and will therefore have their individual effects more greatly masked—and so their frequency dynamics more severely slowed—by the Bulmer effect than alleles at loci with looser recombination relations with other loci $(\bar{r}_h/\bar{r}_h^{(l)} < 1)$.

The expected change in frequency of the minor allele at l across a single generation is

$$\mathbb{E}\left[\Delta p_l \right] = p_l(1 - p_l) \left(p_l - \frac{1}{2} \right) \frac{\left(\alpha_{\text{eff}}^{(l)} \right)^2}{V_S + V_P} = p_l(1 - p_l) \left(p_l - \frac{1}{2} \right) \frac{\alpha^2 \left(1 - \frac{d}{V_g} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}} \right)^2}{V_S + V_P}. \tag{33}$$

Taking the average of Eq. (33) provides a prediction of the average across loci of the change in the minor-allele's frequency. This prediction agrees well with simulations that make use of realistic linkage maps (Fig. 2). In the case of a low-recombination species especially (*Drosophila melanogaster*; $\bar{r}_h \approx 0.07$), in which the LD generated by the Bulmer effect is expected to be particularly strong, Eq. (33) offers a much improved prediction of the allele-frequency dynamics over Eqs. (5) and (10) (Fig. 2B).

We can also check the prediction of Eq. (33) for loci with lower and higher locus-specific harmonic mean recombination rates $\bar{r}_h^{(l)}$. Collecting loci in bins of gradually higher values of $\bar{r}_h^{(l)}$ across the *D. melanogaster* linkage map (which, unlike in high-recombination species like humans, shows a sizeable range of $\bar{r}_h^{(l)}$ values), Fig. 3 plots the average minor-allele frequency in each bin after 250 generations of selection against

Figure 2: Average change in the minor-allele frequency at polymorphic loci affecting a trait under stabilizing selection, when these loci are distributed across the human linkage map (A) and the linkage map of *Drosophila melanogaster* (B). Eq. (33) is seen to predict simulated trajectories of the average minor-allele frequency better than Eqs. (5) and (10), especially for *D. melanogaster*, a low-recombination species. Other than the variable linkage relations among loci, simulation details are identical to Fig. 1.

the average value of $\bar{r}_h^{(l)}$ within each bin. While the prediction of Eq. (33) is a substantial improvement over predictions that do not take into account LD (and therefore also ignore variable linkage relations among loci), Eq. (33) underpredicts the degree of allele-frequency change for the lowest-recombination bins.

The reason has to do with the separation-of-timescales assumption underlying Eq. (33), and in particular the assumption that the quasi-equilibrium degree of LD (Eq. 28) is attained instantly for every pair of loci at the onset of selection. In reality, the expected degree of LD between each pair of loci builds up over time, approaching its equilibrium value at a rate that depends on the recombination fraction between the pair of loci involved: very rapidly for loosely linked locus pairs, but more slowly for tightly linked locus pairs. Therefore, for the most tightly linked locus pairs, the Bulmer effect is, for an appreciable number of generations after the onset of selection, weaker than assumed by Eq. (33), and so the allele-frequency dynamics at these loci are faster in these early generations than predicted by our calculations. The result is that, after a given number of generations, allele frequencies have not changed as much as predicted by Eq. (33), with the disparity greater for more tightly-linked loci—the pattern observed in Fig. 3.

We can, in fact, for every pair of loci, calculate the expected LD in each generation after the onset of selection, and use these value to obtain a more accurate—albeit more complicated—prediction for the overall allele-frequency change at each locus after a given number of generations. Initially, the degree of

Figure 3: Minor-allele frequencies after 250 generations of selection, for loci with different average recombination rates with the rest of the genome. The simulations here are the same as for Fig. 2B, with loci distributed along the *D. melanogaster* linkage map. For the simulated minor-allele frequencies, loci have been binned according to their locus-specific harmonic mean recombination fractions with other loci, $\bar{r}_h^{(l)}$. While the equilibrium-based prediction (Eq. 33) that takes into account the Bulmer effect (blue line) is a substantial improvement over predictions that do not take into account the Bulmer effect (Eqs. 5 and 10; pink and yellow lines), it underpredicts the degree of allele-frequency change, especially for loci with low values of $\bar{r}_h^{(l)}$, since for these values, the approach to equilibrium values of LD is slow. A prediction based on the full sequence of non-equilibrium values of LD (Eq. 34; details in Methods) performs better than the equilibrium-based prediction. Note that the fourth and fifth bins contain only 6 and 2 loci respectively, potentially explaining the discrepancy between the simulated minor-allele frequencies and the predictions based on Eq. (34). Simulation points are averages across 500 replicate trials.

LD between each pair of loci l and l' is zero: $D_{ll'}^{(0)} = 0$. Within each generation t, selection generates an increment to the degree of disequilibrium between l and l' equal to $2K$ (we will determine the value of K shortly). This amount $2K$ is the same in expectation for all pairs of loci, because we have assumed that effect sizes are constant, and is divided equally in expectation between cis-LD (LD among pairs of alleles inherited from the same parent) and trans-LD (LD among alleles inherited from different parents). In transmission to the next generation $t + 1$, recombination between loci l and l' generates an amount $r_{ll'}K$ of new cis-LD from the trans-LD that built up in generation t, while an amount $(1 - r_{ll'})K$ of the cis-LD that built up in generation t is preserved in transmission to the generation $t + 1$, along with an amount $(1 - r_{ll'})D_{ll'}^{(t)}$ of the cis-LD that was already present at the beginning of generation t (before selection acted). Therefore,

$$\mathbb{E}\left[D_{ll'}^{(t+1)} \mid D_{ll'}^{(t)}\right] = r_{ll'}K + (1 - r_{ll'})K + (1 - r_{ll'})D_{ll'}^{(t)} = K + (1 - r_{ll'})D_{ll'}^{(t)}. \tag{34}$$

Starting with zero LD ($D_{ll'}^{(0)} = 0$), Eq. (34) defines a sequence of expected LD values

$$D_{ll'}^{(1)} = K, \quad D_{ll'}^{(2)} = \left(1 + (1 - r_{ll'})\right)K, \quad D_{ll'}^{(3)} = \left(1 + (1 - r_{ll'}) + (1 - r_{ll'})^2\right)K, \quad \ldots, \tag{35}$$

which converges at rate $r_{ll'}$ to an asymptotic value of $K/r_{ll'}$ (Bulmer 1974). This is the equilibrium degree of LD between l and l'; comparison with Eq. (28) therefore reveals that $K = -d\bar{r}_h/2\alpha^2 L(L - 1)$.

For each polymorphic locus l affecting the trait, Eq. (35) thus specifies the expected LD with every other polymorphic locus l' affecting the trait, and so, for each generation t after the onset of selection, we can substitute these into Eqs. (13) and (14) to calculate the mean trait value experienced by the trait-increasing and trait-decreasing alleles at l; from these, we can calculate the expected change in frequency of the alleles at locus l from generation t to generation $t + 1$ (see Methods). Carrying out this calculation, we find that it predicts the cumulative allele-frequency change observed across generations in our simulations more successfully than Eq. (33), especially for loci with especially tight average linkage relations with the other loci in the genome (Fig. 3).

3.3.4 Incorporating variable effect sizes and minor-allele frequencies

We now allow for differences across loci in their effect sizes and minor-allele frequencies. At a focal locus l, the mean phenotypes of the trait-increasing and trait-decreasing alleles are still given by Eqs. (13) and (14), and in particular by the degrees of LD between l and other loci l'. However, now these degrees of LD depend on the effect sizes at l and l'.

Consider first the case where all loci are unlinked. In Appendix A2, we show that in this case the 'effective' effect size at locus l can be written

$$\alpha_{\text{eff}}^{(l)} = \alpha_l \left(1 - \frac{d}{2L\sqrt{p_l(1 - p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle}\right), \tag{36}$$

where d is the quasi-equilibrium reduction in the genetic variance due to the Bulmer effect (specified by Eq. 17) and the angle brackets in the denominator denote an average taken across all loci. The predicted single-generation change in allele frequency at locus l is then

$$\mathbb{E}\left[\Delta p\right] = p(1 - p)\left(p - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P}. \tag{37}$$

If minor-allele frequencies do not differ much across loci, $2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle \approx V_g$, so

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{V_g}\right), \tag{38}$$

as in the case of symmetric effects and minor-allele frequencies across loci (Eq. 24). Therefore, in this case,

$$\mathbb{E}\left[\Delta p\right] = p(1-p)\left(p - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P} \approx p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha^2\left(1 - \frac{d}{V_g}\right)^2}{V_S + V_P}, \tag{39}$$

as in Eq. (25). Eq. (39) accurately predicts the trajectory of the average minor-allele frequency in simulations that allow for variation in effect sizes and initial minor-allele frequencies across loci when all loci are unlinked from one another (Fig. S2).

We now consider the possibility of variable linkage relations among loci. We assume that the recombination fractions between loci are independent of their effect sizes, and that variation in recombination fractions has a relatively weak effect on variation in minor-allele frequencies. Under these assumptions, we show in Appendix A2 that the effective effect size of locus l is approximately

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right), \tag{40}$$

where d is the quasi-equilibrium reduction in the genetic variance due to the Bulmer effect (specified by Eq. 26). The predicted single-generation change in allele frequency at locus l is then

$$\mathbb{E}\left[\Delta p\right] = p(1-p)\left(p - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P}. \tag{41}$$

If minor-allele frequencies do not differ much across loci,

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{V_g}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right), \tag{42}$$

as in the case of symetric effects and minor-allele frequencies across loci (Eq. 32), and so

$$\mathbb{E}\left[\Delta p\right] = p(1-p)\left(p - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P} \approx p(1-p)\left(p - \frac{1}{2}\right)\frac{\alpha^2\left(1 - \frac{d}{V_g}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right)^2}{V_S + V_P}, \tag{43}$$

as in Eq. (33). Eq. (43) accurately predicts changes in minor-allele frequencies in simulations with variable effect sizes and initial minor-allele frequencies across loci, for both the *D. melanogaster* linkage map (Fig. S3) and the human map (Fig. S4).

4   Discussion

Understanding the processes that govern the genetic architecture of complex traits will require the interpretation of genomic data in terms of population genetic models. The richest and finest-scale data come

from genome-wide association studies, which offer estimates of allele effect sizes at thousands of sites throughout the genome (Abdellaoui et al. 2023). Because of the per-site nature of GWAS data, its interpretation in terms of population genetic models will require a detailed understanding of allele-frequency dynamics under these models.

Here, we have provided simple calculations that predict allele-frequency dynamics under stabilizing selection—a common mode of selection on complex traits (Sanjak et al. 2018; Sella and Barton 2019)—substantially more accurately than the formulae that have until now been used for this purpose. To do so, we have incorporated into these formulae the linkage disequilibrium that stabilizing selection rapidly generates between opposite-effect alleles throughout the genome (Bulmer 1971, 1974).

The accuracy of our calculations in predicting simulated allele-frequency trajectories under stabilizing selection (Figs. 1, 2, and 3) suggests that they may make possible more precise quantitative interpretation of GWAS and other genomic data in terms of population genetic models of stabilizing selection. Below, we discuss some of the implications of our results for the interpretation of such data.

4.1 Genetic architecture of complex traits

In an important series of papers, Simons et al. (2018, 2022) have shown that, for many human traits, the joint distribution of allele frequencies and effect sizes (as inferred from GWAS) is consistent with the allele-frequency dynamics expected under stabilizing selection. In demonstrating this consistency, Simons et al. (2018, 2022) made use of Eq. (5) as a description of the allele-frequency dynamics. As we have shown, Eq. (5) overpredicts the rate of allele-frequency change under stabilizing selection, because it ignores the systematically signed LD generated by stabilizing selection (and background phenotypic variance in the trait under selection). Importantly, however, the equations that we have derived for allele-frequency change that do take into account the LD generated by stabilizing selection are qualitatively of the same form as those that ignore this LD, with an 'effective' effect size of each allele substituted for the allele's true effect size (Eqs. 24,32,36,40). Therefore, the results of Simons et al. (2018, 2022) are not qualitatively affected by the more accurate predictions of allele-frequency change under stabilizing selection that we have derived. However, the estimates obtained by Simons et al. (2018, 2022) for the strength of stabilizing selection on human traits must be revised upwards, since for a given strength of stabilizing selection on a trait, the slowdown of the allele-frequency dynamics induced by the Bulmer effect will result in higher minor-allele frequencies, on average, across loci.

4.2 Allelic turnover and the portability of polygenic scores across populations

Once a GWAS has been carried out in a particular sample, it is common to use the effect-size estimates obtained from the GWAS to generate polygenic scores (PGSs)—sums of an individuals' genotypes across trait-associated loci, with each locus weighted in the sum by its estimated effect size—and to measure the accuracy of these PGSs as predictors of trait values both within the original GWAS sample and in other samples. Such exercises have revealed that PGSs are often substantially worse predictors of trait values in samples that are more distantly related to the original GWAS sample (e.g., Martin et al. 2019; Mostafavi et al. 2020; Ding et al. 2023).

Several explanations have been offered for this 'portability' problem, including differences between populations in (i) the effect sizes of causal alleles, (ii) the frequencies of causal alleles, (iii) the degree to which genotyped SNPs 'tag' causal alleles (via LD), and (iv) the environmental and genetic effects with which variation at genotyped SNPs is confounded. Of interest here is the second explanation, that PGSs can suffer reduced portability because of differences between populations in allele frequencies at

causal loci (Wang et al. 2020; Carlson et al. 2022; Yair and Coop 2022). In some cases, an ancestral polymorphism might have been retained in a sample drawn from one population but lost in a sample drawn from another; the locus would then contribute trait variation in the one sample but not in the other.

As Yair and Coop (2022) argue, although turnover of the loci underlying variation in a trait is possible by neutral drift alone, the process is accelerated—and the portability of PGS prediction across populations more rapidly degenerates—if the trait is under stabilizing selection, because stabilizing selection speeds up allele-frequency dynamics relative to neutral drift. Yair and Coop (2022) further recognized that the allele-frequency dynamics under stabilizing selection, and consequently the rate of allelic turnover between populations, would be slowed by the Bulmer effect (especially for ensembles of tightly linked-loci; see Fig S1 in Yair and Coop 2022). However, for simplicity, in quantifying the effect of stabilizing selection on the reduction of trait variance contributed by ancestral polymorphisms, and the consequent degeneration of PGS prediction accuracy across descendent populations, Yair and Coop (2022) ignored the contribution of LD generated by the Bulmer effect.

Since, as we have shown, the impact of the Bulmer effect on allele-frequency dynamics under stabilizing selection can be captured by defining 'effective' effect sizes of causal alleles, the Bulmer effect can be incorporated in the calculations of Yair and Coop (2022) in a simple way. For example, Eq. (2.3) in Yair and Coop (2022) calculates the fraction of trait variance in a generation-t descendent population explained by polymorphisms of effect size α that were present in the ancestral population: $\exp(-F_{ST}[1 + S/4])$, where $F_{ST} = t/2N$ is the value of F_{ST} between the ancestral and descendant populations at neutral loci and $S = 2N\alpha^2/V_S$. This calculation can be modified to take into account the Bulmer effect simply by replacing α with an 'effective' effect size α_{eff}, as defined in Eqs. (24), (32), (36), and (40). The same is true of the analogous calculations developed by Yair and Coop (2022) based on a diffusion approximation.

4.3 Other processes that generate long-range signed LD

While stabilizing selection itself generates systematically signed long-range LD between alleles that affect the trait under selection, with these LDs impacting the frequency dynamics of the alleles in the way we have described, other processes can also generate systematically signed LD, and these LDs too will influence allele-frequency dynamics under stabilizing selection.

Most important among these other processes is assortative mating, which is known to be common for traits in humans (Horwitz et al. 2023; Border et al. 2022a) and other animals (Jiang et al. 2013). Assortative mating for a heritable trait generates positive LD between alleles with like effects on the trait (Wright 1921; Crow and Felsenstein 1968). If the trait is also under stabilizing selection, this positive LD will tend to counteract the negative LD among like alleles generated by stabilizing selection, speeding up the rate at which the minor allele declines in frequency relative to the case where there is no assortment on the trait. Whether the frequency dynamics are ultimately faster or slower than expected in the absence of all LD depends on whether the impact of the negative LD generated by stabilizing selection is outweighed by, or outweighs, the impact of the positive LD generated by assortative mating. Our general calculation of the mean trait values experienced by the two alleles at a locus, as a function of their effects and their degrees of LD with other causal loci (Eqs. 13 and 14), allows the balance to be quantified.

As with the LD generated by stabilizing selection itself, the effect of LD generated by assortative mating on allele-frequency dynamics under stabilizing selection can be captured in 'effective' effect sizes of alleles at polymorphic loci affecting the trait. In contrast to the case of stabilizing selection, under assortative mating, the equilibrium degree of LD expected between a pair of loci does not depend on the recombination fraction between the loci (Crow and Felsenstein 1968). It can be calculated as a simple

function of the correlation of the trait among mates, the trait's heritability, and the effect sizes of the alleles involved (Crow and Felsenstein 1968; Border et al. 2022b; Veller and Coop 2024).

Suppose, for simplicity, that effect sizes α and minor-allele frequencies p are equal across loci. If the heritability of the trait is h^2 and the phenotypic correlation among mates is ρ, then the expected LD between the trait-increasing alleles at loci l and l' due to assortative mating is $D_{ll'}^{\text{assort}} = \frac{h^2\rho}{1-h^2\rho} \cdot \frac{H}{4(L-1)}$, where $H = 2p(1-p)$ is the heterozygosity at each locus (see SI Section S3.1.1 in Veller and Coop 2024). Therefore, for a given locus l, the expected sum of the LDs between its trait-increasing allele and the trait-increasing alleles at other loci, due to assortative mating, is

$$\mathbb{E}\left[\sum_{\substack{l'=1 \\ l'\neq l}}^{L} D_{ll'}^{\text{assort}}\right] = \frac{h^2\rho}{1-h^2\rho} \cdot \frac{H}{4} = \frac{h^2\rho}{1-h^2\rho} \cdot \frac{V_g}{4L\alpha^2}. \tag{44}$$

If, under both assortative mating and stabilizing selection, the LD between a pair of loci is simply the sum of the LDs expected under the two processes separately ($D_{ll'} = D_{ll'}^{\text{assort}} + D_{ll'}^{\text{stab}}$), then, if all loci are unlinked, we combine Eqs. (44) and (21) to find

$$\mathbb{E}\left[\sum_{\substack{l'=1 \\ l'\neq l}}^{L} D_{ll'}\right] = \mathbb{E}\left[\sum_{\substack{l'=1 \\ l'\neq l}}^{L} D_{ll'}^{\text{assort}}\right] + \mathbb{E}\left[\sum_{\substack{l'=1 \\ l'\neq l}}^{L} D_{ll'}^{\text{stab}}\right]$$

$$= \frac{h^2\rho}{1-h^2\rho} \cdot \frac{V_g}{4L\alpha^2} - \frac{d}{2L\alpha^2}$$

$$= \frac{V_g}{4L\alpha^2}\left(\frac{h^2\rho}{1-h^2\rho} - 2\frac{d}{V_g}\right), \tag{45}$$

where d/V_g is given by Eq. (17).

If we allow for variable linkage relations among loci, a similar calculation combining Eqs. (44) and (29) yields

$$\mathbb{E}\left[\sum_{\substack{l'=1 \\ l'\neq l}}^{L} D_{ll'}\right] = \frac{V_g}{4L\alpha^2}\left(\frac{h^2\rho}{1-h^2\rho} - 2\frac{d}{V_g} \cdot \frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right), \tag{46}$$

where d/V_g is now given by Eq. (26). On average across loci, therefore, $\mathbb{E}_l[\mathbb{E}[\sum_{l'\neq l} D_{ll'}]] = \frac{V_g}{4L\alpha^2}\left(\frac{h^2\rho}{1-h^2\rho} - 2\frac{d}{V_g}\right)$, as in the case with no linkage (though with a larger value of d owing to linkage).

Therefore, on average, the impact of stabilizing selection will outweigh that of assortative mating, the 'effective' effect sizes of alleles are attenuated from their true effects, and allele-frequency dynamics under stabilizing selection will be slowed if $2d/V_g > h^2\rho/(1-h^2\rho)$. In contrast, if $2d/V_g < h^2\rho/(1-h^2\rho)$, the impact of assortative mating outweighs that of stabilizing selection, the effective effect sizes of alleles are amplified, and allele-frequency dynamics under stabilizing selection are accelerated.

As an example, human height is under strong assortative mating ($\rho \sim 0.25$; Stulp et al. 2017) and moderately strong stabilizing selection ($V_S/V_P \sim 20\text{--}30$; Stulp et al. 2012; Sanjak et al. 2018). Assuming its heritability to be $h^2 = 0.8$, and making use of the value $\bar{r}_h = 0.464$ we have calculated and the estimate $V_S/V_P \sim 30$ from Sanjak et al. (2018), we find that $2d/V_g \sim 0.054$, while $h^2\rho/(1-h^2\rho) = 0.25$. Therefore, the effective effects of loci that affect height are amplified, on average, because of the relatively strong assortative mating; their frequency dynamics under stabilizing selection are therefore expected to be accelerated, in spite of the Bulmer effect.

Note that patterns of trait-based migration can lead to long-range linkage disequilibria similar in nature to those under assortative mating (Veller and Coop 2024).

4.4 Pleiotropy

Our calculations have ignored the possibility that the alleles underlying variation in the focal trait also affect other traits which might be under stabilizing selection (or, following the discussion above, subject to other processes that generate long-range signed LD). To see how such pleiotropy could affect our conclusions, consider a pair of alleles at distinct loci that both increase the focal trait. All else equal, we would expect stabilizing selection on the focal trait to generate negative LD between these two alleles, to a degree that we can calculate in expectation (e.g., Eqs. 20 and 28). However, suppose that there is another trait under stabilizing selection, and the one allele increases this trait but the other allele decreases it. All else equal, we would expect stabilizing selection on this other trait to generate positive LD between the alleles. Clearly, the sign and magnitude of the LD between the alleles will depend on the size of their effects on the two traits, as well as the relative strengths of selection on the two traits.

While ultimately the degree and nature of pleiotropy is an empirical question, we can suggest some qualitative predictions for how the Bulmer effect will influence the frequency dynamics of alleles that affect multiple traits under stabilizing selection. Following Simons et al. (2018, 2022), we focus on a scenario where we have allele effect-size estimates for a single trait, but where these alleles also affect other, unmeasured traits.

Consider first the 'isotropic' case where the effects of an allele on the various traits are independent. This is the case considered by Simons et al. (2018, 2022), who note that it will always apply in a suitably reconfigured coordinate system for the traits in question. Consider two alleles that increase the focal trait. In the absence of pleiotropy, we would expect these alleles to come into negative LD because of selection on the focal trait. And because, under isotropic pleiotropy, the signs and magnitudes of the alleles' effects on other traits are independent of their effects on the focal trait, the LD between the alleles is unchanged, *in expectation*, by their effects on these other traits. To be clear, the realized value of LD between this particular pair of alleles will change because of pleiotropy, but across such locus pairs, there is no tendency for pleiotropy to systematically increase or decrease the degree of LD from the value expected under selection on the focal trait alone.

It is, however, perhaps more likely that alleles' effects on traits will be correlated. That is, if a pair of alleles both increase trait A, and we learn that one of the alleles decreases trait B, then the other allele probably also decreases trait B (since, for example, the effects of the alleles on traits A and B might in part be mediated through shared pathways in a genetic network). In this case, alleles with like effects on the focal trait we have measured are disproportionately likely to have like effects on other, unmeasured traits. In this case, the magnitude of the LD between a pair of alleles would, in expectation, exceed the value expected based on their effects on the focal trait alone, as calculated, for example, in Eqs. (20) and (28). Therefore, in this perhaps more realistic scenario than the isotropic model, the slowdown in allele-frequency dynamics induced by the Bulmer effect would be more severe than we have calculated.

5 Conclusion

Stabilizing selection is increasingly recognized as a potential driver of the genetic architecture of complex traits (Simons et al. 2018, 2022; Sella and Barton 2019; Yair and Coop 2022). The calculations that we have developed here make possible more precise estimation of its effects on genetic architectures and, conversely, more precise inference of its parameters from genomic data.

As we have noted, there is a rich theoretical literature on the population genetics of stabilizing selection, wherein the Bulmer effect has been extensively treated (e.g., Bulmer 1971, 1974; Lande 1975, 1977; Turelli and Barton 1990, 1994). We see our work as complementary to this literature in its focus on the consequences of the Bulmer effect for single-locus allele-frequency dynamics rather than aggregate quantities such as genetic and phenotypic variance. In this, and in its relative simplicity, our work offers a potential bridge between this prior theoretical literature and practical application to the growing body of population genomic data.

6 Methods

All simulations were run in SLiM 4.0 (Haller and Messer 2023). The model organism is diploid, sexual, and undergoes random mating each generation. In all setups, the population size is $N = 10,000$, and there are initially $L = 1,000$ polymorphic loci, either all unlinked or placed along a linkage map as described below. Each individual's trait value is specified by $Y = \sum_{l=1}^{L} \alpha_l g_l$, where g_l is the number of trait-increasing alleles that the individual carries at locus l minus 1, and α_l is the effect size at the locus, as described in the Model section of the Main Text. For simplicity, in our simulations there is no environmental component of trait variation; i.e., the trait is fully heritable. The fitness of an individual with trait value Y is given by $\phi(Y) = \exp(-Y^2/2V_S)$. In each simulation, V_S was chosen so that $V_S/V_P = 5$, where V_P is the initial variance of the trait (equal to the initial genetic variance V_G, since the trait is by assumption fully heritable).

Fig. 1. The L loci are all unlinked with respect to each other (specified by recombination rate $1/2$ between 'adjacent' loci in SLiM), and the effect size at each locus l is $\alpha_l = 1$. At 500 of these loci, the initial frequency of the trait-increasing allele is $p_l = 0.2$, while at the other 500 loci, it is $p_l = 0.8$. The designation of the loci at which the trait-increasing allele is common versus rare is arbitrary, since all loci are unlinked with respect to each other. Trait-increasing and trait-decreasing alleles are initially assigned randomly to haploid genomes, independently across loci, so that there is no LD among loci in expectation. The initial additive genetic variance, V_G, is therefore equal in expectation to the initial genic variance, $V_g = \sum_{l=1}^{L} 2p_l(1-p_l)\alpha_l^2 = 320$, and so the (inverse) strength of selection is $V_S = 5 \times 320 = 1{,}600$.

The simulation then proceeds for 500 generations, with fitness in each generation specified as described above. In each generation, we measure the frequency of the (initially) minor allele at each locus, and average the frequencies across loci. 1,000 replicate simulations were run, with results averaged across simulations to produce the 'simulation' line in Fig. 1.

The prediction lines in Fig. 1 were constructed by iterating Eqs. (5), (10), and (25), updating only the allele frequency each generation (i.e., holding V_P constant, even though it is slowly decreasing according to the equations themselves).

Fig. 2. The specification of the simulations in Fig. 2 is the same as that in Fig. 1, except that the loci now lie along sex-specific linkage maps. For humans, we used the male and female maps generated by Kong et al. (2010), while for *Drosophila melanogaster*, we used the female map generated by Comeron et al. (2012) (there is no crossing over in male *Drosophila*). In each case, we apportioned the $L = 1,000$ loci to chromosomes proportional to their physical (bp) lengths (as reported in build 38 of the human reference genome[2] and in release 6 of the *D. melanogaster* reference genome[3]). We ignored the sex chromosomes and

---

[2]`https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_000001405.26/`
[3]`https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_029775095.1/`

the fourth 'dot' chromosome of *D. melanogaster*. For each chromosome, we spaced the loci apportioned to it uniformly along its sex-averaged genetic (cM) length. This was done to avoid the occurrence of adjacent loci with a zero recombination fraction, since in such cases the harmonic mean recombination fraction among all locus pairs would be undefined. As is usual in SLiM, there is no crossover interference; recombination fractions r between adjacent loci were therefore calculated from the map distances between these loci d via Haldane's map function, $r(d) = (1 - e^{-2d})/2$.

To plot the prediction line for Eq. (33) in Fig. 2 requires calculating the harmonic mean sex-averaged recombination fraction for each locus, $\bar{r}_h^{(l)}$, and the overall harmonic mean sex-averaged recombination fraction, $\bar{r}_h$. For every pair of loci on the same chromosome, we calculated the recombination fraction from the map distance using Haldane's map function, as specified above; for pairs of loci on different chromosomes, the recombination fraction is $1/2$. The relevant averages of these recombination fractions were then taken. We calculated values of $\bar{r}_h$ for humans and *D. melanogaster* of 0.464 and 0.0744 respectively. Notice that, in averaging Eq. (33) across the $L = 1{,}000$ loci, we must first calculate $\left[1 - (d/V_g)\left(\bar{r}_h/\bar{r}_h^{(l)}\right)\right]^2$ for each locus l separately, and then average this quantity across loci.

Fig. 3. The 'simulation' points in Fig. 3 derive from the same simulations as in Fig. 2 under the *D. melanogaster* linkage map. For each locus, the minor-allele frequency was measured after 250 generations, having started at frequency 0.2. We also calculated each locus l's harmonic mean recombination fraction with all other loci, $\bar{r}_h^{(l)}$—among the 1,000 loci, the minimum and maximum values were 0.063 and 0.133.

Because the dynamics at individual loci are noisy owing to drift, we binned loci according to their locus-specific harmonic mean recombination rates, into bins 0.05–0.07 (486 loci), 0.07–0.09 (388 loci), 0.09–0.11 (118 loci), 0.11–0.13 (6 loci), and 0.13–0.15 (2 loci). The simulation points in Fig. 3 are the average minor-allele frequency in each bin after 250 generations, plotted against the (arithmetic) average value of $\bar{r}_h^{(l)}$ in each bin.

For the predictions in Fig. 3 that take into account non-equilibrium values of LD after the onset of stabilizing selection (purple dots), note that Eq. (34) defines the sequence across generations t of expected LD values between loci l and l', starting with zero LD in generation 0 ($D_{ll'}^{(0)} = 0$):

$$\mathbb{E}\left[D_{ll'}^{(t)}\right] = -\frac{d\bar{r}_h}{2\alpha^2 L(L-1)} \sum_{i=0}^{t-1} (1 - r_{ll'})^i. \tag{47}$$

In each generation t, for each locus l, we calculate its expected LD values with every other locus l' according to Eq. (47); we then sum these values and substitute the sums into Eqs. (15) and (16) to find the mean trait values experienced by the alleles at locus l. These define the effective effect size at locus l in generation t,

$$\alpha_{\text{eff},l}^{(t)} = \alpha \left(1 + \frac{1}{p_l^{(t)}(1 - p_l^{(t)})} \sum_{l' \neq l} \mathbb{E}\left[D_{ll'}^{(t)}\right]\right), \tag{48}$$

which then defines the expected change in allele frequency at locus l from generation t to $t+1$:

$$\mathbb{E}\left[p_l^{(t+1)} - p_l^{(t)}\right] = p_l^{(t)}\left(1 - p_l^{(t)}\right)\left(p_l^{(t)} - 1/2\right)\frac{\left(\alpha_{\text{eff},l}^{(t)}\right)^2}{V_S + V_P}. \tag{49}$$

We calculated the expected allele frequency after 250 generations for each locus, and averaged these predictions within each bin to obtain the purple points plotted in Fig. 3.

Figs. S1–S4. In Figs. S1–S4, minor-allele frequencies and effect sizes vary across loci. Initial minor-allele frequencies were chosen, independently for each locus, from a uniform distribution on $[0.1, 0.3]$, and effect sizes were chosen, independently for each locus and independent of minor-allele frequencies, from a normal distribution with mean 0 and variance 1. Since effect sizes can be negative or positive here, we assigned the chosen effect size to the minor allele at each locus. The configuration results in an expected initial genetic variance of $V_G = 313.3$, and so we chose $V_S = 5V_G = 1566.7$.

Acknowledgements

We are grateful to Jeremy Berg, Graham Coop, Andy Dahl, Marida Ianni-Ravn, Xinyi Li, Pavitra Muralidhar, John Novembre, and Yuval Simons for helpful discussions. Funding was provided by a Branco Weiss fellowship to CV.

References

Abdellaoui, A., Yengo, L., Verweij, K. J., and Visscher, P. M. (2023). 15 years of gwas discovery: realizing the promise. *American Journal of Human Genetics*, 110(2):179–194.

Border, R., Athanasiadis, G., Buil, A., Schork, A. J., Cai, N., Young, A. I., Werge, T., Flint, J., Kendler, K. S., Sankararaman, S., et al. (2022a). Cross-trait assortative mating is widespread and inflates genetic correlation estimates. *Science*, 378(6621):754–761.

Border, R., O'Rourke, S., de Candia, T., Goddard, M. E., Visscher, P. M., Yengo, L., Jones, M., and Keller, M. C. (2022b). Assortative mating biases marker-based heritability estimators. *Nature Communications*, 13(1):660.

Bulmer, M. G. (1971). The effect of selection on genetic variability. *American Naturalist*, 105(943):201–211.

Bulmer, M. G. (1974). Linkage disequilibrium and genetic variability. *Genetics Research*, 23(3):281–289.

Carlson, M. O., Rice, D. P., Berg, J. J., and Steinrücken, M. (2022). Polygenic score accuracy in ancient samples: quantifying the effects of allelic turnover. *PLoS Genetics*, 18(5):e1010170.

Comeron, J. M., Ratnappan, R., and Bailin, S. (2012). The many landscapes of recombination in *Drosophila melanogaster*. *PLoS Genetics*, 8(10):e1002905.

Crow, J. F. and Felsenstein, J. (1968). The effect of assortative mating on the genetic composition of a population. *Eugenics Quarterly*, 15(2):85–97.

Ding, Y., Hou, K., Xu, Z., Pimplaskar, A., Petter, E., Boulier, K., Privé, F., Vilhjálmsson, B. J., Olde Loohuis, L. M., and Pasaniuc, B. (2023). Polygenic scoring accuracy varies across the genetic ancestry continuum. *Nature*, 618(7966):774–781.

Haller, B. C. and Messer, P. W. (2023). SLiM 4: multispecies eco-evolutionary modeling. *The American Naturalist*, 201(5):E127–E139.

Hayward, L. K. and Sella, G. (2022). Polygenic adaptation after a sudden change in environment. *eLife*, 11:e66697.

Horwitz, T. B., Balbona, J. V., Paulich, K. N., and Keller, M. C. (2023). Evidence of correlations between human partners based on systematic reviews and meta-analyses of 22 traits and uk biobank analysis of 133 traits. *Nature Human Behaviour*, 7(9):1568–1583.

Jiang, Y., Bolnick, D. I., and Kirkpatrick, M. (2013). Assortative mating in animals. *American Naturalist*, 181(6):E125–E138.

Kong, A., Thorleifsson, G., Gudbjartsson, D. F., Masson, G., Sigurdsson, A., Jonasdottir, A., Walters, G. B., Jonasdottir, A., Gylfason, A., Kristinsson, K. T., et al. (2010). Fine-scale recombination rate differences between sexes, populations and individuals. *Nature*, 467(7319):1099–1103.

Lande, R. (1975). The maintenance of genetic variability by mutation in a polygenic character with linked loci. *Genetics Research*, 26(3):221–235.

Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution. *Evolution*, 30(2):314–334.

Lande, R. (1977). The influence of the mating system on the maintenance of genetic variability in polygenic characters. *Genetics*, 86(2):485–498.

Martin, A. R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B. M., and Daly, M. J. (2019). Clinical use of current polygenic risk scores may exacerbate health disparities. *Nature Genetics*, 51(4):584–591.

Mostafavi, H., Harpak, A., Agarwal, I., Conley, D., Pritchard, J. K., and Przeworski, M. (2020). Variable prediction accuracy of polygenic scores within an ancestry group. *eLife*, 9:e48376.

Robertson, A. (1956). The effect of selection against extreme deviants based on deviation or on homozygosis. *Journal of Genetics*, 54:236–248.

Sanjak, J. S., Sidorenko, J., Robinson, M. R., Thornton, K. R., and Visscher, P. M. (2018). Evidence of directional and stabilizing selection in contemporary humans. *Proceedings of the National Academy of Sciences*, 115(1):151–156.

Sella, G. and Barton, N. H. (2019). Thinking about the evolution of complex traits in the era of genome-wide association studies. *Annual Review of Genomics and Human Genetics*, 20:461–493.

Simons, Y. B., Bullaughey, K., Hudson, R. R., and Sella, G. (2018). A population genetic interpretation of GWAS findings for human quantitative traits. *PLoS Biology*, 16(3):e2002985.

Simons, Y. B., Mostafavi, H., Smith, C. J., Pritchard, J. K., and Sella, G. (2022). Simple scaling laws control the genetic architectures of human complex traits. *bioRxiv*, doi: https://doi.org/10.1101/2022.10.04.509926.

Stulp, G., Pollet, T. V., Verhulst, S., and Buunk, A. P. (2012). A curvilinear effect of height on reproductive success in human males. *Behavioral Ecology and Sociobiology*, 66:375–384.

Stulp, G., Simons, M. J., Grasman, S., and Pollet, T. V. (2017). Assortative mating for human height: A meta-analysis. *American Journal of Human Biology*, 29(1):e22917.

Turelli, M. (1988). Phenotypic evolution, constant covariances, and the maintenance of additive variance. *Evolution*, 42(6):1342–1347.

Turelli, M. and Barton, N. H. (1990). Dynamics of polygenic characters under selection. *Theoretical Population Biology*, 38(1):1–57.

Turelli, M. and Barton, N. H. (1994). Genetic and statistical analyses of strong selection on polygenic traits: what, me normal? *Genetics*, 138(3):913–941.

Veller, C. and Coop, G. (2024). Interpreting population and family-based genome-wide association studies in the presence of confounding. *PLOS Biology*, 22(4):e3002511.

Veller, C., Kleckner, N., and Nowak, M. A. (2019). A rigorous measure of genome-wide genetic shuffling that takes into account crossover positions and Mendel's second law. *Proceedings of the National Academy of Sciences*, 116(5):1659–1668.

Wang, Y., Guo, J., Ni, G., Yang, J., Visscher, P. M., and Yengo, L. (2020). Theoretical and empirical quantification of the accuracy of polygenic scores in ancestry divergent populations. *Nature Communications*, 11(1):3865.

Wright, S. (1921). Systems of mating. III. Assortative mating based on somatic resemblance. *Genetics*, 6(2):144–161.

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, 16(2):97.

Yair, S. and Coop, G. (2022). Population differentiation of polygenic score predictions under stabilizing selection. *Philosophical Transactions of the Royal Society B*, 377(1852):20200416.

A1 The average fitness of an allele, accounting for phenotypic variance

Suppose that an allele's trait distribution is normal with mean μ and variance V_P. To find the average fitness of this allele, we integrate the fitness function $e^{-y^2/2V_S}$ over the allele's trait distribution[4]:

$$
\bar{w}(\bar{y} = \mu, \sigma^2 = V_P) = \int \overbrace{\frac{1}{\sqrt{2\pi V_P}} e^{-\frac{(y-\mu)^2}{2V_P}}}^{\text{trait distribution}} \cdot \overbrace{e^{-\frac{y^2}{2V_S}}}^{\text{fitness}} dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{V_S(y-\mu)^2 + V_P y^2}{2 V_S V_P}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{(V_S + V_P)y^2 - 2V_S\mu y + V_S\mu^2}{2 V_S V_P}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{y^2 - 2\frac{V_S\mu}{V_S+V_P} y + \frac{V_S\mu^2}{V_S+V_P}}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2 + \frac{V_S\mu^2}{V_S+V_P} - \frac{V_S^2\mu^2}{(V_S+V_P)^2}}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2 + \frac{V_S\mu^2}{V_S+V_P}\left(1 - \frac{V_S}{V_S+V_P}\right)}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2 + \frac{V_S V_P \mu^2}{(V_S+V_P)^2}}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2 + \frac{V_S V_P \mu^2}{(V_S+V_P)^2}}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \frac{1}{\sqrt{2\pi V_P}} \int \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2}{\frac{2 V_S V_P}{V_S+V_P}} - \frac{\mu^2}{2(V_S + V_P)}\right) dy
$$

$$
= \sqrt{\frac{V_S}{V_S + V_P}} e^{-\frac{\mu^2}{2(V_S+V_P)}} \cdot \int \frac{1}{\sqrt{2\pi \frac{V_S V_P}{V_S+V_P}}} \exp\left(-\frac{\left(y - \frac{V_S\mu}{V_S+V_P}\right)^2}{\frac{2 V_S V_P}{V_S+V_P}}\right) dy
$$

$$
= \sqrt{\frac{V_S}{V_S + V_P}} e^{-\frac{\mu^2}{2(V_S+V_P)}}, \tag{A.1}
$$

where the cancellation in the last line follows from the fact that the integrand in the second-to-last line is the probability density function of a normally distributed random variable [with mean $V_S\mu/(V_S+V_P)$ and variance $V_S V_P/(V_S + V_P)$]. Since we are interested only in relative fitness, and because the phenotypic variance is the same for the two alleles at the focal locus, we can ignore the constant at the front of

---

[4]The result of this calculation is standard in the literature; we detail the calculation here only for expository purposes.

Eq. (A.1) and write the relative fitness of the allele with trait mean μ and variance V_P as

$$\bar{w}(\bar{y} = \mu, \sigma^2 = V_P) = e^{-\frac{\mu^2}{2(V_S + V_P)}}.$$ (A.2)

A2 Mean phenotypes of a trait-increasing and trait-decreasing allele, and their subsequent frequency dynamics, when effect sizes and minor-allele frequencies differ across loci

The effect size and frequency of the trait-increasing allele at locus l are α_l and p_l.

Case 1: All loci are unlinked

We first consider the case where all loci are unlinked. The correlation coefficient between the allelic states at loci l and l' (polarized as being between the trait-increasing alleles at the two loci) is proportional to $\alpha_l \alpha_{l'}$, so that the LD between the loci $D_{ll'}$ is, in expectation, $-K\alpha_l\alpha_{l'}\sqrt{p_l(1-p_l)}\sqrt{p_{l'}(1-p_{l'})}$ where K is a constant of proportionality. As before, we denote the total reduction in the genetic variance due to the Bulmer effect by d, with this value specified by Eq. (17) in the Main Text. Then

$$\begin{aligned}
-d &= 2\sum_l \sum_{l' \neq l} D_{ll'} \alpha_l \alpha_{l'} \\
&= -2K \sum_l \sum_{l' \neq l} \alpha_l^2 \alpha_{l'}^2 \sqrt{p_l(1-p_l)}\sqrt{p_{l'}(1-p_{l'})} \\
&= -2KL(L-1) \left\langle \alpha_{l'}^2 \sqrt{p_{l'}(1-p_{l'})} \right\rangle^2,
\end{aligned}$$ (A.3)

where the angle brackets here denote an average across loci. Therefore,

$$K = \frac{d}{2L(L-1) \left\langle \alpha_{l'}^2 \sqrt{p_{l'}(1-p_{l'})} \right\rangle^2}.$$ (A.4)

From Eq. (13), the mean phenotype experienced by the trait-increasing allele at locus l is

$$\mathbb{E}[Y \mid I_l = 1] = (1 - p_l)\left(\alpha_l + \frac{1}{p_l(1 - p_l)}\sum_{l' \neq l} D_{ll'}\alpha_{l'}\right)$$

$$= (1 - p_l)\left(\alpha_l - \frac{K}{p_l(1 - p_l)}\sum_{l' \neq l}\alpha_l\alpha_{l'}^2\sqrt{p_l(1 - p_l)}\sqrt{p_{l'}(1 - p_{l'})}\right)$$

$$= (1 - p_l)\alpha_l\left(1 - \frac{K}{\sqrt{p_l(1 - p_l)}}\sum_{l' \neq l}\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right)$$

$$= (1 - p_l)\alpha_l\left(1 - \frac{d}{\sqrt{p_l(1 - p_l)}2L(L - 1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle^2}\cdot(L - 1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle\right)$$

$$= (1 - p_l)\alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1 - p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle}\right). \tag{A.5}$$

Similarly, the mean phenotype experienced by the trait-decreasing allele at l is

$$\mathbb{E}[Y \mid I_l = 0] = -p_l\alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1 - p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle}\right). \tag{A.6}$$

The 'effective' effect size of locus l is therefore

$$\alpha_{\text{eff}}^{(l)} = \alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1 - p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle}\right), \tag{A.7}$$

and the change in frequency at the locus across a single generation is, in expectation,

$$\mathbb{E}[\Delta p_l] = p_l(1 - p_l)\left(p_l - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P}$$

$$= p_l(1 - p_l)\left(p_l - \frac{1}{2}\right)\frac{\alpha_l^2\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\right)^2}{V_S + V_P}. \tag{A.8}$$

If allele frequencies do not differ much across loci,

$$2L\sqrt{p_l(1 - p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1 - p_{l'})}\right\rangle \approx 2L\left\langle\alpha_{l'}^2 p_{l'}(1 - p_{l'})\right\rangle = V_g, \tag{A.9}$$

and so

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{V_g}\right), \tag{A.10}$$

as in the case of symmetric effects and minor-allele frequencies across loci (Eq. 24).

Case 2: Linkage

If we allow for variable linkage relations among loci, then, in expectation, the degree of LD between loci l and l' is $-K\alpha_l\alpha_{l'}\sqrt{p_l(1-p_l)}\sqrt{p_{l'}(1-p_{l'})}/r_{ll'}$, where $r_{ll'}$ is the recombination fraction between loci l and l' and K is again a constant of proportionality. If d is the total reduction in the genetic variance due to the Bulmer effect, with this value specified by Eq. (26) in the Main Text, then

$$-d = 2\sum_l \sum_{l'\neq l} D_{ll'}\alpha_l\alpha_{l'}$$

$$= -2K\sum_l \sum_{l'\neq l} \frac{\alpha_l^2\alpha_{l'}^2\sqrt{p_l(1-p_l)}\sqrt{p_{l'}(1-p_{l'})}}{r_{ll'}}$$

$$\approx -2KL(L-1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle^2\frac{1}{\bar{r}_h} \tag{A.11}$$

assuming independence of α_l and $\alpha_{l'}$ from $r_{ll'}$ and a weak dependence of p_l and $p_{l'}$ on $r_{ll'}$. Therefore,

$$K \approx \frac{d\bar{r}_h}{2L(L-1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle^2}. \tag{A.12}$$

From Eq. (13), the mean phenotype experienced by the trait-increasing allele at locus l is

$$\mathbb{E}[Y\mid I_l=1] = (1-p_l)\left(\alpha_l + \frac{1}{p_l(1-p_l)}\sum_{l'\neq l}D_{ll'}\alpha_{l'}\right)$$

$$= (1-p_l)\left(\alpha_l - \frac{K}{p_l(1-p_l)}\sum_{l'\neq l}\alpha_l\alpha_{l'}^2\sqrt{p_l(1-p_l)}\sqrt{p_{l'}(1-p_{l'})}\cdot\frac{1}{r_{ll'}}\right)$$

$$= (1-p_l)\alpha_l\left(1 - \frac{K}{\sqrt{p_l(1-p_l)}}\sum_{l'\neq l}\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\cdot\frac{1}{r_{ll'}}\right)$$

$$\approx (1-p_l)\alpha_l\left(1 - \frac{d\bar{r}_h}{\sqrt{p_l(1-p_l)}2L(L-1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle^2\bar{r}_h^{(l)}}\cdot(L-1)\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle\right)$$

$$= (1-p_l)\alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right), \tag{A.13}$$

where the approximation follows from our assumption of a weak dependence of p_l and $p_{l'}$ on $r_{ll'}$. Similarly, the mean phenotype experienced by the trait-decreasing allele at l is

$$\mathbb{E}[Y\mid I_l=0] \approx -p_l\alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right). \tag{A.14}$$

The 'effective' effect size of locus l is therefore

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right), \tag{A.15}$$

and the change in frequency at the locus across a single generation is, in expectation,

$$\mathbb{E}\left[\Delta p_l\right] = p_l(1 - p_l)\left(p_l - \frac{1}{2}\right)\frac{\left(\alpha_{\text{eff}}^{(l)}\right)^2}{V_S + V_P}$$

$$\approx p_l(1 - p_l)\left(p_l - \frac{1}{2}\right)\frac{\alpha_l^2\left(1 - \frac{d}{2L\sqrt{p_l(1-p_l)}\left\langle\alpha_{l'}^2\sqrt{p_{l'}(1-p_{l'})}\right\rangle}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right)^2}{V_S + V_P}.\tag{A.16}$$

If allele frequencies do not differ much across loci,

$$\alpha_{\text{eff}}^{(l)} \approx \alpha_l\left(1 - \frac{d}{V_g}\cdot\frac{\bar{r}_h}{\bar{r}_h^{(l)}}\right),\tag{A.17}$$

as in the case of symmetric effects and minor-allele frequencies across loci (Eq. 32).

Figure S1: Separation of timescales over which stabilizing selection reduces genetic variance by generating linkage disequilibrium (Bulmer effect; difference between genic and genetic variance) and selecting against minor alleles at polymorphic loci affecting the trait (reduction in the genic variance). Here, loci are distributed along the human linkage map. Initial minor-allele frequencies were chosen from a uniform distribution on $[0.1, 0.3]$ and effect sizes were chosen, independently, from a normal distribution with mean 0 and variance 1. The strength of selection V_S was chosen such that $V_S/V_G = 5$ initially.

Figure S2: Predictions of average allele-frequency trajectories, versus simulated trajectories, when effect sizes and initial minor-allele frequencies are allowed to differ, and all loci are unlinked. Initial minor-allele frequencies were chosen from a uniform distribution on $[0.1, 0.3]$ and effect sizes were chosen, independently, from a normal distribution with mean 0 and variance 1. The strength of selection V_S was chosen such that $V_S/V_G = 5$ initially. Otherwise, simulation details are as for Main Text Fig. 1.

Figure S3: Predictions of average allele-frequency trajectories, versus simulated trajectories, when effect sizes and initial minor-allele frequencies are allowed to differ, and loci are distributed along the autosomal linkage map of *Drosophila melanogaster* in the same way as in Main Text Fig. 3B. Initial minor-allele frequencies were chosen from a uniform distribution on $[0.1, 0.3]$ and effect sizes were chosen, independently, from a normal distribution with mean 0 and variance 1. The strength of selection V_S was chosen such that $V_S/V_G = 5$ initially.

Figure S4: Predicted versus simulated minor-allele frequencies after 500 generations of selection when allelic effect sizes vary across loci, plotted as a function of the per-locus squared effect sizes. Loci are distributed along the human linkage map in the same way as in Fig. 3A. Initial minor-allele frequencies were chosen from a uniform distribution on $[0.1, 0.3]$ and effect sizes were chosen, independently, from a normal distribution with mean 0 and variance 1. The strength of selection V_S was chosen such that $V_S/V_G = 5$ initially.