The Amphibian Genomics Consortium: advancing genomic and genetic

resources for amphibian research and conservation

Authors

Tiffany A. Kosch[1][§][*], María Torres-Sánchez[2][§][*], H. Christoph Liedtke[3], Kyle Summers[4],

Maximina H. Yun[5,6], Andrew J. Crawford[7,8], Simon T. Maddock[9,10], Md. Sabbir

Ahammed[11], Victor L. N. Araújo[7], Lorenzo V. Bertola[12], Gary Bucciarelli[13], Albert

Carné[14], Céline M. Carneiro[15], Kin O. Chan[16], Ying Chen[17], Angelica Crottini[18,19,20],

Jessica M. da Silva[21,22], Robert D. Denton[23], Carolin Dittrich[24], Gonçalo Espregueira

Themudo[25], Katherine A. Farquharson[26,27], Natalie J. Forsdick[28], Edward Gilbert[29,30],

Che Jing[31,32], Barbara A. Katzenback[33], Ramachandran Kotharambath[34], Nicholas A.

Levis[35], Roberto Márquez[36], Glib Mazepa[37,38], Kevin P. Mulder[39], Hendrik Müller[40],

Mary J. O'Connell[41], Pablo Orozco-terWengel[42], Gemma Palomar[43,44], Alice

Petzold[45], David W. Pfennig[46], Karin S. Pfennig[46], Michael S. Reichert[47], Jacques

Robert[48], Mark D. Scherz[49], Karen Siu-Ting[50,51,52], Anthony A Snead[53], Matthias

Stöck[54], Adam M. M. Stuckert[55], Jennifer L. Stynoski[56], Rebecca D. Tarvin[57],

Katherina C. Wollenberg Valero[58], and The Amphibian Genomics Consortium (AGC).

[§]contributed equally to the manuscript

*correspondence:

Tiffany A. Kosch (tiffany.kosch@gmail.com)

María Torres-Sánchez (torressanchez.maria@gmail.com)

Affiliations

[1]One Health Research Group, Melbourne Veterinary School, Faculty of Science, University of Melbourne, Werribee, Victoria, Australia

[2]Department of Biodiversity, Ecology, and Evolution, Complutense University of Madrid, 28040 Madrid, Spain

[3]Eco-Evo-Devo Group, Estación Biológica de Doñana CSIC, Seville, Spain

[4]Biology Department, East Carolina University, Greenville, NC, USA 27858

[5]Technische Universität Dresden, CRTD/Center for Regenerative Therapies Dresden, Dresden, Germany

[6]Max Planck Institute for Molecular Cell Biology and Genetics, Dresden, Germany

[7]Department of Biological Sciences, Universidad de los Andes, Bogotá, 111711, Colombia

[8]Museo de Historia Natural C.J. Marinkelle, Universidad de los Andes, Bogotá, 111711, Colombia.

[9]School of Natural and Environmental Sciences, Newcastle University, UK

[10]Island Biodiversity and Conservation Centre, University of Seychelles, Seychelles

[11]Department of Zoology, Jagannath University, Dhaka 1100, Bangladesh

[12]Centre for Tropical Bioinformatics and Molecular Biology, James Cook University, QLD 4810, Australia

[13]Department of Wildlife, Fish, and Conservation Biology, University of California, Davis, USA

[14]Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain

[15]Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA

[16]University of Kansas Biodiversity Institute and Natural History Museum, Lawrence,

Kansas 66045, USA

[17]Biology Department, Queen's University, Ontario, Canada

[18]CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO

Laboratório Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão,

Portugal

[19]Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, rua do

Campo Alegre s/n, 4169– 007 Porto, Portugal

[20]BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus

de Vairão, 4485-661 Vairão, Portugal

[21]Evolutionary Genomics and Wildlife Management, Foundatonal Biodiversity

Science, Kirstenbosch Research Centre, South African National Biodiversity

Institute, Newlands 7735, Cape Town, South Africa

[22]Centre for Evolutionary Genomics and Wildlife Conservation, Department of

Zoology, University of Johannesburg, Auckland Park 2006, Johannesburg, South

Africa

[23]Department of Biology, Marian University, Indianapolis, IN 46222, USA

[24]Rojas Lab, Konrad-Lorenz-Institute of Ethology, Department of Life Science,

University of Veterinary Medicine, Vienna, Austria

[25]CIIMAR Interdisciplinary Centre of Marine and Environmental Research, University

of Porto, Terminal de Cruzeiros do Porto de Leixões, Avenida General Norton de

Matos, S/N, Matosinhos, Portugal

[26]School of Life and Environmental Sciences, The University of Sydney, Sydney,

NSW 2006, Australia

[27]Australian Research Council Centre of Excellence for Innovations in Peptide and

Protein Science, The University of Sydney, Sydney, New South Wales, Australia

[28]Manaaki Whenua – Landcare Research, Auckland, New Zealand

[29]School of Natural Sciences, The University of Hull, Hull, HU6 7RX, United Kingdom

[30]Energy and Environment Institute, The University of Hull, Hull, HU6 7RX, United

Kingdom

[31] Key Laboratory of Genetic Evolution and Animal Models, and Yunnan Key

Laboratory of Biodiversity and Ecological Conservation of Gaoligong Mountain,

Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223,

China

[32] Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences,

Yezin, Nay Pyi Taw 05282, Myanmar

[33]Department of Biology, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1

[34]Herpetology Lab, Dept. of Zoology, Central University of Kerala, Tejaswini Hills,

Kasaragod, Kerala, 671320, India

[35]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[36]Department of Biological Sciences, Virginia Tech. Blacksburg, VA 24060, USA

[37]Department of Ecology and Evolution, University of Lausanne, Biophore, 1015,

Switzerland

[38]Department of Ecology and Genetics, Evolutionary Biology, Norbyvägen 18D,

75236 Uppsala, Sweden

[39]Wildlife Health Ghent, Faculty of Veterinary Medicine, Ghent University, Merelbeke,

Belgium

[40]Central Natural Science Collections, Martin Luther University Halle-Wittenberg, D-

06108 Halle (Saale), Germany

[41]School of Life Sciences, Faculty of Medicine and Health Sciences, University of

Nottingham, UK

[42]School of Biosciences, Cardiff University, Museum Avenue, CF10 3AX Cardiff,

United Kingdom

[43]Department of Genetics, Physiology, and Microbiology; Faculty of Biological

Sciences; Complutense University of Madrid, Madrid, Spain

[44]Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University,

Kraków, Poland

[45]Institute of Biochemistry and Biology, University of Potsdam, Karl-Liebknecht

Str.24-25, 14476 Potsdam, Germany

[46]Department of Biology, University of North Carolina, Chapel Hill, NC 27599, USA

[47]Department of Integrative Biology, Oklahoma State University, Stillwater OK, USA

[48]Department of Microbiology and Immunology, University of Rochester Medical

Center, Rochester, NY, 14642, USA

[49]Natural History Museum of Denmark, University of Copenhagen,

Universitetsparken 15, 2100, Copenhagen Ø, Denmark

[50]School of Biological Sciences, Queen's University Belfast, Belfast, BT7 1NN,

Northern Ireland, United Kingdom

[51]Instituto Peruano de Herpetología, Ca. Augusto Salazar Bondy 136, Surco, Lima,

Peru

[52]Herpetology Lab, The Natural History Museum, London, United Kingdom

[53]Department of Biology, New York University, New York, NY, USA

[54]Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB),

Müggelseedamm 301, D-12587 Berlin, Germany

[55]Department of Biology and Biochemistry, University of Houston, Houston, Texas,

77204, USA

[56]Instituto Clodomiro Picado, Universidad de Costa Rica, San José, Costa Rica

[57]Museum of Vertebrate Zoology and Department of Integrative Biology, University of

California, Berkeley, CA 94720, USA

[58]School of Biology and Environmental Science, University College Dublin, Belfield

Campus, Dublin 4, Ireland

Abstract

Amphibians represent a diverse group of tetrapods, marked by deep divergence

times between their three systematic orders and families. Studying amphibian

biology through the genomics lens increases our understanding of the features of

this animal class and that of other terrestrial vertebrates. The need for amphibian

genomics resources is more urgent than ever due to the increasing threats to this

group. Amphibians are one of the most imperiled taxonomic groups, with

approximately 41% of species threatened with extinction due to habitat loss,

changes in land use patterns, disease, climate change, and their synergistic effects.

Amphibian genomics resources have provided a better understanding of ontogenetic

diversity, tissue regeneration, diverse life history and reproductive modes, anti-

predator strategies, and resilience and adaptive responses. They also serve as

critical models for understanding widespread genomic characteristics, including

evolutionary genome expansions and contractions given they have the largest range

in genome sizes of any animal taxon and multiple mechanisms of genetic sex

determination. Despite these features, genome sequencing of amphibians has

significantly lagged behind that of other vertebrates, primarily due to the challenges

of assembling their large, repeat-rich genomes and the relative lack of societal

support. The advent of long-read sequencing technologies, along with computational

techniques that enhance scaffolding capabilities and streamline computational

workload is now enabling the ability to overcome some of these challenges. To

promote and accelerate the production and use of amphibian genomics research

through international coordination and collaboration, we launched the Amphibian

Genomics Consortium (AGC) in early 2023. This burgeoning community already has

more than 282 members from 41 countries (6 in Africa, 131 in the Americas, 27 in

Asia, 29 in Australasia, and 89 in Europe). The AGC aims to leverage the diverse

capabilities of its members to advance genomic resources for amphibians and bridge

the implementation gap between biologists, bioinformaticians, and conservation

practitioners. Here we evaluate the state of the field of amphibian genomics,

highlight previous studies, present challenges to overcome, and outline how the AGC

can enable amphibian genomics research to "leap" to the next level.

Keywords

Amphibians, Biodiversity conservation, Comparative genomics, Genomics,

Lissamphibia, Metagenomics, Phylogenomics, Population genomics, Taxonomy,

Transcriptomics.

State of the field of amphibian genomics

In 2010, the genome of the Western clawed frog (*Xenopus tropicalis*) was

sequenced, marking the first genome assembly for Class Amphibia [1]. This species

serves as a crucial laboratory model organism for cell biology, molecular genetics,

and developmental biology [2]. The first amphibian genome assembly came years

after the completion of the first genomes for other vertebrate groups: fishes in 2002

(*Fugu rubripes*; [3]), mammals in 2003 (*Homo sapiens*; [4]), birds in 2004 (*Gallus*

gallus; [5]), and reptiles in 2007 (*Anolis carolinensis*; Anolis Genome Project

https://www.broadinstitute.org/anolis/anolis-genome-project). Since then, the

generation and annotation of amphibian reference genomes has dramatically lagged

behind those of other vertebrates [6], even though amphibians represent nearly 22%

of all tetrapods [7]. Nearly 15 years later, amphibians are still the tetrapod class with

the lowest number of sequenced genomes (111 genomes of 8648 described

amphibian species being the tetrapod class with the second lowest proportion after

Reptiles [database records accessed on 1 March 2024], Fig. 1A and Supplementary

File 1). This is likely attributable to the size of amphibian genomes, which are

generally larger than the genomes of other terrestrial vertebrates (Fig. 1B and Fig.

S1; see Supplementary Material for methodological information). Indeed, among all

vertebrates, only the genomes of lungfish are larger (up to 130 Gb) than the largest

amphibian genomes (up to ~120 Gb in *Necturus lewisi*) [8-10].

To lower cost and enhance feasibility, early amphibian genome sequencing projects

tended to select species with comparatively small genomes (Fig. 1B). This has

resulted in the disproportionately fewer sequenced salamander genomes [11]. To

date, the largest amphibian genome assemblies belong to three salamander

species: *Ambystoma mexicanum* (27.3 Gb assembly; [12]), *Pleurodeles waltl* (20.3

Gb; [13]), and *Calotriton arnoldi* (22.8 Gb; [14]). However, these only represent the

lower end of the genome size range for this group, with the genomes of *Necturus*

salamanders exceeding 100 Gb (Fig. 2) [10].

In addition to their large sizes, amphibian genomes have also been challenging to

assemble due to their extensive repeat content (up to 82% [15]). Amphibian

transposable elements have expanded and become highly abundant in younger

clades, posing challenges for the construction of contiguous genome assemblies

[16]. These characteristics of amphibian genomes make sequencing and assembly

both costly and technically challenging (e.g., repetitive regions can often lead to

fragmented assemblies when using short-read sequencing). However, the advent of

new sequencing approaches such as long-read sequencing (e.g., PacBio HiFi and

Oxford Nanopore Duplex), Hi-C scaffolding, along with reduced sequencing costs

have resolved many of these assembly challenges (e.g., *Nanorana parkeri*; [17]).

Thus, the number of amphibian genome assemblies has increased rapidly in recent

years, reaching 111 currently listed as reference genomes at scaffold level or higher

in the National Center for Biotechnology Information (NCBI) genome database (52

for Anura, 55 for Urodela, and four for Gymnophiona; NCBI genome database

records accessed on 1 March 2024). Despite this rapid increase, the quality of

available amphibian genomes varies significantly, only 38 are chromosome-level

assemblies, and among these, only 16 are annotated. This indicates that the majority

of available assemblies are incomplete or partial. For example, several recently

published salamanders genomes of the genus *Desmognathus* have assembly sizes

of ~1 Gb while their genome size estimates based on flow cytometry or image

densitometry average 14 Gb [18, 19]. Furthermore, the gene content values for

many of these incomplete genomes can be as low as 0.7% [15]. Besides the

variation in quality, there are substantial taxonomic gaps in genome representation

across Amphibia. Notably, 48 of the 77 amphibian families (62%) lack a

representative genome assembly in the NCBI genome database (Fig. 2B), indicating

significant gaps in our understanding (see "The AGC's genome sequencing targets"

section and Table 1 for more information about these 48 families).

Due to the difficulty of assembling genomes, most previous genomic research in

amphibians has relied on alternative high-throughput sequencing methodologies,

including RNA sequencing (RNA-seq), reduced representation or target-capture

approaches, or metagenomic methods (Fig. 3 and Supplementary File 2). For

example, RNA sequencing (RNA-Seq) techniques have been used to explore gene

expression across more than 300 different amphibian species. Furthermore, a

substantial number of *de novo* transcriptomes are available through the NCBI

Transcriptome Shotgun Assemblies (TSA) database (79 total: 59 for Anura, 15 for

Urodela, and 5 for Gymnophiona). Various reduced-representation (e.g., ddRADseq)

and targeted-capture sequencing approaches have also been implemented in recent

years to obtain genome-wide sequence information from more than 1,400 amphibian

species. All this information—from whole genomes to gene transcript features—has

advanced the understanding of amphibian biology and directly contributed to

conservation efforts as described below.

Advancing research and conservation through amphibian genomics

Amphibians have many unique characteristics that make them subjects of interest to

a wide variety of scientific disciplines, spanning from developmental biology and

medical research to ecology and evolution. The rapid development of genomic tools

is galvanizing the study of amphibian biology and uncovering important facets of

their biology and conservation [20-22]. We highlight some examples here.

<u>Embryogenesis, developmental and regenerative biology</u>

Amphibians have played a fundamental role in uncovering development principles

[for a detailed review see 23]. Research on anurans has enabled the understanding

of critical mechanisms underlying such as the breaking of egg asymmetry [24], axis

establishment, and nerve transmission [25]. Notably, the availability of genome

assemblies for *Xenopus laevis* and *X. tropicalis* has significantly advanced

embryological and developmental biology. This advancement has enabled gene loss-

of-function research through the combination of transgenesis with RNA interference,

gene editing, and enhanced morpholino design. This has facilitated the in-depth

analysis of regulatory and non-coding genomic influences in developmental

processes [26, 27]. Consequently, these studies have generated thousands of

genomic and transcriptomic resources for these two species [28, 29].

Yet, there is much more to uncover about amphibian development, especially given

the numerous developmental modalities found across amphibian species, which

likely demonstrates the highest diversity among vertebrates [30]. This includes direct

development (egg to froglet; the first genome of a direct-developing amphibian,

Eleutherodactylus coqui, was published in 2024 [31]), and phenotypic plasticity [32,

33].

Sexual development and determination are also diverse in amphibians [34],

encompassing female and male heterogamety, unique microscopically recognizable

sex chromosomes, and species with undifferentiated sex chromosomes. Sex-

determining genes across these systems are starting to be explored through high-

throughput sequencing [6, 35-38]. For instance, a Y-specific non-coding RNA

involved in male sex determination in *Bufotes viridis* was identified through the

application of multiple omics techniques [38]. Strikingly, some salamanders in the

genus *Ambystoma* exist as a single all-female, polyploid lineage that can incorporate

new chromosome sets from up to five other sexual species [39]. Transcriptomes

from these salamanders have shown that gene expression from their divergent

genomes is balanced for some genes but biased for others [40]. Sexual development

in amphibians can result in sexually dimorphic features such as nuptial spines, which

have been explored using comparative genomics approaches in the frog

Leptobrachium leishanense [41]. The increasing availability of amphibian genomes

will enable a deeper understanding of the molecular mechanisms underlying such

ontogenetic diversity.

Metamorphosis sets many amphibian species apart from amniotes. Transcriptomics

has revealed a remarkable turnover in gene expression between larval and adult

stages of both frogs [42-45] and salamanders [46, 47]. This represents genomic

uncoupling of these life history phases with major macroevolutionary implications

[44, 48]. Amphibian omics approaches are rapidly increasing our understanding of

the developmental process of metamorphosis, including the role of methylation in

gene regulation [49]. Amphibians have also been found to respond to environmental

perturbations by altering their behavior or phenotypes in various ways. These include

changing developmental rate [32], hybridization with positive fitness effects [50],

producing novel trophic morphologies [51], and kin detection to avoid cannibalizing

relatives [52-54].

Due to their exceptional tissue repair and regenerative capacities [55, 56],

amphibians are leading models for understanding the mechanisms of regeneration.

This is particularly true for salamanders, which display the most extensive adult

regenerative repertoire among vertebrates, including the ability to regenerate parts of

their eyes, brain, heart, jaws, lungs, spinal cord, tail, and entire limbs [56]. Due to

new genome assemblies for urodele species, *Ambystoma mexicanum* and the

Pleurodeles waltl, regeneration can now be studied with transgenesis, advanced

imaging, and genome editing. Intensive transcriptomic sequencing for these two

salamander species has facilitated gene expression studies, including investigations

into regeneration processes and characterization of other genomic features [57].

Additionally, a novel mechanism of telomere length maintenance and elongation has

recently been described in *P. waltl* [58], potentially linking regenerative capability with

longevity. Other amphibian species have also contributed to genomic research on

regeneration. For example, databases compiled from gene expression resources of

Notophthalmus viridescens have enabled comparative studies [59].

<u>Ecology and evolution</u>

Modern amphibians are the sister lineage of all amniotes, making them a valuable

resource for studying species relationships and trait evolution. This is exemplified by

studies that explore the rapid diversification of frogs [60], the evolution of vision [61],

hybridogenesis [62-64], and the evolution of limblessness [65]. Amphibian

phylogenomics has addressed many longstanding questions in amphibian evolution

[66-69]. Comparative genomic analyses including amphibian groups have also

revealed important gaps in our understanding of tetrapod molecular evolution such

as chromosomal rearrangements and group-specific gene families that remain

unclassified to date [65, 70, 71]. In this section, we explore how genomics is being

applied to understand the diverse ecological and evolutionary features unique to

amphibians.

Like mammals, birds, and reptiles [72-74], some amphibians have evolved the ability

to live in high-elevation environments such as the Andes (up to 5400 m) [75, 76] and

the Tibetan Plateau (4478 m) [17]. However, unlike other groups, amphibians lack

fur, feathers, or scales to protect them from physiological stressors such as UV

exposure. This vulnerability makes them an intriguing model for studying the effects

of UV radiation, which is relevant not only to humans [17] but also to species

impacted by climate change. Amphibians have evolved multiple mechanisms of

resisting UV including increasing antioxidant efficiency and gene regulatory changes

in defense pathways [17, 77]. There is also evidence from Tibetan anurans that

genes that impact other high-elevation traits (e.g., hypoxia resistance, immunity, cold

tolerance) have evolved convergently across distantly related families (e.g.,

Dicroglossidae, Bufonidae, Megophryidae) [78], and that intraspecific divergence in

many of these genes correlates with elevation [79, 80]. Additional genomic

signatures of elevation adaptations, including genes regulating resistance to cold,

hypoxia, immunity, and reproduction, have been described in ranid species of

western North America inhabiting elevations from the sea level to nearly 3000 m [81].

While we are beginning to understand the genetic mechanisms of high-elevation

adaptation in some Asian and North American frogs, this has yet to be investigated in

other high-elevation amphibians, including Andean anurans (e.g., *Telmatobius culeus*

[82]) and high-elevation salamanders, such as *Pseudoeurycea gadovii* [83]).

The ability to produce toxins has evolved across all three amphibian orders, where it

primarily serves as an anti-predation mechanism. The source of amphibian toxins

varies: some species are capable of synthesizing poisonous compounds (e.g.,

bufonids, myobatrachids), whereas others sequester toxic substances from their diet

(e.g., dendrobatids, mantellids) [84-87] or microbial symbionts (e.g., newts) [88].

Since dendrobatid frogs sequester their toxins from prey (e.g., mites and ants), they

lack genes encoding these toxins [89, 90]. However, they require genes to facilitate

the transport of these toxins to the skin. Recent genomic and proteomic research

has identified candidate genes coding for proteins that may serve dual roles in toxin

transport and resistance [91-93]. Comparative genomic research has identified

specific substitutions that allow toxic amphibian species to effectively mitigate the

effects of the sequestered toxins on their own tissues [94-96]. Skin transcriptomes

have also proven to be a rich source for data mining and the identification of

candidate toxins and antimicrobial peptides in various amphibians, including

caecilians [97-101].

Interactions between toxic amphibians and their predators have resulted in a

fascinating variety of co-evolutionary arms races. These include well-characterized

systems of toxicity resistance mechanisms in amphibian predators [102-106] and

aposematism and mimicry in toxic species [107, 108]. Research on aposematism

and mimicry has utilized whole genome, exome capture, and transcriptome

sequencing to elucidate the genes underlying the vast diversity of color patterns

across populations and species in dendrobatids [109-114]. These approaches have

yielded a goldmine of information that can be used to understand the genes, gene

networks, and biochemical pathways that underlie variation in coloration in other

amphibian groups including highly diverged aposematic taxa such as Australian

myobatrachid frogs (e.g., *Pseudophryne corroboree)*, Malagasy poison frogs

(Mantellidae), caecilians (e.g., *Schistometopum thomense*), and salamanders (e.g.,

Salamandra salamandra). Indeed, these methods have already enabled the

identification of genes and loci involved in coloration in the salamander *S.*

salamandra bernardezi [115].

Despite the numerous advances made with amphibian omics in elucidating

evolutionary and ecological mechanisms, fully unraveling their genetic basis requires

generating a vast number of genomes due to the comparative nature of these fields.

Some of the exciting research avenues in amphibians include parental care [116,

117], gliding ability [118], lunglessness [119, 120], unusual defense mechanisms,

such as the ability of some newts to pierce their ribs through toxin glands in their skin

[121, 122], milk production or skin feeding in caecilians [123, 124], and spatial

navigation [125].

<u>Conservation</u>

Amphibians are the most endangered class of vertebrates with current estimates

suggesting that more than 40% of species are threatened with extinction [126]. The

threats amphibians are facing continue to increase [126], creating a clear need to

develop innovative and effective methods to conserve them. Paradoxically, current

rates of amphibian species description are exponential, and numerous candidate

species are being flagged worldwide. This suggests that we are still far from

overcoming the amphibian Linnean shortfall, especially in tropical regions [127, 128].

Hence, numbers of threatened species are likely underestimated, as undescribed

species cannot be assessed and are more likely to become extinct [129]. Further, the

conservation status of many amphibians remains unknown, especially for tropical

species [130] and for a number of soil-dwelling caecilians known only from a few

specimens [131]. Generating genomic data is one method to address this challenge,

as it can be used to estimate both evolutionary potential and extinction risk [132].

Genomes are also vital for understanding species boundaries and the geographic

distribution of genetic diversity within species, and for identifying populations under

higher risk due to anthropogenic pressures or climate change [20, 21, 133, 134].

These features make genomic resources invaluable for developing species

conservation action plans [135].

Amphibian conservation efforts should leverage population genetic theory and the

burgeoning field of conservation genomics. This approach enables the quantification

of both neutral and adaptive diversity across genomes, thereby facilitating the

promotion of adaptive potential or genetic rescue through translocation programs

[136-139]. Typically, these studies begin with the genomic characterization of

populations across various environmental conditions, assessing population genetic

health and disease risk [140, 141]. Omics resources are becoming increasingly

important in species detection (e.g., via eDNA [142-144]). They can also support

monitoring and surveillance efforts by identifying populations most at risk of declines

due to potential genetic threats such as maladaptive alleles, genetic load, inbreeding

and outbreeding depression, hybridization, and/or genetic incompatibility [136, 145].

Increased monitoring and maintenance of genomic diversity are key targets of many

national and international recommendations such as the US Endangered Species

Act [146], the Kunming-Montreal Global Biodiversity Monitoring Framework [147],

and the Amphibian Conservation Action Plan [135].

A more specific application of amphibian genomics for conservation requires

understanding the genetic basis of traits that impact fitness, such as disease

resistance or climate change tolerance. This information can be used to promote

adaptation using approaches like Targeted Genetic Intervention (TGI), which aims to

increase the frequency of adaptive alleles with approaches such as selective

breeding, genome editing, or targeted gene flow [148]. Considerable effort has been

invested in understanding the genetic basis of resistance to the devastating

amphibian disease chytridiomycosis, which has resulted in the identification of

multiple candidate genes [149-151] that could be targeted to increase

chytridiomycosis resistance with TGI. Additionally, the efficacy of TGI at increasing

chytridiomycosis resistance has already been demonstrated in North American

mountain yellow-legged frogs (*Rana muscosa* and *R. sierrae*) where translocation of

resistant individuals increased recipient population persistence [152]. Despite the

obvious appeal of using genetic intervention approaches for conservation, these

methods should be evaluated in contained facilities whenever possible and

accompanied by long-term monitoring to ensure their efficacy and rule out any

unintended impacts [148, 153-155]. Although such conservation interventions require

extensive resources, this may be the only effective method for restoring some

species to the wild, especially in those threatened by intractable threats such as

chytridiomycosis [156].

Challenges for amphibian genomic research and ways forward

A major challenge for amphibian omics research, which will be anchored by high-

quality reference genomes, is gaining access to starting material (e.g., tissue, blood).

Common logistic challenges include: 1) obtaining research funding, 2) collaborating

equitably with local or Indigenous communities on the development and execution of

biodiversity genomics research, 3) obtaining collection and research permits from the

state, 4) obtaining samples from difficult-to-access regions, and 5) obtaining high-

quality samples and maintaining quality during transport.

With increasingly easy access to genomic data, researchers and industry need to be

even more aware of the principles of fair and equitable access to genetic resources,

as stipulated by Convention on Biological Diversity (CBD) and expanded upon by the

Nagoya Protocol https://www.cbd.int/abs/default.shtml). Indigenous peoples and

local communities (IPLC) are often custodians of genetic resources (physical

material) sought by researchers, requiring that all parties enter into collaborative and

equitable agreements on access and benefit-sharing (ABS) before embarking on a

genomics project [157-161]. As a negative example from amphibians, *Phyllomedusa*

bicolor skin secretions traditionally used by Amazonian indigenous peoples were

patented by actors in the US, Japan, Russia and elsewhere, promoting the 'legal' but

unfair appropriation of genetic resources and potentially the traditional knowledge

itself from the Matses and other Indigenous tribes [162]. To promote better practices,

researchers should budget the time and money required to engage in prior

consultation as part of planning field work, and consult their National Focal Points on

ABS. How the concept of ABS will or could be applied to the downstream use of the

digital sequence information (DSI) generated has yet to be resolved (although there

are currently developments underway, https://www.cbd.int/dsi-gr) but must also be

considered going forward [see for example 163]. Moreover, voucher specimens and

duplicate tissue samples should be deposited in local museums or preferred partners

of the local communities [164, 165].

Amphibian fieldwork often involves overcoming numerous hurdles such as

navigating socio-political conflicts and accessing remote field sites. These

challenges may be exacerbated in developing economies or for researchers with

limited access to high volume funding streams. In addition, inadequate infrastructure

for accessing field sites, and the need to time of fieldwork to coincide with the often

highly seasonal and cryptic activities of amphibians [166] can further complicate the

task. Attention should also be directed towards overcoming inequities that may pose

additional obstacles to fieldwork for underrepresented groups [161].

Once amphibians are collected, selecting the tissue sample to obtain sufficient high-

molecular-weight DNA (HMW, reaching 100 Kb or ultra HMW, reaching 1 Mb) can be

challenging due to the small body sizes of most amphibians (e.g., < 30 g). For

generating reference genomes, blood is one of the most recommended sample

types for amphibian genomics [167]. However, the blood volume of most amphibians

is too small for non-lethal collection [168]. This presents a significant challenge

because obtaining the required quantity of HMW DNA often necessitates lethal

sampling, which may not always be legally permitted or ethically advisable, especially

for threatened species or those in captive collections [169]. Non-lethal sampling

approaches, such as buccal swabs or toe or tail clips, are increasingly viable for

various genomic applications, including low-coverage whole genome sequencing or

targeted sequencing approaches [170, 171]. Another alternative is to use tadpoles

instead of adults as was done to generate the genome of *Taudactylus pleione* [172].

Most tissue sampling protocols for reference genomes or transcriptome sequencing

recommend harvesting samples from fresh tissue, followed immediately by flash

freezing in liquid nitrogen (LN2) and storing at -80°C until extraction

(https://www.vertebrategenomelab.org/resources/guidelines). Maintaining ultracold

storage and the cold chain during transport from remote collection sites can be

challenging. After being adequately charged with LN2, dry vapor shippers can keep

samples ultra-cold for a week or more in the field. While LN2 is the gold standard,

other preservatives like 95% EtOH, 20-25% DMSO in EDTA may adequately

preserve DNA for long-read sequencing, although efficacy may vary depending on

the tissue type [167, 173]. Other options include lysis buffers or commercial products

such as Zymo DNA/RNA shield (Zymo Research, USA). However, it remains

essential to test the impact of these preservatives on sequencing outcomes

beforehand, specifically for the given taxon and tissue type, since preservation

methods may inhibit downstream approaches, such as Hi-C library construction [167,

174].

While these challenges apply to any researcher or organization, they become

particularly challenging when the sequencing work occurs outside the species'

country of origin. This difficulty arises not only because genetic material is prone to

degradation but also because of regulations on moving biological samples across

political borders [175, 176]. The global genomics community should strive to ensure

that sequencing projects occur within the country of origin of the samples and

discourage 'parachute' or 'helicopter science' [177, 178]. Oxford Nanopore

Technology (ONT) may be promising solution, providing comparatively affordable

access to equipment and reagents for ultra-long read sequencing that can even be

done directly in the field [179]. However, working with non-model organisms requires

prior optimization, and the startup costs for this infrastructure remain prohibitive for

many scientists from low-income countries. Moving forward, the goal should be to

apply these technologies in collaboration with local researchers. For example,

programs like the In Situ Laboratories Initiative (https://insitulabs.org/hubs/) aim to

provide affordable access to high-tech laboratories in remote biodiverse areas).

Such collaboration should proceed from finding shared interests, developing ideas,

realizing the shared benefits from research outputs, and focusing on capacity-

building efforts [180].

Working with museum specimens [the burgeoning field of "museomics"; 181] is

another promising avenue of research allowing to access to past amphibian

biodiversity. However, there are several additional challenges associated with DNA

degradation, preservation methods, and contamination that need to be overcome

[182-184]. This is particularly relevant for wet-preserved amphibian specimens, as

retrieving DNA can be challenging due to the often unknown fixation and

preservation methods that can alter nucleotide integrity. Methodological advances in

laboratory protocols [e.g., 185, 186, 187] and the development of sequencing

strategies, such as 'Barcode Fishing' [188, 189], have made significant progress in

addressing these challenges, including the ability to sequence extinct species [190,

191]. In the current era, even limited sequences from taxonomic type specimens are

of unparalleled importance, especially for species identification using genetic data,

by those applying methods like eDNA and metagenomics [192]. Museomics has also

revolutionized amphibian taxonomy by integrating DNA from name-bearing types,

overcoming impediments like uncertainty in taxonomic names, species complexes,

and cryptic species [188, 189, 193, 194].

As mentioned earlier, the large genome sizes of many amphibians affect

sequencing, computing, and storage costs, as well as computational requirements

for data analysis [6, 195]. However, the greater challenges lie in the methodological

and theoretical limitations of assembling large genomes. Additionally, access to

cluster computing for genome assembly remains limited for many scientists,

particularly those from low-income countries. Genome size in amphibians is

correlated with increased intron lengths and repeat content [196], posing challenges

for assembly, especially when using with short-read sequencing technologies or

during contig joining processes. Regions misassembled due to low complexity have

previously resulted in a significant loss of sequence information (e.g., by as much as

16%) through the collapsing of repetitive sequences [197]. Repetitive regions can

also result in the formation of problematic chimeras during assembly, where two

distant contigs are erroneously joined due to a shared repeat sequence [198]. These

problems are exacerbated when repeats are longer than sequencing reads. Further

challenges are posed by polyploidy [199], which has evolved repeatedly in

amphibians [200, 201]. This can make developing haplotype-specific assemblies

challenging and may require dramatically increased sequencing and computational

efforts [199, 202]. Thus, the development of long-read and 3C technology (i.e., Hi-C

scaffolding) is especially important for assembling amphibian genomes [198, 203].

Recent technological advances, including long-read techniques such as PacBio HiFi,

ONT, as well as scaffolding methods such as Hi-C and optical mapping, now make it

feasible and affordable to generate reference genomes for most frogs, caecilians,

and increasingly for salamanders [203].

Gene and repeat annotation resources in amphibians are also under-developed.

Although this may not pose a significant issue for well-conserved genes, where

researchers can retrieve annotation information from orthologous regions using

existing databases like UniProt [204], it often results in missed or poorly annotated

genes, particularly for highly polymorphic genes or genes lacking representation in

model taxa. Hence, the ability to study features such as gene evolution [67], repeats

[15, 196], immune genes [205], and genetic sex determination [206] is still limited

and requires caution to prevent overinterpretation.

Although key tools for understanding functional genomics exist, tools and protocols

for gene editing, transgenesis, *in vitro* fertilization, are rare or non-existent for most

amphibians [148, 207], with the exception of some model species (e.g., *Xenopus*

laevis, *X. tropicalis*, and *Ambystoma mexicanum*) [56, 208-211]. Immortal cell lines

have been successfully generated for some amphibians [212] and protocols have

been established to facilitate the initiation of spontaneously arising cell lines for a

subset of anurans [213]. However, establishing cell cultures for most species

requires extensive problem-solving and expertise [212].

Aims, priorities, and structure of the Amphibian Genomics Consortium (AGC)

The AGC was launched in March 2023 to address the aforementioned knowledge

gaps through technological advances and international cooperation. The mission of

the AGC is to increase collaboration and communication among amphibian

researchers internationally and across scientific disciplines, to increase amphibian

genomics resources, and to apply genomic data and functional resources to help

bridge the implementation gaps between genome biologists, other scientists and

conservation practitioners. The leadership structure of the AGC consists of a director,

two co-directors, and a 10-member board. The board was carefully chosen to ensure

gender equality, diversity of scientific disciplines, career stages, and representation

from various geographic regions.

Currently the AGC organizes monthly virtual meetings and a monthly virtual seminar

series. They maintain a website (https://mvs.unimelb.edu.au/amphibian-genomics-

consortium), facilitate active discussion groups, and have hosted social events at

several scientific conferences. Additionally, the AGC is preparing to host a full-day

symposium titled "Beyond the reference: genomics for amphibian research and

conservation" at the World Congress of Herpetology in August 2024 in Kuching,

Malaysia.

The first actions of the AGC include raising funds for genome sequencing,

developing technical resources and best practices guidelines, improving amphibian

genome annotation, supporting travel for students and early career researchers, and

conducting virtual and in-person computational workshops. The AGC plans to secure

funding to sequence high-priority amphibian species (see The AGC's genome

sequencing targets section and Table 1). Additionally, the AGC aims to facilitate

amphibian sample collection for broader taxonomic consortia. The AGC is already

affiliated with the Earth BioGenome Project (EBP) and AmphibiaWeb

(https://amphibiaweb.org), reinforcing its commitment to advancing amphibian

genomics and conservation efforts.

<u>AGC membership</u>

At the time of the submission of this work, the AGC had 282 members from 41

countries (Fig. 4). Although the membership is geographically diverse, there remains

some disparity across regions. Efforts will be intensified to attract members from

underrepresented countries, particularly regions known for high amphibian diversity

and/or endemism such as Central and South America, and Southeast Asia. We

promote equity between members by providing additional support and opportunities

to those from developing countries and underrepresented groups. This includes

eliminating membership fees, scheduling online meetings at alternating times to

accommodate global time zones, facilitating discussion groups on the cloud-based

collaboration platform Discord, and translating AGC correspondence into multiple

languages. Furthermore, we are also committed to fostering knowledge and skills

transfer to all emerging scientists worldwide, and we actively encourage early career

researchers to join the initiative and participate in governance.

<u>Current use and perception of genomics technologies by members of the AGC</u>

The AGC leadership designed a 23-question survey to investigate consortium

members' experiences in amphibian genomics (questions can be found in

Supplementary Table S1). The survey was distributed using the Qualtrics XM

platform and remained active from the 4th of March to the 27th of December 2023.

We collected responses from a total of 133 AGC members from 32 countries with

different expertise in sequencing approaches and bioinformatics techniques, who

primarily work on the ecology and evolution of anurans. Overall, respondents

emphasized the urgency of filling knowledge gaps in amphibian genomics due to the

current conservation crisis, pinpointing the necessity to expand the number of high-

quality chromosome-level amphibian genomes. Additionally, there was strong

agreement among survey respondents that the generation of new genomics

resources needs to be coupled with the improvement and accessibility of annotation

processes. A better development of sharing computational expertise among

members and resources internationally was also underscored. More than half of the

survey participants said they use sequencing technologies for their studies (70 of the

133). About half of the respondents said their main work activities were "genomics

lab work" or "computational analyses" (48% and 57%, respectively).

To evaluate consortium members' experience in amphibian genomics, we applied a

principal components analysis to the quantitative responses. Bioinformatic

competencies and perceived challenges of the AGC respondents were grouped in

two dimensions, respectively (Fig. 5A and Fig. S2; see Supplementary Material for

methodological information). To explain the variation of these two new variables, we

used the scientific expertise of AGC members, the funding success, and two

variables related to the country of main affiliation of the respondent: the number of

amphibian species and gross domestic expenditure on R&D (GERD) per capita, as

explanatory variables. Amphibian genomics expertise and identified challenges

varied substantially among respondents. The number of amphibian species and

GERD per capita of the respondent's main affiliation country did not capture this

variation (Fig. 5B and Fig. S3; see Supplementary Material for methodological

information). Instead, genomics funding success and years of scientific expertise

were, as expected, positively correlated and both variables and were associated with

a reduction in the perceived challenges associated with amphibian genomics.

<u>The AGC's genome sequencing targets</u>

Following the efforts of genomics consortia for other tetrapod groups [e.g., 214], and

previous research efforts [21], we identified amphibian families without any

representative genomes and selected one representative species per family for our

sequencing priority list (Fig. 2B and Table 1). We propose 48 candidate species

based on their IUCN Red List category, ecological and evolutionary distinctiveness,

and the availability of other genomics records, especially transcriptomics. This list

includes 38 anurans, four urodeles, and six caecilians. We aim to build upon the

efforts of existing genomics consortia such as the Vertebrate Genomes Project

(VGP), hence, we included in our sequencing target list species with draft genomes

in the GenomeArk (https://www.genomeark.org/).

Conclusion and outlook

Amphibians are declining faster than they can be discovered [215]. Our hope is that

the recent advancements in technology (e.g., long-read sequencing, computational

tools) and integration of the research community to from the Amphibian Genomics

Consortium (AGC) will ignite research to revolutionize amphibian conservation and

our understanding of their fascinating biology, ecology and evolution. By supporting

amphibian genomics research and uniting amphibian researchers worldwide, the

AGC aims to propel amphibian genomics research into the future.

Moving forward, the AGC is committed to supporting amphibian sequencing

initiatives worldwide, with a particular emphasis on families lacking representation

and species from biodiverse countries (Table 1). Local sequencing initiatives will be

given priority whenever feasible to promote the development of *in situ* research

efforts and facilities. Additionally, we aim to provide funding and training opportunities

to facilitate collaboration among underrepresented groups, molecular and organismal

biologists, bioinformaticians, and conservation practitioners. We aspire to stimulate

public and scientific interest in amphibian research and enhance conservation efforts

for this intriguing and highly endangered group of vertebrates.

List of Abbreviations

ABS: access and benefit-sharing

AGC: Amphibian Genomics Consortium

CBD: Convention on Biological Diversity

DSI: digital sequence information

EBP: Earth BioGenome Project

GERD: gross domestic expenditure on research and development

GoaT: Genomes on a Tree

HMW: High molecular weight DNA

IPLC: Indigenous peoples and local communities

IUCN: International Union for Conservation of Nature

ONT: Oxford Nanopore Technology

VGP: Vertebrate Genomes Project

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

The authors declare no competing interests.

Funding

T.A.K. was supported by Australian Research Council grants (FT190100462 and LP200301370). M.T.-S. was supported by María Zambrano fellowship from Complutense University of Madrid and NextGenerationEU. The *Xenopus laevis* Research Resource for Immunobiology is supported by the National Institute of Health (R24-AI-05983).

Authors' contributions

T.A.K. and M.T.-S. drafted the manuscript. T.A.K., M.T.-S., H.C.L., K.S., M.H.Y., S.T.M., A.J.C. contributed text to the first draft, M.T.-S. and T.A.K. analyzed the data and created the figures, members of the Amphibian Genomics Consortium (AGC) reviewed later drafts. All authors reviewed the manuscript.

Acknowledgments

Contributing members of the Amphibian Genomics Consortium (AGC) in alphabetical

order: Aldemar A. Acevedo, Steven J. R. Allain, Lisa N. Barrow, M. Delia Basanta,

Roberto Biello, Gabriela B. Bittencourt-Silva, Amaël Borzée, Ian G. Brennan, Rafe M.

Brown, Natalie Calatayud, Hugo Cayuela, Jing Chai, Ignacio De la Riva, Lana J.

Deaton, Khalid A. E. Eisawi, Kathryn R. Elmer, W. Chris Funk, Giussepe Gagliardi-

Urrutia, Wei Gao, Mark J. Goodman, Sandra Goutte, Melissa Hernandez Poveda,

Tomas Hrbek, Oluyinka A. Iyiola, Gregory F.M. Jongsma, J. Scott Keogh, Tianming

Lan, Pablo Lechuga-Paredes, Emily Moriarty Lemmon, Stephen C. Lougheed, Thom

A. Lyons, Mariana L. Lyra, Jimmy A. McGuire, Marco A. Mendez, Hosne Mobarak,

Edina Nemesházi, Tao T. Nguyen, Michaël P.J. Nicolaï, Lotanna M. Nneji, John B.

Owens, Hibraim Pérez-Mendoza, Nicolas Pollet, Megan L. Power, Mizanur Rahman,

Hans Recknagel, Ariel Rodríguez, Santiago R. Ron, Joana Sabino-Pinto, Yongming

Sang, Suman Sapkota, Rosio G. Schneider, Laura Schulte, Ana Serra Silva, Lee F.

Skerratt, Nicholas Strowbridge, Karthikeyan Vasudevan, Govindappa Venu, Lucas

Vicuña, David R. Vieites, Judit Vörös, Matt West, Mark Wilkinson, Guinevere O. U.

Wogan. We also appreciate the helpful suggestions from Jia-Tang Li, Jun Li, Wei

Wu, and Hua Wu.

Table 1. Amphibian Genomics Consortium (AGC) sequencing priority list. Table of

amphibian families without any sequenced genomes. For each family, AGC

proposed a candidate species based on its IUCN Red List category (LC: Least

Concern, NT: Near Threatened, VU: Vulnerable, EN: Endangered, CR: Critically

Endangered, and NA: Not evaluated), ecological and evolutionary distinctiveness,

and availability of other genomic records. This table shows the amphibian order to

which each family belongs and its number of genera (#G) and described extant

species (#S) as well as distribution region. *Species with available draft genome

assemblies in the GenomeArk (https://www.genomeark.org/).

Family	Region	#G	#S	Candidate species	IUCN	Other motives
Anura: Allophrynidae	South America	1	3	*Allophryne relicta*	EN	
Anura: Alsodidae	South America	3	26	*Alsodes gargola*	LC	High altitude adaptation
Anura: Arthroleptidae	Africa	8	151	*Leptopelis vermiculatus*	EN	
Anura: Ascaphidae	North America	1	2	*Ascaphus montanus**	LC	High altitude adaptation
Anura: Batrachylidae	South America	4	13	*Batrachyla leptopus*	LC	
Anura: Brachycephalidae	South America	2	79	*Brachycephalus pitanga*	LC	Transcriptomic resources
Anura: Brevicipitidae	Africa	5	36	*Breviceps fuscus*	LC	Burrowing adaptation
Anura: Caligophrynidae	South America	1	1	*Caligophryne doylei*	NA	Pantepui endemism
Anura: Calyptocephalellidae	South America	2	5	*Telmatobufo bullocki*	EN	
Anura: Centrolenidae	Central & South America	12	166	*Centrolene pipilata*	CR	Gigantism
Anura: Ceratobatrachidae	Southeast Asia	4	103	*Platymantis spelaeus*	EN	Cave-dweller

Family	Region	#G	#S	Candidate species	IUCN	Other motives
Anura: Ceratophryidae	South America	3	12	*Lepidobatrachus laevis*	LC	Transcriptomic resources
Anura: Ceuthomantidae	South America	2	6	*Ceuthomantis cavernibardus*	LC	Cave-dweller
Anura: Conrauidae	Africa	1	8	*Conraua goliath*	EN	Giantism
Anura: Craugastoridae	Central America	3	136	*Craugastor fitzingeri*	LC	Transcriptomic resources
Anura: Cycloramphidae	South America	3	37	*Cycloramphus granulosus*	CR	
Anura: Heleophrynidae	South Africa	2	6	*Heleophryne rosei*	CR	
Anura: Hemiphractidae	Central & South America	6	123	*Gastrotheca cornuta*	CR	
Anura: Hemisotidae	Sub-Saharan Africa	1	9	*Hemisus marmoratus*	LC	Transcriptomic resources
Anura: Hylodidae	South America	4	49	*Phantasmarana massarti*	EN	
Anura: Hyperoliidae	Sub-Saharan Africa & Madagascar	17	236	*Hyperolius thomensis*	EN	Population genomic resources
Anura: Leiopelmatidae	New Zealand	1	3	*Leiopelma archeyi*	CR	
Anura: Mantellidae	Madagascar	12	272	*Mantidactylus betsileanus*	LC	Transcriptomic resources
Anura: Micrixalidae	India	1	24	*Micrixalus mallani*	EN	Transcriptomic resources
Anura: Nasikabatrachidae	India	1	2	*Nasikabatrachus sahyadrensis*	NT	EDGE target species
Anura: Neblinaphrynidae	South America	1	1	*Neblinaphryne mayeri*	NA	Pantepui endemism
Anura: Nyctibatrachidae	India & Sri Lanka	3	37	*Nyctibatrachus grandis*	EN	Transcriptomic resources
Anura: Odontobatrachidae	Tropical West Africa	1	5	*Odontobatrachus fouta*	EN	
Anura: Odontophrynidae	South America	3	54	*Proceratophrys redacta*	EN	
Anura: Petropedetidae	Sub-Saharan tropical Africa	3	13	*Petropedetes perreti*	CR	
Anura: Phrynobatrachidae	Africa	1	99	*Phrynobatrachus guineensis*	LC	Tree-hole breeder
Anura: Ranixalidae	India	2	19	*Indirana chiravasi*	LC	Transcriptomic

Family	Region	#G	#S	Candidate species	IUCN	Other motives
						resources
Anura: Rhacophoridae	Eastern Asia	22	444	*Buergeria otai*	LC	Transcriptomic resources
Anura: Rhinodermatidae	South America	1	3	*Rhinoderma darwinii*	EN	Targeted sequencing resources
Anura: Rhinophrynidae	Central America	1	1	*Rhinophrynus dorsalis* *	LC	Targeted sequencing resources
Anura: Sooglossidae	Seychelles Islands	2	4	*Sooglossus sechellensis*	EN	EDGE target species
Anura: Strabomantidae	South America	19	792	*Oreobates cruralis*	LC	Transcriptomic resources
Anura: Telmatobiidae	South America	1	63	*Telmatobius simonsi*	CR	
Gymnophiona: Caeciliidae	Central & South America	2	49	*Caecilia tentaculata*	LC	Transcriptomic resources
Gymnophiona: Chikilidae	India	1	4	*Chikila gaiduwani*	LC	
Gymnophiona: Grandisoniidae	Africa, Seychelles & India	7	24	*Hypogeophis montanus*	NA	
Gymnophiona: Herpelidae	Sub-Saharan Africa	2	11	*Boulengerula niedeni*	EN	EDGE target species
Gymnophiona: Scolecomorphidae	Africa	2	6	*Crotaphatrema lamottei*	CR	
Gymnophiona: Typhlonectidae	South America	5	14	*Typhlonectes compressicauda*	LC	Transcriptomic resources
Urodela: Cryptobranchidae	Asia & North America	2	6	*Cryptobranchus alleganiensis*	VU	Transcriptomic resources
Urodela: Dicamptodontidae	North America	1	4	*Dicamptodon tenebrosus*	LC	Giantism
Urodela: Hynobiidae	Eastern Asia	9	98	*Hynobius vandenburghi*	VU	Transcriptomic resources
Urodela: Rhyacotritonidae	North America	1	4	*Rhyacotriton olympicus*	NT	

766
767
768

769 **Figures**
770

Figure 1. **Estimated genome size across tetrapod classes in relation to**

sequenced genomes. (A) Mosaic plot representing the percentage of species with

sequenced genomes as a proportion of the number of described species for each

tetrapod class (Yes: % species with sequenced genome; No: % species without

sequenced genome). (B) Combined box and density plot with raw data as points

comparing genome size of species with sequenced genome (gray; genome sizes

from NCBI genome assemblies) versus a subset of species without a sequenced

genome (red; genome sizes from the Animal Genome Size Database) for each

tetrapod class. The y-axis is log-transformed to facilitate visualization. Information

about sequenced genomes and genome sizes was obtained from the NCBI Genome

Browser, the Animal Genome Size database, and amphibian records from [11, 19].

Figure 2. Estimated genome size across amphibian orders in relation to

sequenced genomes. (A) Combined box and density plot with raw data as points

showing genome size of species with sequenced genome (gray color; genome sizes

from NCBI genome assemblies) versus a subset of species without available

genome assembly (red color; genome sizes from the Animal Genome Size

Database) for each amphibian order. The y-axis is logarithmic transformed to

facilitate visualization. Information about sequenced genomes and genome sizes

was obtained from the NCBI Genome Browser, the Animal Genome Size database

[19], and amphibian records from [11]. (B) Amphibian phylogenetic tree was adapted

from [66], which includes species with genome size estimates from Genomes on a

Tree (GoaT) [18]. Branches are color coded to represent families without any

genomic record (in red) and families with at least a representative genome

sequenced (in gray). Bar plots around the phylogeny indicate relative genome sizes.

Figure 3. Main sequencing techniques applied to amphibian genomics studies.

Yearly cumulative number of amphibian BioProjects split and color-coded by

sequencing technique (DNA accessibility Sequencing includes ATAC-Seq and

Mnase-Seq; Immunoprecipitation Sequencing includes: ChIP-Seq and RIP-Seq;

Amplicon sequencing was included with Targeted-Capture Sequencing; Noncoding

RNA Sequencing includes: miRNA-Seq and ncRNA-Seq). BioProject information

was obtained from the NCBI Sequence Read Archive (SRA, accessed 1 March

2024).

Figure 4. Amphibian Genomics Consortium (AGC) membership by country.

Inset map showing the size of each country scaled by number of members in the

AGC.

Figure 5. Sequencing competencies and identified challenges of the members of the Amphibian Genomics Consortium (AGC). (A) Representation of the contribution of the AGC survey quantitative questions to the first dimensions after computing a principal component analysis (PCA). Bioinformatic competencies and perceived challenges were grouped into dimensions one and two, respectively. (B) Scatter plot showing PCA scores for each AGC survey respondent. Respondent answers are coded by the qualitative question about funding success for amphibian genomics projects using shape; number of amphibian species of the respondent main affiliation country by size, and gross domestic expenditure on R&D (GERD) per capita of the respondent main affiliation country by gray-scale color coded. Information about the number of amphibian species per country was obtained from AmphibiaWeb. GERD per capita was calculated using information from the UNESCO and World Bank websites from the information about the most recent year for each country.

References

1. Hellsten U, Harland RM, Gilchrist MJ, Hendrix D, Jurka J, Kapitonov V, Ovcharenko I, Putnam NH, Shu S, Taher L *et al*: **The Genome of the Western Clawed Frog *Xenopus tropicalis***. *Science* 2010, **328**(5978):633-636.

2. Band MR, Larson JH, Rebeiz M, Green CA, Heyen DW, Donovan J, Windish R, Steining C, Mahyuddin P, Womack JE *et al*: **An ordered comparative map of the cattle and human genomes**. *Genome Res* 2000, **10**(9):1359-1368.

3. Aparicio S, Chapman J, Stupka E, Putnam N, Chia JM, Dehal P, Christoffels A, Rash S, Hoon S, Smit A *et al*: **Whole-genome shotgun assembly and analysis of the genome of *Fugu rubripes***. *Science* 2002, **297**(5585):1301-1310.

4. Collins FS, Green ED, Guttmacher AE, Guyer MS, on behalf of the USNHGRI: **A vision for the future of genomics research**. *Nature* 2003, **422**(6934):835-847.

5. Hillier LW, Miller W, Birney E, Warren W, Hardison RC, Ponting CP, Bork P, Burt DW, Groenen MAM, Delany ME *et al*: **Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution**. *Nature* 2004, **432**(7018):695-716.

6. Stöck M, Kratochvíl L, Kuhl H, Rovatsos M, Evans BJ, Suh A, Valenzuela N, Veyrunes F, Zhou Q, Gamble T *et al*: **A brief review of vertebrate sex evolution with a pledge for integrative research: towards 'sexomics'**. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2021, **376**(1832):20200426.

7. Pillay R, Venter M, Aragon-Osejo J, González-del-Pliego P, Hansen AJ, Watson JE, Venter O: **Tropical forests are home to over half of the world's vertebrate species**. *Front Ecol Environ* 2022, **20**(1):10-15.

8. Gregory TR: **The evolution of the genome**: Elsevier; 2011.

9. Biscotti MA, Carducci F, Olmo E, Canapa A: **Vertebrate Genome Size and the Impact of Transposable Elements in Genome Evolution**. In: *Evolution, Origin of Life, Concepts and Methods*. Edited by Pontarotti P. Cham: Springer International Publishing; 2019: 233-251.

10. Weisrock DW, Hime PM, Nunziata SO, Jones KS, Murphy MO, Hotaling S, Kratovil JD: **Surmounting the large-genome "problem" for genomic data generation in salamanders**. *Population genomics: wildlife* 2021:115-142.

11. Liedtke HC, Gower DJ, Wilkinson M, Gomez-Mestre I: **Macroevolutionary shift in the size of amphibian genomes and the role of life history and climate**. *Nature Ecology & Evolution* 2018, **2**(11):1792-1799.

12. Nowoshilow S, Schloissnig S, Fei J-F, Dahl A, Pang AWC, Pippel M, Winkler S, Hastie AR, Young G, Roscito JG *et al*: **The axolotl genome and the evolution of key tissue formation regulators**. *Nature* 2018.

13. Brown T, Elewa A, Iarovenko S, Subramanian E, Araus AJ, Petzold A, Susuki M, Suzuki K-iT, Hayashi T, Toyoda A *et al*: **Sequencing and chromosome-scale assembly of the giant *Pleurodeles waltl* genome**. *bioRxiv* 2022:2022.2010.2019.512763.

14. Talavera A, Palmada-Flores M, Burriel-Carranza B, Valbuena-Ureña E, Mochales-Riaño G, Adams DC, Tejero-Cicuéndez H, Soler-Membrives A, Amat F, Guinart D *et al*: **Genomic insights into the Montseny brook newt (*Calotriton arnoldi*), a Critically Endangered glacial relict**. *iScience* 2024, **27**(1):108665.

15. Kosch TA, Crawford AJ, Mueller RL, Wollenberg Valero KC, Rodríguez A, O'Connell LA, Young ND, Skerratt LF: **Comparative analysis of amphibian genomes: an emerging resource for basic and applied research**. *bioRxiv* 2023:2023.2002.2027.530355.

16. Sun C, Shepard DB, Chong RA, López Arriaza J, Hall K, Castoe TA, Feschotte C, Pollock
DD, Mueller RL: **LTR retrotransposons contribute to genomic gigantism in**
plethodontid salamanders. *Genome Biol Evol* 2012, **4**(2):168-183.
17. Fu T-T, Sun Y-B, Gao W, Long C-B, Yang C-H, Yang X-W, Zhang Y, Lan X-Q, Huang S, Jin
J-Q *et al*: **The highest-elevation frog provides insights into mechanisms and**
evolution of defenses against high UV radiation. *Proceedings of the National*
Academy of Sciences 2022, **119**(46):e2212406119.
18. Challis R, Kumar S, Sotero-Caio C, Brown M, Blaxter M: **Genomes on a Tree (GoaT): A**
versatile, scalable search engine for genomic and sequencing project metadata
across the eukaryotic tree of life [version 1; peer review: 2 approved]. *Wellcome*
Open Research 2023, **8**(24).
19. Gregory TR: **Animal genome size database**. 2024.
20. Calboli FCF, Fisher MC, Garner TWJ, Jehle R: **The need for jumpstarting amphibian**
genome projects. *Trends Ecol Evol* 2011, **26**(8):378-379.
21. Funk WC, Zamudio KR, Crawford AJ: **Advancing Understanding of Amphibian**
Evolution, Ecology, Behavior, and Conservation with Massively Parallel Sequencing.
In: *Population Genomics: Wildlife*. Edited by Hohenlohe PA, Rajora OP. Cham:
Springer International Publishing; 2018: 211-254.
22. Sun Y-B, Zhang Y, Wang K: **Perspectives on studying molecular adaptations of**
amphibians in the genomic era. *Zool Res* 2020, **41**(4):351.
23. Callery EM: **There's more than one frog in the pond: A survey of the Amphibia and**
their contributions to developmental biology. In: *Semin Cell Dev Biol: 2006*: Elsevier;
2006: 80-92.
24. Weaver C, Kimelman D: **Move it or lose it: axis specification in *Xenopus***.
Development 2004, **131**(15):3491-3499.
25. Burggren WW, Warburton S: **Amphibians as animal models for laboratory research**
in physiology. *ILAR journal* 2007, **48**(3):260-269.
26. Naert T, Van Nieuwenhuysen T, Vleminckx K: **TALENs and CRISPR/Cas9 fuel**
genetically engineered clinically relevant *Xenopus tropicalis* tumor models. *Genesis*
2017, **55**(1-2):e23005.
27. Guille M, Grainger R: **Genetics and Gene Editing Methods in *Xenopus laevis* and**
Xenopus tropicalis. *Cold Spring Harbor Protocols* 2023, **2023**(6):pdb. top107045.
28. Horb M, Wlizla M, Abu-Daya A, McNamara S, Gajdasik D, Igawa T, Suzuki A, Ogino H,
Noble A, France CdRBXti: ***Xenopus* resources: transgenic, inbred and mutant**
animals, training opportunities, and web-based support. *Frontiers in physiology*
2019, **10**:387.
29. Fisher M, James-Zorn C, Ponferrada V, Bell AJ, Sundararaj N, Segerdell E, Chaturvedi P,
Bayyari N, Chu S, Pells T: **Xenbase: key features and resources of the *Xenopus* model**
organism knowledgebase. *Genetics* 2023, **224**(1):iyad018.
30. Liedtke HC, Wiens JJ, Gomez-Mestre I: **The evolution of reproductive modes and life**
cycles in amphibians. *Nat Commun* 2022, **13**(1):7039.
31. Bredeson JV, Mudd AB, Medina-Ruiz S, Mitros T, Smith OK, Miller KE, Lyons JB, Batra
SS, Park J, Berkoff KC *et al*: **Conserved chromatin and repetitive patterns reveal slow**
genome evolution in frogs. *Nat Commun* 2024, **15**(1):579.
32. Liedtke HC, Harney E, Gomez-Mestre I: **Cross-species transcriptomics uncovers**
genes underlying genetic accommodation of developmental plasticity in spadefoot
toads. *Mol Ecol* 2021, **30**(10):2220-2234.

33.   Isdaner AJ, Levis NA, Pfennig DW: **Comparative transcriptomics reveals that a novel**
form of phenotypic plasticity evolved via lineage-specific changes in gene
expression. *Ecol Evol* 2023, **13**(10):e10646.
34.   Nemesházi E, Bókony V: **HerpSexDet: the herpetological database of sex**
determination and sex reversal. *Scientific Data* 2023, **10**(1):377.
35.   Roco ÁS, Olmstead AW, Degitz SJ, Amano T, Zimmerman LB, Bullejos M: **Coexistence**
of Y, W, and Z sex chromosomes in *Xenopus tropicalis*. *Proceedings of the National*
Academy of Sciences 2015, **112**(34):E4752-E4761.
36.   Jeffries DL, Lavanchy G, Sermier R, Sredl MJ, Miura I, Borzée A, Barrow LN, Canestrelli
D, Crochet P-A, Dufresnes C *et al*: **A rapid rate of sex-chromosome turnover and**
non-random transitions in true frogs. *Nat Commun* 2018, **9**(1):4088.
37.   Cauret CM, Jordan DC, Kukoly LM, Burton SR, Anele EU, Kwiecien JM, Gansauge M-T,
Senthillmohan S, Greenbaum E, Meyer M: **Functional dissection and assembly of a**
small, newly evolved, W chromosome-specific genomic region of the African
clawed frog *Xenopus laevis*. *PLoS Genet* 2023, **19**(10):e1010990.
38.   Kuhl H, Tan WH, Klopp C, Kleiner W, Koyun B, Ciorpac M, Feron R, Knytl M, Kloas W,
Schartl M *et al*: **A candidate sex determination locus in amphibians which evolved**
by structural variation between X- and Y-chromosomes. *Nat Commun* 2024,
15(1):4781.
39.   Bogart JP, Bi K, Fu J, Noble DW, Niedzwiecki J: **Unisexual salamanders (genus**
***Ambystoma*) present a new reproductive mode for eukaryotes**. *Genome* 2007,
50(2):119-136.
40.   McElroy KE, Denton RD, Sharbrough J, Bankers L, Neiman M, Gibbs HL: **Genome**
expression balance in a triploid trihybrid vertebrate. *Genome Biol Evol* 2017,
9(4):968-980.
41.   Li Y, Ren Y, Zhang D, Jiang H, Wang Z, Li X, Rao D: **Chromosome-level assembly of the**
mustache toad genome using third-generation DNA sequencing and Hi-C analysis.
GigaScience 2019, **8**(9):giz114.
42.   Das B, Cai L, Carter MG, Piao Y-L, Sharov AA, Ko MS, Brown DD: **Gene expression**
changes at metamorphosis induced by thyroid hormone in *Xenopus laevis* tadpoles.
Dev Biol 2006, **291**(2):342-355.
43.   Schott RK, Bell RC, Loew ER, Thomas KN, Gower DJ, Streicher JW, Fujita MK:
Transcriptomic evidence for visual adaptation during the aquatic to terrestrial
metamorphosis in leopard frogs. *BMC Biol* 2022, **20**(1):138.
44.   Wollenberg Valero KC, Garcia-Porta J, Rodríguez A, Arias M, Shah A, Randrianiaina
RD, Brown JL, Glaw F, Amat F, Künzel S *et al*: **Transcriptomic and macroevolutionary**
evidence for phenotypic uncoupling between frog life history phases. *Nat Commun*
2017, **8**(1):15213.
45.   Zhao L, Liu L, Wang S, Wang H, Jiang J: **Transcriptome profiles of metamorphosis in**
the ornamented pygmy frog *Microhyla fissipes* clarify the functions of thyroid
hormone receptors in metamorphosis. *Sci Rep* 2016, **6**(1):27310.
46.   Palacios-Martinez J, Caballero-Perez J, Espinal-Centeno A, Marquez-Chavoya G,
Lomeli H, Salas-Vidal E, Schnabel D, Chimal-Monroy J, Cruz-Ramirez A: **Multi-organ**
transcriptomic landscape of *Ambystoma velasci* metamorphosis. *Dev Biol* 2020,
466(1-2):22-35.
47.   Sanchez E, Küpfer E, Goedbloed DJ, Nolte AW, Lüddecke T, Schulz S, Vences M,
Steinfartz S: **Morphological and transcriptomic analyses reveal three discrete**

primary stages of postembryonic development in the common fire salamander,
Salamandra salamandra. *Journal of Experimental Zoology Part B: Molecular and*
Developmental Evolution 2018, **330**(2):96-108.

48.   Goedert D, Calsbeek R: **Experimental evidence that metamorphosis alleviates**
genomic conflict. *The American Naturalist* 2019, **194**(3):356-366.

49.   Kyono Y, Raj S, Sifuentes CJ, Buisine N, Sachs L, Denver RJ: **DNA methylation**
dynamics underlie metamorphic gene regulation programs in *Xenopus* **tadpole**
brain. *Dev Biol* 2020, **462**(2):180-196.

50.   Pfennig KS: **Facultative mate choice drives adaptive hybridization.** *Science* 2007,
318(5852):965-967.

51.   Levis NA, Pfennig DW: **Innovation and diversification via plasticity-led evolution.** In:
Phenotypic plasticity & evolution. CRC Press; 2021: 211-240.

52.   Pfennig DW, Collins JP: **Kinship affects morphogenesis in cannibalistic salamanders.**
Nature 1993, **362**(6423):836-838.

53.   Pfennig DW, Reeve HK, Sherman PW: **Kin recognition and cannibalism in spadefoot**
toad tadpoles. *Anim Behav* 1993, **46**(1):87-94.

54.   DeVore JL, Crossland MR, Shine R, Ducatez S: **The evolution of targeted cannibalism**
and cannibal-induced defenses in invasive populations of cane toads. *Proceedings*
of the National Academy of Sciences 2021, **118**(35):e2100765118.

55.   Brockes JP, Kumar A: **Comparative aspects of animal regeneration.** *Annu Rev Cell Dev*
Biol 2008, **24**:525-549.

56.   Cox BD, Yun MH, Poss KD: **Can laboratory model systems instruct human limb**
regeneration? *Development* 2019, **146**(20):dev181016.

57.   Torres-Sánchez M: **Variation under domestication in animal models: the case of the**
Mexican axolotl. *BMC Genomics* 2020, **21**(1):827.

58.   Yu Q, Gates PB, Rogers S, Mikicic I, Elewa A, Salomon F, Lachnit M, Caldarelli A,
Flores-Rodriguez N, Cesare AJ *et al*: **Telomerase-independent maintenance of**
telomere length in a vertebrate. *bioRxiv* 2022:2022.2003.2025.485759.

59.   Bruckskotten M, Looso M, Reinhardt R, Braun T, Borchardt T: **Newt-omics: a**
comprehensive repository for omics data from the newt *Notophthalmus*
viridescens. *Nucleic Acids Res* 2012, **40**(Database issue):D895-900.

60.   Feng Y-J, Blackburn DC, Liang D, Hillis DM, Wake DB, Cannatella DC, Zhang P:
Phylogenomics reveals rapid, simultaneous diversification of three major clades of
Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proceedings of the*
National Academy of Sciences 2017, **114**(29):E5864-E5870.

61.   Schott RK, Fujita MK, Streicher JW, Gower DJ, Thomas KN, Loew ER, Bamba Kaya AG,
Bittencourt-Silva GB, Guillherme Becker C, Cisneros-Heredia D: **Diversity and**
Evolution of Frog Visual Opsins: Spectral Tuning and Adaptation to Distinct Light
Environments. *Mol Biol Evol* 2024, **41**(4):msae049.

62.   Rancilhac L, Irisarri I, Angelini C, Arntzen JW, Babik W, Bossuyt F, Künzel S, Lüddecke T,
Pasmans F, Sanchez E *et al*: **Phylotranscriptomic evidence for pervasive ancient**
hybridization among Old World salamanders. *Mol Phylogen Evol* 2021, **155**:106967.

63.   Dubey S, Maddalena T, Bonny L, Jeffries DL, Dufresnes C: **Population genomics of an**
exceptional hybridogenetic system of *Pelophylax* **water frogs.** *BMC Evol Biol* 2019,
19(1):164.

64. Dufresnes C, Ambu J, Galán P, Sequeira F, Viesca L, Choda M, Álvarez D, Alard B,
Suchan T, Künzel S: **Delimiting phylogeographic diversity in the genomic era:**
application to an Iberian endemic frog. *Zool J Linn Soc* 2023:zlad170.
65. Ovchinnikov V, Uliano-Silva M, Wilkinson M, Wood J, Smith M, Oliver K, Sims Y,
Torrance J, Suh A, McCarthy SA *et al*: **Caecilian Genomes Reveal the Molecular Basis**
of Adaptation and Convergent Evolution of Limblessness in Snakes and Caecilians.
Mol Biol Evol 2023, **40**(5).
66. Jetz W, Pyron RA: **The interplay of past diversification and evolutionary isolation**
with present imperilment across the amphibian tree of life. *Nature Ecology &*
Evolution 2018, **2**(5):850-858.
67. Siu-Ting K, Torres-Sánchez M, San Mauro D, Wilcockson D, Wilkinson M, Pisani D,
O'Connell MJ, Creevey CJ: **Inadvertent Paralog Inclusion Drives Artifactual**
Topologies and Timetree Estimates in Phylogenomics. *Mol Biol Evol* 2019,
36(6):1344-1356.
68. Hime PM, Lemmon AR, Lemmon ECM, Prendini E, Brown JM, Thomson RC, Kratovil
JD, Noonan BP, Pyron RA, Peloso PL: **Phylogenomics reveals ancient gene tree**
discordance in the amphibian tree of life. *Syst Biol* 2021, **70**(1):49-66.
69. Portik DM, Streicher JW, Blackburn DC, Moen DS, Hutter CR, Wiens JJ: **Redefining**
Possible: Combining Phylogenomic and Supersparse Data in Frogs. *Mol Biol Evol*
2023, **40**(5).
70. Sun Y-B, Xiong Z-J, Xiang X-Y, Liu S-P, Zhou W-W, Tu X-L, Zhong L, Wang L, Wu D-D,
Zhang B-L *et al*: **Whole-genome sequence of the Tibetan frog *Nanorana parkeri* and**
the comparative evolution of tetrapod genomes. *Proceedings of the National*
Academy of Sciences 2015, **112**(11):E1257-E1262.
71. Torres-Sánchez M, Gower DJ, Alvarez-Ponce D, Creevey CJ, Wilkinson M, San Mauro
D: **What lies beneath? Molecular evolution during the radiation of caecilian**
amphibians. *BMC Genomics* 2019, **20**(1):354.
72. Li J-T, Gao Y-D, Xie L, Deng C, Shi P, Guan M-L, Huang S, Ren J-L, Wu D-D, Ding L:
Comparative genomic investigation of high-elevation adaptation in ectothermic
snakes. *Proceedings of the National Academy of Sciences* 2018, **115**(33):8406-8411.
73. Storz JF: **High-Altitude Adaptation: Mechanistic Insights from Integrated Genomics**
and Physiology. *Mol Biol Evol* 2021, **38**(7):2677-2691.
74. Yu L, Wang G-D, Ruan J, Chen Y-B, Yang C-P, Cao X, Wu H, Liu Y-H, Du Z-L, Wang X-P:
Genomic analysis of snub-nosed monkeys (*Rhinopithecus*) identifies genes and
processes related to high-altitude adaptation. *Nat Genet* 2016, **48**(8):947-952.
75. Seimon TA, Seimon A, Daszak P, Halloy SR, Schloegel LM, Aguilar CA, Sowell P, Hyatt
AD, Konecky B, E Simmons J: **Upward range extension of Andean anurans and**
chytridiomycosis to extreme elevations in response to tropical deglaciation. *Global*
Change Biol 2007, **13**(1):288-299.
76. Acosta-Galvis AR: **Ranas, salamandras y caecilias (Tetrapoda: Amphibia) de**
Colombia. *Biota colombiana* 2000, **1**(3).
77. Yang W, Qi Y, Fu J: **Genetic signals of high-altitude adaptation in amphibians: a**
comparative transcriptome analysis. *BMC Genet* 2016, **17**(1):134.
78. Lu B, Jin H, Fu J: **Molecular convergent and parallel evolution among four high-**
elevation anuran species from the Tibetan region. *BMC Genomics* 2020, **21**:1-14.

79. Wang G-D, Zhang B-L, Zhou W-W, Li Y-X, Jin J-Q, Shao Y, Yang H-c, Liu Y-H, Yan F, Chen
H-M: **Selection and environmental adaptation along a path to speciation in the**
Tibetan frog *Nanorana parkeri*. *Proc Natl Acad Sci USA* 2018, **115**(22):E5056-E5065.
80. Sun Y-B, Fu T-T, Jin J-Q, Murphy RW, Hillis DM, Zhang Y-P, Che J: **Species groups**
distributed across elevational gradients reveal convergent and continuous genetic
adaptation to high elevations. *Proceedings of the National Academy of Sciences*
2018, **115**(45):E10634-E10641.
81. Cayuela H, Dorant Y, Forester BR, Jeffries DL, Mccaffery RM, Eby LA, Hossack BR,
Gippet JM, Pilliod DS, Chris Funk W: **Genomic signatures of thermal adaptation are**
associated with clinal shifts of life history in a broadly distributed frog. *J Anim Ecol*
2022, **91**(6):1222-1238.
82. Hutchison VH, Haines HB, Engbretson G: **Aquatic life at high altitude: Respiratory**
adaptations in the lake titicaca frog, *Telmatobius culeus*. *Respiration Physiology*
1976, **27**(1):115-129.
83. Dunn ER: **The salamanders of the family Plethodontidae**, vol. 7: Smith College; 1926.
84. Daly J, Garraffo H, Pannell L, Spande T, Severini C, Erspamer V: **Alkaloids from**
Australian frogs (Myobatrachidae): pseudophrynamines and pumiliotoxins. *J Nat*
Prod 1990, **53**(2):407-421.
85. Darst CR, Cummings ME: **Predator learning favours mimicry of a less-toxic model in**
poison frogs. *Nature* 2006, **440**(7081):208-211.
86. Hayes RA, Piggott AM, Dalle K, Capon RJ: **Microbial biotransformation as a source of**
chemical diversity in cane toad steroid toxins. *Bioorganic & Medicinal Chemistry*
Letters 2009, **19**(6):1790-1792.
87. Tóth Z, Kurali A, Móricz ÁM, Hettyey A: **Changes in Toxin Quantities Following**
Experimental Manipulation of Toxin Reserves in *Bufo bufo* Tadpoles. *J Chem Ecol*
2019, **45**(3):253-263.
88. Vaelli PM, Theis KR, Williams JE, O'Connell LA, Foster JA, Eisthen HL: **The skin**
microbiome facilitates adaptive tetrodotoxin production in poisonous newts. *eLife*
2020, **9**:e53898.
89. Daly JW, Martin Garraffo H, Spande TF, Jaramillo C, Stanley Rand A: **Dietary source**
for skin alkaloids of poison frogs (Dendrobatidae)? *J Chem Ecol* 1994, **20**:943-955.
90. Darst CR, Menéndez-Guerrero PA, Coloma LA, Cannatella DC: **Evolution of dietary**
specialization and chemical defense in poison frogs (Dendrobatidae): a
comparative analysis. *The American Naturalist* 2005, **165**(1):56-69.
91. Caty SN, Alvarez-Buylla A, Byrd GD, Vidoudez C, Roland AB, Tapia EE, Budnik B,
Trauger SA, Coloma LA, O'Connell LA: **Molecular physiology of chemical defenses in**
a poison frog. *J Exp Biol* 2019, **222**(12):jeb204149.
92. Alvarez-Buylla A, Fischer M-T, Moya Garzon MD, Rangel AE, Tapia EE, Tanzo JT, Soh
HT, Coloma LA, Long JZ, O'Connell LA: **Binding and sequestration of poison frog**
alkaloids by a plasma globulin. *eLife* 2023, **12**:e85096.
93. Abderemane-Ali F, Rossen ND, Kobiela ME, Craig RA, Garrison CE, Chen Z, Colleran
CM, O'Connell LA, Du Bois J, Dumbacher JP: **Evidence that toxin resistance in poison**
birds and frogs is not rooted in sodium channel mutations and may rely on "toxin
sponge" proteins. *J Gen Physiol* 2021, **153**(9):e202112872.
94. Márquez R, Ramírez-Castañeda V, Amézquita A: **Does batrachotoxin autoresistance**
coevolve with toxicity in *Phyllobates* poison-dart frogs? *Evolution* 2019, **73**(2):390-
400.

95. Tarvin RD, Borghese CM, Sachs W, Santos JC, Lu Y, O'Connell LA, Cannatella DC, Harris
RA, Zakon HH: **Interacting amino acid replacements allow poison frogs to evolve**
epibatidine resistance. *Science* 2017, **357**(6357):1261-1266.
96. Tarvin RD, Santos JC, O'Connell LA, Zakon HH, Cannatella DC: **Convergent**
substitutions in a sodium channel suggest multiple origins of toxin resistance in
poison frogs. *Mol Biol Evol* 2016, **33**(4):1068-1080.
97. Shibao PYT, Cologna CT, Morandi-Filho R, Wiezel GA, Fujimura PT, Ueira-Vieira C,
Arantes EC: **Deep sequencing analysis of toad** *Rhinella schneideri* **skin glands and**
partial biochemical characterization of its cutaneous secretion. *J Venom Anim Tox*
incl Trop Dis 2018, **24**(1):36.
98. Torres-Sánchez M, Wilkinson M, Gower DJ, Creevey CJ, San Mauro D: **Insights into**
the skin of caecilian amphibians from gene expression profiles. *BMC Genomics*
2020, **21**(1):515.
99. Liscano Martinez Y, Arenas Gómez CM, Smith J, Delgado JP: **A tree frog (***Boana***
pugnax) dataset of skin transcriptome for the identification of biomolecules with**
potential antimicrobial activities. *Data in Brief* 2020, **32**:106084.
100. Lan Y, He L, Dong X, Tang R, Li W, Wang J, Wang L, Yue B, Price M, Guo T *et al*:
Comparative transcriptomes of three different skin sites for the Asiatic toad (*Bufo***
gargarizans)**. *PeerJ* 2022, **10**:e12993.
101. Lv Y, Li Y, Wen Z, Shi Q: **Transcriptomic and gene-family dynamic analyses reveal**
gene expression pattern and evolution in toxin-producing tissues of Asiatic toad
(*Bufo gargarizans***)**. *Frontiers in Ecology and Evolution* 2022, **10**:924248.
102. Mohammadi S, Herrera-Álvarez S, Yang L, Rodriguez-Ordonez MdP, Zhang K, Storz JF,
Dobler S, Crawford AJ, Andolfatto P: **Constraints on the evolution of toxin-resistant**
Na, K-ATPases have limited dependence on sequence divergence. *PLoS Genet* 2022,
18(8):e1010323.
103. Mohammadi S, Yang L, Harpak A, Herrera-Álvarez S, del Pilar Rodríguez-Ordoñez M,
Peng J, Zhang K, Storz JF, Dobler S, Crawford AJ: **Concerted evolution reveals co-**
adapted amino acid substitutions in Na+ K+-ATPase of frogs that prey on toxic
toads. *Curr Biol* 2021, **31**(12):2530-2538. e2510.
104. Hutchinson DA, Mori A, Savitzky AH, Burghardt GM, Wu X, Meinwald J, Schroeder FC:
Dietary sequestration of defensive steroids in nuchal glands of the Asian snake
Rhabdophis tigrinus. *Proceedings of the National Academy of Sciences* 2007,
104(7):2265-2270.
105. Brodie III ED, Brodie Jr ED: **Tetrodotoxin resistance in garter snakes: an evolutionary**
response of predators to dangerous prey. *Evolution* 1990, **44**(3):651-659.
106. Mancuso M, Zaman S, Maddock ST, Kamei RG, Salazar-Valenzuela D, Wilkinson M,
Roelants K, Fry BG: **Resistance Is Not Futile: Widespread Convergent Evolution of**
Resistance to Alpha-Neurotoxic Snake Venoms in Caecilians (Amphibia:
Gymnophiona). *International Journal of Molecular Sciences* 2023, **24**(14):11353.
107. Symula R, Schulte R, Summers K: **Molecular phylogenetic evidence for a mimetic**
radiation in Peruvian poison frogs supports a Müllerian mimicry hypothesis.
Proceedings of the Royal Society of London Series B: Biological Sciences 2001,
268(1484):2415-2421.
108. Daly JW, Brown GB, Mensah-Dwumah M, Myers CW: **Classification of skin alkaloids**
from neotropical poison-dart frogs (Dendrobatidae). *Toxicon* 1978, **16**(2):163-188.

109. Stuckert AMM, Moore E, Coyle KP, Davison I, MacManes MD, Roberts R, Summers K: **Variation in pigmentation gene expression is associated with distinct aposematic color morphs in the poison frog *Dendrobates auratus*.** *BMC Evol Biol* 2019, **19**(1):85.

110. Stuckert AM, Chouteau M, McClure M, LaPolice TM, Linderoth T, Nielsen R, Summers K, MacManes MD: **The genomics of mimicry: gene expression throughout development provides insights into convergent and divergent phenotypes in a Müllerian mimicry system.** *Mol Ecol* 2021, **30**(16):4039-4061.

111. Twomey E, Johnson JD, Castroviejo-Fisher S, Van Bocxlaer I: **A ketocarotenoid-based colour polymorphism in the Sira poison frog *Ranitomeya sirensis* indicates novel gene interactions underlying aposematic signal variation.** *Mol Ecol* 2020, **29**(11):2004-2015.

112. Twomey E, Kain M, Claeys M, Summers K, Castroviejo-Fisher S, Van Bocxlaer I: **Mechanisms for color convergence in a mimetic radiation of poison frogs.** *The American Naturalist* 2020, **195**(5):E132-E149.

113. Linderoth T, Aguilar-Gómez D, White E, Twomey E, Stuckert A, Bi K, Ko A, Graham N, Rocha JL, Chang J *et al*: **Genetic basis of aposematic coloration in a mimetic radiation of poison frogs.** *bioRxiv* 2023:2023.2004.2020.537757.

114. Stuckert AM, Freeborn L, Howell KA, Yang Y, Nielsen R, Richards-Zawacki C, MacManes MD: **Transcriptomic analyses during development reveal mechanisms of integument structuring and color production.** *Evol Ecol* 2023:1-22.

115. Burgon JD, Vieites DR, Jacobs A, Weidt SK, Gunter HM, Steinfartz S, Burgess K, Mable BK, Elmer KR: **Functional colour genes and signals of selection in colour-polymorphic salamanders.** *Mol Ecol* 2020, **29**(7):1284-1299.

116. Fischer EK, Roland AB, Moskowitz NA, Tapia EE, Summers K, Coloma LA, O'Connell LA: **The neural basis of tadpole transport in poison frogs.** *Proceedings of the Royal Society B: Biological Sciences* 2019, **286**(1907):20191084.

117. Crump ML: **Parental Care among the Amphibia.** In: *Advances in the Study of Behavior.* Edited by Rosenblatt JS, Snowdon CT, vol. 25: Academic Press; 1996: 109-144.

118. Wu W, Gao YD, Jiang DC, Lei J, Ren JL, Liao WB, Deng C, Wang Z, Hillis DM, Zhang YP *et al*: **Genomic adaptations for arboreal locomotion in Asian flying treefrogs.** *Proc Natl Acad Sci U S A* 2022, **119**(13):e2116342119.

119. Blackburn DC, Gray JA, Stanley EL: **The only "lungless" frog has a glottis and lungs.** *Curr Biol* 2024, **34**(10):R492-R493.

120. Lewis ZR, Kerney R, Hanken J: **Developmental basis of evolutionary lung loss in plethodontid salamanders.** *Sci Adv* 2022, **8**(33):eabo6108.

121. Heiss E, Natchev N, Salaberger D, Gumpenberger M, Rabanser A, Weisgram J: **Hurt yourself to hurt your enemy: new insights on the function of the bizarre antipredator mechanism in the salamandrid *Pleurodeles waltl*.** *J Zool* 2010, **280**(2):156-162.

122. Brodie Jr ED, Nussbaum RA, DiGiovanni M: **Antipredator adaptations of Asian salamanders (Salamandridae).** *Herpetologica* 1984:56-68.

123. Mailho-Fontana PL, Antoniazzi MM, Coelho GR, Pimenta DC, Fernandes LP, Kupfer A, Brodie ED, Jared C: **Milk provisioning in oviparous caecilian amphibians.** *Science* 2024, **383**(6687):1092-1095.

124. Kupfer A, Müller H, Antoniazzi MM, Jared C, Greven H, Nussbaum RA, Wilkinson M:
Parental investment by skin feeding in a caecilian amphibian. *Nature* 2006,
440(7086):926-929.
125. Liu Y, Jones CD, Day LB, Summers K, Burmeister SS: **Cognitive phenotype and**
differential gene expression in a hippocampal homologue in two species of frog.
Integr Comp Biol 2020, **60**(4):1007-1023.
126. Luedtke JA, Chanson J, Neam K, Hobin L, Maciel AO, Catenazzi A, Borzée A, Hamidy A,
Aowphol A, Jean A *et al*: **Ongoing declines for the world's amphibians in the face of**
emerging threats. *Nature* 2023, **622**(7982):308-314.
127. Vacher JP, Chave J, Ficetola FG, Sommeria-Klein G, Tao S, Thébaud C, Blanc M,
Camacho A, Cassimiro J, Colston TJ: **Large-scale DNA-based survey of frogs in**
Amazonia suggests a vast underestimation of species richness and endemism. *J*
Biogeogr 2020, **47**(8):1781-1791.
128. Oliver PM, Bower DS, McDonald PJ, Kraus F, Luedtke J, Neam K, Hobin L, Chauvenet
AL, Allison A, Arida E: **Melanesia holds the world's most diverse and intact insular**
amphibian fauna. *Communications biology* 2022, **5**(1):1182.
129. Liu J, Slik F, Zheng S, Lindenmayer DB: **Undescribed species have higher extinction**
risk than known species. *Conservation Letters* 2022, **15**(3):e12876.
130. Re:wild, Earth S, Group ISAS: **State of the World's Amphibians: The Second Global**
Amphibian Assessment. In. Texas, USA: Re:wild; 2023.
131. Gower DJ, San Mauro D, Giri V, Bhatta G, Govindappa V, Kotharambath R, Oommen
OV, Fatih FA, Mackenzie-Dodds JA, Nussbaum RA: **Molecular systematics of caeciliid**
caecilians (Amphibia: Gymnophiona) of the Western Ghats, India. *Mol Phylogen*
Evol 2011, **59**(3):698-707.
132. Forester BR, Beever EA, Darst C, Szymanski J, Funk WC: **Linking evolutionary**
potential to extinction risk: applications and future directions. *Front Ecol Environ*
2022, **20**(9):507-515.
133. Nemesházi E, Bókony V: **Interplay of genotypic and thermal sex determination**
shapes climatic distribution in herpetofauna. *bioRxiv* 2024:2024.2004.2021.589911.
134. Wollenberg Valero KC, Marshall JC, Bastiaans E, Caccone A, Camargo A, Morando M,
Niemiller ML, Pabijan M, Russello MA, Sinervo B *et al*: **Patterns, Mechanisms and**
Genetics of Speciation in Reptiles and Amphibians. *Genes (Basel)* 2019, **10**(9):646.
135. IUCN SSC ASG: **Draft for Open Consultation. The Amphibian Conservation Action**
Plan (ACAP): A status review and roadmap for global amphibian conservation.
EcoEvoRxiv 2022.
136. Hogg CJ: **Translating genomic advances into biodiversity conservation**. *Nat Rev*
Genet 2023, **25**(5):362-373.
137. Whiteley AR, Fitzpatrick SW, Funk WC, Tallmon DA: **Genetic rescue to the rescue**.
Trends Ecol Evol 2015, **30**(1):42-49.
138. Pabijan M, Palomar G, Antunes B, Antoł W, Zieliński P, Babik W: **Evolutionary**
principles guiding amphibian conservation. *Evol Appl* 2020, **13**(5):857-878.
139. Theissinger K, Fernandes C, Formenti G, Bista I, Berg PR, Bleidorn C, Bombarely A,
Crottini A, Gallo GR, Godoy JA *et al*: **How genomics can help biodiversity**
conservation. *Trends Genet* 2023, **39**(7):545-559.
140. Trumbo D, Hardy B, Crockett H, Muths E, Forester B, Cheek R, Zimmerman S, Corey-
Rivas S, Bailey L, Funk WC: **Conservation genomics of an endangered montane**

amphibian reveals low population structure, low genomic diversity, and selection pressure from disease. *Mol Ecol* 2023, **32**(24):6777-6795.

141. Torres-Sánchez M, Longo AV: **Linking pathogen-microbiome-host interactions to explain amphibian population dynamics**. *Mol Ecol* 2022, **31**(22):5784-5794.

142. Lopes CM, Baêta D, Valentini A, Lyra ML, Sabbag AF, Gasparini JL, Dejean T, Haddad CFB, Zamudio KR: **Lost and found: Frogs in a biodiversity hotspot rediscovered with environmental DNA**. *Mol Ecol* 2021, **30**(13):3289-3298.

143. Cutajar TP, Rowley JJL: **Surveying frogs from the bellies of their parasites: Invertebrate-derived DNA as a novel survey method for frogs**. *Global Ecology and Conservation* 2020, **22**:e00978.

144. Villacorta-Rath C, Hoskin CJ, Strugnell JM, Burrows D: **Long distance (> 20 km) downstream detection of endangered stream frogs suggests an important role for eDNA in surveying for remnant amphibian populations**. *PeerJ* 2021, **9**:e12013.

145. Allendorf FW, Funk WC, Aitken SN, Byrne M, Luikart G: **Conservation and the genomics of populations**: Oxford University Press; 2022.

146. Fischman RL, Ruhl JB, Forester BR, Lama TM, Kardos M, Rojas GA, Robinson NA, Shirey PD, Lamberti GA, Ando AW *et al*: **A landmark environmental law looks ahead**. *Science* 2023, **382**(6677):1348-1355.

147. Programme UE: **Kunming-Montreal Global Biodiversity Framework — CBD/COP/15/L25**. In. Edited by Diversity UCoB. Montreal Canada; 2022.

148. Kosch TA, Waddle AW, Cooper CA, Zenger KR, Garrick DJ, Berger L, Skerratt LF: **Genetic approaches for increasing fitness in endangered species**. *Trends Ecol Evol* 2022, **37**(4):332-345.

149. Kosch T, Silva C, Brannelly L, Roberts A, Lau Q, Marantelli G, Berger L, Skerratt L: **Genetic potential for disease resistance in critically endangered amphibians decimated by chytridiomycosis**. *Anim Conserv* 2019, **22**(3):238-250.

150. Savage AE, Gratwicke B, Hope K, Bronikowski E, Fleischer RC: **Sustained immune activation is associated with susceptibility to the amphibian chytrid fungus**. *Mol Ecol* 2020, **29**(15):2889-2903.

151. Bataille A, Cashins SD, Grogan L, Skerratt LF, Hunter D, McFadden M, Scheele B, Brannelly LA, Macris A, Harlow PS: **Susceptibility of amphibians to chytridiomycosis is associated with MHC class II conformation**. *Proc R Soc Lond, Ser B: Biol Sci* 2015, **282**(1805):20143127.

152. Knapp RA, Wilber MQ, Byrne AQ, Joseph MB, Smith TC, Rothstein AP, Grasso RL, Rosenblum EB: **Reintroduction of resistant frogs facilitates landscape-scale recovery in the presence of a lethal fungal disease**. *bioRxiv* 2023:2023.005.2022.541534.

153. Byrne PG, Silla AJ: **An experimental test of the genetic consequences of population augmentation in an amphibian**. *Conserv Sci Pract* 2020, **2**(6):e194.

154. Liddell E, Sunnucks P, Cook CN: **To mix or not to mix gene pools for threatened species management? Few studies use genetic data to examine the risks of both actions, but failing to do so leads disproportionately to recommendations for separate management**. *Biol Conserv* 2021, **256**:109072.

155. Kyriazis CC, Wayne RK, Lohmueller KE: **Strongly deleterious mutations are a primary determinant of extinction risk due to inbreeding depression**. *Evolution Letters* 2021, **5**:33-47.

156. Berger L, Skerratt LF, Kosch TA, Brannelly LA, Webb RJ, Waddle AW: **Advances in**
Managing Chytridiomycosis for Australian Frogs: *Gradarius Firmus* **Victoria**. *Annu*
Rev Anim Biosci 2024, **12**(1):113-133.
157. Collier-Robinson L, Rayne A, Rupene M, Thoms C, Steeves T: **Embedding indigenous**
principles in genomic research of culturally significant species. *N Z J Ecol* 2019,
43(3):1-9.
158. Rayne A, Blair S, Dale M, Flack B, Hollows J, Moraga R, Parata RN, Rupene M, Tamati-
Elliffe P, Wehi PM *et al*: **Weaving place-based knowledge for culturally significant**
species in the age of genomics: Looking to the past to navigate the future. *Evol Appl*
2022, **15**(5):751-772.
159. Mc Cartney AM, Head MA, Tsosie KS, Sterner B, Glass JR, Paez S, Geary J, Hudson M:
Indigenous peoples and local communities as partners in the sequencing of global
eukaryotic biodiversity. *npj Biodiversity* 2023, **2**(1):8.
160. Carroll S, Garba I, Figueroa-Rodríguez O, Holbrook J, Lovett R, Materechera S, Parsons
M, Raseroka K, Rodriguez-Lonebear D, Rowe R: **The CARE principles for indigenous**
data governance. *Data Science Journal* 2020, **19**(43):1-21.
161. Ramírez-Castañeda V, Westeen EP, Frederick J, Amini S, Wait DR, Achmadi AS,
Andayani N, Arida E, Arifin U, Bernal MA *et al*: **A set of principles and practical**
suggestions for equitable fieldwork in biology. *Proceedings of the National Academy*
of Sciences 2022, **119**(34):e2122667119.
162. Feres MVC: **Biodiversity, traditional knowledge and patent rights: The case study of**
Phyllomedusa bicolor. *Revista Direito GV* 2022, **18**:e2205.
163. Te Aika B, Liggins L, Rye C, Perkins EO, Huh J, Brauning R, Godfery T, Black MA:
Aotearoa genomic data repository: An āhuru mōwai for taonga species sequencing
data. *Mol Ecol Resour* 2023, **00**:1-14.
164. Buckner JC, Sanders RC, Faircloth BC, Chakrabarty P: **The critical importance of**
vouchers in genomics. *eLife* 2021, **10**:e68264.
165. Golan J, Riddle K, Hudson M, Anderson J, Kusabs N, Coltman T: **Benefit sharing: Why**
inclusive provenance metadata matter. *Front Genet* 2022, **13**:1014044.
166. Duellman WE, Trueb L: **Biology of Amphibians**: JHU press; 1994.
167. Dahn HA, Mountcastle J, Balacco J, Winkler S, Bista I, Schmitt AD, Pettersson OV,
Formenti G, Oliver K, Smith M *et al*: **Benchmarking ultra-high molecular weight DNA**
preservation methods for long-read and long-range sequencing. *GigaScience* 2022,
11.
168. Forzán MJ, Heatley J, Russell KE, Horney B: **Clinical pathology of amphibians: a**
review. *Veterinary Clinical Pathology* 2017, **46**(1):11-33.
169. IUCN: **IUCN policy statement on research involving species at risk of extinction**. In.
Gland, Switzerland: International Union for the Conservation of Nature.; 1989.
170. Ambu J, Dufresnes C: **Buccal swabs for amphibian genomics**. *Amphibia-Reptilia*
2023, **44**(2):249-255.
171. Oyler-McCance SJ, Ryan MJ, Sullivan BK, Fike JA, Cornman RS, Giermakowski JT,
Zimmerman SJ, Harrow RL, Hedwall SJ, Hossack BR *et al*: **Genetic connectivity in the**
Arizona toad (*Anaxyrus microscaphus*): implications for conservation of a stream
dwelling amphibian in the arid Southwestern United States. *Conserv Genet* 2024,
25(3):835-848.

172. Farquharson KA, McLennan EA, Belov K, Hogg CJ: **The genome sequence of the**
critically endangered Kroombit tinkerfrog (*Taudactylus pleione***)**. *F1000Research*
2023, **12**:845.
173. Minich JJ, Moore ML, Allsing NA, Aylward A, Murray ER, Tran L, Michael TP:
Generating high-quality plant and fish reference genomes from field-collected
specimens by optimizing preservation. *Communications Biology* 2023, **6**(1):1246.
174. Blom MPK: **Opportunities and challenges for high-quality biodiversity tissue**
archives in the age of long-read sequencing. *Mol Ecol* 2021, **30**(23):5935-5948.
175. Alves RJV, Weksler M, Oliveira JA, Buckup PA, SANTANA HR, PERACCHI AL, KELLNER
AW, ALEIXO A, LANGGUTH A, ALMEIDA A: **Brazilian legislation on genetic heritage**
harms Biodiversity Convention goals and threatens basic biology research and
education. In., vol. 90: SciELO Brasil; 2018: 1279-1284.
176. Alexander GJ, Tollev KA, Maritz B, McKechnie A, Manger P, Thomson RL, Schradin C,
Fuller A, Meyer L, Hetem RS: **Excessive red tape is strangling biodiversity research in**
South Africa. *S Afr J Sci* 2021, **117**(9-10):1-4.
177. Haelewaters D, Hofmann TA, Romero-Olivares AL: **Ten simple rules for Global North**
researchers to stop perpetuating helicopter research in the Global South. *PLoS*
Comp Biol 2021, **17**(8):e1009277.
178. de Vos A, Schwartz MW: **Confronting parachute science in conservation**. *Conserv Sci*
Pract 2022, **4**(5):e12681.
179. Watsa M, Erkenswick GA, Pomerantz A, Prost S: **Portable sequencing as a teaching**
tool in conservation and biodiversity research. *PLoS Biol* 2020, **18**(4):e3000667.
180. Armenteras D: **Guidelines for healthy global scientific collaborations**. *Nature*
Ecology & Evolution 2021, **5**(9):1193-1194.
181. Fong JJ, Blom MP, Aowphol A, McGuire JA, Sutcharit C, Soltis PS: **Recent advances in**
museomics: revolutionizing biodiversity research. *Frontiers in Ecology and Evolution*
2023, **11**:1188172.
182. Raxworthy CJ, Smith BT: **Mining museums for historical DNA: advances and**
challenges in museomics. *Trends Ecol Evol* 2021, **36**(11):1049-1060.
183. Roycroft E, Moritz C, Rowe KC, Moussalli A, Eldridge MDB, Portela Miguez R, Piggott
MP, Potter S: **Sequence Capture From Historical Museum Specimens: Maximizing**
Value for Population and Phylogenomic Studies. *Frontiers in Ecology and Evolution*
2022, **10**:931644.
184. Speer KA, Hawkins MTR, Flores MFC, McGowen MR, Fleischer RC, Maldonado JE,
Campana MG, Muletz-Wolz CR: **A comparative study of RNA yields from museum**
specimens, including an optimized protocol for extracting RNA from formalin-fixed
specimens. *Frontiers in Ecology and Evolution* 2022, **10**:953131.
185. Dabney J, Knapp M, Glocke I, Gansauge M-T, Weihmann A, Nickel B, Valdiosera C,
García N, Pääbo S, Arsuaga J-L: **Complete mitochondrial genome sequence of a**
Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments.
Proceedings of the National Academy of Sciences 2013, **110**(39):15758-15763.
186. Gansauge M-T, Gerber T, Glocke I, Korlević P, Lippik L, Nagel S, Riehl LM, Schmidt A,
Meyer M: **Single-stranded DNA library preparation from highly degraded DNA using**
T4 DNA ligase. *Nucleic Acids Res* 2017, **45**(10):e79-e79.
187. Straube N, Lyra ML, Paijmans JL, Preick M, Basler N, Penner J, Rödel MO, Westbury
MV, Haddad CF, Barlow A: **Successful application of ancient DNA extraction and**

library construction protocols to museum wet collection specimens. *Mol Ecol*
Resour 2021, **21**(7):2299-2315.

188. Rancilhac L, Bruy T, Scherz MD, Pereira EA, Preick M, Straube N, Lyra ML, Ohler A,
Streicher JW, Andreone F: **Target-enriched DNA sequencing from historical type**
material enables a partial revision of the Madagascar giant stream frogs (genus
***Mantidactylus)**. *J Nat Hist* 2020, **54**(1-4):87-118.

189. Scherz MD, Rasolonjatovo SM, Köhler J, Rancilhac L, Rakotoarison A, Raselimanana
AP, Ohler A, Preick M, Hofreiter M, Glaw F: **'Barcode fishing' for archival DNA from**
historical type material overcomes taxonomic hurdles, enabling the description of a
new frog species. *Sci Rep* 2020, **10**(1):19109.

190. Dalén L, Heintzman PD, Kapp JD, Shapiro B: **Deep-time paleogenomics and the limits**
of DNA survival. *Science* 2023, **382**(6666):48-53.

191. Ramírez JP, Jaramillo CA, Lindquist ED, Crawford AJ, Ibáñez R: **Recent and Rapid**
Radiation of the Highly Endangered Harlequin Frogs (*Atelopus*) into Central
America Inferred from Mitochondrial DNA Sequences. *Diversity* 2020, **12**(9):360.

192. Renner SS, Scherz MD, Schoch CL, Gottschling M, Vences M: **DNA sequences from**
type specimens and type strains—how to increase their number and improve their
annotation in NCBI GenBank and related databases. *Syst Biol* 2023:syad068.

193. Rakotoarison A, Scherz MD, Mullin KE, Crottini A, Petzold A, Ranjanaharisoa FA,
Maheritafika HMR, Rafanoharana JM, Raherinjatovo H, Andreone F: **Gray versus**
yellow ventral coloration: Identity, distribution, color polymorphism and molecular
relationships of the microhylid frog *Platypelis mavomavo* Andreone, Fenolio &
Walvoord, 2003. *Zootaxa* 2023, **5352**(2):221-234.

194. Evans BJ, Gansauge M-T, Stanley EL, Furman BLS, Cauret CMS, Ofori-Boateng C,
Gvoždík V, Streicher JW, Greenbaum E, Tinsley RC *et al*: ***Xenopus fraseri*: Mr. Fraser,**
where did your frog come from? *PLOS ONE* 2019, **14**(9):e0220892.

195. Wong K-C: **Big data challenges in genome informatics**. *Biophysical Reviews* 2019,
11(1):51-54.

196. Zuo B, Nneji LM, Sun Y-B: **Comparative genomics reveals insights into anuran**
genome size evolution. *BMC Genomics* 2023, **24**(1):379.

197. Alkan C, Sajjadian S, Eichler EE: **Limitations of next-generation genome sequence**
assembly. *Nat Methods* 2011, **8**(1):61-65.

198. Treangen TJ, Salzberg SL: **Repetitive DNA and next-generation sequencing:**
computational challenges and solutions. *Nat Rev Genet* 2012, **13**(1):36-46.

199. Sun Y, Shang L, Zhu Q-H, Fan L, Guo L: **Twenty years of plant genome sequencing:**
achievements and challenges. *Trends Plant Sci* 2022, **27**(4):391-401.

200. Mable B, Alexandrou M, Taylor M: **Genome duplication in amphibians and fish: an**
extended synthesis. *J Zool* 2011, **284**(3):151-182.

201. Schmid M, Evans BJ, Bogart JP: **Polyploidy in Amphibia**. *Cytogenet Genome Res* 2015,
145(3-4):315-330.

202. Session AM, Uno Y, Kwon T, Chapman JA, Toyoda A, Takahashi S, Fukui A, Hikosaka A,
Suzuki A, Kondo M *et al*: **Genome evolution in the allotetraploid frog *Xenopus***
laevis. *Nature* 2016, **538**(7625):336-343.

203. Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, Uliano-Silva M, Chow
W, Fungtammasan A, Kim J *et al*: **Towards complete and error-free genome**
assemblies of all vertebrate species. *Nature* 2021, **592**(7856):737-746.

204. Consortium TU: **UniProt: the Universal Protein Knowledgebase in 2023**. *Nucleic*
Acids Res 2022, **51**(D1):D523-D531.
205. Dimitrakopoulou D, Khwatenge CN, James-Zorn C, Paiola M, Bellin EW, Tian Y,
Sundararaj N, Polak EJ, Grayfer L, Barnard D *et al*: **Advances in the *Xenopus***
immunome: Diversification, expansion, and contraction. *Dev Comp Immunol* 2023,
145:104734.
206. Bertola LV, Hoskin CJ, Jones DB, Zenger KR, McKnight DT, Higgie M: **The first linkage**
map for Australo-Papuan Treefrogs (family: Pelodryadidae) reveals the sex-
determination system of the Green-eyed Treefrog (*Litoria serrata*). *Heredity* 2023,
131(4):263-272.
207. Silla AJ, Byrne PG: **The Role of Reproductive Technologies in Amphibian**
Conservation Breeding Programs. *Annu Rev Anim Biosci* 2019, **7**(1):499-519.
208. Banach M, Edholm E-S, Robert J: **Exploring the functions of nonclassical MHC class**
Ib genes in *Xenopus laevis* by the CRISPR/Cas9 system. *Dev Biol* 2016, **426**(2):261-
269.
209. Grainger RM: ***Xenopus tropicalis* as a model organism for genetics and genomics:**
past, present, and future. *Xenopus Protocols: Post-Genomic Approaches* 2012:3-15.
210. Sousounis K, Courtemanche K, Whited JL: **A Practical Guide for CRISPR-Cas9-Induced**
Mutations in Axolotls. In: *Salamanders: Methods and Protocols.* Springer; 2022: 335-
349.
211. Wang F, Shi Z, Cui Y, Guo X, Shi Y-B, Chen Y: **Targeted gene disruption in *Xenopus***
***laevis* using CRISPR/Cas9**. *Cell & bioscience* 2015, **5**:1-5.
212. Douglas AJ, Todd LA, Katzenback BA: **The amphibian invitrome: Past, present, and**
future contributions to our understanding of amphibian immunity. *Dev Comp*
Immunol 2023, **142**:104644.
213. Bui-Marinos MP, Todd LA, Douglas AJ, Katzenback BA: **So, you want to create a frog**
cell line? A guide to establishing frog skin cell lines from tissue explants. *MethodsX*
2022, **9**:101693.
214. Stiller J, Feng S, Chowdhury A-A, Rivas-González I, Duchêne DA, Fang Q, Deng Y,
Kozlov A, Stamatakis A, Claramunt S *et al*: **Complexity of avian evolution revealed by**
family-level genomes. *Nature* 2024, **629**(818):851-860.
215. Lees AC, Pimm SL: **Species, extinct before we know them?** *Curr Biol* 2015,
25(5):R177-R180.
1464