B cell receptor dependent enhancement of dengue virus infection

Chad Gebo[1], Céline S.C. Hardy[1], Benjamin D. McElvany[2], Nancy R. Graham[2], Joseph Lu[1,3], Shima Moradpour[1], Jeffrey R Currier[4], Heather Friberg[4], Gregory D Gromowski[4], Stephen J. Thomas[1,3], Gary C. Chan[1], Sean A. Diehl[2], Adam T. Waickman[1,3,*]

[1]Department of Microbiology and Immunology, State University of New York Upstate Medical University, Syracuse, NY 13210, USA.

[2]Department of Microbiology and Molecular Genetics, The University of Vermont Larner College of Medicine, Vaccine Testing Center, Burlington, Vermont, USA.

[3]Global Health Institute, State University of New York Upstate Medical University, Syracuse, NY 13210, USA.

[4]Viral Diseases Branch, Walter Reed Army Institute of Research, Silver Spring, MD, 20910, USA

*Corresponding author: waickmaa@upstate.edu

Key words: dengue virus, immune mediated enhancement, B cells, B cell receptor

Abstract

Dengue virus (DENV) is the causative agent of dengue, a mosquito-borne disease that represents a significant and growing public health burden around the world. A unique pathophysiological feature of dengue is immune-mediated enhancement, wherein preexisting immunity elicited by a primary infection can enhance the severity of a subsequent infection by a heterologous DENV serotype. A leading mechanistic explanation for this phenomenon is antibody dependent enhancement (ADE), where sub-neutralizing concentrations of DENV-specific IgG antibodies facilitate entry of DENV into $Fc\gamma R$ expressing cells such as monocytes, macrophages, and dendritic cells. Accordingly, this model posits that phagocytic mononuclear cells are the primary reservoir of DENV. However, analysis of samples from individuals experiencing acute DENV infection reveals that B cells are the largest reservoir of infected circulating cells, representing a disconnect in our understanding of immune-mediated DENV tropism. In this study, we demonstrate that the expression of a DENV-specific B cell receptor (BCR) renders cells highly susceptible to DENV infection, with the infection-enhancing activity of the membrane-restricted BCR correlating with the ADE potential of the IgG version of the antibody. In addition, we observed that the frequency of DENV-infectable B cells increases in previously flavivirus-naïve volunteers after a primary DENV infection. These findings suggest that BCR-dependent infection of B cells is a novel mechanism immune-mediated enhancement of DENV-infection.

INTRODUCTION

Dengue virus (DENV) is a prevalent arboviral pathogen that poses a significant global public health burden. Transmitted by the bite of infected *Aedes* family mosquitos, DENV co-circulates as four genetically and antigenically distinct types: DENV-1 to -4. While the majority of DENV infections resolve without the need for medical intervention, dengue can quickly progress in some patients to dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS) [1-3]. The risk factors associated with progressing to severe dengue are complex and incompletely understood. However, the risk of developing severe disease increases significantly in individuals experiencing secondary/heterologous infections [4, 5].

The leading mechanistic explanation for this phenomenon is antibody-dependent enhancement (ADE). This model posits that DENV-specific IgG antibodies elicited by a prior heterotypic DENV infection can opsonize DENV without neutralizing infectivity, facilitating uptake by FcγR bearing phagocytes [6-9]. This mechanism is supported by strong epidemiological data and *in vitro* infection studies and has been a foundational model of dengue immunopathogenesis for over 40 years [10-12]. However, an underappreciated facet of the DENV lifecycle is that B cells are the most frequently infected cell type observed in circulation during acute DENV infection [13-17]. This observation has been substantiated by multiple groups using a wide array of assays including flow cytometry, qRT-PCR, mosquito-inoculation, and scRNAseq [13-17]. While B cells are known to be phagocytic [18, 19] their antigen-specific receptors provide an additional, specialized means of internalizing extracellular antigens. Accordingly, we hypothesize that binding of a DENV-specific BCR to DENV could facilitate virion uptake in a manner analogous to ADE.

In this study, we demonstrate that the expression of a DENV-specific B cell receptor renders cells highly susceptible to DENV infection. We posit that this process of BCR-dependent enhancement (BDE) of DENV infection represents an additional mechanism of immune-mediated enhancement which may fill some critical gaps in our current understanding of dengue immunopathogenesis.

RESULTS

Expression of a DENV-specific BCR renders cells susceptible to DENV infection. The motivation for this study was the appreciation that the structure of an FcγR receptor engaged with a DENV/IgG complex resembles that of a DENV-specific B cell receptor bound to DENV (**Figure 1A**). Accordingly, we hypothesized that the expression of a DENV envelope-specific BCR would render a cell susceptible to DENV infection, representing an alternative method of DENV-specific ADE. To test this hypothesis, we generated a panel of transmembrane Ig (tmIg) constructs encoding three well-characterized DENV-specific monoclonal antibodies as well as a control HA-specific tmIg antibody (**Figure 1B, Figure S1A, Figure S1B**). Cells expressing DENV-specific tmIgG constructs retained their DENV-E protein binding activity, while no DENV-specific binding activity was observed for the HA-specific tmIg construct (**Figure 1B**).

To evaluate the contribution of DENV-specific tmIgG expression on cellular susceptibility to DENV infection, we exposed tmIgG expressing cells to a DENV-1 pseudotyped flavivirus reporter virus particle (RVP) encoding a dsRed florescent reporter [20, 21], alongside cells transfected with the canonical DENV-entry receptor DC-SIGN (**Figure 1C and 1D**). A reporter virus platform was utilized

139 in this analysis to allow for the accurate discrimination of true cellular infection from tmIgG-mediated
140 binding of virions to the exterior of the target cell, as expression of the dsRed fluorescent reporter
141 encoded by the RVP will only occur following intracellular replication and translation of the RVP
142 genome.
143
Expression of DENV-specific tmIgG receptors significantly increased susceptibility to infection by the
DENV-1 RVP, to levels similar to cells expressing DC-SIGN (**Figure 1C and 1D**). The ability of
DENV-specific tmIg constructs to facilitate infection was not limited to IgG isotype BCRs, as a similar
pattern was observed in cells expressing tmIgM and tmIgA versions of DENV-specific antibodies
(**Supplemental Figure 2**). Notably, the expression of a virus-specific tmIgG is not a universal
mechanism for enhancing the susceptibility cells to viral infection, as the expression of HCMV-specific
tmIgG constructs did not increase the susceptibility of cells to infection by HCMV.(**Supplemental**
Figure 3).
152
While the expression of all DENV envelope-specific tmIgG constructs increased the susceptibility of
cells to DENV infection, there was some heterogeneity observed between the constructs. To ascertain if
this was due to the specificity of the antibodies for their cognate antigens, we compared the rate of BCR-
dependent infection observed in our assay to the rate of ADE achieved with soluble IgG versions of the
same mAbs. Strikingly, we observed the same rank-order of ADE and BDE activity of the antibodies
when analyzed using a conventional K562 ADE assay (**Figure 1D, 1E, 1F**). Given that ADE and BDE
were in the same rank-order (**Figure 1G**), these results suggest that many of the features of IgG isotype
antibodies that are associated with ADE risk may be directly translatable to BDE risk.
161
DENV-specific B cells are susceptible to DENV infection. To extend our results showing that
transgene-expressed DENV-specific BCRs increase the susceptibility of a permissive cell line to DENV
infection, we next examined this in human B cells with endogenous BCR expression. To this end, we
leveraged a panel of human memory B cell lines derived from flavivirus-immune donors that were
immortalized via genetic reprogramming with BCL-6/BCL-x$_L$ encoding retroviruses and maintained
with the T-cell derived stimuli CD40L and IL-21 (**Supplemental Figure 4A**) [22]. BCL6/BCL-x$_L$-
immortalized human memory B cells maintain surface Ig expression, ability to bind their cognate
antigen, engage proximal BCR signaling, as well the ability to secrete soluble immunoglobulin [22].
Two cell lines were selected for this analysis: clone 7B9 - which expresses an IgG isotype BCR that
binds (but poorly neutralizes) DENV-1 to -4 and ZIKV – and clone 2F3 – which expresses a ZIKV-
specific IgG isotype BCR but exhibits no DENV-binding activity (**Figure 2A, 2B, 2C, Supplemental**
Figure 4B, 4C, 4D).
174
To determine the impact of BCR specificity on the susceptibility of these cells to DENV infection, both
the 7B9 (DENV^{react}) and 2F3 (ZIKV^{react}) cell lines were exposed to live DENV-2 for 24 hours, followed
by extensive washing and additional incubation for 24 hours to allow for viral replication. Cells were
then fixed/permeabilized and analyzed by flow cytometry to quantify the abundance of DENV-infected
cells within the culture (**Supplemental Figure 4E, 4F**). Consistent with the results obtained with
transgenic expression of DENV-specific BCRs, the DENV-specific 7B9 cell line exhibited a significant
increase in the frequency of DENV-infected cells relative to the ZIKV-specific 2F3 line (**Figure 2D**). To
better characterize the nature of DENV-infection with this B cell system, scRNAseq analysis was
performed on both control and DENV-2 exposed 7B9 (DENV^{react}) cells (**Supplemental Figure 5,**
Supplemental Tables 1-4). This analysis revealed a similar infection rate (defined as the presence of

DENV-2 RNA) within the DENV-2 exposed culture as was observed by flow cytometric analysis of the
same culture (**Figure 2E**). Furthermore, both positive- and negative-sense DENV-2 RNA was detected
within the cells containing the most significant quantity of DENV-2 RNA, indicating active replication
in DENV-specific B cells. (**Figure 2F**). Differential gene expression analysis between DENV-2 infected
cells and uninfected cells demonstrated a transcriptional signature consistent with acute lymphocyte
activation and cytokine production (**Supplemental Figure 5J, 5K**).
191
To assess whether primary human B cells are susceptible to DENV infection, we developed a BCR-
dependent DENV infection assay that bypasses the need for cells to express a DENV-specific BCR. In
this experiment, DENV is bound to an enhancing IgM (VDB33-IgM) and these DENV/Ig immune
complexes are crosslinked to the BCR of polyclonal B cells using an anti-IgM secondary antibody
(**Supplemental Figure 6A)** and infection is assessed by intracellular DENV antigen with the pan-
DENV mouse mAb 4G2. We first tested this in Raji cells, which are poorly susceptible to DENV
infection even at high MOI yet are highly permissive if transfected with a scavenging receptor such as
DC-SIGN [23]. We observed minimal DENV infection in non-transfected Raji cells in all culture
conditions except when all components of the DENV/Ig/BCR crosslinking cocktail were present
(**Supplemental Figure 6B, C**). To survey the infectibility of primary human B cells we exposed
purified B cells from a flavivirus-naïve donor to the same infection conditions described above and
assessed the frequency of B cell infection by flow cytometry. As was observed for the Raji cells,
significant infection of the primary B cells was only observed in conditions containing the full
DENV/Ig/BCR crosslinking cocktail (**Figure 2G, Figure 2H, Supplemental Figure 6D**).
206
Frequency of DENV-infectible B cells increases following primary DENV Infection. A key
prediction of the BDE model described above is that the frequency of DENV-infectible B cells in a
flavivirus naïve individual should be quite low, but that this frequency should increase following a
primary DENV infection due to the accumulation of DENV-specific memory B cells. To address this
hypothesis, we analyzed PBMC collected from a dengue human challenge study, wherein flavivirus
naïve volunteers were exposed to the attenuated DENV-3 strain CH53489 (**Figure 3A, Figure 3B**). We
exposed PBMC collected on days 0, 28, and 90 post DENV-3 infection from 8 volunteers to a DENV-1
RVP, followed by overnight incubation and flow cytometric analysis (**Figure 3A, Supplemental Figure
7**). We observed a significant increase in the number of DENV-infectable B cells on days 28 and 90 post
DENV-3 infection relative to pre-infection (**Figure 3C, 3D**). No such increase in DENV-infectability
was observed in either the T cell (**Figure 3E, 3F**) or monocyte compartment (**Figure 3G, 3H**). These
results suggest that the accumulation of DENV-specific immunologic memory can alter the
susceptibility of an individual's B cells to DENV infection.
220
221   <u>**DISCUSSION**</u>
222
In this study, we demonstrate that the expression of a DENV envelope-specific B cell receptor renders
cells highly susceptible to DENV infection, and that the frequency of DENV-infectible B cells within a
given individual can be modified based on DENV immune status. While an extensive body of literature
supports the concept that B cells are a significant circulating reservoir of infectious virus during acute
dengue [13-17] – and CD300a has been posited to be a B cell entry receptor for DENV [17] – we
believe these are the first results to suggest that B cell susceptibility to DENV infection can be
dynamically impacted by prior DENV infections.
230

Although numerous other viruses exhibit tropism for B cells, we posit that several fundamental features of DENV uniquely position it as amenable to BCR-mediated infection. First, DENV does not have – or require - a single cognate entry receptor. Rather, DENV can utilize a broad range of non-specific scavenging receptors or immunoreceptors to gain access to acidified endosomes of phagocytes, resulting in the viral/endosomal membrane fusion and release of genomic material. Indeed, this is exemplified by prior work which demonstrated that simply tethering DENV to the surface of permissive cells using bi-specific antibodies was sufficient to cause efficient infection, similar in magnitude to what can be achieved via ADE in cells expressing FcγRs [24]. These prior results suggest that the expression of a DENV-specific BCR may render a cell susceptible to DENV infection simply by tethering the virion to the surface of the cell, thereby facilitating viral entry during the process of normal membrane recycling without the need to engage full BCR signaling. However, it is also possible that that BCR-mediated entry of DENV could activate the B cell via multivalent ligation of the BCR complex, thereby increasing protein production and increasing the efficiency of infection in a fashion analogous to what has been described as "intrinsic ADE" for FcγR-mediated entry. Additional work will be required to disentangle these two non-mutually exclusive processes.

An intriguing implication of this model of BCR-mediated entry of DENV is that it may offer some insight into the heterogeneity of risk associated with post-primary DENV exposures. Increased risk for severe disease following a secondary, heterotypic infection has been associated with a narrow range of low level preexisting DENV antibody titers – but only a small fraction of these individuals actually develop severe disease [5, 25, 26]. Given the extraordinary stability of memory B cells [27], one potential immunopathological mechanism contributing to severe dengue disease may be the relative abundance of DENV-specific antibodies and DENV virion-specific memory B cells circulating after a primary DENV infection. This may result in a situation where the population of highly-suseptible DENV-specific memory B cells are protected from infection with DENV until neutralizing antibody titers wane below a protective threshold. Additional analyses are required to determine if – and how – this mechanism may contribute to risk or protection from DENV infection.

Materials and Methods

Cells and Cell lines. PBMC for flow cytometric analysis were obtained from a previously described phase 1, open-label dengue human challenge study [28]. This DENV-3 human challenge study and all associated analysis was approved by the State University of New York Upstate Medical University (SUNY-UMU) and the Department of Defense's Human Research Protection Office. 293T cell line were maintained DMEM media (Gibco, # 11965092) supplemented to 10% FBS, 1% penicillin/streptomycin, and 1% L-glutamine. K562, and U937 DC-SIGN cell line RPMI. Raji cell line was purchased through ATCC and maintained in complete IMDM media (Gibco, # 12440053).

Generation of Immortalized B cell lines. 7B9 cells were generated from a flavivirus-naïve volunteer who was vaccinated with DENV1Δ30 and challenged 270 days later with DENV2Δ30 Tonga (NCT02392325). B cells were isolated 90 days following DENV2Δ30 challenge. This study was approved by the Johns Hopkins University Institutional Review Board (IRB). 2F3 cells were obtained from a DENV-exposed volunteer (DT165) about 1 year following a secondary ZIKV infection (approved by University of Vermont IRB). Both 7B9 and 2F3 cells were generated by transduction of IgM–CD27+ memory B cells with a retrovirus encoding human B cell lymphoma (BCL)-6 and BCL-xL and enhanced green fluorescent protein (GFP) and culture on CD40L-expressing L cell fibroblasts and

277 rhIL-21 (25 ng/mL, Peprotech) as described [23]. Surface IgG expression was performed on live B cells
278 (near-IR fluorescent reactive dye, Invitrogen, cat no. L34975A), that were positive for GFP and CD19-
279 PerCP.Cy5.5 (Clone HIB-19, Biolegend, #302230) using APC anti-human IgG Fc (Biolegend, cat. no.
280 410712) on a Beckman CytoFlex flow cytometer.
281
Viruses and RVPs. DENV-1 (strain Wp74) dsRed Reporter virus (RVP) was prepared as previously
described [20, 21]. RVPs were titrated on U937 DC-SIGN cells, infectious units (IU) calculated based
on the dilution of virus needed to infect 10-15% of cells as previously described [29]. DENV-4 (strain
D4-1036) stock was prepared by propagating a low-passage inoculum in C6/36 cells. DENV-2 (strain
Thailand/NGS-C/1944) stock was prepared by propagating a low-passage inoculum in Vero cells. RVP
infection assays were performed using an MOI of 0.1, with cells incubated with the RVP for 6hrs,
followed by and an additional 18hrs of incubation prior to analysis by flow cytometry. Surface staining
was performed in PBS + 2% FBS. Aqua Live/Dead (ThermoFisher, L34957) or ZombieUV (Biolegend,
423107) PBMC infection experiments. Data collection was performed on a BD LSRII or Fortessa flow
cytometer and analyzed using FlowJo v10.2 (Becton Dickinson). ADE assays were performed using
K562 cells as previously described [30, 31]. In short, serial dilutions of monoclonal antibodies were
incubated with the DENV-1 RVP (in sufficient amounts to infect 10%–15% of U937-DC-SIGN cells) at
a 1:1 ratio for 1 h at 37 °C. This mixture was then added to a 96-well plate containing 5×10^4 K562 cells
per well in duplicate. Cells were cultured for 18–20 hr overnight in a 37 °C, 5% CO2, humidified
incubator, follow by analysis by flow cytometry.
297
ELISA to detect DENV-specific antibodies. The assays were performed as previously described
[PMID: 30857944]. Briefly, virus was captured by plate-adsorbed mouse cross-reactive anti-DENV
envelope (E) protein monoclonal antibody 4G2. IgG-containing cell culture supernatant was then added
and positive DENV binding was detected by alkaline phosphatase-conjugated goat anti-human IgG (Fc)
antibody (Millipore Sigma) and p-nitrophenyl phosphate substrate (Millipore Sigma). Reaction color
change, indicating DENV-binding, was measured by spectrophotometry as OD_{405}.
304
305 Serum neutralizing antibody titers against DENV1-4 and ZIKV were determined by plaque reduction
306 neutralization test (PRNT), using lowest serum dilution that yielded a 50% reduction in viral plaques
307 ($PRNT_{50}$) as previously described (pmids: 27089205 and 30996133).
308
Generation and expression of transmembrane antibody constructs. Sequences for each monoclonal
antibody were appended with an Ig transmembrane domain appropriate to the mAb isotype and cloned
into pcDNA3.1(+) expression constructs. Transient expression of all tmIg constructs was performed in
293T cells using Lipofectamine (Invitrogen, L3000001). Cells were incubated for 18hrs before use in
binding and infection assays. Antigen-specificity of the DENV-specific BCR was assessed by staining
with biotin-labeled DENV-2 E protein pre-incubated with FITC-tagged streptavidin.
315
316 **hCMV infection.** hCMV strain TB40E-GFP [32] was diluted in complete RPMI media then incubated
317 with transfected cells at an MOI of 1 at 37C for 6 hours. Cells were then washed before incubating at
318 37C for an additional 18 hours before quantification of infection by flow cytometry.
319
Single cell RNA sequencing: Samples were prepared for single-cell RNA sequencing according to the
10x Genomics RNA-seq protocol. Cells were resuspended at a concentration of 1400 cells/uL in PBS
and loaded for a target of 69000 cells per reaction. Cells were loaded for Gel emulsion bead (GEM)

generation and barcoding. Construction of 5' gene expression libraries was performed using the Next
GEM Single Cell 5' reagent kit, Library Construction Kit, and The i7 Multiplex Kit (10x Genomics,
CA). was used for reverse transcription, complementary DNA amplification and construction of gene
expression libraries. The quality of gene expression libraries was assessed using an Agilent 4200
TapeStation with High Sensitivity D500 ScreenTape Assay and Qubit Fluorometer (Thermo Fisher
Scientific) according to the manufacturer's recommendations. Sequencing of 5' gene expression
libraries was performed on an Illumina NextSeq 2000 (Illumina) using P3 reagent kits (100 cycles).
Parameters for sequencing were set at 26 cycles for Read1, 10 cycles for Index1, and 90 cycles for
Read2.
332
5' gene expression analysis and visualization: Gene expression alignment, sample demultiplexing,
alignment, and barcode/UMI filtering was performed using the Cell Ranger software package (10x
Genomics, CA) and bcl2fastq (Illumina, CA) using the commands mkfastq and count. The human
reference genome (Ensembl GRCh38.93) was combined with the DENV2 genome as an additional
chromosome (NC 001474.2). Sequenced transcripts were aligned to a human reference library created
using the Cell Ranger mkref command, combined human and DENV2 reference genome, and custom
Ensembl GRCh38 DENV2 GTF. Reads were mapped to both the positive and negative sense DENV2
genome. Multi-sample integration, data normalization, dimensional reduction, visualization and
differential gene expression were performed in R studio (v4.3.2) using R package Seurat (v4.4.0). The
datasetfiltered to exclude genes expressed in fewer than 3 cells, and to contain cells with less than 22%
mitochondrial RNA content and between 500-7,500 unique features. The resulting dataset was
normalized and scaled using the Seurat functions NormalizeData(), ScaleData(), and
FindVariableFeatures(). Data was normalized and scaled, and principal component analysis was
performed using RunPCA(). Clustering was based on the first 10 principal components and a resolution
parameter of 0.1 using FindNeighbors() and FindClusters(), respectively.
348
Cells labelled as infected were identified by the presence of both DENV (+) and (-) RNA with
expression of both transcripts >0. Differentially expressed genes were identified by applying the
FindMarkers() command and the Wilcoxon rank-sum test to the normalized gene expression dataset. A
default minimum logFC value of 0.1 and min.pct of 0.01 were used. Genes with a corrected p-value by
Bonferroni correction of <0.05 were considered significant. Ingenuity Pathway Analysis (IPA, Qiagen)
was performed using genes with a corrected p-value <0.01 and log fold change <-0.25 or >0.25.
355
DENV/Ig/BCR crosslinking infection. DENV/antibody immune complexes were generated by mixing
DENV-4 (6.25 x 10^7 IU/mL) using an MOI of 62 with 1ug/mL of VDB33-IgM and anti-IgM in 96-well
plate. 5 x 10^4 Raji cells or primary B cells isolated from flavivirus-naïve healthy donors using MojoSort
Human Pan B cell Isolation Kit (BioLegend, 480082) were then added and incubated for 24 hours.
Supernatants were harvest at 24hrs for cytokine analysis, while cells were washed and incubated an
additional 24 hours before analysis. Cells were intracellularly stained with FITC-labelled DENV E-
reactive monoclonal 4G2 antibody at 4µg/mL to quantify DENV infection.
363
364 **Statistical analysis.** Statistical analyses were performed using GraphPad Prism 9 (La Jolla, CA) with a
365 p-value < 0.05 considered significant.
366
367 **Data availability:** scRNAseq data are publicly available through the Gene Expression Omnibus,
368 accession number GSE276766. Code used to perform scRNAseq analysis in R is available upon request.

369
<u>Acknowledgment</u>

We gratefully acknowledge excellent technical assistance provided by Lisa Phelps of the SUNY Upstate
Medical University Flow Cytometry Core. We also wish to thank Ted Pierson and Alan Rothman for the
constructs and protocols for reporter virus production, and Saravanan Thangamani for the DENV-2
utilized in this study. The opinions or assertions contained herein are the private views of the authors
and are not to be construed as reflecting the official views of the US Army or the US Department of
Defense. Material has been reviewed by the Walter Reed Army Institute of Research. There is no
objection to its presentation and/or publication. The investigators have adhered to the policies for
protection of human participants as prescribed in AR 70-25. We acknowledge the Harry Hood Bassett
Flow Cytometry and Small Particles Detection facility (RRID:SCR_022147) at the University of
Vermont Larner College of Medicine.

Funding: Funding for this research was provided by the State of New York and the Department of
Defense, Medical Research and Material Command. Additional funding was provided by NIH
U01AI141997-06 and P20GM125498 (S.A.D.). Clinical trial NCT02392325 was funded by the Bill and
Melinda Gates Foundation (OPP1109415).

Author contributions. Conceptualization: C.G., J.R.C., H.F., G.D.G., A.T.W. Formal analysis: C.G.,
C.S.C.H., J.Q.L, A.T.W. Funding acquisition: S.J.T., A.T.W., S.A.D. Investigation: C.G., C.S.C.H.,
B.D.M., N.R.G., S.A.D., A.T.W., J.Q.L., S.M. Resources: S.J.T., G.C.C., A.T.W. Visualization: C.G.,
A.T.W. Writing – original draft: C.G. and A.T.W. Writing – review & editing: all authors

Competing interests: All authors: No reported conflicts of interest.

Data and material availability. The authors declare that all data supporting the findings of this study
are available within this article and its Supplementary Information files, or from the corresponding
author upon reasonable request.

References

1.   Organization WH. Dengue and severe dengue World Health Organization [updated March 17, 2023;
cited 2024 April 7]. Available from: https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-
dengue.
2.   Tantracheewathorn T, Tantracheewathorn S. Risk factors of dengue shock syndrome in children. J Med
Assoc Thai. 2007;90(2):272-7. PubMed PMID: 17375631.
3.   Thein S, Aung MM, Shwe TN, Aye M, Zaw A, Aye K, et al. Risk factors in dengue shock syndrome. Am J
Trop Med Hyg. 1997;56(5):566-72. doi: 10.4269/ajtmh.1997.56.566. PubMed PMID: 9180609.
4.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and
burden of dengue. Nature. 2013;496(7446):504-7. Epub 2013/04/09. doi: 10.1038/nature12060. PubMed PMID:
23563266; PubMed Central PMCID: PMCPMC3651993.
5.   Salje H, Cummings DAT, Rodriguez-Barraquer I, Katzelnick LC, Lessler J, Klungthong C, et al.
Reconstruction of antibody dynamics and infection histories to evaluate dengue risk. Nature.
2018;557(7707):719-23. Epub 20180523. doi: 10.1038/s41586-018-0157-4. PubMed PMID: 29795354; PubMed
Central PMCID: PMCPMC6064976.

6.   Halstead SB, O'Rourke EJ. Dengue viruses and mononuclear phagocytes. I. Infection enhancement by
non-neutralizing antibody. J Exp Med. 1977;146(1):201-17. doi: 10.1084/jem.146.1.201. PubMed PMID: 406347;
PubMed Central PMCID: PMCPMC2180729.
7.   Halstead SB, O'Rourke EJ, Allison AC. Dengue viruses and mononuclear phagocytes. II. Identity of blood
and tissue leukocytes supporting in vitro infection. J Exp Med. 1977;146(1):218-29. doi: 10.1084/jem.146.1.218.
PubMed PMID: 195000; PubMed Central PMCID: PMCPMC2180735.
8.   Bournazos S, Gupta A, Ravetch JV. The role of IgG Fc receptors in antibody-dependent enhancement.
Nat Rev Immunol. 2020;20(10):633-43. Epub 2020/08/13. doi: 10.1038/s41577-020-00410-0. PubMed PMID:
32782358; PubMed Central PMCID: PMCPMC7418887.
9.   Wang TT, Sewatanon J, Memoli MJ, Wrammert J, Bournazos S, Bhaumik SK, et al. IgG antibodies to
dengue enhanced for FcgammaRIIIA binding determine disease severity. Science. 2017;355(6323):395-8. doi:
10.1126/science.aai8128. PubMed PMID: 28126818; PubMed Central PMCID: PMCPMC5557095.
10.   Halstead SB. Neutralization and antibody-dependent enhancement of dengue viruses. Adv Virus Res.
2003;60:421-67. Epub 2003/12/24. doi: 10.1016/s0065-3527(03)60011-4. PubMed PMID: 14689700.
11.   Halstead SB, Rojanasuphot S, Sangkawibha N. Original antigenic sin in dengue. Am J Trop Med Hyg.
1983;32(1):154-6. Epub 1983/01/01. doi: 10.4269/ajtmh.1983.32.154. PubMed PMID: 6824120.
12.   Kliks SC, Nimmanitya S, Nisalak A, Burke DS. Evidence that maternal dengue antibodies are important in
the development of dengue hemorrhagic fever in infants. Am J Trop Med Hyg. 1988;38(2):411-9. Epub
1988/03/01. doi: 10.4269/ajtmh.1988.38.411. PubMed PMID: 3354774.
13.   Srikiatkhachorn A, Wichit S, Gibbons RV, Green S, Libraty DH, Endy TP, et al. Dengue viral RNA levels in
peripheral blood mononuclear cells are associated with disease severity and preexisting dengue immune status.
PLoS One. 2012;7(12):e51335. Epub 2013/01/04. doi: 10.1371/journal.pone.0051335. PubMed PMID: 23284680;
PubMed Central PMCID: PMCPMC3526575.
14.   Zanini F, Robinson ML, Croote D, Sahoo MK, Sanz AM, Ortiz-Lasso E, et al. Virus-inclusive single-cell RNA
sequencing reveals the molecular signature of progression to severe dengue. Proc Natl Acad Sci U S A.
2018;115(52):E12363-E9. Epub 2018/12/12. doi: 10.1073/pnas.1813819115. PubMed PMID: 30530648; PubMed
Central PMCID: PMCPMC6310786.
15.   Sanborn MA, Li T, Victor K, Siegfried H, Fung C, Rothman AL, et al. Analysis of cell-associated DENV RNA
by oligo(dT) primed 5' capture scRNAseq. Sci Rep. 2020;10(1):9047. Epub 2020/06/05. doi: 10.1038/s41598-020-
65939-5. PubMed PMID: 32493997; PubMed Central PMCID: PMCPMC7270085 the conduct of the study. A.L.R.
reports grants from National Institute of Allergy and Infectious Diseases, during the conduct of the study. All
other authors declare no competing interests.
16.   Ghita L, Yao Z, Xie Y, Duran V, Cagirici HB, Samir J, et al. Global and cell type-specific immunological
hallmarks of severe dengue progression identified via a systems immunology approach. Nat Immunol.
2023;24(12):2150-63. Epub 2023/10/24. doi: 10.1038/s41590-023-01654-3. PubMed PMID: 37872316.
17.   Upasani V, Vo HTM, Auerswald H, Laurent D, Heng S, Duong V, et al. Direct Infection of B Cells by
Dengue Virus Modulates B Cell Responses in a Cambodian Pediatric Cohort. Front Immunol. 2020;11:594813.
Epub 2021/03/02. doi: 10.3389/fimmu.2020.594813. PubMed PMID: 33643283; PubMed Central PMCID:
PMCPMC7907177.
18.   Zhu Q, Zhang M, Shi M, Liu Y, Zhao Q, Wang W, et al. Human B cells have an active phagocytic capability
and undergo immune activation upon phagocytosis of Mycobacterium tuberculosis. Immunobiology.
2016;221(4):558-67. Epub 2016/01/01. doi: 10.1016/j.imbio.2015.12.003. PubMed PMID: 26719096.
19.   Martinez-Riano A, Bovolenta ER, Mendoza P, Oeste CL, Martin-Bermejo MJ, Bovolenta P, et al. Antigen
phagocytosis by B cells is required for a potent humoral response. EMBO Rep. 2018;19(9). Epub 2018/07/11.
doi: 10.15252/embr.201846016. PubMed PMID: 29987136; PubMed Central PMCID: PMCPMC6123646.
20.   Pierson TC, Sanchez MD, Puffer BA, Ahmed AA, Geiss BJ, Valentine LE, et al. A rapid and quantitative
assay for measuring antibody-mediated neutralization of West Nile virus infection. Virology. 2006;346(1):53-65.
Epub 2005/12/06. doi: 10.1016/j.virol.2005.10.030. PubMed PMID: 16325883.

21. Ansarah-Sobrinho C, Nelson S, Jost CA, Whitehead SS, Pierson TC. Temperature-dependent production
of pseudoinfectious dengue reporter virus particles by complementation. Virology. 2008;381(1):67-74. Epub
2008/09/20. doi: 10.1016/j.virol.2008.08.021. PubMed PMID: 18801552; PubMed Central PMCID:
PMCPMC3428711.
22. Kwakkenbos MJ, Diehl SA, Yasuda E, Bakker AQ, van Geelen CM, Lukens MV, et al. Generation of stable
monoclonal antibody-producing B cell receptor-positive human memory B cells by genetic programming. Nat
Med. 2010;16(1):123-8. Epub 2009/12/22. doi: 10.1038/nm.2071. PubMed PMID: 20023635; PubMed Central
PMCID: PMCPMC2861345.
23. Martin NC, Pardo J, Simmons M, Tjaden JA, Widjaja S, Marovich MA, et al. An immunocytometric assay
based on dengue infection via DC-SIGN permits rapid measurement of anti-dengue neutralizing antibodies. J
Virol Methods. 2006;134(1-2):74-85. Epub 2006/01/19. doi: 10.1016/j.jviromet.2005.12.002. PubMed PMID:
16417930.
24. Mady BJ, Erbe DV, Kurane I, Fanger MW, Ennis FA. Antibody-dependent enhancement of dengue virus
infection mediated by bispecific antibodies against cell surface molecules other than Fc gamma receptors. J
Immunol. 1991;147(9):3139-44. Epub 1991/11/01. PubMed PMID: 1680925.
25. Salje H, Alera MT, Chua MN, Hunsawong T, Ellison D, Srikiatkhachorn A, et al. Evaluation of the extended
efficacy of the Dengvaxia vaccine against symptomatic and subclinical dengue infection. Nat Med.
2021;27(8):1395-400. Epub 2021/06/26. doi: 10.1038/s41591-021-01392-9. PubMed PMID: 34168334; PubMed
Central PMCID: PMCPMC8364868.
26. Katzelnick LC, Gresh L, Halloran ME, Mercado JC, Kuan G, Gordon A, et al. Antibody-dependent
enhancement of severe dengue disease in humans. Science. 2017;358(6365):929-32. Epub 2017/11/04. doi:
10.1126/science.aan6836. PubMed PMID: 29097492; PubMed Central PMCID: PMCPMC5858873.
27. Crotty S, Felgner P, Davies H, Glidewell J, Villarreal L, Ahmed R. Cutting edge: long-term B cell memory in
humans after smallpox vaccination. J Immunol. 2003;171(10):4969-73. Epub 2003/11/11. doi:
10.4049/jimmunol.171.10.4969. PubMed PMID: 14607890.
28. Waickman AT, Newell K, Lu JQ, Fang H, Waldran M, Gebo C, et al. Low-dose dengue virus 3 human
challenge model: a phase 1 open-label study. Nat Microbiol. 2024;9(5):1356-67. Epub 2024/04/02. doi:
10.1038/s41564-024-01668-z. PubMed PMID: 38561497.
29. Lambeth CR, White LJ, Johnston RE, de Silva AM. Flow cytometry-based assay for titrating dengue virus.
J Clin Microbiol. 2005;43(7):3267-72. Epub 2005/07/08. doi: 10.1128/JCM.43.7.3267-3272.2005. PubMed PMID:
16000446; PubMed Central PMCID: PMCPMC1169137.
30. Wegman AD, Fang H, Rothman AL, Thomas SJ, Endy TP, McCracken MK, et al. Monomeric IgA
Antagonizes IgG-Mediated Enhancement of DENV Infection. Front Immunol. 2021;12:777672. Epub 2021/12/14.
doi: 10.3389/fimmu.2021.777672. PubMed PMID: 34899736; PubMed Central PMCID: PMCPMC8654368.
31. Waickman AT, Gromowski GD, Rutvisuttinunt W, Li T, Siegfried H, Victor K, et al. Transcriptional and
clonal characterization of B cell plasmablast diversity following primary and secondary natural DENV infection.
EBioMedicine. 2020;54:102733. Epub 2020/04/22. doi: 10.1016/j.ebiom.2020.102733. PubMed PMID:
32315970; PubMed Central PMCID: PMCPMC7170960.
32. O'Connor CM, Murphy EA. A myeloid progenitor cell line capable of supporting human cytomegalovirus
latency and reactivation, resulting in infectious progeny. J Virol. 2012;86(18):9854-65. Epub 2012/07/05. doi:
10.1128/JVI.01278-12. PubMed PMID: 22761372; PubMed Central PMCID: PMCPMC3446554.

Figure Legends

Figure 1. Expression of DENV-specific BCRs renders cells susceptible to DENV-infection. **A)**
Schematic representation of DENV/IgG immune complex binding to an FcgR (left), and a DENV virion
binding to a DENV-specific BCR (right). **B)** Expression and DENV E protein binding activity of 293T
cells transfected with the indicated tmIgG constructs. **C)** DENV-1 RVP infection of 293T cells
transfected with the indicated tmIgG constructs. DC-SIGN and tmIgG conditions gated on receptor-
positive cells. **D)** Quantification of DENV-1 RVP infection of 293T cells transfected with DC-SIGN or
the indicated tmIgG constructs. Error bars +/- SEM. Experiments were conducted across three biological
replicates in triplicate. **E)** ADE activity of the indicated DENV-specific IgG mAbs at a 1ug/ml
concentration in K562 cells utilizing a DENV-1 RVP. **F)** Titration of ADE activity of the indicated
DENV-specific IgG mAbs in K562 cells utilizing a DENV-1 RVP. **G)** Pearson correlation analysis of
the ADE and BDE results described above. Error bars +/- SEM

Figure 2. DENV-specific B cells are susceptible to *ex vivo* DENV infection. A) Whole virus binding
profiles of IgG antibody (1 µg) from cross-reactive 7B9 and ZIKV-specific 2F3 mAb to DENV1-4 and
ZIKV by ELISA. **B)** Surface IgG surface expression of immortalized 7B9 and **C)** 2F3 B cells. **D)**
Quantification of DENV-2 infection of immortalized 7B9 (DENV^{react}) and 2F3 (ZIKV^{react}) B cells at
MOI of 10. Error bars +/- SEM. Experiments were conducted across two biological replicates in
triplicate **E)** Pie charts of DENV-2 infected 7B9 B cell proportions in control and DENV-2 inoculated
conditions as assessed by scRNAseq. **F)** Correlation plot of DENV (+) and DENV (-) sense RNA
expression in DENV-2 exposed 7B9 B cells as assessed by scRNAseq. **G)** Representative flow
cytometry plot showing the frequency of DENV-4 infected CD19+ B cells under the indicated culture
conditions. Detection of DENV-infected cells was performed by staining fixed/permeabilized cells with
a FITC-conjugated 4G2 antibody. **H)** Quantification of DENV-4 infected CD19+ B cells under the
indicated culture conditions. Error bars +/- SEM. **** $p < 0.001$, One-way ANOVA with Dunnett's
multiple comparisons test, with a single pooled variance. Experiments were conducted across three
biological replicates in triplicate.

Figure 3. Frequency of DENV-infectible B cells increases following primary DENV Infection. A)
Schematic representation of the dengue human infection model and *in vitro* DENV-1 RVP infection
assay. **B)** Key DHIM study characteristic and performance parameters. **C)** Representative flow
cytometry plots showing the frequency of DENV-1 RVP infectible viable B cells (CD3-CD19+) in
PBMC samples obtained 0-, 28-, and 90-days post DENV-3 infection. **D)** Quantification of DENV-1
RVP infectable B cells from all subjects included in this analysis. **E)** Representative flow cytometry
plots showing the frequency of DENV-1 RVP infectible viable T cells (CD3+CD19-) in PBMC samples
obtained 0-, 28-, and 90-days post DENV-3 infection. **F)** Quantification of DENV-1 RVP infectable T
cells from all subjects included in this analysis. **G)** Representative flow cytometry plots showing the
frequency of DENV-1 RVP infectible viable monocytes (CD3-CD19-CD14+) in PBMC samples
obtained 0-, 28-, and 90-days post DENV-3 infection. **H)** Quantification of DENV-1 RVP infectable
monocytes from all subjects included in this analysis. Error bars +/- SEM. ** $p < 0.01$, paired one-way
ANOVA with correction for multiple comparisons (Friedman test with Dunn's multiple comparisons
test).

Figure 1

Figure 2

Figure 3