B cell receptor dependent enhancement of dengue virus infection

Chad Gebo[1], Joseph Lu[1,2], Shima Moradpour[1], Jeffrey R Currier[3], Heather Friberg[3], Gregory D Gromowski[3], Stephen J. Thomas[1,2], Gary C. Chan[1], Adam T. Waickman[1,2,*]

[1]Department of Microbiology and Immunology, State University of New York Upstate Medical University, Syracuse, NY 13210, USA.

[2]Institute for Global Health and Translational Sciences, State University of New York Upstate Medical University, Syracuse, NY 13210, USA.

[3]Viral Diseases Branch, Walter Reed Army Institute of Research, Silver Spring, MD, 20910, USA

*Corresponding author: waickmaa@upstate.edu

Key words: dengue virus, immune mediated enhancement, B cells, B cell receptor

Abstract

Dengue virus (DENV) is the causative agent of dengue, a mosquito-borne disease that represents a significant and growing public health burden around the world. A unique pathophysiological feature of dengue is immune-mediated enhancement, wherein preexisting immunity elicited by a primary infection can enhance the severity of a subsequent infection by a heterologous DENV serotype. A leading mechanistic explanation for this phenomenon is antibody dependent enhancement (ADE), where sub-neutralizing concentrations of DENV-specific IgG antibodies facilitate entry of DENV into $Fc\gamma R$ expressing cells such as monocytes, macrophages, and dendritic cells. Accordingly, this model posits that phagocytic mononuclear cells are the primary reservoir of DENV. However, analysis of samples from individuals experiencing acute DENV infection reveals that B cells are the largest reservoir of infected circulating cells, representing a disconnect in our understanding of immune-mediated DENV tropism. In this study, we demonstrate that the expression of a DENV-specific B cell receptor (BCR) renders cells highly susceptible to DENV infection, with the infection-enhancing activity of the membrane-restricted BCR correlating with the ADE potential of the IgG version of the antibody. In addition, we observed that the frequency of DENV-infectable B cells increases in previously flavivirus-naïve volunteers after a primary DENV infection, and that the direct infection of B cells by DENV results in a pro-inflammatory cytokine response. These findings suggest that BCR-dependent infection of B cells is a novel mechanism of viral entry during DENV infection, and could contribute to immune-mediated enhancement of dengue pathogenesis.

INTRODUCTION

Dengue virus (DENV) is a prevalent arboviral pathogen that poses a significant global public health burden. Transmitted by the bite of infected *Aedes* family mosquitos, DENV co-circulates as four genetically and antigenically distinct types: DENV-1 to -4. While the majority of DENV infections resolve without the need for medical intervention, dengue can quickly progress in some patients to dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS) [1-3]. The risk factors associated with progressing to severe dengue are complex and incompletely understood. However, the risk of developing severe disease increases significantly in individuals experiencing secondary/heterologous infections [4, 5].

The leading mechanistic explanation for this phenomenon is antibody-dependent enhancement (ADE). This model posits that DENV-specific IgG antibodies elicited by a prior heterotypic DENV infection can opsonize DENV without neutralizing infectivity, facilitating uptake by FcγR bearing phagocytes [6-10]. This mechanism is supported by strong epidemiological data and *in vitro* infection studies and has been a foundational model of dengue immunopathogenesis for over 40 years [11-13]. However, an underappreciated facet of the DENV lifecycle is that B cells are the most frequently infected cell type observed in circulation during acute DENV infection [14-18]. This observation has been substantiated by multiple groups using a wide array of assays. While B cells are known to be phagocytic [19, 20] their antigen-specific receptors provide an additional, specialized means of internalizing extracellular antigens. Accordingly, we hypothesize that binding of a DENV-specific BCR to DENV could facilitate virion uptake in a manner analogous to ADE

In this study, we demonstrate that the expression of a DENV-specific B cell receptor renders cells highly susceptible to DENV infection. We posit that this process of BCR-dependent enhancement (BDE) of DENV infection represents an additional mechanism of immune-mediated enhancement which may fill some critical gaps in our current understanding of dengue immunopathogenesis.

RESULTS

Expression of a DENV-specific BCR renders cells susceptible to DENV infection. The motivation for this study was the appreciation that the structure of an FcγR receptor engaged with a DENV/IgG complex closely mirrored that of a DENV-specific B cell receptor bound to DENV (**Figure 1A**). Accordingly, we hypothesized that the expression of a DENV envelope-specific BCR would render a cell susceptible to DENV infection, representing an alternative method of DENV-specific ADE. To test this hypothesis, we generated a panel of transmembrane Ig (tmIg) constructs encoding three well-characterized DENV-specific monoclonal antibodies as well as a control HA-specific tmIg antibody (**Figure 1B, Figure S1A, Figure S1B**). We observed that the cells expressing DENV-specific tmIgG constructs retained their DENV-E protein binding activity, while no DENV-specific binding activity was observed for the HA-specific tmIg construct (**Figure 1B**).

To evaluate the contribution of DENV-specific tmIgG expression to susceptibility to DENV infection, we exposed tmIgG expressing cells to a DENV-1 RVP, alongside cells transfected with the canonical DENV-entry receptor DC-SIGN (**Figure 1C and 1D**). Expression of DENV-specific tmIgG receptors significantly increased susceptibility to infection by the DENV-1 RVP, to similar levels achieved in cells expressing DC-SIGN (**Figure 1C and 1D**). The ability of DENV-specific tmIg constructs to

139 facilitate infection was not limited to IgG isotype BCRs, as a similar pattern was observed in cells
140 expressing tmIgM and tmIgA versions of DENV-specific antibodies (**Supplemental Figure 2**). Notably,
141 the expression of a virus-specific tmIgG is not a universal mechanism for enhancing the susceptibility
142 cells to viral infection, as the expression of HCMV-specific tmIgG constructs did not increase the
143 susceptibility of cells to infection by this virus (**Supplemental Figure 3**).
144
While the expression of all DENV envelope-specific tmIgG constructs increased the susceptibility of
cells to DENV infection, there was some heterogeneity observed between the constructs. To ascertain if
this was due to the specificity of the antibodies for their cognate antigens, we compared the rate of BCR-
dependent infection observed in our assay to the rate of ADE achieved with soluble IgG versions of the
same mAbs. Strikingly, we observed the same rank-order of ADE and BDE activity of the antibodies
when analyzed using a conventional K562 ADE assay (**Figure 1D, 1E, 1F**). These results suggest that
many of the features of IgG isotype antibodies that are associated with ADE risk may be directly
translatable to BDE risk.
153
Frequency of DENV-infectible B cells increases following primary DENV Infection. A key
prediction of the BDE model described above is that the frequency of DENV-infectible B cells in a
flavivirus naïve individual should be quite low, but that this frequency should increase following a
primary DENV infection due to the accumulation of DENV-specific memory B cells. To address this
hypothesis, we analyzed PBMC collected from a dengue human challenge study, wherein flavivirus
naïve volunteers were exposed to the attenuated DENV-3 strain CH53489 (**Figure 2A, Figure 2B**). We
exposed PBMC collected on days 0, 28, and 90 post DENV-3 infection from 8 volunteers to a DENV-1
RVP, followed by overnight incubation and flow cytometric analysis (**Figure 2A, Supplemental Figure
4**). We observed a significant increase in the number of DENV-infectable B cells on days 28 and 90 post
DENV-3 infection relative to pre-infection (**Figure 2A, 2B**). No such increase in DENV-infectability
was observed in either the T cell (**Figure 2E, 2F**) or monocyte compartment (**Figure 2G, 2H**). These
results suggest that the accumulation of DENV-specific immunologic memory fundamentally and
specifically alters the susceptibility of an individual's B cells to DENV infection.
167
DENV infection of B cells increases pro-inflammatory cytokine production. The low frequency of
DENV-infectable B cells observed in even DENV-immune individuals significantly complicates the
evaluation of BDE-associated immunopathogenesis. Therefore, we sought to develop a BCR-dependent
DENV infection assay that bypasses the need for cells to express a DENV-specific BCR. To this end,
we utilized an DENV/Ig/BCR crosslinking strategy, wherein DENV/Ig immune complexes were
crosslinked to the BCR of polyclonal B cells using an anti-Ig secondary antibody (**Supplemental Figure
5**). The effectiveness and specificity of this infection strategy was first assessed using Raji cells, which
are poorly susceptible to DENV even at high MOI. We observed minimal DENV infection in the Raji
cells in all culture conditions except when all components of the DENV/Ig/BCR crosslinking cocktail
were present in the inoculum (**Figure 3A, 3B, Supplemental Figure 5**).
178
Having successfully established an infection assay that facilitates the direct infection of a B cell line in a
BCR-dependent fashion, we sought to determine if this assay could be utilized to infect primary B cells
from a flavivirus-naïve donor. To this end, we exposed purified primary human B cells to the same
infection conditions described above and assessed the frequency of B cell infection by flow cytometry.
As was observed for the Raji cells, significant infection of the primary B cells was only observed in
conditions containing the full DENV/Ig/BCR crosslinking cocktail (**Figure 3C, 3D, Supplemental**

Figure 5). Furthermore, B cell cultures containing the full DENV/Ig/BCR crosslinking cocktail
demonstrated significant IL-8 production, a cytokine associated with acute DENV infection [21], but
minimal TNFα production.
188
189 **DISCUSSION**
190
191 In this study, we demonstrate that the expression of a DENV envelope-specific B cell receptor renders
192 cells highly susceptible to DENV infection. Furthermore, we observed that the frequency of DENV-
193 infectible B cells within a given individual can be modified based on DENV immune status, and that
194 BCR-dependent infection of B cells results in the production of pro-inflammatory cytokines known to be
195 associated with DENV infection in human patients.
196
An intriguing implication of this model is that it may offer some insight into the heterogeneity of risk
associated with post-primary DENV exposure. The risk of developing severe dengue after a secondary
heterotypic infection is associated with having DENV-specific antibody levels from the first infection
within a narrow range of [5, 22, 23]. However, only a small fraction of individuals experiencing
secondary DENV infection with antibody titers within this window of risk actually experience severe
disease [5, 22, 23]. Given the extraordinary stability of memory B cells [24], one potential
immunopathological mechanism contributing to severe dengue may be the ratio of serum neutralizing
antibody titers to DENV envelope-specific memory B cells circulating after a primary DENV infection.
Additional analysis is required to determine if – and how – this mechanism may contribute to risk or
protection from DENV infection.
207
208 **Materials and Methods**
209
Cells and Cell lines. PBMC for flow cytometric analysis were obtained from a previously described
phase 1, open-label dengue human challenge study [21]. 293T cell line was purchased through ATCC
(CRL-3216) and maintained in DMEM media (Gibco, # 11965092) supplemented to 10% FBS, 1%
penicillin/streptomycin, and 1% L-glutamine. U937 DC-SIGN cell line was supplied by WRAIR. Raji
cell line was purchased through ATCC (CCL-86) and maintained in complete RPMI media. K562 cell
line was purchased through ATCC and maintained in complete IMDM media (Gibco, # 12440053).
216
Viruses and RVPs. DENV-1 (strain Wp74) dsRed Reporter virus (RVP) was prepared as previously
described [25, 26]. RVPs were titrated on U937 DC-SIGN cells, infectious units (IU) calculated based
on the dilution of virus needed to infect 10-15% of cells. DENV-4 (strain D4-1036) stock was prepared
by propagating a low-passage inoculum in C6/36 cells. RVP infection assays were performed using an
MOI of 0.1, with cells incubated with the RVP for 6hrs, followed by and an additional 18hrs of
incubation prior to analysis by flow cytometry. Surface staining was performed in PBS + 2% FBS. Aqua
Live/Dead (ThermoFisher, L34957) or ZombieUV (Biolegend, 423107) PBMC infection experiments.
Data collection was performed on a BD LSRII or Fortessa flow cytometer and analyzed using FlowJo
v10.2 (Becton Dickinson). ADE assays were performed using K562 cells as previously described [27,
28].
227
Generation and expression of transmembrane antibody constructs. Sequences for each monoclonal
antibody were appended with an Ig transmembrane domain appropriate to the mAb isotype and cloned
into pcDNA3.1(+) expression constructs. Transient expression of all tmIg constructs was performed in

293T cells using Lipofectamine (Invitrogen, L3000001). Cells were incubated for 18hrs before use in
binding and infection assays. Antigen-specificity of the DENV-specific BCR was assessed by staining
with biotin-labeled DENV-2 E protein pre-incubated with FITC-tagged streptavidin.
234
235 **hCMV infection.** hCMV strain TB40E-GFP [29] was diluted in complete RPMI media then incubated
236 with transfected cells at an MOI of 5 at 37C for 6 hours. Cells were then washed before incubating at
237 37C for an additional 18 hours before quantification of infection by flow cytometry.
238
DENV/Ig/BCR crosslinking infection. DENV/antibody immune complexes were generated by mixing
DENV-4 (6.25×10^7 IU/mL) using an MOI of 62 with 1ug/mL of VDB33-IgM and anti-IgM in 96-well
plate. 5×10^4 Raji cells or primary B cells were then added and incubated for 24 hours. Supernatants
were harvest at 24hrs for cytokine analysis, while cells were washed and incubated an additional 24
hours before analysis. Cytokine production was assessed using the MSD human U-PLEX Viral Combo
1 kit according to the manufacturer's recommendation. Cells were intracellularly stained with FITC-
labelled DENV E-reactive monoclonal 4G2 antibody at 4μg/mL to quantify DENV infection.
246
247 **Statistical analysis.** Statistical analyses were performed using GraphPad Prism 9 (La Jolla, CA) with a
248 p-value < 0.05 considered significant.
249
250 Acknowledgment

251
We gratefully acknowledge excellent technical assistance provided by Lisa Phelps of the SUNY Upstate
Medical University Flow Cytometry Core. We also wish to thank Ted Pierson and Alan Rothman for the
constructs and protocols for reporter virus production. The opinions or assertions contained herein are
the private views of the authors and are not to be construed as reflecting the official views of the US
Army or the US Department of Defense. Material has been reviewed by the Walter Reed Army Institute
of Research. There is no objection to its presentation and/or publication. The investigators have adhered
to the policies for protection of human participants as prescribed in AR 70-25.
259
260 **Funding:** Funding for this research was provided by the State of New York and the Department of
261 Defense, Medical Research and Material Command.

262
263 **Author contributions.** Conceptualization: C.G., J.R.C., H.F., G.D.G., A.T.W. Formal analysis: C.G.,
264 J.Q.L, A.T.W. Funding acquisition: S.J.T., A.T.W. Investigation: C.G., A.T.W., J.Q.L., S.M. Resources:
265 S.J.T., G.C.C., A.T.W. Visualization: C.G., A.T.W. Writing – original draft: C.G and A.T.W. Writing –
266 review & editing: all authors
267
268 **Competing interests:** All authors: No reported conflicts of interest.
269
270 **Data and material availability.** The authors declare that all data supporting the findings of this study
271 are available within this article and its Supplementary Information files, or from the corresponding
272 author upon reasonable request.
273
274 References

1.   Organization WH. Dengue and severe dengue World Health Organization [updated March 17, 2023;
cited 2024 April 7]. Available from: https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-
dengue.
2.   Tantracheewathorn T, Tantracheewathorn S. Risk factors of dengue shock syndrome in children. J Med
Assoc Thai. 2007;90(2):272-7. PubMed PMID: 17375631.
3.   Thein S, Aung MM, Shwe TN, Aye M, Zaw A, Aye K, et al. Risk factors in dengue shock syndrome. Am J
Trop Med Hyg. 1997;56(5):566-72. doi: 10.4269/ajtmh.1997.56.566. PubMed PMID: 9180609.
4.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and
burden of dengue. Nature. 2013;496(7446):504-7. Epub 2013/04/09. doi: 10.1038/nature12060. PubMed PMID:
23563266; PubMed Central PMCID: PMCPMC3651993.
5.   Salje H, Cummings DAT, Rodriguez-Barraquer I, Katzelnick LC, Lessler J, Klungthong C, et al.
Reconstruction of antibody dynamics and infection histories to evaluate dengue risk. Nature.
2018;557(7707):719-23. Epub 20180523. doi: 10.1038/s41586-018-0157-4. PubMed PMID: 29795354; PubMed
Central PMCID: PMCPMC6064976.
6.   <B cell inhibitory receptors and autoimmunity.pdf>.
7.   Halstead SB, O'Rourke EJ. Dengue viruses and mononuclear phagocytes. I. Infection enhancement by
non-neutralizing antibody. J Exp Med. 1977;146(1):201-17. doi: 10.1084/jem.146.1.201. PubMed PMID: 406347;
PubMed Central PMCID: PMCPMC2180729.
8.   Halstead SB, O'Rourke EJ, Allison AC. Dengue viruses and mononuclear phagocytes. II. Identity of blood
and tissue leukocytes supporting in vitro infection. J Exp Med. 1977;146(1):218-29. doi: 10.1084/jem.146.1.218.
PubMed PMID: 195000; PubMed Central PMCID: PMCPMC2180735.
9.   Bournazos S, Gupta A, Ravetch JV. The role of IgG Fc receptors in antibody-dependent enhancement.
Nat Rev Immunol. 2020;20(10):633-43. Epub 2020/08/13. doi: 10.1038/s41577-020-00410-0. PubMed PMID:
32782358; PubMed Central PMCID: PMCPMC7418887.
10.   Wang TT, Sewatanon J, Memoli MJ, Wrammert J, Bournazos S, Bhaumik SK, et al. IgG antibodies to
dengue enhanced for FcgammaRIIIA binding determine disease severity. Science. 2017;355(6323):395-8. doi:
10.1126/science.aai8128. PubMed PMID: 28126818; PubMed Central PMCID: PMCPMC5557095.
11.   Halstead SB. Neutralization and antibody-dependent enhancement of dengue viruses. Adv Virus Res.
2003;60:421-67. Epub 2003/12/24. doi: 10.1016/s0065-3527(03)60011-4. PubMed PMID: 14689700.
12.   Halstead SB, Rojanasuphot S, Sangkawibha N. Original antigenic sin in dengue. Am J Trop Med Hyg.
1983;32(1):154-6. Epub 1983/01/01. doi: 10.4269/ajtmh.1983.32.154. PubMed PMID: 6824120.
13.   Kliks SC, Nimmanitya S, Nisalak A, Burke DS. Evidence that maternal dengue antibodies are important in
the development of dengue hemorrhagic fever in infants. Am J Trop Med Hyg. 1988;38(2):411-9. Epub
1988/03/01. doi: 10.4269/ajtmh.1988.38.411. PubMed PMID: 3354774.
14.   Srikiatkhachorn A, Wichit S, Gibbons RV, Green S, Libraty DH, Endy TP, et al. Dengue viral RNA levels in
peripheral blood mononuclear cells are associated with disease severity and preexisting dengue immune status.
PLoS One. 2012;7(12):e51335. Epub 2013/01/04. doi: 10.1371/journal.pone.0051335. PubMed PMID: 23284680;
PubMed Central PMCID: PMCPMC3526575.
15.   Zanini F, Robinson ML, Croote D, Sahoo MK, Sanz AM, Ortiz-Lasso E, et al. Virus-inclusive single-cell RNA
sequencing reveals the molecular signature of progression to severe dengue. Proc Natl Acad Sci U S A.
2018;115(52):E12363-E9. Epub 2018/12/12. doi: 10.1073/pnas.1813819115. PubMed PMID: 30530648; PubMed
Central PMCID: PMCPMC6310786.
16.   Sanborn MA, Li T, Victor K, Siegfried H, Fung C, Rothman AL, et al. Analysis of cell-associated DENV RNA
by oligo(dT) primed 5' capture scRNAseq. Sci Rep. 2020;10(1):9047. Epub 2020/06/05. doi: 10.1038/s41598-020-
65939-5. PubMed PMID: 32493997; PubMed Central PMCID: PMCPMC7270085 the conduct of the study. A.L.R.
reports grants from National Institute of Allergy and Infectious Diseases, during the conduct of the study. All
other authors declare no competing interests.

17.   Ghita L, Yao Z, Xie Y, Duran V, Cagirici HB, Samir J, et al. Global and cell type-specific immunological
hallmarks of severe dengue progression identified via a systems immunology approach. Nat Immunol.
2023;24(12):2150-63. Epub 2023/10/24. doi: 10.1038/s41590-023-01654-3. PubMed PMID: 37872316.
18.   Upasani V, Vo HTM, Auerswald H, Laurent D, Heng S, Duong V, et al. Direct Infection of B Cells by
Dengue Virus Modulates B Cell Responses in a Cambodian Pediatric Cohort. Front Immunol. 2020;11:594813.
Epub 2021/03/02. doi: 10.3389/fimmu.2020.594813. PubMed PMID: 33643283; PubMed Central PMCID:
PMCPMC7907177.
19.   Zhu Q, Zhang M, Shi M, Liu Y, Zhao Q, Wang W, et al. Human B cells have an active phagocytic capability
and undergo immune activation upon phagocytosis of Mycobacterium tuberculosis. Immunobiology.
2016;221(4):558-67. Epub 2016/01/01. doi: 10.1016/j.imbio.2015.12.003. PubMed PMID: 26719096.
20.   Martinez-Riano A, Bovolenta ER, Mendoza P, Oeste CL, Martin-Bermejo MJ, Bovolenta P, et al. Antigen
phagocytosis by B cells is required for a potent humoral response. EMBO Rep. 2018;19(9). Epub 2018/07/11.
doi: 10.15252/embr.201846016. PubMed PMID: 29987136; PubMed Central PMCID: PMCPMC6123646.
21.   Waickman AT, Newell K, Lu JQ, Fang H, Waldran M, Gebo C, et al. Low-dose dengue virus 3 human
challenge model: a phase 1 open-label study. Nat Microbiol. 2024;9(5):1356-67. Epub 2024/04/02. doi:
10.1038/s41564-024-01668-z. PubMed PMID: 38561497.
22.   Salje H, Alera MT, Chua MN, Hunsawong T, Ellison D, Srikiatkhachorn A, et al. Evaluation of the extended
efficacy of the Dengvaxia vaccine against symptomatic and subclinical dengue infection. Nat Med.
2021;27(8):1395-400. Epub 2021/06/26. doi: 10.1038/s41591-021-01392-9. PubMed PMID: 34168334; PubMed
Central PMCID: PMCPMC8364868.
23.   Katzelnick LC, Gresh L, Halloran ME, Mercado JC, Kuan G, Gordon A, et al. Antibody-dependent
enhancement of severe dengue disease in humans. Science. 2017;358(6365):929-32. Epub 2017/11/04. doi:
10.1126/science.aan6836. PubMed PMID: 29097492; PubMed Central PMCID: PMCPMC5858873.
24.   Crotty S, Felgner P, Davies H, Glidewell J, Villarreal L, Ahmed R. Cutting edge: long-term B cell memory in
humans after smallpox vaccination. J Immunol. 2003;171(10):4969-73. Epub 2003/11/11. doi:
10.4049/jimmunol.171.10.4969. PubMed PMID: 14607890.
25.   Pierson TC, Sanchez MD, Puffer BA, Ahmed AA, Geiss BJ, Valentine LE, et al. A rapid and quantitative
assay for measuring antibody-mediated neutralization of West Nile virus infection. Virology. 2006;346(1):53-65.
Epub 2005/12/06. doi: 10.1016/j.virol.2005.10.030. PubMed PMID: 16325883.
26.   Ansarah-Sobrinho C, Nelson S, Jost CA, Whitehead SS, Pierson TC. Temperature-dependent production
of pseudoinfectious dengue reporter virus particles by complementation. Virology. 2008;381(1):67-74. Epub
2008/09/20. doi: 10.1016/j.virol.2008.08.021. PubMed PMID: 18801552; PubMed Central PMCID:
PMCPMC3428711.
27.   Wegman AD, Fang H, Rothman AL, Thomas SJ, Endy TP, McCracken MK, et al. Monomeric IgA
Antagonizes IgG-Mediated Enhancement of DENV Infection. Front Immunol. 2021;12:777672. Epub 2021/12/14.
doi: 10.3389/fimmu.2021.777672. PubMed PMID: 34899736; PubMed Central PMCID: PMCPMC8654368.
28.   Waickman AT, Gromowski GD, Rutvisuttinunt W, Li T, Siegfried H, Victor K, et al. Transcriptional and
clonal characterization of B cell plasmablast diversity following primary and secondary natural DENV infection.
EBioMedicine. 2020;54:102733. Epub 2020/04/22. doi: 10.1016/j.ebiom.2020.102733. PubMed PMID:
32315970; PubMed Central PMCID: PMCPMC7170960.
29.   O'Connor CM, Murphy EA. A myeloid progenitor cell line capable of supporting human cytomegalovirus
latency and reactivation, resulting in infectious progeny. J Virol. 2012;86(18):9854-65. Epub 2012/07/05. doi:
10.1128/JVI.01278-12. PubMed PMID: 22761372; PubMed Central PMCID: PMCPMC3446554.

Figure Legends

Figure 1. Expression of DENV-specific BCRs renders cells susceptible to DENV-infection. **A)** Schematic representation of DENV/IgG immune complex binding to an FcgR (left), and a DENV virion binding to a DENV-specific BCR (right). **B)** Expression and DENV E protein binding activity of 293T cells transfected with the indicated tmIgG constructs. **C)** DENV-1 RVP infection of 293T cells transfected with the indicated tmIgG constructs. DC-SIGN and tmIgG conditions gated on receptor-positive cells. **D)** Quantification of DENV-1 RVP infection of 293T cells transfected with DC-SIGN or the indicated tmIgG constructs. Error bars +/- SEM. **E)** ADE activity of the indicated DENV-specific IgG mAbs at a 1ug/ml concentration in K562 cells utilizing a DENV-1 RVP. **F)** Titration of ADE activity of the indicated DENV-specific IgG mAbs in K562 cells utilizing a DENV-1 RVP. **G)** Pearson correlation analysis of the ADE and BDE results described above. Error bars +/- SEM

Figure 2. Frequency of DENV-infectible B cells increases following primary DENV Infection. A) Schematic representation of the dengue human infection model and *in vitro* DENV-1 RVP infection assay. **B)** Key DHIM study characteristic and performance parameters. **C)** Representative flow cytometry plots showing the frequency of DENV-1 RVP infectible viable B cells (CD3-CD19+) in PBMC samples obtained 0-, 28-, and 90-days post DENV-3 infection. **D)** Quantification of DENV-1 RVP infectable B cells from all subjects included in this analysis. **E)** Representative flow cytometry plots showing the frequency of DENV-1 RVP infectible viable T cells (CD3+CD19-) in PBMC samples obtained 0-, 28-, and 90-days post DENV-3 infection. **F)** Quantification of DENV-1 RVP infectable T cells from all subjects included in this analysis. **G)** Representative flow cytometry plots showing the frequency of DENV-1 RVP infectible viable monocytes (CD3-CD19-CD14+) in PBMC samples obtained 0-, 28-, and 90-days post DENV-3 infection. **H)** Quantification of DENV-1 RVP infectable monocytes from all subjects included in this analysis. Error bars +/- SEM. ** $p < 0.01$, paired one-way ANOVA with correction for multiple comparisons (Friedman test with Dunn's multiple comparisons test).

Figure 3. DENV infection of B cells increases pro-inflammatory cytokine production. DENV/Ig/BCR crosslinking can facilitate DENV infection of polyclonal B cells. **A)** Representative flow cytometry plot showing the frequency of DENV-4 infected Raji cells under the indicated culture conditions. Detection of DENV-infected cells was performed by staining fixed/permeabilized cells with a FITC-conjugated 4G2 antibody. **B)** Quantification of DENV-4 infected Raji cells under the indicated culture conditions. **C)** Representative flow cytometry plot showing the frequency of DENV-4 infected CD19+ B cells under the indicated culture conditions. Detection of DENV-infected cells was performed by staining fixed/permeabilized cells with a FITC-conjugated 4G2 antibody. **D)** Quantification of DENV-4 infected CD19+ B cells under the indicated culture conditions. **E)** Quantification of IL-8 production from DENV-4 infected B cells under the indicated culture conditions. **F)** Quantification of TNFα production from DENV-4 infected B cells under the indicated culture conditions. **G)** Schematic representation of the contribution of BCR-dependent enhancement to DENV infection and immunopathogenesis. Error bars +/- SEM. **** $p < 0.001$, One-way ANOVA with Dunnett's multiple comparisons test, with a single pooled variance

Figure 1

Figure 2

A)

Dengue human infection model

Flavivirus naive volunteers   Start   Day 0   DENV-3 inoculation   Day 28   Day 90

PBMC Samples Collected

6 Hours   Wash   18 Hours

DENV-1 Reporter Virus

Assess RVP infection in B cells, T cells, monocytes

B)

DHIM-3 study parameters

Challenge virus	CH53489
Frequency of infection	100%
Frequency of seroconversion	100%
Window of viremia	D3 – D12

C) CD3- CD19+ B cells

Day 0   Day 28   Day 90

Mock   0   2.61E-3   0

+ RVP   0   0.056   0.036

dsRed (DENV-1 RVP)

D)

Frequency of DENV-1 infectible B cells

% dsRed+ (DENV-1 RVP)

**   **

Days post infection

E) CD3+ CD19- T cells

Day 0   Day 28   Day 90

Mock   2.30E-4   0   0

+ RVP   0   0   0

dsRed (DENV-1 RVP)

F)

Frequency of DENV-1 infectible T cells

% dsRed+ (DENV-1 RVP)

ns   ns

Days post infection

G) CD3- CD19- CD14+ monocytes

Day 0   Day 28   Day 90

Mock   0.010   0.016   2.29E-3

+ RVP   0.013   0.024   3.38E-3

dsRed (DENV-1 RVP)

H)

Frequency of DENV-1 infectible monocytes

% dsRed+ (DENV-1 RVP)

ns   ns

Days post infection

Figure 3