An applicable and efficient retrograde monosynaptic circuit mapping tool for larval zebrafish

Tian-Lun Chen[1,2,5], Qiu-Sui Deng[1,2,5], Kun-Zhang Lin[3], Xiu-Dan Zheng[1], Xin Wang[1,2], Yong-Wei Zhong[1], Xin-Yu Ning[1], Ying Li[1], Fu-Qiang Xu[3], Jiu-Lin Du[1,2,4*], Xu-Fei Du[1,2*]

[1]Institute of Neuroscience, State Key Laboratory of Neuroscience, Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, 320 Yue-Yang Road, Shanghai 200031, China

[2]University of Chinese Academy of Sciences, 19A Yu-Quan Road, Beijing 100049, China

[3]The Brain Cognition and Brain Disease Institute (BCBDI), Shenzhen Key Laboratory of Viral Vectors for Biomedicine, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

[4]School of Life Science and Technology, ShanghaiTech University, 319 Yue-Yang Road, Shanghai 200031, China

[5]These authors contributed equally

*Correspondence: xufeidu@ion.ac.cn, forestdu@ion.ac.cn

Abstract

The larval zebrafish is a vertebrate model for *in vivo* monitoring and manipulation of whole-brain neuronal activities. Tracing its neural circuits still remains challenging. Here we report an applicable methodology tailored for larval zebrafish to achieve efficient retrograde trans-monosynaptic tracing from genetically defined neurons via EnvA-pseudotyped glycoprotein-deleted rabies viruses. By combinatorially optimizing multiple factors involved, we identified the CVS strain trans-complemented with advanced expression of N2cG at 36°C as the optimal combination. It yielded a tracing efficiency of up to 20 inputs per starter cell. Its low cytotoxicity enabled the viable labeling and calcium imaging of infected neurons 10 days post-infection, spanning larval ages commonly used for functional examination. Cre-dependent labeling was further developed to enable input cell-type-specific tracing and circuit reconstruction. We mapped cerebellar circuits and uncovered the ipsilateral preference and subtype specificity of granule cell-to-Purkinje cell connections. Our method offers an efficient way for tracing neural circuits in larval zebrafish.

Introduction

Given the small transparent brain and accessible genetic manipulations, the larval zebrafish has been emerging as a promising animal model for systems neuroscience. Large-scale molecular, light- and electron-microscopical (LM and EM) morphological, and physiological datasets have been gathered over the past decade to decipher the identity and connectivity of its whole-brain cells[1–7]. Whole-brain neurons' activities can be monitored and manipulated in awake as well as behaving zebrafish larvae, providing an unprecedented research paradigm for dissecting neural mechanisms of brain functions[8–14]. Further mechanistic insights into the roles of different cell types in brain-wide activities requires effective tools for dissecting underlying neural circuits[15].

Virus-based tracing tools have opened new avenues for disentangling neural circuits[16,17]. Among them, the EnvA-pseudotyped glycoprotein (G)-deleted rabies virus (RV) (RV*d*G[EnvA]) has achieved most success[18–20]. Its success derives from being a native retrograde transsynaptic tracer engineered for cell-type-specific targeting through the recognition between EnvA and its receptor TVA[21] and crossing only one synaptic step via trans-complementation of rabies G[18,19]. This tool has revolutionized the structural analysis of neural circuits in rodents[15,22] and has been continually evolved to adapt to functional studies[23–25]. However, its application in other species including zebrafish is retarded due to lack of tailored methodologies.

The G-deleted RV (RV*d*G) infection of neurons in zebrafish was first demonstrated in 2009. Ten years later, RV*d*G[EnvA] was applied to trace input neurons in the adult zebrafish cerebellum, but the efficiency showed very low[26]. Recently, the herpes simplex virus (HSV1) was used as a helper virus to deliver TVA and rabies G, leading to an increased efficiency of around one input per starter neuron in the brain of adult zebrafish[27]. For larval zebrafish, efforts have been focused on the vesicular stomatitis virus (VSV), another rhabdovirus similar to RV but with anterograde spread[28–30]. However, these reported VSV tools for larval zebrafish appeared relatively high cytotoxicity and have not been iterated to utilize the EnvA-TVA system to achieve initial infection specificity[28–30]. Therefore, developing applicable efficient viral tracing methods is still an important mission in the field, and will boost the application of larval zebrafish in systems neuroscience research.

In the present study, we established applicable methodologies to implement RV*d*G[EnvA] for

efficient retrograde trans-monosynaptic tracing from genetically defined neurons in larval zebrafish. We first demonstrated that the transient co-expression of the helper proteins TVA and rabies G in specific neurons through one-cell-stage microinjection of the GAL-UAS binary DNA plasmids could successfully help the RV*d*G[EnvA] microinjected later on to infect and spread from targeted starter cells. This simple two-round microinjection procedure ensures the high feasibility of our method. Then via practicing in the cerebellar circuit, we iteratively tested key factors that influence the tracing efficiency and cytotoxicity, which include the virus strain, G protein type, G expression level, and working temperature. We discovered an optimal combinatory condition: CVSdG trans-complemented with advanced expression of N2cG at 36°C. The optimality lies in the high tracing efficiency and low cytotoxicity, embodied by a 20-fold increase in efficiency compared to previous studies in zebrafish[26,27] and a long enough time window (final fish age up to 2 - 3 weeks old) for conducting functional studies in larval zebrafish. Furthermore, by using Cre-expressing RV*d*G, we developed a transgenic reporter framework to enable input cell-type-specific tracing and circuit reconstruction based on single-neuron morphology. As an application demonstration, we applied the method to map the monosynaptic inputs to Purkinje cells (PCs) from granule cells (GCs) in the cerebellum and revealed the wiring relationship between morphological subtypes of GCs and PCs. These endeavors prove the applicability and efficiency of our method in dissecting interested circuits in larval zebrafish.

Results

Implementation of RV*d*G[EnvA]-based neural circuit tracing in larval zebrafish
To implement RV*d*G[EnvA]-based retrograde tracing in larval zebrafish, we first verified that larval zebrafish themselves do not express endogenous TVA. In wild type (WT) larvae without exogenous TVA expression, the injected SAD*d*G-mCherry[EnvA], a type of RV*d*G[EnvA], did not infect any cells (Figure 1—figure supplement 1A). The observed mCherry-positive puncta co-localized with microglia (Figure 1—figure supplement 1B), suggesting the clearance of uninfected virus particles by microglia. These data demonstrate that, similar to mammals[31], zebrafish do not express endogenous TVA.

Then we specifically expressed TVA in neurons (Figures 1A and B), and found that injected SAD*d*G-mCherry[EnvA] infected TVA-expressing cells around the injection site (Figure 1C and D, dashed yellow circles). The targeted expression of TVA and the fluorescent indicator enhanced green fluorescent protein (EGFP) in neurons was achieved through transient transgene of GAL4-UAS constructs in which the GAL4 activator is driven by *elavl3*, a pan-neuronal marker in zebrafish (Figure 1A, and "*UGT*" in Figure 1B). As the G protein is necessary for viral spread, SAD*d*G-mCherry[EnvA] did not spread beyond the initially infected neurons (i.e. starter cells) (Figure 1C,E-G).

We next examined whether viral particles can spread to other cells by complementing the G protein in starter neurons. As the efficiency of virus spread is reported to be positively correlated with the level of G expression in starter neurons[32,33], we generated two helper plasmids, *UAS:EGFP-P2A-TVA-T2A-B19G* (*UGTB*) and *UAS:EGFP-P2A-B19G-T2A-TVA* (*UGBT*), to search for a plasmid design with higher G expression while ensuring the co-expression of TVA and G (Figure 1B, see Methods). The position of the RV G-protein-coding gene SAD B19G was varied in the tri-cistronic 2A constructs for adjusting its expression level[34]. As shown in Figure

1D, besides starter cells (EGFP[+] and mCherry[+], dashed yellow circles) and uninfected TVA-

expressing cells (EGFP[+] only), we observed mCherry[+] only cells (arrows) of which the number

increased over time, indicating the virus spread beyond the starter cells by trans-complementation

with G. Notably, the traced cells included both neurons and radial glia (hereafter referred to as

glia) identified by their distinct morphology (Figure 1D, red and gray arrows, respectively).

Notably, the trans-infection of glia was also observed in mice[35,36], and its mechanism remains

unknown.

Convergence index (CI), defined as the number of trans-infected cells divided by that of

starter cells, was then used to estimate the tracing efficiency[19,32]. We calculated the CI of traced

neurons and glia separately, and found a higher tracing efficiency in larvae expressing *UGBT* than

those expressing *UGTB* (Figure 1E-G and Figure 1,2—table supplement 1; CI_{neuron}: 1.55 ± 0.29 *vs*

0.44 ± 0.11, $P < 0.01$; CI_{glia}: 4.41 ± 1.48 *vs* 2.50 ± 0.65, $P = 0.30$; mean ± SEM), suggesting that

UGBT is better for G expression and tracing efficiency. Taken together, these results demonstrate

that in larval zebrafish, transneuronal tracing from genetically defined cell types can be

implemented by combining the injection of RV*d*G[EnvA] with the co-expression of helper

proteins mediated by the GAL4-UAS system.

Optimization and verification of the retrograde monosynaptic tracing

To optimize this system for high tracing efficiency and application feasibility in circuit research,

we iterated it step-by-step in the cerebellum via testing different combinations of RV strains

(SAD-B19 *vs* CVS-N2c), G proteins (SAD B19G *vs* CVS N2cG), and rearing temperatures (28°C

vs 36°C) (Figure 2A). The cerebellar structure is evolutionarily conserved in zebrafish comparing

with mammals[37,38]. In zebrafish, Purkinje cells (PCs), the central cell type in the cerebellar circuit,

receive excitatory inputs from parallel fibers and climbing fibers originated respectively from

granule cells (GCs) and inferior olivary cells (IOCs), and local inhibitory inputs from stellate cells

(SCs)[37,38] (Figure 2B). The helper proteins G and TVA were co-expressed in PCs by using the

plasmid in *UGBT* mode, driven by *cpce*, a *ca8* promoter-derived PC-specific enhancer element[37,39]

(Figure 2A).

With the same G protein SAD B19G and rearing temperature at 28°C, CVS*d*G[EnvA]

exhibited around 6-fold increase in neuron tracing efficiency compared to SAD*d*G[EnvA] (Figure

2C,D, Figure 2—figure supplement 1A-C and Figure 1,2—table supplement 1; CI_{neuron}: 0.41 ±

0.07 *vs* 0.06 ± 0.03, $P < 0.001$). After replacing SAD B19G with CVS N2cG to complement

CVS*d*G[EnvA], the tracing efficiency was further increased by ~ 2 times (Figure 2C,D and Figure

1,2—table supplement 1; CI_{neuron}: 1.02 ± 0.18 *vs* 0.41 ± 0.07, $P < 0.01$). In addition,

CVS*d*G[EnvA] showed a greater tendency to spread to neurons rather than glia in comparison

with SAD*d*G[EnvA] (Figure 2C,E). In all experiments, virus injection was performed at 4 - 6 days

post-fertilization (dpf). The labeling of input neurons usually emerged around 2 days post

injection (dpi), and all CIs were calculated at 6 - 10 dpi, during which the viral transfer was

relatively stable (Figure 2—figure supplement 1C).

After elevating the temperature from 28°C to 36°C for rearing virus-injected larvae, the

neuron tracing efficiency of SAD*d*G[EnvA] became obvious (Figure 2C,D, Figure 2—figure

supplement 1D, and Figure 1,2—table supplement 1; CI_{neuron}: 4.96 ± 1.01 *vs* 0.06 ± 0.03, $P <$

0.001); for CVS*d*G[EnvA] trans-complemented with N2cG, the efficiency further increased by

one order of magnitude (Figure 2C,D, Figure 2—figure supplement 1E and Figure 1,2—table

supplement 1; CI_{neuron}: 11.61 ± 2.43 *vs* 1.02 ± 0.18, *P* < 0.001). In addition, the elevated

temperature also significantly increased the proportion of initially infected larvae (Figure 2F) and

the tendency of viral spread to neurons (Figure 2E). This efficient trans-synaptic viral transfer

from PCs allowed clear labeling of GC parallel fibers and long-range contralateral inputs from

IOCs (Figure 2G-I). We then generated a *Tetoff* element-based advanced helper plasmid (*UGNT-*

A) to enhance the expression level of CVS N2cG (A-N2cG) (Figure 2—figure supplement 1F, see

Methods), and this optimization resulted in a nearly two-fold increase in the neuron tracing

efficiency (Figure 2C,D, Figures 2—figure supplement 1F,G and Figure 1,2—table supplement 1;

CI_{neuron}: 20.35 ± 3.45 *vs* 11.61 ± 2.43, *P* < 0.05). The improvement was also observed for tracing

glia (Figure 2C, Figure 2—figure supplement 1H,I and Figure 1,2—table supplement 1).

To further functionally test the synaptic specificity of RV retrograde spread in the cerebellar

circuit, we conducted *in vivo* Ca^{2+} imaging of starter PCs while simultaneously activating single

traced GCs via electrical stimulation (Figure 2J, see Methods). To monitor the activity of starter

PCs, the fluorescent protein in the helper plasmid was replaced with the Ca^{2+} indicator GCaMP6s

(*UG6sNT-A*, Methods). GC stimulation-induced Ca^{2+} activities in starter PCs were abolished after

two-photon laser-based ablation of the GC stimulated (Figure 2K,L). These results indicate that

RV can retrogradely spread to GCs which are functionally connected with starter PCs, confirming

the capability of RV to trace monosynaptic inputs in larval zebrafish.

Time window for normal neuronal health after RV infection

In sum, we have identified the optimal conditions for efficient retrograde monosynaptic tracing in

zebrafish larvae: CVS*d*G[EnvA] trans-complemented with A-N2cG via the GAL4-UAS system at

36°C. The high efficiency of these optimal tracing conditions prompted concerns regarding the

potential cytotoxicity of CVS*d*G and CVS N2cG. In line with previous reports in mice[20], we found

lower toxicity of CVS*d*G compared with SAD*d*G. At 36°C, the survival rate of larvae infected

with CVS*d*G was significantly higher than those infected with SAD*d*G (Figure 3A; 75% ± 3% *vs*

52% ± 7%, *P* < 0.05). Furthermore, by using the time-lapse imaging as shown in Figure S2, we

often observed that early emerged starter cells disappeared in larvae infected with SAD*d*G (Figure

3B). As new starter cells continued to appear from 2 to 6 dpi (Figure 3C), we utilized starter cells

emerged at 2 dpi to quantify their lifetime. Consistently, starter cells infected by CVS*d*G survived

longer than those infected by SAD*d*G (Figure 3D; 10.0 ± 0 days *vs* 7.5 ± 0.48 days, *P* < 0.001).

Notably, the viability of starter cells decreased when CVS*d*G was trans-complemented with A-

N2cG (Figure 3D; 10.0 ± 0 days *vs* 8.7 ± 0.48 days, *P* < 0.001).

To further test whether physiological functions of starter cells were impaired by CVS*d*G

trans-complemented with A-N2cG, we conducted *in vivo* whole-cell recordings on starter PCs at

around 10 dpi (Figure 3E, see Methods). Infected PCs exhibited normal spontaneous excitatory

postsynaptic currents and potentials (sEPSCs and sEPSPs), and current injection-evoked firing

patterns (Figures 3F,G)[40]. Then we examined Ca^{2+} activities of infected (i.e., starter cells) and

uninfected PCs (i.e., non-starter cells) in respond to complex visual stimulation[41] (see Methods).

No significant differences were detected between starter and non-starter PCs (Figure 3H-L). Taken

together, these results indicate that under the optimal tracing conditions, infected neurons can

remain relatively healthy up to 10 days after infection in larval zebrafish, allowing for functional

probing of traced neural circuits.

Input cell-type-specific tracing and circuit reconstruction in the cerebellum

One important application of neural circuit tracing is to map the structural connectivity of specific
neuron types to build the brain connectome atlas. As shown above, even under the optimal tracing
conditions, glia still constituted around 47% on average of the traced cells (see Figure 2C,E). This
will certainly affect discrimination of traced neurons and their neural fibers. To confine retrograde
tracing to neurons instead of glia, we generated a Cre-Switch[42] transgenic reporter fish
Tg(elavl3:Tetoff-DO_DIO-Hsa.H2B-mTagBFP2_tdTomato-CAAX). This reporter expresses
pan-neuronally nuclear-localized blue fluorescent protein (mTagBFP2) by default, which serves as
a reference for image registration and data integration. It also achieves Cre-dependent neuron-
specific labeling with tdTomato-CAAX via the *elavl3* promoter (Figure 4A, see Methods). We
employed this reporter together with Cre-expressing CVS*d*G (CVS*d*G-Cre[EnvA]) to map
neuronal inputs to PCs in the cerebellum (Figure 4B). The transgenic labeling of traced neurons
showed much brighter and sharper fluorescence compared with virus-assisted labeling (Figure
4C). Among all 13 fish examined, we detected tdTomato-CAAX signals exclusively in neurons
without any labeling of glia (Figure 4C), facilitating the reconstruction of traced neural circuits
(Figure 4D). We noticed a relatively lower tracing efficiency of this Cre-dependent tracing system
(Figures 4C, 2C; CI_{neuron}: 5.20 ± 1.09 *vs* 11.61 ± 2.43). Ongoing efforts are performed to increase
the expression completeness of the transgenic reporter and improve the efficiency of Cre-mediated
recombination.

To demonstrate the feasibility of our tracing system in building and analyzing brain input
maps, we then generated a three-dimensional (3D) reference template brain from the transgenic
reporter fish at 17 dpf (see Methods). To incorporate cerebellar expression patterns, we generated

the transgenic lines *Tg(2×en.cpce:tdTomato-CAAX)* and *Tg(cbln12:GAL4FF)* to label PCs and

GCs, respectively. By utilizing H2B-mTagBFP2 expression in the reporter fish as a bridge, we
mapped the two cerebellar expression patterns and circuit tracing results of 13 fish mentioned
above onto the common coordinate space of the reference template (Figure 4B,E, see Methods).
We delineated the cerebellar brain area and reconstructed each starter PC (n = 24) and its input
GCs (n = 58) with clearly visible morphology (Figure 4E,F), resulting in a digital cellular-
resolution input atlas of cerebellar PCs (Figure 4F).

This input atlas clearly shows an ipsilateral preference of afferents from GCs to PCs (Figure
4G), which was in accordance with the summarized results of non-Cre-dependent viral tracing
data (Figures 4H and 2C,D). Interestingly, all reconstructed GCs also innervated contralaterally by
crossing the midline. We found two morphological subtypes for both GCs and PCs in the
reconstruction, and each included one local subtype with axonal projections within the cerebellar
area and one long projection subtype with axons extending caudally to the dorsal hindbrain[43,44]
(Figure 4I). Interestingly, although a single PC can receive inputs from the two subtypes of GCs
(Figure 4J), the two subtypes of PCs prefer inputs from different subtypes of GCs (Figure 4K,L).
These results demonstrate how cellular-level connectivity can advance our understanding of
circuit wiring patterns.

Together, the development of the Cre-dependent tracing tool empowers the RV*d*G-based
circuit tracing method, enabling input-specific tracing, cellular-level circuit reconstruction, and
integrated mapping and analysis of afferent connections in larval zebrafish.

Discussion

Understanding brain functions requires systematic dissection of the involved neural circuits. Here,
we developed a highly feasible and efficient implementation of the viral tracing tool RV*d*G[EnvA]
for circuit studies in larval zebrafish (Figure 1-4—table supplement 1). Through step-by-step
improvement, we identified an optimal tracing condition for larval zebrafish, which includes the
CVS*d*G strain, the native G protein of the CVS (i.e., N2cG), an enhanced expression system for
the G protein, and elevated rearing temperature. Our method exhibited a 20-fold increase in
tracing efficiency compared to previous studies in adult zebrafish[26,27], and demonstrated low
cytotoxicity. Moreover, we established a custom-designed transgenic reporter framework, which
enables genetic access to specific input cell types and allows for cellular-resolution reconstruction
and alignment of traced circuits. We demonstrated the application of this framework to decipher
cell-type specific connections from GCs to PCs in the cerebellum.

Helper protein expression design that guarantees high feasibility of the tracing method
Well-controlled spatial expression of helper proteins (i.e. TVA and G) in starter cells is critical for
the application of RV*d*G[EnvA]-based circuit tracing. Previous studies on zebrafish expressed
TVA and G through generating transgenic fish lines[26,27] or injecting helper viral vectors[27,29,30].
Both approaches allow for genetic access to a specific types or groups of neurons through the
GAL4-UAS system. The transgene typically results in expression in the majority of neurons of the
targeted type that usually distribute in many brain areas. Virus-assisted expression depends on the
distribution territory of injected virus. Restricting to specific sub-regions of the brain is quite
challenging due to the small compact brain of zebrafish larvae. Our method employs the transient
transgene technique in zebrafish to express helper proteins. This involves temporarily introducing
GAL4-UAS binary DNA plasmids into the fish's genome through microinjection into fertilized
eggs (see Figures 1A and 2A). It can achieve sparse expression of TVA and G in genetically
defined cell types, even in a single neuron. Tracing from a single neuron can help cell-subtype-
specific circuit dissection. Furthermore, we linked the two helper protein genes with the 2A
element to co-express TVA and G in the same neuron (see Figures 1A and 2A). Co-expression
guarantees that the tracing is monosynaptically restricted. This is because if G-expressing-only
neurons exist, they may act as input cells for initially infected starter cells, resulting in further
trans-complementation and spread of RV*d*G across the second and even more synaptic steps[32]
Previous studies on zebrafish expressed TVA and G separately using either two transgenic lines or
two viral vectors[26,27], potentially resulting in separate expression of them in different neurons.

Our study demonstrated that the transient expression of helper proteins effectively facilitates
subsequent viral infection and trans-complementation, offering a time-efficient and cost-effective
alternative to generating stable transgenic lines. It also addresses the potential cytotoxicity effects
associated with the use of helper viral vectors.

With commercially available recombinant RV tools, our method can be easily implemented
by zebrafish research laboratories using three commonly used techniques: plasmid construction,
microinjection (DNA constructs and recombinant RV with a 3 - 6 day interval) and live optical
imaging. Additional enhancements for the helper plasmid could include the implementation of
light-responsive inducible systems[45] to achieve both temporal and spatial control of helper protein
expression.

Key factors that influence tracing efficiency
To enhance RV*d*G tracing efficiency, we introduced the CVS-N2c rabies strain in zebrafish for the

first time. It has been reported that the CVS-N2c strain exhibits significantly higher tracing efficiency in mice compared to the SAD-B19 strain, with an improvement of almost tenfold[20]. In zebrafish larvae, CVS*d*G, when trans-complemented with its native N2cG, also showed an average increase close to tenfold compared to SAD*d*G trans-complemented with its native B19G (see Figure 2D and Figure 1,2—table supplement 1, CI_{neuron}). This improvement may be attributed to the reduced replication and protein expression of the CVS-N2c[20]. This could result in decreased toxicity to host cells, longer reproductive times, and ultimately higher spread efficiency. Consistently, in zebrafish larvae, we found that neurons infected with CVS-N2c displayed lower fluorescence intensity and longer survival times compared to those infected with SAD-B19. Notably, N2cG exhibited higher tracing efficiency than B19G when they were used to trans-complement the same CVS*d*G (see Figure 2D and Figure 1,2—table supplement 1). This could be attributed to the greater neurotropism of N2cG compared to B19G, or potential mismatches between the cytoplasmic domain of B19G and the CVS-N2c capsid. Further investigation into the impact of different G proteins on RV*d*G tracing efficiency in zebrafish would benefit from the construction of chimeric G proteins for comparison[46].

Besides the virus strain, temperature also plays a pivotal role. Under the optimal temperature for virus replication at 36°C, the tracing efficiency of CVS-N2c increased more than tenfold in comparison with that under the standard rearing temperature for zebrafish at 28°C. This effect was more pronounced for SAD-B19, showing an increase of over 80 times (see Figure 2D and Figure 1,2—table supplement 1). Importantly, in line with previous reports[27], we did not observe any noteworthy increase in mortality or behavioral abnormalities among zebrafish larvae exposed to elevated temperatures.

The expression level of G proteins is another determinant for efficient rabies tracing[46]. Building upon the codon optimization of the G protein, we further increased the expression of N2cG by inserting the Tetoff element upstream of the N2cG gene (A-N2cG) in the UAS helper plasmid (see Figure 2-figure supplement 1F, Figure 2D and Figure 1,2—table supplement 1). This integration enables dual amplification of N2cG and TVA expression by both the GAL4-UAS and tTA-TRE binary systems. Combining the three key optimal conditions: CVS-N2c strain, 36°C temperature, and A-N2cG, we have achieved the highest tracing efficiency in zebrafish to date, with a 20-fold increase compared to previous reports[27].

Monosynaptic specificity of the tracing

The RV*d*G-based transneuronal labeling of radial glial cells was commonly observed in larval zebrafish[29,30] (see Figure 1D,G and Figure 2—figure supplement 1H). The viral spread to glial cells may cause concerns about the synaptic specificity of RV*d*G spread between neurons. By employing the cerebellar circuit, we examined the complete pattern of traced neurons from PCs in larval zebrafish, and found that CVS*d*G was able to trace well-known presynaptic neurons of PCs, including both intra-cerebellar GCs and extra-cerebellar IOCs (see Figure 2G-I). Importantly, IOCs were the only labeled cells outside of the cerebellum, demonstrating the retrograde and monosynaptic specificity of this viral tracer in the zebrafish brain. We further provided functional evidence that CVS*d*G, even at its highest efficiency with A-N2cG, spreads retrogradely between neurons through synaptic connections (see Figure 2H-L). In all experiments involving the successful activation of PCs in response to electrical stimulation of single GCs, we consistently did not observe PC activation after ablating the GCs.

Low cytotoxicity that enables functional experiments in a wide time window

We found that the CVS-N2c strain did not significantly affect the neuronal health and physiology of starter cells for at least 10 days after infection (see Figure 3D-L). Meanwhile, consistent with previous reports in mice[20], the SAD-B19 strain exhibited higher cytotoxicity than CVS-N2c, resulting in more cell death of starter cells (see Figure 3D). It indicates that CVS-N2c is more suitable for application in zebrafish. We could observe the labeling of starter cells even one month later after CVS-N2c infection, and the infected zebrafish could be raised to adulthood. It is worth noting that enhanced G protein expression achieved through the introduction of the Tetoff element (i.e. A-N2cG) in the helper plasmid improved tracing efficiency, but also resulted in elevated levels of cytotoxicity, though lower than that induced by SAD-B19 (see Figure 3D). This indicates an association between the expression levels of G proteins and cytotoxicity[47,48]. Given the transferred RV is deficient in G, it is expected that retrogradely infected input cells will have lower cytotoxicity levels than starter cells. Therefore, input cells will survive for longer periods of time. Based on our experience, the CVS-N2c can be microinjected into zebrafish larvae between 3 and 6 dpf, and stable and viable labeling of traced circuits can be observed within 6 to 16 dpi. Therefore, the observation time window ranges from 9 to 22 dpf, spanning the larval ages commonly used for whole-brain functional imaging studies[14,49].

Functions and expandability of the transgenic framework

In the transgenic reporter framework (i.e. *elavl3DoDioBR*) (see Figure 4A), we designed three expansion motifs: 1) a promoter (*elavl3*) motif that allows genetic access to the expression of reporter cassettes in specific cell types, e.g. the *vglut2a* and *gad1b* gene promoters to restrict excitatory and inhibitory targeting, respectively; 2) a Cre-dependent "Switch-On" reporter (tdTomato-CAAX) motif that allows the expression of various structural and functional tools in traced neurons after viral spread, such as photoconvertible fluorescent proteins and optogenetic tools; 3) a Cre-dependent "Switch-Off" reporter (H2B-mTagBFP2) motif that allows a default reporter to be turned off specifically in infected neurons. In addition, the helper plasmid can serve as additional expansion motif that works together with the Cre-dependent transgenic reporter, allowing for full genetic visualization and dissection of neural circuits[15].

In this study, we used the *elavl3* promoter to restrict the fluorescence reporter to neurons and the "Switch-On" reporter tdTomato-CAAX to label the membrane of traced neurons. This enables clear neuron morphology labeling (see Figure 4C), facilitating the reconstruction of traced neural circuits based on single-neuron morphology. We used H2B-mTagBFP2 as the default "Switch-Off" reporter to provide a reference expression pattern for registering and integrating traced circuit images from different individual fish and experiments. As an application demonstration, we applied the transgenic framework to map the monosynaptic inputs to PCs from GCs in the cerebellum of larval zebrafish at ~ 2.5 weeks old (see Figure 4). Interestedly, we uncovered two properties of the cerebellar circuit: 1) the ipsilateral preference of GC-to-PC connections; 2) the subtype specificity of GC-to-PC connections. Further investigation is required to elucidate the functional implications of subtype-specific GC-to-PC connections and the apparent contradiction between the bilateral projection of GC parallel fibers and the GCs' preference for ipsilateral connections to PCs. The reference expression pattern can serve as a reference atlas to integrate other modal data from brain-wide images with information on gene expression, neurochemistry and neuronal activity. This will largely enhance our understanding of neural wiring patterns in

362 relation to other neural characteristics[50].

Conclusion

364 Our study established an applicable and effective viral circuit tracing system for larval zebrafish.

365 We believe that this tracing system will offer an avenue for the integrated structural and functional

366 dissection of neural circuits underlying a wide range of brain functions in larval zebrafish.

Methods

Animals

Adult zebrafish (*Danio rerio*) were reared under standard laboratory conditions, with temperatures set at 28°C, and a light/dark cycle of 14 hr/10 hr. The larval zebrafish used were in *nacre* background and at 3 - 21 dpf. The sex of the larvae at this stage is not determined. All experimental procedures were approved by the Animal Care and Use Committee of the Center for Excellence in Brain Science and Intelligence Technology (CEBSIT), Chinese Academy of Sciences (NA-046-2023).

Generation of DNA constructs and transgenic lines

The GAL4-UAS binary plasmids were applied to express fluorescent protein, TVA protein, and viral G in specific neuronal types. To randomly target neurons, we used *elavl3:GAL4-VP16*[51]. To target Purkinje cells (PCs), we generated a construct to drive the *GAL4FF* under the control of *cpce* (*ca8* promoter derived PC-specific enhancer element) fused to *E1B*[39]. The *cpce* sequence was PCR-amplified from the zebrafish genome DNA, and the *cpce-E1B:GAL4FF* was flanked by a minimal *Tol2* transposase recognition sites[52]. We generated seven helper plasmids (abbreviated names are in parentheses): *14×UAS-E1B:EGFP-P2A-TVA* (*UGT*), *14×UAS-E1B:EGFP-P2A-TVA-T2A-B19G* (*UGTB*), *14×UAS-E1B/5×UAS-hsp:EGFP-P2A-B19G-T2A-TVA* (*UGBT*, the *14×* and *5×UGBT* were used with *elavl3:GAL4-VP16* and *cpce-E1B:GAL4FF*, respectively), *5×UAS-hsp:sfGFP-P2A-N2cG-P2A-TVA* (*UGNT*), *5×UAS-hsp:sfGFP-P2A-Tetoff-N2cG-P2A-TVA* (*UGNT-A*), and *5×UAS-hsp:GCaMP6s-P2A-Tetoff-N2cG-P2A-TVA* (*UG6sNT-A*). The *sfGFP*, *N2cG*, *B19G*, *TVA*, and *Tetoff* were codon optimized for zebrafish and *de novo* synthesized; the *GCaMP6s* was PCR-amplified from *elavl3:GCaMP6s* plasmid (gift from M. B. Ahrens, Janelia Research Campus, USA). All these elements were sequentially assembled with 2A sequences and the linearized *miniTol2-UAS* backbone to generate polycistronic vectors through In-Fusion cloning. Sparse expression of helper proteins was achieved by the microinjection of a mixture (1 nl) of *elavl3* or *cpce*-driven GAL4 plasmid, UAS-driven helper plasmid, and *Tol2* transposase mRNA into one-cell stage zebrafish embryos at a concentration of 30 (10+10+10) ng/μl using an air-puffed pressure injector (MPPI-3, ASI).

To generate *Tg(2×en.cpce-E1B:tdTomato-CAAX)*, we created a construct to drive *tdTomato* with a 3' membrane tag encoding the *CAAX* box of human Harvey Ras (*CTGAACCCTCCTGATGAGAGTGGCCCCGGCTGCATGAGCTGCAAGTGTGTGCTCTCC*) under the control of two tandem *cpce* fused to E1B. To generate *Tg(cbln12:GAL4FF)* for labeling granule cells (GCs) in the cerebellum, we amplified a ~2-kbp regulatory element from zebrafish genome DNA by PCR for the *cbln12* (*cerebellin12*) gene and subcloned it into the linearized *miniTol2-GAL4FF* backbone. To generate *elavl3DoDioBR*, we first *de novo* synthesized a zebrafish codon-optimized *mTagBFP2* and constructed an intermediate plasmid containing *Tetoff-DO_DIO-Hsa.H2B-mTagBFP2_tdTomato-CAAX* using In-Fusion cloning. Subsequently, we subcloned this target fragment into the linearized mini*Tol2-elavl3* backbone to generate the final vector. Transgenic zebrafish were prepared using the *Tol2* transposase-based approach. Purified target plasmid DNA was microinjected with the *Tol2* transposase mRNA into one-cell stage zebrafish embryos at a concentration of 60 (30+30) ng/μl. Injected embryos were raised to sexual maturity to identify founder fish. The *Tg(cbln12:GAL4FF)* was crossed with *Tg(5×UAS:EGFP)* to

410   visualize GCs.

Production of EnvA-pseudotyped rabies virus

The EnvA-pseudotyped rabies viruses (RV), including SAD*d*G-mCherry[EnvA], CVS*d*G-
tdTomato[EnvA], and CVS*d*G-mCherry-2A-Cre[EnvA], were rescued and prepared following the
previously reported method[53,54]. Rabies vectors were stored at –80°C, and the titers were diluted to
2×10^8 infectious units per ml (IU/ml) with phosphate-buffered saline (PBS) before use. We found
that mCherry fluorescence after CVSdG-Cre[EnvA] infection *in vivo* was much weaker than that
of transgenetically expressed tdTomato-CAAX, and thus did not affect the observation and
reconstruction of neuronal morphology.

Virus application

At 3 - 6 dpf, larvae expressing helper proteins in specific neurons were anesthetized using 0.03%
tricaine methanesulfonate (MS-222) and mounted laterally in 1.0% low-melting agarose (Sigma)
on a glass-bottom dish filled with the extracellular solution consisted of (in mM): 134 NaCl, 2.9

KCl, 2.1 $CaCl_2 \cdot 2H_2O$, 1.2 $MgCl_2 \cdot 6H_2O$, 10 HEPES, and 10 glucose (pH = 7.8, osmolality =

290 mOsm). Under an inverted fluorescence microscope, 2 - 5 nl of virus suspension was injected
near the target cells (revealed by sfGFP or GCaMP6s fluorescence) in one cerebellar hemisphere
using a microinjector (Nanoliter2010, WPI) moved with a motorized controller (MC1000e,
Siskiyou). After injection, the fish were allowed to rest for one minute before withdrawing the
pipette. Then, they were placed in standard tanks in an incubator set at temperatures of 28°C or
36°C.

Confocal and two-photon imaging

Larval fish were mounted dorsal-side up in 1.5% low-melting agarose on a custom-made chamber.
Structural live imaging of virus-injected fish and transgenic lines was performed using an upright

confocal laser scanning microscope (FV3000, Olympus) equipped with a 20× water-immersion

objective (N.A., 0.9) at a 1-μm step size. For virus-injected fish, the imaging dimension was based
on the specific region of viral infection and spread. For transgenic lines, the entire brain was
imaged by multiple volumes, which were then stitched together using a custom ImageJ
(http://fiji.sc) macro based on the Grid/Collection stitching plugin[55]. Functional Ca^{2+} imaging of
PCs expressing GCaMP6s was performed using an upright two-photon microscope (FVMPE-RS,

Olympus) equipped with a 25× water-immersion objective (N.A., 1.05) at a wavelength of 920

nm. The calcium activity was recorded at ~5 Hz on single optical plane.

Electrical stimulation of single granule cell

Within 6 to 10 days after virus injection, larvae were paralyzed with pancuronium dibromide
(PCD, 1mM, Tocris 0693) for 1 min and then mounted in 1.5% low-melting agarose on a chamber
filled with the extracellular solution. The agarose and skin over the caudal side of the cerebellum
were removed for the access of stimulation electrodes. Stimulation microelectrodes (1 - 2 μm tip
opening) were created using borosilicate theta glass capillaries (BT-150-10, Sutter Instrument)
through a micropipette puller (P-97, Sutter Instrument). Then a silver wire was inserted into the
electrode filled with the extracellular solution. The electrode was positioned at and attached to the

cell body of the target GC, which was identified by the fluorescence and morphology using
confocal imaging. The stimulation was delivered by a flexible stimulus isolator (ISO-Flex, AMPI),
consisting of 5 pulse trains (6 - 8 V, 10 ms duration for one pulse, 10 pulses, 50 Hz) with inter-trial
intervals of 40 s. The electrical stimulation of single GC was synchronized with the Ca^{2+} imaging
of PCs by a stimulator (Master-8, AMPI).

Two-photon laser-based ablation of granule cells
A two-photon laser (795 nm) was used to ablate the GC after electrical stimulation. Confirmation
of a successful ablation involved the identification of a bulb-like structure in the bright-field image
and the absence of the tdTomato fluorescence signal from the targeted GC body. The efficiency of
the ablation was further validated one hour later, before performing post hoc electrical stimulation
at the same position using the bulb-like structure and adjacent fluorescent cells as references.

In vivo **whole-cell recording of Purkinje cells**
Within 6 to 15 days after virus injection, larvae were paralyzed with α-bungarotoxin (1mM) for 1
min, then mounted in 1.5% low-melting agarose on a chamber filled with the extracellular
solution. The agarose and skin over the caudal side of the cerebellum were removed for the access
of recording pipettes. A recording micropipette (15 - 20 MΩ, 1 - 1.5 µm tip opening) filled with
internal solution was approached to the target PC by a motorized micromanipulator (MP-225,
Sutter Instruments). The internal solution consisted of (in mM): 100 K-gluconate, 10 KCl, 2
$CaCl_2 \cdot 2H_2O$, 2 $Mg_2 \cdot ATP$, 0.3 $GTP \cdot Na_4$, 2 phosphocreatine, 10 EGTA, and 10 HEPES (pH = 7.4,
osmolality = 270 mOsm). The target PC was identified by double fluorescence (sfGFP[+] and
tdTomato[+]) using confocal imaging and was recorded at whole-cell mode with similar procedures
as our previous work[6]. The EPSP recording and current-step delivery were conducted in current-
clamp mode, and the EPSC was assessed in voltage-clamp mode. Data were acquired using a
patch-clamp amplifier (MultiClamp 700B, Axon Instruments) and a digitizer (Digidata 1440A,
Axon Instruments), and signals were sampled at 10 kHz with Clampex 10.6 software (Molecular
Devices).

Visual stimulation
Larvae examined were not anesthetized and were fully embedded in low-melting agarose. Visual
stimuli were generated by a custom program written in Python and presented using an LCD screen
covered with red filter paper to avoid spectrum interference. The visual stimuli basically followed
a prior study[41]. Briefly, three trials were presented and each trial consisted of three types of stimuli
in the following order: (1) four-directional whole-field gratings in black and white (including three
forward, one backward, one leftward, and one rightward directions); (2) windmill patterns in black
and white rotating at 0.2 Hz and varying in velocity following a sine function (including six
whole, two right-half, and two left-half filed stimuli presented in alternating clockwise and
counterclockwise directions); (3) whole-field flashes of 1-second duration (alternating between
white and black, repeated 4 times). Except for the flash, each stimulus lasted 5 seconds, with a 5-
second inter-stimulus interval.

Quantification of RV*d*G[EnvA]-based circuit tracing
Cells were counted by manually annotating the center of their cell bodies in image stacks using the
Cell Counter plugin in ImageJ. This was done for cells showing green (sfGFP/GCaMP6s; i.e.,
infected and uninfected TVA-expressing cells), red (tdTomato/mCherry; i.e., traced input cells and

starter cells), or yellow (both green and red; i.e., initially infected starter cells) fluorescence
separately. To count the traced neurons and glia separately, we also manually identified and
marked the neurons based on their significant morphological distinctions with glia in larval
zebrafish. The number of traced glia was then calculated by subtracting the number of starter cells
and traced neurons from the total number of cells with red fluorescence. To determine virus
tracing efficiency, a convergence index (CI) was calculated as the number of traced inputs
(neurons, glia, or all) divided by the number of starter cells.

Analysis of calcium imaging data

The acquired time-lapse images were aligned first, and individual regions of interest (ROIs) were
manually circled on the averaged image. Each ROI corresponds to either a starter cell or a non-
starter cell, and the mean grayscale value within each ROI over time was represented as F. In the
electrical stimulation experiment, the baseline value F0 was calculated as the average value of F
from a 10-second interval before each stimulus. In the visual stimulation experiment, F0 was
calculated as the average value of F over a 30-second period preceding the stimuli. The relative
change in the intensity of each ROI was calculated using the formula $(F - F0)/F0$. The area under
the curve (AUC) and the number of Ca^{2+} activity events were determined by applying a threshold
of 2× standard deviation (SD) of a 60-second baseline activity. All of the mentioned calculations
were performed using custom Python scripts.

Template generation and registration

We first generated two two-channel templates (17 dpf), *Tg(2×en.cpce:tdTomato-*
CAAX);elavl3DoDioBR and *Tg(cbln12:GAL4FF);Tg(5×UAS:EGFP);elavl3DoDioBR*, by running
a shell script antsMultivariateTemplateConstruction2.sh (-d 3 -b 1 -g 0.1 -i 10 -c 2 -j 6 -k 2 -w 1x1
-f 12x8x4x2 -s 4x3x2x1vox -q 400x200x200x20 -r 1 -n 0 -m CC -t SyN ${image-inputs}) on the
CEBSIT's Linux computing cluster using the open-source Advanced Normalization Tools (ANTs)
software[56]. Confocal images were acquired at $1 \times 1 \times 1$ μm resolution. Both templates were shape-
based average representations from 6 transgenic larvae after 7 iterations. Using normalized cross
correlation (NCC) as a metric to evaluate the registration accuracy between *elavl3DoDioBR*
templates and individual *elavl3DoDioBR* scans of larvae used for circuit tracing, we determined
the *elavl3DoDioBR* in *Tg(2×en.cpce:tdTomato-CAAX);elavl3DoDioBR* template space as the
common coordinate reference for registration. The expression pattern of
Tg(cbln12:GAL4FF);Tg(5×UAS:EGFP) and circuit tracing results were transformed into the
common coordinate space by using *elavl3DoDioBR* as a bridge. For image registration, we used
antsRegistration and antsApplyTransforms functions in ANTs, following the registration
parameters for live fish scans reported in previous work[57]. Neuron reconstructions (SWC files[58])
were aligned using antsApplyTransformsToPoints function in ANTs.

Cerebellum delineation and neuron reconstruction

527 The 3D cerebellum region was delineated manually using 3D Slicer software
528 (https://www.slicer.org/) based on the expression patterns of *elavl3DoDioBR*,
529 *Tg(2×en.cpce:tdTomato-CAAX)*, and *Tg(cbln12:GAL4FF);Tg(5×UAS:EGFP)* in the common
530 coordinate space. The boundary delineation was performed on *elavl3DoDioBR*, assisted by
531 overlaying the other two patterns.

To reconstruct the morphology of neurons in virus-traced circuits, neurons expressing membrane-bound tdTomato were semi-automatically traced using the simple neurite tracer (SNT)[59] plugin in ImageJ. The classification of dendritic and axonal compartments of neurites was based on structural knowledge. Neurons with low fluorescence intensity and ambiguous morphology were excluded during reconstruction. The resulting individual neuron skeletons were saved in SWC format. Tracings containing undistinguished several neuron skeletons were also saved as individual SWC files.

The co-volume rendering of template cerebellum and cerebellum region (see Figure 4E) was performed using ParaView[60] (https://www.paraview.org). The co-visualization of the cerebellar region and registered neuron reconstructions (see Figure 4F,I,J) was performed using custom python scripts based on NAVis package[61] (https://github.com/navis-org/navis).

Statistics

All statistical tests and graphs were performed with GraphPad Prism or Python software. Data normality was first examined with Shapiro-Wilk test. Comparisons between two groups with normal and non-normal distributions were made using two-tailed unpaired Student's *t* test and nonparametric two-tailed Mann-Whitney test, respectively. Differences between groups involving multiple factors were analyzed using two-way ANOVA. *P* value < 0.05 achieved statistical significance. Data are represented as mean ± SEM. The sample size and/or the number of replicates for each experiment are reported in the text, figures or figure legends.

Acknowledgments

We are grateful to Drs. M. B. Ahrens for sharing DNA plasmids, F. Peri and K. Kawakami for sharing zebrafish lines. We thank X.L. Shen, L. Sun, S. Li, and K. Wang for their kind help on virus injection, calcium imaging, visual stimulation, and plasmid construction experiments, respectively. This work was supported by STI2030-Major Projects (2021ZD0204500 and 2021ZD0204502), National Key R&D Program of China (2018YFA0801000 and 2018YFA0801001), National Natural Science Foundation of China (32321003), and Shanghai Municipal Science and Technology Major Project (18JC1410100 and 2018SHZDZX05). X.F.D. is also supported by the Youth Innovation Promotion Association of CAS.

Author contributions

X.F.D. and J.L.D. conceptualized and supervised the project. T.L.C. and Q.S.D. developed the methodology and designed the experiments with input from X.F.D. and J.L.D.. T.L.C. and Q.S.D. performed the experiments with input from X.W. and X.D.Z.. X.D.Z. created the cerebellum-related transgenic fish. K.Z.L. produced the rabies viruses under the supervision of F.Q.X.. T.L.C. analyzed data and made the illustrations with input from Q.S.D., Y.W.Z., and X.Y.N.. Y.L. helped to establish the virus injection in zebrafish. T.L.C., Q.S.D., X.F.D., and J.L.D. wrote the manuscript.

Data availability

Plasmids and transgenic zebrafish lines generated in this study, along with the microscopy and neuron reconstruction data reported in this paper, will be made available upon request. All original code has been deposited at https://github.com/soaringdu/Proj-

RVCT and is publicly available as of the date of publication. Any additional information
required to reanalyze the data reported in this paper is available upon request.

References

1.   Raj, B., Farrell, J.A., Liu, J., El Kholtei, J., Carte, A.N., Navajas Acedo, J., Du, L.Y.,
McKenna, A., Relić, Đ., Leslie, J.M., et al. (2020). Emergence of neuronal diversity during
vertebrate brain development. Neuron *108*, 1058-1074.e6.
2.   Zhang, H., Wang, H., Shen, X., Jia, X., Yu, S., Qiu, X., Wang, Y., Du, J., Yan, J., and He, J.
(2021). The landscape of regulatory genes in brain-wide neuronal phenotypes of a vertebrate
brain. eLife *10*, e68224.
3.   Shainer, I., Kuehn, E., Laurell, E., Kassar, M.A., Mokayes, N., Sherman, S., Larsch, J., Kunst,
M., and Baier, H. (2023). A single-cell resolution gene expression atlas of the larval zebrafish
brain. Sci. Adv. *9(8)*, eade9909.
4.   Kunst, M., Laurell, E., Mokayes, N., Kramer, A., Kubo, F., Fernandes, A.M., Förster, D., Dal
Maschio, M., and Baier, H. (2019). A cellular-resolution atlas of the larval zebrafish brain.
Neuron *103*, 21-38.e5.
5.   Svara, F., Förster, D., Kubo, F., Januszewski, M., Dal Maschio, M., Schubert, P.J., Kornfeld,
J., Wanner, A.A., Laurell, E., Denk, W., et al. (2022). Automated synapse-level reconstruction
of neural circuits in the larval zebrafish brain. Nat. Methods *19*, 1357–1366.
6.   Mu, Y., Li, X., Zhang, B., and Du, J. (2012). Visual input modulates audiomotor function via
hypothalamic dopaminergic neurons through a cooperative mechanism. Neuron *75*, 688–699.
7.   Yao, Y., Li, X., Zhang, B., Yin, C., Liu, Y., Chen, W., Zeng, S., and Du, J. (2016). Visual cue-
discriminative dopaminergic control of visuomotor transformation and behavior selection.
Neuron *89*, 598–612.
8.   Vanwalleghem, G.C., Ahrens, M.B., and Scott, E.K. (2018). Integrative whole-brain
neuroscience in larval zebrafish. Curr. Opin. Neurobiol. *50*, 136–145.
9.   Loring, M.D., Thomson, E.E., and Naumann, E.A. (2020). Whole-brain interactions
underlying zebrafish behavior. Curr. Opin. Neurobiol. *65*, 88–99.
10.   Naumann, E.A., Fitzgerald, J.E., Dunn, T.W., Rihel, J., Sompolinsky, H., and Engert, F.
(2016). From whole-brain data to functional circuit models: the zebrafish optomotor response.
Cell *167*, 947-960.e20.
11.   Mu, Y., Bennett, D.V., Rubinov, M., Narayan, S., Yang, C.-T., Tanimoto, M., Mensh, B.D.,
Looger, L.L., and Ahrens, M.B. (2019). Glia Accumulate Evidence that Actions Are Futile
and Suppress Unsuccessful Behavior. Cell *178*, 27-43.e19.
12.   Kappel, J.M., Förster, D., Slangewal, K., Shainer, I., Svara, F., Donovan, J.C., Sherman, S.,
Januszewski, M., Baier, H., and Larsch, J. (2022). Visual recognition of social signals by a
tectothalamic neural circuit. Nature *608*, 146–152.
13.   Petrucco, L., Lavian, H., Wu, Y.K., Svara, F., Štih, V., and Portugues, R. (2023). Neural
dynamics and architecture of the heading direction circuit in zebrafish. Nat. Neurosci. *26*,
765–773.
14.   Shang, C.-F., Wang, Y.-F., Zhao, M.-T., Fan, Q.-X., Zhao, S., Qian, Y., Xu, S.-J., Mu, Y., Hao,
J., and Du, J.-L. (2024). Real-time analysis of large-scale neuronal imaging enables closed-
loop investigation of neural dynamics. Nat. Neurosci. *27*, 1014–1018.
15.   Luo, L., Callaway, E.M., and Svoboda, K. (2018). Genetic dissection of neural circuits: a

decade of progress. Neuron *98*, 256–281.

16. Xu, X., Holmes, T.C., Luo, M.-H., Beier, K.T., Horwitz, G.D., Zhao, F., Zeng, W., Hui, M.,
Semler, B.L., and Sandri-Goldin, R.M. (2020). Viral vectors for neural circuit mapping and
recent advances in trans-synaptic anterograde tracers. Neuron *107*, 1029–1047.

17. Nectow, A.R., and Nestler, E.J. (2020). Viral tools for neuroscience. Nat. Rev. Neurosci. *21*,
669–681.

18. Wickersham, I.R., Lyon, D.C., Barnard, R.J.O., Mori, T., Finke, S., Conzelmann, K.-K.,
Young, J.A.T., and Callaway, E.M. (2007). Monosynaptic restriction of transsynaptic tracing
from single, genetically targeted neurons. Neuron *53*, 639–647.

19. Miyamichi, K., Amat, F., Moussavi, F., Wang, C., Wickersham, I., Wall, N.R., Taniguchi, H.,
Tasic, B., Huang, Z.J., He, Z., et al. (2011). Cortical representations of olfactory input by
trans-synaptic tracing. Nature *472*, 191–196.

20. Reardon, T.R., Murray, A.J., Turi, G.F., Wirblich, C., Croce, K.R., Schnell, M.J., Jessell, T.M.,
and Losonczy, A. (2016). Rabies virus CVS-n2c ΔG strain enhances retrograde synaptic
transfer and neuronal viability. Neuron *89*, 711–724.

21. Barnard, R.J.O., Elleder, D., and Young, J.A.T. (2006). Avian sarcoma and leukosis virus-
receptor interactions: from classical genetics to novel insights into virus–cell membrane
fusion. Virology *344*, 25–29.

22. Yao, S., Wang, Q., Hirokawa, K.E., Ouellette, B., Ahmed, R., Bomben, J., Brouner, K., Casal,
L., Caldejon, S., Cho, A., et al. (2023). A whole-brain monosynaptic input connectome to
neuron classes in mouse visual cortex. Nat. Neurosci. *26*, 350–364.

23. Chatterjee, S., Sullivan, H.A., MacLennan, B.J., Xu, R., Hou, Y., Lavin, T.K., Lea, N.E.,
Michalski, J.E., Babcock, K.R., Dietrich, S., et al. (2018). Nontoxic, double-deletion-mutant
rabies viral vectors for retrograde targeting of projection neurons. Nat. Neurosci. *21*, 638–
646.

24. Jin, L., Sullivan, H.A., Zhu, M., Lea, N.E., Lavin, T.K., Fu, X., Matsuyama, M., Hou, Y.,
Feng, G., and Wickersham, I.R. (2023). Third-generation rabies viral vectors allow nontoxic
retrograde targeting of projection neurons with greatly increased efficiency. Cell Rep.
Methods *3*, 100644.

25. Jin, L., Sullivan, H.A., Zhu, M., Lavin, T.K., Matsuyama, M., Fu, X., Lea, N.E., Xu, R., Hou,
Y., Rutigliani, L., et al. (2024). Long-term labeling and imaging of synaptically connected
neuronal networks in vivo using double-deletion-mutant rabies viruses. Nat. Neurosci. *27*,
373–383.

26. Dohaku, R., Yamaguchi, M., Yamamoto, N., Shimizu, T., Osakada, F., and Hibi, M. (2019).
Tracing of afferent connections in the zebrafish cerebellum using recombinant rabies virus.
Front. Neural Circuits *13*, 30.

27. Satou, C., Neve, R.L., Oyibo, H.K., Zmarz, P., Huang, K.-H., Arn Bouldoires, E., Mori, T.,
Higashijima, S., Keller, G.B., and Friedrich, R.W. (2022). A viral toolbox for conditional and
transneuronal gene expression in zebrafish. eLife *11*, e77153.

28. Mundell, N.A., Beier, K.T., Pan, Y.A., Lapan, S.W., Göz Aytürk, D., Berezovskii, V.K., Wark,
A.R., Drokhlyansky, E., Bielecki, J., Born, R.T., et al. (2015). Vesicular stomatitis virus
enables gene transfer and transsynaptic tracing in a wide range of organisms. J. Comp.
Neurol. *523*, 1639–1663.

29. Ma, M., Kler, S., and Pan, Y.A. (2020). Structural neural connectivity analysis in zebrafish

with restricted anterograde transneuronal viral labeling and quantitative brain mapping. Front. Neural Circuits *13*, 85.

30. Kler, S., Ma, M., Narayan, S., Ahrens, M.B., and Pan, Y.A. (2021). Cre-dependent anterograde transsynaptic labeling and functional imaging in zebrafish using VSV with reduced cytotoxicity. Front. Neuroanat. *15*, 758350.

31. Federspiel, M.J., Bates, P., Young, J.A., Varmus, H.E., and Hughes, S.H. (1994). A system for tissue-specific gene targeting: transgenic mice susceptible to subgroup a avian leukosis virus-based retroviral vectors. Proc. Natl. Acad. Sci. *91*, 11241–11245.

32. Callaway, E.M., and Luo, L. (2015). Monosynaptic circuit tracing with glycoprotein-deleted rabies viruses. J. Neurosci. *35*, 8979–8985.

33. Miyamichi, K., Shlomai-Fuchs, Y., Shu, M., Weissbourd, B.C., Luo, L., and Mizrahi, A. (2013). Dissecting local circuits: parvalbumin interneurons underlie broad feedback control of olfactory bulb output. Neuron *80*, 1232–1245.

34. Liu, Z., Chen, O., Wall, J.B.J., Zheng, M., Zhou, Y., Wang, L., Ruth Vaseghi, H., Qian, L., and Liu, J. (2017). Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector. Sci. Rep. *7*, 2193.

35. Marshel, J.H., Mori, T., Nielsen, K.J., and Callaway, E.M. (2010). Targeting single neuronal networks for gene expression and cell labeling in vivo. Neuron *67*, 562–574.

36. Beier, K.T. (2021). The Serendipity of Viral Trans-Neuronal Specificity: More Than Meets the Eye. Front. Cell. Neurosci. *15*, 720807.

37. Bae, Y.-K., Kani, S., Shimizu, T., Tanabe, K., Nojima, H., Kimura, Y., Higashijima, S., and Hibi, M. (2009). Anatomy of zebrafish cerebellum and screen for mutations affecting its development. Dev. Biol. *330*, 406–426.

38. Hibi, M., and Shimizu, T. (2012). Development of the cerebellum and cerebellar neural circuits. Dev. Neurobiol. *72*, 282–301.

39. Namikawa, K., Dorigo, A., Zagrebelsky, M., Russo, G., Kirmann, T., Fahr, W., Dübel, S., Korte, M., and Köster, R.W. (2019). Modeling neurodegenerative spinocerebellar ataxia type 13 in zebrafish using a purkinje neuron specific tunable coexpression system. J. Neurosci. *39*, 3948–3969.

40. Sengupta, M., and Thirumalai, V. (2015). AMPA receptor mediated synaptic excitation drives state-dependent bursting in purkinje neurons of zebrafish larvae. eLife *4*, e09158.

41. Knogler, L.D., Kist, A.M., and Portugues, R. (2019). Motor context dominates output from purkinje cell functional regions during reflexive visuomotor behaviours. eLife *8*, e42138.

42. Saunders, A., Johnson, C., and Sabatini, B. (2012). Novel recombinant adeno-associated viruses for cre activated and inactivated transgene expression in neurons. Front. Neural Circuits *6*, 47.

43. Takeuchi, M., Yamaguchi, S., Yonemura, S., Kakiguchi, K., Sato, Y., Higashiyama, T., Shimizu, T., and Hibi, M. (2015). Type IV collagen controls the axogenesis of cerebellar granule cells by regulating basement membrane integrity in zebrafish. PLOS Genet. *11*, e1005587.

44. Matsui, H., Namikawa, K., Babaryka, A., and Koster, R.W. (2014). Functional regionalization of the teleost cerebellum analyzed in vivo. Proc. Natl. Acad. Sci. *111*, 11846–11851.

45. Morikawa, K., Furuhashi, K., de Sena-Tomas, C., Garcia-Garcia, A.L., Bekdash, R., Klein, A.D., Gallerani, N., Yamamoto, H.E., Park, S.-H.E., Collins, G.S., et al. (2020).

Photoactivatable Cre recombinase 3.0 for in vivo mouse applications. Nat. Commun. *11*, 2141.

46. Kim, E.J., Jacobs, M.W., Ito-Cole, T., and Callaway, E.M. (2016). Improved Monosynaptic Neural Circuit Tracing Using Engineered Rabies Virus Glycoproteins. Cell Rep. *15*, 692–699.

47. Faber, M., Pulmanausahakul, R., Hodawadekar, S.S., Spitsin, S., McGettigan, J.P., Schnell, M.J., and Dietzschold, B. (2002). Overexpression of the rabies virus glycoprotein results in enhancement of apoptosis and antiviral immune response. J. Virol. *76*, 3374–3381.

48. Ohara, S., Sato, S., Oyama, K., Tsutsui, K.-I., and Iijima, T. (2013). Rabies virus vector transgene expression level and cytotoxicity improvement induced by deletion of glycoprotein gene. PLoS One *8*, e80245.

49. Cong, L., Wang, Z., Chai, Y., Hang, W., Shang, C., Yang, W., Bai, L., Du, J., Wang, K., and Wen, Q. (2017). Rapid whole brain imaging of neural activity in freely behaving larval zebrafish (danio rerio). eLife *6*, e28158.

50. Leergaard, T.B., and Bjaalie, J.G. (2022). Atlas-based data integration for mapping the connections and architecture of the brain. Science *378*, 488–492.

51. Du, X., Xu, B., Zhang, Y., Chen, M., and Du, J. (2018). A transgenic zebrafish model for in vivo long-term imaging of retinotectal synaptogenesis. Sci. Rep. *8*, 14077.

52. Kawakami, K. (2005). Transposon tools and methods in zebrafish. Dev. Dyn. *234*, 244–254.

53. Osakada, F., and Callaway, E.M. (2013). Design and generation of recombinant rabies virus vectors. Nat. Protoc. *8*, 1583–1601.

54. Lin, K.-Z., Li, L., Ma, W.-Y., Yang, X., Han, Z.-P., Luo, N.-S., Wang, J., and Xu, F.-Q. (2022). A rabies virus– based toolkit for efficient retrograde labeling and monosynaptic tracing. Neural Regen. Res. *0*, 0.

55. Preibisch, S., Saalfeld, S., and Tomancak, P. (2009). Globally optimal stitching of tiled 3D microscopic image acquisitions. Bioinformatics *25*, 1463–1465.

56. Avants, B.B., Yushkevich, P., Pluta, J., Minkoff, D., Korczykowski, M., Detre, J., and Gee, J.C. (2010). The optimal template effect in hippocampus studies of diseased populations. Neuroimage *49*, 2457–2466.

57. Marquart, G.D., Tabor, K.M., Horstick, E.J., Brown, M., Geoca, A.K., Polys, N.F., Nogare, D.D., and Burgess, H.A. (2017). High-precision registration between zebrafish brain atlases using symmetric diffeomorphic normalization. GigaScience *6*.

58. Cannon, R.C., Turner, D.A., Pyapali, G.K., and Wheal, H.V. (1998). An on-line archive of reconstructed hippocampal neurons. J. Neurosci. Methods *84*, 49–54.

59. Arshadi, C., Günther, U., Eddison, M., Harrington, K.I.S., and Ferreira, T.A. (2021). SNT: a unifying toolbox for quantification of neuronal anatomy. Nat. Methods *18*, 374–377.

60. Ahrens, J., Geveci, B., and Law, C. (2005). ParaView: An End-User Tool for Large-Data Visualization. In Visualization Handbook (Elsevier), pp. 717–731.

61. Schlegel, P., Barnes, C., Champion, A., Dokato, Jagannathan, S., Court, R., Pedigo, B., Newstein, P., Azatian, Y., and Antonio (2023). Navis-org/navis: version 1.5.0. Version v1.5.0 (Zenodo).

Figures

Figure 1. Implementation of RV*d*G[EnvA]-based neural circuit tracing in larval zebrafish

(A) Schematic of the method used to implement RV*d*G[EnvA]-based circuit tracing from

genetically defined cell types in larval zebrafish.

(B) Schematic of the three *UAS*-driven plasmids used to express the helper proteins TVA and G.

(C and D) Time-lapse confocal images of the hindbrain of larvae expressing EGFP and TVA using

UGT (C) or EGFP, TVA, and B19G using *UGTB* (D) in randomly sparse neurons (*elavl3*[+]) and

infected by SAD*d*G-mCherry[EnvA] (magenta) injected into the hindbrain. Dashed yellow circles,

starter neurons (EGFP[+]/mCherry[+]); arrows, transneuronally labeled neurons (red) and radial glia

(gray) (mCherry[+]/EGFP[−]); dashed white lines, hindbrain boundaries. C, caudal; R, rostral. Scale

bars, 50 μm.

(E) Plots of the numbers of starter neurons, transneuronally traced neurons and glia in each larva

expressing one of the three helper plasmids shown in (B) and injected with SAD*d*G-

mCherry[EnvA]. Data were collected between 6 and 10 dpi. Numbers in brackets indicate the

number of larvae examined.

(F and G) Boxplots of the convergence index for the connection of transneuronally traced neurons

(F) and glia (G) to starter neurons shown in (E). Center, median; bounds of box, first and third

quartiles; whiskers, minimum and maximum values.

**$P < 0.01$, ***$P < 0.001$ (nonparametric two-tailed Mann-Whitney test).

Figure 2. Optimization and verification of RV*d*G[EnvA]-based retrograde monosynaptic tracing

(A) Schematic of the method used to trace inputs to PCs with RV*d*G[EnvA] in larval zebrafish. One of the three helper plasmids was selected to be combined with subsequent injection of SAD*d*G-mCherry[EnvA] or CVS*d*G-tdTomato[EnvA]. The virus-injected larvae were raised at either 28 or 36°C, and the resulting labeling was monitored from 2 to 16 dpi.

(B) Schematic of the zebrafish cerebellar circuit related to inputs to PCs.

(C) Top, examples of *in situ* complementation in PCs under six tracing conditions. Confocal images and corresponding schematic illustrations of one cerebellar hemisphere are shown. Bottom, plots of the numbers of starter neurons, transneuronally traced neurons and glia in each

larva under one of the six tracing conditions at 6 - 10 dpi. Dashed lines, the cerebellar boundaries.
CI, convergence index of traced neurons. Numbers in brackets indicate the number of larvae
examined.
(D-F) CI for the connection of transneuronally traced neurons to starter PCs (D), the ratio of the
number of traced neurons to that of glia (E), and the proportion of infected fish (F, 3 to 5
independent experiments) under the six tracing conditions shown in (C, bottom).
(G), Confocal image of a larva infected with CVSdG-tdTomato[EnvA] that was trans-
complemented with CVS N2cG at 36°C. Dashed yellow circle, starter PC; dashed white line, the
cerebellar boundary.
(H and I) Enlarged view of the boxed regions in (G), showing transneuronally traced parallel
fibers of GCs (H) and two IOCs (I). Blue arrows, somata of IOCs.
(J) Schematic of the experiment for verifying the functional connection between traced GCs and
the starter PC in larvae infected with CVSdG-tdTomato[EnvA] that was trans-complemented with
advanced CVS N2cG at 36°C. Ca^{2+} imaging was conducted on the starter PC in response to
electrical stimulation (ES) at the GC position before and after two-photon laser ablation of the GC.
(K) Average Ca^{2+} response (5 trials) of the same starter PC before (red) and after (black) GC
ablation. The shadowed area represents SEM.
(L) Quantification of the amplitude of PC Ca^{2+} responses before and after GC ablation. The
number in bracket indicate the number of GC-PC pairs examined.
Scale bars, 50 μm. SAD/CVS|B19G/N2cG/A-N2cG|28/36°C, the tracing conditions written as the
G-deleted RV strain that was injected | the helper glycoprotein used for trans-complementation |
fish rearing temperature after virus injection. C, caudal; L, lateral; R, rostral. Boxplots in (D and
L): center, median; bounds of box, first and third quartiles; whiskers, minimum and maximum
values. n.s., not-significant; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (nonparametric two-tailed Mann-
Whitney test in (D-F); two-tailed unpaired Student's t test in (L)). Error bars denote SEM..
804

Figure 3. Neuronal health and physiology appeared normal for at least 10 days after RV
infection

(A) Survival rate of larvae injected with SAD*d*G-mCherry[EnvA] or CVS*d*G-tdTomato[EnvA].
Five independent experiments were repeated.

(B) Time-lapse confocal images showing a starter cell (unfilled white arrowhead) present at 2 dpi
disappears by 8 dpi, and two starter cells (white arrowheads) that remain present from 2 to 8 dpi.
Boxed regions are enlarged on the right. Dashed lines indicate the cerebellar boundaries.

(C) Cumulative proportion of starter cells that have appeared at 2, 4, 6 and 10 dpi to the total
number of observed starter cells under the three tracing conditions, showing the continuous
appearing of starter cells at 2 - 6 dpi. Numbers in brackets indicate the number of larvae
examined. The shadowed area represents SEM.

(D) Summary of the lifetime of starter cells appeared at 2 dpi under the three tracing conditions.
The color scheme follows the same conventions shown in (C). Numbers in brackets indicate the
number of starter cells examined.

(E) Schematic illustration of *in vivo* whole-cell recording on a starter PC at 11 dpi.

(F) Example traces of spontaneous EPSCs (left) and EPSPs (right) from starter PCs.

(G) Example trace of spikes in response to current-step injection into a starter PC.

(H) Confocal image (top) and Ca^{2+} responses (bottom) of a starter PC (dashed orange circle) and
non-starter PC (dashed blue circle) to visual stimuli at 10 dpi.

(I-L) Boxplots (I and J) and distributions (K and L) of Ca^{2+} activity event number (I and K) and
area under curve (AUC) of Ca^{2+} activity events (J and L) of starter PCs (n = 88) and non-starter
PCs (n = 127) in larvae (N = 8) at 6 - 10 dpi.

Scale bars, 5 μm (H), 10 μm (B), 20 μm (E). SAD/CVS|B19G/N2cG/A-N2cG|28/36°C, the tracing

conditions written as the G-deleted RV strain that was injected | the helper glycoprotein used for

trans-complementation | fish rearing temperature after virus injection. L, lateral; R, rostral.

Boxplots in (D, I and J): center, median; cross symbol, mean; bounds of box, first and third

quartiles; whiskers, minimum and maximum values. n.s., not-significant; *$P < 0.05$; ***$P < 0.001$

(two-tailed unpaired Student's t test). Error bars denote SEM.

Figure 4. Input cell-type-specific tracing and circuit reconstruction in the cerebellum

(A) A single Cre-Switch[42] transgene (*elavl3:Tetoff-DO_DIO-Hsa.H2B-mTagBFP2_tdTomato-CAAX*; abbr., *elavl3DoDioBR*) was generated to achieve a reference transgene (*elavl3:H2B-mTagBFP2*) for image registration and the Cre-dependent reporter expression (*tdTomato-CAAX*) specifically in neurons (*elavl3+*). Two ORFs flanked by two pairs of LOX sites oriented oppositely

with the activation of forward-oriented *H2B-mTagBFP2* ORF by the *elavl3* promotor in global neurons. After Cre-expressing virus infection, Cre acts on either pair of LOX sites (LOXP (gray triangles) or LOXP2272 (black triangles)), resulting in stable inversion of the two ORFs and the activation of *tdTomato-CAAX* ORF by *elavl3* promotor in initially infected starter neurons and transneuronally infected input neurons.

(B) Schematic of building PC input atlas in larval zebrafish. The *elavl3DoDioBR* reporter fish (pan-neuronal H2B-mTagBFP2$^+$) expressing helper proteins in PCs (sfGFP$^+$/mTagBFP2$^+$) were injected with Cre-expressing CVS*d*G[EnvA], which were trans-complemented in starter PCs and spread to direct inputs, resulting in Cre-dependent expression of the reporter protein (tdTomato-CAAX) in *elavl3$^+$* starter PCs (sfGFP$^+$/tdTomato$^+$/mTagBFP2$^-$) and input GCs (sfGFP$^-$/tdTomato$^+$/mTagBFP2$^-$). The pan-neuronally-expressing *H2B-mTagBFP2* in *elavl3DoDioBR* serves as the reference and bridge for the registration of reconstructed circuits. BCs, Bergmann glia; GCs, granule cells; PCs, Purkinje cells.

(C) Example of *in situ* complementation (11 dpi) in WT larvae (left) or *elavl3DoDioBR* reporter larvae (right) infected with CVS*d*G that was trans-complemented with CVS N2cG at 36°C, showing exclusive labeling of neurons in tdTomato$^+$-only traced cells when using Cre-dependent *elavl3DoDioBR* reporter fish. Confocal images (top) and corresponding schematic illustrations (bottom) are shown. Dashed yellow circles, starter PCs; dashed lines, the cerebellar boundaries. Black arrows indicate the reconstructed neurons shown in (D).

(D) Dorsal 3D view of the reconstructed starter PC and two input GCs. Dashed line indicates the cerebellar boundary. The color scheme follows the same conventions as in (F).

(E) Left, single horizontal section view of *Tg(2×en.cpce-E1B:tdTomato-CAAX)* (red), *Tg(cbln12:GAL4FF);Tg(5×UAS:EGFP)* (green), and *elavl3DoDioBR* (blue) templates aligned in a common coordinate space. Right, volume rending of the cerebellar area (dashed white rectangle on the left) of the templates labeling PCs (red) and GCs (green) in outline of cerebellum region (gray).

(F) Dorsal (left) and lateral (right) 3D view of reconstructed starter PCs (n = 24) and input GCs (n = 58) in the cerebellum (gray). Different subcellular parts of PCs and GCs are color-coded. Data were collected from 13 larvae at 10 - 16 dpi (15 - 21 dpf).

(G and H) Ipsilateral *vs* contralateral proportions of GC inputs to PCs in the left and right cerebellum. Analysis were performed using the reconstructed data in (E) and the viral tracing data shown in Figures 2C and 2D for (G) and (H), respectively. Numbers in brackets indicate the number of larvae examined which only had starter PCs on one cerebellar hemisphere.

(I) Subtypes of reconstructed PCs and GCs. Two typical cells are shown for each subtype.

(J) Example showing three GCs traced from a single PC.

(K) Summed input proportion of two subtypes of GCs to two subtypes of PCs. Cases with only one subtype of PC and clearly identifiable GCs were included. The cell numbers are as follows: PC1, 12; PC2, 6; GC1, 19; GC2, 18.

(L) Schematic representation of subtype-specific connectivity patterns from GCs to PCs. High saturation of color indicates a higher input strength. CCe, corpus cerebelli; EG, eminentia granularis; LCa, lobus caudalis cerebelli; Va, valvula cerebelli.

Scale bars, 50 μm. A, anterior; C, caudal; L, lateral; R, rostral; V, ventral. Dashed vertical lines indicate the midline of the cerebellum.

Figure 1—figure supplement 1. No infectiousness of RV*d*G[EnvA] in the absence of TVA

(A) Injection of SAD*d*G-mCherry[EnvA] into the brain of WT larvae. Left, schematic; Right,

time-lapse confocal images of the optic tectum of a WT larva injected with the virus.

(B) Time-lapse confocal images of the optic tectum of a *Tg(apoeb:LY-EGFP)* larva injected with

SAD*d*G-mCherry[EnvA], showing the location of mCherry-positive puncta in EGFP-positive

microglia.

Scale bars, 50 μm. Arrowheads, fluorescent-positive punctate structure with no discernible cellular

morphology under high laser excitation; dashed lines, boundaries of the cell body layer of the

optic   tectum.   C,   caudal;   R,   rostral.

Figure 2—figure supplement 1. Time-lapse images of trans-complemented tracing from
Purkinje cells in the cerebellum under different tracing conditions
(A, B, and D-F) Time-lapse (2, 4, and 6 dpi) confocal images of *in situ* complementation in
Purkinje cells (PCs) under tracing conditions of SAD|B19G|28°C (A), CVS|B19G|28°C (B),
SAD|B19G|36°C (D), CVS|N2cG|36°C (E), and CVS|A-N2cG|36°C (F). The schematic drawing
of the advanced helper plasmid (*UGNT-A*) is shown in (F). Dashed yellow circles, starter PCs;
dashed white lines, the cerebellar boundaries. L, lateral; R, rostral. Scale bars, 20 μm.
(C and G) Summary of CI for the connection of traced neurons, glia or all cells to starter PCs at 2,
4, 6 and10 dpi under the first two tracing conditions at 28°C (C) and the last three tracing
conditions at 36°C (G). Significant intergroup differences exist in CI for traced neurons, glia or all
cells (see the *P* values). There are no statistical differences between 6 and 10 dpi in any of the
groups (n.s.). Two-way ANOVA and nonparametric two-tailed Mann-Whitney test for intergroup

and intragroup statistics, respectively. Error bars denote SEM. Numbers in brackets indicate the
number of larvae examined.

(H and I) Boxplots of CI for the connection of traced glia (H) or all cells (I) to starter PCs under
six tracing conditions at 6 - 10 dpi. See also Figures 2C-2E. Center, median; bounds of box, first
and third quartiles; whiskers, minimum and maximum values. n.s., not-significant; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$ (nonparametric two-tailed Mann-Whitney test).

914 **Figure 1,2—table supplement 1. The efficiency of viral transfer under different tracing**
915 **conditions in larval zebrafish.** The convergence index (CI) was calculated as the number of
916 traced cells divided by the number of starter cells.

Promoter	RV Strain	Helper plasmid	Glyco-protein	Temp (°C)	CI_{neuron}	CI_{glia}	CI_{all}
elavl3 (Pan-neuron)	SAD	*UGT*	No	28	0	0	0
	SAD	*UGTB*	B19G	28	0.44 ± 0.11	2.50 ± 0.65	2.94 ± 0.72
	SAD	*UGBT*	B19G	28	1.55 ± 0.29	4.41 ± 1.48	5.97 ± 1.69
cpce (Purkinje cell)	SAD	*UGBT*	B19G	28	0.06 ± 0.03	0.10 ± 0.05	0.16 ± 0.06
	CVS	*UGBT*	B19G	28	0.41 ± 0.07	1.58 ± 0.23	1.99 ± 0.26
	CVS	*UGNT*	N2cG	28	1.02 ± 0.18	4.20 ± 0.94	5.22 ± 1.05
	SAD	*UGBT*	N2cG	36	4.96 ± 1.01	14.91 ± 2.49	19.87 ± 3.14
	CVS	*UGNT*	N2cG	36	11.61 ± 2.43	13.21 ± 2.44	24.82 ± 4.58
	CVS	*UGNT-A*	N2cG	36	20.35 ± 3.45	27.03 ± 4.37	47.38 ± 7.25

918   **Figure 1-4—table supplement 1. Properties of currently available viral circuit tracing**
919   **systems for zebrafish.**

	This work	Dohaku et al. 2019 [26]	Satou et al. 2022 [27]		Ma et al. 2020 [29]	Kler et al. 2021 [30]
Age	3 - 21 dpf	Adult	Adult	7 - 14 dpf	2.5 - 9 dpf	3 - 9 dpf
Region being studied	Brain	Brain	Brain	Spinal cord	Brain	Brain
Virus strains	RV-SAD*d*G[EnvA]; RV-CVS*d*G[EnvA]	RV-SAD*d*G[EnvA]	RV-SAD*d*G[EnvA]	RV-SAD*d*G[EnvA]	rVSV*d*G[VSVG]	rVSV(M51R)*d*G[VSVG]
Glycoprotein	SAD-B19G; CVS-N2cG	SAD-B19G	SAD-B19G	SAD-B19G	VSVG	VSVG
Spread direction	Retrograde	Retrograde	Retrograde	Retrograde	Anterograde	Anterograde
Highest Tracing efficiency (CI)	~20	N.A.	~1	N.A.	N.A.	N.A.
Genetic targeting of starter cells?	Y (EnvA-TVA)	Y (EnvA-TVA)	Y (EnvA-TVA)	Y (EnvA-TVA)	N	N
Methods for helper proteins expression	Transient expression using single plasmid	Transgenesis	HSV1	HSV1	Lentivirus	Lentivirus
Co-expression of TVA & G?	Y	N	N	N	N	N
Synaptic specificity verification?	Y	N	N	N	N	N
Input or output selectivity?	Y	N	N	N	N	Y
Neuron viability test?	Y	N	Y	N	N	N
Life span of infected cells (d)	10 - 14 (starter cells)	N.A.	~6 (starter cells)	N.A.	<6 (traced cells)	<6 (traced cells)
Rearing temperature for virus-injected fish (°C)	28 & 36	~35	36	36	28	28
Experimental procedure feasibility ■1.Helper protein expression ■Plasmid microinjection ■Transgenic fish ■Helper virus ■2.Virus injection ■3.Fluorescence signal amplification ■Brain slicing ■Immunohistochemistry ■4.Imaging	■■■	■■■■■■	■■■■■■	■■■■■	■■■	■■■

920   **Y, yes; N, no; N.A., not available**