# Make-or-break prime editing for bacterial genome engineering

Monica Rengifo-Gonzalez[1,#], Maria-Vittoria Mazzuoli[1,#], Axel B. Janssen[1,#], Anne-Stéphanie Rueff[1], Xue Liu[2,*], Jan-Willem Veening[1,*]

[1]Department of Fundamental Microbiology, Faculty of Biology and Medicine, University of Lausanne, Biophore Building, CH-1015 Lausanne, Switzerland

[2] Department of Pathogen Biology, Base for International Science and Technology Cooperation: Carson Cancer Stem Cell Vaccines R&D Center，International Cancer Center，Shenzhen University Medical School, Shenzhen 518055, Guangdong, China.

[#]These authors contributed equally

*Correspondence to Xue Liu (xueliu@szu.edu.cn) or Jan-Willem Veening (Jan-Willem.Veening@unil.ch), tel: +41 (0)21 6925625

## Abstract

**CRISPR-Cas9 has revolutionized genome engineering by allowing precise introductions of DNA double-strand breaks (DSBs). However, genome engineering in bacteria is still a complex, multi-step process requiring a donor DNA template for homology-directed repair of DSBs. Prime editing circumvents this need as the repair template is provided within the prime editing guide RNA (pegRNA). Here, we developed make-or-break Prime Editing (mbPE) that allows for precise and effective genetic engineering in the opportunistic human pathogen *Streptococcus pneumoniae*. In contrast to traditional prime editing in which a nicking Cas9 is employed, mbPE harnesses wild type Cas9 in combination with a pegRNA that destroys the seed region or protospacer adjacent motif. Since most bacteria poorly perform template-independent end joining, correctly genome-edited clones are selectively enriched during mbPE. We show that mbPE can be used to introduce point mutations, deletions and targeted insertions, including protein tags such as a split luciferase, at selection efficiencies of over 93%. mbPE enables sequential genome editing, is scalable, and can be used to generate pools of mutants in a high-throughput manner. The mbPE system and pegRNA design guidelines described here will ameliorate future bacterial genome editing endeavors.**

## Introduction

The use of CRISPR/Cas9 technology has facilitated genetic engineering for a wide range of bacteria, including difficult or laborious to transform species such as *Staphylococcus aureus*, *Clostridium* spp. and *Streptomyces* spp.[1–5]. The basis for CRISPR/Cas9-based bacterial genome editing is the targeting of *Streptococcus pyogenes* Cas9 to the genetic location of interest by a single guide RNA (sgRNA) followed by production of a double-stranded break (DSB)[6]. By co-transforming a donor template DNA, the DSB can be repaired by homology-directed repair, resulting in the desired genome edit[1,7]. This is particularly efficient in bacteria, as most strains lack (or poorly perform) template-independent end joining mechanisms such as non-homologous end joining (NHEJ) or alternative end joining[2,8,9]. Thus, only clones edited at the DSB and thereby disrupting the protospacer adjacent motif (PAM) or sgRNA seed region will survive. The major drawback of CRISPR/Cas9-based genome editing in bacteria is that several steps are involved such as the cloning of a sgRNA and the production or cloning of a template DNA[1,10]. Especially with hard-to-transform or slow growing bacteria, each of these steps can take several days, if not weeks.

Recently, prime editors such as PE2 were described in which a Cas9 nickase is fused to an engineered reverse transcriptase[7,11]. PE2 has the advantage that it does not introduce DSBs, which, in

eukaryotes typically leads to error-prone end-joining via NHEJ and has the risk of inducing
chromosomal rearrangements[7]. In addition, prime editing uses a pegRNA for both targeting of the PE2
complex as well as a template for DNA repair in which the desired mutation is encoded. The pegRNA
contains a spacer sequence to direct targeting of the prime editor to the PAM site, a primer binding site
(PBS) that facilitates the hybridization of the 3′-end of the nicked DNA strand to the pegRNA and
includes a reverse transcriptase template (RTT) that carries the intended genetic modification[11]. Because
of these advantages, prime editing has rapidly gained traction and has been successfully applied to
engineer the genomes in various eukaryotes including plants[12–15]. However, prime editing has not yet
been widely applied for bacterial genome editing, probably because of limitations in the selection
efficiency during the introduction of PE2-mediated mutations[16,17].

Here, we describe make-or-break prime editing (mbPE) as an efficient genome editing method
in the opportunistic human pathogen *Streptococcus pneumoniae*. mbPE is a nuclease-based prime
editing system that uses wild type *S. pyogenes* Cas9 fused to a bacterial codon-optimized reverse
transcriptase under inducible control. Using mbPE, we were able to make point mutations, deletions
over 100 bps as well as insertions up to 52 bps in a targeted fashion at high selection efficiencies. Using
a pooled high-throughput approach, we established mbPE design rules and show that an optimal
pegRNA, to insert a single adenine base, consists of a PBS of 16 nt and a RTT of 15 nt. We provide
pegRNA vectors with different antibiotic selection markers allowing for sequential pneumococcal
genome edits. The mbPE system developed here represents one of the most efficient and scalable
bacterial genome engineering platforms currently available.

## Results

### Prime editing in *S. pneumoniae*

Several Cas9-based tools have been developed for *S. pneumoniae*[18–22], but prime editing has not been
implemented yet. To establish a practical prime editing system for *S. pneumoniae*, we first focused on
the prime editing tool PE2. PE2 is a fusion of the *S. pyogenes* Cas9$^{\text{H840A}}$ nickase domain and an
engineered Moloney murine leukemia virus reverse transcriptase domain (MMLVRT)[11]. Previously,
we described tight inducible control by the addition of tetracycline-group molecules (including
anhydrotetracycline (aTc)) of a catalytically inactive Cas9 (dCas9: D10A, H840A) expressed as a single
copy from the non-essential chromosomal CEP locus (P$_{tet}$-*dcas9*) for CRISPR interference (CRISPRi)
in *S. pneumoniae*[21]. In a first step to a functioning PE2 for *S. pneumoniae*, we mutated the alanine
residue at position 10 of the P$_{tet}$-*dcas9* construct back to an aspartic acid to create P$_{tet}$-*cas9*$^{\text{H840A}}$. The
engineered MMLVRT domain and a linker domain from PE2 were codon optimized for optimal

expression in low GC-rich bacteria such as *S. pneumoniae* and cloned to the C-terminus of the Cas9

nickase resulting in strain VL4199 ($P_{tet}$-*cas9*[H840A]-MMLVRT or $P_{tet}$-PE2[S.pn]) (Fig. 1A).

Next, a pegRNA construction and *S. pneumoniae* integration vector was designed in which the

pegRNA is constitutively expressed from the constitutive P3 promoter including its +1 adenine

transcription start site to ensure that all cloned pegRNAs are efficiently transcribed and do not contain

undesirable uridines at the 5' end[22–24]. For easy cloning, an *mCherry* cassette flanked by Esp3I

(isoschizomer of BsmBI) sites is present allowing for rapid, oligo-based, Golden Gate cloning of

pegRNAs (Fig. 1B). Sequence elements facilitating one-step library preparation for NGS were also

introduced[25] resulting in pegRNA cloning vectors pVL4132, pVL4133 and pVL4393 (carrying 7

thymine nucleotides, a *S. pneumoniae* terminator or a *S. pneumoniae tRNA-asp-1* after the extension

sequence, respectively). pegRNAs consist of four elements: 1) a 20 nt spacer sequence for targeting of

the prime editor, 2) a scaffold sequence for binding to Cas9, 3) a primer binding site (PBS) that allows

the 3′-end of the nicked DNA strand to hybridize to the pegRNA, and 4) a reverse transcriptase template

(RTT) containing the desired edit[11]. Since bacterial transcription termination leaves nucleotides at the

3'end of the pegRNA, which may interfere with efficient prime editing downstream, we cloned *S.*

*pneumoniae tRNA-asp-1* directly following the pegRNA (Fig. 1B). This strategy has been previously

used to produce multiple individual sgRNAs from a single polycistronic operon as the endogenous

tRNA-processing system precisely cleaves both ends of the *tRNA-asp-1* precursor[26]. To test whether

the PE2[S.pn] is functional, we designed several pegRNAs using PrimeDesign[27] to introduce either a

premature stop codon in the firefly luciferase gene (*luc*) expressed in pneumococcal strain VL4200, or

to introduce a repairing mutation in strain VL4297 in which *luc* has a premature stop codon (Fig. 1C).

Interestingly, induction of PE2[S.pn] together with a targeting pegRNA did not result in significant growth

delay, suggesting that *S. pneumoniae* can tolerate and readily repair ssDNA breaks (Fig. S1). As shown

in Fig. 1D, approximately 2% of clones in which PE2[S.pn] was induced demonstrated the desired

phenotype. Sanger sequencing confirmed the bioluminescence observations (Fig. S1B).

Recently, it was shown that a structured 3'-end of a pegRNA increases stability and improves

editing efficiency in human cells[28]. In addition, a functional mismatch repair system was shown to

negatively affect editing efficiencies[29,30]. To test whether the addition of a structured RNA motif at the

3'-end would also improve PE2 editing efficiency in *S. pneumoniae*, we replaced the tRNA sequence

for the structured and highly efficient *S. pneumoniae rpsT* transcriptional terminator[31] (SpTer) or a

stretch of 7 thymine nucleotides (7 Ts) (Fig. 1E). As shown in Fig. 1D, both pegRNAs showed improved

efficiency of inserting a single A in *luc* (~3%) through PE2[S.pn]-mediated editing, compared to the tRNA-

Asp1 system. Repairing an internal stop codon (TAA to GCT) in *luc* was approximately 3 times more

efficient with the SpTer terminator compared to the tRNA-Asp1 construct (Fig. 1D). Deleting the

pneumococcal mismatch repair gene *hexA*[32] further enhanced PE2[S.pn] editing efficiency in the constructs

with either the 7T or SpTer terminators (Fig. 1F). These results demonstrate that prime editing using PE2$^{S,pn}$ in combination with a structured pegRNA works in *S. pneumoniae* and can be used to incorporate insertions and mutations, albeit at a relatively low selection efficiency.

**Figure 1. Prime editing 2 (PE2) in *S. pneumoniae*.** (**A**) Schematic overview of PE2$^{S,pn}$. (**B**) Schematic overview of the general cloning strategy of pegRNAs (shown is the *tRNA-asp-1* containing vector pVL4393 as backbone). (**C**) Editing strategy used to test efficiency of PE2$^{S,pn}$ in pneumococcal strains expressing either a functional luciferase (VL4200) or a *luc* gene interrupted by a premature stop codon (VL4297). (**D**) Induction of PE2$^{S,pn}$ by aTc in combination with expression of a pegRNA without 3' leader, by processing of the *tRNA-asp-1*, leads to accurate gene editing in approximately 2% of *S. pneumoniae* cells. Induction of PE2$^{S,pn}$ in combination with 3' extended pegRNAs provides an editing efficiency of approximately 3%. Cells were grown in C+Y medium containing luciferin substrate. The maximal bioluminescence (relative light units; RLU) reached, normalized by the optical density (OD), of each clone is shown (AU, arbitrary units). RLU/OD values were normalized against the baseline level of medium luminescence obtained for each experiment. (**E**) Schematic overview of pegRNAs

133    containing the *S. pneumoniae rpsT* terminator (SpTer) or a stretch of 7 Ts. (**F**) Prime editing is slightly enhanced

134    in the absence of mismatch repair in *S. pneumoniae*.

## Make-or-break prime editing (mbPE) in *S. pneumoniae*

Since prime editing depends on successful hybridization of the 3'end of the nicked DNA to the

pegRNA[11], we wondered if prime editing would also be functional when, instead of a nick, a DSB is

introduced. Recent work in human HEK293T cells showed efficient prime editing utilizing Cas9-

induced DSBs that largely depend on NHEJ[33–35]. To test if DSB-driven prime editing (here after called

make-or-break prime editing: mbPE) would also work in bacteria lacking NHEJ, we first mutated the

alanine residue at position 840 within the Cas9 nickase moiety of $PE2^{S.pn}$ back to a histidine. This led

to full restoration of wild type Cas9, resulting in strain VL5149 ($P_{tet}$-*cas9*-MMLVRT or $mbPE^{S.pn}$) (Fig.

2A). This construct was combined with *luc* expression (strain VL5155) and the same set of pegRNAs

containing the SpTer or 7 Ts as previously used to verify the efficiency of the nickase-based $PE2^{S.pn}$

(Fig. 1E). All pegRNAs were designed to either disrupt the PAM or the seed region within *luc*. Thus,

only correctly edited bacteria are expected to survive the expression of $mbPE^{S.pn}$, as unedited repairs

will lead to continued DSBs, hence the name make-or-break prime editing. Induction of $mbPE^{S.pn}$ with

aTc, in the presence of a targeting pegRNA, caused severe growth defects indicating that DSBs were

formed (Fig. S2A). Strikingly, surviving colonies (approximately 1 - 5%, depending on the pegRNA),

predominantly displayed the correct luciferase phenotype (Fig. 2B). Sanger sequencing confirmed that

over 93% of colonies had the precise edit as encoded by the pegRNA template (Fig. S2B). Colonies

that did not display the correct *luc*-edit either had suppressor mutations in the Cas9 domain, or deletions

in the $P_{tet}$ promoter driving the expression of $mbPE^{S.pn}$ (Fig. S2C). Note that, if desired, strains can

easily be cured from the prime editing system by transforming with plasmid pPEPY to exchange the

$mbPE^{S.pn}$ cassette with a kanamycin resistance marker[36]. PacBio HiFi long read whole genome

sequencing of 2 edited clones did not reveal signs of off-target editing nor any genomic rearrangements

(SRA accession number PRJNA1062582).

**Figure 2. Make-or-break prime editing selectively enriches correctly edited clones.** (**A**) Schematic overview
of mbPE$^{S.pn}$. Binding of the spacer sequence to the target region leads to formation of DSBs by the Cas9 domain.
After, the PBS can hybridize to its complementary sequence. The reverse transcriptase domain of mbPE$^{S.pn}$
subsequently synthesizes cDNA using the RTT as template. (**B**) Schematic depicting the mechanism leading to
selective survival of edited clones. The hybridization of the 3'-flap containing the edit is in equilibrium with the
unedited strand. Repair with the edited region will prevent repeated DSBs by Cas9. (**C**) Induction of mbPE$^{S.pn}$ by
aTc in combination with expression of a structured pegRNA leads to accurate gene editing. Approximately 93%
of surviving *S. pneumoniae* cells after induction are accurately edited. Cells were grown in C+Y medium
containing luciferin substrate. The maximal bioluminescence reached of each clone is shown (RLU: relative light
units. OD: optical density).

**Scalable pooled pegRNA library construction shows efficient installment of base pair
substitutions**

To explore the potential of mbPE$^{S.pn}$, we generated a pool of different pegRNAs targeting *luc* (Table
S1), followed by Golden Gate cloning (Fig. 3A). A total of 873 oligos were designed and synthesized
as a pool, containing different PBS and RTT sequences to introduce various genome edits. The pegRNA
pool was cloned into plasmid pVL4134 that already contains the *luc*-targeting spacer sequence. The

resulting pool was subsequently used to transform strain VL5155 carrying *luc* and the aTc-inducible

mbPE*S.pn* (see Methods). The library was plated on agar lacking or containing 20 ng/ml aTc to induce

mbPE*S.pn*. Quadruplicate induced and uninduced experiments, each with a minimum of 40000 colonies,

were pooled, and their pegRNA contents and edited *luc* region were sequenced (Fig. 3A). In the

uninduced pool, all but 12 of the designed pegRNAs were present, and we did not observe some of the

pegRNAs encoding the longest insertions. Strikingly, most pegRNAs were also present in the induced

pool albeit at varying abundances. By counting and comparing the number of reads coming of each

pegRNA in both conditions, we could estimate an editing efficiency score for each pegRNA for a

specific mutation (Fig. 3A). By also sequencing the *luc* gene, we confirmed that most of the designed

edits were correctly made in the pool of mutants (see Methods).

First, we examined whether mbPE*S.pn* was able to install various substitutions (a>t, a>c, a>g,

t>a, t>c, t>g, c>a, c>t, c>g) of targets within a seed region. We designed pegRNAs that have the same

20 nt *luc* spacer and 15 nt PBS, but varying RTTs encoding the different substitutions (and the necessary

*luc* seed disruptions) (Fig. 3B, Table S1). As shown in Fig. 3C, pegRNAs encoding all substitutions

were present at high frequencies under induced conditions suggesting that the installment of single base

pair substitutions is highly efficient with mbPE*S.pn*. Next, we wondered if mbPE*S.pn* could introduce

multiple substitutions simultaneously. Indeed, all tested substitutions, up to the 6 adjacent substitutions

tested (CCTCTA> GGAGAT), could readily be introduced (Fig. 3D).

**Figure 3. mbPE$^{S.pn}$ is highly scalable and efficient in making base pair substitutions. (A)** Cloning strategy to produce a library of pegRNAs targeting *luc* from a ssDNA oligo pool. The oligo pool was amplified by PCR as well as the backbone of pVL4134 that includes the *luc* spacer sequence and assembled by Golden Gate digestion/ligation followed by transformation to *S. pneumoniae* containing *luc* and mbPE2$^{S.pn}$. Chromosomal DNA was isolated from colonies grown in absence or presence of aTc (to induce mbPE2$^{S.pn}$) and PCR amplified using staggered primers followed by illumina sequencing (see Methods). 2FAST2Q[37] was used to extract and count pegRNAs and *luc* alleles. **(B)** Schematic overview of the pegRNA design encoding for different base pair substitutions. **(C-D)** Single base pair substitutions and multiple base pair substitutions are installed with high selection efficiency. **(C)** pegRNA abundance of +/- aTc samples are shown. **(D)** The fraction of *luc* reads carrying the desired mutation is shown for induced and non-induced samples on the left Y-axis. The average fraction of pegRNA reads for the substitutions is shown as dots on the right Y-axis.

**Efficient DNA insertion and deletion with mbPE**

After establishing that mbPE$^{S.pn}$ is highly efficient at introducing point mutations, we examined editing efficiencies for DNA insertions or deletions (Fig. 4A, 4C). To test DNA insertion efficiencies, we designed 181 different insertion pegRNAs targeting *luc* with a constant 20 nt spacer, a constant 15 nt PBS, a constant 17 nt RTT scaffold followed by an RTT insertion of varying length (between 1 and 181 nt) that would insert in the *luc* seed region thereby preventing Cas9 retargeting (Fig. 4A, Table S1). We analyzed pegRNA abundances of uninduced and induced libraries, as well as sequenced the *luc* genes in the pools. We observed insertions up to 52 bps, indicating that there is a maximum length that can be installed through mbPE in *S. pneumoniae* (Fig. 4B). Note that longer pegRNAs were less well represented in the insertion pegRNA pool likely because of reduced abundance due to uneven synthesis and/or cloning efficiencies (Fig. 4B). This could have influenced our ability to observe the mutations encoded in these larger pegRNAs.

A similar strategy was employed to assess deletion efficiency inside *luc*, and 200 deletion pegRNAs were designed each containing a constant 20 nt *luc* spacer and constant 15 nt PBS. The RTT's were 15 nt long, each shifted by one base pair relative to each other to create varying deletion sizes in *luc* whilst simultaneously disrupting the seed region (Fig. 4C). We observed a decrease in editing efficiency as deletion length increased, with the longest deletion observed being 184 bp (Fig. 4D). Indeed, we could readily obtain deletions at high selection efficiency when reconstructing a set of deletion pegRNAs encoding the 10, 63 or 127 base pair deletion (ranging between 80-90% efficiency). However, only 3 out of 28 tested clones showed reduced luciferase activity with the 184 bp deletion pegRNA and Sanger sequencing indicated that these were not correctly edited at the 184 bp position (Fig. 4E).

**Figure 4. mbPE for DNA insertions and genome deletions.** (**A**, **C**) Schematic overview of the approach used to test insertion and deletion efficiency in *luc* using mbPE. For pegRNAs encoding insertions, additional nucleotides were added between the RTT and PBS. For deletions, the RTT portion of the pegRNA was progressively moved within the *luc* gene, increasing deletion size, whilst the RTT remained constant. (**B**, **D**) Possibility of making insertions (**B**), and deletions (**D**) using mbPE. For each modification size, the fraction of *luc* reads carrying that mutation is displayed in bars on the left axis with bars, whilst the fraction of pegRNA reads from uninduced samples carrying the mutation is displayed with dots mapping to the right axis. (**E**) Verification of the ability to make deletions in *luc*, through targeted deletion of 10, 63, 127, or 184 bp.

## pegRNA design rules for optimal mbPE

Previous work in human cells has shown that PE2 gene editing efficiencies highly depend on RTT and PBS length[11]. To determine the optimal RTT and PBS length for mbPE$^{S.pn}$ to introduce a single adenine nucleotide insertion in *S. pneumoniae*, we designed 336 pegRNAs, each encoding the same 20-nucleotide *luc* spacer sequence but varying in RTT length (6-26 nucleotides) and in PBS length (5-20 nucleotides). Each RTT contained the insertion of a single adenine nucleotide to disrupt the *luc* seed sequence, thereby preventing retargeting of Cas9 (Fig. 5A). Differential enrichment analysis of pegRNA abundance with and without mbPE$^{S.pn}$ inducer showed relatively more surviving clones with a PBS longer than 16 nts, and an RTT longer than 11. The most enriched pegRNAs for inserting a single adenine in *luc* contained a combination of a RTT length of 14 or 15 with a PBS length of 16 or 17 nts (Fig. 5B). Together, these experiments provide guidelines for optimal pegRNA design and shows that

mbPE is a versatile editing tool capable of introducing single point mutations, multiple simultaneous substitutions, insertions and deletions.

**Figure 5. pegRNA design rules for mbPE$^{S.pn}$.** (**A**) Schematic overview of the approach used to test mbPE$^{S.pn}$ efficiency at inserting a single adenine in *luc*. (**B**) Combinations of RTT and PBS with statistically significant differences in pegRNA abundance between uninduced vs induced are outlined and overrepresented combinations show higher log2 fold changes (log$_2$FC) and are colored bright yellow. Most overrepresented pegRNAs of the induced library compared to uninduced suggest that a 16 or 17 nts for the PBS and 14 or 15 nts for the RTT yield the highest editing efficiency. Statistical significance was defined as an |log$_2$FC| > 1, and an adjusted p-value < 0.05.

## Exploring the editing distance from the seed region

The strategies employed in this study guide the Cas9 nuclease moiety of mbPE$^{S.pn}$ to cut within the seed region near the PAM motif as dictated by the spacer sequence, while the reverse transcriptase (RT) component of mbPE$^{S.pn}$ introduces the desired genome edit. This approach prevents Cas9 from retargeting the same location. The dependency on a PAM site is, however, a possible limitation of the mbPE system. For mbPE to be broadly applicable for the introduction of mutations, even in genomic

regions without a nearby PAM sequence, editing should also be possible at sites distal from the PAM.

To determine the maximum distance from the Cas9 cut site at which a single nucleotide substitution is

possible, we designed a set of 100 pegRNAs with identical spacers. Each pegRNA contained RTT

sequences to mutate the seed region (preventing retargeting) while introducing a SNP at varying

distances from the PAM site (1-100 nts from the restriction site; Fig. 6A). As shown in Fig. 6B, edits

were observed as far as 91 bp from the PAM site.

Several pegRNAs were individually cloned, each with a mutation that destroys the seed region

to insert a silent mutation in *luc* and a substitution further from the PAM site. This showed that SNPs

could be efficiently introduced at a distance of 33 base pairs from the PAM (Fig. 6C). Editing was also

possible at further distances, but with increasingly reduced efficiency, similar to what was observed in

the pooled screen (Fig. 6B). Together, this demonstrates that mbPE$^{S.pn}$ is highly versatile and can, in

principle, be used to edit nearly every base in the pneumococcal genome (see Discussion).

**Figure 6. mbPE can introduce edits at sites distal from the PAM.** (**A**) Schematic overview of the RTT design

strategy used to edit bases distal from the PAM site. (**B**) Fraction of correctly edited *luc* alleles and pegRNA read

counts of DNA extracted from cells grown in the absence (control) or presence of aTc. Edits are observed up to

91 bp from the PAM site. (**C**) Validation of the maximum editing distance through individually cloned pegRNAs.

The mutations conferred by the pegRNAs encoding a mutation at distances of 33, and 45 bps from the PAM site,

confer a premature stop codon, whilst the mutations conferred by the pegRNAs encoding mutation either 72 or

91 bps from the PAM site resulted in an glycine to cysteine (72 bps) or isoleucine to asparagine (91 bps)

substitution. The percentages indicate the fraction of clones that were phenotypically observed to be edited. The

mutation at 91 nts (*) could only be confirmed by sequencing individual clones, as the amino acid change does

not affect luciferase function. This showed an editing efficiency of 4.8 % (1 out of 21 tested colonies) with the
correct genome edit.

**mbPE to introduce a split-*luc* tag for protein-protein interactions**

Our pooled pegRNA assays indicated that DNA up to 52 bps can be inserted into the pneumococcal

genome using mbPE (Fig. 4B). To test whether mbPE could be used to introduce protein tags, we

considered the split luciferase tag. The use of split luciferase has emerged as a powerful approach to

identify protein-protein interactions in live bacterial cells in which the bait is fused to the large luciferase

component (LgBit) and the target to the small moiety (SmBit). When both parts of the split-luciferase

are in close proximity, luciferase activity is restored and light is produced in the presence of furimazine-

based (NanoLuc) substrate[38,39] (Fig. 7A). An outstanding question in the pneumococcal cell biology

field is whether the muramidase MpgA and the bifunctional class A penicillin binding protein PBP1a

are part of the so called elongasome[40–43], the machinery responsible for cell elongation. To address this

question, we fused RodZ, a known component of the elongasome, as well as MpgA, to the LgBit. We

designed a pegRNA to insert a sequence encoding a 2-amino acid flexible linker (SG), followed by the

SmBit fragment (VTGYRLFEEIL), at the C-terminus of PBP1a (Fig. 7B). Of the 12 clones tested after

selection on aTc, 1 contained the correct fusion, signifying a selection efficiency of 8% for introduction

of a 39 bps-long tag, in line with the pooled library sequencing results (Fig. 4B). After transforming the

mbPE[S.pn]-created *pbp1A-SmBit* strain with *mpgA-LgBit* or *rodZ-LgBit*, respectively, cells were grown

in microtiter plates in the presence of NanoLuc luciferase substrate and bioluminescence recorded. As

shown in Fig. 7C, we could detect a strong and reproducible signal in both strains, providing evidence

that PBP1a, MpgA and RodZ are in close proximity in space and time in live pneumococcal cells.

**Sequential genome editing with mbPE[S.pn]**

The ability to introduce several genome edits sequentially would greatly accelerate genetic engineering

projects in *S. pneumoniae*. To facilitate this, we constructed a second pegRNA cloning vector with

identical integration sites as plasmid pVL4133 but a different antibiotic resistance marker, conferring

resistance against chloramphenicol instead of spectinomycin (pVL7227) (Fig. 7D). In this way, after

the first genome edit has been made, the pegRNA can be replaced by transformation of the second

vector containing a different pegRNA. As a proof of concept, we first introduced a 63 bps deletion in

*luc* in strain VL6307 using pegRNA plasmid pVL6304, resulting in strain VL6926 (Δ63-*luc*). Next, we

designed and generated pegRNA constructs that would introduce stop mutations in either the capsule

gene *cps2A* or the major autolysin *lytA*. Both genes encode major pneumococcal virulence factors (Fig.

7D). By sequential transformation of strain VL6926 with plasmids pVL7230 (chloramphenicol

resistance, pegRNA- Δ*cps2A-stop*) and pVL5524 (spectinomycin resistance, Δ*lytA-stop*), we could

readily obtain strain VL7292 containing the three desired mutations (Δ63-*luc*, Δ*cps2A-stop*, Δ*lytA-stop*). Whole genome PacBio resequencing of two of the triple mutants verified the correct edits and demonstrated the absence of off-target edits or genomic rearrangements (SRA accession number PRJNA1062582).

**Figure 7. PBP1a is in a complex with MpgA and RodZ and sequential genome editing using mbPE.** (**A**) Schematic overview of the principle of the split *luc* system. (**B**) Schematic overview of the pegRNA used to fuse a SmBit tag to *pbp1A* using mbPE. (**C**) PBP1a interacts with MpgA and RodZ. Each dot represents the average of 15 measurements of a technical replicate, with the size of the dot representing the SEM. Control strains only expressing *pbp1A-SmBit* (D39V) or *pbp1A-SmBit* together with the abundant DNA-binding protein HU-LgBit did not demonstrate bioluminescence. (**D-E**) Sequential gene editing in *S. pneumoniae* using mbPE. Schematic overview of cloning regime with the two alternative pegRNA vector backbones (pVL4133, *spec*; pVL7227, *cat*) to generate a triple genome edited *S. pneumoniae* strain (Δ63-*luc*, Δ*cps2A-stop*, Δ*lytA-stop*) in a sequential manner.

## Discussion

The principal contribution of this work is the development of make-or-break prime editing (mbPE), a highly effective bacterial genome editing system. In contrast to traditional prime editing in which a nickase is used, mbPE utilizes a Cas9 nuclease fused to a reverse transcriptase. Combined with the high recombinogenic potential of *S. pneumoniae*[19], Cas9-induced DSBs are efficiently repaired using the edit encoded within the pegRNA that also disrupts the PAM or seed region. Since Cas9 cannot re-target correctly edited cells, as they would lack a PAM or seed region, they are efficiently selected, while unedited clones die due to repeated Cas9-induced DNA breaks. Recently developed nuclease-based prime editing systems in eukaryotic cells have also shown improved selection efficiencies[33–35,44–46]. However, these systems often produce more undesired by-products, such as insertions and deletions (indels), than the intended edits[47]. Probably because *S. pneumoniae* does not, or very poorly, performs template-independent end joining (like most other bacteria[2,8,9]), we have not observed off-target indels with mbPE[S.pn] when editing efficiencies were high. Bystander mutations, if present, were rather found near the Cas9 cleavage site that are typically also found during regular PE2 prime editing[7] or within mbPE[S.pn] itself (Fig.S2C). Long-read PacBio sequencing of triply sequential edited strains (Fig. 7) also did not show evidence of genomic rearrangements or untargeted indels.

By screening pooled libraries containing a suite of pegRNAs with different RTT and PBS lengths to introduce point mutations, we established that ideal PBS and RTT lengths for a single adenine insertion are 16 nts and 15 nts, respectively (Fig. 5). In addition, mbPE was able to make all tested nucleotide substitutions as well as insertions up to 52 bps (Figs. 3-4). As a proof of principle, we used mbPE to tag the important amoxicillin-resistance determinant Pbp1A[48] with the SmBit split luciferase moiety and unequivocally showed that Pbp1A is in close spatial and temporal proximity to MpgA and RodZ. Deletions up to 127 bps could also be efficiently introduced (Fig. 4C-E). By creating pegRNA vectors with various antibiotic resistance markers, sequential genome editing is a straightforward procedure and after every consecutive edit the pegRNA is simply replaced by the next (Fig. 7D-E). Together, this work demonstrates that mbPE is a flexible, scalable and efficient genome editing tool for *S. pneumoniae.*

**Limitations and perspectives of mbPE**

Due to its natural competence, *S. pneumoniae* has been a cornerstone of molecular biology[49]. This feature also enabled us to optimize mbPE for genome editing. It will now be interesting to test whether mbPE is also effective in other bacteria, particularly in strains that are difficult to transform. Another advantage of *S. pneumoniae* is that its genome, which is 2.04 Mb in size in case of strain D39V[31],

contains 145,892 *S. pyogenes* PAM sites (NGG) distributed fairly evenly across its sequence. This

results in one PAM site every 14 base pairs on average. If a distance of 50 bp from a PAM site is

considered as the maximum distance to be efficiently edited by mbPE$^{S.pn}$ (Fig. 6B), 99.845% of the

D39V genome would be in range for efficient substitution editing through this system, leaving only

3158 bp unavailable. Although SNPs can be efficiently introduced at locations near a PAM site, the

efficiency decreases as the distance from the PAM site increases (Fig. 6). This is thus not a major issue

for *S. pneumoniae*, but could be problematic for other bacterial species, particularly those with very low

GC content. To obtain more genome editing flexibility, Cas9 variants with relaxed or different PAM

requirements[50] could be used as nuclease within the mbPE. For instance, highly promising mbPE

nucleases would be SpRY and SpRYc, which are engineered near-PAMless *S. pyogenes* Cas9

variants[51,52].

While the *selection* efficiency of mbPE is greater than 93%, if the mutation being introduced

concerns a simple edit taking place near the PAM site (Fig. 2), the actual *editing* efficiency is typically

less than 5% (Fig. 1). This low editing efficiency thus hinders multiplex genome editing to introduce

multiple edits simultaneously. Another limitation of prime editing are the restrictions in insertion as

well as deletion lengths. Therefore, future make-or-break prime editors would also benefit from

improvements in reverse transcriptases and pegRNA design driven by genetic screens, advances in

machine learning and/or structural insights[53,54]. Future optimized mbPEs with PAMless nucleases and

efficient reverse transcriptases with optimal pegRNA designs would enable multiplex editing of every

base in a genome, increase the efficiency of large genomic deletions, and facilitate the integration of

larger DNA segments.

While not yet perfect, the here-described mbPE system still represents one of the most powerful

and versatile bacterial genome editor currently available due to its scalability, and we foresee many

exciting and important applications, especially since the quality (and price) of oligo pools has

significantly improved in the last years. For instance, by designing a pool of pegRNAs, mbPE could be

employed to perform 'all codon scanning' on single genes of interest to gain new important insights

into protein function, protein interaction, or identify temperature-sensitive alleles. Another application

could be to examine gene essentiality at a genome-wide scale. Since mbPE can easily introduce

mutations without polar effects, large, genome-wide stop-codon or start-codon mutant libraries could

be made. Testing such libraries, with and without mbPE inducer under various environmental

conditions, followed by pegRNA sequencing and quantification of their abundance, would give precise

genome-wide fitness estimates in a single experiment. This could provide insights into gene essentiality

without the downsides of polar effects that methods such as Tn-Seq and CRISPRi-seq suffer from[25,55].

Alternatively, inducible promoters could be inserted by mbPE in front of every gene in a genome using

a pool of pegRNAs to test gene fitness in presence or absence of inducer. Another approach using

pooled pegRNAs would be to encode protein-tags such as the split-luc tag here described (Fig. 7A-C)

or any short immunogenic epitope (e.g. FLAG, HA, 6xHIS). pegRNAs could be designed to tag every

gene in a bacterial genome. This will, for instance, allow testing of protein-protein interactions at the

genome-wide scale or the preparation of bait pools in protein-purification strategies. To reduce the

number of false positives, wild type Cas9 or a second copy of the mbPE can be simultaneously

expressed with the prime editor to further enhance selection of correctly edited clones and reduce the

selection of mutants in the editor itself.

**Concluding Remarks**

The here-described mbPE system will facilitate genome-editing approaches in the significant human

pathogen *S. pneumoniae* and may contribute to the discovery of new insights with regards to

pneumococcal cell biology as well as lead to innovative therapeutic approaches. The described methods

and approaches may serve as a roadmap to develop mbPE for other bacteria.

# Methods

**Bacterial strains, culture conditions and transformation**

All strains, plasmids and primers used in this study are listed in Tables S2 and S3. Pneumococcal strains

in this study are derivate of *S. pneumoniae* D39V[30], unless specified otherwise, and are listed in Table

S2. Construction of strains is described in the Supplementary information. Strains were grown in liquid

semi-defined C+Y medium[55] (pH 6.8) or Columbia Agar Base (CAB) supplemented with 5% sheep

blood at 37°C. For transformation, pneumococci were grown in C+Y medium to an optical density

($OD_{595nm}$) of 0.1. Competence was activated by adding 100 ng/mL (CSP-1) (synthetic competence-

stimulating peptide 1) for 12 minutes at 37°C. The donor DNA was added to the culture and incubated

20 minutes at 30°C. The cultures were then diluted 10 times in C+Y and incubated at 37°C for 1.5

hours. Transformants were selected in CAB containing 5% blood and the appropriate antibiotic. Re-

streaked and confirmed clones were grown in C+Y to an $OD_{595nm}$ of 0.3 and stored at -80°C in C+Y

containing 17% glycerol.

*E. coli* Stbl3 (Thermofisher) was used for plasmid cloning. Strains were cultured in LB or LBA

with appropriate antibiotics.

**Cloning of pegRNAs by Golden Gate Assembly**

The pPEPZ_pegRNA vector was digested with Esp3I and purified from agarose gel to ensure removal

of the *mCherry* fragment. The two oligonucleotides designed for the spacer, the scaffold and the

extension sequences of each pegRNA were annealed in TEN buffer (10 mM Tris, 1 mM EDTA, 100 mM NaCl, pH 8) in a thermocycler at 95°C for 5 minutes and then slowly cooled down at 25°C. Equimolar ratios of each annealed pegRNA pieces were phosphorylated with T4 polynucleotide kinase (New England Biolabs). The annealed and phosphorylated pieces were ligated with the pPEPZ_pegRNA digested vector with T4 DNA ligase. The ligation products were transformed in *E. coli* Stbl3 and the transformants were selected on LBA plates with the appropriate antibiotic. Details about each pegRNA plasmid construction are indicated in Supplementary Information.

**Induction of PE2 and mbPE$^{S.pn}$**

Strains containing the PE2 nickase prime editor or the mbPE$^{S.pn}$ prime editor were grown in C+Y to an $OD_{595nm}$ of 0.3 ($3 \times 10^8$). Serial dilutions were prepared in C+Y medium and plated in CAB plates containing 5% blood and 50 ng/mL aTc and incubated overnight at 37°C. A control without aTc was included for the $1 \times 10^{-4}$ dilution to compare the colony number with and without inducer. Single colonies were picked and cultured in C+Y for further experiments.

**Growth assays and luciferase activity assays**

Single colonies from PE2 or mbPE induction plates were picked and pre-cultured in 200 µl of C+Y with no inducer and incubated at 37°C. After 2 hours of incubation, 50 µL of the colony suspension were transferred to white, clear bottom 96-well plates containing 250 µL of C+Y plus 0.12 mg/mL luciferin (final luciferin concentration, 0.1 mg/mL; BioSynth). For each assay D39V, uninduced colonies and media were included as controls in duplicates. Absorbance ($OD_{595}$) and luminescence (relative luminesce units, RLU) were measured for 20 hours with an interval time of 10 minutes, using a Tecan (Spark) plate reader. RLU/OD values were normalized against the baseline level of medium luminescence obtained for each experiment and data analyzed using GraphPad Prism.

**Nano-Glo Dual-Luciferase assay**

Strains were pre-cultured in C+Y to an $OD_{595}$ of 0.3. Luminescence (relative luminesce units, RLU) and Absorbance ($OD_{595}$) of sample triplicates supplemented with NanoGlo Live Cell Assay System (Promega) as recommended by the manufacturer, were monitored using a Tecan (Spark) plate reader during 15 minutes without intervals.

**pegRNA pool cloning**

The backbone for the cloning of the pegRNA pool (pVL4134) was linearized by XbaI digestion. The linearized vector was then amplified with primers OVL7743 and OVL7744 to introduce flanking Esp3I cleavage sites. Template plasmid was removed by digestion with DpnI. The ssDNA pegRNA pool (Table S1, Twist Bioscience) was amplified using primers OVL7966 and OVL6481 and the PCR

product was excised from a 2% agarose gel using the Monarch DNA cleanup and gel extraction kit

(NEB). One-step Golden Gate Assembly (GGA) was used to ligate the vector and the pegRNA pool

using Esp3I (molar ratio vector insert 1:10). The GGA product was transformed in *E. coli* Stbl3 and

transformants were selected on LBA plates containing 100 µg/mL spectinomycin. After overnight

incubation at 37°C, several single colonies were picked to check the cloning efficiency and the rest of

the colonies were scraped-off from the plates to prepare glycerol stocks with the suspension. The

plasmid pegRNA pool was purified and transformed into *S. pneumoniae* VL5155 and transformants

were plated on CAB plates supplemented with 5% blood and 100 µg/mL spectinomycin and incubated

overnight at 37°C. The cells were scraped-off the plates and the bacterial suspension was stored at -

80°C in C+Y containing 17% glycerol (pool name: VL5315).

**pegRNA pool screening**

To determine the best conditions for the pegRNA pool screening, a pilot mbPE induction experiment

was performed. 100 µL of VL5355 were resuspended in 4 mL of C+Y containing 100 µg/mL

spectinomycin, in triplicates. Starting $OD_{595}$ was 0.17 and the cells were grown to an $OD_{595}$ of 0.34.

Serial dilutions of the bacterial suspension (up to $10^{-5}$) were prepared in C+Y containing 100 µg/mL

spectinomycin and plated onto 75 mL CAB + 5% blood plates with and without 50 ng/mL aTc and

incubated overnight at 37°C. The CFU were counted, and it was determined that the selection efficiency

was about 10%. The dilutions to be used for the screening were chosen to be $10^{-2}$ for the inducer

condition and $10^{-3}$ for the no inducer condition (single colonies could be still observed and the plates

had a good coverage). The pegRNA pool screening was performed in quadruplicates with the same

conditions previously tested in the pilot experiment. All colonies grown with and without inducer were

scraped-off using C+Y + spectinomycin and stored with 17% glycerol at -80°C. Chromosomal DNA

from the pegRNA pool screening samples was extracted for Illumina library preparation using the

Promega kit as described previously[38].

**Targeted *luc* modification site sequencing and differential enrichment analysis**

Custom oligonucleotide primers (Table S3) consisting of P5 or P7 Illumina sequencing adapters,

Illumina sequencing barcodes (N501-504/N701-704), read 1 or read 2 sequencing primer binding sites,

and portions binding to the chromosomally located *luc* were used to amplify the targeted *luc*

modification site using Phanta high-fidelity polymerase (Vazyme). Amplified fragments were purified

from gel and sequenced on an Illumina MiSeq, using a 2 x 300 (600 cycles) paired-end v3 sequencing

kit. Illumina sequencing reads were checked for quality using FastQC (v0.11.8, available from

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and paired reads were merged using

PEAR (v0.9.6, available from https://www.h-its.org/software/pear-paired-end-read-merger/). PEAR

merged reads were subsequently used to count the number of occurrences of each pegRNA-encoded mutation, using 2FAST2Q (v2.5.3)[37] in counter mode with consideration for the minimal Phred quality score (all set to 20), and the mismatches allowed in upstream, downstream, and queried sequences (all 0). Counts from 2FAST2Q were imported into R, and normalized using DESeq2 (v1.40.2). DESeq2[56] was subsequently used to perform the differential enrichment analysis between the control and aTc induced samples. Statistical significance for differential presence was defined as an absolute log2FoldChange > 1, and an adjusted p-value < 0.05. Full results of the *luc* mutation analysis can be found in Table S4. Note that because of the high number of unedited, wild type *luc* alleles present in the amplified pools, we were unable to accurately quantify the substitution identities and their editing efficiencies for single base pair substitutions. Instead, we quantified the abundance of the pegRNAs in presence and absence of aTc as a proxy for editing efficiency of single base pair substitutions.

**Sequencing of pegRNA and differential enrichment analysis**

Custom oligonucleotide primers (Table S3) consisting of P5 or P7 Illumina sequencing adapters, Illumina sequencing barcodes (N501-504/N701-704), read 1 or read 2 sequencing primer binding sites, and portions binding to the chromosomally integrated pegRNA expression were used to amplify the pegRNAs using Phanta polymerase. Amplified fragments were purified and sequenced using an Illumina MiniSeq instrument, with a 2 x 150 (300 cycles) paired-end mid output sequencing kit. Raw Illumina sequencing reads were checked for quality using FastQC (v0.11.8, available from https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and paired reads were merged using PEAR (v0.9.6, available from https://www.h-its.org/software/pear-paired-end-read-merger/). PEAR merged reads were subsequently used to count the number of occurrences of each pegRNA-encoded mutation, using 2FAST2Q (v2.5.3)[37] in counter mode with consideration for the minimal Phred quality score (all set to 20), and the mismatches allowed in upstream, downstream, and queried sequences (all 0). Counts from 2FAST2Q were imported into R, and normalized using DESeq2 (v1.40.2). DESeq2[56] was subsequently used to perform the differential enrichment analysis between the control and aTc induced samples. Statistical significance for differential presence was defined as an absolute log2FoldChange > 1, and an adjusted p-value < 0.05. Full results of the pegRNA abundance analysis can be found in Table S5.

**Quantifications and statistical analysis**

548 Data analysis was performed using R and Prism (Graphpad). Data shown are represented as mean of at
549 least three replicates ± SEM if data came from one experiment with replicated measurement, and ± SD
550 if data came from separate experiments.

### Genome sequencing and analysis

Genome sequencing was performed using PacBio HiFi sequencing on a Pacbio sequel II machine.
PacBio HiFi reads were assembled into circular genomes using Flye[57]. Flye-produced assemblies were
rotated for *dnaA* using circlator[58]. Assemblies were assessed for structural rearrangements and large
indels through progressiveMauve[59], by mapping them to an *in silico* constructed reference genome
containing the known mutations and insertions. For SNPs and small indel calling, the raw reads from
the PacBio sequencing were mapped to the synthetic genomes using minimap2[60], and filtered for
secondary and split alignments. Resulting alignments were sorted using SAMtools, and mutations were
called using bcftools. Coverage at each position was assessed using SAMtools[61].

### Data availability

563 Sequencing data from sequencing reactions of the luciferase gene and the pegRNAs, and the PacBio
564 sequencing, are available in the Sequence Read Archive under BioProject accession code
565 PRJNA1062582.

# Acknowledgements
Work in the Veening lab is supported by the Swiss National Science Foundation (SNSF) (project grants
310030_192517, 310030_200792 and 'AntiResist' 51NF40_180541) and ERC consolidator grant
771534-PneumoCaTChER. A.B.J. is supported through a Postdoctoral Fellowship grant
(TMPFP3_210202) from the SNSF. X.L. was supported by Shenzhen University 2035 Program for
Excellent Research (86901-00000216). The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.

# Authors Contributions
576 J.W.V. wrote the paper with input from all authors. M.R.G.., M.V.M. A.B.J. and A.S.R. performed the
577 experiments. M.G.R., A.B.J., X.L., M.V.M., A.S.R. and J.W.V designed, analyzed and interpreted the
578 data. All authors reviewed and approved the final version of the manuscript.

## Transparency declarations

M.R.G.: None

M.V.M.: None

A.B.J. : None

A.S.R. : None

X.L. : None

J.W.V.: None

## ORCID and email addresses

| Name | Email address | ORCID ID |
|---|---|---|
| Monica Rengifo-Gonzalez | Monica.Rengifo@unil.ch | |
| Maria Vittoria Mazzuoli | MariaVittoria.Mazzuoli@unil.ch | 0000-0002-7071-1698 |
| Axel B. Janssen | Absalom.Janssen@unil.ch | 0000-0002-9865-447X |
| Anne-Stephanie Rueff | Anne-Stephanie.Rueff@unil.ch | |
| Xue Liu | XueLiu@szu.edu.cn | 0000-0001-6485-1865 |
| Jan-Willem Veening | Jan-Willem.Veening@unil.ch | 0000-0002-3162-6634 |

## References

1. Ramachandran, G. & Bikard, D. Editing the microbiome the CRISPR way. *Philos. Trans. R. Soc.*
*B Biol. Sci.* **374**, 20180103 (2019).

2. Shapiro, R. S., Chavez, A. & Collins, J. J. CRISPR-based genomic tools for the manipulation of
genetically intractable microorganisms. *Nat. Rev. Microbiol.* **16**, 333–339 (2018).

3. Penewit, K. *et al.* Efficient and Scalable Precision Genome Editing in Staphylococcus aureus
through Conditional Recombineering and CRISPR/Cas9-Mediated Counterselection. *mBio* **9**,
e00067-18 (2018).

4. McAllister, K. N. & Sorg, J. A. CRISPR Genome Editing Systems in the Genus Clostridium: a
Timely Advancement. *J. Bacteriol.* **201**, e00219-19 (2019).

5. Tong, Y. *et al.* CRISPR-Cas9, CRISPRi and CRISPR-BEST-mediated genetic manipulation in
streptomycetes. *Nat. Protoc.* **15**, 2470–2502 (2020).

6. Jinek, M. *et al.* A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821 (2012).

7. Anzalone, A. V., Koblan, L. W. & Liu, D. R. Genome editing with CRISPR-Cas nucleases, base editors, transposases and prime editors. *Nat. Biotechnol.* **38**, 824–844 (2020).

8. Finger-Bou, M., Orsi, E., van der Oost, J. & Staals, R. H. J. CRISPR with a Happy Ending: Non-Templated DNA Repair for Prokaryotic Genome Engineering. *Biotechnol. J.* **15**, 1900404 (2020).

9. Shuman, S. & Glickman, M. S. Bacterial DNA repair by non-homologous end joining. *Nat. Rev. Microbiol.* **5**, 852–861 (2007).

10. Jiang, W. & Marraffini, L. A. CRISPR-Cas: New Tools for Genetic Manipulations from Bacterial Immunity Systems. *Annu. Rev. Microbiol.* **69**, 209–228 (2015).

11. Anzalone, A. V. *et al.* Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature* **576**, 149–157 (2019).

12. Scholefield, J. & Harrison, P. T. Prime editing - an update on the field. *Gene Ther.* **28**, 396–401 (2021).

13. Hua, K., Han, P. & Zhu, J.-K. Improvement of base editors and prime editors advances precision genome engineering in plants. *Plant Physiol.* **188**, 1795–1810 (2022).

14. Villiger, L. *et al.* CRISPR technologies for genome, epigenome and transcriptome editing. *Nat. Rev. Mol. Cell Biol.* (2024) doi:10.1038/s41580-023-00697-6.

15. Zhao, Z., Shang, P., Mohanraju, P. & Geijsen, N. Prime editing: advances and therapeutic applications. *Trends Biotechnol.* **41**, 1000–1012 (2023).

16. Tong, Y., Jørgensen, T. S., Whitford, C. M., Weber, T. & Lee, S. Y. A versatile genetic engineering toolkit for E. coli based on CRISPR-prime editing. *Nat. Commun.* **12**, 5206 (2021).

17. Zhang, H. *et al.* BacPE: a versatile prime-editing platform in bacteria by inhibiting DNA exonucleases. *Nat. Commun.* **15**, 825 (2024).

18. Bikard, D. *et al.* Programmable repression and activation of bacterial gene expression using an engineered CRISPR-Cas system. *Nucleic Acids Res.* **41**, 7429–7437 (2013).

19. Jiang, W., Bikard, D., Cox, D., Zhang, F. & Marraffini, L. A. RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat. Biotechnol.* **31**, 233–239 (2013).

20. Synefiaridou, D. & Veening, J.-W. Harnessing CRISPR-Cas9 for Genome Editing in Streptococcus pneumoniae D39V. *Appl. Environ. Microbiol.* **87**, e02762-20 (2021).

21. Liu, X. *et al.* Exploration of Bacterial Bottlenecks and Streptococcus pneumoniae Pathogenesis by CRISPRi-Seq. *Cell Host Microbe* **29**, 107-120.e6 (2021).

22. Liu, X. *et al.* High-throughput CRISPRi phenotyping identifies new essential genes in Streptococcus pneumoniae. *Mol. Syst. Biol.* **13**, 931 (2017).

23. Sorg, R. A., Kuipers, O. P. & Veening, J.-W. Gene expression platform for synthetic biology in the human pathogen Streptococcus pneumoniae. *ACS Synth. Biol.* **4**, 228–239 (2015).

24. van Gestel, J., Hawkins, J. S., Todor, H. & Gross, C. A. Computational pipeline for designing guide RNAs for mismatch-CRISPRi. *STAR Protoc.* **2**, 100521 (2021).

25. de Bakker, V., Liu, X., Bravo, A. M. & Veening, J.-W. CRISPRi-seq for genome-wide fitness quantification in bacteria. *Nat. Protoc.* **17**, 252–281 (2022).

26. Xie, K., Minkenberg, B. & Yang, Y. Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 3570–3575 (2015).

27. Hsu, J. Y. *et al.* PrimeDesign software for rapid and simplified design of prime editing guide RNAs. *Nat. Commun.* **12**, 1034 (2021).

28. Nelson, J. W. *et al.* Engineered pegRNAs improve prime editing efficiency. *Nat. Biotechnol.* **40**, 402–410 (2022).

29. Chen, P. J. *et al.* Enhanced prime editing systems by manipulating cellular determinants of editing outcomes. *Cell* **184**, 5635-5652.e29 (2021).

30. Ferreira da Silva, J. *et al.* Prime editing efficiency and fidelity are enhanced in the absence of mismatch repair. *Nat. Commun.* **13**, 760 (2022).

31. Slager, J., Aprianto, R. & Veening, J.-W. Deep genome annotation of the opportunistic human pathogen Streptococcus pneumoniae D39. *Nucleic Acids Res.* **46**, 9971–9989 (2018).

32. Martin, B., Prats, H. & Claverys, J. P. Cloning of the hexA mismatch-repair gene of Streptococcus pneumoniae and identification of the product. *Gene* **34**, 293–303 (1985).

33. Tao, R. *et al.* WT-PE: Prime editing with nuclease wild-type Cas9 enables versatile large-scale genome editing. *Signal Transduct. Target. Ther.* **7**, 108 (2022).

34. Peterka, M. *et al.* Harnessing DSB repair to promote efficient homology-dependent and -independent prime editing. *Nat. Commun.* **13**, 1240 (2022).

35. Adikusuma, F. *et al.* Optimized nickase- and nuclease-based prime editing in human and mouse cells. *Nucleic Acids Res.* **49**, 10785–10795 (2021).

36. Keller, L. E., Rueff, A.-S., Kurushima, J. & Veening, J.-W. Three New Integration Vectors and Fluorescent Proteins for Use in the Opportunistic Human Pathogen Streptococcus pneumoniae. *Genes* **10**, (2019).

37. Bravo, A. M., Typas, A. & Veening, J.-W. 2FAST2Q: a general-purpose sequence search and counting program for FASTQ files. *PeerJ* **10**, e14041 (2022).

38. Gallay, C. *et al.* CcrZ is a pneumococcal spatiotemporal cell cycle regulator that interacts with FtsZ and controls DNA replication by modulating the activity of DnaA. *Nat. Microbiol.* **6**, 1175–1187 (2021).

39. Oliveira Paiva, A. M. *et al.* The Bacterial Chromatin Protein HupA Can Remodel DNA and Associates with the Nucleoid in Clostridium difficile. *J. Mol. Biol.* **431**, 653–672 (2019).

40. Perez, A. J. *et al.* Elongasome core proteins and class A PBP1a display zonal, processive movement at the midcell of Streptococcus pneumoniae. *BioRxiv Prepr. Serv. Biol.* 2024.01.10.575112 (2024) doi:10.1101/2024.01.10.575112.

41. Gibson, P. S. & Veening, J.-W. Gaps in the wall: understanding cell wall biology to tackle amoxicillin resistance in Streptococcus pneumoniae. *Curr. Opin. Microbiol.* **72**, 102261 (2023).

42. Straume, D., Piechowiak, K. W., Kjos, M. & Håvarstein, L. S. Class A PBPs: It is time to rethink traditional paradigms. *Mol. Microbiol.* **116**, 41–52 (2021).

43. Lenoir, C. *et al.* The morphogenic protein CopD controls the spatio-temporal dynamics of PBP1a and PBP2b in Streptococcus pneumoniae. *mBio* **14**, e0141123 (2023).

44. Li, X. *et al.* Development of a versatile nuclease prime editor with upgraded precision. *Nat. Commun.* **14**, 305 (2023).

45. Jiang, T., Zhang, X.-O., Weng, Z. & Xue, W. Deletion and replacement of long genomic sequences using prime editing. *Nat. Biotechnol.* **40**, 227–234 (2022).

46. Kweon, J. *et al.* Targeted genomic translocations and inversions generated using a paired prime editing strategy. *Mol. Ther. J. Am. Soc. Gene Ther.* **31**, 249–259 (2023).

47. Zeng, H. *et al.* Recent advances in prime editing technologies and their promises for therapeutic applications. *Curr. Opin. Biotechnol.* **86**, 103071 (2024).

48. Gibson, P. S., Bexkens, E., Zuber, S., Cowley, L. A. & Veening, J.-W. The acquisition of clinically relevant amoxicillin resistance in Streptococcus pneumoniae requires ordered horizontal gene transfer of four loci. *PLoS Pathog.* **18**, e1010727 (2022).

49. Hiller, N. L. & Orihuela, C. J. Biological puzzles solved by using Streptococcus pneumoniae: a historical review of the pneumococcal studies that have impacted medicine and shaped molecular bacteriology. *J. Bacteriol.* **0**, e00059-24 (2024).

50. Collias, D. & Beisel, C. L. CRISPR technologies and the search for the PAM-free nuclease. *Nat. Commun.* **12**, 555 (2021).

51. Walton, R. T., Christie, K. A., Whittaker, M. N. & Kleinstiver, B. P. Unconstrained genome targeting with near-PAMless engineered CRISPR-Cas9 variants. *Science* **368**, 290–296 (2020).

52. Zhao, L. *et al.* PAM-flexible genome editing with an engineered chimeric Cas9. *Nat. Commun.* **14**, 6175 (2023).

53. Shuto, Y. *et al.* Structural basis for pegRNA-guided reverse transcription by a prime editor. *Nature* (2024) doi:10.1038/s41586-024-07497-8.

54. Chen, P. J. & Liu, D. R. Prime editing for precise and highly versatile genome manipulation. *Nat. Rev. Genet.* **24**, 161–177 (2023).

55. Cain, A. K. *et al.* A decade of advances in transposon-insertion sequencing. *Nat. Rev. Genet.* **21**, 526–540 (2020).

56. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

57. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using
repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

58. Hunt, M. *et al.* Circlator: automated circularization of genome assemblies using long sequencing
reads. *Genome Biol.* **16**, 294 (2015).

59. Darling, A. E., Mau, B. & Perna, N. T. progressiveMauve: multiple genome alignment with gene
gain, loss and rearrangement. *PloS One* **5**, e11147 (2010).

60. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinforma. Oxf. Engl.* **34**, 3094–
3100 (2018).

61. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**, 2078–
2079 (2009).