Title page

Title

IGF2BP3 promotes the progression of gastric cancer by activating cGMP-PKG signaling pathway via targeting FBXO32

Authors' names

Yi Si[1#], Bo Tian[1#], Rui Zhang[1], Mingda Xuan[1], Kunyi Liu[1], Jiao Jiao[1], Shuangshuang Han[1], Hongfei Li[1], Yanhong Hu[1], Hongyan Zhao[1], Wenjing He[2], Jia Wang[3*], Ting Liu[1*], Weifang Yu[1*]

[1] Department of Endoscopy Center, The First Hospital of Hebei Medical University, Shijiazhuang, China

[2] Department of Endoscopy Center, Kuancheng manchu autonomous county hospital, Chengde, China

[3] Department of Infectious Diseases, The First Hospital of Hebei Medical University, Shijiazhuang, China

# These authors contributed equally

*The corresponding authors.

Running title

IGF2BP3 promotes GC by regulating FBXO32 and cGMP-PKG signaling pathway

Corresponding author: Weifang Yu, M.D., Professor, No. 89 Donggang Road, Shijiazhuang, Hebei, China. E-mail: yuweifang@hebmu.edu.cn. Ting Liu, No. 89 Donggang Road, Shijiazhuang, Hebei, China. E-mail: 59103696@hebmu.edu.cn. Jia

Wang, Professor, No. 89 Donggang Road, Shijiazhuang, Hebei, China. E-mail:

57101040@hebmu.edu.cn

Keywords: Gastric cancer; N6-methyladenosine (m^{6}A); IGF2BP3; FBXO32; cGMP-

PKG

Abstract

N6-methyladenosine (m⁶A) represents the most prevalent chemical modification on eukaryotic mRNA, with an accumulating body of literature indicating its pivotal significance in the pathogenesis of human cancers. Nevertheless, the precise molecular interplay between the m⁶A reader protein IGF2BP3 and gastric cancer remains to be thoroughly delineated. Our study uncovered that the expression of IGF2BP3 in gastric cancer tissues is markedly elevated in comparison to adjacent normal tissues, and this upregulation is tightly correlated with the incidence of lymph node metastasis, more advanced TNM stages, and deeper invasion depth of tumor in patients. *In vitro* experiments demonstrated that IGF2BP3 potentiates the proliferative, migratory, and invasive capacities of gastric cancer cells, while concurrently inhibiting apoptosis and augmenting the intracellular levels of aerobic glycolysis. *In vivo* experiments revealed that IGF2BP3 contributes to the growth of gastric cancer. Mechanistically, IGF2BP3 can increase the expression of FBXO32 protein by recognizing and binding to the m⁶A binding site on FBXO32 mRNA and further activate the downstream cGMP-PKG signaling pathway, thereby modulating various biological functions of gastric cancer cells and ultimately promoting the progression of gastric cancer. In summary, our findings suggest that IGF2BP3 upregulates the expression of FBXO32 protein in an m⁶A dependent manner and subsequently activates the cGMP-PKG signaling pathway, ultimately leading to the onset and progression of gastric cancer. Consequently, the targeting of the IGF2BP3/FBXO32/cGMP-PKG axis emerges as a promising therapeutic modality for the treatment of gastric cancer.

Introduction

Gastric cancer (GC) is a heterogeneous disease characterized by significant differences in epidemiology and histopathology across countries and is one of the leading causes of cancer-related deaths[1]. The morbidity and mortality rates of GC are high worldwide and particularly in China, posing a serious threat to human health[2]. Typically, there are no obvious clinical symptoms in the early stage of GC, and most patients are diagnosed at an advanced stage. The recurrence rate of advanced GC is high, leading to a poor overall prognosis for GC patients[3]. To date, the specific molecular mechanisms of GC have not been fully elucidated. This may be attributed to its complex etiology, which primarily involves the interactions between oncogenes and tumor suppressor genes[4], as well as genetic, epigenetic, and environmental factors[5]. Therefore, further exploration of the molecular mechanisms underlying the occurrence and development of GC will provide potential new targets for the clinical treatment of GC, which is of significant clinical importance.

In recent years, the role of epigenetics in the pathogenesis of GC has attracted extensive attention from scholars worldwide. Epigenetics is generally defined as the study of chemical modifications to DNA and RNA, which mainly include DNA and RNA methylation, histone modification, non-coding RNA modification, and chromatin rearrangement[6]. N6-methyladenosine (m6A) modification is an integral part of epigenetics research. m6A modification is the most common internal modification of eukaryotic mRNA, regulating the stability, translation efficiency, alternative splicing and localization of RNA at the post-transcriptional level. It participates in various

biological processes such as initiation, proliferation, differentiation, metastasis and metabolic reprogramming of tumor cells[7, 8]. m6A related proteins are mainly divided into methyltransferase, demethylase and reader proteins, involving 21 related genes[9]. By analyzing the expression of m6A related proteins in GC and adjacent normal tissues in the TCGA and GEO (GSE54129 and GSE66229 data sets) databases, we screened out four 'readers', YTHDF1, YTHDF2, IGF2BP1 and IGF2BP3, whose expression levels were higher in cancer tissues than in adjacent normal tissues in the above databases. Among these, the expression difference of IGF2BP3 was the most significant, and there are few studies about IGF2BP3 and GC, so IGF2BP3 was selected as the research target. It is reported that the main function of m6A 'readers' is to interpret RNA methylation modification information and regulate the translation and degradation of its downstream RNA[10, 11].

Insulin like growth factor II mRNA binding protein 3 (IGF2BP3), an m6A 'reader', is located on human chromosome 7p15.3 and contains two RNA recognition motifs and four K homologous domains[12]. Overexpression of IGF2BP3 was first observed in pancreatic cancer[13]. Its primary role is to recognize m6A binding sites through the K homology domain, thereby increasing the stability and translation efficiency of target mRNAs[14]. IGF2BP3 is implicated in the development of various cancers, playing distinct roles in each, such as, IGF2BP3 enhances the stability of TMBIM6 mRNA by identifying the m6A site, thus promoting the progression of laryngeal squamous cell carcinoma[15]; Similarly, IGF2BP3 stabilizes lnc-CTHCC by recognizing its m6A site, which increases lnc-CTHCC expression in hepatocellular carcinoma, ultimately

facilitating the disease's progression[16]; In colorectal cancer, IGF2BP3 stabilizes CCND1 mRNA by identifying its m^6A site, promoting CCND1 protein expression, regulating the cell cycle, and thus promoting the proliferation and growth of cancer cells[17]; Furthermore, Wang et al. found that m^6A 'reader' IGF2BP2 enhances the stability of GLUT1 mRNA, a key gene in aerobic glycolysis, in an m^6A dependent manner, significantly promoting the aerobic glycolysis of colorectal cancer cells[18]. Therefore, IGF2BP3 may be a critical oncogene. We observed that IGF2BP3 was highly expressed in GC and was closely related to poor prognosis in GC patients. IGF2BP3 reinforces the malignant phenotype of GC cells and promotes the growth of GC *in vivo*. Through omics analysis and various experiments, we identified for the first time that the downstream gene with m^6A modification of IGF2BP3 in GC is FBXO32, which is also highly expressed in GC and interacts with IGF2BP3. IGF2BP3 upregulates the expression of FBXO32 protein in an m^6A dependent manner and activates the cGMP-PKG signaling pathway, which further enhances the proliferation, migration, and invasion capabilities of GC cells, and ultimately promotes the progression of GC.

Materials and Methods

Patients and samples

63 pairs of GC tissues and adjacent normal tissues were obtained from the First Hospital of Hebei Medical University. None of the patients had received radiation or chemotherapy prior to surgery and they had signed informed consent forms. The above tissue samples were promptly placed in liquid nitrogen after isolation, and then

transferred to the Biological Sample Bank of the First Hospital of Hebei Medical University for storage. The cancer tissues used in this study were pathologically confirmed as gastric adenocarcinoma/signet-ring cell carcinoma. This study was approved by the Ethics Committee of the First Hospital of Hebei Medical University (Approval letter No. S00180) and adheres to the principles of the Declaration of Helsinki.

Cell culture

Human gastric mucosa cell (GES-1) and GC cells (HGC-27, MKN7, MKN74) were obtained from Wuhan Pricella Biotechnology Co., Ltd., and were verified by STR analysis. The cells were cultured in RPMI 1640 medium (Gibco, Gaithersburg, MD, USA) supplemented with 10% fetal bovine serum (FBS; Gibco) and 1% penicillin-streptomycin solution (Solarbio Sciences & Technology Co. Ltd, Beijing, China). They were maintained in a constant temperature and humidity incubator with 5% carbon dioxide at 37℃, and routine mycoplasma detection was performed monthly using PCR. In all experiments, the cells were used for a maximum of no more than 25 passages.

Plasmids, siRNAs, shRNAs, Transfections and Stable cell lines

Transfection was performed according to Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) when the cell confluence reached 80-90% in a cell culture flask or dish. IGF2BP3 siRNA and negative control (GenePharma Co., Ltd., Shanghai, China) were transfected into HGC-27 and MKN74 cells, while an IGF2BP3 overexpression plasmid and a negative control (GenePharma) were transfected into MKN7 and GES-1 cells. Additionally, IGF2BP3 shRNA and a negative control (GenePharma) were transfected

into HGC-27 cells. FBXO32 siRNA and a negative control (GenePharma) were

transfected into MKN7 cells, and an FBXO32 overexpression plasmid and a negative

control (GenePharma) were transfected into HGC-27 cells. Follow-up experiments

were conducted 48 hours after transfection. To eatablish stable transfected cell lines

(HGC-27 shNC, HGC-27 shIGF2BP3), the cell culture medium was changed to include

neomycin at a final concentration of 600 μg/mL (without other antibiotics) 48 hours

after transfection with IGF2BP3 shRNA in HGC-27 cells. Once the clones have formed,

the neomycin concentration was adjusted to 300 μg/mL, continue culturing until day

10.

RNA extraction and qRT-PCR

Total RNA was extracted from cells and tissues using RNA-Easy Isolation Reagent

(Vazyme Biotech Co.,Ltd., Nanjing, China). The RNA was then reverse-transcribed

into complementary DNA (cDNA) following the instructions of PrimeScript RT reagent

kit (Takara, Beijing, China). qRT-PCR was performed using the AceQ Universal SYBR

qPCR Master Mix (Vazyme), with β-actin serving as the endogenous control mRNA.

Each sample was tested in triplicate. The mRNA expression level was determined based

on the CT (cyclic threshold) value of each sample, and the relative expression level of

each sample was calculated using the $2^{-\Delta\Delta CT}$ method. The sequence of primers are

detailed in Supplementary Table 1.

Immunohistochemistry (IHC)

Tissues from patients and nude mice were fixed with 4% paraformaldehyde, followed

by paraffin embedding (Shanghai YiYang Instrument Co., Ltd., Shanghai, China),

sectioning, and staining. The Rabbit/Mouse two-step detection kit (Beijing ZSBG-Bio

Co., Ltd., Beijing, China) was used for immunohistochemical analysis according to the

manufacturer's instructions. Anti-IGF2BP3 (diluted 1:150; Proteintech, Wuhan, Hubei,

P.R.C; Catalog number: 14642-1-AP), Anti-Ki67 (diluted 1:100; Proteintech; Catalog

number: 27309-1-AP) and Anti-FBXO32 (diluted 1:100; Proteintech; Catalog number:

67172-1-Ig) antibodies were used for immunohistochemical staining. The staining

results were analyzed by using a double-blind scoring method and the average optical

density (AOD) method.

Western blot analysis and antibodies

Protein was extracted using RIPA lysis buffer (Solarbio) and the protease/phosphatase

inhibitor (Solarbio) at a ratio of 100:1. The proteins were separated electrophoretically

on a 10% SDS polyacrylamide gel (Bio-Rad Laboratories Inc, Hercules, CA, USA) and

transferred to a polyvinylidene fluoride membrane (Merck Millipore, Billerica, MA,

USA). The following antibodies were used in this study: IGF2BP3 antibody (diluted

1:1000; Abcam, Cambridge, MA, USA; Catalog number: ab177477), β-actin antibody

(diluted 1:1500; ZSBG-Bio; Catalog number: TA-09), FBXO32 antibody (diluted

1:500; Abcam; Catalog number: ab168372) and GAPDH antibody (diluted 1:500;

Goodhere Co. Ltd, Hangzhou, China; Catalog number: AB-P-R 001), PRKG1 (PKG1)

Polyclonal antibody (diluted 1:100; Proteintech; Catalog number: 21646-1-AP),

Phospho-VASP (Ser239) Antibody (diluted 1:100; Affinity; Catalog number: AF3338),

VASP Polyclonal antibody (diluted 1:100; Proteintech; Catalog number: 13472-1-AP).

The immunoreactive protein bands were detected by the Odyssey Scanning System (LI-

COR Biosciences, Lincoln, NE, USA).

Cell proliferation assay

For CCK-8 assay, the cells of each group were uniformly distributed in 96-well plates with a density of 1×10^3 cells per well, and cultured with complete culture medium. At 24, 48, 72, and 96 hours after distribution, 10 μL of Cell Counting Kit -8 (CCK-8; Dojindo, Tokyo, Japan) reagent was added to each well. Then, the cells were incubated in the incubator for an additional 2 hours. The absorbance of each well at 450 nm was measured using a Promega GloMax luminescence detector (Promega, Madison, WI, USA).

For the colony formation assay, cells of each group were uniformly distributed in 6-well plates at a density of 500 cells per well. Fresh culture medium was replaced every 3 days. Cells were cultured for 7-14 days, then washed twice with PBS, treated with 4% paraformaldehyde for 30 minutes, and stained with 0.1% crystal violet for 20 minutes.

Cell migration assay

For the wound healing assay, the cells of each group were uniformly distributed in 6-well plates. When the cell confluence reached or was close to 100%, a 200 μL pipette tip was used to draw two straight lines on each well to simulate the wound. The cells were then washed twice with PBS and cultured with a culture medium without FBS. Each well was photographed at 0 hours and 48 hours after the wound was created, and the migration rate was expressed as the ratio of gap width measured at 48 hours to the gap width measured at 0 hour.

For the Transwell migration assay, 3×10^4 (HGC-27, GES-1) /1×10^5 (MKN7, MKN74)

cells were added to the upper chamber (Corning Incorporated, Corning, NY, USA) in

200 μL FBS-free medium. To induce downward migration, 700 μL of complete medium

was added to the lower chamber. After 48 hours of culture, cells on the upper chamber

side of the polycarbonate membrane were removed using a swab. The chamber was

cleaned twice with PBS, then treated with 4% paraformaldehyde for 30 minutes,

followed by 0.1% crystal violet for 20 minutes. The excess crystal violet was cleaned

again with PBS. Five visual fields were randomly selected for photography, and the

stained cells were counted using ImageJ software (National Institutes of Health,

Bethesda, MD, USA).

Cell invasion assay

For the Transwell invasion assay, cells were resuspended in 100 μL FBS-free medium

and added to the upper chamber, while 500 μL of complete medium was added to the

lower chamber to induce downward invasion of the cells in the upper chamber. All other

experimental procedures were the same as those for the Transwell migration assay.

Cell apoptosis assay

Cells from each group were collected 48 hours after transfection and stained using the

Annexin V-FITC/PI apoptosis assay kit (NeoBioscience, Shenzhen, China). Flow

cytometry (BD Biosciences, San Jose, CA, USA) was used to analyze the apoptosis of

1×10^5 cells, and FlowJo software was used to calculate the percentage of apoptotic

cells.

Cell glucose metabolism assay

MKN7/HGC-27 cells that overexpressed or have knockdown of IGF2BP3 were

collected and processed according to the kit instructions (Glucose Assay kit, Abcam,

ab65333; L- Lactate Assay kit, Abcam, ab65331; ATP Assay Kit, Abcam, ab83355). In

brief, the standard solutions of glucose, lactic acid and ATP were prepared in a gradient

dilution. The reaction mixtures were then prepared, and the diluted standard solutions

were mixed with the reaction mixtures for incubation (37℃ or room temperature) for

30 minutes. The spectrophotometer was adjusted to the corresponding wavelength to

read the absorbance value, and the standard curve for glucose, lactic acid and ATP were

generated. The corresponding reagents from the kits were added to the lysed cells, and

the cell lysate was deproteinized using Deproteinizing Sample Preparation Kit - TCA

(Abcam; Catalog number: ab204708). The deproteinized cell lysates were then mixed

with the reaction mixtures for incubation (37℃ or room temperature) for 30 minutes,

and the absorbance values were compated to the standard curves to calculate the

concentrations of glucose, lactic acid and ATP.

Animal studies

Ten male BALB/c nude mice aged 4-6 weeks, 16-20 g (procured from Beijing

Huafukang Biotechnology Co., LTD.) were randomly divided into two groups (five

mice per group) and housed in a pathogen-free environment with sufficient water and

food. The room temperature was maintained at approximately 22℃ and a 12/12-hour

light/dark cycle. In a clean bench, the skin of the left hind limb flank of the nude mice

was disinfected, and HGC-27 cells (5×10^6 cells per mice) stably transfected with

shRNA-NC and shRNA-IGF2BP3 were subcutaneously injected using a sterile syringe.

Starting from day 3 post-injection, tumor size was evaluated every two days using the

formula: volume = (long diameter × short diameter ^{2})/2. When the tumor volume

approached 1000 mm^3, the nude mice were euthanized humanely. Tumor specimens

were collected, photographed, weighed, and fixed in formalin. The tumor samples were

then paraffin-embedded and sectioned for subsequent HE and IHC analysis. The above

experiments were approved by the Ethics Committee of the First Hospital of Hebei

Medical University and carried out in accordance with the experimental animal care

and use system.

RNA sequencing (RNA-seq)

Total RNA was extracted from HGC-27 cells transfected with siRNA-NC and siRNA-

IGF2BP3 for 24 hours. The RNA library construction and sequencing analysis were

entrusted to Beijing Novogene Technology Co., LTD. In short, a total of six samples

(siNC × 3, siIGF2BP3 × 3) were tested using Illumina's NovaSeq 6000 platform.

Quality control and filtering are carried out on the output data from each sample to

obtain high-quality data. HISAT software was used to align high-quality data to

reference genomes and to quantify gene or transcript expression levels in the sample.

DESeq2 software was used for differential expression analysis, with differential genes

defined as those with log$_2$|Fold Change| > 1.0 and $P < 0.05$ were selected.

Methylated RNA immunoprecipitation sequencing (MeRIP-seq)

264 Total RNA was extracted from HGC-27 cells transfected with siRNA-NC and siRNA-

265 IGF2BP3 for 24 hours. The RNA library construction and sequencing analysis were

266 entrusted to Novogene. In short, the original data was processed using fastp (version

267 0.19.11) to obtain high quality data. The reference genome index was constructed using

BWA (version 0.7.12), and the high-quality data was aligned with the reference genome

using BWA mem (version 0.7.12). After mapping reads to the reference genome, the

m^6A peaks within each group were identified using the exomePeak R package (version

2.16.0). The genes corresponding to each peak were presumed to be peak-related genes.

The m^6A enrichment motif of each group was identified by HOMER (version 4.9.1).

The peak distribution in the 5' UTR, CDS, 3' UTR and other functional regions on

mRNA transcripts was statistically analyzed.

RNA immunoprecipitation and high-throughput sequencing assay (RIP-seq)

According to the instructions of the RNA Immunoprecipitation (RIP) Kit（BersinBio,

Guangzhou, China; Catalog number: Bes5101）, HGC-27 cells transfected with siRNA-

NC and siRNA-IGF2BP3 for 48 hours were collected and lysed. An IGF2BP3 specific

antibody (10 μg; Proteintech; Catalog number: 14642-1-AP) was employed to bind the

endogenous expression of IGF2BP3 protein in the cells, precipitate the target protein-

RNA complex, and subsequently isolate and purify the RNA within the complex.

Novogene was commissioned to perform ribosomal RNA removal, RNA library

construction and high-throughput sequencing analysis. In brief, RNA fragment

distribution and concentration after immunoprecipitation and ribosomal RNA removal

were measured using an Agilent 2100 bioanalyzer (Agilent) and a simpliNano

spectrophotometer (GE Healthcare). The library was constructed using the NEB

Next®Ultra™RNA Library Preparation Kit (New England Biolabs). The library quality

was assessed using the Agilent Bioanalyzer 2100 system and then sequenced using the

Illumina Novaseq platform. The fastp software was utilized to filter the original data

and obtain high-quality data. The reference genome index was established using BWA

(version 0.7.12), and the filtered data was compared with the reference genome using

BWA mem (version 0.7.12). After the comparison was completed, the IP enrichment

region was identified against the background using MACS2(version 2.1.0) peak call

software. After peak calling was completed, the results were analyzed for chromosome

distribution, peak width, fold abundance, significance level and peak number per peak.

Homer (version 4.9.1) software was used to identify m^6A enrichment motifs in each

group. Peak Annotator software was used to identify peak-related genes. Differential

peak analysis was carried out based on the different enrichment times of the peaks in

the results, and then the genes related to the difference peak were identified.

Co-Immunofluorescent (Co-IF) assay

HGC-27 cells were evenly distributed on coverslips in a 6-well plate. When the cell

fusion rate reached 20-30%, the cells were gently rinsed with PBS for three times.

Subsequently, the cells were fixed with 4% paraformaldehyde, permeated with 0.2%

Triton X-10 (Solarbio; Catalog number: T8200), and blocked with 2% bovine serum

albumin (BSA-V; Solarbio; Catalog number: A8020). The cells were then incubated

overnight at 4℃ with IGF2BP3 antibody (diluted 1:50; Proteintech; Catalog number:

14642-1-AP) and FBXO32 antibody (diluted 1:50; Proteintech; Catalog number:

67172-1-Ig). Following this, the cells were incubated with fluorescent secondary

antibody (diluted 1:500; Cy3 anti-rabbit IgG, FITC anti-mouse IgG; Beyotime

Biotechnology Co., Shanghai, China) at room temperature for 1 hour in a dark

environment. Finally, 4 '6' -diamino-2-phenylindole dihydrochloride (DAPI; Beyotime

Biotechnology Co.) was used to stain the nuclei. Representative images were captured

using a laser scanning confocal microscope (ZEISS, Oberkochen, Germany).

Bioinformatics analysis

The expression of IGF2BP3 mRNA in cancer and adjacent tissues of patients with GC

was obtained from TCGA (https://www.aclbi.com/static/index.html#/tcga) and GEO

(https://www.aclbi.com/static/index.html#/geo) databases. The online website Kaplan-

Meier Plotter (https://kmplot.com/analysis/) was used to analyze the influence of gene

expression level on the prognosis of GC patients. GEPIA2 (http://gepia2.cancer-

pku.cn/#index) was used to analyze gene expression levels in GC and adjacent normal

tissues, while GeneCards (https://www.genecards.org/) provided insight into gene

sublocalization within cells. PrimerBank (https://pga.mgh.harvard.edu/primerbank/)

and the Primer designing tool (https://www.ncbi.nlm.nih.gov/tools/primer-blast/) were

employed for primer prediction. BioRender (https://www.biorender.com/) facilitated

the creation of scientific schematic diagrams, and catRAPID omics

(http://s.tartaglialab.com/page/catrapid_group) was used to predict binding site motifs

between the IGF2BP3 protein and FBXO32 mRNA. The iGV software was utilized for

the analysis of genomic m^6A methylation levels.

Statistical analysis

In this study, three independent replicates were performed for each group of

experiments. The results of normal distribution data were presented as mean ± standard

deviation, while non-normally distributed data were presented as median and

interquartile distance. Statistical analysis in this study was processed using GraphPad

Prism 9.5 (GraphPad Software, La Jolla, CA, USA) and SPSS 26.0 (IBM, Armonk, NY,

USA). Statistical methods including Student's *t*-test, one-way ANOVA and two-way

ANOVA were employed. A significance level of $P<0.05$ was considered statistically

significant.

Data Availability

The data generated in this study are available upon request from the corresponding

author. The data analyzed in this study were obtained from The Cancer Genome Atlas

(TCGA) and Gene Expression Omnibus (GEO) at GSE54129 and GSE66229.

Results

IGF2BP3 is upregulated in GC patients and is associated with poor prognosis

To investigate the role of m^{6}A related genes in the pathogenesis of GC, we first analyzed

the expression of 21 m^{6}A related genes in GC and adjacent normal tissues in the TCGA

database and the GSE54129 and GSE66229 datasets (Figure 1A-C). Four m^{6}A reader

proteins (YTHDF1, YTHDF3, IGF2BP1 and IGF2BP3) with higher expression levels

in GC than in adjacent normal tissues were screened. Among them, IGF2BP3 exhibited

the most significant differential expression, and due to the limited research on its

mechanism in GC, we selected IGF2BP3 as our research focus. Kaplan-Meier survival

analysis suggested that GC patients with high IGF2BP3 expression had worse overall

survival compared to those with low IGF2BP3 expression (Figure 1D). We then

assessed IGF2BP3 mRNA expression in 50 pairs of fresh frozen GC tissues and

adjacent normal tissues using qRT-PCR. Consistent with the TCGA and GEO data, the

expression of IGF2BP3 was significantly higher in GC tissues compared to adjacent normal tissues, with a statistically significant difference ($P<0.01$) (Figure 1E). IHC was employed to detect IGF2BP3 protein expression in these samples. The results showed that IGF2BP3 protein was primarily localized in the cytoplasm of GC cells, and was expressed at significantly higher levels than in adjacent normal tissues ($\chi^2 =$ 22.415), with a statistically significant difference ($P<0.001$) (Figure 1F, Table 1). Finally, we analyzed the correlation between IGF2BP3 expression and the clinical characteristics of GC patients. The results suggest that IGF2BP3 expression is significantly associated with lymph node metastasis ($P<0.001$), TNM staging ($P =$ 0.001), and invasion range ($P = 0.001$), while it shows no correlation with gender, age, tumor size, or tumor differentiation (Table 2). These findings suggest that the m^6A reader protein IGF2BP3 is highly expressed in GC and is closely associated with poor prognosis in GC patients.

IGF2BP3 knockdown suppress the progression of GC *in vitro*

In order to explore the mechanism of IGF2BP3 in GC, we first downregulated IGF2BP3 expression in HGC-27 and MKN74 cell lines and observed the effects on various cellular functions. Following transfection with siRNA targeting IGF2BP3 (siIGF2BP3) and a siRNA negative control (siNC), both the mRNA (Figure 2A, B) and protein (Figure 2C, D) levels of IGF2BP3 were significantly reduced. The CCK-8 assay revealed that cells proliferation was markedly inhibited after knockdown of IGF2BP3 (Figure 2E, F). The wound healing assay (Figure 2G, H) and Transwell migration assay (Figure 2I, J) demonstrated that the downregulation of IGF2BP3 significantly

suppressed cells migration. Furthermore, the Transwell invasion assay indicated that

reducing IGF2BP3 expression significantly impaired the invasion capability of the cells

(Figure 2K, L). Flow cytometry analysis showed that silencing IGF2BP3 expression

promoted the apoptosis of HGC-27 and MKN74 cells (Figure 2M, N). These results

suggest that downregulation of IGF2BP3 inhibits proliferation, migration and invasion

of GC cells and promotes apoptosis.

Overexpression of IGF2BP3 can promote the progression of GC *in vitro*

We upregulated the expression of IGF2BP3 in MKN7 and GES-1 cell lines to observe

its effect on cellular biological functions. After transfecting the cells with pcDNA3.1-

IGF2BP3 (IGF2BP3) and pcDNA3.1-vector (Vector), we obversed a significant

increase in IGF2BP3 mRNA (Figure 3A, B) and protein (Figure 3C, D) levels. The

CCK-8 assay demonstrated a notable enhancement in cell proliferation following after

the overexpression of IGF2BP3 (Figure 3E, F). Both the wound healing assay (Figure

3G, H) and the Transwell migration assay (Figure 3I, J) indicated that upregulating

IGF2BP3 significantly strengthened the cells' migration capability. Furthermore, the

Transwell invasion assay revealed that elevated IGF2BP3 expression significantly

heightened the invasion ability of cells (Figure 3K). The flow cytometry analysis

revealed that increased IGF2BP3 expression led to a reduction in apoptosis levels in

both MKN7 and GES-1 cell lines (Figure 3L, M). Despite multiple experiments, GES-

1 cells did not demonstrate the ability to penetrate the matrix glue at the base of the

Transwell chambers, due to the inherently weaker invasion ability of normal epithelial

cells. In summary, these results show that overexpression of IGF2BP3 can promote the

proliferation, migration and invasion of GC cells and inhibit their apoptosis, suggesting that IGF2BP3 may play the role of a pro-oncogene in GC.

IGF2BP3 promotes glucose metabolism in GC cells

Tumor cells rapidly produce a large amounts of energy through a unique aerobic glycolysis pathway to meet their surging metabolic demands, thereby maintaining normal cellular functions. We monitored the levels of glucose, lactic acid and ATP in GC cells after modulating the expression of IGF2BP3 to explore its role in glucose metabolism in these cells. The results showed that after down-regulating or up-regulating the expression level of IGF2BP3 in HGC-27 and MKN7 cells, the intracellular levels of glucose (Figure S1A), lactic acid (Figure S1B) and ATP (Figure S1C) were correspondingly decreased or increased. These findings suggest that IGF2BP3 may play a significant role in regulating glucose metabolism in GC cells. In the near future, we will continue to examine changes in the oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) in GC cells after altering IGF2BP3 expression, to further explore the effects of IGF2BP3 on the glycolytic capability of these cells.

IGF2BP3 deficiency inhibits the growth of GC *in vivo*

In order to further investigate the mechanism of IGF2BP3 in GC *in vivo*, we established tumor bearing nude mice models with low expression of IGF2BP3 and corresponding negative controls. HGC-27 cell lines stably transfected with shRNA-IGF2BP3 and shRNA-negative control were successfully established (shNC/shIGF2BP3) and subsequently injected subcutaneously into two groups of nude mice, respectively. The

tumor-bearing models, along with the curves of tumor volume, tumor photographs and

weights were presented in Figure 4A-D. The results demonstrated a significant

reduction in the growth rate, volume, and weight of tumors in the shIGF2BP3 group

compared to those in the shNC group. HE staining was utilized to evaluate the

morphological characteristics of the xenograft tissues, and the results indicated that the

cell morphology and structure were similar to those of GC tissues (Figure 4E). Ki67

immunohistochemical staining indicated that the proliferation ability of cells in the

shIGF2BP3 group was significantly inhibited compared with the shNC group (Figure

4F). These findings suggest that IGF2BP3 knockdown can inhibit the growth of GC *in*

vivo.

FBXO32 is the m^{6}A modification target of IGF2BP3 in GC

In order to further explore the potential mechanism of IGF2BP3 in the pathogenesis of

GC, we performed RNA-seq, MeRIP-seq, and RIP-seq in HGC-27 cells of siNC group

and siIGF2BP3 group, respectively. The RNA-seq results indicated that the expression

levels of a total of 388 genes were significantly changed after IGF2BP3 expression was

knocked down, including 181 down-regulated genes and 207 up-regulated genes

(Figure 5A, B). MeRIP-seq results showed that 25,331 m^{6}A peaks were identified in

the siNC group and 26,188 m^{6}A peaks in the siIGF2BP3 group (Figure 5C). When

mapping the m^{6}A methyl group in HGC-27 cells, the m^{6}A consistent sequence motif

(GGAC) could be identified, indicating successful enrichment of mRNA with m^{6}A

modifications (Figure 5D). In addition, m^{6}A modification was observed to be mainly

enriched in the CDS and 3' UTR of mRNA (Figure 5E). Then, the genes corresponding

to the m^{6}A peaks of the siNC group and siIGF2BP3 group were intersected, resulting

in the identification of 10,514 genes (Figure 5F). Analysis of the RIP-seq results

showed that m^{6}A modification was also primarily enriched in the CDS and 3' UTR of

mRNA (Figure 5G). Specifically, 1,928 and 810 m^{6}A peaks were identified in siNC

group and siIGF2BP3 group, respectively (Figure 5H). After removing the same genes

from the m^{6}A peaks corresponding genes of both group, 73 unique genes were

identified (Figure 5I). Subsequent unified analysis of the genes identified by RNA-seq,

MeRIP-seq and RIP-seq showed that 19 genes bound to the IGF2BP3 protein were

simultaneously labeled by m^{6}A, and none of them had changes in transcription level

after IGF2BP3 knockdown. These genes were identified as IGF2BP3 candidate target

genes (Figure 5J). Further analysis of these genes included examing their expression in

GC tissues, subcellular localization, survival of GC patients associated with these genes,

and the abundance and loci of m^{6}A on their mRNA. The results suggested that FBXO32

was highly expressed in GC tissues according to the TCGA database (Figure 5K),

predominantly distributed in the cytoplasm and nucleus (Figure 5L), and the overall

survival of patients with high expression of FBXO32 was poor (Figure 5M), and the

IGF2BP3 binding site was coincide with the m^{6}A modification site (Figure 5N).

Additionally, RIP-qPCR demonstrated that the IGF2BP3 protein could significantly

enrich FBXO32 mRNA (Figure 5O). Moreover, as predicted by the catRAPID website,

there is a direct binding site between the IGF2BP3 protein and FBXO32 mRNA, as

shown in the motif (Figure 5P). Based on the above analysis results, FBXO32 was

identified as the m^{6}A modification target of IGF2BP3 in GC.

FBXO32 is highly expressed in GC and directly interacts with IGF2BP3

Next, we initially verified the expression of FBXO32 in GC cell lines and GC tissues. The results indicated that the level of FBXO32 mRNA in GC cell lines was higher than that in normal gastric mucosa cells (Figure 6A), and in GC tissues, it was higher in adjacent tissues (34 pairs of GC paired tissues) (Figure 6B). The expression of FBXO32 protein was highest in MKN7 cells (Figure 6C), and higher in GC tissues than in adjacent tissues (19 pairs of GC paired tissues) (Figure 6D). These results suggest that FBXO32 may acts as an oncogene in GC pathogenesis. Subsequently, the interaction between IGF2BP3 and FBXO32 was verified by Co-IF experiments. The Co-IF results showed that IGF2BP3 and FBXO32 were mainly located in the cytoplasm and nucleus of GC cells, and they have co-localization (Figure 6E). Finally, to further confirm the regulation of FBXO32 expression by IGF2BP3 in GC cells, we examined the transcriptional and translational changes of FBXO32 after IGF2BP3 knockdown or overexpression. After down-regulating the expression level of IGF2BP3 in HGC-27 cells, the expression level of FBXO32 mRNA did not change significantly (Figure 6F), while the expression level of its protein was down-regulated (Figure 6G). Similarly, when IGF2BP3 expression was upregulated in MKN7 cells, the expression level of FBXO32 mRNA was not significantly changed (Figure 6H), while its protein expression level was upregulated (Figure 6I). Theses phenomena indicate that IGF2BP3, as an m^6A reader, only regulates the expression of downstream genes at the translation level, not at the transcriptional level, which is consistent with the existing research conclusions on other m^6A readers. In conclusion, these results indicate that IGF2BP3

and FBXO32 interact with each other in GC, and IGF2BP3 can regulate the expression

of FBXO32 protein.

The oncogenic effect of IGF2BP3 depends on targeting FBXO32 and activating

the cGMP-PKG signaling pathway

Four siRNA-FBXO32 sequences were transfected simultaneously into MKN7 cells

with the highest FBXO32 expression, and one of the siRNA sequences with the most

significant knockdown efficiency (si-FBXO32-910) was screened for subsequent

experiments (Figure S2A). Next, we transfected siRNA-negative control/siRNA-

IGF2BP3 and pcDNA3.1-vector/pcDNA3.1-FBXO32 into HGC-27 and pcDNA3.1-

vector/pcDNA3.1-IGF2BP3 and siRNA-negative control/siRNA-FBXO32 into MKN7

cells, respectively or simultaneously. The changes of IGF2BP3 expression in each

group were detected by qRT-PCR. The results suggest that the knockdown effect of

IGF2BP3 in HGC-27 cells could be attenuated by the overexpression of FBXO32

(Figure S2B), and the up-regulation effect of IGF2BP3 in MKN7 cells could also be

weakened by the knockdown of FBXO32 (Figure S2C). These findings indicate that

FBXO32 can affect the expression of IGF2BP3 in GC cells. To further explore the

downstream molecular mechanisms of FBXO32, we selected the top 1000 genes co-

expressed with FBXO32 in gastric adenocarcinoma from the TCGA database and

analyzed the signaling pathways controlled by them. The results suggested that genes

in the cGMP-PKG signaling pathway were significantly enriched (Figure 7A).

Subsequently, we tested whether FBXO32 could rescue the inhibitory effects on cGMP-

PKG signaling pathway activation induced by the knockdown of IGF2BP3. The results

suggest that when the expression level of IGF2BP3 is downregulated in HGC-27 cells, the expression levels of downstream effectors of cGMP (PKG1, pVASP) are also significantly downregulated. Concurrently, overexpression of FBXO32 can rescue the downregulatory effects on these effectors caused by the knockdown of IGF2BP3 (Figure 7B). These findings suggest that FBXO32 is involved in the regulation of IGF2BP3＇s activation of the cGMP-PKG signaling pathway. Next, we further explored whether the effects of IGF2BP3 on GC cell function would be influenced by FBXO32. The results showed that the attenuation of cell proliferation, migration, and invasion induced by knockdown of IGF2BP3 in HGC-27 cells could be rescued by the up-regulation of FBXO32 expression (Figure 7C-G). The enhancement of cell proliferation, migration, and invasion caused by overexpression of IGF2BP3 in MKN7 cells could be attenuated by the down-regulation of FBXO32 expression (Figure 7D-L). In summary, FBXO32 serves as a crucial downstream target for IGF2BP3 in promoting the progression of GC, and the oncogenic effect of IGF2BP3 in GC is dependent on FBXO32 and the cGMP-PKG signaling pathway. Therefore, we propose that increased IGF2BP3 can bind to more FBXO32 mRNA by recognizing m^{6}A sites on the FBXO32 mRNA, leading to an increase in FBXO32 protein expression and subsequently activating the cGMP-PKG signaling pathway, ultimately promoting tumorigenesis in GC cells (Figure 7M).

Discussion

In this study, we report for the first time that IGF2BP3 binds to FBXO32 mRNA by recognizing the m^{6}A site, resulting in increased intracellular level of FBXO32 protein,

which then activates the cGMP-PKG signaling pathway, ultimately promotes the progression of GC. Currently, no optimal prognostic biomarker for GC has been identified. This study also confirmed that the expression of IGF2BP3 is upregulated in GC, which is associated with poor prognosis in patients and alterations in cellular glucose metabolism. It has unveiled the clinical significance and potential value of the IGF2BP3/FBXO32/cGMP-PKG axis in the therpy of GC.

Through literature review and database analysis, we found that the expression of IGF2BP3 was most significantly different in GC tissues compared to adjacent normal tissues. However, the relationship between IGF2BP3 and GC remains to be further explored. Previous studies indicate that the expression level of IGF2BP3 in laryngeal squamous cell carcinoma (LSCC) was higher than that in adjacent normal tissues[15]. Additionally, the expression of IGF2BP3 is elevated in gallbladder cancer and is associated with poor prognosis of patients, which can enhance multiple malignant phenotypes of gallbladder cancer cells, and knockdown of IGF2BP3 can inhibit the tumor growth *in vivo*[19]. The expression level of IGF2BP3 in colon cancer is also significantly higher than that in adjacent normal tissues and is associated with worse overall survival. Altering the expression level of IGF2BP3 can regulate the cell cycle and affect cell proliferation, and knocking it down can inhibit the tumor growth of colon cancer *in vivo*[17]. Consistent with these findings, our study revealed that the expression levels of IGF2BP3 mRNA and protein in GC tissues were higher than those in adjacent normal tissue. The increased expression level of IGF2BP3 was closely related to lymph node metastasis, more advanced TNM stage and deeper invasion depth. In our

554 experiments, knocking down the expression level of IGF2BP3 in GC cells significantly

555 inhibited, cell proliferation, migration and invasion, and increased apoptosis.

556 Conversely, upregulating IGF2BP3 expression in GC cells had the opposite effect.

557 Furthermore, we observed that IGF2BP3 deficiency inhibited the growth of GC *in vivo*.

558 These findings suggest that IGF2BP3 may be an important oncogene in GC.

559 Additionally, Wang J et al. found that IGF2BP2 enhanced the stability of GLUT1

560 mRNA, a key gene of aerobic glycolysis, in an m^6A dependent manner, significantly

561 promoting the aerobic glycolysis in colorectal cancer cells[18]. Our results also showed

562 that after up-regulating or down-regulating the expression level of IGF2BP3 in GC cells,

563 the contents of glucose, lactic acid and ATP in the cells correspondingly

564 increased/decreased. Thus, the effects of IGF2BP3 on the biological function of GC

565 cells may be closely related to the regulation of glucose metabolism in cells, which

566 warrants further exploration.

567 Through RNA-seq, MeRIP-seq, RIP-seq and bioinformatics analysis, we identified

568 FBXO32 as the target of IGF2BP3 with m^6A modification in GC for the first time. F-

569 Box Protein 32 (FBXO32), originally identified as a muscle-specific gene involved in

570 muscle atrophy, is an E3 ubiquitin ligase located on chromosome 8q24.13[20, 21].

571 Previous studies have reported that FBXO32 is involved in various processes such as

572 epithelial-mesenchymal transformation and tumorigenesis[22]. The overexpression of

573 FBXO32 is conducive to the proliferation and migration of melanoma cells, while its

574 downregulation has the opposite effect[23]. FBXO32 is abnormally upregulated in

575 pancreatic ductal adenocarcinoma, promoting the migration and invasion of pancreatic

cancer cells *in vitro*, and promoting tumor growth and metastasis *in vivo*[24]. FBXO32

enhances the PI3K/AKT/mTOR pathway by degrading PTEN, thereby promoting the

progression of lung adenocarcinoma[25]. In this study, the expression levels of FBXO32

mRNA and protein in GC tissues were higher than those in adjacent normal tissues, and

the expression levels of FBXO32 mRNA and protein in GC cells were also higher than

those in normal gastric mucosa cells. These findings suggest that FBXO32 may be an

important oncogene in GC. However, there are some results that contradict our

conclusions. For example, the expression of FBXO32 is down-regulated in multiple

myeloma[26]. Overexpression of FBXO32 can inhibit tumorigenicity of ovarian cancer

cells both *in vivo* and *in vitro*[27]. Therefore, the role of FBXO32 in cancer cannot be

generalized, and more extensive and in-depth studies are needed to elucidate its

mechanisms. Subsequently, Co-IF experiments confirmed that IGF2BP3 and FBXO32

had a direct interaction relationship in GC cells, and this conclusion has not been

reported in existing studies. In addition, as predicted by bioinformatics analysis, there

was a binding site between IGF2BP3 protein and FBXO32 mRNA, which further

supports our conclusion about their interaction. To explore the regulatory mechanism

of IGF2BP3 on FBXO32 in GC, we demonstrated that IGF2BP3 regulates FBXO32

expression at the translational level but not at the transcriptional level. Previous studies

have reported that IGF2BP3 can enhance the translation of COPS7B mRNA and

promote the growth and metastasis of colorectal cancer[28]. IGF2BP3 promotes EGFR

mRNA translation in an m^6A-dependent manner, contributing to CRC progression and

cetuximab resistance[29]. Thus, it is possible that IGF2BP3 binds FBXO32 mRNA in an

m[6]A-dependent manner and promotes its translation, ultimately contributing to GC progression.

To further elucidate the mechanism by which IGF2BP3 targets FBXO32 to promote GC progression, we analyzed the signaling pathways controlled by the top 1000 genes co-expressed with FBXO32 in gastric adenocarcinoma from the TCGA database. The results suggest that these genes are significantly enriched in the cGMP-PKG signaling pathway. The cGMP-PKG signaling pathway was originally reported to participate in the inhibition of platelet activation and the phosphorylation of VASP regulated by this pathway is an integral part of an efficient and sensitive signaling cascade[30]. Many recent studies have shown that the cGMP-PKG signaling pathway is also involved in the development of various cancers. For example, cGMP can activate PKG and downstream MAPK pathways, resulting in the increased tumor cell stemness and metastasis[31]; DARS-AS1 can accelerate the progression of cervical cancer by activating the cGMP-PKG pathway[32]; NO-cGMP-PKG can increase the proliferation of colon cancer cells while inhibiting apoptosis by activating ERK-1/2 and AP-1[33]. Subsequently, we confirmed through WB that the inhibitory effect on cGMP-PKG signaling pathway activation caused by the knockdown of IGF2BP3 can be rescued by FBXO32, and FBXO32 can also activate the cGMP-PKG signaling pathway. Finally, we also conducted rescue studys to further confirm that the IGF2BP3/FBXO32/cGMP-PKG axis can affect the proliferation, migration, and invasion capabilities of GC cells. These results indicate that the IGF2BP3/FBXO32/cGMP-PKG axis is involved in the occurrence and progression of GC and represents a promising new therapeutic target

for clinical treatment of GC.

Admittedly, our research currently has certain limitations. For example, the human

tissue samples used for clinical validation were all from the same hospital, and there

was a lack of multi-center clinical data. In addition, our findings about the effect of

changes in IGF2BP3 expression level on cellular glucose metabolism in GC cells are

worthy of further exploration to elucidate the in-depth mechanism of GC and glucose

metabolism.

In conclusion, our study demonstrated that IGF2BP3 is a oncogenic protein in GC,

which promotes the translation of FBXO32 in an m^6A dependent manner and activates

the downstream cGMP-PKG signaling pathway, thereby modulating multiple

biological functions of GC cells and ultimately promoting the progression of GC.

Consequently, targeting the IGF2BP3/FBXO32/cGMP-PKG axis may offer a novel

approach to the clinical treatment of GC.

Acknowledgments

This work was supported by Hebei Provincial Government-funded Provincial Medical

Excellent Talent Project (ZF2023025, ZF2024134, and LS202008), Hebei Natural

Science Foundation (H2022206292, H2024206140), Key R&D Program of Hebei

Province (223777103D and 223777113D), Prevention and treatment of geriatric

diseases by Hebei Provincial Department of Finance (LNB202202, LNB201809 and

LNB201909), Spark Scientific Research Project of the First Hospital of Hebei Medical

University (XH202312 and XH201805), Hebei Province Medical Applicable

Technology Tracking Project (G2019035) and other projects of Hebei Province (1387 and SGH201501).

CRediT authorship contribution statement

Yi Si: Conceptualization, Data curation, Formal analysis, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing. **Bo Tian:** Conceptualization, Data curation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing. **Rui Zhang:** Visualization, Writing – original draft. **Mingda Xuan:** Methodology, Writing – original draft. **Kunyi Liu:** Visualization, Writing – original draft. **Jiao Jiao**: Data curation, Writing – original draft. **Shuangshuang Han:** Formal analysis, Visualization, Writing – original draft. **Hongfei Li:** Formal analysis, Writing – original draft. **Yanhong Hu:** Formal analysis, Writing – original draft. **Hongyan Zhao:** Investigation, Writing – original draft. **Wenjing He:** Investigation, Writing – original draft. **Jia Wang:** Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing – review & editing. **Ting Liu:** Conceptualization, Funding acquisition, Project administration, Validation, Writing – review & editing. **Weifang Yu:** Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing – review & editing.

Conflict Of Interest Statement

The authors have no conflicts of interest to declare.

References

1. Chia NY, Tan P. Molecular classification of gastric cancer[J]. Annals of oncology : official journal of the European Society for Medical Oncology, 2016, 27(5): 763-769.

2. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, *et al*. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries[J]. CA: a cancer journal for clinicians, 2024, 74(3): 229-263.

3. Song Z, Wu Y, Yang J, Yang D, Fang X. Progress in the treatment of advanced gastric cancer[J]. Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine, 2017, 39(7): 1010428317714626.

4. Fukagawa T, Katai H, Mizusawa J, Nakamura K, Sano T, Terashima M, *et al*. A prospective multi-institutional validity study to evaluate the accuracy of clinical diagnosis of pathological stage III gastric cancer (JCOG1302A)[J]. Gastric cancer : official journal of the International Gastric Cancer Association and the Japanese Gastric Cancer Association, 2018, 21(1): 68-73.

5. Arnold M, Karim-Kos HE, Coebergh JW, Byrnes G, Antilla A, Ferlay J, *et al*. Recent trends in incidence of five common cancers in 26 European countries since 1988: Analysis of the European Cancer Observatory[J]. European journal of cancer (Oxford, England : 1990), 2015, 51(9): 1164-1187.

6. Tang SY, Zhou PJ, Meng Y, Zeng FR, Deng GT. Gastric cancer: An epigenetic view[J]. World journal of gastrointestinal oncology, 2022, 14(1): 90-109.

7. Han X, Wang L, Han Q. Advances in the role of m(6)A RNA modification in cancer

metabolic reprogramming[J]. Cell & bioscience, 2020, 10: 117.

8. Xie Q, Wu TP, Gimple RC, Li Z, Prager BC, Wu Q, *et al*. N(6)-methyladenine DNA

Modification in Glioblastoma[J]. Cell, 2018, 175(5): 1228-1243.e1220.

9. Yang Y, Hsu PJ, Chen YS, Yang YG. Dynamic transcriptomic m(6)A decoration:

writers, erasers, readers and functions in RNA metabolism[J]. Cell research, 2018,

28(6): 616-624.

10. Lee M, Kim B, Kim VN. Emerging roles of RNA modification: m(6)A and U-tail[J].

Cell, 2014, 158(5): 980-987.

11. Jacobs BL, Lee CT, Montie JE. Bladder cancer in 2010: how far have we come?[J].

CA: a cancer journal for clinicians, 2010, 60(4): 244-272.

12. Bell JL, Wächter K, Mühleck B, Pazaitis N, Köhn M, Lederer M, *et al*. Insulin-like

growth factor 2 mRNA-binding proteins (IGF2BPs): post-transcriptional drivers of

cancer progression?[J]. Cellular and molecular life sciences : CMLS, 2013, 70(15):

2657-2675.

13. Müeller-Pillasch F, Lacher U, Wallrapp C, Micha A, Zimmerhackl F, Hameister H,

et al. Cloning of a gene highly overexpressed in cancer coding for a novel KH-domain

containing protein[J]. Oncogene, 1997, 14(22): 2729-2733.

14. Liu X, Chen J, Chen W, Xu Y, Shen Y, Xu X. Targeting IGF2BP3 in Cancer[J].

International journal of molecular sciences, 2023, 24(11).

15. Wang X, Tian L, Li Y, Wang J, Yan B, Yang L, *et al*. RBM15 facilitates laryngeal

squamous cell carcinoma progression by regulating TMBIM6 stability through

IGF2BP3 dependent[J]. Journal of experimental & clinical cancer research : CR, 2021,

40(1): 80.

16. Xia A, Yuan W, Wang Q, Xu J, Gu Y, Zhang L, *et al*. The cancer-testis lncRNA lnc-

CTHCC promotes hepatocellular carcinogenesis by binding hnRNP K and activating

YAP1 transcription[J]. Nature cancer, 2022, 3(2): 203-218.

17. Yang Z, Wang T, Wu D, Min Z, Tan J, Yu B. RNA N6-methyladenosine reader

IGF2BP3 regulates cell cycle and angiogenesis in colon cancer[J]. Journal of

experimental & clinical cancer research : CR, 2020, 39(1): 203.

18. Wang J, Zhu M, Zhu J, Li J, Zhu X, Wang K, *et al*. HES1 promotes aerobic

glycolysis and cancer progression of colorectal cancer via IGF2BP2-mediated GLUT1

m6A modification[J]. Cell death discovery, 2023, 9(1): 411.

19. Qin J, Cui Z, Zhou J, Zhang B, Lu R, Ding Y, *et al*. IGF2BP3 drives gallbladder

cancer progression by m6A-modified CLDN4 and inducing macrophage

immunosuppressive polarization. Transl Oncol. 2023 Nov;37:101764.

20. Sahu SK, Tiwari N, Pataskar A, Zhuang Y, Borisova M, Diken M, *et al*. FBXO32

promotes microenvironment underlying epithelial-mesenchymal transition via CtBP1

during tumour metastasis and brain development. Nature communications, 2017, 8(1):

1523.

21. Mei Z, Zhang D, Hu B, Wang J, Shen X, Xiao W. FBXO32 Targets c-Myc for

Proteasomal Degradation and Inhibits c-Myc Activity. The Journal of biological

chemistry, 2015, 290(26): 16202-16214.

22. Wang H, Liu H, Zhao L, Luo S, Akinyemiju T, Hwang S, *et al*. Association of

genetic variants of FBXO32 and FOXO6 in the FOXO pathway with breast cancer risk.

Molecular carcinogenesis, 2021, 60(10): 661-670.

23. Li Y, Gao S, Xue W, Ma Y, Meng Y, Zhang D. miR-19a-3p Functions as an Oncogene by Regulating FBXO32 Expression in Multiple Myeloma. Balkan medical journal, 2021, 38(1): 43-49.

24. Su D, Wang R, Chen G, Ding C, Liu Y, Tao J, *et al*. FBXO32 Stimulates Protein Synthesis to Drive Pancreatic Cancer Progression and Metastasis. Cancer Res. 2024 May 22.

25. Wu J, Wen T, Marzio A, Song D, Chen S, Yang C, *et al*. FBXO32-mediated degradation of PTEN promotes lung adenocarcinoma progression. Cell Death Dis. 2024 Apr 20;15(4):282.

26. Guo W, Zhang M, Shen S, Guo Y, Kuang G, Yang Z, *et al*. Aberrant methylation and decreased expression of the TGF-β/Smad target gene FBXO32 in esophageal squamous cell carcinoma. Cancer, 2014, 120(16): 2412-2423.

27. Habel N, El-Hachem N, Soysouvanh F, Hadhiri-Bzioueche H, Giuliano S, Nguyen S, *et al*. FBXO32 links ubiquitination to epigenetic reprograming of melanoma cells. Cell death and differentiation, 2021, 28(6): 1837-1848.

28. Tang J, Wang S, Weng M, Guo Q, Ren L, He Y, *et al*. The IGF2BP3-COPS7B Axis Facilitates mRNA Translation to Drive Colorectal Cancer Progression. Cancer Res. 2023 Nov 1;83(21):3593-3610.

29. Chen LJ, Liu HY, Xiao ZY, Qiu T, Zhang D, Zhang LJ, *et al*. IGF2BP3 promotes the progression of colorectal cancer and mediates cetuximab resistance by stabilizing

EGFR mRNA in an m6A-dependent manner. Cell Death Dis. 2023 Sep 1;14(9):581.

30. Eigenthaler M, Nolte C, Halbrügge M, Walter U. Concentration and regulation of cyclic nucleotides, cyclic-nucleotide-dependent protein kinases and one of their major substrates in human platelets. Estimating the rate of cAMP-regulated and cGMP-regulated protein phosphorylation in intact cells[J]. European journal of biochemistry, 1992, 205(2): 471-481.

31. Lv Y, Wang X, Li X, Xu G, Bai Y, Wu J, Piao Y, Shi Y, Xiang R, Wang L. Nucleotide de novo synthesis increases breast cancer stemness and metastasis via cGMP-PKG-MAPK signaling pathway[J]. PLoS biology, 2020, 18(11): e3000872.

32. Kong X, Wang JS, Yang H. Upregulation of lncRNA DARS-AS1 accelerates tumor malignancy in cervical cancer by activating cGMP-PKG pathway[J]. Journal of biochemical and molecular toxicology, 2021, 35(6): 1-11.

33. Babykutty S, Suboj P, Srinivas P, Nair AS, Chandramohan K, Gopala S. Insidious role of nitric oxide in migration/invasion of colon cancer cells by upregulating MMP-2/9 via activation of cGMP-PKG-ERK signaling pathways[J]. Clinical & experimental metastasis, 2012, 29(5): 471-492.

Table 1 The expression level of IGF2BP3 protein in two groups

Group	Cases	IGF2BP3 expression		Positive Rate(%)	χ^2	P
		High	Low			
Gastric cancer tissues	63	38	25	60%		
					22.415	< 0.001***
Adjacent normal tissues	63	12	51	19%		

Table 2 The relationship between the expression of IGF2BP3 in gastric cancer tissue and the clinicoapathological data of patients

Group	Cases	IGF2BP3 expression		χ^2	P
		High	Low		
Gender					
Male	48	33	15		
Female	15	7	8	2.404	0.121
Age (years)					
$\leqq 60$	16	11	5		
> 60	47	29	18	0.256	0.613
Tumor size (cm)					
< 5	34	20	14		
≥ 5	27	20	7	1.550	0.213#
Differentiation					
Well	27	14	13		
Poorly	36	26	10	2.762	0.097
Lymphnode metastasis					
Negative	18	8	13		
Positive	45	35	10	13.867	< 0.001***
TNM Staging					
0	2	0	2		
I	4	0	4		
II	17	9	8	17.037	0.001**
III	30	21	9		
IV	10	10	0		
Invasion range					
Tis	2	0	2		
T1	3	0	3		
T2	8	5	3	14.165	0.001**
T3	5	1	4		
T4	45	34	11		

2 cases of carcinoma in situ were excluded.

Figure 1

Figure 1. Increased IGF2BP3 expression is associated with poor prognosis in GC patients. (A-C) Expression of 21 m^{6}A related genes in the TCGA database, as well as the GSE54129 and GSE66229 datasets. **(D)** Kaplan-Meier analysis of IGF2BP3 expression levels and overall survival in GC patients. **(E)** The expression levels of IGF2BP3 mRNA in 50 pairs of GC and adjacent normal tissue samples were detected by qRT-PCR. **(F)** Representative images showing IGF2BP3 protein expression levels in 63 pairs of GC and adjacent normal tissue samples, as detected by IHC. Scale bar, 20 μm. Data are presented as means ± SD. **P <0.01, ***P <0.001

Figure 2

Figure 2. Knockdown of IGF2BP3 in GC cells suppresses cell proliferation,

migration and invasion *in vitro*. (A, B) The expression of IGF2BP3 mRNA in HGC-

27 and MKN74 cells was detected by qRT-PCR. **(C, D)** The expression of IGF2BP3

protein in HGC-27 and MKN74 was detected by WB. **(E, F)** The effect of IGF2BP3

knockdown on the proliferation of HGC-27 and MKN74 cells was determined by the

CCK-8 assay. **(G, H)** The effect of IGF2BP3 knockdown on the migration of HGC-27

and MKN74 cells was assessed by the wound healing assay. Scale bar, 200 μm. **(I, J)**

The effect of IGF2BP3 knockdown on the migration of HGC-27 and MKN74 cells was

evaluated using the Transwell migration assay. Scale bar, 100 μm. **(K, L)** The effect of

IGF2BP3 knockdown on the invasion of HGC-27 and MKN74 cells was assessed using

Transwell invasion assay. Scale bar, 100 μm. **(M, N)** The effect of IGF2BP3 knockdown

on the apoptosis of HGC-27 and MKN74 cells was determined by flow cytometry. Data

are presented as means ± SD. *P <0.05, **P <0.01, ***P <0.001, ****P <0.0001

Figure 3

Figure 3. Overexpression of IGF2BP3 in GC cells and human gastric mucosa cells

promotes cell proliferation, migration and invasion *in vitro*. (A, B) The expression

of IGF2BP3 mRNA in MKN7 and GES-1 was detected by qRT-PCR. **(C, D)** The

expression of IGF2BP3 protein in MKN7 and GES-1 was detected by WB. **(E, F)** The

effect of IGF2BP3 overexpression on the proliferation of MKN7 and GES-1 cells was

assessed using the CCK-8 assay. **(G, H)** The effect of IGF2BP3 overexpression on the

migration of MKN7 and GES-1 cells was evaluated through a wound healing assay.

Scale bar, 200 μm. **(I, J)** The effect of IGF2BP3 overexpression on the migration of

MKN7 and GES-1 cells was determined utilizing a Transwell migration assay. Scale

bar, 100 μm. **(K)** The effect of IGF2BP3 overexpression on the invasion of MKN7 cells

was assessed via a Transwell invasion assay. Scale bar, 100 μm. **(L, M)** The effect of

IGF2BP3 overexpression on the apoptosis of MKN7 and GES-1 cells was analyzed by

flow cytometry. Data are presented as means ± SD. ***P* <0.01, ****P* <0.001, *****P*

<0.0001

Figure 4

Figure 4. IGF2BP3 regulates the growth of GC *in vivo*. (A) Nude mouse tumor-

bearing model. **(B)** Tumor growth curve. **(C, D)** Images (C) and weight (D) of

xenografted tumors were analyzed. **(E)** Representative images of tumor tissue sections

stained with HE. Scale bar, 50 μm. **(F)** Representative image of Ki67 IHC staining in

tumor tissue sections. Scale bar, 50 μm. Data are shown as means ± SD. **P <0.01,

****P <0.0001

Figure 5. FBXO32 is the m^{6}A modification target of IGF2BP3. (A) Differential

expression genes (DEGs) heat map identified by RNA-seq. **(B)** The number of DEGs

identified by RNA-seq. **(C)** The number of m^{6}A peaks identified by MeRIP-seq. **(D)**

m^{6}A motif identified by MeRIP-seq. **(E)** The enrichment regions of m^{6}A peaks on

mRNA identified by MeRIP-seq. **(F)** The common genes between the siNC group and

the siIGF2BP3 group in MeRIP-seq. **(G)** The enrichment regions of m^{6}A peaks on

mRNA identified by RIP-seq. **(H)** The number of m^{6}A peaks identified by RIP-seq. **(I)**

Genes identified by RIP-seq. **(J)** Overlap analysis of MeRIP-seq, RIP-seq and RNA-

seq identified genes. **(K)** Expression of FBXO32 in GC paired tissue cohort from the

TCGA database. **(L)** Subcellular localization of FBXO32 from GeneCards. **(M)**

Kaplan-Meier analysis of FBXO32 expression levels and overall survival in patients

with GC. **(N)** The m^{6}A site and its abundance on FBXO32 transcripts were analyzed

using IGV software. **(O)** The interaction between IGF2BP3 and FBXO32 was detected

by RIP-qPCR. **(P)** Schematic diagram of the catRAPID website predicting the binding

site between IGF2BP3 protein and FBXO32 mRNA.

Figure 6

Figure 6. Expression of FBXO32 and its interaction with IGF2BP3 in GC. (A) The

expression of FBXO32 mRNA in GC cell lines and GES-1 was detected by qRT-PCR.

(B) The expression of FBXO32 mRNA in 34 pairs of GC paired tissues was detected

by qRT-PCR. **(C)** The basal expression of FBXO32 protein in cell lines was detected

by WB. **(D)** The expression of FBXO32 protein in 19 pairs of GC paired tissues was

detected by IHC. Scale bar, 50 μm. **(E)** The distribution of IGF2BP3 and FBXO32 in

HGC-27 cells was confirmed by Co-IF. Scale bar, 10 μm. **(F)** qRT-PCR was used to

detect FBXO32 mRNA expression after IGF2BP3 knockdown in HGC-27 cells. **(G)**

WB was used to detect FBXO32 protein expression after IGF2BP3 knockdown in

HGC-27 cells. **(H)** qRT-PCR was used to detect FBXO32 mRNA expression after

IGF2BP3 overexpression in MKN7 cells. **(I)** WB was used to detect FBXO32 protein

expression after IGF2BP3 overexpression in MKN7 cells. Data are presented as means

± SD. *P <0.05, **P <0.01, ****P <0.0001

Figure 7

Figure 7. Oncogenic function of IGF2BP3 depends on FBXO32 and cGMP-PKG

signaling pathway. (A) KEGG enrichment result of the top 1000 genes coexpressed

with FBXO32 in gastric adenocarcinoma in TCGA database. **(B)** The expression levels

of cGMP-PKG signaling pathway related proteins were detected by WB after

instantaneous co-transfection of IGF2BP3 siRNA and FBXO32 overexpression

plasmid in HGC-27. **(C-G)** The effects of FBXO32 on IGF2BP3 in HGC-27 cells were

detected by CCK-8 assay (C), colony formation assay (D), wound healing assay (E),

Scale bar, 200 μm, Transwell migration assay (F) and Transwell invasion assay (G).

Scale bar, 100 μm. **(H-L)** The effects of FBXO32 on IGF2BP3 in MKN7 cells were

detected by CCK-8 assay (H), colony formation assay (I), wound healing assay (J),

Scale bar, 200 μm, Transwell migration assay (K) and Transwell invasion assay (L).

Scale bar, 100 μm. **(M)** Schematic illustration of the molecular mechanism of IGF2BP3

in GC. Data are presented as means ± SD. *P <0.05, **P <0.01, ***P <0.001, ****P

<0.0001

Supplementary Table

Supplementary Table 1 The primer sequences of IGF2BP3, FBXO32, β-actin.

Supplementary Figures

Supplementary Figure 1. IGF2BP3 affects glucose metabolism in GC cell lines. (A)

The effect of IGF2BP3 expression level on glucose uptake in GC cells was determined

using a glucose uptake assay. **(B)** The effect of IGF2BP3 expression level on lactic acid

production in GC cells was determined by utilizing a lactic acid production assay. **(C)**

An ATP content assay showed the effect of IGF2BP3 expression level on ATP content

in GC cells.

Supplementary Figure 2. The knockdown efficiency of FBXO32 and the

expression level of IGF2BP3 were detected by qRT-PCR. (A) The knockdown

efficiency of four siRNA-FBXO32 sequences in MKN7 cells was assessed by qRT-

PCR. **(B, C)** The expression of IGF2BP3 mRNA in HGC-27 (B) and MKN7 (C) cells

was measured by qRT-PCR after co-transfection of respective siRNAs and

overexpression plasmids.

Supplementary Figure 3. Protein interaction network diagram.