1

2 **Giant transposons promote strain heterogeneity in a major fungal pathogen**

Emile Gluck-Thaler[1,2,3]*, Adrian Forsythe[4], Charles Puerner[5], Cecilia Gutierrez-Perez[5,6], Jason E.
Stajich[7], Daniel Croll[1], Robert A. Cramer[5], Aaron A. Vogan[4]

[1] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel,
Neuchâtel 2000, Switzerland
[2] Department of Plant Pathology, University of Wisconsin-Madison, Madison, WI 53706, USA
[3] Wisconsin Institute for Discovery, Madison, WI 53706, USA
[4] Systematic Biology, Department of Organismal Biology, Uppsala University, Uppsala, 752 36,
Sweden
[5] Department of Microbiology and Immunology, Geisel School of Medicine, Dartmouth College,
Hanover, NH 03755 USA
[6] Present address: Department of Molecular Microbiology, Washington University School of
Medicine, St. Louis, MO 63110, USA
[7] Department of Microbiology and Plant Pathology, University of California-Riverside, Riverside,
CA 92521, USA

* Corresponding author: gluckthaler@wisc.edu

Running title: *Starships* generate variation in a fungal pathogen

ORCIDs
Emile Gluck-Thaler: https://orcid.org/0000-0003-0438-7495
Adrian Forsythe: https://orcid.org/0000-0003-1966-9946
Charles Puerner: https://orcid.org/0000-0002-4402-0295
Cecilia Gutierrez-Perez: https://orcid.org/0000-0002-9808-0701
Jason E. Stajich: https://orcid.org/0000-0002-7591-0020
Daniel Croll: https://orcid.org/0000-0002-2072-380X
Robert A. Cramer: https://orcid.org/0000-0001-5503-5006
Aaron A. Vogan https://orcid.org/0000-0003-2013-7445

Key words: transposable element, transposon, Aspergillosis, Aspergillus fumigatus, strain
heterogeneity, secondary metabolism

Abstract and Importance word count: 311
Main text word count: 9,079

Abstract

Fungal infections are difficult to prevent and treat in large part due to strain heterogeneity which confounds diagnostic predictability. Yet the genetic mechanisms driving strain-to-strain variation remain poorly understood. Here, we determined the extent to which *Starships*—giant transposons capable of mobilizing numerous fungal genes—generate genetic and phenotypic variability in the opportunistic human pathogen *Aspergillus fumigatus*. We analyzed 519 diverse strains, including 11 newly sequenced with long-read technology and multiple isolates of the same reference strain, to reveal 20 distinct *Starships* that are generating genomic heterogeneity over timescales relevant for experimental reproducibility. *Starship*-mobilized genes encode diverse functions, including known biofilm-related virulence factors and biosynthetic gene clusters, and many are differentially expressed during infection and antifungal exposure in a strain-specific manner. These findings support a new model of fungal evolution wherein *Starships* help generate variation in genome structure, gene content and expression among fungal strains. Together, our results demonstrate that *Starships* are a previously hidden mechanism generating genotypic and, in turn, phenotypic heterogeneity in a major human fungal pathogen.

Importance

No "one size fits all" option exists for treating fungal infections in large part due to genetic and phenotypic variability among strains. Accounting for strain heterogeneity is thus fundamental for developing efficacious treatments and strategies for safeguarding human health. Here, we report significant progress towards achieving this goal by uncovering a previously hidden mechanism generating heterogeneity in the human fungal pathogen *Aspergillus fumigatus*: giant transposons called *Starships* that span dozens of kilobases and mobilize fungal genes as cargo. By conducting a systematic investigation of these unusual transposons in a single fungal species, we demonstrate their contributions to population-level variation at the genome, pangenome and transcriptome levels. The *Starship* compendium we develop will not only help predict variation introduced by these elements in laboratory experiments but will serve as a foundational resource for determining how *Starships* impact clinically-relevant phenotypes, such as antifungal resistance and pathogenicity.

Introduction

Infectious diseases caused by fungi pose a grave threat to human health and society. The World Health Organization recently coordinated a global effort to prioritize research among fungal pathogens based on unmet research needs and public health importance (1). Among the pathogens deemed most important for research include *Aspergillus fumigatus*, a globally distributed opportunistic human pathogen causing disease in an estimated 4 million people yearly (2, 3). Several infectious diseases are caused by *A. fumigatus*, including invasive pulmonary aspergillosis, which manifests primarily in immunocompromised individuals with mortality rates of up to 85% (2, 4). The treatment of *A. fumigatus* is complicated by its remarkable variation in virulence, resistance to antifungals, metabolism, and other infection-relevant traits (3, 5–8). Strain heterogeneity confounds "one size fits all" therapies and poses a significant challenge for developing efficacious disease management strategies (9). Recent evaluations of the *A. fumigatus* pangenome have revealed extensive genetic variability underlying variation in clinically-relevant traits such as antifungal resistance and virulence, yet in many cases, the origins of such variation remain unexplained (10–12). Determining the genetic drivers of strain heterogeneity will help accelerate the development of strain-specific diagnostics and targeted therapies.

Mobile genetic elements (MGEs) are ubiquitous among microbial genomes and their activities profoundly shape the distribution of phenotypic variation (13). MGE transposition generates structural variation that directly impacts gene regulation and function (14–16), and many elements also modulate genome content by acquiring genes as "cargo" and transposing them within and between genomes, facilitating gene gain and loss (17, 18). The ability to generate contiguous genome assemblies with long-read sequencing technologies has dramatically enhanced our ability to find new lineages of MGEs (13, 19). For example, we recently discovered the *Starships*, a new superfamily of MGEs found across hundreds of filamentous fungal taxa (20–22). *Starships* are fundamentally different from other fungal MGEs because they are typically 1-2 orders of magnitude larger (~20-700 kb versus 1-8 kb) and carry dozens of protein-coding genes encoding fungal phenotypes (23). In addition to flanking short direct repeats, all *Starships* possess a "captain" tyrosine site-specific recombinase at their 5' end that is both necessary and sufficient for transposition (21). Several *Starships* mediate the evolution of adaptive phenotypes: for example, *Horizon* and *Sanctuary* carry the ToxA virulence factor that facilitates the infection of wheat, and *Hephaestus* and *Mithridate* encode resistance against heavy metals and formaldehyde, respectively (24–27). *Starships* have been experimentally shown to horizontally transfer between fungal species (28), and an increasing number of comparative studies implicate them in the horizontal dissemination and repeated evolution of diverse traits (20, 21, 25–27). *Starships* may also fail to re-integrate into the genome during a transposition event, leading to the loss of the element, its cargo and its encoded phenotypes (21). Through horizontal transfer and failed re-integration events, *Starships* contribute directly to the generation of rapid gene gain and loss polymorphisms across individuals. However, we know little about how *Starships* drive genetic and phenotypic variation in fungal species relevant for human health and disease.

Here, we conduct the first systematic assessment of *Starship* activity and expression in a human
fungal pathogen to test the hypothesis that these unusual transposons are a source of strain
heterogeneity. We reveal that *Starships* are responsible in part for generating key instances of
previously unexplainable variation in genome content and structure. Our interrogation of 508
diverse clinical and environmental strains of *A. fumigatus,* combined with highly contiguous
assemblies of 11 newly sequenced strains, enabled the unprecedented quantification of *Starship*
diversity within a single species and even within the same "reference" strain. We leveraged the
wealth of functional data available for *A. fumigatus* to draw links between *Starship*-mediated
genetic variation and phenotypic heterogeneity in secondary metabolite and biofilm production
contributing to pathogen survival and virulence. We analyzed multiple transcriptomic studies and
determined that variation in *Starship* cargo expression arises from strain- and treatment-specific
effects. By revealing *Starships* as a previously unexamined mechanism generating phenotypic
variation, our work sheds light on the origins of strain heterogeneity and establishes a predictive
framework to decipher the intertwining impacts of transposons on fungal pathogenesis and human
health.

Results and Discussion

Data curation and long-read sequencing

Starships are difficult (but not impossible) to detect in short-read assemblies due to their large
size and frequent localization in regions with high repetitive content (20). The wealth of publicly
available short-read genomes is still of high value, however, due to the breadth of sampling they
provide combined with the still relevant possibility of finding *Starships*, especially in higher quality
short-read assemblies such as those available for *A. fumigatus*. We therefore deployed a hybrid
sampling strategy that leveraged both short- and long-read genome assemblies, combined with
short-read mapping to genotype *Starship* presence/absence, to enable an accurate and precise
accounting of how *Starships* impact strain heterogeneity. We downloaded 508 publicly available
A. fumigatus assemblies (mostly assembled from short-read data) that span the known genetic
diversity of this species isolated from environmental and clinical sources (51.7% clinical, 47.9%
environmental, 0.4% unknown isolation source) and used Oxford Nanopore to sequence 11
additional strains with long-read technology (66.7% clinical, 25% environmental, 8.3% unknown),
for a combined total of 519 assemblies (10, 11). The 11 isolates selected for long-read sequencing
were chosen because they represent commonly utilized laboratory strains, clinical isolates and
environmental strains from three major *A. fumigatus* clades (10). We performed additional short-
read Illumina sequencing of the 11 strains to provide additional support and error-correct the long-
read assemblies. The long-read assemblies are of reference quality, representing nearly full
chromosome assemblies with an L50 range of 4-5 and a N50 range of 1.9-4.7 Mb, and are thus
ideal complements to the available short-read assemblies for investigating the connection
between *Starship*s and strain heterogeneity (Table S1).

At least 20 active and distinct *Starships* vary among *A. fumigatus* strains

We began evaluating the impact of *Starships* on strain heterogeneity by systematically annotating them in the 519 assemblies using the starfish workflow (22). We identified a total of 787 individual *Starships* and validated these predictions by manually annotating a subset of 86 elements (Table S2, Table S3, Methods) (10, 11). As expected, more *Starships* were recovered in total from short-read assemblies, but on average more *Starships* were recovered from each reference-quality long-read assembly, highlighting the utility of sampling both short- and long-read assemblies (Fig. S1). Importantly, unlike many other fungal transposons, we found that with few exceptions, if a *Starship* is present in a strain, it is present at most in a single copy. Some other species appear to readily accumulate multiple copies of the same *Starship* (20), but this is not the case for *A. fumigatus*. In other fungi, genome defense mechanisms such as repeat-induced point mutations (RIP) are capable of degrading multi-copy *Starships* and may act to limit their copy number expansion (21); however, it remains unclear whether such mechanisms also explain the lack of copy number variation observed in *A. fumigatus* .

To determine how many different *Starships* actively generate variability in *A. fumigatus*, we grouped *Starships* into discrete types following a hierarchical framework that incorporates amino acid sequence comparisons of captain tyrosine recombinases and nucleotide sequence comparisons of the entire element (23). Following Gluck-Thaler and Vogan 2024, we assigned each *Starship* element to a family (based on similarity to a reference library of captain tyrosine recombinases), a *navis* (Latin for "ship"; based on orthology among the *A. fumigatus* captains) and a *haplotype* (based on k-mer similarity scores of the cargo sequences; Methods). For example, *Starship Osiris h4* belongs to *haplotype* 4 within the *Osiris navis*, which is part of the Enterprise-family. The most reasonable conclusion when observing the same transposon bounded by identical direct repeats at two different loci in two individuals is that this transposon is presently or recently active (21). Using a threshold that required observing the same *navis-haplotype* (hereafter *Starship*) at two or more sites, we identified 20 high-confidence *Starships* that met these criteria and 34 medium-confidence *Starships* that did not. These data reveal a phylogenetically and compositionally diverse set of *Starships* actively transposing within *A. fumigatus* that together make up this species' *Starship* compartment (Methods, Table 1, Fig. 1, Table S4, Table S5) (23). For each of the 20 high-confidence elements, we aligned representative copies present in different genomic regions to highlight how precisely and repeatedly element boundaries are conserved across transposition events (Fig. S2).

After using all available genome assembly data to define the *Starship* compartment with automated methods, we explored a variety of other approaches for augmenting and validating our predictions, including manual annotation, BLAST-based detection, and short-read mapping (Methods). Certain detection methods were more appropriate for particular analyses, resulting in the partitioning of our findings into the high-confidence, the expanded, and the genotyping datasets. To analyze *Starship* features, like gene content and pangenome distributions (see below), we limited our analyses to the set of 459 high-confidence elements for which we have end-to-end boundaries and evidence of transposition (high-confidence dataset). For analyses that examine the presence/absence of elements across genomic regions, we used an expanded set

of 1818 elements that consists of the 787 high and medium-confidence elements as well as 1031
elements whose presence was detected through BLAST searches, because these methods
maximize our ability to correctly predict the presence/absence of an element in a given region
(expanded dataset; Table S4, Table S5). For genotyping analyses that focus on detecting the
presence/absence of an element regardless of where it is found in the genome, we took a
conservative approach by limiting our analysis to 13 reference quality assemblies (reference
strains CEA10 and Af293 in addition to the 11 newly sequenced long-read strains) plus 466 *A.*
fumigatus strains with publicly available paired-end short-read Illumina data that were used to
validate *Starship* presence/absence by short-read mapping to a reference *Starship* library
(genotyping dataset; Table S21).

Starships distributions are partially explained by strain relatedness but overall heterogeneous

All high-confidence *Starships* show polymorphic presence/absence variation, representing a
previously unexamined source of genetic heterogeneity across *A. fumigatus* strains (Fig. 1A,
Table S6). For example, the two commonly used reference strains Af293 and CEA10 carry 11
different high-confidence *Starships* but share only 4 in common (or only 2, using a more
conservative threshold of not counting fragments or degraded elements; Fig. 1D, Table S7). The
number of high-confidence *Starships* per long-read isolate ranges from 0-11 (median = 3, std.
dev. = 3.39) (29, 30). To identify the underlying drivers of variation in *Starship* distributions, we
investigated the relationship between *Starship* repertoires and strain relatedness by testing if
phylogenetic signal underlies *Starship* distributions. We genotyped *Starship* presence/absence
using short-read mapping in the best sampled clade of *A. fumigatus* with well resolved
phylogenetic relationships (Clade 1 *sensu* Lofgren et al., 2022, n = 151) and found that isolate
relatedness is weakly but significantly correlated with similarity in *Starship* repertoire, which is
indicative of weak phylogenetic signal in *Starship* distributions (Fig. S3, Table S8, Mantel r =
0.163; *P* = 0.001, using the genotyping dataset). The relatively weak but significant correlation
indicates that although closely related isolates tend to share similar *Starships*, relatedness alone
cannot explain all observed variation in *Starship* repertoires. Thus, we conclude *Starships* are a
source of genetic heterogeneity among even closely related strains.

While no sequence-based or phylogenetic method to our knowledge would enable us to
unequivocally differentiate horizontal transfer events from incomplete lineage sorting, loss or
sexual and parasexual recombination at the within species level, we hypothesize that horizontal
transfer within *A. fumigatus* is at least partially responsible for generating the observed patchy
distributions of *Starships*. *Starship* horizontal transfer has occurred repeatedly between
individuals of different genera, and barriers to transfer are likely lower within species (25, 26).
Starship horizontal transfer has recently been experimentally demonstrated under simple co-
culture conditions between *A. fumigatus* and *Paecilomyces variotii*, which are separated by
approximately ~100 million years of evolutionary divergence (28). Surveys of the *Paecilomyces*
fungi further estimate that ~1/3 of *Starships* in this genus have evidence of horizontal transfer in

natural populations (28). The dynamics of *Starship* horizontal transfer among *A. fumigatus* strains
must be determined with future laboratory experiments.

Starships actively transpose in isolates of the same laboratory strain

We examined the potential for *Starships* to introduce unwanted variation into laboratory
experiments by comparing isolates of the same reference strain used by different research
groups. First, we found that *Starship* insertions in the reference Af293 assembly used in our study
coincide with 4/8 putative structural variants inferred in other Af293 strains using RNAseq
expression data by Collabardini et al. 2022 (Table S9). This suggests that in addition to other
mechanisms, *Starship* transposition and/or loss contributes to structural variation observed
among isolates of what should otherwise be clones of the same strain (6).

We gathered additional evidence supporting our hypothesis by comparing independently
collected DNA sequencing data from different isolates of the same or derived strains. We
compared the locations of *Starships* between the long-read CEA10=CBS144.89 reference
assembly (31) and 6 other CEA10 and CEA10-derived isolates from various labs, including our
own, using short-read mapping and de-novo assembly comparisons (Methods). We found that
Starship Nebuchadnezzar h1 has jumped from chromosome 6 in the long-read reference CEA10
and A1160 assemblies and some A1163 strains to chromosome 4 in the A1163 isolate used by
our lab (A1163-Cramer_SM), likely at some point after we acquired this isolate from its original
culture collection stock (Fig. 1E). A1163 is a commonly used laboratory strain derived from
CEA10; the only differences between A1163 compared with CEA10 should be that its native *pyrG*
has a nonsense mutation and that it harbors an ectopic insertion of the *A. niger pyrG* (32).
Similarly, we evaluated *Starship* heterogeneity in three additional strains, S02-30 (=AF100-9B),
TP9 (=08-19-02-30) and ATCC46645, that met the criteria of having independently collected DNA
sequencing datasets and long read assemblies. We found that ATCC46645 showed signs of
Starship heterogeneity among isolates sequenced by different research groups. Specifically, a
small number of reads supported precise excision events for *Nebuchadnezzar h1*, suggesting this
Starship has transposed or has been lost in a subset of the nuclei within the short-read sequenced
isolate (Fig. S4). This suggests that some *Starships* appear to be active under laboratory
conditions, which is directly relevant for experimental design in this case as *Nebuchadnezzar h1*
carries the HAC gene cluster known to impact biofilm-associated virulence (see below) and genes
on *Starships* may be lost through failed transposition events (21, 33).

Together, *Starship*-mediated variation among isolates of the same "clonal" *A. fumigatus* strain
suggests these transposons cause genomic instability over short-enough timescales to potentially
impact routine laboratory work and experimental reproducibility. It is interesting to note that not
all strains or closely related strains demonstrate intraspecific variation: for example, although we
did not detect variation in *Starship* location or presence/absence between the long-read CEA10
and A1160 assemblies, which have been separated for decades (31), we did detect variation
between this CEA10 strain and the A1163 isolate that we routinely use in our lab, indicating that
much remains to be known about the signals and triggers promoting within-strain *Starship*
variation.

Starships mobilize upwards of 16% of the accessory genome that differs across strains

A. fumigatus strains differ extensively in the combinations of genes found in their genomes, resulting in a large and diverse pangenome (10, 11). To quantify how much pangenomic variation is attributable to actively transposing *Starships,* we estimated the total proportion of genomic content present in the 20 high-confidence *Starships* (Fig. 2, Methods). We examined these *Starships* in the 13 reference-quality assemblies and found that between 0-2.4% of genomic nucleotide sequence (and 0-2.1% of all genes per genome) is mobilized as *Starship* cargo (Fig. 2A). We then built a pangenome with all 519 strains and extracted all orthogroups (i.e., genes) found in the 13 reference-quality genomes to gain insight into the distribution of *Starship*-associated gene content (Table S10). Across the 13 isolate pangenome, 2.9% of all genes, which corresponds to 9.7% of all accessory and singleton orthogroups, have at least 1 member carried as *Starship* cargo (Fig. 2B). Accessory and singleton orthogroups are overrepresented >3 fold in *Starships*, with ~92% of *Starship*-associated orthogroups being either accessory or singleton compared with 24.6% of non-*Starship* associated orthogroups. We determined the upper bounds of this conservative estimate by examining the 13 isolate pangenome in the context of all 54 high- and medium-confidence *Starships*, and found that 4.8% of all orthogroups, representing 16% of all accessory and singleton orthogroups, have at least 1 member carried as *Starship* cargo (Fig. S5, Table S11). Drawing parallels from decades of observations of bacterial MGEs (17) and recently published experimental data from *Starships* (21, 28), we hypothesize that localization on a *Starship* promotes a gene's rate of gain and loss through *Starship*-mediated horizontal transfer and failed re-integration events (21). Together, these data reveal a previously hidden association between *Starships* and the making of *A. fumigatus'* accessory pangenome.

Starships differ in their potential to mediate gain and loss of unique sequence

We further determined the potential for *Starships* to introduce variation into *A. fumigatus* strains by testing if *Starship* genes are uniquely found in those elements or found elsewhere in the genome. We calculated the degree of association between a given cargo gene and a *Starship* by identifying the genomic locations of best reciprocal BLAST hits to genes in the 20 high-confidence type elements (Fig. 2C, Fig. S6, Table S12, Table S13, Methods). Across the 13 reference-quality assemblies, we found that the vast majority of *Starship* genes display presence/absence variation between strains (i.e., are accessory or singleton) although several genes are present in conserved regions in these particular strains (while seemingly counter-intuitive, these genes do vary in their presence/absence and are *Starship*-associated when examining the larger 519 strain population, as expected). We found that many genes are almost always carried as cargo in active *Starships* when present in a given genome (e.g., the majority of cargo on *Tardis h1*), while others have weaker associations and are found in both *Starships* and non-*Starship* regions across different strains (e.g., *Gnosis h1*). Variation in the degree to which genes associate with active *Starships* suggests elements differ in capacity to generate accessory sequence variation among strains, highlighting the importance of investigating *Starships* at the individual element level.

Starship activity generates structural variation at a genome-wide scale

We next asked where *Starship*-mediated variation occurs in the genome to better understand the implications of *Starship* activity for genome organization. We identified the genomic locations where *Starships* introduce structural variation by sorting all 787 high and medium-confidence elements, along with elements detected by BLASTn, into homologous genomic regions (Table S14, Table S15, Table S16; expanded dataset; Methods). This enabled us to genotype individuals in the 519 strain population for segregating *Starship* insertions that are polymorphic across strains (Fig. 3, Fig. S7; Methods). Across all strains and chromosomes, we found 79 regions that contain at least 1 segregating "empty" insertion site, for a total of 154 sites distributed across all eight chromosomes (a single region can have >1 insertion site if they are located close to each other). The average number of empty sites per genome is 44.5 (range = 9-63, std. dev. = 13.88), indicating that each strain harbors dozens of sites with structural variation introduced by *Starships*. For example, the Af293 reference strain has a total of 6 full-length *Starships* with annotated boundaries and 2 *Starship* fragments, and a total of 56 segregating empty sites (Fig. 3A). We found a significant linear relationship between the number of genomic regions a given *Starship* is present in (a proxy for transposition activity) and the total copy number of that element in the 519 strain population, suggesting active *Starships* contribute more to strain heterogeneity compared with less active elements (Fig. 3C; y = 14.7x - 1.24; $P = 6.5e^{-4}$; R^2 adj = 0.46; expanded dataset). Predicted *Starship* boundaries are precisely conserved across copies present in different genomic regions, suggesting transposition, and not translocation or segmental duplication, is the major mechanism responsible for generating variation in *Starship* location (Fig. 3D).

Given that *Starships* are mobilized by a tyrosine site-specific recombinase (21), we attempted to predict each *Starship*'s target site to gain insight into the genomic sequences that are susceptible to *Starship* insertions. We manually curated the sequence features of a large sample of high-confidence *Starships* and found 19/20 have identifiable direct repeats (DRs) ranging from 2-12 bp in length while 18/20 have terminal inverted repeats (TIRs; Table 1, Table S3). DRs typically reflect a portion of the element's target site, while TIRs are predicted to facilitate transposition (34). While precise target site motifs must be confirmed experimentally, the target site TTACA(N_5)AAT that we recovered for *Nebuchadnezzar* elements, which belong to the Hephaestus-family, resembles the canonical *Hephaestus* target site TTAC(N_7)A, demonstrating the utility of our approach as a first step towards understanding what sequence features promote the gain and loss of *Starship*-mediated variation (21).

Generally, DRs >6 bp are associated with targeting of the 5S rDNA gene (e.g. *Osiris*), presumably because this is a relatively stable target site (22). In contrast to this, *Tardis* has a highly conserved 9 - 11 bp DR that does not correspond to the 5S gene. A k-mer analysis shows that the 10 bp consensus motif CTACGGAGTA is strongly overrepresented in the genome of Af293 (>99.99th percentile of k-mers; Fig. S8), indicating that this motif may represent some other type of highly conserved genomic sequence, such as a transcription factor binding site. This motif overlaps with predicted motifs associated with C6 zinc cluster factors in *A. nidulans* (https://jaspar.elixir.no/matrix/UN0291.2/) (35) and this DR sequence is further conserved among

366   other *Eurotiomycetes* (22); however, the exact functions of these predicted motifs must be
367   confirmed experimentally. This result highlights the fact that studying *Starship* elements in detail
368   can reveal other important aspects of a species' biology, and characterizing the *Tardis* motif is of
369   interest for future research.

Starships mediate variation at an idiomorphic biosynthetic gene cluster

371
Several genomic regions harbor multiple types of *Starships*, raising the possibility that strain
heterogeneity arises in part from the formation of *Starship* insertion hotspots. We investigated the
genomic region with the highest density of *Starship* insertions to define the upper bounds of
Starship-mediated structural variation at a single locus (Fig. 4C, Methods). This region spans an
average of 498.76 Kb (range = 158.07-781.54 kb; std. dev. = 83.02 kb) across the n = 210 strains
for which we could detect it, and ranges from position 584,521-1,108,409 on Chromosome 3 in
the Af293 reference assembly (representing 12.84% of the entire chromosome). By
supplementing starfish's automated genotyping with manual annotation of nine strains with
distinct alleles at this region, we found this region contains at least 7 distinct *Starships* with
identifiable DRs and 1 degraded *Starship* that range from 47.52-151.62 kb long and are inserted
into 6 independent segregating sites (Fig. 4C). The majority (75%) of the *Starships* in this region
are inserted into 5S rDNA coding sequences, which effectively fragments that copy of the 5S
rDNA gene. Total 5S rDNA copy number varies between 28-33 in the 13 reference quality
assemblies (median = 31, std. dev. = 1.4; Table S17), but between 6-8 intact copies are typically
present at this single locus, representing a ~10-fold enrichment of the 5S rDNA sequence relative
to background expectations given the length of this region (using 5S rDNA frequencies in the
Af293 genome). Thus, the enrichment of 5S rDNA at this locus potentiates *Starship*-mediated
variation and the generation of a *Starship* hotspot.

390
We were surprised to find that in addition to multiple segregating *Starships*, the Chromosome 3
region contains an idiomorphic biosynthetic gene cluster (BGC) that was previously noted to be
a recombination hotspot (Cluster 10 *sensu* Lind et al. 2017) (36, 37). The idiomorphic BGC locus
is polymorphic for upwards of 4 smaller BGC modules and is upstream of a fifth BGC encoding
the biosynthesis of fusarinine C (BGC8 *sensu* Bignell et al. 2016) that itself is embedded within a
larger stretch of sequence containing NRPS and KR-PKS core genes. While the idiomorphic
modules were not predicted by antiSMASH, manual inspection revealed that each contains a
different core SM biosynthesis gene and numerous other genes involved in metabolic processes,
suggesting they are part of a larger BGC or represent different cryptic BGCs. The NR-PKS BGC
module at this locus is carried by *Starship Osiris h4,* and its presence/absence is directly
associated with the presence/absence of the element, implicating *Starships* as a mechanism
generating idiomorphic BGCs. Furthermore, some BGC modules are disrupted by the insertion of
a *Starship* (e.g., the NRPS-like BGC module). Together, these findings support the assertion that
Starship activity helps generate selectable variation in BGC genotypes. The implication of this
variation for the expression and diversification of natural products warrants further investigation.

Starships encode clinically-relevant phenotypes and are enriched in clinical
strains

Starships encode genes with diverse functions typically associated with fungal fitness, including
carbohydrate active enzymes, BGCs and metal detoxification genes (Table 1, Table S5). To
investigate broad trends in how *Starships* might contribute to phenotypic heterogeneity, we
compared the predicted cargo functions among the 20 high-confidence *Starships* (Table S18;
high-confidence dataset; Methods). As expected, *Starship* types differ from each other in their
predicted functional content, but functional diversity also varies to some extent within elements
from the same type, indicating that both inter- and intra-specific *Starship* variation may contribute
to functional heterogeneity among *A. fumigatus* strains. In this sense, different copies of the
"same" *Starship* in different individuals have much greater potential to differ from each other at
the sequence level compared with much smaller transposons. We found that 49.7% of COG-
annotated genes in the high-confidence *Starships* are "Poorly Categorized"; 28.8% belong to the
"Metabolism" category, 14.7% belong to "Cellular Processes and Signaling" and 6.8% belong to
"Information Storage and Processing".

Given the importance of metabolic processes for diverse fungal traits, we examined granular
classifications within the Metabolism category and found that *Starship* types differ specifically in
their contributions to metabolic heterogeneity (Fig. S9). For example, nearly all 75 copies of
Gnosis h1 carry at least 1 gene with a predicted role in "Coenzyme transport and metabolism,"
and none carry genes with predicted roles in "Lipid transport and metabolism", while the exact
opposite is true for the 119 copies of *Tardis h1*. The most frequently mobilized COG metabolism
categories are "Carbohydrate transport and metabolism" (present in 371 individual elements
belonging to 13 high-confidence *Starships*), "Secondary metabolite transport and metabolism"
(present in 329 elements belonging to 16 high-confidence *Starship* types) and "Energy production
and conversion" (present in 281 elements belonging to 9 high-confidence *Starship* types), each
of which has known associations with clinically-relevant pathogen phenotypes. We found no
evidence that any metabolism-associated COG category was enriched in *Starships* compared
with background frequencies in Af293, indicating it is likely the specific mobilized functions, as
opposed to an enrichment of certain classes of functions, that defines *Starship* contributions to
fungal phenotypes (data not shown).

While investigating associations between *Starships* and characterized pathogen phenotypes, we
found three notable examples of cargo genes that encode known traits important for pathogen
survival and virulence (Table S19, Methods) (10, 11, 36, 38–44). The BGC encoding
fumihopaside A (AFUA_5G00100-AFUA_5G00135) is carried by *Starship Gnosis h2* (Fig. 4A).
fumihopaside A is a triterpenoid glycoside that increases fungal spore survival under heat and UV
stress exposure (38). Similarly, the BGC encoding the polyketide fumigermin (AFUA_1G00970-
AFUA_1G01010), which inhibits bacterial spore germination, is carried by *Lamia h2* and *h4* (Fig.
4B) (45). Both the fumihopaside A and fumigermin BGCs were previously identified as "mobile
gene clusters" based on their presence/absence at different chromosomal locations in *A.*

fumigatus strains, and our results reveal that *Starships* are the mechanism underpinning their
mobility (Cluster 1 and 33, respectively from Lind et al., 2017) (36).

Finally, a cluster of three genes (AFUA_5G14900/*hrmA*, AFUA_5G14910/*cgnA*,
AFUA_5G14915/*bafA*, collectively referred to as the <u>h</u>rm*A*-<u>a</u>ssociated <u>c</u>luster or HAC) carried by
Nebuchadnezzar h1 increases virulence and low oxygen growth, and its expression modulates
colony level morphological changes associated with biofilm development (Fig. 4D) (33, 46). The
genes *bafB* and *bafC*, which are homologs of *bafA,* also mediate colony and submerged biofilm
morphology and are each carried by up to 5 additional *Starships*, indicating a sustained
association between this gene family and *Starships* (Fig. 4D, Fig. S10) (33). Mobilization of
biosynthetic gene clusters and biofilm-related loci by *Starships* has potential to contribute to the
rapid evolution of these traits. For example, *Starships* are known to mediate rapid gene gain and
loss through horizontal *Starship* transfer, which has recently been experimentally demonstrated
using *A. fumigatus* as a recipient (28), and *Starship* loss, which has been experimentally
demonstrated to occur through failed re-integration events (21). Thus, it is reasonable to predict
that *Starship*-mediated gain and loss effectively increases the capacity of BGC and biofilm-
associated traits to rapidly increase or decrease in frequency within populations; however, this
hypothesis remains to be tested experimentally. Minimally, we propose that by mobilizing genes
contributing to biofilm formation, *Starships* help drive heterogeneity in clinically-relevant
phenotypes at a population level (33).

To identify *Starships* that could be relevant in either environmental or clinical settings, we tested
for an association between the high-confidence *Starships* and strain isolation source (genotyping
dataset; Methods). We found that 5 high-confidence *Starships* are significantly enriched (P_{adj}
<0.05) in strains from either clinical or environmental isolation sources across the 475/479
genotyped strains for which source data exists (Table S2, Table S20, Table S21, Fig. S11). *Lamia*
h3 (P_{adj} = 0.013) and *Janus h3* (P_{adj} = 0.034) are enriched in environmental strains. *Osiris h4* (P_{adj}
= 0.022), *Nebuchadnezzar h1* (P_{adj} = 0.015; carries *bafA*, see above) and J*anus h2* (P_{adj} = 0.013)
are enriched in clinical strains, providing complementary evidence to recently published analyses
of *Starship* captain enrichment in clinical isolates across several other fungal species (47).
Although clinical and environmental strains were sampled in roughly equal proportions, we can't
completely rule out biases in our sampling (e.g. phylogenetic and ecological biases) that would
contribute to these enrichment patterns. Nevertheless, for each of the 5 enriched *Starships*,
strains were often isolated from different countries (between 3-10 countries) and often belong to
different phylogenetic groups (between 1-3 clades *sensu* Lofgren et al., 2022 and 1-6
phylogenetic clusters *sensu* Barber et al., 2021), suggesting these *Starships* are excellent
candidates to experimentally test how giant transposons contribute to fitness in environmental
and clinical settings.

Starship expression contributes to heterogeneity in a strain-, treatment- and
strain by treatment-dependent manner

The annotation of *A. fumigatus' Starship* compartment next allowed us to test the hypothesis that
Starship cargo is expressed under clinically-relevant conditions. We analyzed patterns of
differential *Starship* cargo gene expression in 14 publicly available transcriptomic studies from
three commonly used reference strains (Af293, CEA10, A1163). In total, 177 transcriptome
samples were included in this meta-analysis. Samples were split into broad treatment categories
and corrected for batch effects to allow for comparison of cargo gene expression between studies
(Methods). Out of 596 *Starship* genes total (carried by the high-confidence *Starships* present in
the three reference strains) and across all treatments, we identified 459 differentially expressed
genes (DEGs) that included both captain tyrosine recombinases and cargo (Supplementary
Data).
499
To identify experimental conditions that may impact *Starship* transposition and the subsequent
generation of *Starship*-mediated heterogeneity, we first examined patterns of transcript
abundance for the captain genes of the 12 high-confidence *Starships* in Af293, CEA10 and A1163
(Fig. 5A, Fig. S12). All captain genes have evidence for constitutive gene expression (a minimum
median transcript coverage of 1 across the gene body for biological replicates) in at least one
experimental condition (Fig. S13). Overall, 8 of the 12 captain genes from high-confidence
Starships are differentially expressed in at least one study for one or more reference strains,
revealing many opportunities for *Starship* transposition under lab conditions (Supplemental
information). Differences in experimental conditions and/or strain backgrounds have no consistent
positive or negative influence on captain gene expression, indicating that the intricacies of captain
expression remain to be elucidated.
511
In order to identify conditions where *Starships* have potential to impact phenotypes, we next
examined patterns of differential expression for *Starship*-mobilized cargo genes. Overall, we did
not find a significant enrichment of DEGs within *Starships* compared to the genomic background
of any strain (Fisher's Exact Test *P*-values > 0.05). However, the expression and significance of
Starship DEGs vary across *Starship* naves, fungal strains, and experimental conditions, indicative
of pervasive strain-specific, treatment-specific and strain by treatment interactions impacting
Starship cargo expression (Fig. 5B, Fig. S14). For example, we compared transcript abundance
patterns for genes within the *hrmA*- and *baf*-associated cluster (46) between treatment categories
and found *cgnA* and *bafA* to be down-regulated in the majority of samples assigned to infection
treatments but not in control or nutrient treatment samples (Fig. 5A, Fig. S12). Together, fungal
strain identity, treatment condition, and non-additive interactions between fungal strains and
treatments generate variation in *Starship* cargo expression.
524
To gain further insight into *Starship*-mediated heterogeneity in gene expression and regulation,
we constructed weighted gene co-expression networks (WGCNA) which visualize correlations in
gene expression. Genes are grouped into modules based on their level of shared co-expression
patterns across all samples (Methods). We found five modules as being significantly associated
with "antifungal"-, "infection"-, or "nutrient"-based studies (Fig. 5E). We identified two major trends
in the transcriptional network properties of *Starship* cargo. First, multiple *Starship*-associated
genes are integrated into modules containing many other genes from the genome, underscoring
the possibility that phenotypes emerging from these networks are the product of regulatory

interactions between *Starship* cargo and non-*Starship* genes (Fig. 5D, Fig. S15). In particular,
connections involving captain genes hint at candidate loci involved in regulatory interactions
between *Starships* and the *A. fumigatus* genome and are prime targets for future investigations
into the regulation of *Starship* transposition (Fig. 5E). Second, we also identified modules
composed of mainly *Starship* cargo, indicative of modular networks specific to particular *Starships*
(Table S24). Together, our network analyses paint a nuanced picture of how *Starships* interact
with the broader regulatory networks of the cell and implicate them in the generation of
transcriptional variation.

Relevance and Outlook

542
Phenotypic heterogeneity among fungal pathogen strains poses a major challenge for combating
infectious diseases, yet we often know little about the genetic basis of this variation. We
hypothesized that a newly discovered group of unusual transposons, the *Starships*, make
important contributions to strain heterogeneity. We tested this hypothesis by systematically
characterizing *Starship* presence/absence and expression in *A. fumigatus*, an important human
fungal pathogen of critically high research priority (1, 48). Our work provides fundamental insight
into the mode and tempo of fungal evolution by revealing that strain variation emerges not only
from well-studied genetic mechanisms (e.g., single nucleotide polymorphisms, copy number
variation, chromosomal aneuploidies, and other types of structural variants) (10, 11) but from the
previously hidden activity of giant *Starship* transposons (Fig. 6).

553
Although much remains unknown about the fundamentals of *Starship* biology, *Starships* in *A.*
fumigatus likely generate variation on a scale approaching MGE-mediated evolution in bacteria,
which is a major mechanism driving bacterial adaptation. The median number of *Starship*-borne
genes per *A. fumigatus* isolate is 0.8% (range = 0 - 2.1%), while between 2.9-4.8% of the
pangenome (corresponding to between 9.7-16% of the accessory genome) is *Starship*-
associated. These observations approach analogous measurements of bacterial plasmids in the
Enterobacteriaceae, where the median number of plasmid-borne genes per genome is estimated
at 3.3% (range = 0 - 16.5%) and where between 12.3 - 21.5% of the pangenome is plasmid-
associated (49). Given our requirements that a high-confidence *Starship* be found in 2 different
locations in 2 different strains, our estimates represent a relatively conservative assessment of
the mobile fraction of *A. fumigatus'* genome. Mobility underpins prokaryotic genome dynamics,
yet current models of fungal pathogen biology and evolution do not take gene mobility through
transposition and horizontal transfer into account. These data highlight the need to understand
the population level impacts of *Starships* across different species and to integrate these findings
into a new predictive framework for fungal biology. Our *Starship* compendium thus provides
practical insights for elucidating *A. fumigatus* biology and establishes a roadmap for deciphering
the origins of strain heterogeneity across the fungal tree of life.

Materials and Methods

Long-read genome sequencing, assembly and annotation

For DNA extraction strains were grown at 37 °C for 24 hours in shaking liquid cultures using 1% glucose minimal media (50). Biomass was collected by gravity filtration through Miracloth (Millipore). High molecular weight DNA extraction followed the Fungal CTAB DNA extraction protocol (51). Biomass was ground in liquid nitrogen and incubated at 65 °C in a lysis buffer composed of 650 µl of Buffer A (0.35 M sorbitol, 0.1 M Tris-HCl pH 9, 5 mM EDTA pH 8), 650 µl Buffer B (0.2 M Tris-HCl pH 9, 0.05 M EDTA pH8, 2M NaCl, 2% CTAB), 260 µl of Buffer C (5% Sarkosyl), 0.01% PVP, 0.2 mg proteinase K for 30 minutes. Potassium acetate (280 µl of 5M solution) was added and incubated on ice for 5 minutes. DNA was extracted by phenol:chloroform:isoamyl alcohol extraction followed by a secondary extraction with chloroform:isoamyl alcohol. The supernatant was RNAse treated (2.5µg) and precipitated using sodium-acetate (0.3M) and isopropanol. DNA was finally purified using 70% ethanol, dried and rehydrated in TE (pH 9). To preserve long strands of DNA suitable for long-read sequencing samples were gently mixed by inversion and transferred using large bore pipette tips. DNA quality was assessed using NanoDrop and concentration quantified using Qubit dsDNA BR assay kit. Finally, DNA fragment size was assessed on 1% agarose gel, looking for large quantities of DNA to remain in the well after 1 hour of running at 100v. DNA was sent to SeqCoast for Oxford Nanopore Sequencing. We used previously sequenced Illumina short read data for polishing the assemblies with Pilon (10). Two strains not previously sequenced were sequenced with Illumina at SeqCoast.

The long-read genome assemblies were assembled with Canu v2.2 and polished with five rounds of Pilon v1.24 wrapped with AAFTF v0.3.0 (52–54). Summary statistics for each assembly were computed with AAFTF and BUSCO v5.7 using eurotiomycetes_odb10 marker set (55). Genome annotations (including for the publicly available CEA10 assembly) were performed using Funannotate v1.8.10 which trained gene predictors with PASA using RNA-Seq generated for each strain (Puerner et al., forthcoming), and predicted genes de novo in each assembly and produced a consensus gene prediction set for each strain (56, 57).

Functional predictions

We annotated all predicted genes with Pfam domains, InterPro domains and GO terms using InterproScan v5.61-93.0 (-appl Pfam --iprlookup --goterms) (58). We annotated carbohydrate active enzymes using dbcan v3.0.4 (--tools all) and EggNOG orthogroups and COG categories using eggnog-mapper v2.1.7 (--sensmode very-sensitive --tax_scope Fungi)(59, 60). We annotated biosynthetic gene clusters using antiSMASH v6.0.1 (--taxon fungi --clusterhmmer --tigrfam) and 5S rDNA using infernal v1.1.4 (--rfam -E 0.001)(61, 62). On average, 35.7% of genes within *Starships* have no known PFAM or Interpro domain, GO term, COG category or eggNOG orthology (range = 0-67.6%, std. dev. = 14.8%) but approximately half of all mobilized genes could be assigned to a COG (e.g., 50% of 11,900 genes total, across 459 high-confidence elements). To investigate associations between *Starships* and pathogen phenotypes, we searched for the

611 presence of 798 genes known to be associated with virulence or infection-relevant phenotypes
612 on *Starships* (Table S19, Methods)(10, 11, 36, 38–44).

SNP analysis

We calculated pairwise kinship among strains by analyzing a previously generated VCF of all
high-quality, filtered, biallelic single nucleotide polymorphisms (SNPs) from the Lofgren et al. 2022
population with plink v1.9 (--distance square ibs flat-missing --double-id --allow-extra-chr; Fig. S3)
(63, 64). We calculated Jaccard similarity in *Starship* repertoire among all pairwise combinations
of strains using genotyping data for the set of 20 high-confidence *Starships* and the following
formula: *Starships* shared between strain 1 and 2 / (*Starships* unique to strain 1 + *Starships*
unique to strain 2 + *Starships* shared between strain 1 and 2).

Pangenome construction

We constructed a pangenome for the 13 reference-quality strains (our 11 long-read assemblies
plus the Af293 and CEA10 references) combined with the 506 publicly available short-read
assemblies (10, 11) using Orthofinder v2.5.4 (--only-groups -S diamond), and then extracted all
orthogroups with a sequence from at least 1 of the 13 reference-quality strains (Fig. 2B)(65). Core
orthologs are defined as being present in all 13 strains; accessory orthologs are present in
between 2-12 strains; singleton orthologs are present in 1 isolate. We included all 519 strains in
the Orthofinder analysis to ensure that ortholog groups are consistent across different population
subsets.

BLAST analysis

We determined the genomic locations of all best reciprocal blast hits to cargo genes carried by
the 20 high-confidence *Starships* using BLASTp v2.13 (Fig. 1C) (66). For each cargo gene, we
retrieved the highest scoring hit with ≥90% identity and ≥90% query coverage in each of the 13
reference-quality assemblies and determined whether that hit was found in the same *Starship*, a
different *Starship*, or not in a *Starship* at all. If no hit was retrieved, that gene was marked as
missing.

Within strain analysis

To assess *Starship* transposition among isolates of the same strain, we compared 6
independently sequenced datasets of CEA10 and CEA10-derived strains to the long-read CEA10
reference assembly. Genome codes and NCBI accessions are as follows: CEA10-1:
SRR7418934, CEA10-2: ERR232423, CEA10-MCH: ERR232426, A1160: SAMN28487500,
A1163: SRR068950, A1163-Cramer_SM: SRX27799936. We evaluated three additional strains
that also met the criteria of having a reference quality long-read assembly and at least 1
independently generated sequencing dataset generated by other research groups: ATCC46645,
S02-30 (=AF100-9B), and TP9 (=08-19-02-30). Short-read sequence data were mapped to our
long-read assemblies with bwa-mem2 and visually inspected in IGV to determine if any *Starships*
were deleted/transposed (67). Short-read data from select isolates with evidence of transposition

were *de novo* assembled with SPAdes and existing gene annotations were mapped from long-
read assemblies using liftoff (68, 69).

Starship annotation

We systematically annotated *Starships* in the 12 newly sequenced long-read genomes plus 507
publicly available *A. fumigatus* genomes by applying the starfish workflow v1.0.0 (default settings)
in conjunction with metaeuk v14.7e284, Mummer v4.0, CNEFinder and BLASTn (Table S2) (22,
66, 70–72). Briefly, captain tyrosine recombinase genes were *de novo* predicted with starfish's
Gene Finder Module and full-length elements associated with captains were predicted with
starfish's Element Finder Module using pairwise BLAST alignments to find empty and occupied
insertion sites. We filtered out all elements that were <15kb in length, which we have found to
correspond to indels of captain genes but never to the transposition of a full-length *Starship* (22).
We then manually examined alignments between each putative *Starship* and its corresponding
insertion site and filtered out all "low confidence" poorly supported alignments indicative of a false
positive insertion. Poor alignments were observed most often when *Starships* inserted into smaller
transposons; when an inversion breakpoint occurred within a putative *Starship*, resulting in an
over-estimation of *Starship* length; or when the insertion site was on a very small contig resulting
in small flanking region alignments.
665
We verified and supplemented these automated *Starship* predictions with manual annotations of
the 3 reference strains Af293, CEA10 and A1163, along with a set of 86 additional elements
(Table S7, Table S3). Insertion sites and DRs were manually verified by generating alignments of
the elements plus the 50kb flanks to a corresponding putative insertion site, as determined by
starfish (Table S4). MAFFT v7.4 was used to generate alignments with default parameters (73).
The DRs were visually assessed and the insertion regions were examined for the presence of
TEs, which could lead to erroneous target site, DR and TIR determinations. For *Starships* in the
three reference genomes, the set of reference *Starships* was used as a query with BLAST to
verify their start and end coordinates, and insertion sites. Additionally, the starfish output was
examined for additional elements which did not meet our initial strict cutoffs to attempt to capture
the entire *Starship* repertoire of these strains.

Starship classification

Starships were then grouped into *naves* (singular: *navis*, latin for "ship") by clustering captain
sequences in ortholog groups using mmseqs easy-cluster (--min-seq-id 0.5 -c 0.25 --alignment-
mode 3 --cov-mode 0 --cluster-reassign)(74). *Starship* sequences were then grouped into
haplotypes using the MCL clustering algorithm in conjunction with sourmash sketch (-p k=510
scaled=100 noabund) that calculated pairwise k-mer similarities over the entire sequence length,
as implemented in the commands "starfish sim" and "starfish group"(75, 76). These automated
and systematic predictions yielded a core set of 787 individual elements grouped into 54 distinct
navis-haplotype combinations.
686
We identified segregating insertion sites associated with the 787 elements across all 519 strains
using the command starfish dereplicate (--restrict --flanking 6 --mismatching 2) in conjunction with

the Orthofinder orthogroups file (see above) that was filtered to contain groups absent in at most
517 strains and present in at most 8 copies per isolate. Starfish dereplicate enables the
identification of independently segregating insertions by grouping *Starships* and their insertion
sites into homologous genomic regions using conserved combinations of orthogroups between
individuals. We genotyped the presence of each *navis-haplotype* combination within each region
for each isolate. If an isolate did not have any *Starships* within a given region, it was assigned
either an "empty" genotype (if the set of orthogroups that define the upstream region flank were
adjacent to the set of orthogroups that define the downstream region flank) or a "fragmented"
genotype (if additional orthogroups were in between the upstream and downstream region flanks).
698
To define high-confidence *Starships,* we used a threshold that required observing the same *navis-*
haplotype in ≥2 independent segregating sites in different strains, which ensured an accurate and
precise determination of each element's boundaries and the sequences therein (22). We found
18 *navis-haplotype* combinations meeting these criteria. We considered 2 additional *navis-*
haplotype combinations of interest (*Osiris h4* and *Lamia h4*) to be high-confidence after manually
annotating their boundaries and finding evidence for flanking direct repeats that are signatures of
Starship boundaries. Each high-confidence *Starship* is represented by a type element (similar in
concept to a type specimen for defining a biological species) that constitutes the longest element
assigned to that *navis-haplotype*. The type elements of the reference *Starships* range in size from
16,357-443,866 bp and collectively represent 459 individual elements with an average length of
71,515 bp (Table S4, Table S5). The remaining 328 elements have an average length of 50,249
bp and are represented by 34 "medium-confidence" *navis-haplotypes* that were only observed at
a single segregating site. We predicted the target sites of each high-confidence *Starship* by
manually aligning multiple empty insertion site sequences to each other and identifying columns
within 25bp of the core motif that had conserved nucleotides in at least 80% of sequences (Table
1).
715
Each *navis* associated with a high-confidence element was assigned a charismatic name (e.g.,
Tardis) and a new, sequentially named *haplotype* (*h1, h2*, etc.) to distinguish it from the
automatically generated *haplotype* codes (*var14, var09*, etc.). All other 34 *navis-haplotype*
combinations with evidence of only a single segregating site were classified as "medium-
confidence". Medium-confidence *Starships* with captains not belonging to the high-confidence
naves kept their automatically generated *navis* and *haplotype* codes (e.g., *navis01-var09*), while
medium-confidence *Starships* with captains belonging to the high-confidence naves were
assigned to that *navis* but kept their automatically assigned *haplotype* (e.g., *Osiris-var04*).

Starship genotyping

Starfish requires a well-resolved empty insertion site in order to annotate the boundaries of a
contiguous *Starship* element. It will therefore not annotate any element whose corresponding
empty site is missing, or any element that is partially assembled or not assembled at all. We
therefore supplemented starfish output using a combination of BLASTn and short-read mapping
to the library of reference *Starship* sequences to decrease the false negative rate for
presence/absence genotyping. First, we recovered full-length and partial elements using the set
of high-confidence *Starships* as input to the command "starfish extend". This command uses

BLASTn to align full-length *Starship* elements to the sequence downstream from all tyrosine
recombinase genes not affiliated with a full-length *Starship* element.

Separately, we downloaded publicly available paired-end short read Illumina sequencing data for
all 466 strains for which these data were available (10, 11) and mapped them to the library of 20
high-confidence *Starships* using the command "starfish coverage" and the aligner strobealign
(77), ensuring that we would correctly genotype the presence/absence of a *Starship* even if it is
not present in an assembly or partially assembled. By mapping short reads directly to a library of
known *Starship* elements, we overcame some of the drawbacks associated with working with
short-read assemblies. We considered any *Starship* with a minimum of 5 mapped reads at each
position across 95% of its entire length to be present. We considered a *Starship* to be present in
a given individual if either the main starfish workflow, BLASTn extension, or short-read mapping
identified it as present; otherwise we considered it absent.

Meta-Analysis of *A. fumigatus* RNAseq Datasets

We collected transcriptomes from a total of 434 publicly available paired-end libraries of *A.*
fumigatus strains Af293, A1163, and CEA10/CEA17 (Table S22). We surveyed the available
metadata from each BioProject and binned the samples into general categories based on the type
of treatment applied in each study. The categories that were the most well represented across
multiple BioProjects include exposure to antifungals ("antifungal"), *in vivo* and *in vitro* infection
experiments ("infection"), or supplemented growth media with specific nutrients ("nutrient"; Table
S22, Table S23).

RNAseq reads were retrieved from the NCBI SRA using fasterq-dump from the SRA toolkit (last
accessed June 26, 2024: https://github.com/ncbi/sra-tools) and were trimmed for quality and
sequencing artifacts using TrimGalore (last accessed June 26, 2024:
https://github.com/FelixKrueger/TrimGalore). Transcripts were quantified using salmon quant,
which employs a reference-free pseudo-mapping based approach to quantify transcripts. In
addition to the pseudo-BAM files created by salmon v1.10.1 (78), we created separate sets of
BAM files to assess transcript coverage across Starship genes by mapping to the transcriptome
with STAR v2.7.11a (79). We assessed transcript coverage across the core of Starship genes
using the bamsignals R package (last accessed June 26, 2024:
https://github.com/lamortenera/bamsignals. Genes with a median core-transcript coverage less
than 1 were considered to have insufficient evidence of being expressed. We applied an
abundance filter on transcript abundances, removing genes represented by fewer than 10
transcripts in at least 3 samples. We corrected for batch effects between BioProjects using
CombatSeq (80) and excluded BioProjects with inconsistent/unclear metadata, or those that
remained as outliers in the PCA after batch-correction. DESeq2 v1.45.0 (81) was used to perform
a series of differential expression tests using the corrected transcript counts. These tests were
performed individually for each BioProject, with each test comparing the differences in expression
between all control and treatment samples. Differences in treatment level were included as a
cofactor in the model, where applicable.

To summarize expression patterns across BioProjects which tested similar conditions, we
employed a random effects model (REM) using the R package metaVolcanoR (82). This method
identifies differentially expressed genes (DEGs) as genes that are significantly differentially
expressed in all studies and accounts for the variation in the expression for each DEG observed
in multiple BioProjects. The output from the meta-volcano REM are summarized log2 fold-change,
a summary p-value, and confidence intervals for each DEG. In addition, a measure of "sign-
consistency" is used to evaluate the consistency of DEG, which is expressed as a count of the
number of BioProjects where a DEG was observed with the same directional change (+/-),
centered around 0.
783
We performed weighted gene co-expression network analysis (WGCNA) using PyWGCNA v1.72-
5 (83) to construct gene co-expression networks for *A. fumigatus* reference strain Af293. A single
network was constructed using the batch-corrected TPM values from the collection of samples
belonging to "antifungal", "infection", and "nutrient" treatment categories. PyWGCNA
automatically estimates an appropriate soft-power threshold based on the lowest power for fitting
scale-free topology and identifies modules of co-expressed genes through hierarchical clustering
of the network and performing a dynamic tree cut based on 99% of the dendrogram height range.
The topological overlap matrix is then computed and a correlation matrix is constructed to produce
the final network. Distributions of TOM scores were generated based on edges in the network
between genes within *Starships*, between *Starships*, between *Starships* and the rest of the
genome, and between non-*Starship* genes in the genome. We compared these distributions using
a Wilcoxon test and anova, adjusting p-values with the Holm–Bonferroni method. Modules in the
network that were significantly associated with samples from a specific treatment category were
identified using the pairwise distances between observations using pdist from the python module
scipy (84). To highlight the genes within each module that have the most strongly correlated
expression profiles, sub-networks were constructed using only the edges 10 highest TOM scores
made between either *Starship* captain genes or any non-*Starship* gene that was present in the
module. The collections of genes within these modules were also tested for functional enrichment
using gprofiler2 (85).

Statistical tests, alignments and data visualization

Enrichment tests were performed with either the Binomial test (binom.test; alternative = "greater")
or the Fisher's exact test (fisher.test; alternative = "greater") implemented in base R, using the
p.adjust function (method = Benjamini-Hochberg) to correct for multiple comparisons. Mantel tests
for matrix correlations were performed in R (method = "pearson", permutations = 999). We
conducted nucleotide alignments among *Starships* and their associated genomic regions (e.g.
Fig. 1E, Fig. 3D) using Mummer 4.0 with the nucmer command with delta-filter options -m -l 5000
-i 95 (70). We visualized all alignments and insertion site distributions using circos and
gggenomes (86, 87) with the help of wrapper script functions implemented in the function starfish
locus-viz in the starfish package v1.1 (22). Step-by-step tutorials for *Starship* data visualization
are maintained on the starfish github repository available at
https://github.com/egluckthaler/starfish/wiki. All other figures were generated in R using ggplot2
(88).

Data availability

Sequencing reads and genomic assemblies for the 11 isolates sequenced with Oxford Nanopore have been deposited and accessioned at NCBI (Table S1). Short-read sequence data for the A1163-Cramer_SM isolate are deposited at NCBI under Bioproject PRJNA1228067. We downloaded the Af293 and CEA10 reference assemblies (NCBI accessions: GCF_000002655.1 and SAMN28487501) as well as 252 assemblies and annotations from Barber et al. 2021 from NCBI, and 256 assemblies and annotations from Lofgren et al. 2022 from the associated Zenodo repository (64). We subsequently reformatted the contig and gene identifiers to include genome codes generated as part of this study (Table S2). All scripts (including raw data for figure generation), genomic data used for this study, and results from differential expression tests are available through the following Figshare repository DOI: 10.6084/m9.figshare.26049703.

Acknowledgements

We thank Jacob Steenwyk for insightful feedback on an earlier draft of this manuscript.

Funding

This work was supported by the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin-Madison with funding from the Wisconsin Alumni Research Foundation to EGT; by funding from start-up funds from the Department of Plant Pathology at the University of Wisconsin-Madison and from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement to EGT (grant number 890630); by the Swedish Research Council Formas (grant number 2019-01227) and the Swedish Research Council VR (grant number 2021-04290) to AAV. JES, CP, and RAC are supported by funding from the National Institutes of Health (grant no. 2R01AI130128), to RAC. JES is a CIFAR Fellow in the program Fungal Kingdom: Threats and Opportunities. Analyses and data management at UC Riverside was performed on the High-Performance Computing Center at the University of California, Riverside, in the Institute of Integrative Genome Biology, supported by grants from NSF (DBI-2215705) and NIH (S10-OD016290).

References

1. 2022. WHO fungal priority pathogens list to guide research, development and public health action. World Health Organization, Geneva.

2. Denning DW. 2024. Global incidence and mortality of severe fungal disease. Lancet Infect Dis S1473-3099(23)00692–8.

3. Latgé J-P, Chamilos G. 2019. Aspergillus fumigatus and Aspergillosis in 2019. Clin Microbiol Rev 33:10.1128/cmr.00140-18.

4. Lin S-J, Schranz J, Teutsch SM. 2001. Aspergillosis Case-Fatality Rate: Systematic Review of the Literature. Clin Infect Dis 32:358–366.

5. Kowalski CH, Beattie SR, Fuller KK, McGurk EA, Tang Y-W, Hohl TM, Obar JJ, Cramer RA. 2016. Heterogeneity among Isolates Reveals that Fitness in Low Oxygen Correlates with Aspergillus fumigatus Virulence. mBio 7:e01515-16.

6. Colabardini AC, Wang F, Dong Z, Pardeshi L, Rocha MC, Costa JH, dos Reis TF, Brown A, Jaber QZ, Fridman M, Fill T, Rokas A, Malavazi I, Wong KH, Goldman GH. 2022. Heterogeneity in the transcriptional response of the human pathogen Aspergillus fumigatus to the antifungal agent caspofungin. Genetics 220:iyab183.

7. Horta MAC, Steenwyk JL, Mead ME, dos Santos LHB, Zhao S, Gibbons JG, Marcet-Houben M, Gabaldón T, Rokas A, Goldman GH. 2022. Examination of Genome-Wide Ortholog Variation in Clinical and Environmental Isolates of the Fungal Pathogen Aspergillus fumigatus. mBio 13:e01519-22.

8. Keller NP. 2017. Heterogeneity Confounds Establishment of "a" Model Microbial Strain. mBio 8:10.1128/mbio.00135-17.

9. Steenwyk JL, Rokas A, Goldman GH. 2023. Know the enemy and know yourself: Addressing cryptic fungal pathogens of humans and beyond. PLoS Pathog 19:e1011704.

10. Lofgren LA, Ross BS, Cramer RA, Stajich JE. 2022. The pan-genome of Aspergillus fumigatus provides a high-resolution view of its population structure revealing high levels of lineage-specific diversity driven by recombination. PLOS Biol 20:e3001890.

11. Barber AE, Sae-Ong T, Kang K, Seelbinder B, Li J, Walther G, Panagiotou G, Kurzai O. 2021. Aspergillus fumigatus pan-genome analysis identifies genetic variants associated with human infection. 12. Nat Microbiol 6:1526–1536.

12. Rhodes J, Abdolrasouli A, Dunne K, Sewell TR, Zhang Y, Ballard E, Brackin AP, van Rhijn N, Chown H, Tsitsopoulou A, Posso RB, Chotirmall SH, McElvaney NG, Murphy PG, Talento AF, Renwick J, Dyer PS, Szekely A, Bowyer P, Bromley MJ, Johnson EM, Lewis White P, Warris A, Barton RC, Schelenz S, Rogers TR, Armstrong-James D, Fisher MC. 2022. Population genomics confirms acquisition of drug-resistant Aspergillus fumigatus infection by humans from the environment. Nat Microbiol 7:663–674.

13. Hall JPJ, Harrison E, Baltrus DA. 2022. Introduction: the secret lives of microbial mobile genetic elements. Philos Trans R Soc B Biol Sci 377:20200460.

14. Aubert D, Naas T, Héritier C, Poirel L, Nordmann P. 2006. Functional characterization of IS1999, an IS4 family element involved in mobilization and expression of beta-lactam resistance genes. J Bacteriol 188:6506–6514.

15. Krishnan P, Meile L, Plissonneau C, Ma X, Hartmann FE, Croll D, McDonald BA, Sánchez-Vallet A. 2018. Transposable element insertions shape gene regulation and melanin production in a fungal pathogen of wheat. BMC Biol 16:78.

16. Hottes AK, Freddolino PL, Khare A, Donnell ZN, Liu JC, Tavazoie S. 2013. Bacterial Adaptation through Loss of Function. PLOS Genet 9:e1003617.

17. Weisberg AJ, Chang JH. 2023. Mobile Genetic Element Flexibility as an Underlying Principle to Bacterial Evolution. Annu Rev Microbiol 77:null.

18. Baquero F. 2004. From pieces to patterns: evolutionary engineering in bacterial pathogens. Nat Rev Microbiol 2:510–518.

893   19.   Benler S, Faure G, Altae-Tran H, Shmakov S, Zhang F, Koonin E. 2021. Cargo Genes of
894   Tn7-Like Transposons Comprise an Enormous Diversity of Defense Systems, Mobile
895   Genetic Elements, and Antibiotic Resistance Genes. mBio 12:e02938-21.
896   20.   Gluck-Thaler E, Ralston T, Konkel Z, Ocampos CG, Ganeshan VD, Dorrance AE, Niblack
897   TL, Wood CW, Slot JC, Lopez-Nicora HD, Vogan AA. 2022. Giant Starship Elements
898   Mobilize Accessory Genes in Fungal Genomes. Mol Biol Evol 39:msac109.
899   21.   Urquhart AS, Vogan AA, Gardiner DM, Idnurm A. 2023. Starships are active eukaryotic
900   transposable elements mobilized by a new family of tyrosine recombinases. Proc Natl
901   Acad Sci U S A 120:e2214521120.
902   22.   Gluck-Thaler E, Vogan AA. 2024. Systematic identification of cargo-mobilizing genetic
903   elements reveals new dimensions of eukaryotic diversity. Nucleic Acids Res gkae327.
904   23.   Urquhart A, Vogan AA, Gluck-Thaler E. 2024. Starships: a new frontier for fungal biology.
905   Trends Genet 40:1060–1073.
906   24.   Gourlie R, McDonald M, Hafez M, Ortega-Polo R, Low KE, Abbott DW, Strelkov SE, Daayf
907   F, Aboukhaddour R. 2022. The pangenome of the wheat pathogen Pyrenophora tritici-
908   repentis reveals novel transposons associated with necrotrophic effectors ToxA and ToxB.
909   BMC Biol 20:239.
910   25.   Urquhart AS, Chong NF, Yang Y, Idnurm A. 2022. A large transposable element mediates
911   metal resistance in the fungus Paecilomyces variotii. Curr Biol 32:937-950.e5.
912   26.   Urquhart AS, Gluck-Thaler E, Vogan AA. 2023. Gene acquisition by giant transposons
913   primes eukaryotes for rapid evolution via horizontal gene transfer. bioRxiv
914   https://doi.org/10.1101/2023.11.22.568313.
915   27.   Bucknell A, Wilson HM, Santos KCG do, Simpfendorfer S, Milgate A, Germain H, Solomon
916   PS, Bentham A, McDonald MC. 2024. Sanctuary: A Starship transposon facilitating the
917   movement of the virulence factor ToxA in fungal wheat pathogens. bioRxiv
918   https://doi.org/10.1101/2024.03.04.583430.
919   28.   Urquhart AS, O'Donnell S, Gluck-Thaler E, Vogan AA. 2025. A natural mechanism of
920   eukaryotic horizontal gene transfer. bioRxiv https://doi.org/10.1101/2025.02.28.640899.
921   29.   Pain A, Woodward J, Quail MA, Anderson MJ, Clark R, Collins M, Fosker N, Fraser A,
922   Harris D, Larke N, Murphy L, Humphray S, O'Neil S, Pertea M, Price C, Rabbinowitsch E,
923   Rajandream M-A, Salzberg S, Saunders D, Seeger K, Sharp S, Warren T, Denning DW,
924   Barrell B, Hall N. 2004. Insight into the genome of Aspergillus fumigatus: analysis of a 922
925   kb region encompassing the nitrate assimilation gene cluster. Fungal Genet Biol FG B
926   41:443–453.
927   30.   Monod M, Paris S, Sarfati J, Jaton-Ogay K, Ave P, Latgé JP. 1993. Virulence of alkaline
928   protease-deficient mutants of Aspergillus fumigatus. FEMS Microbiol Lett 106:39–46.
929   31.   Bowyer P, Currin A, Delneri D, Fraczek MG. 2022. Telomere-to-telomere genome
930   sequence of the model mould pathogen Aspergillus fumigatus. Nat Commun 13:5394.
931   32.   Bertuzzi M, van Rhijn N, Krappmann S, Bowyer P, Bromley MJ, Bignell EM. 2021. On the
932   lineage of Aspergillus fumigatus isolates in common laboratory use. Med Mycol 59:7–13.
933   33.   Kowalski CH, Morelli KA, Stajich JE, Nadell CD, Cramer RA. 2021. A Heterogeneously
934   Expressed Gene Family Modulates the Biofilm Architecture and Hypoxic Growth of
935   Aspergillus fumigatus. mBio 12:e03579-20.
936   34.   Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P,
937   Morgante M, Panaud O, Paux E, SanMiguel P, Schulman AH. 2007. A unified classification
938   system for eukaryotic transposable elements. Nat Rev Genet 8:973–982.
939   35.   Lambert SA, Yang AWH, Sasse A, Cowley G, Albu M, Caddick MX, Morris QD, Weirauch
940   MT, Hughes TR. 2019. Similarity regression predicts evolution of transcription factor
941   sequence specificity. Nat Genet 51:981–989.
942   36.   Lind AL, Wisecaver JH, Lameiras C, Wiemann P, Palmer JM, Keller NP, Rodrigues F,
943   Goldman GH, Rokas A. 2017. Drivers of genetic diversity in secondary metabolic gene

clusters within a fungal species. PLoS Biol 15:e2003583.

37. Abdolrasouli A, Rhodes J, Beale MA, Hagen F, Rogers TR, Chowdhary A, Meis JF,
Armstrong-James D, Fisher MC. 2015. Genomic Context of Azole Resistance Mutations in
Aspergillus fumigatus Determined Using Whole-Genome Sequencing. mBio 6:e00536-15.

38. Ma K, Zhang P, Tao Q, Keller NP, Yang Y, Yin W-B, Liu H. 2019. Characterization and
Biosynthesis of a Rare Fungal Hopane-Type Triterpenoid Glycoside Involved in the
Antistress Property of Aspergillus fumigatus. Org Lett 21:3252–3256.

39. Brown NA, Goldman GH. 2016. The contribution of Aspergillus fumigatus stress responses
to virulence and antifungal resistance. J Microbiol 54:243–253.

40. Steenwyk JL, Mead ME, de Castro PA, Valero C, Damasio A, dos Santos RAC, Labella
AL, Li Y, Knowles SL, Raja HA, Oberlies NH, Zhou X, Cornely OA, Fuchs F, Koehler P,
Goldman GH, Rokas A. 2021. Genomic and Phenotypic Analysis of COVID-19-Associated
Pulmonary Aspergillosis Isolates of Aspergillus fumigatus. Microbiol Spectr
9:10.1128/spectrum.00010-21.

41. Raffa N, Osherov N, Keller NP. 2019. Copper Utilization, Regulation, and Acquisition by
Aspergillus fumigatus. 8. Int J Mol Sci 20:1980.

42. Bignell E, Cairns TC, Throckmorton K, Nierman WC, Keller NP. 2016. Secondary
metabolite arsenal of an opportunistic pathogenic fungus. Philos Trans R Soc B Biol Sci
371:20160023.

43. Abad A, Fernández-Molina JV, Bikandi J, Ramírez A, Margareto J, Sendino J, Hernando
FL, Pontón J, Garaizar J, Rementeria A. 2010. What makes Aspergillus fumigatus a
successful pathogen? Genes and molecules involved in invasive aspergillosis. Rev
Iberoam Micol 27:155–182.

44. Auxier B, Zhang J, Marquez FR, Senden K, Heuvel J van den, Aanen DK, Snelders E,
Debets AJM. 2022. Identification of heterokaryon incompatibility genes in Aspergillus
fumigatus highlights a narrow footprint of ancient balancing selection. bioRxiv
https://doi.org/10.1101/2022.11.25.517501.

45. Stroe MC, Netzker T, Scherlach K, Krüger T, Hertweck C, Valiante V, Brakhage AA. 2020.
Targeted induction of a silent fungal gene cluster encoding the bacteria-specific
germination inhibitor fumigermin. eLife 9:e52541.

46. Kowalski CH, Kerkaert JD, Liu K-W, Bond MC, Hartmann R, Nadell CD, Stajich JE,
Cramer RA. 2019. Fungal biofilm morphology impacts hypoxia fitness and disease
progression. Nat Microbiol 4:2430–2441.

47. O'Donnell S, Rezende G, Vernadet J-P, Snirc A, Ropars J. 2024. Harbouring Starships:
The accumulation of large Horizontal Gene Transfers in Domesticated and Pathogenic
Fungi. bioRxiv https://doi.org/10.1101/2024.07.03.601904.

48. Rokas A, Mead ME, Steenwyk JL, Oberlies NH, Goldman GH. 2020. Evolving moldy
murderers: Aspergillus section Fumigati as a model for studying the repeated evolution of
fungal pathogenicity. PLOS Pathog 16:e1008315.

49. Shaw LP, Chau KK, Kavanagh J, AbuOun M, Stubberfield E, Gweon HS, Barker L, Rodger
G, Bowes MJ, Hubbard ATM, Pickford H, Swann J, Gilson D, Smith RP, Hoosdally SJ,
Sebra R, Brett H, Peto TEA, Bailey MJ, Crook DW, Read DS, Anjum MF, Walker AS,
Stoesser N, REHAB CONSORTIUM. 2021. Niche and local geography shape the
pangenome of wastewater- and livestock-associated Enterobacteriaceae. Sci Adv
7:eabe3868.

50. Shimizu K, Keller NP. 2001. Genetic involvement of a cAMP-dependent protein kinase in a
G protein signaling pathway regulating morphological and chemical transitions in
Aspergillus nidulans. Genetics 157:591–600.

51. Carter-House D, Stajich J, Unruh S, Kurbessoian T. 2020. Fungal CTAB DNA Extraction.

52. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.

Genome Res 27:722–736.

53.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,
Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive
microbial variant detection and genome assembly improvement. PloS One 9:e112963.

54.   Palmer JM, Stajich JE. 2022. Automatic assembly for the fungi (AAFTF): genome
assembly pipeline (v0.4.1). Zenodo.

55.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for
Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol 38:4647–4654.

56.   Palmer JM, Stajich J. 2020. Funannotate v1.8.1: Eukaryotic genome annotation (v1.8.1).
Zenodo.

57.   Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith RK, Hannick LI, Maiti R, Ronning
CM, Rusch DB, Town CD, Salzberg SL, White O. 2003. Improving the Arabidopsis
genome annotation using maximal transcript alignment assemblies. Nucleic Acids Res
31:5654–5666.

58.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-
Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30:1236–1240.

59.   Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-
mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the
Metagenomic Scale. Mol Biol Evol https://doi.org/10.1093/molbev/msab293.

60.   Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y. 2023. dbCAN3: automated
carbohydrate-active enzyme and substrate annotation. Nucleic Acids Res 51:W115–W121.

61.   Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, van Wezel GP, Medema MH, Weber
T. 2021. antiSMASH 6.0: improving cluster detection and comparison capabilities. Nucleic
Acids Res 49:W29–W35.

62.   Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics 29:2933–2935.

63.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P,
de Bakker PIW, Daly MJ, Sham PC. 2007. PLINK: A Tool Set for Whole-Genome
Association and Population-Based Linkage Analyses. Am J Hum Genet 81:559–575.

64.   Lofgren LA, Ross B, Cramer Jr RA, Stajich JE. 2021. Aspergillus fumigatus pangenome
dataset (v2021-12-12). Zenodo.

65.   Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol 16:157.

66.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. BMC Bioinformatics 10:421.

67.   Vasimuddin Md, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of
BWA-MEM for Multicore Systems, p. 314–324. *In* 2019 IEEE International Parallel and
Distributed Processing Symposium (IPDPS).

68.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De
Novo Assembler. Curr Protoc Bioinforma 70:e102.

69.   Shumate A, Salzberg SL. 2021. Liftoff: accurate mapping of gene annotations.
Bioinformatics 37:1639–1643.

70.   Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A
fast and versatile genome alignment system. PLOS Comput Biol 14:e1005944.

71.   Levy Karin E, Mirdita M, Söding J. 2020. MetaEuk—sensitive, high-throughput gene
discovery, and annotation for large-scale eukaryotic metagenomics. Microbiome 8:48.

72.   Ayad LAK, Pissis SP, Polychronopoulos D. 2018. CNEFinder: finding conserved non-
coding elements in genomes. Bioinforma Oxf Engl 34:i743–i747.

73. Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol 30:772–780.
74. Hauser M, Steinegger M, Söding J. 2016. MMseqs software suite for fast and deep
clustering and searching of large protein sequence sets. Bioinformatics 32:1323–1330.
75. Brown CT, Irber L. 2016. sourmash: a library for MinHash sketching of DNA. J Open
Source Softw 1:27.
76. Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale
detection of protein families. Nucleic Acids Res 30:1575–1584.
77. Sahlin K. 2022. Strobealign: flexible seed size enables ultra-fast and accurate read
alignment. Genome Biol 23:260.
78. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods 14:417–419.
79. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.
80. Zhang Y, Parmigiani G, Johnson WE. 2020. ComBat-seq: batch effect adjustment for
RNA-seq count data. NAR Genomics Bioinforma 2:lqaa078.
81. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol 15:550.
82. MetaVolcanoR. Bioconductor. http://bioconductor.org/packages/MetaVolcanoR/. Retrieved
26 June 2024.
83. Rezaie N, Reese F, Mortazavi A. 2023. PyWGCNA: a Python package for weighted gene
co-expression network analysis. Bioinformatics 39:btad415.
84. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,
Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ,
Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore
EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris
CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P. 2020. SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nat Methods 17:261–272.
85. Kolberg L, Raudvere U, Kuzmin I, Vilo J, Peterson H. 2020. gprofiler2 -- an R package for
gene list functional enrichment analysis and namespace conversion toolset g:Profiler.
F1000Research 9:ELIXIR-709.
86. Krzywinski MI, Schein JE, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra
MA. 2009. Circos: An information aesthetic for comparative genomics. Genome Res
https://doi.org/10.1101/gr.092759.109.
87. Hackl T, Ankenbrand M, van Adrichem B. 2023. gggenomes: A Grammar of Graphics for
Comparative Genomics (R package version 0.9.9.9000).
88. Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis, 3rd ed. Springer
International Publishing, Cham. http://link.springer.com/10.1007/978-3-319-24277-4.
Retrieved 10 October 2023.
89. McInnes L, Healy J, Saul N, Großberger L. 2018. UMAP: Uniform Manifold Approximation
and Projection. J Open Source Softw 3:861.

1087

Figure legends

Figure 1: At least 20 distinct *Starships* carrying hundreds of protein-coding genes vary in their presence/absence across *Aspergillus fumigatus* strains. A) Top: a SNP-based maximum likelihood tree of 220 *A. fumigatus* strains from Lofgren et al. 2022 (visualized as a cladogram for clarity), color-coded according to phylogenetic clade as defined by Lofgren et al. 2022 (10). Bottom: A heatmap depicting the presence (gray) and absence (white) of 20 high-confidence *Starships* in each isolate. B) Schematics and sequence alignments of the type elements from 20 high-confidence *Starships*, where links between schematics represent alignable regions ≥500bp and ≥80% nucleotide sequence identity and arrows represent predicted coding sequences. C) A donut chart summarizing the number of distinct types of *Starships* by the quality of their prediction. D) A Venn diagram indicating the number of shared and unique high-confidence *Starships* in the reference strains Af293 and CEA10 (Table S7). E) Pairwise genome alignments between the CEA10 isolate from BioSample SAMN28487501 and the CEA10-derived A1163 isolate sequenced in this study demonstrating *Starship* movement. *Nebuchadnezzar h1* is present on Chromosome 6 in CEA10 (CP097568.1: 3645813-3698867) and absent at the corresponding locus in A1163 (NODE10: 188873-188877). Conversely, *Nebuchadnezzar h1* is present on Chromosome 4 in A1163 (NODE6: 77737-130777) but absent from the corresponding locus in CEA10 (CP097566.1:1344957-1344961).

Figure 2: *Starships* are enriched in accessory genes whose presence/absence and genomic location vary across *Aspergillus fumigatus* strains. All panels visualize data derived from 13 reference-quality *A. fumigatus* assemblies and the 20 high-confidence *Starships* (n = 459 elements total). A) Box-and-whisker plots summarizing the total percentage of nucleotide sequence and predicted genes carried by *Starships* per genome. B) Donut charts summarizing the percentages of gene orthogroups in the core, accessory, singleton, and *Starship*-associated compartments of the *A. fumigatus* pangenome (Table S10). C) Iceberg plots summarizing the genomic locations of the single best BLASTp hits (≥90% identity, ≥33% query coverage) to the cargo genes from the type elements of all 20 high-confidence *Starships* (left) and two individual *Starships* (right; Table S12). Each column in the iceberg plot represents a cargo gene and is color-coded according to the genomic location of hits (full dataset in Fig. S5).

Figure 3: *Starships* and their insertion sites are distributed across all major chromosomes in the *Aspergillus fumigatus* genome. A) A Circos plot summarizing all inserted *Starships* (in red) and all genomic regions containing either an empty insertion site or a fragmented *Starship* (in black, along perimeter and labeled with the *r* prefix) in the 8 chromosomes of the *A. fumigatus* reference strain Af293 (Table S14). All genomic regions contain a *Starship* insertion in some other individual from the 519 strain population. B) Barcharts summarizing the genotypes of segregating genomic regions associated with the four most active high-confidence elements in the 519 strain population (Table S16; full dataset in Fig. S7). If an isolate did not have any *Starships* within a given region, it was assigned either an "empty" or "fragmented" genotype (Methods). C) A scatterplot summarizing the relationship between the number of genomic regions containing a given *Starship* and the total number of copies of that *Starship* in the 519 strain population, where each point represents one of the 20 high-confidence *Starships.* A line derived from a linear regression model

1131   is superimposed, with shaded 95% confidence intervals drawn in gray. D) Alignments of *Tardis*
1132   *h1* copies +/- 50kb of flanking sequence across 14 genomic regions. Links between schematics
1133   represent alignable regions ≥1000bp and ≥90% nucleotide sequence identity.

1134

Figure 4: *Starships* mobilize adaptive traits and generate allelic diversity among *Aspergillus*
fumigatus strains. Schematics and alignments of: A) *Starship Lamia h2* and *h4*, which carry the
biosynthetic gene cluster (BGC) encoding the polyketide secondary metabolite fumigermin (45),
shown inserted at 3 independent sites. B) *Starship Gnosis h2* carrying the BGC encoding the
terpene secondary metabolite fumihopaside A (38), shown inserted at 2 independent sites. C) a
large region on Chromosome 3 previously identified as an idiomorphic BGC (36) containing
multiple segregating *Starship* insertions and various combinations of putative BGCs, including a
a non-reducing polyketide synthase (NR-PKS) BGC carried by *Starship Osiris h4*. *Starships* from
the *Osiris navis* specifically insert in 5S rDNA sequence and are predicted to fragment it. D) Eight
Starships in the reference strains Af293 and CEA10 that all carry homologs of *biofilm architecture*
factor A (*bafA*)(33). *Starship Nebuchadnezzar h1 (Neb. h1)* carries *bafA* as part of the H_AAC
(*hrmA*-associated gene cluster), while *Starship navis10-var35* carries *bafB* as part of H_BAC and
Starship Osiris h3 carries *bafC* as part of H_CAC (33). Only a portion of *Starship Lamia h4* is
visualized for figure legibility. All data for A-D was collected from the 519 *A. fumigatus* strain
population (Table S5). Links between schematics represent alignable regions ≥5000bp and ≥95%
nucleotide sequence identity and arrows represent predicted coding sequences. Abbreviations:
polyketide synthase (PKS); non-ribosomal peptide synthetase (NRPS); highly reducing (HR).

1152

Figure 5: A) A heatmap of transcript abundances (log_{10}TPM) of *Starship* captain tyrosine
recombinase genes (top) and genes within the *hrmA*-associated cluster (HAC; bottom), collapsed
across treatment replicates, for 11 RNAseq studies from *A. fumigatus* Af293. B) Results from
differential expression tests for select *Starships*, treatment categories, and strain combinations.
Differentially expressed genes (DEGs) based on a single study are shown as log_2 fold-change
(log_2FC) in black with standard error bars, whereas DEGs identified across multiple studies are
represented with summarized log_2FC values from a random effects model (REM) and coloured
by "sign-consistency", the number of studies that reported DEGs in the positive (+1) or negative
(-1) direction, centered around 0. Labels in bold font represent DEGs that are significantly
differentially expressed in more than one study. C) The results of a weighted gene co-expression
network analysis (WGCNA) constructed from 14 "antifungal", "infection", and "nutrient" RNAseq
studies represented using UMAP clustering (89) based on the co-expression eigengene values
for all genes in the *A. fumigatus* Af293 genome. Non-*Starship* genes present within modules of
interest are shown in blue, while all *Starship* genes are shown in red, with those genes within
modules of interest having a black outline. D) Genes within modules that were significantly
associated with samples from specific treatment categories or studies were used to construct sub-
networks which contained the top 10 edges in the network that were made between any pair of
genes or any gene and a *Starship* captain. The connections between genes (edges) are based
on the topological overlap matrix (TOM) for each module, and have been 0-1 scaled. E) Boxplots
of eigengene values from WGCNA, akin to a weighted average expression profile, indicate the
extent of co-expression (correlation) of the genes present within each module. Pairwise

comparisons of module eigengene values determined if a module was significantly associated
with samples from specific treatment categories or studies (Fig. S16).

1176

1177 Figure 6: Contributions of *Starships* to strain heterogeneity in *Aspergillus fumigatus*.

1178

Supplemental figure legends

Figure S1: Counts of starfish-predicted *Starships* per *Aspergillus fumigatus* strain (n = 519), broken up by assembly project (either the 12 Oxford Nanopore Assemblies generated by study plus the AF293 reference genome, or from Lofgren et al 2022 or Barber et al 2021; Table S2). A) Counts of *Starships* with either 'insert' or 'flank' boundaries, which are derived directly from pairwise genome alignments against a putative insertion site. B) Counts of *Starships* with 'insert' or 'flank' boundaries plus those with 'extend' boundaries, which are derived from aligning genomic sequences to known *Starship* sequences. Insert and flank boundaries are associated with full-length *Starship* elements, while "extend" boundaries may be associated with either full-length *Starship* elements or element fragments.

Figure S2: Nucleotide alignments of representative *Starship* copies, +/- 100kb of flanking sequence, across all genomic regions where that *Starship* is found. Predicted functions of interest are annotated above the corresponding gene. Links between schematics represent alignable regions ≥1000bp and ≥90% nucleotide sequence identity. A) *Tardis*. B) *Gnosis*. C) *Janus*. D) *Osiris*. E) *Lamia*. F) *Logos*. G) *Nebuchadnezzar.*

Figure S3: A scatterplot depicting pairwise comparisons of single nucleotide polymorphism (SNP)-based Identity by State (IBS) and Jaccard similarity in high-confidence Starship presence/absence profiles between 151 Clade 1 strains from Lofgren et al. 2022 (Table S8).

Figure S4: Screenshots from the IGV genome browser showing deletions of *Nebuchadnezzar h1* among different isolates of the ATCC46645 strain sequenced by different research groups (regions depicted in red denote *Nebuchadnezzar h1*). Illumina short-reads of strain ATCC46645 sequenced by a different research group (accession: SRR7418935) mapped to the ATCC46645 genome sequenced with long read technology in this study. A zoom in of the genomic location of *Nebuchadnezzar h1* in the ATCC46645 long read assembly is shown (contig accession: ATCC46645-lr_scaffold15). Three short-read tracks indicate a deletion of the *Starship* (above). Note that more short reads are mapped than shown in the image. Track and color descriptions can be found in the IGV manual.

Figure S5: Donut charts summarizing the percentages of gene orthogroups in the core, accessory, singleton, and *Starship*-associated compartments of the *Aspergillus fumigatus* pangenome, derived from the 13 reference-quality *A. fumigatus* strains and the expanded set of 54 high and medium-confidence *Starships* (n = 1818 elements total; Table S11)*.*

Figure S6: Iceberg plots summarizing the genomic locations of the single best BLASTp hits (≥90% identity, ≥33% query coverage) to the cargo genes from the type elements of the 20 high-confidence *Starships* in 519 *Aspergillus fumigatus* assemblies (Table S13). Each column represents a cargo gene. A) Results broken down by *Starship*, with columns arranged according to gene order within each *Starship*. B) Compiled results across all 20 *Starships*, with columns arranged according to the number of strains with BLASTp hits. Bars are colored according to the genomic location in which the BLASTp hits are found.

1221

Figure S7: Barcharts summarizing the genotypes of segregating genomic regions associated with
the 20 high-confidence elements in the *Aspergillus fumigatus* 519 strain population (Table S16).
If an isolate did not have any Starships within a given region, it was assigned either an "empty"
or "fragmented" genotype (Methods).

Figure S8: The putative target site of *Starship Tardis* (indicated with an arrow) occurs more often
than you would expect by chance in the *Aspergillus fumigatus* genome. We estimated the copy
numbers of all k-mers of length 10 in the Af293 reference genome and found that the k-mer of
length 10 that corresponds to the putative target site of *Tardis* is present in high copy numbers
that exceed the expected genome-wide frequency of k-mers of this length (>99.99th percentile).

1231

1232 Figure S9: *Starships* mobilize genes encoding diverse metabolic functions. Barcharts
1233 summarizing the presence of genes with metabolism-related COG (Clusters of Orthologous
1234 Groups) annotations in the 20 high-confidence *Starships* in the 519 *Aspergillus fumigatus* strain
1235 population (Table S18). The X-axis measures the percentage of copies of a given *Starship* that
1236 carry at least 1 gene with an annotation of interest.

1237

Figure S10: The Biofilm Architecture Factor (*baf*) gene family is closely associated with diverse
Starships in *Aspergillus fumigatus*. A maximum likelihood tree of *baf* sequences from the 519
Aspergillus fumigatus strain population. Branches with ≥80% SH-ALRT and ≥95% ultrafast
bootstrap support are in bold. *Baf* sequences found in *Starships* have the corresponding *Starship*
identification number appended to their right (Table S5). Sequences are color-coded according
to 6 corresponding *baf* clades of interest, and a summary of all the *Starship* types found
associated with each clade is printed on the right.

1245

Figure S11: *Starships* are enriched in environmental and clinical strains. Barcharts summarizing
the proportion of strains from 475/479 *Aspergillus fumigatus* genotyped strains with known
isolation sources that have the 20 high-confidence *Starships*, broken up by isolation source.
Fisher's exact test P values for *Starship* enrichment across isolation source categories are shown
above each bar (adjusted for multiple comparisons using the Benjamini Hochberg procedure;
Table S20).

1252

1253 Figure S12: A heatmap of transcript abundances ($\log_{10}$ TPM) of *Starship* captain genes (A, C)
1254 and genes within the *hrmA*-associated cluster (HAC; B, D), collapsed across treatment replicates,
1255 for RNAseq studies from *A. fumigatus* CEA10 (A, B) and A1163 (C, D).

1256

1257 Figure S13: Heatmap of binary core transcript coverage across genes in *Starships* in strains
1258 Af293 (A), CEA10 (B) and A1163 (C). Genes that have a median core transcript coverage greater
1259 than 1 are shown here in blue, and values below 1 are shown in red.

1260

Figure S14: Differentially expressed *Starship*-mobilized genes (DEGs) displayed in volcano plots
across combinations of *A. fumigatus* strains and treatment categories. Differential expression
based on singleton studies are shown as $\log_2$ fold-change ($\log_2$FC) in black with standard error

bars, whereas DEGs identified across multiple studies are represented with summarized log$_2$FC
values from a random effects model (REM) and coloured by values "sign-consistency", the
number of studies that reported differential gene expression in the positive (+1) or negative (-1)
direction, centered around 0.

1268

Figure S15: Distributions of TOM scores from the WGCNA constructed from Af293 *A. fumigatus*
studies. TOM scores were separated into distinct categories for edges that connected any two
genes within a *Starships*, between different *Starships*, between *Starships* and the rest of the
genome, or between any two non-*Starship* genes in the genome were compared. These
distributions were compared using a Wilcoxon test (p > 0.05 = "ns"; p < 0.05 = "*", p < 0.01 = "**",
p < 0.001 = "***", p < 0.0001 = "****").

1275

1276   Figure S16: Heatmap displaying the correlation values for eigengenes in each module and their
1277   association with samples from a specific study (labeled with BioProject IDs), or different levels of
1278   experimental treatment category across studies. Cells of the heatmap are coloured by the
1279   correlation value, as well as labeled in each cell. P-values from a pairwise comparison between
1280   observations (*pdist*) are shown in parentheses.

1281

Supplemental table legends

Table S1: Genome assembly statistics for the 12 strains sequenced with Oxford Nanopore long-read technology

Table S2: Metadata for publically available *Aspergillus fumigatus* strains

Table S3: Manual annotation data of 86 Starship elements

Table S4: Metadata for all starfish-predicted Starships

Table S5: Sequence features of all starfish-predicted Starships in the 519 strain population

Table S6: Frequencies of the 20 high-confidence Starships in the 519 strain population using the expanded dataset

Table S7: Manual annotation of Starship elements in three *Aspergillus fumigatus* reference strains

Table S8: Pairwise comparisons between strains of SNP identity by state and Starship repertoire similarity

Table S9: Comparison of Starship coordinates with structural variants identified by Colabardini et al 2022 (doi: 10.1371/journal.pgen.1010001)

Table S10: Orthogroup frequencies in the 519 strain population

Table S11: Orthogroup frequencies in the 13 reference-quality strains

Table S12: BLAST recovery of Starship cargo genes from the 13 reference-quality strains

Table S13: BLAST recovery of Starship cargo genes from the 519 strain population

Table S14: Genotyping of genomic regions with segregating Starship insertions

Table S15: Starships in genotyped genomic regions with segregating Starship insertions

Table S16: Summary of genotyping data for genomic regions with segregating Starship insertions

Table S17: Coordinates of predicted 5s rDNA sequences in the 519 strain population

Table S18: Proportion of elements carrying at least one gene annotated with various COG categories

1325   Table S19: List of putative and published virulence and stress resistance genotypes in *Aspergillus*
1326   *fumigatus*
1327
1328   Table S20: Fisher's exact test statistics for Starship enrichment by strain isolation source
1329
1330   Table S21: Presence/absence genotyping data for the 20 high-confidence Starships in the 519
1331   strain population using the genotyping dataset
1332
1333   Table S22: Metadata from RNASeq studies collected from NCBI used in the meta-analyses of
1334   *Starship* gene expression.
1335
1336   Table S23: Summary of RNASeq studies collected from NCBI used in the meta-analyses of
1337   *Starship* gene expression.
1338
1339   Table S24: Summary of modules assigned, and genes within them, from the WGCNA constructed
1340   from Af293 samples.
1341
1342   Table S25: Significantly enriched functional terms from a series of enrichment tests (Fisher's
1343   Exact Test) conducted on the genes within WGCNA modules.
1344
1345
1346

 # Supplemental information and results

Starship frequencies

The population-level frequency of high-confidence *Starships* ranges from 1-60%, with 13 of the
high-confidence *Starships* present in <10% of strains and no *Starship* present in more than 60%
of strains (Table S6). Each high-confidence *Starship* is present across multiple individuals;
however, any given *Starship* element is only present at most once in a genome, with only 3
exceptions that likely involve recently degraded copies, suggesting some mechanism for
controlling their copy number (e.g., repeat-induced point mutation (21)). Individual elements
typically segregate at low frequencies, with each segregating site containing on average an
inserted element in just 0.4% of strains across the 20 high-confidence *Starships* (range = 0.2-
52.3%, std. dev. = 6.5%, n = 154; Fig. 3B). However, multiple insertion sites often exist for a given
element such that each high-confidence *Starship* is present on average in 4.35 different regions
(range = 1-15, std. dev. = 3.92).

Differential expression of *Starship* captains and cargo

Patterns of transcript coverage, transcript abundance, and differential expression of *Starship*-
associated genes were generally heterogeneous, with various strain-, treatment- and strain by
treatment interactions. The transcriptional profile for many *Starships* is patchy, and not all
Starship-mobilized genes are expressed in response to experimental conditions. Only certain
cargo genes have the minimum required evidence for constitutive expression: a median transcript
coverage of 1 across the gene body. Nearly all genes in *A. fumigatus* CEA10 *Starships* surpass
this threshold (Fig. S13B, across all samples from the various studies we included in our analyses.
One notable exception being the captain gene in *Janus h2*, which is sparsely expressed across
most studies. In comparison, genes present in *Starships* from Af293 (Fig. S13A) and A1163 (Fig.
S13C) have fewer genes that surpass this threshold, with certain experimental conditions having
more consistent expression of the *Starship* genes in these strains. For example, specific studies
conducted on Af293 or CEA10, testing exposure to heatshock (PRJDB6203), hypoxia
(PRJNA144647), *in vitro* infections (PRJEB1583, PRJNA399754, PRJNA399754). and various
gene knock-out experiments (PRJNA390719, PRJNA396210, and PRJNA601094) have
consistently reported expression of all genes across the *Starship* element.
1376
Captain genes of multiple *Starships* from Af293 and CEA10 had low transcript abundance within
"infection" studies. Furthermore, samples of a study performing an experimental infection using a
murine model (PRJNA421149) had near-zero transcript abundances of captain genes from Af293
Starships Janus h2, *Osiris h3*, and *Gnosis h1*. However, this was not found within samples of a
different study of an Af293 infection of human pneumocyte cell lines (PRJNA399754). Near-zero
abundances were also found for a different set of captain genes, within CEA10 *Starships Tardis*
h1, Logos h2, and *Gnosis h2*, from an *in vitro* infection study using human cell lines (Fig. S12),
suggesting that the transcriptional response of *Starship* captains within the infection environment
differs between strains.
1386

We used tests of differential expression using existing *A. fumigatus* RNAseq studies to identify which environments or conditions influence the expression of genes within the *Starship* compartment (Fig. 5B, Fig. S14, Table S23). In particular, the change in expression of *Starship* captain genes within studies that applied similar types of treatments may have implications for what conditions may be more conducive for transposition. In general, multiple Af293 RNAseq "infection" studies have a prevailing negative impact on the expression of captain genes. Starships *Lamia h4*, *Tardis h1*, and *Gnosis h1* all had significantly lower expression within captain genes compared to their respective control samples (Figure 5B). We also observed a negative impact on captain expression within "antifungal" studies, which consisted of multiple studies conducted on Af293 and CEA10 with exposure to the antifungal caspofungin. Treatments of caspofungin were found to decrease the expression of Starship captains within Af293 Nebuchadnezzar h1 and CEA10 *Logos h2*. However, caspofungin was also found to be associated with increased expression of Af293 *Tardis h1* and *Gnosis h1* captain genes. Specific "nutrient" studies were also found to have a positive influence on captain gene expression. This includes a single study exposing A1163 to glucose/xylose media (PRJNA634102), which significantly increased the expression of captain genes within *Logos h1* and *Nebuchadnezzar h1*, compared to the control group grown on minimal media. Together, the differential expression results across these gene expression datasets demonstrate the context-dependent expression of captain genes, which we speculate would lead to variation in *Starship* transposition rates across different environments and strain backgrounds.

Similar to captain genes, the expression patterns of cargo genes within *Starships* are heterogeneous. The transcriptional profile of *Starships* is punctuated with expression of cargo genes (Fig. 5B, Fig. S14, Table S23), which can have sufficient transcript coverage even in the absence of captain expression, suggesting that the transcriptional regulation of these cargo genes are decoupled from transposition, and could be integrated into the existing genomic transcriptional networks. Of particular note are the genes that form biosynthetic gene clusters (BGC) which reside within *Starships*. In Af293, 63 BGC genes across seven Starships (*Gnosis h1, Lamia h4, Nebuchadnezzar h1, Nebuchadnezzar h2, Osiris h4, Tardis h1*, and *nav10 var35*) are differentially expressed in one or more of the treatment categories investigated in this study. At least one gene from the BGCs across these seven *Starships* are differentially expressed in response to caspofungin exposure, from a specific Af293 study (PRJNA472460). Furthermore, multiple genes present in Osiris h4 which are associated with another BGC (AFUA_3G02580, AFUA_3G02620, AFUA_3G02630, AFUA_3G02640) have significantly increased expression in nutrient studies compared to controls. Interestingly, generally fewer genes in BGCs are differentially expressed in Af293 "nutrient" studies.

As an opportunistic human pathogen, we were also interested in *A. fumigatus* genes that have specific relevance to virulence, such as genes within the *hrmA*-associated cluster (HAC). Two HAC genes present in *Nebuchadnezzar h1*, *cgnA* and *bafA*, have near-zero transcript abundances within samples from an Af293 study of an experimental infection of a murine model (PRJNA421149). However, genes within the HAC cluster were only differentially expressed within certain CEA10 antifungal studies and a single infection study: we found *bafA* in *Nebuchadnezzar*

1430 *h1* (CEA10_g5945) and *bafZ* in *Logos h1* (CEA10_g7202) to both have decreased expression in
1431 these studies, respectively (Fig. 5B, Fig. S14).

1432

Certain cargo genes within Starships demonstrate patterns of differential expression in response
to various experimental conditions, suggesting that these genes play a functional role in metabolic
or enzymatic activity of the cell in response to specific stimuli. In Af293, we identified several
Starship cargo genes with condition-specific expression patterns. Within *Nebuchadnezzar h1*, a
proteoglycan 4 gene (AFUA_5G14930) showed increased expression during infection.

A group of cargo genes within Af293 *Lamia h4* displayed complex expression patterns across
various treatment categories in this study. Within "infection" studies, multiple genes with diverse
functions had decreased expression, including genes relevant to metabolic functions (
AFUA_1G00800; flavin-containing amine oxidoreductase, AFUA_1G00340; thioesterase, and
AFUA_1G00490; dihydrolipoamide acetyltransferase component of pyruvate dehydrogenase
complex). One gene relevant to post-transcriptional modification also had decreased expression
(AFUA_1G00730; DNAj domain protein). Few genes within Lamia h4 that serve potential roles in
redox processes had increased expression within "infection" or "nutrient" treatments
(AFUA_1G00440; DUF895 domain membrane protein and AFUA_1G00350; FAD-binding
domains, respectively).

In CEA10, several genes showed increased expression with "antifungal" treatments, including
genes containing NAD(P)-binding (CEA10_g9263; *Janus h2*) and PAS superfamily protein
(CEA10_g5959; *Nebuchadnezzar h1*), and phosphotyrosine protein phosphatase
(CEA10_g5954; *Nebuchadnezzar h1*). While several cargo genes were consistently
downregulated in CEA10 *Logos h2*, including those encoding WD40 repeat proteins
(CEA10_g5566), P-loop containing nucleoside triphosphate hydrolase (CEA10_g5567),
homeodomain-like superfamily proteins (CEA10_g5587), and serine/threonine and tyrosine
kinase (CEA10_g5577).

In response to "nutrient" conditions, several cargo genes within A1163 *Logos h1* were differentially
expressed. with increased expression of serine threonine-protein (A1163_007845), DUF3723,
MADS-box transcription factor (A1163_007852), and freB (A1163_007853). Conversely, a BTB
domain-containing gene (A1163_007857), potentially involved in transcriptional regulation
through chromatin structure control, showed decreased expression under the same conditions.

Additional cargo genes of interest may convey resistance to environmental stressors, including
genes that provide resistance to environmental toxins. Genes incorporated in the arsenic
detoxification pathway are present within *A. fumigatus* Starships *Nebuchadnezzar h1* within
strains Af293 and CEA10. Arsenite methyltransferase genes were found to have significantly
decreased expression in the Af293 copy (AFUA_5G15000) and significantly increased expression
in the CEA10 copy (CEA10_g5956) with exposure to caspofungin (Fig. 5B). Collectively, these
findings highlight the diverse functional roles of Starship cargo genes in responding to
environmental stressors and nutrient availability, suggesting their importance in adaptability and
survival of *A. fumigatus*.

Starship co-expression networks

We performed weighted gene co-expression network analysis (WGCNA) using *PyWGCNA* to construct gene co-expression networks for *A. fumigatus* reference strain Af293. Overall, we found that genes within or between *Starships* are more strongly co-expressed together than with non-*Starship* genes, based on pairwise comparisons of the distribution of scores in the topological overlap matrices (TOM) (corrected p-values < 0.01 for all comparisons; Fig. S15). This suggests that a tighter transcriptional relationship exists between genes mobilized by *Starships* in Af293 compared to co-expression with other genes in the genome.

Genes in the *A. fumigatus* Af293 co-expression network were resolved into 27 modules (Table S24). The majority of genes found within each module are not contained within a *Starship*, yet almost all modules contain one or more *Starship* genes, including those previously identified as DEGs (Table S24). Modules were selected for further analyses based on the correlation of their gene expression profiles (WGCNA eigengenes) and their association with a treatment category ("antifungal", "infection", or "nutrient") or specific study (Fig. 5E, Fig. S16).

To investigate functional compartmentalization of WGCNA modules, we performed enrichment tests and identified which GO/KEGG terms are significantly overrepresented within each module. Genes in module "2" are significantly enriched in functions including the production of antimicrobial secondary metabolites (fumagillin), the secretion of *A. fumigatus* mycotoxins (helvolic acid), proteins for heme binding, and synthesis of immunosuppressive compounds (endocrcoin) (Table S25).

To identify which genes are most strongly co-expressed with *Starship* cargo or captain genes, we subsetted the co-expression network to keep only the top 10 edges that were made between any pair of genes or any gene and a *Starship* captain. Two modules within the Af293 co-expression network ("2" and "15") are significantly more commonly co-expressed within samples from a single infection study of Af293 (PRJEB1583; Fig. 5E). The connections with the highest TOM scores in module "2" include those between an IBR finger domain protein within the *Starship Lamia h4* (AFUA_1G00150) and genes involved in RNA binding pathways (AFUA_6G12070), as well as genes for alpha-amylase (AFUA_2G03230) and anthrone oxygenase (AFUA_4G00225) (Fig. 5D). Genes within the module "15" include those tightly connected to the expression of F-box domain protein within *Lamia h4*. Genes within module "15" are enriched in glycerophospholipid metabolism (Table S25). These co-expression relationships provide insight into how *Starships*-mobilized genes integrate into the existing transcriptional network in *A. fumigatus* strains.

Module "3" is significantly associated with increased expression within a single study of an experimental infection in mice (PRJNA693756) (Fig. 5E, S16). Genes within module "3" are enriched for genes in ribosome biogenesis in eukaryotes, carbon metabolism, and pyruvate metabolism (Table S25). The connections with the highest TOM scores in module "3" include genes co-expressed with *Gnosis h1* captain gene: genes with predicted catalytic activity (AFUA_1G12370), zinc-containing alcohol dehydrogenase (AFUA_2G00970), mitochondrial respiration (AFUA_2G06020), and exonuclease activity/DNA-directed DNA polymerase activity/role in mitochondrial DNA replication and mitochondrion localization (AFUA_5G12640).

1514

Module "7" is significantly associated with two studies testing supplementation with 5,8-
dihydroxyoctadecadienoic acid (PRJNA658306) and lipo-chitooligosaccharides (PRJNA642658)
in Af293 (Fig. 5E, S16). The focus of both of these studies is to understand how these
supplementations impact the regulation of fungal growth and development. The connections with
the highest TOM scores in the module "7", include genes co-expressed with captain genes of
Tardis h1 and *Osiris h3. Osiris h3* captain co-expressed with predicted RNA binding, ribonuclease
III activity and role in RNA processing (AFUA_3G03050). The connections made in this module
with *Starship* genes present themselves as good candidates for future research to understand
which genes are expressed along with transposition.

1524
1525

Figure 1: At least 20 distinct *Starships* carrying hundreds of protein-coding genes vary in their presence/ absence across *Aspergillus fumigatus* strains. A) Top: a SNP-based maximum likelihood tree of 220 *A. fumigatus* strains from Lofgren et al. 2022 (visualized as a cladogram for clarity), color-coded according to phylogenetic clade as defined by Lofgren et al. 2022. Bottom: A heatmap depicting the presence (gray) and absence (white) of 20 high-confidence *Starships* in each isolate. B) Schematics and sequence alignments of the type elements from 20 high-confidence *Starships*, where links between schematics represent alignable regions ≥500bp and ≥80% nucleotide sequence identity and arrows represent predicted coding sequences. C) A donut chart summarizing the number of distinct types of *Starships* by the quality of their prediction. D) A Venn diagram indicating the number of shared and unique high-confidence *Starships* in the reference strains Af293 and CEA10 (Table S7). E) Pairwise genome alignments between the CEA10 isolate from BioSample SAMN28487501 and the CEA10-derived A1163 isolate sequenced in this study demonstrating *Starship* movement. *Nebuchadnezzar h1* is present on Chromosome 6 in CEA10 (CP097568.1: 3645813-3698867) and absent at the corresponding locus in A1163 (NODE10: 188873-188877). Conversely, *Nebuchadnezzar h1* is present on Chromosome 4 in A1163 (NODE6: 77737-130777) but absent from the corresponding locus in CEA10 (CP097566.1:1344957-1344961).

Figure 2: *Starships* are enriched in accessory genes whose presence/absence and genomic location vary across *Aspergillus fumigatus* strains. All panels visualize data derived from 13 reference-quality *A. fumigatus* assemblies and the 20 high-confidence *Starships* (n = 459 elements total). A) Box-and-whisker plots summarizing the total percentage of nucleotide sequence and predicted genes carried by *Starships* per genome. B) Donut charts summarizing the percentages of gene orthogroups in the core, accessory, singleton, and *Starship*-associated compartments of the *A. fumigatus* pangenome (Table S10). C) Iceberg plots summarizing the genomic locations of the single best BLASTp hits (≥90% identity, ≥33% query coverage) to the cargo genes from the type elements of all 20 high-confidence *Starships* (left) and two individual *Starships* (right; Table S12). Each column in the iceberg plot represents a cargo gene and is color-coded according to the genomic location of hits (full dataset in Fig. S5).

Figure 3: *Starships* and their insertion sites are distributed across all major chromosomes in the *Aspergillus fumigatus* genome. A) A Circos plot summarizing all inserted *Starships* (in red) and all genomic regions containing either an empty insertion site or a fragmented *Starship* (in black, along perimeter and labeled with the r prefix) in the 8 chromosomes of the *A. fumigatus* reference strain Af293 (Table S14). All genomic regions contain a *Starship* insertion in some other individual from the 519 strain population. B) Barcharts summarizing the genotypes of segregating genomic regions associated with the four most active high-confidence elements in the 519 strain population (Table S16; full dataset in Fig. S7). If an isolate did not have any *Starships* within a given region, it was assigned either an "empty" or "fragmented" genotype (Methods). C) A scatterplot summarizing the relationship between the number of genomic regions containing a given *Starship* and the total number of copies of that *Starship* in the 519 strain population, where each point represents one of the 20 high-confidence *Starships*. A line derived from a linear regression model is superimposed, with shaded 95% confidence intervals drawn in gray. D) Alignments of *Tardis h1* copies +/- 50kb of flanking sequence across 14 genomic regions. Links between schematics represent alignable regions ≥1000bp and ≥90% nucleotide sequence identity.

Figure 4: *Starships* mobilize adaptive traits and generate allelic diversity among *Aspergillus fumigatus* strains. Schematics and alignments of: A) *Starship Lamia h2* and *h4*, which carry the biosynthetic gene cluster (BGC) encoding the polyketide secondary metabolite fumigermin, shown inserted at 3 independent sites. B) Starship *Gnosis h2* carrying the BGC encoding the terpene secondary metabolite fumihopaside A, shown inserted at 2 independent sites. C) a large region on Chromosome 3 previously identified as an idiomorphic BGC containing multiple segregating *Starship* insertions and various combinations of putative BGCs, including a a non-reducing polyketide synthase (NR-PKS) BGC carried by *Starship Osiris h4*. *Starships* from the *Osiris* navis specifically insert in 5S rDNA sequence and are predicted to fragment it. D) Eight *Starships* in the reference strains Af293 and CEA10 that all carry homologs of biofilm architecture factor A (bafA). *Starship Nebuchadnezzar h1* (*Neb. h1*) carries bafA as part of the H$_A$AC (hrmA-associated gene cluster), while *Starship navis10-var35* carries bafB as part of H$_B$AC and *Starship Osiris h3* carries bafC as part of H$_C$AC. Only a portion of *Starship Lamia h4* is visualized for figure legibility. All data for A-D was collected from the 519 *A. fumigatus* strain population (Table S5). Links between schematics represent alignable regions ≥5000bp and ≥95% nucleotide sequence identity and arrows represent predicted coding sequences. Abbreviations: polyketide synthase (PKS); non-ribosomal peptide synthetase (NRPS); highly reducing (HR).

Figure 5: A) A heatmap of transcript abundances (log10TPM) of *Starship* captain tyrosine recombinase genes (top) and genes within the *hrmA*-associated cluster (HAC; bottom), collapsed across treatment replicates, for 11 RNAseq studies from *A. fumigatus* Af293. B) Results from differential expression tests for select *Starships*, treatment categories, and strain combinations. Differentially expressed genes (DEGs) based on a single study are shown as log2 fold-change (log2FC) in black with standard error bars, whereas DEGs identified across multiple studies are represented with summarized log2FC values from a random effects model (REM) and coloured by "sign-consistency", the number of studies that reported DEGs in the positive (+1) or negative (-1) direction, centered around 0. Labels in bold font represent DEGs that are significantly differentially expressed in more than one study. C) The results of a weighted gene co-expression network analysis (WGCNA) constructed from 14 "antifungal", "infection", and "nutrient" RNAseq studies represented using UMAP clustering (90) based on the co-expression eigengene values for all genes in the *A. fumigatus* Af293 genome. Non-*Starship* genes present within modules of interest are shown in blue, while all *Starship* genes are shown in red, with those genes within modules of interest having a black outline. D) Genes within modules that were significantly associated with samples from specific treatment categories or studies were used to construct sub-networks which contained the top 10 edges in the network that were made between any pair of genes or any gene and a *Starship* captain. The connections between genes (edges) are based on the topological overlap matrix (TOM) for each module, and have been 0-1 scaled. E) Boxplots of eigengene values from WGCNA, akin to a weighted average expression profile, indicate the extent of co-expression (correlation) of the genes present within each module. Pairwise comparisons of module eigengene values determined if a module was significantly associated with samples from specific treatment categories or studies (Fig. S16).

Figure 6: Contributions of *Starships* to strain heterogeneity in *Aspergillus fumigatus*.