Density-dependent network structuring within and across wild animal systems

Gregory F Albery*[1,2,3], Daniel J Becker[4], Josh A Firth[5,6], Delphine De Moor[7], Sanjana Ravindran[8], Matthew Silk[8,9,10], Amy R Sweeny[8,11], Eric Vander Wal[12], Quinn Webber[13], Bryony Allen[14,15], Simon A Babayan[16], Sahas Barve[17,18], Mike Begon[14], Richard J Birtles[19], Theadora A Block[20,21], Barbara A Block[22], Janette E Bradley[23], Sarah Budischak[24], Christina Buesching[25], Sarah J Burthe[26], Aaron B Carlisle[27], Jennifer E Caselle[28], Ciro Cattuto[29], Alexis S Chaine[30], Taylor K Chapple[31], Barbara J Cheney[32], Timothy Clutton-Brock[33], Melissa Collier[2], David J Curnick[34], Richard J Delahay[35], Damien R Farine[36,37,38,39], Andy Fenton[14], Francesco Ferretti[40], Laura Feyrer[41,42], Helen Fielding[43,44], Vivienne Foroughirad[2,45], Celine Frere[46], Michael G Gardner[47], Eli Geffen[48], Stephanie S Godfrey[49], Andrea L Graham[50], Phil S Hammond[51], Maik Henrich[52], Marco Heurich[52,53,54], Paul Hopwood[10], Amiyaal Ilany[55], Joseph A Jackson[19], Nicola Jackson[46], David MP Jacoby[56], Ann-Marie Jacoby[57], Miloš Ježek[58], Lucinda Kirkpatrick[59,60], Alisa Klamm[61], James A Klarevas-Irby[36,37,62,63], Sarah Knowles[5], Lee Koren[55], Ewa Krzyszczyk[60], Jillian M Kusch[64,65], Xavier Lambin[66], Jeffrey E Lane[65], Herwig Leirs[59], Stephan T Leu[67], Bruce E Lyon[20], David W MacDonald[68], Anastasia E Madsen[69,70], Janet Mann[2], Marta Manser[71], Joachim Mariën[59], Apia Massawe[72], Robbie A McDonald[44], Kevin Morelle[58,73], Johann Mourier[74,75,76], Chris Newman[68], Kenneth Nussear[77], Brendah Nyaguthii[38,39,62,78], Mina Ogino[36,37,62], Laura Ozella[79], Craig Packer[80], Yannis P Papastamatiou[81], Steve Paterson[14], Eric Payne[82], Amy B Pedersen[8], Josephine M Pemberton[8], Noa Pinter-Wollman[83], Serge Planes[75,76], Aura Raulo[5], Rolando Rodríguez-Muñoz[10], Lauren Rudd[5], Christopher Sabuni[72], Pratha Sah[2], Robert J Schallert[84], Ben C Sheldon[5], Daizaburo Shizuka[85], Andrew Sih[82], David L Sinn[86], Vincent Sluydts[59], Orr Spiegel[48], Sandra Telfer[66], Courtney A Thomason[40,87], David M Tickler[88], Tom Tregenza[10], Kimberley VanderWaal[89], Sam Walmsley[41], Eric L Walters[17], Klara M Wanelik[14,90], Hal Whitehead[41], Elodie Wielgus[52], Jared Wilson-Aggarwal[10,44], Caroline Wohlfeil[47], Shweta Bansal[2]

*Corresponding author: gfalbery@gmail.com

Affiliations can be found at the end of the document.

Summary

High population density should drive individuals to more frequently share space and interact, producing better-connected spatial and social networks [1–4]. Although this theory is fundamental to our understanding of disease dynamics [2,5–8], it remains unconfirmed how local density generally drives individuals' positions within their networks, which reduces our ability to understand and predict density-dependent processes [4,9,10]. Here we provide the first general evidence that density drives greater network connectedness at fine spatiotemporal scales, at the scale of individuals within wild animal populations. We analysed 36 datasets of simultaneous spatial and social behaviour in >58,000 individual animals, spanning 30 species of fish, reptiles, birds, mammals, and insects. 80% of systems exhibited strong positive relationships between local density and network centrality. However, >80% of relationships were nonlinear and 75% became shallower at higher values, signifying that demographic and behavioural processes counteract density's effects,

thereby producing saturating trends [11–15]. Density's effect was much stronger and less
saturating for spatial than social networks, such that individuals become disproportionately
spatially connected rather than socially at higher densities. Consequently, ecological
processes that depend on spatial connections (e.g. indirect pathogen transmission, resource
competition, and territory formation) are likely more density-dependent than those involving
social interactions (e.g. direct pathogen transmission, aggression, and social learning).
These findings reveal fundamental ecological rules governing societal structuring, with
widespread implications. Identifying scaling rules based on processes that generalise across
systems, such as these patterns of density dependence, might provide the ability to predict
network structures in novel systems.

Keywords: Behavioural ecology, Spatial ecology, Disease ecology, Epidemiology,
Population dynamics, Social network structure, Network analysis, Spatial analysis

Introduction

The number of individuals occupying a given space – i.e., population density – is a central
factor governing social systems. At higher densities, individuals are expected to more
frequently share space, associate, and interact, producing more-connected spatial and
social networks and thereby influencing downstream processes such as mating, learning,
and competition. In particular, density-driven increases in network connectedness should
provide more opportunities for parasites [1–4,9] or information [16] to spread between
hosts[1–4,9] Despite the fundamental nature of such density-dependent processes,
evidence is relatively limited that individuals inhabiting higher-density areas have more
spatial and social connections. Furthermore, density effects should differ for asynchronous
space sharing (e.g. home range overlap) *versus* social associations (e.g. den sharing or
grouping) or interactions (e.g. mating or fighting). While several studies have compared
animal populations at different densities to demonstrate variation in social association rates
among populations (e.g., [15,17,18]) or groups (e.g., [11–13]), attempts to identify such
density effects *within* continuous populations of individuals are rarer (but see [14,15,19–21]),
and their findings have never been synthesised or compared for spatial and social
behaviours. We therefore have an incomplete understanding of how density, as a
fundamental ecological parameter, determines socio-spatial dynamics within and across
systems. This inhibits our ability to identify and predict how changes in density – e.g. through
culling, natural mortality, dispersal, or population booms – influence downstream processes
that depend on shared space and social interactions.

The rate at which an individual interacts with conspecifics depends on its spatial and social
behaviour within the context of the surrounding environment and population. Adding more
individuals into the same space should cause them to more frequently spatially overlap and
socially associate or interact (Figure 1). Often, individuals are modelled as randomly moving
and interacting molecules ("mass action" or "mean field"). In this conceptualisation, direct
contact between two molecules is analogous to a social interaction or association; rates of
such interactions are often assumed to increase with density ("density-dependent"; e.g.,
[22]), and/or to be homogenous in space (e.g., [13]). In reality individuals are unlikely to
behave and interact randomly in space, and instead will be influenced by spatially varying
factors including local density [10] and competition for resources [15]. Changes in density
may cause individuals to alter their foraging behaviour [23–25], dispersal [26,27], social

preference or avoidance [14,28], mating behaviour [29], or preferred group size [18]. In some
cases, density may have no effect on interaction rates, because individual animals alter their
behaviour in a density-dependent manner to maintain a desired interaction rate [30]. These
and related processes might produce strong nonlinearities in density-interaction
relationships, which can complicate the predictions of density dependence models of
pathogen transmission, for example [2,3,9]. For example, individuals or groups can learn to
avoid where competitors might go, resulting in greater spatial partitioning under higher
densities [31]. Nevertheless, nonlinearities such as these are poorly understood and rarely
considered.

Several wild animal studies have suggested relationships between density and social
association rates are often nonlinear and saturating [11–15]. Such relationships imply that
association rates do not increase passively with density, but rather that behavioural or
demographic processes likely change as density increases, with the ultimate consequence
of slowing association rates. However, these nonlinearities are difficult to examine between
populations or between species because they introduce a range of confounders and have
few replicates along the density axis [9]. On the other end, lower densities may provide less
ability to exert social preferences, but low-density populations may be harder to study due to
(for example) low return on sampling investment; alternatively, the failure to achieve
sufficient interaction rates may result in Allee effects and ultimately drive populations toward
decline [32,33].

Characterising gradients of density across individuals within a population offers a
workaround to these problems, and facilitates an appreciation of the fact that interactions
occur between individuals rather than at the population level. Examining between-individual
variation is one reason that social network analysis – which allows characterisation and
analysis of individual-level social traits, amongst other things – has become so popular in
animal ecology in recent years [34–38]. Additionally, recent years have seen a substantial
growth in understanding of socio-spatial behaviours, including harmonising the concepts of
spatial and social density [9,10,39]. Applying network analyses coupled with this socio-
spatial understanding of density could provide an individual-level picture of density's effects
on spatial and social connectedness, offering far higher resolution and statistical power and
greater ability to detect within-system nonlinearities and between-system differences [9]. By
providing new understanding of the correlates and emergent consequences of variation in
density, this expansion could help to identify general rules underlying social structuring and
network scaling in space.

Critically, different types of interactions or associations should show different relationships
with density: for example, the need to compete for food at higher densities could drive a
disproportionate increase in aggression [40], but this is unlikely to be true of mating
interactions. In contrast, higher density and food scarcity should lead to lower exclusivity in
resources and more overlapping home ranges, thus enhancing the effect of density on
spatial network [41]. This rationale is well-understood in disease ecology, as differences in
density-contact relationships are thought to drive differences in density dependence of
infection – where "contact" is defined as an interaction or association that could spread a
pathogen (Figure 1). "Contacts" then form the basis of spatial and social networks used to
investigate pathogen transmission dynamics, which should likewise diverge with density just
as contacts do. For example, density should drive greater transmission of respiratory

pathogens but not sexually transmitted pathogens [1,42]. Establishing these density-contact relationships is integral to understanding disease dynamics and developing control measures [1,43], but we still have a poor understanding of how different interactions (and therefore contact events for different pathogens) are driven by density. This direct/indirect interaction dichotomy is most fundamental to disease ecology [39,44], but given building interest in the spatial-social interface and relationships between spatial and social networks in behavioural ecology [10], the framework is readily related to other fields (e.g. direct versus indirect cues that can lead to social learning [45]). Previously established density-interaction relationships are diverse and include feral dog bites [19], ant antennations [46] and trophallaxis [30], ungulate group memberships [20,23], rodent co-trapping [11,47], and agamid association patterns [14,21], but no study has yet synthesised how the rates of multiple interaction or association types relate to density, within or across systems.

Identifying the general rules underlying density dependence requires quantifying density's relationship with proxies of interaction rates at fine scales across a diversity of systems, then identifying the factors determining their slope and shape. To this end, we collate a meta-dataset of over 58,000 individual animals across 36 wildlife systems globally (Figure 2) to ask how within-population variation in density determines between-individual interaction rates based on connectedness in spatial and social networks. We fit multiple competing linear and nonlinear relationships to identify the slope and shape of density effects within each system, and we use meta-analyses to investigate general rules determining their slope and shape across systems. In particular, we focus on comparing space sharing with social interactions and associations as a cross-system case study. Ultimately, we present a *de novo* cross-system analysis of individuals' social and spatial behaviour that traverses fields of behavioural, population, and disease ecology, which could help to inform general rules governing the structure of social systems, and eventually shape management and conservation decisions in a wide range of systems.

160 *Figure 1: Schematic detailing the rationale underlying this study, outlining how population density*
161 *drives the formation of spatial and social networks. This depiction uses the Wytham Wood great tits*
162 *as an example. In panel C, the different purple shades correspond to different individuals' home*
163 *ranges. In panel D, red lines depict connections among individuals, with each individual located at*
164 *their centroid. Ultimately, one of our main aims is to ask whether spatial or social connections*
165 *generally show a stronger relationship with density, partly functioning as a proxy for indirect and direct*
166 *contact events with the potential to transmit pathogens. This framework moves between concepts of*
167 *network and contact formation traversing behavioural ecology, spatial and social network ecology,*
168 *and disease ecology.*

170
171

Figure 2: The phylogenetic (A) and geographic (B) distribution of our 36 examined datasets of spatial
and social behaviour, with (C) schematic depicting the methodology for deriving local density values,
using the Isle of Rum red deer data as an example. The X and Y axes are bivariate spatial
coordinates. The panels within (C) show raw observations of individuals in space that we then
average at the individual level to make centroids; we use the centroids to generate annual density
distributions, which are then assigned to individuals in the form of local density measures. Animal
silhouettes are from phylopic.org; a list of attributions is in the supplement (Supplementary Table 2).
NB the Potomac dolphins are now defined as Tursiops erebennus; they are currently incorporated in
Panel A as T. truncatus, following the Open Tree of Life nomenclature.

Figure 3: Meta-analysis revealed drivers of variation in linear density effects on individual network connectedness across systems. A) Our fitted linear model estimates of density effects on network strength. Each point represents the mean estimate from a given system; the error bars denote 95% confidence intervals. Opaque error bars were significant (i.e., do not overlap with 0); transparent ones were not. The estimates are in units of standard deviations for both density and network strength. The colour of the point denotes whether the network being examined was defined using spatial or social connections. B) Meta-analyses revealed that centrality in spatial networks (i.e., home range overlap; red points) had a significantly steeper relationship with density than social networks (blue points). C) We fitted linear models separately to two portions of the data within each study population ("first" and "last" represent values below and above the median). The slopes for the latter portion (pink points) were generally less positive than the former portion (purple points), implying a general saturation shape. In panels B) and C), each coloured point represents a study replicate fitted to the strength estimate; points are sized according to sample size, and jittered slightly on the x axis to reduce overplotting. The large black points represent the mean slope estimated from the meta-analysis, and the error bars represent 95% confidence intervals.

Figure 4: Relationships between density and network connectedness varied substantially across animal systems. Density in individuals per area is on the x axis; network connectedness (strength centrality) is on the y axis. Both values have been standardised to have a mean of zero and a standard deviation of 1 within each system; the axis ticks are in units of 1 standard deviation. Each point represents an individual-year-behaviour replicate; the lines portray the model fit from our generalised additive models (GAMs). Red lengths of the smooth=significantly positive; grey=not significantly different from zero; blue=significantly negative. Points are semi-transparent to enhance visibility. Panels are arranged phylogenetically following the tree displayed in Figure 2A; GOG=gambit of the group; HRO=home range overlap. Animal silhouettes are from phylopic.org; a set of links and attributions are in the Supplement.

208 Results and Discussion

We compiled a comparative meta-dataset of over ten million observations of individual
animals' spatial and social behaviour, across a wide range of ecological systems and
taxonomic groups of animals. We then ran a standardised pipeline to align their spatial and
social observations, identifying strong and predictable relationships between local density
and network connectedness at the individual level.

We observed strong positive relationships between individuals' local population density and
their connectedness in spatial and social networks across a wide range of wild animals: of
our 64 replicates, 51 (78%) were significantly positive when analysed using linear models
(Figure 3A). Meta-analyses identified a highly significant positive mean correlation between
density and connectedness, both for social networks (Estimate 0.22; 95% CI 0.17, 0.27) and
spatial networks (0.45; 0.36, 0.53; Figure 3B). Our study therefore provides fundamental
evidence that high local population density broadly drives greater connectedness within
ecological systems, at the individual level. Slopes were highly variable across systems for
both spatial and social networks (Figure 3A; Q-test of heterogeneity across systems: Q_{37} =
5627.33 and Q_{25} = 1281.83, both P<0.0001), indicating that quantifying these slopes within
and between multiple systems and comparing them is important for understanding animal
socio-spatial structure. That is, relationships between density and individual connectedness
differ substantially between populations, and the biological mechanisms underlying these
divergent trends are likely important. As well as adding resolution and allowing comparisons
of density effects across systems, our methodology facilitated fitting of nonlinear
relationships (using generalised additive models (GAMs); see below). This approach has
only rarely been applied before, and then at much coarser resolution (see [11,12,19]). As
such, this study fills an important empirical gap by providing insights into the slope and
shape of density-connectedness relationships for a diverse variety of animal groups and
their social and spatial behaviours (Figure 4). Nevertheless, despite this diversity, we were
able to identify several further general trends in our data.

Remarkably, density's effect more than doubled in size for spatial compared to social
networks (Figure 3B; r=0.45 *versus* 0.22); there was a difference of 0.26 (CI 0.16, 0.36,
P<0.0001) for this effect when we meta-analysed the two contact types together. This finding
indicates that as density increases, wild animals are more likely to share space with each
other, but that social connections increase at a much slower rate. Similarly, we discovered
that saturating shapes were extremely common: as density increased, its effect on
connectedness decreased, such that 48/64 systems (75%) had a steeper slope at low
density values than at high ones. This effect was strong for both social networks (effect on r=
-0.11; CI -0.19, -0.03; P=0.01) and for spatial networks, with substantial overlap between
their estimates (-0.22; -0.37, -0.07; P=0.0042). Due to the greater overall effect for space
sharing, the latter half of density's spatial effect was still higher than the first half of its social
effect (Figure 3C). Together, these observations suggest that density-dependent processes
act to limit the increase in social connectedness with density, but without limiting spatial
overlaps to the same extent. Consequently, higher-density areas are characterised
disproportionately by individuals asynchronously sharing space rather than socially
associating, while in lower-density areas individuals are disproportionately more socially
connected proportional to their shared space.

There are many possible social reasons for saturating nonlinearity in density-dependent
network structuring: for example, individuals in higher density areas may begin to avoid each
other, seeking to avoid competition or aggression [40] or exposure to infectious disease [48].
For instance, Eastern water dragons (*Intellagama lesueurii*) show greater avoidance at
higher densities [14], supporting avoidance-related mechanisms. Alternatively, in species
with high social cognition or stable bonds, saturation could reflect lower social effort or ability
to keep track of social affiliates at higher densities [49]. In general, individuals likely have a
preferred social interaction rate or group size – a preference that they may increasingly exert
at higher densities [18]. It remains to be seen how this preference varies among individuals,
and whether individuals vary in their preferred social network position given a certain
density. Given that individuals vary in their movement and spatial phenotypes [50–52], and
social phenotypes [52–54] in ways that should manifest for density-dependent behaviours
specifically [10], it seems likely that these slopes could vary between individuals as they do
between populations. Future analyses might fit variable density-connectedness slopes
among individuals to identify socio-spatial syndromes across systems, as has been done
previously in single systems including caribou (*Rangifer tarandus*) [55] and American red
squirrels (*Tamiasciurus hudsonicus*) [56]. Additionally, we could dissect the social network
and its relationship to the spatial network to identify levels of attraction [57,58] or avoidance
[59] and how they depend on density.

We considered that density-dependent changes in spatial behaviours might explain these
trends: for example, density could create greater competition over resources and therefore
reduce energy to roam (and contact others). Individuals may partition their niches [60], or
reduce their territory or home range sizes [56,61,62], potentially driven by years of plentiful
resources supporting higher densities alongside smaller home ranges sufficiently providing
ones' resource needs, which could drive lower association rates. However, our findings do
not seem to support explanations related to small home ranges, because such explanations
should produce an equivalent or stronger reduction in (relative) spatial connectedness. In
contrast, we observed that density drove individuals to become spatially connected faster
than socially, such that the underlying mechanisms likely involve behaviours and
demographic processes that specifically affect social collocation in space and time. Testing
the precise underlying mechanism will likely require finer-scale behavioural observations, as
described below. Regardless of mechanism, these saturating density-connectedness
relationships strongly support the idea that examining density effects at the individual level –
rather than between populations – is highly informative. For many systems, "mean field"
expectations of homogenous interactions under increasing density likely produce an
inaccurate (i.e., inflated) picture of density's effects. Importantly, our study included many
examples of proximity-based social networks – most notably "gambit of the group" measures
[63] – but relatively few "direct" interactions such as mating, grooming, or fighting. It is
interesting that these differences manifested even among two ostensibly spatially-defined
contact metrics (gambit of the group and home range overlap). This observation supports
the assertion that social association metrics defined by spatiotemporal proximity are valuable
for informing on social processes separately from more spatial behaviours *sensu stricto* such
as ranging behaviour [20]; we expect that "more direct" interactions could show even further
differences in relationships with density. Incorporating a larger number of "direct" metric-
based systems could help to address this question (see Supplementary Discussion).

The fact that spatial networks show stronger and more linear density dependence than
social networks could heavily influence the ecology of animal systems. For example,
indirectly transmitted (i.e., environmentally latent) parasites may exhibit greater density
dependence than directly transmitted ones, given that individuals likely experience
disproportionately more indirect contact at higher densities. This observation contrasts with
orthodoxy that directly transmitted parasites are most likely to be density dependent [64],
and supports the value of investigating nonlinear changes in socio-spatial behaviour and
grouping patterns in response to density when considering density dependence. Saturating
density-connectedness functions further have implications for disease modelling and control.
Specifically, our findings lend behavioural support to the growing consensus that many
diseases are density-dependent at lower densities, but not at higher densities (i.e., that the
slope flattens with density) [22,65]. Rather than assuming constant behavioural mixing at
higher densities, epidemiological models could benefit from incorporating density-dependent
shifts in behaviours and demography that influence direct and indirect interaction
frequencies, as previously suggested empirically and by epidemiological theory [22]. These
relationships could influence our targets for culling or vaccination coverage [5]. Given that
animals at high density seem likely to have a relatively shallow relationship between density
and contact rates, reducing population density – for example by culling – might therefore be
ineffective at reducing pathogen transmission initially, particularly when considering socially
transmitted pathogens, where contact rates are particularly likely to have become saturated
(Figure 3C). Similar problems with culling have already been acknowledged in specific
systems – e.g. in canine rabies [43,66,67] – but our study implies that shallow nonlinear
density-contact trends could be more general than previously thought and could be driven by
flexible density-dependent changes in behaviour and demography. Conversely, culling could
be disproportionately effective at intermediate densities and identifying the inflection points
of the curve might help to design optimal management strategies. Future studies should
investigate whether the divergence in spatial and social connectedness with density drives a
concurrent divergence in the prevalence of directly and indirectly transmitted parasites, as
well as addressing several other biases in our selection of systems (e.g. [68]; see
Supplementary Discussion).

Beyond these general trends, we ran generalised additive models (GAMs) that revealed that
52/64 density effects on network connectedness (81%) were significantly nonlinear
(ΔAIC>2); these relationships took a wide variety of shapes, representing a range of
nonlinear functions that are hard to generalise (Figure 4). Notably, while many GAM
smooths were eventually significantly negative (Figure 4), the vast majority of linear models
fitted to the second half of the data were positive (Figure 3C); this result is likely an artefact
of restricted model fitting, rather than true downturns in connectedness with density.
Nonlinearity did not cluster according to connection type definitions, or according to animal
group. These observations were largely corroborated by our meta-analytical models, which
found no factors influencing the slope and shape of density effects overall (P>0.05;
Supplementary Table 3), including no clear phylogenetic signal (ΔAIC=2.71). This
observation speaks to the complexity of these relationships within and across systems, while
accentuating that simple functional relationships are often likely to be complicated by
contravening ecological factors like habitat selection [69,70], group formation [15], parasite
avoidance [71], and demographic structuring [72]. While we were unable to identify specific
between-system predictors of nonlinearity of density-connectedness relationships, the

finding that most such relationships are strongly nonlinear is an important consideration for
future work.

Density is a universal factor underlying the dynamics of animal populations, and its linear
and nonlinear effects on spatial and social network structure are likely to impact myriad
processes in behaviour, ecology, and evolution. Similar to other studies that have reported
general scaling patterns in network analysis [73] and in food web ecology [74], the patterns
we report strongly suggest that animal systems generally become more connected spatially
than socially under increasing density. These trends might extrapolate to human networks,
given that other scaling patterns in animal networks do [73]. As these patterns seemingly
manifest regardless of animal group and interaction type, they may reflect a generalisable
rule governing the socio-spatial structure of ecological systems. Further refining and
implementing these models could facilitate prediction of network structure in novel systems.

Finally, this study is relatively unique in conducting an expansive meta-analysis of
behavioural data from individual animals across a diverse selection of systems. As datasets
accumulate comparative analyses are increasing in frequency in social network ecology [75],
but often revolve around analysing whole networks rather than individuals [76], and never (to
our knowledge) in conjunction with analyses of spatial behaviour. These analyses therefore
hold exceptional promise for disentangling spatial and social behaviour across diverse
systems; for example, given that our dataset includes many repeatedly sampled known
individuals, future analyses could investigate individual-level repeatability or multi-behaviour
"behavioural syndromes" across a variety of different taxa and environments [10,77].
Additionally, capitalising on the wide range of methodological approaches to behavioural
data collection (e.g. censuses, trapping, and GPS telemetry), the methodological constraints
of socio-spatial analyses could be tested in this wide meta-dataset as they have been in
other recent comparative analyses of wild ungulates [78]. As well as being diverse, our
meta-dataset had several replicate examples of (for example) marine mammals and trapped
rodents, which could be used for finer-scale and more targeted comparative analyses within
these smaller taxonomic groupings. For now, it is highly encouraging that we uncovered
general trends across these disparate animal systems, and further explorations of these
socio-spatial patterns may help to inform a wide range of exciting and longstanding
questions at the spatial-social interface [10].

374 Methods

375 Data standardisation and behavioural pipeline

Data were manipulated and analysed using R version 4.2.3 [79], and all R code is available
at https://github.com/gfalbery/DensityMetaAnalysis. Our 36 datasets each involved at least
one continuous uninterrupted spatial distribution of observations in a single population; some
datasets comprised multiple such populations; all systems had at least one social network
measure, and two had two different types of social interaction. These datasets covered 30
different animal species, including sharks, carnivores, cetaceans, ungulates, rodents,
elephants, birds, reptiles, and one orthopteran insect (Figure 2). In one case (The Firth of
Tay and Moray Dolphins) we used two distinct replicates despite being composed of
overlapping groups of individuals, because of their distinct spatial distributions, which made
it difficult to fit a coherent density distribution.

To standardise the timescale across studies, all systems were analysed as annual replicates
– i.e., social and spatial networks were summarised within each year. Our analyses used 64
system-behaviour replicates, listed in Supplementary Table 1, and totalled 151,835 unique
system-individual-year-behaviour data points.

All spatial coordinates were converted to the scale of kilometres or metres to allow
comparison across systems. To provide an approximation of local density, following prior
methodology [20,80], we took each individual's average location across the year (their
centroid) and created a spatial density kernel using the `adehabitathr` package [81], which
provides a probabilistic distribution of population density across each study system based on
the local frequencies of observed individuals. Each individual was assigned an annual
estimate of local density based on their centroid's location within this spatial density
distribution. We made these density distributions as comparable as possible between
systems by incorporating the density raster using metre squares; however, there were large
differences in density across populations that were difficult to resolve and put on the same
scale (e.g. interactions per individual/km^2 unit of density). Consequently, we scaled and
centred density to have a mean of zero and a standard deviation of one within each
population, which allowed us to focus on differences in relative slope and shape across
systems.

To validate the local density measures estimated using the kernel density approach, we also
estimated local density for individuals across all populations based on the locations of
individual annual centroids within a designated area. To do so, we first estimated the area of
the minimum bounding box (MBB) within which all individuals were censused during the
study period based on their annual centroids. For each individuals mean location, we then
estimated a circular boundary of radius r=1/20 * area of MBB. We then calculated the
number of individuals present within this boundary as an individual's local density measure.
We estimated the Pearson correlation coefficients between the local density measures
derived using the KDE approach and the proportional area - based approach
(Supplementary Figure 1).

To provide a measure of asynchronous space sharing, we constructed home range overlap
(HRO) networks based on proportional overlap of two individuals' minimum convex polygon
(MCP; i.e., the bounding polygon around all observations of each individual in a given year).
These HRO networks were restricted to only individuals with five or more observations in a
given year to allow us to create convex polygons effectively; 10/36 (28%) systems did not
have sufficient sampling for this analysis. We also repeated our analyses with a series of
higher sampling requirements for observation numbers to ensure that our findings were
robust to this assumption. The MCP approach is relatively low-resolution, and assumes
uniform space use across an individual's home range; however, this approach is less data
intensive – and less sensitive to assumptions – than density kernel-based approaches that
would estimate variation in space use across the home range, allowing us to apply the
models across more systems, more generalisably, and more conservatively.

To provide a measure of social connectedness, we built social networks using various
approaches as defined by the original studies: direct observations of dyadic interactions (e.g.
fighting or mating); gambit of the group (GoG; i.e., membership of the same group) [63]; co-
trapping (i.e., trapped together or in adjacent traps within a given number of trapping

sessions); or direct contact measured by proximity sensors (defined by a certain distance-
based detection threshold). Notably some analyses use indirect interactions – i.e., spatial
overlap – to *approximate* direct interactions, which requires spatiotemporal coincidence,
which we caution against particularly when modelling pathogen transmission [39,82]. While
the two do often correlate, here we are not using HRO to approximate direct interaction
rates, but rather as a measure of indirect interactions (e.g., indicative of transmission of
environmental parasites).

For each social network, we scaled connection strength relative to the number of
observations of each individual in a dyad (i.e., simple ratio index [83]). Our response variable
therefore took the form of strength centrality, scaled to between 0-1 for each dyad, for each
social and spatial network. We focus on comparing density effects on social interactions and
associations with density's effects on space sharing.

Density-connectedness models: what forms do density effects take?

We developed a novel workflow to allow us to derive and compare density's effects on
connectedness – and their drivers – in a standardised way across our animal systems. We
fitted models with three main forms: **linear models** fitted to the whole dataset, nonlinear
Generalised additive models fitted to the whole dataset, and linear **saturation models**
fitted separately to low- and high-density subsets of each dataset.

Linear models: For each system-behaviour replicate, we first fitted a linear model using the
`lm` function in R, fitting scaled density as an explanatory variable to estimate linear density
effect slopes. The linear fits are displayed in the supplement (Supplementary Figure 2), as
are the residuals (Supplementary Figure 3).

Generalised additive models (GAMs): We fitted GAMs in the `mgcv` package [84] to
identify whether each density effect was better described by a linear or nonlinear
relationship, and to identify the shape of these nonlinear relationships. For each model, we
fitted a default thin plate spline with k=4 knots. This knot number was selected to reduce
overfitting in our models, which formed several fits to the data that were difficult to reconcile
with functional formats. To assess whether nonlinear models fit better than linear models, we
used Akaike Information Criterion (AIC), with a contrast of 2ΔAIC designated to distinguish
between models.

Saturation models: To quantify whether density effects were generally saturating (i.e., that
density had steeper relationships with individuals' connectedness at lower density values),
we split the data into two portions: all values below the median density value, and all values
above the median. We then re-ran linear models examining the relationship between density
and strength in each portion. We attempted to investigate nonlinear patterns (especially
saturating effects) across all our systems using a range of other methods (e.g., comparing
specific functional relationships with nonlinear least squares), but found that they were
generally incapable of fitting well to the data in a standardised way across the many datasets
(i.e., non-convergence of nonlinear least squares using semi-automated starting estimates
across systems). As such, this approach represented a tradeoff between tractable,
generalisable model fitting, interpretability, and accurate representation of the relationship's
shape. All else being equal, we posit that investigating the relative slopes of two otherwise-

identical portions of the data is a conservative and informative method of identifying
saturation, which was our main hypothesis for the expected shape of density effects.

Heteroskedasticity and log-log models: To ensure that our estimates were robust to non-
normality and to provide another source of information concerning possible saturation
effects, we also conducted tests of heteroskedasticity on our linear models and
accompanied them with simulations and fitted log-log linear models. First, we carried out a
simple simulation study to test how: a) the skew in residuals; b) a saturating relationship; and
c) heteroscedasticity impact whether we may under- or overestimate the slope of an
assumed linear relationship between density and strength (See Supplementary Methods -
Heteroskedasticity Simulations). These demonstrated that our models were resilient to skew
and saturating effects, but that heteroskedasticity in residuals could drive overestimated
linear effects in our models.

To examine this possibility further, we derived the Breusch-Pagan statistic for each linear
model as a measure of heteroskedasticity, and then plotted it against the meta-analysis
covariates and fixed effects. There was no evidence that the density effect was being
skewed to be greater for spatial behaviours due to heteroskedasticity, and neither were the
second portions of the data more heteroskedastic, which would be expected if this was
driving the saturating effect (Supplementary Figure 4). Finally, we fitted log-log linear models
with the same formulations as our main linear models defined above, but with both density
and strength log(X+1)-transformed, rather than scaled to have a mean of 0 and a standard
deviation of 1 (Supplementary Figure . Our results showed broadly identical findings of
greater estimates for spatial behaviours, and the fact that the slopes were largely under 1 is
indicative of a saturating effect. As such, these tests strongly support our findings' resilience
to uneven data distributions.

Meta-analysis: what factors determine the slope of density-connectedness relationships?

To characterise the typical relative slope of density effects across systems and identify the
factors influencing their variation, we fitted hierarchical meta-analytical models using the
`metafor` package in R. The response variable was the standardised slope of the linear
density effect; because both individual network strength and density were scaled to have
mean of zero and standard deviation of one in the linear regression, this is equivalent to the
correlation coefficient (r) [85]. We converted all correlation coefficients into Fisher's Z (Z_r)
and computed associated sampling variance.

For our hierarchical meta-analysis models, we used an initial model that nested observations
within a system-level random effect to account for within- and between-system heterogeneity
[86], as 26/36 systems had more than one density effect. We used another random effect for
species to account for repeat observations per animal species.

We then added a separate random effect for animal phylogeny [87]. This effect used a
phylogenetic correlation matrix of our 30 animal species derived from the Open Tree of Life
via the `rotl` package [88], with the `ape` package used to resolve multichotomies and
provide branch lengths [89].

We then fitted intercept-only models using the `rma.mv()` function with restricted maximum
likelihood (REML), weighted by inverse sampling variance, and used variance components
to quantify I^2, the contribution of true heterogeneity to the total variance in effect size. We
used Cochran's Q to test whether such heterogeneity was greater than that expected by
sampling error alone.

We next fitted models with the same random effects structure that included explanatory
variables. To detect whether some animals were more likely to experience density effects,
we fitted **Animal group** as a factor with six categories, representing a combination of
species' taxonomy and general ecology: aquatic (fish and dolphins), birds, large herbivores
(elephants and ungulates), small mammals (rodents and hyraxes), carnivores, and
ectotherms (insects and reptiles). We also fitted several explanatory variables indicative of
greater statistical power that might increase the strength of density effects: **Geographic
area** (km^2, log$_{10}$-transformed), **Number of years** of study, and **Number of individuals**, all of
which we fitted as continuous covariates. Broadly, the animal group model was highly
uninformative and competed with the other effects, and we expected that the phylogeny
would be more informative, so we report the results of the model without the animal group
effect fitted.

We ran several different versions of these meta-analyses: first, we fitted meta-analytical
models to the **overall linear models** of spatial and social interaction types separately, and
then together, to investigate differences between the spatial and social networks in terms of
their mean density slope. Next, we fitted duplicated versions of these models, but with the
saturation models. These models were identical, but each system replicate had two linear
estimates: one taken from the first 50% of the data (up to the median), and one from the
latter 50%. By fitting a binary fixed effect of "data portion" to the meta-analytical models, this
model would tell us whether the slopes were generally higher in the first portion of the data
than the last (and therefore showed generally saturating shapes). We were unable to fit
meta-analytical models to our GAMMs, as methods for the meta-analysis of nonlinear
estimates are not yet well defined.

542   Acknowledgements

GFA and SB were supported by NSF grant number DEB-2211287. GFA acknowledges
support from a College for Life Sciences Fellowship at the Wissenschaftskolleg zu Berlin and
WAI (C-2023-00057). DJB acknowledges support from the Edward Mallinckrodt, Jr.
Foundation and NSF BII 2213854. JAF acknowledges funding from BBSRC
(BB/S009752/1), NERC (NE/S010335/1 and NE/V013483/1), and WAI (C-2023-00057). For
a list of system-specific acknowledgements, please see the supplement.

550   Code availability

551   All R code is available at https://github.com/gfalbery/DensityMetaAnalysis.

Affiliations

1. School of Natural Sciences, Trinity College Dublin, Dublin, Ireland
2. Department of Biology, Georgetown University, Washington, DC, USA
3. Wissenschaftskolleg zu Berlin, Berlin, Germany
4. School of Biological Sciences, University of Oklahoma, Norman, OK, USA
5. Department of Biology, University of Oxford, Oxford, UK
6. School of Biology, University of Leeds, Leeds, UK
7. Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany
8. Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, UK
9. CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
10. Centre for Ecology and Conservation, University of Exeter, Penryn, UK
11. School of Biosciences, University of Sheffield, Sheffield, UK
12. Department of Biology, Memorial University of Newfoundland, 45 Arctic Avenue, St. John's NL
13. University of Guelph, Guelph, Canada
14. Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK
15. Institute of Zoology, Zoological Society of London, Regents Park, London, UK
16. University of Glasgow, Glasgow, UK
17. Department of Biological Sciences, Old Dominion University, Norfolk, VA, USA
18. Avian Ecology Program, Archbold Biological Station, Venus, FL, USA
19. School of Science, Engineering and Environment, University of Salford, Salford, UK
20. Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA
21. Canine Companions, 2965 Dutton Ave, Santa Rosa, CA, USA
22. Hopkins Marine Station, Stanford University, Stanford, CA, USA
23. School of Life Sciences, University of Nottingham, Nottingham, UK
24. Pitzer and Scripps Colleges, Claremont, CA, USA
25. The University of British Columbia, Irving K. Barber Faculty of Sciences, Okanagan Department of Biology, Kelowna, British Columbia, Canada
26. UK Centre for Ecology and Hydrology, Bush Estate, Midlothian, UK
27. University of Delaware, Newark, DE, USA
28. Marine Science Institute, UC Santa Barbara, Santa Barbara, CA, USA
29. ISI Foundation, Torino, Italy
30. Evolutionary Ecology Group, Station d'Ecologie Theorique et Experimentale du CNRS, Moulis, France
31. Oregon State University, Corvallis, OR, USA
32. Lighthouse Field Station, School of Biological Sciences, University of Aberdeen, UK
33. Department of Zoology, University of Cambridge, Cambridge, UK
34. Institute of Zoology, Zoological Society of London, London, UK

35. National Wildlife Management Centre, Animal and Plant Health Agency, Sand Hutton, York, UK

36. Department of Collective Behavior, Max Planck Institute for Animal Behavior, 78464 Konstanz, Germany

37. Department of Evolutionary Biology and Environmental Studies, University of Zurich, 8057 Zurich, Switzerland

38. Department of Ornithology, National Museum of Kenya, P.O. Box 40241-001000, Nairobi, Kenya

39. Division of Ecology and Evolution, Research School of Biology, Australian National University, 46 Sullivan Creek Road, Canberra, ACT 2600, Australia

40. Virginia Tech, Blacksburg, VA, USA

41. Department of Biology, Dalhousie University, Halifax, Nova Scotia, B3H 4R2, Canada

42. Department of Fisheries and Oceans, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, B2Y 4A2, Canada

43. The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK

44. Environment and Sustainability Institute, University of Exeter, Penryn, UK

45. Department of Marine Biology, Texas AandM University at Galveston, USA

46. School of the Environment, University of Queensland, Brisbane, Queensland, Australia

47. College of Science and Engineering, Flinders University, Adelaide, Australia

48. School of Zoology, Faculty of Life Science, Tel Aviv University, Tel Aviv, Israel

49. Department of Zoology, University of Otago, Otago, New Zealand

50. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

51. Sea Mammal Research Unit, Scottish Oceans Institute, University of St Andrews, UK

52. Department of Visitor Management and National Park Monitoring, Bavarian Forest National Park, Grafenau, Germany

53. Chair of Wildlife Ecology and Wildlife Management, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany

54. Department of Forestry and Wildlife Management, Faculty of Applied Ecology, Agricultural Sciences and Biotechnology, Campus Evenstad, Inland Norway University of Applied Sciences, Postboks 400 Vestad, 2418 Elverum, Norway

55. Faculty of Life Sciences, Bar Ilan University, Ramat Gan, Israel

56. Lancaster University, Lancaster, UK

57. Division of Marine Science and Conservation, Nicholas School of the Environment, Duke University Marine Laboratory, Beaufort, NC, USA

58. Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech Republic

59. Evolutionary Ecology Group, EVECO, Universiteit Antwerpen, Antwerp, Belgium

60. School of Environment and Natural Science, Bangor University, Bangor, Wales

61. Department of Nature Conservation and Research, Hainich National Park,
Bad Langensalza, Germany
62. Mpala Research Center, P.O Box 555-10400, Nanyuki, 10400, Kenya
63. Centre for the Advanced Study of Collective Behaviour, University of
Konstanz, 78464 Konstanz, Germany
64. Department of Biology, Memorial University of Newfoundland, St John's, NL,
Canada
65. Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada
66. School of Biological Sciences, University of Aberdeen, Aberdeen, UK
67. School of Animal and Veterinary Sciences, The University of Adelaide,
Adelaide, Australia
68. Wildlife Conservation Research Unit, The Recanati-Kaplan Centre,
Department of Zoology, University of Oxford, Oxford, UK
69. School of Biological Sciences, University of Nebraska-Lincoln, NE, USA
70. Department of Ecology, Behavior, and Evolution, University of California, San
Diego, CA
71. Department of Evolutionary Biology and Environmental Studies, University of
Zurich, Zurich, Switzerland
72. Institute of Pest Management, Sokoine University of Agriculture, Morogoro,
Tanzania
73. Max Planck Institute of Animal Behavior, Radolfzell, Germany
74. MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France
75. PSL Research University, Université de Perpignan, Perpignan, France
76. Laboratoire d'Excellence "CORAIL", Papetoai, Moorea, French Polynesia
77. University of Nevada, Reno, NV, USA
78. University of Eldoret, School of Natural Resource Management, Department
of Wildlife, 1125-30100 Eldoret, Kenya
79. Department of Veterinary Sciences, University of Turin, Grugliasco (TO), Italy
80. Department of Ecology, Evolution and Behavior, University of Minnesota
81. Institute of Environment, Department of Biological Science, Florida
International University, FL, USA
82. Department of Environmental Science and Policy, University of California
Davis, Davis, CA, USA
83. Department of Ecology and Evolutionary Biology, University of California, Los
Angeles, CA, USA
84. Stanford University, Stanford, CA, USA
85. School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE,
USA
86. Department of Wildlife, California State Polytechnic University, Humboldt,
Arcata, CA, USA
87. Texas Tech University, Lubbock, TX, USA
88. University of Western Australia, Crawley WA 6009, Australia
89. Department of Veterinary Population Medicine, University of Minnesota,
Minneapolis, US
90. School of Biosciences, University of Surrey, Guildford, UK
688

References

1. Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latto J, Sanchez MS, Smith AB, Swei A. 2005 Should we expect population thresholds for wildlife disease? *Trends Ecol Evol* **20**, 511–519. (doi:10.1016/j.tree.2005.07.004)

2. Begon M, Bennett M, Bowers RG, French NP, Hazel SM, Turner J. 2002 A clarification of transmission terms in host-microparasite models: Numbers, densities and areas. *Epidemiol Infect* **129**, 147–153. (doi:10.1017/S0950268802007148)

3. Ferrari MJ, Perkins SE, Pomeroy LW, Bjrnstad ON. 2011 Pathogens, social networks, and the paradox of transmission scaling. *Interdiscip Perspect Infect Dis* **2011**. (doi:10.1155/2011/267049)

4. Hopkins SR, Fleming-Davies AE, Belden LK, Wojdak JM. 2020 Systematic review of modeling assumptions and empirical evidence: does parasite transmission increase nonlinearly with host density? *Methods Ecol Evol* **11**, 0–2. (doi:10.1111/2041-210x.13361)

5. McCallum H, Barlow N, Hone J. 2001 How should pathogen transmission be modelled? *Trends Ecol Evol* **16**, 295–300. (doi:10.1016/S0169-5347(01)02144-9)

6. Anderson RM, May RM. 1991 *Infectious Diseases of Humans*. Oxford, UK: Oxford University Press. See https://global.oup.com/academic/product/infectious-diseases-of-humans-9780198540403?cc=gb&lang=en&.

7. Antonovics J, Iwasa Y, Hassell MP. 1995 A generalized model of parasitoid, venereal, and vector-based transmission processes. *American Naturalist* **145**, 661–675. (doi:10.1086/285761)

8. Fenton A, Fairbairn JP, Norman R, Hudson PJ. 2002 Parasite transmission: reconciling theory and reality. *Journal of Animal Ecology* **71**, 893–905. (doi:10.1046/J.1365-2656.2002.00656.X)

9. Albery GF. 2022 Density dependence and disease dynamics: moving towards a predictive framework. *EcoEvoRxiv Preprints* (doi:10.32942/OSF.IO/GAW49)

10. Webber QMR, Albery GF, Farine DR, Pinter-wollman N, Sharma N, Vander Wal E, Manlove K. 2023 Behavioural ecology at the spatial-social interface. *Biological Reviews* **3**, 868–886.

11. Borremans B, Reijniers J, Hens N, Leirs H. 2017 The shape of the contact–density function matters when modelling parasite transmission in fluctuating populations. *R Soc Open Sci* **4**. (doi:10.1098/rsos.171308)

727   12.   Davis S, Abbasi B, Shah S, Telfer S, Begon M. 2015 Spatial analyses of wildlife
728   contact networks. *J R Soc Interface* **12**, 20141004. (doi:10.1098/rsif.2014.1004)

729   13.   Carson BD, Orians CM, Crone EE. 2023 When does spatial clustering in
730   caterpillars inuence the relationship between population density and contact
731   rates? (doi:10.21203/rs.3.rs-2842538/v1)

732   14.   Strickland K, Patterson EM, Frère CH. 2018 Eastern water dragons use
733   alternative social tactics at different local densities. *Behav Ecol Sociobiol* **72**.
734   (doi:10.1007/s00265-018-2563-x)

735   15.   Rouhbakhsh A, Wright AN, Ferguson JM. 2025 Ecological Interactions Drive a
736   Power-Law Relationship Between Group Size and Population Density in Social
737   Foragers. *Ecol Lett* **28**, 70111. (doi:10.1111/ele.70111)

738   16.   Cantor M, Chimento M, Smeele SQ, He P, Papageorgiou D, Aplin LM, Farine
739   DR. 2021 Social network architecture and the tempo of cumulative cultural
740   evolution. *Proceedings of the Royal Society B: Biological Sciences* **288**.
741   (doi:10.1098/rspb.2020.3107)

742   17.   Webber QMR, Vander Wal E. 2020 Heterogeneity in social network
743   connections is density-dependent: implications for disease dynamics in a
744   gregarious ungulate. *Behav Ecol Sociobiol* **74**, 1–13.

745   18.   Beauchamp G. 2011 Functional relationship between group size and
746   population density in Northwest Atlantic seabirds. *Mar Ecol Prog Ser* **435**, 225–
747   233. (doi:10.3354/meps09239)

748   19.   Mancy R *et al.* 2022 Rabies shows how scale of transmission can enable acute
749   infections to persist at low prevalence. *Science (1979)* **376**, 512–516.

750   20.   Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH,
751   Firth JA. 2021 Multiple spatial behaviours govern social network positions in a
752   wild ungulate. *Ecol Lett* **24**, 676–686. (doi:10.1101/2020.06.04.135467)

753   21.   Strickland K, Frère CH. 2019 Individual Variation in the Social Plasticity of
754   Water Dragons. *Am Nat* **194**, 194–206. (doi:10.5061/dryad.5j5gh63)

755   22.   Hu H, Nigmatulina K, Eckhoff P. 2013 The scaling of contact rates with
756   population density for the infectious disease models. *Math Biosci* **244**, 125–
757   134. (doi:10.1016/j.mbs.2013.04.013)

758   23.   Albon SD, Staines HJ, Guinness FE, Clutton-Brock TH. 1992 Density-
759   Dependent Changes in the Spacing Behaviour of Female Kin in Red Deer.
760   *Journal of Animal Ecology* **61**, 131–137.

761   24.   Vas J, Andersen IL. 2015 Density-dependent spacing behaviour and activity
762   budget in pregnant, domestic goats (Capra hircus). *PLoS One* **10**, e0144583.
763   (doi:10.1371/journal.pone.0144583)

25. Connor EF, Cargain MJ. 1994 Density-related foraging behaviour in
Closterocerus tricinctus, a parasitoid of the leaf-mining moth, Cameraria
hamadryadella. *Ecol Entomol* **19**, 327–334. (doi:10.1111/J.1365-
2311.1994.TB00249.X)

26. Matthysen E. 2005 Density-dependent dispersal in birds and mammals.
Ecography **28**, 403–416. (doi:10.1111/j.0906-7590.2005.04073.x)

27. Avgar T, Betini GS, Fryxell JM. 2020 Habitat selection patterns are density
dependent under the ideal free distribution. *Journal of Animal Ecology* **89**,
2777–2787. (doi:10.1111/1365-2656.13352)

28. Vander Wal E, Laforge MP, Mcloughlin PD, Wal E Vander, Laforge MP,
Mcloughlin PD, Vander Wal E, Laforge MP, Mcloughlin PD. 2014 Density
dependence in social behaviour: Home range overlap and density interacts to
affect conspecific encounter rates in a gregarious ungulate. *Behav Ecol*
Sociobiol **68**, 383–390. (doi:10.1007/s00265-013-1652-0)

29. Kokko H, Rankin DJ. 2006 Lonely hearts or sex in the city? Density-dependent
effects in mating systems. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **361**, 319–334. (doi:10.1098/rstb.2005.1784)

30. Modlmeier AP, Colman E, Hanks EM, Bringenberg R, Bansal S, Hughes DP.
2019 Ant colonies maintain social homeostasis in the face of decreased
density. *Elife* **8**, 1–17. (doi:10.7554/eLife.38473)

31. Ogino M, Farine DR. 2024 Collective intelligence facilitates emergent resource
partitioning through frequency-dependent learning. *Philosophical Transactions*
of the Royal Society B: Biological Sciences **379**.
(doi:10.1098/RSTB.2023.0177;JOURNAL:JOURNAL:RSTB1887;WEBSITE:W
EBSITE:RSJ-SITE;WGROUP:STRING:PUBLICATION)

32. Stephens PA, Sutherland WJ. 1999 Consequences of the Allee effect for
behaviour, ecology and conservation. *Trends Ecol Evol* **14**, 401–405.
(doi:10.1016/S0169-5347(99)01684-5)

33. Courchamp F, Clutton-Brock T, Grenfell B. 1999 Inverse density dependence
and the Allee effect. *Trends Ecol Evol* **14**, 405–410. (doi:10.1016/S0169-
5347(99)01683-3)

34. Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting
animal social network analysis. *Journal of Animal Ecology* **84**, 1144–1163.
(doi:10.1111/1365-2656.12418)

35. Farine DR. 2024 Modelling animal social networks: New solutions and future
directions. *Journal of Animal Ecology* **93**, 250–253. (doi:10.1111/1365-
2656.14049)

36. Webber QMR, Vander Wal E. 2019 Trends and perspectives on the use of
animal social network analysis in behavioural ecology: a bibliometric approach.
Anim Behav **149**, 77–87. (doi:10.1016/j.anbehav.2019.01.010)

37. Pinter-Wollman N *et al.* 2014 The dynamics of animal social networks:
Analytical, conceptual, and theoretical advances. *Behavioral Ecology* **25**, 242–
255. (doi:10.1093/beheco/art047)

38. Sah P, Mann J, Bansal S. 2018 Disease implications of animal social network
structure: A synthesis across social systems. *Journal of Animal Ecology* **87**,
546–558. (doi:10.1111/1365-2656.12786)

39. Albery GF, Kirkpatrick L, Firth JA, Bansal S. 2021 Unifying spatial and social
network analysis in disease ecology. *Journal of Animal Ecology* **90**, 1–17.
(doi:10.1111/1365-2656.13356)

40. Cubaynes S, Macnulty DR, Stahler DR, Quimby KA, Smith DW, Coulson T.
2014 Density-dependent intraspecific aggression regulates survival in northern
Yellowstone wolves (Canis lupus). *Journal of Animal Ecology* **83**, 1344–1356.
(doi:10.1111/1365-2656.12238)

41. Jetz W, Carbone C, Fulford J, Brown JH. 2004 The scaling of animal space
use. *Science (1979)* **306**, 266–268.
(doi:10.1126/SCIENCE.1102138/SUPPL_FILE/JETZ.SOM.PDF)

42. Lloyd-Smith JO, Getz WM, Westerhoff H V. 2004 Frequency-dependent
incidence in models of sexually transmitted diseases: Portrayal of pair-based
transmission and effects of illness on contact behaviour. *Proceedings of the*
Royal Society B: Biological Sciences **271**, 625–634.
(doi:10.1098/rspb.2003.2632)

43. Morters MK, Restif O, Hampson K, Cleaveland S, Wood JLN, Conlan AJK.
2013 Evidence-based control of canine rabies: A critical review of population
density reduction. *Journal of Animal Ecology* **82**, 6–14. (doi:10.1111/j.1365-
2656.2012.02033.x)

44. Dougherty ER, Seidel DP, Carlson CJ, Spiegel O, Getz WM. 2018 Going
through the motions: incorporating movement analyses into disease research.
Ecol Lett **21**, 588–604. (doi:10.1111/ele.12917)

45. Hoppitt W, Laland KN. 2008 Chapter 3 Social Processes Influencing Learning
in Animals: A Review of the Evidence. *Adv Study Behav* **38**, 105–165.
(doi:10.1016/S0065-3454(08)00003-X)

46. Gordon DM, Paul RE, Thorpe K. 1993 What is the function of encounter
patterns in ant colonies? *Anim Behav* **45**, 1083–1100.
(doi:10.1006/ANBE.1993.1134)

47.   Davis S, Abbasi B, Shah S, Telfer S, Begon M, Davis S. 2015 Spatial analyses
839   of wildlife contact networks. *Journal of the Royal Society, Interface / the Royal
840   Society* **12**, 20141004. (doi:10.1098/rsif.2014.1004)

841   48.   Gibson AK, Amoroso CR. 2022 Evolution and Ecology of Parasite Avoidance.
842   *Annu Rev Ecol Evol Syst* **53**, 47–67.

843   49.   Colman E, Colizza V, Hanks EM, Hughes DP, Bansal S. 2021 Social fluidity
844   mobilizes contagion in human and animal populations. *Elife* **10**.
845   (doi:10.7554/ELIFE.62177)

846   50.   Michelangeli M, Payne E, Spiegel O, Sinn D, Leu ST, Gardner M, Sih A. 2022
847   Personality, spatiotemporal ecological variation, and resident/explorer
848   movement syndromes in the sleepy lizard. *Journal of Animal Ecology* **91**, 210–
849   223. (doi:10.1111/1365-2656.13616)

850   51.   Spiegel O, Leu ST, Bull CM, Sih A. 2017 What's your move? Movement as a
851   link between personality and spatial dynamics in animal populations. *Ecol Lett*
852   **20**, 3–18. (doi:10.1111/ele.12708)

853   52.   Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014 Individual-level personality
854   influences social foraging and collective behaviour in wild birds. *Proceedings of
855   the Royal Society of London B* **281**, 20141016. (doi:10.1098/rspb.2014.1016)

856   53.   Hunt ER, Mi B, Fernandez C, Wong BM, Pruitt JN, Pinter-Wollman N. 2018
857   Social interactions shape individual and collective personality in social spiders.
858   *Proceedings of the Royal Society B: Biological Sciences* **285**, 20181366.
859   (doi:10.1098/rspb.2018.1366)

860   54.   Firth JA, Cole EF, Ioannou CC, Quinn JL, Aplin LM, Culina A, McMahon K,
861   Sheldon BC. 2018 Personality shapes pair bonding in a wild bird social system.
862   *Nat Ecol Evol* **2**, 1696–1699. (doi:10.1038/s41559-018-0670-8)

863   55.   Webber Q, Laforge M, Bonar M, Wal E Vander. 2024 The adaptive value of
864   density-dependent habitat specialization and social network centrality. *Nat
865   Commun* **15**, 4423. (doi:10.22541/AU.165210484.46672609/V1)

866   56.   Webber QMR, Dantzer B, Lane JE, Boutin S, McAdam AG. 2023 Density-
867   dependent plasticity in territoriality revealed using social network analysis.
868   *Journal of Animal Ecology* **92**, 207–221. (doi:10.1111/1365-2656.13846)

869   57.   Spiegel O, Leu ST, Sih A, Bull CM. 2016 Socially interacting or indifferent
870   neighbours □? Randomization of movement paths to tease apart social
871   preference and spatial constraints. *Methods Ecol Evol* , 971–979.
872   (doi:10.1111/2041-210X.12553)

873   58.   Gahm K, Nguyen R, Acácio M, Anglister N, Vaadia G, Spiegel O, Pinter-
874   Wollman N. 2024 A wrap-around movement path randomization method to
875   distinguish social and spatial drivers of animal interactions. *Philosophical

Transactions of the Royal Society B: Biological Sciences **379**.
(doi:10.1098/RSTB.2022.0531,)

59. Strickland K, Levengood A, Foroughirad V, Mann J, Krzyszczyk E, Frère CH.
2017 A framework for the identification of long-term social avoidance in
longitudinal datasets. *R Soc Open Sci* **4**. (doi:10.1098/rsos.170641)

60. Sheppard CE, Inger R, McDonald RA, Barker S, Jackson AL, Thompson FJ,
Vitikainen EIK, Cant MA, Marshall HH. 2018 Intragroup competition predicts
individual foraging specialisation in a group-living mammal. *Ecol Lett* **21**, 665–
673. (doi:10.1111/ELE.12933)

61. Sanchez JN, Hudgens BR. 2015 Interactions between density, home range
behaviors, and contact rates in the Channel Island fox (Urocyon littoralis). *Ecol*
Evol **5**, 2466–2477. (doi:10.1002/ece3.1533)

62. Wilkin TA, Garant D, Gosler AG, Sheldon BC. 2006 Density effects on life-
history traits in a wild population of the great tit Parus major: Analyses of long-
term data with GIS techniques. *Journal of Animal Ecology* **75**, 604–615.
(doi:10.1111/j.1365-2656.2006.01078.x)

63. Franks DW, Ruxton GD, James R. 2010 Sampling animal association networks
with the gambit of the group. *Behav Ecol Sociobiol* **64**, 493–503.
(doi:10.1007/s00265-009-0865-8)

64. Cote IM, Poulin R. 1995 Parasitism and group size in social animals: a meta-
analysis. *Behavioral Ecology* **6**, 159–165. (doi:10.1093/beheco/6.2.159)

65. Ketwaroo FR, Matechou E, Silk M, Delahay R. 2025 Modeling Disease
Dynamics From Spatially Explicit Capture-Recapture Data. *Environmetrics* **36**,
e2888. (doi:10.1002/ENV.2888)

66. Rajeev M, Metcalf CJE, Hampson K. 2020 Modeling canine rabies virus
transmission dynamics. In *Rabies*, pp. 655–670. Elsevier. (doi:10.1016/b978-0-
12-818705-0.00020-0)

67. Townsend SE *et al.* 2013 Designing Programs for Eliminating Canine Rabies
from Islands: Bali, Indonesia as a Case Study. *PLoS Negl Trop Dis* **7**, e2372.
(doi:10.1371/JOURNAL.PNTD.0002372)

68. Albery GF, Sweeny AR, Corripio-Miyar Y, Evans MJ, Hayward A, Pemberton
JM, Pilkington JG, Nussey DH. 2024 Local and global density have distinct and
parasite-dependent effects on infection in wild sheep. *bioRxiv* ,
2024.04.30.591781. (doi:10.1101/2024.04.30.591781)

69. Tardy O, Massé A, Pelletier F, Mainguy J, Fortin D. 2014 Density-dependent
functional responses in habitat selection by two hosts of the raccoon rabies
virus variant. *Ecosphere* **5**, 1–16. (doi:10.1890/ES14-00197.1)

70. Wilber MQ, Yang A, Boughton R, Manlove KR, Miller RS, Pepin KM, Wittemyer
G. 2022 A model for leveraging animal movement to understand
spatio-temporal disease dynamics. *Ecol Lett* , 1–15. (doi:10.1111/ele.13986)

71. Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary
Consequences of Parasite Avoidance. *Trends Ecol Evol* **33**, 619–632.
(doi:10.1016/j.tree.2018.05.001)

72. Lion S, Boots M. 2010 Are parasites '"prudent"' in space? *Ecol Lett* **13**, 1245–
1255. (doi:10.1111/j.1461-0248.2010.01516.x)

73. Rocha LEC, Ryckebusch J, Schoors K, Smith M. 2021 Scaling of social
interactions across animal species. *Sci Rep* **11**, 12584.

74. Lafferty KD, DeLeo G, Briggs CJ, Dobson AP, Gross T, Kuris AM. 2015 A
general consumer-resource population model. *Science (1979)* **349**, 854–857.
(doi:10.1126/SCIENCE.AAA6224/SUPPL_FILE/LAFFERTY.SM.PDF)

75. Albery GF, Bansal S, Silk MJ. 2024 Comparative approaches in social network
ecology. *Ecol Lett* **27**. (doi:10.1111/ele.14345)

76. Collier M, Albery GF, McDonald GC, Bansal S. 2022 Pathogen transmission
modes determine contact network structure, altering other pathogen
characteristics. *Proceedings of the Royal Society B* **289**.
(doi:10.1098/RSPB.2022.1389)

77. Stuber EF, Carlson BS, Jesmer BR. 2022 Spatial personalities: A meta-
analysis of consistent individual differences in spatial behavior. *Behavioral
Ecology*. **33**, 477–486. (doi:10.1093/beheco/arab147)

78. Kaur P *et al.* 2024 A protocol for assessing bias and robustness of social
network metrics using GPS based radio-telemetry data. *Mov Ecol* **12**.
(doi:10.1186/s40462-024-00494-6)

79. R Development Core Team R. 2011 R: A Language and Environment for
Statistical Computing. *R Foundation for Statistical Computing*. **1**, 409.
(doi:10.1007/978-3-540-74686-7)

80. Albery GF, Newman C, Bright Ross J, Macdonald DW, Bansal S, Buesching
CD. 2020 Negative density-dependent parasitism in a group-living carnivore.
Proceedings of the Royal Society B: Biological Sciences **287**, 20202655.
(doi:10.1101/2020.06.15.153726)

81. Calenge C. 2011 Home range estimation in R: the adehabitatHR package. See
https://cran.r-project.org/web/packages/adehabitatHR/index.html (accessed on
10 March 2020).

82. Wanelik KM, Farine DR. 2022 A new method for characterising shared space
use networks using animal trapping data. *Behav Ecol Sociobiol* **76**.
(doi:10.1007/s00265-022-03222-5)

83. Cairns SJ, Schwager SJ. 1987 A comparison of association indices. *Anim Behav* **35**, 1454–1469. (doi:10.1016/S0003-3472(87)80018-0)

84. Wood SN. 2011 Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J R Stat Soc Series B Stat Methodol* **73**, 3–36. (doi:10.1111/j.1467-9868.2010.00749.x)

85. Koricheva J, Gurevitch J, Mengersen KL. 2013 Handbook of meta-analysis in ecology and evolution. , 498.

86. Konstantopoulos S. 2011 Fixed effects and variance components estimation in three-level meta-analysis. *Res Synth Methods* **2**, 61–76. (doi:10.1002/jrsm.35)

87. Cinar O, Nakagawa S, Viechtbauer W. 2022 Phylogenetic multilevel meta-analysis: A simulation study on the importance of modelling the phylogeny. *Methods Ecol Evol* **13**, 383–395. (doi:10.1111/2041-210X.13760)

88. Michonneau F, Brown JW, Winter DJ. 2016 rotl: an R package to interact with the Open Tree of Life data. *Methods Ecol Evol* **7**, 1476–1481. (doi:10.1111/2041-210X.12593)

89. Paradis E, Schliep K. 2019 ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528. (doi:10.1093/BIOINFORMATICS/BTY633)

Supplement: Albery et al. (2024): Density-dependent network structuring within and across wild animal systems

System	Species	Host Group	Contact Type	Behaviour	N	NID	Area	NYrs
AcornWoodpeckers [1]	*Melanerpes formicivorus*	Bird	GOG	Proximity	143	86	7.627757322	3
AcornWoodpeckers [1]	*Melanerpes formicivorus*	Bird	HRO	Indirect	56	37	5.006385251	3
AfricanDogs [2]	*Canis lupus*	Carnivore	HRO	Indirect	297	223	75.63528521	2
AfricanDogs [2]	*Canis lupus*	Carnivore	ProximityCollar	Proximity	297	223	75.63528521	2
BeakedWhales [3]	Hyperoodon ampullatus	Cetacean	Census	Proximity	942	526	351.0723	15
BeakedWhales [3]	Hyperoodon ampullatus	Cetacean	HRO	Indirect	136	109	184.109	13
ChagosSharks [4]	*Carcharhinus amblyrhynchos*	Shark	GOG	Proximity	103	71	1.290740108	3
DesertTortoises [5]	*Gopherus agassizii*	Ectotherm	DenSharing	Proximity	6864	3544	1.055393893	4
DesertTortoises [5]	*Gopherus agassizii*	Ectotherm	HRO	Indirect	3181	1519	0.627186899	4
EuropeanBoar [6]	*Sus scrofa*	LargeHerbivore	HRO	Indirect	46	46	1058.606722	1
EuropeanBoar [6]	*Sus scrofa*	LargeHerbivore	ProximityCollar	Proximity	46	46	1058.606722	1
FalkirkMice [7]	*Apodemus sylvaticus*	SmallMammal	TrapProximity	Proximity	168	168	0.0377755	1
FieldingCows [8]	*Bos taurus*	LargeHerbivore	HRO	Indirect	365	365	0.00788627	1
FieldingCows [8]	*Bos taurus*	LargeHerbivore	ProximityCollar	Proximity	365	365	0.00788627	1

Study	Species	Type	Method	Mode				
GoldenCrownedSparrows [9]	*Zonotrichia atricapilla*	Bird	GOG	Proximity	688	341	0.113428125	5
GoldenCrownedSparrows [9]	*Zonotrichia atricapilla*	Bird	HRO	Indirect	545	285	0.103413426	5
KalahariMeerkats [10]	*Suricata suricatta*	Carnivore	GOG	Proximity	6418	2734	2.3798173	22
KalahariMeerkats [10]	*Suricata suricatta*	Carnivore	HRO	Indirect	2412	1357	2.110591914	19
KenyanElephants [11]	*Loxodonta africana*	LargeHerbivore	GOG	Proximity	215	151	11677.13864	2
KenyanElephants [11]	*Loxodonta africana*	LargeHerbivore	HRO	Indirect	77	55	10647.43727	2
KielderVoles [12]	*Microtus agrestis*	SmallMammal	HRO	Indirect	545	544	0.028251875	6
KielderVoles [12]	*Microtus agrestis*	SmallMammal	TrapProximity	Proximity	10029	9140	0.050075	6
LiverpoolMice [13]	*Apodemus sylvaticus*	SmallMammal	HRO	Indirect	243	243	0.01890773	4
LiverpoolMice [13]	*Apodemus sylvaticus*	SmallMammal	TrapProximity	Proximity	4110	4054	0.0284	4
MooreaSharks [14]	*Carcharhinus melanopterus*	Shark	GOG	Proximity	269	105	11.74987669	3
MorayDolphins [15]	*Tursiops truncatus*	Cetacean	Census	Proximity	2849	687	1894.792195	32
MorayDolphins [15]	*Tursiops truncatus*	Cetacean	HRO	Indirect	1061	224	744.7757738	27
MountainLake [16]	*Peromyscus leucopus*	SmallMammal	TrapProximity	Proximity	319	272	0.0245	2
MultimammateMice [17]	*Mastomys natalensis*	SmallMammal	HRO	Indirect	323	321	0.000234789	20
MultimammateMice [17]	*Mastomys natalensis*	SmallMammal	TrapProximity	Proximity	21142	18631	0.000261	25
PalmyraSharks [18]	*Carcharhinus amblyrhynchos*	Shark	GOG	Proximity	121	41	51.38758347	4

Study	Species	Type	Method	Interaction			Value	
PalmyraSharks [18]	*Carcharhinus amblyrhynchos*	Shark	HRO	Indirect	88	36	48.65584545	4
PotomacDolphins [19]	*Tursiops erebennus*	Cetacean	GOG	Proximity	925	861	227.787706	3
PrairieDogs [20]	*Cynomys ludovicianus*	SmallMammal	GOG	Proximity	49	49	3990.296949	1
ReticulatedGiraffes [21]	*Giraffa camelopardalis*	LargeHerbivore	GOG	Proximity	214	214	189.1271192	1
ReticulatedGiraffes [21]	*Giraffa camelopardalis*	LargeHerbivore	HRO	Indirect	204	204	134.6231999	1
RockHyraxes [22]	*Procavia capensis*	SmallMammal	GOG	Proximity	399	196	0.093825	14
RumDeer [23]	*Cervus elaphus*	LargeHerbivore	Dominance	Direct	2173	391	16.91608696	22
RumDeer [23]	*Cervus elaphus*	LargeHerbivore	GOG	Proximity	6807	987	19.63333333	47
RumDeer [23]	*Cervus elaphus*	LargeHerbivore	HRO	Indirect	8251	889	17.77714286	47
SerengetiLions [24]	Panthera leo	Carnivore	Census	Proximity	1990	1990	3096.55	30
SerengetiLions [24]	Panthera leo	Carnivore	HRO	Indirect	1478	1478	2778.931	30
SharkBayDolphins [25]	*Tursiops aduncus*	Cetacean	Census	Proximity	4139	770	470.1272	12
SharkBayDolphins [25]	*Tursiops aduncus*	Cetacean	HRO	Indirect	1257	328	306.2262	12
SilwoodMice [26]	*Apodemus sylvaticus*	SmallMammal	CoTrapping	Proximity	542	497	0.000204	2
SilwoodMice [26]	*Apodemus sylvaticus*	SmallMammal	HRO	Indirect	93	92	0.000177	2
SleepyLizards [27]	*Tiliqua rugosa*	Ectotherm	GPSProximity	Proximity	546	199	1.678341	9
SleepyLizards [27]	*Tiliqua rugosa*	Ectotherm	HRO	Indirect	546	199	1.678341	9

					N	NID	Area	NYears
SoaySheep [28]	*Ovis aries*	LargeHerbivore	GOG	Proximity	15683	5670	2.145238	34
SoaySheep [28]	*Ovis aries*	LargeHerbivore	HRO	Indirect	13807	5006	1.775595	34
TayDolphins [29]	*Tursiops truncatus*	Cetacean	Census	Proximity	1208	266	1468.99	13
TayDolphins [29]	*Tursiops truncatus*	Cetacean	HRO	Indirect	201	94	266.2977	13
VulturineGuineafowl [30]	*Acryllium vulturinum*	Bird	GPSProximity	Proximity	3681	49	7.122599	5
VulturineGuineafowl [30]	*Acryllium vulturinum*	Bird	HRO	Indirect	3681	49	7.122599	5
WaterDragons [31]	*Intellagama lesueurii*	Ectotherm	CensusProximity	Proximity	3685	1254	0.054415	11
WaterDragons [31]	*Intellagama lesueurii*	Ectotherm	HRO	Indirect	2376	796	0.049776	11
WildCrickets [32]	*Gryllus campestris*	Ectotherm	Fight	Direct	1334	1334	0.000839	10
WildCrickets [32]	*Gryllus campestris*	Ectotherm	HRO	Indirect	1112	556	0.000537	10
WildCrickets [32]	*Gryllus campestris*	Ectotherm	Mate	Direct	1334	1334	0.000839	10
WoodchesterBadgers [33]	*Meles meles*	Carnivore	CoTrapping	Proximity	477	477	6.900926	8
WythamBadgers [34]	*Meles meles*	Carnivore	CoTrapping	Proximity	4993	1585	5.3095	32
WythamMice [35]	*Apodemus sylvaticus*	SmallMammal	CoTrapping	Proximity	274	274	0.00038	1
WythamTits [36]	*Parus major*	Bird	GOG	Proximity	2874	1966	12.29117	3
WythamTits [36]	*Parus major*	Bird	HRO	Indirect	1070	857	9.520905	3

974   Supplementary Table 1: List of study system replicates and their associated traits, with an
975   example reference from each. Also available as a supplementary file. N=number of
976   individual-by-year values; NID=number of unique individuals. Area=study extent in Km^2.
977   NYears=number of years covered by the study.

System	Species	Image credit	License link
VulturineGuineafowl	*Acryllium vulturinum*	James Klarevas	https://creativecommons.org/licenses/by/4.0/
WythamMice	*Apodemus sylvaticus*	Anthony Caravaggi	https://creativecommons.org/licenses/by-nc-sa/3.0/
SilwoodMice	*Apodemus sylvaticus*	NA	Public domain
LiverpoolMice	*Apodemus sylvaticus*	NA	Public domain
FalkirkMice	*Apodemus sylvaticus*	NA	Public domain
FieldingCows	*Bos taurus*	NA	Public domain
AfricanDogs	*Canis lupus*	NA	Public domain
ChagosSharks	*Carcharhinus amblyrhynchos*	NA	Public domain
PalmyraSharks	*Carcharhinus amblyrhynchos*	Russell Engelman	https://creativecommons.org/licenses/by/4.0/
MooreaSharks	*Carcharhinus melanopterus*	Ignacio Contreras	https://creativecommons.org/licenses/by/3.0/
RumDeer	*Cervus elaphus*	NA	Public domain

PrairieDogs	*Cynomys ludovicianus*	NA	Public domain
ReticulatedGiraffes	*Giraffa camelopardalis*	Cathy	https://creativecommons.org/licenses/by-nc-sa/3.0/
DesertTortoises	*Gopherus agassizii*	Andrew A. Farke, shell lines added by Yan Wong	https://creativecommons.org/licenses/by/3.0/
WildCrickets	*Gryllus campestris*	NA	Public domain
WaterDragons	*Intellagama lesueurii*	NA	Public domain
KenyanElephants	*Loxodonta africana*	Agnello Picorelli	https://creativecommons.org/licenses/by-nc-sa/3.0/
Multimammate Mice	*Mastomys natalensis*	David Liao	https://creativecommons.org/licenses/by-sa/3.0/
AcornWoodpeckers	*Melanerpes formicivorus*	NA	Public domain
WoodchesterBadgers	*Meles meles*	NA	Public domain
WythamBadgers	*Meles meles*	Anthony Caravaggi	https://creativecommons.org/licenses/by-nc-sa/3.0/
KielderVoles	*Microtus agrestis*	NA	Public domain
SoaySheep	*Ovis aries*	Gabriela Palomo-Munoz	https://creativecommons.org/licenses/by-nc/3.0/
WythamTits	*Parus major*	NA	Public domain

MountainLake	*Peromyscus leucopus*	Nina Skinner	https://creativecommons.org/licenses/by/3.0/
RockHyraxes	*Procavia capensis*	NA	Public domain
KalahariMeerkats	*Suricata suricatta*	NA	Public domain
GermanBoar	*Sus scrofa*	NA	Public domain
SleepyLizards	*Tiliqua rugosa*	CNZdenek	https://creativecommons.org/licenses/by-nc/3.0/
PotomacDolphins	*Tursiops truncatus*	Chris huh	https://creativecommons.org/licenses/by-sa/3.0/
MorayDolphins	*Tursiops truncatus*	NA	Public domain
AberdeenDolphins	*Tursiops truncatus*	Chris huh	https://creativecommons.org/licenses/by-sa/3.0/
SharkBayDolphins	*Tursiops truncatus*	Chris huh	https://creativecommons.org/licenses/by-sa/3.0/
GoldenCrownedSparrows	*Zonotrichia atricapilla*	NA	Public domain
SerengetiLions	*Panthera leo*	NA	Public domain
BeakedWhales	*Hyperoodon ampullatus*	Chris huh	https://creativecommons.org/licenses/by-sa/3.0/

978
979   Supplementary Table 2:The phylopic.org images used in the main text and their associated
980   licenses.

Variable	Estimate	P value	CI_Lower	CI_Upper
Intercept	0.191	7.09E-05	0.097	0.285
Behaviour: Spatial	0.263	5.56E-07	0.16	0.366
NID	0	0.217148	0	0
Area	0.017	0.213823	-0.01	0.043
NYears	0.005	0.060808	0	0.009

983
984
985   Supplementary Table 3: Meta-analysis effect estimates from the full meta-analytical model,
986   providing the estimate, P value, and 95% confidence intervals.
987

Supplementary figures

Supplementary Figure 1: Relationships between two measures of local population density
across animal systems. Density on the x axis was calculated using a kernel density
estimation approach; density on the y axis was calculated based on the number of
individuals located within a radius defined by the study system's area. X axis values have
been standardised to have a mean of zero and a standard deviation of 1 within each site; the
axis ticks are in units of 1 standard deviation. Each point represents an individual-year-
behaviour replicate; the lines portray the model fit from a generalised additive model (GAM).
Correlation coefficients were calculated using a Spearman's rank correlation.

Supplementary Figure 2: Linear relationships between density and network connectedness across animal systems. Density in individuals per area is on the x axis; network connectedness (strength centrality) is on the y axis. Both values have been standardised to have a mean of zero and a standard deviation of 1 within each system; the axis ticks are in units of 1 standard deviation. Each point represents an individual-year-behaviour replicate; the lines portray the model fit from our linear models for meta-analysis. Points are semi-transparent to enhance visibility. Panels are arranged phylogenetically following the tree displayed in Figure 2A; GOG=gambit of the group; HRO=home range overlap.

1008

1009 Supplementary Figure 3: Distributions of residuals from the linear models displayed in Figure
1010 2 above.

1011

1012
1013   Supplementary Figure 4: distribution of the Breusch-Pagan Test Statistic as a measure of
1014 heteroskedasticity (y axis) according to the type of contact being analysed (panel A) and the
1015   magnitude of the linear effect estimate (panel B). The y axis is log-transformed. These
1016 findings demonstrate no substantial difference in the levels of heteroskedasticity in the two
1017   contact types that might explain spatial models exhibiting a steeper density effect.
1018

1019
Supplementary Figure 5: Log-linear relationships between density and network
connectedness across animal systems. Density in individuals per area is on the x axis;
network connectedness (strength centrality) is on the y axis, both of which are then log(x+1)-
transformed. Each point represents an individual-year-behaviour replicate; the lines portray
the model fit from our linear models for meta-analysis. Points are semi-transparent to
enhance visibility. Panels are arranged phylogenetically following the tree displayed in
Figure 2A; GOG=gambit of the group; HRO=home range overlap.

Supplementary discussion

Operational implications: Operationally, the common nature of saturating density effects will impact researchers' ability to detect density dependence: that is, density dependence could be harder to observe in higher-density areas given the shallower slopes we observed. Most of the systems in this study are relatively long-term studies of known individuals; these populations tend to be in carefully selected, high-density areas that make it convenient to study the focal animal with relatively low operational costs. For example, it has previously been noted that the badgers of Wytham Wood, the red deer of the Isle of Rum, and the Soay sheep of St Kilda are all at high densities for their respective species [37–39]. As such, we may be inherently investigating the upper end of density-connectedness relationships in the wild, and it could be difficult and costly to investigate the effects of low density so widely.

Limitations: We acknowledge several limitations of our study, which we nevertheless believe could be remedied in the future. First, many of our social networks were formed of general spatiotemporal associations, and relatively few from specific social interactions – particularly those involving direct physical contact (only 3/36 social networks). Our current dataset could therefore benefit from supplementation with a broader range of direct interactions, particularly involving antagonism or bonding. For example, datasets concerning aggression or dominance interactions (e.g. [40]) or grooming alongside spatial behaviour could inform how density dependence affects the transmission of certain parasites such as mycobacteria [41] or tattoo skin disease [42]. The meta-dataset was also unevenly distributed across animal taxa (Figure 1): there were no primates or bony fishes and only one invertebrate, while rodents and ungulates were over-represented. These biases likely emerge through differences in data collection approaches: for example, although primate social behaviour is often studied with observations of direct interactions that could augment our data as described above (e.g. [43]), the spatial data required to build density distributions are rarely collected in these systems. This is linked to their social structures: our workflow was best suited to studies of fission-fusion societies or relatively asocial animals, rather than those with wide-ranging fixed social groups that are more common in primate systems. Finally, given that our data were observational, we could not account for (or estimate) bidirectional causality between density and social relationships (point 4 in Figure 1): that is, as well as encountering more conspecifics in areas of high density, individuals may be drawn to conspecifics, *creating* areas of high density [44]. To do so might require creating in-depth, high-resolution models of animal movement and group formation (e.g. [45]), potentially making use of telemetry approaches and drawing from large-scale open movement repositories like Movebank [46]. Using remote and next-generation approaches may help to access and incorporate more remote areas, which could also help to ameliorate the substantial geographic biases in our meta-dataset (Figure 2).

Analytical expansions: Aside from incorporating more specific interaction types, there exist a range of potential extensions to our analysis. For example, density dependence often varies between age or sex classes (e.g. [47,48]), and age effects on infection are common and ecologically important [49,50], as are sex differences [51]. We chose not to analyse how individual animals' traits alter the shape or slope of density's effects for brevity and simplicity; however, given that many of the systems nevertheless include these data, future analyses could make use of this meta-dataset to investigate how density affects connectedness of different classes of hosts. Further, researchers could investigate other behavioural questions

such as the role of observation biases; the factors influencing the correlation between spatial
and social networks; and the role of environmental drivers and spatial autocorrelation in
driving observed patterns of connectivity [44,52]. Finally, as our analysis approximated
density-contact relationships and not host-parasite interactions specifically, important future
work could investigate whether contact rates (as approximated by network connectedness)
ultimately translate to greater infection risk or parasite burden. Although some previous
investigations have linked density-related metrics to aspects of infection [53,54], density
covaries with a range of other factors including nutrition, cooperation, and competition, all of
which could complicate density-driven increases in exposure [55]. For example, in the case
of ectoparasite transmission, although contact rates in general would likely increase with
density, so too might grooming behaviours that remove parasites; in cases such as these,
density's overall effect on ectoparasite infection may be neutral. In the future, verifying that
within- and between-population variation in density-contact relationships translate to
variation in infection – and whether these trends might be influenced by flexible avoidance
behaviours [56] – will be a vital part of understanding and predicting density-dependent
disease dynamics.

Supplementary acknowledgements

Acorn woodpeckers

Sahas Barve and Eric L Walters were supported by funding from the National Science Foundation (Grant IOS-1455900). We would like to thank Walt Koenig, Jen Hunter, and many graduate students and field assistants who have helped contribute to this project and the Hastings Natural History Reservation for logistical support.

African dogs

This system acknowledges support from The Carter Center and the Natural Environment Research Council. Ciro Cattuto and Laura Ozella acknowledge partial support from the Lagrange Project of the ISI Foundation funded by CRT Foundation.

Bottlenose whales

Fieldwork and data management for this population were supported by Fisheries and Oceans Canada.

Chagos sharks

This population was funded by the Bertarelli Foundation. Ciro Cattuto and Laura Ozella acknowledge partial support from the Lagrange Project of the ISI Foundation funded by CRT Foundation.

Cornish cattle

This project acknowledges support from the Biotechnology and Biological Sciences Research Council.

European boar

Alisa Klamm (AK) is grateful to the Thuringian Ministry of Environmental, Energy and Nature Conservation and the Thuringian Ministry for Infrastructure and Agriculture for funding the research project on wild boar in the Hainich National Park. AK also thanks the people involved in the fieldwork and the copartners: Landesjagdverband Thüringen e.V. and ThüringenForst-Anstalt öffentlichen Rechts. Kevin Morelle (KM) was supported by the H2020 project VACDIVA (grant ID: 862874).

Golden-crowned sparrows

The research was supported by an NSF Graduate Research Fellowship to Theadora A Block, a CAREER grant (NSF IOS-1750606) and National Geographic grant (WW-R012-17) to Dai Shizuka, and a Laboratoire d'Excellence grant (TULIP, ANR-10-LABX-41) and ANR-SoCo to Alexis S Chaine.

Firth of Tay and Moray Dolphins

BJC and PSH thank all the colleagues who have helped collect and analyse the Moray Firth and Firth of Tay bottlenose dolphin data. Funding to support these long-term projects was provided by NatureScot, Beatrice Offshore Windfarm Ltd., Moray East and Moray West Offshore Wind Farms, Vattenfall, Seagreen, Inch Cape Offshore Ltd, NnG Offshore Wind, Marine Scotland, The Crown Estate, Highlands and Islands Enterprise, BES, ASAB, Greenpeace Environmental Trust, Scottish Government, Whale and Dolphin Conservation, Talisman Energy (UK) Ltd., Department of Energy and Climate Change, Chevron, Natural Environment Research Council and the Universities of Aberdeen and St Andrews.

Kalahari meerkats

This analysis uses data on meerkats collected on a project supported by European Research Council Grants numbers 742808 and 294494.

Kenyan elephants

We would like to thank the Kenyan Office of the President and Kenya Wildlife Service (KWS) for permission to conduct this research and Felix Mdamu and David Korir for help collecting the data. KWS funded radio collars, tracking equipment, and aerial surveys. The research was approved by the Institutional Animal Care and Use Committee (IACUC) of the University of California, Davis, protocol #10087 and received the permission of the Kenya Ministry of Education, Science, and Technology (Permit No. MOEST 13/001/34C 301). Funding for this work was provided by The Lincoln Park Zoo, International Elephant Foundation, and UC Davis Animal Behavior Graduate Group.

Liverpool wood mice

Data were collected under funding from NERC (grants: NE/G007349/1, NE/G006830/1, NE/I024038/1 and NE/I026367/1), and data collectors: Becci Barber, Trevor Jones, Łukasz Łukomski, Stephen Price, Susan Withenshaw, Niamh Quinn and many undergraduate assistants.

Mountain Lake mice

Andrea Graham was funded by Defense Advanced Research Projects Agency.

Potomac dolphins

This population was funded by the Earth Commons Institute, Georgetown University, and under National Geographic Grant WW-022ER-17.

Prairie dogs

We would like to respectfully acknowledge that our research on prairie dogs occurred on Treaty 4 territory, the traditional territory of the Očhéthi Šakówiŋ, Niitsítpiis-stahkoii ᖹᑳᖹᓇᒣᒪᖽᐧ ᖹᖿᖹᒪ, and the homeland of the Michif Piyii (Métis) Nation. We are very thankful to our field technicians for assistance with data collection. We thank S. Liccioli, H. Facette, and the staff at Grasslands National Park for their assistance with permitting and logistical

support. This research was supported by a Natural Sciences and Engineering Research Council Discovery Grant RGPIN 04093-2014 (J.E.L.), Parks Canada Contribution Agreement GC-794 (J.E.L.), Saskatchewan Fish and Wildlife Development Fund Student Research Award (J.M.K., and Colleen Crill Matzke), Nature Saskatchewan (J.M.K), and the University of Saskatchewan.

Rum deer

The authors thank NatureScot for permission to work on the Isle of Rum National Nature Reserve. The authors thank F. Guinness, M. Baker, A. Morris, S. Morris, and many others for collecting field data, and S. Albon and L Kruuk for their contributions to the long-term project. The project has been funded principally by the UK Natural Environment Research Council.

Serengeti lions

This study population acknowledges NSF grant DEB-1020479.

Shark Bay dolphins

This study population was funded by NSF grants 2146995, 2106909, 1755229, and 1559380.

Sleepy lizards

We thank Ron and Leona Clark and Chris Mosey for access to their land and use of the homestead at Bundey Bore Station. We thank Jess Clayton and Dale Burzacott for logistical support, and we are especially grateful to the late Professor Michael Bull and the late Dale Burzacott, who established the sleepy lizard monitoring project. We are also grateful for many students that worked on the sleepy lizard system along the years. (not sure naming them all is helpfull here, so i suggest leaving as is). This work was funded by National Science Foundation of the United States Grant DEB-1456730 and Australian Research Council Grants DP0877384 and DP130100145.

Soay sheep

The authors thank the National Trust for Scotland for permission to work on St Kilda, and QinetiQ and Kilda Cruises for logistical support. The authors thank J. Pilkington, I. Stevenson and many others for collecting the field data and D. Nussey, J. Slate and M. Morrissey for their contributions to the long-term project. The study on St Kilda has been funded principally by the UK Natural Environment Research Council.

Vulturine guineafowl

The vulturine guineafowl work was funded the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement number 850859) and Eccellenza Professorship Grant of the Swiss National Science Foundation (grant number PCEFP3_187058) awarded to DRF.

Wild crickets

WildCrickets would like to thank Luis Rodríguez and María del Carmen Muñoz for unconditional support and providing access to facilities including the WildCrickets study meadow. The following people contributed to video processing and data recording: David Fisher, Ian Skicko, Xing P. Liu, Thor Veen, Carlos Rodríguez del Valle, Alan Rees, Sophie Haugland Pedersen, Hannah Hudson, Jasmine Jenkin, Lauren Morse, Emma Rogan, Emelia Hiorns, Sarah Callow, Jamie Barnes, Chloe Mnatzaganian, Olivia Pearson, Adèle James, Robin Brown, Chris Shipway, Luke Meadows and Peter Efstratiou. The WildCrickets project was supported by the Natural Environment Research Council (NERC); standard grants: NE/E005403/1, NE/H02364X/1, NE/L003635/1, NE/R000328/1, NE/V000772/1 and the Leverhulme Trust.

Woodchester badgers

This study was funded by the UK Department for Environment, Farming and Rural Affairs, Animal and Plant Health Agency, and the Natural Environmental Research Council.

Wytham great tits

The Wytham great tit data were funded by ERC AdG 250164, NERC NE/S010335/1 and NE/D011744/1, and collected by the members of the Wytham Social Network Group.

Supplement references

1. Shizuka D, Barve S, Johnson AE, Walters EL. 2021 Constructing social networks from automated telemetry data: a worked example using within- and across-group associations in cooperatively breeding birds. *Methods Ecol Evol* , 0–2. (doi:10.1111/2041-210x.13737)

2. Wilson-Aggarwal JK, Ozella L, Tizzoni M, Cattuto C, Swan GJF, Moundai T, Silk MJ, Zingeser JA, McDonald RA. 2019 High-resolution contact networks of free-ranging domestic dogs Canis familiaris and implications for transmission of infection. *PLoS Negl Trop Dis* **13**, e0007565. (doi:10.1371/journal.pntd.0007565)

3. Gowans S, Whitehead H, Hooker SK. 2001 Social organization in northern bottlenose whales, Hyperoodon ampullatus: not driven by deep-water foraging? *Anim Behav* **62**, 369–377. (doi:10.1006/ANBE.2001.1756)

4. Jacoby DMP *et al.* 2020 Shark movement strategies influence poaching risk and can guide enforcement decisions in a large, remote marine protected area. *Journal of Applied Ecology* **57**, 1782–1792. (doi:10.1111/1365-2664.13654)

5. Sah P, Nussear KE, Esque TC, Aiello CM, Hudson PJ, Bansal S. 2016 Inferring social structure and its drivers from refuge use in the desert tortoise, a relatively solitary species. *Behav Ecol Sociobiol* **70**, 1277–1289. (doi:10.1007/s00265-016-2136-9)

6. Wielgus E, Klamm A, Conraths FJ, Dormann CF, Henrich M, Kronthaler F, Heurich M. 2023 First-Passage Time Analysis Based on GPS Data Offers a New Approach to Estimate Restricted Zones for the Management of Infectious Diseases in Wildlife: A Case Study Using the Example of African Swine Fever. *Transbound Emerg Dis* **2023**, 4024083. (doi:10.1155/2023/4024083)

7. Sweeny AR, Clerc M, Pontifes PA, Venkatesan S, Babayan SA, Pedersen AB. 2021 Supplemented nutrition decreases helminth burden and increases drug efficacy in a natural host-helminth system. *Proceedings of the Royal Society B: Biological Sciences* **288**. (doi:10.1098/rspb.2020.2722)

8. Fielding HR, Silk MJ, McKinley TJ, Delahay RJ, Wilson-Aggarwal JK, Gauvin L, Ozella L, Cattuto C, McDonald RA. 2021 Spatial and temporal variation in proximity networks of commercial dairy cattle in Great Britain. *Prev Vet Med* **194**, 105443. (doi:10.1016/j.prevetmed.2021.105443)

9. Madsen AE, Lyon BE, Chaine AS, Block TA, Shizuka D. 2023 Loss of flockmates weakens winter site fidelity in golden-crowned sparrows (Zonotrichia atricapilla). *Proc Natl Acad Sci U S A* **120**, e2219939120. (doi:10.1073/PNAS.2219939120/SUPPL_FILE/PNAS.2219939120.SAPP.PDF)

10. Bateman AW, Lewis MA, Gall G, Manser MB, Clutton-Brock TH. 2015 Territoriality and home-range dynamics in meerkats, Suricata suricatta: a mechanistic modelling approach. *Journal of Animal Ecology* **84**, 260–271. (doi:10.1111/1365-2656.12267)

11. Pinter-Wollman N, Isbell LA, Hart LA. 2009 The relationship between social behaviour
and habitat familiarity in African elephants (Loxodonta africana). *Proceedings of the*
Royal Society B: Biological Sciences **276**, 1009–1014. (doi:10.1098/rspb.2008.1538)

12. Wanelik KM, Farine DR. 2022 A new method for characterising shared space use
networks using animal trapping data. *Behav Ecol Sociobiol* **76**. (doi:10.1007/s00265-
022-03222-5)

13. Sweeny AR, Albery GF, Venkatesan SV, Fenton A, Pedersen AB. 2021
Spatiotemporal variation in drivers of parasitism in a wild wood mouse population.
Funct Ecol , 1–11. (doi:10.1111/1365-2435.13786)

14. Mourier J, Planes S. 2021 Kinship does not predict the structure of a shark social
network. *Behavioral Ecology* **32**, 211–222. (doi:10.1093/beheco/araa119)

15. Fisher DN, Cheney BJ. 2023 Dolphin social phenotypes vary in response to food
availability but not the North Atlantic Oscillation index. *Proceedings of the Royal*
Society B **290**. (doi:10.1098/RSPB.2023.1187)

16. Sweeny AR, Thomason CA, Carbajal EA, Hansen CB, Graham AL, Pedersen AB.
2020 Experimental parasite community perturbation reveals associations between Sin
Nombre virus and gastrointestinal nematodes in a rodent reservoir host. *Biol Lett* **16**.
(doi:10.1098/RSBL.2020.0604)

17. Mariën J *et al.* 2020 Density dependence and persistence of Morogoro arenavirus
transmission in a fluctuating population of its reservoir host. *Journal of Animal Ecology*
89, 506–518. (doi:10.1111/1365-2656.13107)

18. Jacoby DMP, Papastamatiou YP, Freeman R. 2016 Inferring animal social networks
and leadership: applications for passive monitoring arrays. *J R Soc Interface* **13**.
(doi:10.1098/RSIF.2016.0676)

19. Collier MA *et al.* 2023 Breathing in sync: how a social behavior structures respiratory
epidemic risk in bottlenose dolphins. *bioRxiv* (doi:10.1101/2023.12.01.569646)

20. Kusch JM, Lane JE. 2021 Determinants of social structure in a northern population of
black- tailed prairie dogs , Cynomys ludovicianus. *Anim Behav* **178**, 1–10.
(doi:10.1016/j.anbehav.2021.05.017)

21. VanderWaal KL, Wang H, McCowan B, Fushing H, Isbell LA. 2014 Multilevel social
organization and space use in reticulated giraffe (Giraffa camelopardalis). *Behavioral*
Ecology **25**, 17–26. (doi:10.1093/beheco/art061)

22. Barocas A, Ilany A, Koren L, Kam M, Geffen E. 2011 Variance in centrality within rock
hyrax social networks predicts adult longevity. *PLoS One* **6**, 1–8.
(doi:10.1371/journal.pone.0022375)

23. Albery GF, Clutton-brock TH, Morris A, Morris S, Pemberton JM, Nussey DH, Firth JA.
2022 Ageing red deer alter their spatial behaviour and become less social. *Nat Ecol*
Evol **6**, 1231–1238. (doi:10.1038/s41559-022-01817-9)

24. Rudd LF, Packer C, Biro D, Firth JA, Albery GF. 2024 Sex-specific social aging in wild
African lions. *Current Biology* (doi:10.1016/j.cub.2024.07.040)

25. Leu ST, Sah P, Krzyszczyk E, Jacoby A, Mann J. 2020 Sex, synchrony, and skin
contact: integrating multiple behaviors to assess pathogen transmission risk.
Behavioral Ecology , 1–10. (doi:10.1093/beheco/araa002)

26. Raulo A, Allen BE, Troitsky T, Husby A, Firth JA, Coulson T, Knowles SCL. 2021
Social networks strongly predict the gut microbiota of wild mice. *ISME J* , 1–13.
(doi:10.1038/s41396-021-00949-3)

27. Sih A, Spiegel O, Godfrey S, Leu S, Bull CM. 2018 Integrating social networks, animal
personalities, movement ecology and parasites: a framework with examples from a
lizard. *Anim Behav* **136**, 195–205. (doi:10.1016/j.anbehav.2017.09.008)

28. Sweeny AR, Corripio-miyar Y, Bal X, Hayward A, Pilkington JG. 2021 Seasonal
dynamics and repeatability of gastrointestinal parasite burdens in wild Soay sheep. ,
1–28.

29. Arso Civil M, Quick NJ, Cheney B, Pirotta E, Thompson PM, Hammond PS. 2019
Changing distribution of the east coast of Scotland bottlenose dolphin population and
the challenges of area-based management. *Aquat Conserv* **29**, 178–196.
(doi:10.1002/AQC.3102)

30. Papageorgiou D, Farine DR. 2020 Group size and composition influence collective
movement in a highly social terrestrial bird. , 1–16.

31. Piza-Roca C, Strickland K, Schoeman D, Frere CH. 2018 Eastern water dragons
modify their social tactics with respect to the location within their home range. *Anim
Behav* **144**, 27–36. (doi:10.1016/j.anbehav.2018.08.001)

32. Fisher DN, Rodríguez-Muñoz R, Tregenza T. 2019 Dynamic networks of fighting and
mating in a wild cricket population. *Anim Behav* **155**, 179–188.
(doi:10.1016/j.anbehav.2019.05.026)

33. Silk MJ, Weber NL, Steward LC, Hodgson DJ, Boots M, Croft DP, Delahay RJ,
McDonald RA. 2018 Contact networks structured by sex underpin sex-specific
epidemiology of infection. *Ecol Lett* **21**, 309–318. (doi:10.1111/ele.12898)

34. Newman C, Macdonald DW, Anwar MA. 2001 Coccidiosis in the European badger,
Meles meles in Wytham Woods: Infection and consequences for growth and survival.
Parasitology **123**, 133–142. (doi:10.1017/S0031182001008265)

35. Raulo A *et al.* 2024 Social and environmental transmission spread different sets of gut
microbes in wild mice. *Nature Ecology & Evolution 2024 8:5* **8**, 972–985.
(doi:10.1038/s41559-024-02381-0)

36. Firth JA, Sheldon BC. 2016 Social carry-over effects underpin trans-seasonally linked
structure in a wild bird population. *Ecol Lett* **19**, 1324–1332. (doi:10.1111/ele.12669)

37.   Macdonald DW, Newman C. 2022 *The Badgers of Wytham Woods: a model for*
1325   *behaviour, ecology, and evolution.* Oxford University Press.

1326   38.   Clutton-Brock TH, Guinness FE, Albon SD. 1982 *Red Deer: Behavior and Ecology of*
1327   *Two Sexes.* Chicago, IL: University of Chicago Press. See
1328   https://books.google.co.uk/books/about/Red_Deer.html?id=x4SGuA3t-NoC&pgis=1.

1329   39.   Clutton-Brock TH, Pemberton JM. 2004 *Soay sheep*□*: population dynamics and*
1330   *selection on St. Kilda.* Cambridge, UK: Cambridge University Press.

1331   40.   Strauss ED, Decasien AR, Galindo G, Hobson EA, Shizuka D, Curley JP. 2022
1332   DomArchive: a century of published dominance data. *Philosophical Transactions of*
1333   *the Royal Society B* **377**. (doi:10.1098/RSTB.2020.0436)

1334   41.   Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in
1335   wild meerkats. *Proceedings. Biological sciences / The Royal Society* **277**, 633–42.
1336   (doi:10.1098/rspb.2009.1775)

1337   42.   Powell SN, Wallen MM, Miketa ML, Krzyszczyk E, Foroughirad V, Bansal S, Mann J.
1338   2021 Sociality and tattoo skin disease among bottlenose dolphins in Shark Bay,
1339   Australia. *Behavioral Ecology* **31**, 459–466. (doi:10.1093/BEHECO/ARZ207)

1340   43.   Testard C *et al.* 2020 Rhesus Macaques Build New Social Connections after a Natural
1341   Disaster. *Current Biology* , 1–11. (doi:10.2139/ssrn.3741230)

1342   44.   Webber QMR, Albery GF, Farine DR, Pinter-wollman N, Sharma N, Vander Wal E,
1343   Manlove K. 2023 Behavioural ecology at the spatial-social interface. *Biological*
1344   *Reviews* **3**, 868–886.

1345   45.   Spiegel O, Leu ST, Sih A, Bull CM. 2016 Socially interacting or indifferent
1346   neighbours□? Randomization of movement paths to tease apart social preference
1347   and spatial constraints. *Methods Ecol Evol* , 971–979. (doi:10.1111/2041-
1348   210X.12553)

1349   46.   Kays R *et al.* 2021 The Movebank system for studying global animal movement and
1350   demography. *Methods Ecol Evol* **2021**, 1–13. (doi:10.1111/2041-210X.13767)

1351   47.   Batsuren E, Zhang X, Song M, Wan X, Li G, Liu J, Huang S, Zhang Z. 2022 Density-
1352   dependent changes of mating system and family structure in Brandt's voles
1353   (*Lasiopodomys brandtii*). *Ecol Evol* , 1–16. (doi:10.1002/ece3.9199)

1354   48.   Vander Wal E, van Beest FM, Brook RK. 2013 Density-Dependent Effects on Group
1355   Size Are Sex-Specific in a Gregarious Ungulate. *PLoS One* **8**, e53777.
1356   (doi:10.1371/JOURNAL.PONE.0053777)

1357   49.   Ashby B, Bruns E. 2018 The evolution of juvenile susceptibility to infectious disease.
1358   *Proceedings of the Royal Society B: Biological Sciences* **285**, 20180844.
1359   (doi:10.1098/rspb.2018.0844)

50. Albery GF, Sweeny AR, Webber Q. In press. How behavioural ageing affects
infectious disease.

51. Zuk M. 2009 The sicker sex. *PLoS Pathog* **5**, 1–3. (doi:10.1371/journal.ppat.1000267)

52. Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, Firth
JA. 2021 Multiple spatial behaviours govern social network positions in a wild
ungulate. *Ecol Lett* **24**, 676–686. (doi:10.1101/2020.06.04.135467)

53. Arneberg P, Skorping A, Grenfell B, Read AF. 1998 Host densities as determinants of
abundance in parasite communities. *Proceedings of the Royal Society B: Biological
Sciences* **265**, 1283–1289. (doi:10.1098/rspb.1998.0431)

54. Arneberg P. 2001 An ecological law and its macroecological consequences as
revealed by studies of relationships between host densities and parasite prevalence.
Ecography **24**, 352–358. (doi:10.1111/j.1600-0587.2001.tb00208.x)

55. Albery GF. 2022 Density dependence and disease dynamics: moving towards a
predictive framework. *EcoEvoRxiv Preprints* (doi:10.32942/OSF.IO/GAW49)

56. Albery GF, Newman C, Bright Ross J, Macdonald DW, Bansal S, Buesching CD.
2020 Negative density-dependent parasitism in a group-living carnivore. *Proceedings
of the Royal Society B: Biological Sciences* **287**, 20202655.
(doi:10.1101/2020.06.15.153726)

1378

1379

1380

1381

Supplementary methods – Study System Details

Acorn woodpeckers

We studied acorn woodpeckers at Hastings Natural History Reservation (36.387° N, 121.551° W) in central coastal California, USA. Adults on their natal territory with their social (and genetic) parents were categorized as nonbreeding helpers. Group members living outside their natal territories, or living with birds of the opposite sex that were nonrelatives, were considered putative breeders[1]. Extra-group mating, as well as incestuous mating, is rare in this study population[1,2]. From 1973 to 2021, the majority of the woodpecker population was color-banded (N = 6572 total individuals) and monitored continuously for group size and composition. Each year, territory quality was assigned to each acorn woodpecker social group territory based on the size of the group's granary (1: <1000 storage holes [low quality], 2: 1000–2500 [medium quality], 3: >2500 [high quality])[3].

Acorn woodpeckers were caught opportunistically and fitted with dorsally mounted solar-powered nanotags[4] with leg loop harnesses adjusted for body size[5]. All tags weighed less than 1% of each bird's body mass. Radio-tagged birds (N=87) were detected by an array of 43 permanently installed autonomous, solar-powered base stations during daylight hours[6] for a total of 497 days between August 2019 and July 2021.

Tags produced an encoded 64-bit, 2.5 ms radio ping every 1.5 s during the day, even in cloudy weather. Each detection of an individual at the base station was accompanied with a date, time, and signal-strength stamp. All detections were stored in files created every 15 min and stored on removable memory drives[7].

The raw data were first cleaned to retain only those detections that occurred with high signal strength (within ~100 m of the receiver). Next, we calculated time spent by each bird (N=87) at each receiver station (N=43). For this, we first partitioned each day into 1-min bins based on the first and the last detection of the day. Next, we assigned a bird to a receiver for a given 1-min bin if we detected that bird more than 10 times at a particular receiver within that minute. This resulted in a cumulative 821,262 minutes of bird presence at receivers.

We classified birds as "associating" if they were detected in the same 1-min bin at the same receiver on a given day. We classified associations as at "home" if the receiver was within 200 m of the bird's territory or "away" if the bird was detected with high signal strength at a receiver farther than 200 m from its own territory. We detected 175,368 minutes of association between birds (range 2–16 birds associating). Most common associations were between two birds (60,299 minutes of association) and least common association was among 16 birds (only one instance detected). Seventy-eight birds were detected at territories away from their home territory. Distance away from home was highly variable (mean ± SD 483.5 m ± 419 m, range 207–2898 m).

References

1. Koenig, W.D., Haydock, J., and Stanback, M.T. Reproductive roles in the cooperatively breeding acorn woodpecker: incest avoidance versus reproductive competition. Am. Nat. 151, 243–255 (1998).

2. Dickinson, J., Haydock, J., Koenig, W., Stanback, M., and Pitelka, F. Genetic monogamy
in single-male groups of acorn woodpeckers, Melanerpes formicivorus. Mol. Ecol. 4, 765–
769 (1995).

3. Koenig, W.D., Walters, E.L., and Haydock, J. Variable helper effects, ecological
conditions, and the evolution of cooperative breeding in the acorn woodpecker. Am. Nat.
178, 145–158 (2011).

4. Stanback, M.T., and Koenig, W.D. Techniques for capturing birds inside natural cavities.
J. Field Ornithol. 65, 70–75 (1994).

5. Rappole, J.H., and Tipton, A.R. New harness design for attachment of radio transmitters
to small passerines. J. Field Ornithol. 62, 335–337 (1991).

6. Barve, S., Hagemeyer, N.D., Winter, R.E., Chamberlain, S.D., Koenig, W.D., Winkler,
D.W., and Walters, E.L. Wandering woodpeckers: foray behavior in a social bird. Ecology
101, e02943 (2020).

7. Barve, S., Lahey, A.S., Brunner, R.M., Koenig, W.D., and Walters, E.L. Tracking the
warriors and spectators of acorn woodpecker wars. Curr. Biol. 30, R982–R983 (2020).

1437

African dogs

Field site: Dogs (*Canis familiaris*) in rural Chad were collared during the dry (5th March - 17th May 2018) and wet seasons (3rd August – 17th October 2018). Dog owners from six villages participated in the study; Medegue (11°01'48.8" N; 15°26'37.7" E), Ngakedji (9°11'16.5" N; 18°18'10.7" E), Kira (9°10'50.8"N; 18°17'00.3"E), Bembaya (9°11'33.6"N; 18°17'42.3"E), Marabodokouya (9°19'42.3" N; 18°43'20.0" E) and Tarangara; (9°08'19.8" N; 18°42'00.9" E)[1,2].

Spatial data: Collars were fitted with an iGotU GT600 GPS logger (Mobile Action Technology, New Taipei City, Taiwan). The fix interval was set to 10 minutes. In both the dry and wet field season, an initial two-week deployment was immediately followed by a second, longer deployment using new collars fitted with a modified GT600 unit with a larger battery. Spatial data were cleaned by removing locations taken up to 12 h after the collar was deployed and 12 h before collar recovery. Fixes with speeds greater than 20 km/h were removed and data were discarded for periods when dogs were known to be tied up.

Proximity sensor data: Dog collars were also fitted with a proximity sensor developed by the OpenBeacon project (http://www.openbeacon.org/) and the SocioPatterns collaboration consortium (http://www.sociopatterns.org/). The sensors exchange multiple radio packets per second and proximity is calculated by the attenuation; difference between the received and transmitted power[3–5]. An attenuation threshold of −70dbm was used, detecting encounters within 1–1.5m[5,6]. Inter-logger variability was assessed by comparing the number of packets emitted and received for all pairs of sensors. Data were discarded if there was evidence for deviations from the expected linear relationship between emitted and received radio packets. A contact event was identified when sensors exchanged radio packets for a minimum of 20 consecutive seconds. A contact event ended when the exchange of radio packets ceased in the subsequent 20s period.

References

1.   Wilson-Aggarwal, J. K. et al. Spatial and temporal dynamics of space use by free-ranging domestic dogs Canis familiaris in rural Africa. Ecological Applications 31, e02328 (2021).

2.   Wilson-Aggarwal, J. K. et al. Spatial-temporal dynamics of contact among free-ranging domestic dogs Canis familiaris in rural Africa. 2024.06.26.600798 Preprint at https://doi.org/10.1101/2024.06.26.600798 (2024).

3.   Cattuto, C. et al. Dynamics of Person-to-Person Interactions from Distributed RFID Sensor Networks. PLOS ONE 5, e11596 (2010).

4.   Isella, L. et al. What's in a crowd? Analysis of face-to-face behavioral networks. Journal of Theoretical Biology 271, 166–180 (2011).

5.   Wilson-Aggarwal, J. K. et al. High-resolution contact networks of free-ranging domestic dogs Canis familiaris and implications for transmission of infection. PLOS Neglected Tropical Diseases 13, e0007565 (2019).

1477   6.   Voirin, N. et al. Combining High-Resolution Contact Data with Virological Data to
1478   Investigate Influenza Transmission in a Tertiary Care Hospital. Infection Control & Hospital
1479   Epidemiology 36, 254–260 (2015).

1480

Bottlenose whales

Fieldwork was conducted during most summers between 1988 and 2023, and was nearly always done from a 12m sailing vessel, though other small vessels were used for brief periods of time. The vessel carried out non-systematic surveys of key northern bottlenose whale (*Hyperoodon ampullatus*) habitat on the Scotian Shelf, mostly focusing on the Gully, a large submarine canyon ~200km from the coast of Nova Scotia. The geographical distribution of effort within the Gully was haphazard, and varied between years. In a subset of years, smaller amounts of time were also spent in the adjacent Shortland and Haldimand canyons. For this analysis we consider observations from the Gully only, as data for adjacent canyons is likely too limited to estimate meaningful variation in spatial behaviour across individuals.

Latitude and longitude for each encounter with northern bottlenose whales, or "sighting", were measured using various technologies over the course of the long-term study. Loran-C was used from 1988-1992, accurate within ~400 m[1], followed by various iterations of increasingly accurate GPS. During sightings, northern bottlenose whales were photographed and photo-identification of individuals was done based on distinctive markings of the dorsal fin. Film photography was used initially, with digital cameras being phased in starting in 2007. IDs are generally side-specific (i.e., left or right), except for those individuals with very distinctive fins that can be unambiguously recognized from either side. For this and most other analyses, we restrict our focus to left-sided IDs only, not considering any right-sided photographs. All observations included in this dataset were from high-quality photographs (quality ratings "3" or "4" out of 4). Additional information can be found in the publicly available photo-identification guide for the northern bottlenose whale project[2].

Sightings of northern bottlenose whales were considered independent social events. Photo-identifications were linked to the nearest sighting by date and time, from which we drew latitude and longitude data. Any photographs lacking a sighting record within 60 minutes were excluded from further analysis. This primarily included a smaller sets of photographs provided by other research vessels for which we lacked comparable location data. This resulted in 1199 groups over 23 years and a total of 615 unique individuals. Many individuals were seen just once (N = 199). On average, individuals were observed in 5 groups, range 1-66. Observations of groups lasted 10.8 minutes on average (time between first and last photo-identification), with a range of 0-107 minutes.

References

1. Hooker, S.K., Whitehead, H., Gowans, S., Baird, R.W. Fluctuations in distribution and patterns of individual range use of northern bottlenose whales. Marine Ecology Progress Series 225, 287–297 (2002).

1517   2. Feyrer, L. & Walmsley, S.F. A guide to northern bottlenose whale photo ID: Version 4.0.

1518   Open Science Framework doi:10.17605/osf.io/xd7ek (2024).

Chagos sharks

Grey reef sharks (Carcharhinus amblyrhynchos) have a broad Indo-Pacific distribution and are one of the most common elasmobranch predators on the reefs found across the Chagos Archipelago (6°00′S 71°30′E / 6.000°S 71.500°E). This study utilised acoustic telemetry to track the movements and distribution of sharks across 93 monitoring locations centred around areas of ecological interest (e.g. islands, islets, atolls and seamounts). Information on the monitoring array and tagging procedure can be found in Jacoby et al. 2020[1].

While tracking of this species occurred continuously between 2013 – 2021, data for this analysis were taken from 2014-2016 when tags at liberty, and thus density of individuals monitored were highest. Tags (Innovasea V16, 69 kHz coded transmitters) acoustically transmit a unique ID code at regular intervals (nominal delay of either 30–90 s or 60–180 s) for the duration of their battery life (~10 years). Tagged animals were detected whenever they came within range (~500 m) of an acoustic receiver. Spatio-temporal co-occurrences of tagged sharks were extracted from the telemetry data stream using a Gaussian mixture modelling approach [2], and implemented using the 'gmmevents' function in the R package asnipe [3]. Group-by-individual (GBI) matrices that reflected all associations between tagged individuals within a year, were extracted from the model, alongside information on the timing and location of group to the nearest acoustic receiver location. The response variable of social network position and the associated geographic location of social behaviour were derived from these GBI matrices. A total of 142 individuals could contribute to a group throughout the duration of the study.

References

1.   Jacoby, D. M. P. et al. Shark movement strategies influence poaching risk and can guide enforcement decisions in a large, remote Marine Protected Area. J. Appl. Ecol. 1–11 (2020) doi:10.1111/1365-2664.13654.

2.   Jacoby, D. M. P., Papastamatiou, Y. P. & Freeman, R. Inferring animal social networks and leadership: applications for passive monitoring arrays. J. R. Soc. Interface 13, 20160676 (2016).

3.   Farine, D. R. Animal social network inference and permutations for ecologists in R using asnipe. Methods Ecol. Evol. 4, 1187–1194 (2013).

Cornish cattle

Farms and cattle management

Proximity and GPS data were collected from 8 groups of dairy cows on 8 commercial farms in South-West England in Summer and Autumn of 2018 for approximately 7 days per farm. One group consisted of dry cows, whereas all other groups were lactating cows. Some devices malfunctioned and therefore we were not able to obtain data for all cattle in the group (60-91% of the group recorded[1]). Grazing management included rotational grazing, strip grazing, set stocking and free range[1]. Mostly, groups were kept on pasture and brought into buildings only for milking (twice daily), except one group were housed at night, one group were allowed access to buildings at all times and were kept in for two nights and days in the middle of the study period due to inclement weather and one group were allowed free access to all pasture, cubicle housing, and the automated milking system at all times during the study.

Equipment

Nylon cattle collars with a plastic clasp (Suevia Haiges, Germany) were fitted with a proximity device and a GPS logger such that one device lay at either side of the animal's neck. The GPS loggers (i-GotU GT-120 and GT-600 devices, Mobile Action Technology Inc., Taiwan) recorded fixes every ten minutes. The proximity device is based on a design by the OpenBeacon project (http://www.openbeacon.org/) and the SocioPatterns collaboration consortium (http://www.sociopatterns.org/) and has been used in contact studies of humans, horses, dogs, and sheep [2–5]. The proximity sensors exchange low-power radio packets in a peer-to-peer fashion, and this exchange of radio packets is used as a proxy for the proximity of individuals wearing the sensors. To estimate distance between devices, the attenuation of the signals with distance is computed as the difference between the received and transmitted power [2]. A contact event occurs if at least one data packet is exchanged during a continuous 20-second time window, and a contact is considered broken if no packets are exchanged in a 20-second period [6], therefore, contact durations were measured in 20-second blocks.

References

1.   Fielding, H. R. et al. Spatial and temporal variation in proximity networks of commercial dairy cattle in Great Britain. Prev Vet Med 194, 105443 (2021).

2.   Cattuto, C. et al. Dynamics of person-to-person interactions from distributed RFID sensor networks. PLoS One 5, e11596 (2010).

3.   Milwid, R. M., O'Sullivan, T. L., Poljak, Z., Laskowski, M. & Greer, A. L. Comparing the effects of non-homogenous mixing patterns on epidemiological outcomes in equine populations: A mathematical modelling study. Scientific Reports 9, 3227 (2019).

4.   Ozella, L. et al. The effect of age, environment and management on social contact patterns in sheep. Applied Animal Behaviour Science 225, 104964 (2020).

5.   Wilson-Aggarwal, J. K. et al. High-resolution contact networks of free-ranging
domestic dogs Canis familiaris and implications for transmission of infection. PLoS
Neglected Tropical Diseases 13, 1–19 (2019).

6.   Kiti, M. C., Melegaro, A., Cattuto, C. & Nokes, D. J. Study design and protocol for
investigating social network patterns in rural and urban schools and households in a coastal
setting in Kenya using wearable proximity sensors [version 2; peer review: 2 approved].
Wellcome Open Research 4, 1–17 (2019).

1597

1598

1599

Desert tortoises

We monitored tortoise movements using radio telemetry across multiple study sites spanning a 15-year period. Datasets were collected from nine study sites across desert tortoise habitat in the Mojave Desert of California, Nevada, and Utah from 1996 to 2014. Each site was monitored over multiple years, though not all sites were surveyed every year. Individuals were tracked at least weekly during their active season and at least monthly during winter months. Tortoises were fitted with very high frequency (VHF) radio-transmitters (e.g., Model RI-2B [13.8–15.0 g], Holohil Systems Ltd., Carp, Ontario, Canada, or AVM models G3, SB2, or SB2-RL, AVM Colfax CA for older studies), which were attached following established methods (Boarman et al. 1998). Locations were determined using hand-held VHF receivers (e.g., Telonics TR-2, Mesa, AZ, or ICOM RC 10) and recorded with GPS units (Universal Transverse Mercator, North American Datum 1983). Periods of intensive tracking (i.e., multiple relocations per day) were conducted to obtain detailed habitat use data during peak activity for some studies.

Each tortoise was uniquely identified with a numbered paper tag sealed with clear epoxy and permanent notches on the marginal scutes (Cagle 1939). During each encounter, we recorded the individual's identification number, date and time of observation, GPS location, microhabitat type (e.g., vegetation, pallet, or burrow), and any visible signs of injury or disease. Burrow use was documented by assigning a unique identification number to each burrow, with new IDs assigned when previously unmarked burrows were encountered. Den sharing was used as the contact type, where observations of individuals witnessed within the same den in the same sampling date were taken to be connected.

The dataset was based on monitoring tagged individuals, so data collection was not blind. While most tortoises were monitored consistently throughout the study, logistical constraints and equipment failures occasionally altered the telemetry schedule.

References

Boarman, W. I., T. Goodlett , G. Goodlett, and P. Hamilton. 1998. Review of radio transmitter attachment techniques for turtle research and recommendations for improvement. Herpetological Review 29: 26-27

Cagle, F. R. 1939. A system for marking turtles for future identification. Copeia 1939: 170-173.

Sah, P.*, Nussear, K.E., Esque, T.C., Aiello, C.M., Hudson, P.J., Bansal, S., 2016. Inferring social structure and its drivers from refuge use in the desert tortoise, a relatively solitary species. Behavioral Ecology and Sociobiology 70, 1277–1289. https://doi.org/10.1101/025494

Ein Gedi hyraxes

The rock hyrax (Procavia capensis) population has been studied since 1999 at the Ein Gedi Nature Reserve (31°28′N, 35°24′E) in Israel (e.g. 1–4). Data used in this study were collected continuously for the years 2000–2020, totaling approximately 75,000 observations. The research focuses on two deep gorges, David and Arugot, located on the western side of the Dead Sea. Each field season, beginning in March and lasting between three and six months, rock hyraxesare trapped and observed daily. Tomahawk live box traps were deployed in secure, shaded locations, baited with cabbage and kohlrabi. Given that rock hyraxes are diurnal, the traps are opened during a fixed morning interval. Once captured, the animals are anesthetized with ketamine hydrochloride (0.1 mg/kg) and receive a subcutaneous microchip (DataMars SA) along with either an ear tag (for pups and juveniles) or a lightweight collar (weighing less than 5 g) bearing identification marks. Each individual is then sexed, weighed, and measured, and is given 90 to 150 minutes to recover from the anesthesia. Animals that are recaptured are not anesthetized; they are simply weighed and are promptly released.

Hyrax activity was monitored each day during the field season using 10×42 binoculars and a telescope with 50–100× magnification. Observations were conducted in the early morning, from first light until noon, the time when the hyraxes retreat to their shelters. Every day, a randomly chosen focal group was followed. Although hyraxes primarily spend their time foraging and resting, which allows for the simultaneous monitoring of several individuals, limitations in visibility (due to rocks, trees, and bushes) and the challenge of following up to ten individuals meant that we could not capture the precise duration of every pairwise interaction. Consequently, we recorded interactions on a daily basis: if two individuals were seen interacting at any point during a day, that day was noted as an interaction event regardless of its length. In addition, we recorded the location of each observed hyrax, up to 5m accuracy.

Our annual marking efficiency was high, with about 95% ± 0.5 of group members identified, thus minimizing bias in assessing each group's social structure. Social interactions that involved unmarked individuals were excluded from analysis. We define positive interactions as those involving direct physical contact (such as huddling or sharing a sleeping burrow) or coordinated behavior (moving together closely or sitting side by side).

References

1.   Bar Ziv, E. et al. Individual, social, and sexual niche traits affect copulation success in a polygynandrous mating system. Behav Ecol Sociobiol 70, 901–912 (2016).

2.   Barocas, A., Ilany, A., Koren, L., Kam, M. & Geffen, E. Variance in Centrality within Rock Hyrax Social Networks Predicts Adult Longevity. PLoS ONE 6, e22375 (2011).

3.   Ilany, A., Barocas, A., Koren, L., Kam, M. & Geffen, E. Structural balance in the social networks of a wild mammal. Animal Behaviour 85, 1397–1405 (2013).

1676   4.   Bordes, C. N. M., Beukeboom, R., Goll, Y., Koren, L. & Ilany, A. High-resolution
1677   tracking of hyrax social interactions highlights nighttime drivers of animal sociality. Commun
1678   Biol 5, 1–12 (2022).

1679

1680

1681

1682

European boar

Behavioral data for wild boar were collected from three European study areas: Bavarian Forest National Park (BFNP, 48°59′N, 13°23′E, Germany), Hainich National Park (HNP, 51°04′N, 10°26′E, Germany), and Kostelec nad Černými lesy (KNC, 50°0′N, 14°50′E, Czech Republic).

BFNP data were collected between October 2021 and January 2023 as part of a research project on movement ecology and African swine fever (ASF) transmission dynamics 1. The study spanned the 250 km² BFNP and the adjacent 684 km² Šumava National Park in the Czech Republic, with elevations ranging from 570 to 1453 m. The landscape consists of mixed coniferous and mountain forests. A total of 42 wild boar were GPS-collared (Vertex Plus, Vectronic Aerospace GmbH, Berlin, Germany) to track movement patterns, home range sizes, and habitat use. Animals were captured using ~30 m² wood-clad corral traps equipped with live-monitoring cameras and counterweight-triggered gates. The captured individuals were restrained in a net tunnel, handled for approximately 5 min, and released. Ethical approval was granted by the Upper Bavaria government (permit ROB-55.2-2532.Vet-02-20-149).

HNP data were collected between April 2017 and August 2019 to investigate the effects of no-hunting zones on wild boar movement and space use 2. The 25,000 ha study area comprises 54.6% agricultural land, 34.8% forests, 7.3% open land, 3.2% anthropogenic areas, and 0.4% water bodies, with elevations ranging from 180 to 500 m a.s.l. HNP is 75 km² large, with 33 km² designated as a no-hunting zone, including UNESCO-listed primeval beech forests and former military sites where hunting is prohibited. A total of 63 wild boar were GPS-collared (Vertex Lite, Vectronic Aerospace GmbH, Berlin, Germany), with GPS fixes recorded at 30-minute intervals.

KNC data were collected between April 2019 and October 2022 as part of a study on wild boar movement ecology, social behavior, and responses to ASF control measures (including hunting pressure, artificial food supply, and movement barriers–electric and odor fences, and sound traps) 3,4. The 2,900 ha study area, located ~30 km east of Prague, comprises mixed forests, agricultural land, water bodies, and built-up areas, with an average elevation of 430 m a.s.l. Managed by the Czech University of Life Sciences Prague (Lesy ČZU), this area is heavily frequented for tourism, forestry, and hunting. A total of 84 wild boar were captured using wooden corral traps baited with corn and immobilized via anesthetic darts delivered by airguns. Each individual was GPS/GSM-collared (Vectronic Aerospace GmbH), with fixes recorded every 30–60 min. Ethical approval was granted by the Ministry of the Environment of the Czech Republic (permit MZP/2019/630/361).

For all three datasets, social networks were constructed using the gambit of the group approach, where individuals were considered to be associating if they were within 100 m of each other within a 10-minute interval. GPS data were processed using the spatsoc package 5,6 in R, with group_times used to assign temporal groupings and edge_dist used to calculate pairwise distances between individuals, generating an edge list of proximity-based associations.

References

1. Wielgus, E. et al. First-Passage Time Analysis Based on GPS Data Offers a New Approach to Estimate Restricted Zones for the Management of Infectious Diseases in Wildlife: A Case Study Using the Example of African Swine Fever. Transboundary and Emerging Diseases 2023, 4024083 (2023).

2. Klamm, A. et al. Entwicklung Und Raumnutzung Eines Schwarzwild-Bestandes in Abhängigkeit von Den Naturräumlichen Gegebenheiten Des Buchenwald-Nationalparks Hainich Und Dessen Intensiv Landwirtschaftlich Genutzten Umfeldes. Abschlussbericht Dezember 2020. (2020).

3. Olejarz, A., Augustsson, E., Kjellander, P., Ježek, M. & Podgórski, T. Experience shapes wild boar spatial response to drive hunts. Sci Rep 14, 19930 (2024).

4. Faltusová, M. et al. Wild Boar Proves High Tolerance to Human-Caused Disruptions: Management Implications in African Swine Fever Outbreaks. Animals 14, 2710 (2024).

5. Robitaille, A. L., Webber, Q. M. R. & Wal, E. V. Conducting social network analysis with animal telemetry data: Applications and methods using spatsoc. Methods in Ecology and Evolution 10, 1203–1211 (2019).

6. Robitaille, A. L., Webber, Q. & Wal, E. V. spatsoc: Group Animal Relocation Data by Spatial and Temporal Relationship. (2023).

Falkirk wood mice

Trapping for this study took place from 2015-2017 in Callendar Wood (55.990470, -3.766636; Falkirk, Scotland), a 100ha broadleaf woodland, which contain a populations of wood mice, which are naturally exposed to and infected with a wide range of parasites and pathogens. The experiment had three temporal replicates; all of which took place during the wood mouse breeding season: (i) May - July 2015 (ii) June - August 2016 and (iii) July-November 2017. Experimental design for these experiments included randomised nutrition supplementation (high quality food pellets vs unmanipulated) at the grid level and randomised parasite treatment (anthelmintic treatment vs water control) at the individual level within grids. We used a weight-adjusted of 2ml/g dose of both Pyrantel pamoate (Strongid-P, 100 mg/kg) and Ivermectin (Eqvalan, 9.4mg/kg). In 2017, treatment was re-administered 4 weeks after initial dose for all individuals still in the experiment. All animal work was carried out under the approved UK Home Office Project License 70/8543 in accordance with the UK Home Office.

Trapping grids were set up with trapping stations 10m apart, and two traps at each stations, as follows: 2015: 3 grids with 8x8 trapping stations; 2016: 4, 6x5 grids; 2017: 4, 7x5 grids. In each year we live-trapped mice for 3 nights/week using Sherman live traps (H.B. Sherman 2x2.5x6.5 inch folding trap, Tallahassee, FL, USA). Trapping protocols followed (Sweeny, Clerc, et al., 2021). Each tagged individual was followed for a period of 12-16 days (2015-2016) or up to 8 weeks (2017). During this time, we collected morphometric data, blood samples, and faecal samples at every capture. Movement was minimal was minimal between grids.

A total of 261 individuals were captured 882 times across this 3 year study. We used the easting and northing location of the trapping stations where individuals were trapped to estimate density and social network metrics. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night.

References

Sweeny, A. R., Clerc, M., & Pontifes, P. A. (2021). Supplemented nutrition decreases helminth burden and increases drug efficacy in a natural host–helminth system. Proceedings of the Royal Society B: Biological Sciences. https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2020.2722

Golden-crowned sparrow

Golden-crowned sparrow flocks have been monitored at the University of California-Santa Cruz Arboretum since 2009 as part of a long-term research study on their winter social behavior (Shizuka et al. 2014; Madsen et al. 2023). Each fall, first-year migrants were fitted with colored plastic and numbered metal leg bands in unique color combinations. Foraging flocks were surveyed throughout the Arboretum by recording the identities of all banded birds in a flock, which was defined as a group of birds foraging within a 5 m radius. Locations of flocks were recorded using a reference map of 10 m x 10 m grid cells (Shizuka et al. 2014). Observations at seed piles were not considered in this definition of foraging flocks and were excluded from our analysis of flocking relationships. To ensure we were observing winter flocking behaviors rather than interactions between transitory individuals on migration, we limited observations to between November 1 (when most birds had arrived from breeding grounds) and March 1 (when winter flocks begin to break apart and birds begin migration). We further limited our sample to birds that had been sighted 3 or more times throughout this period to remove transient individuals. We then inferred social associations using the 'gambit of the group'. from co-occurrences in foraging flocks and calculated the simple ratio index (SRI) to represent association strength (see Shizuka et al. 2014 for further details).

References

Shizuka, D., Chaine, A.S., Anderson, J., Johnson, O., Laursen, I.M., Lyon, B.E., 2014. Across-year social stability shapes network structure in wintering migrant sparrows. Ecol Lett 17, 998–1007. https://doi.org/10.1111/ele.12304

Madsen, A.E., Lyon, B.E., Chaine, A.S., Block, T.A., Shizuka, D., 2023. Loss of flockmates weakens winter site fidelity in golden-crowned sparrows (Zonotrichia atricapilla). Proc. Natl. Acad. Sci. U.S.A. 120, e2219939120. https://doi.org/10.1073/pnas.2219939120

Kalahari meerkats

An individual-based study of meerkats has been running at the Kuruman River Reserve in South Africa since 1993 (26°58□S, 21°49□E, Clutton-Brock and Manser 2016). The study area, which covers approximately 50–60 km2, includes a diverse landscape of dry pans, vegetated sand dunes, and arid bushveld that is typical of a South African savannah, where livestock and game farming form the principal land use. Since the start of the project, approximately 15 groups (range□=□6 to 21 groups) and around 200 individuals were followed at any one time (mean ± SD per month□=□214.5 ± 59.4, range□=□101 to 359). Most individuals were born into the study population and were habituated from birth to allow close behavioural observation. Groups were visited 3–4 times per week in the morning and afternoon throughout the study, with data collected on the composition of groups and on the behaviour, reproductive status, social status, and health status of individuals, so that pregnancies, births, deaths, emigrations and immigrations could be enumerated (summarised in Clutton-Brock and Manser 2016). Most individuals in the population were also weighed at each visit by enticing them onto electronic scales with small amounts of hard-boiled egg or water, and while foraging, GPS locations were taken from the center of the group at 15 min intervals, which we use to estimate home ranges. Though the project began in 1993, all the above data were only collected on multiple groups (> 5) simultaneously from 1998 onwards. Most of our analyses therefore cover the breeding seasons from 1998–2023. The only exception is the GPS data, which were collected in the form needed for our analyses from 2002 onwards. Individuals witnessed within the same group in the same sampling date were taken to be connected.

References

Clutton-Brock, Tim H; Manser, Marta B (2016). Meerkats: Cooperative breeding in the Kalahari. In: Koenig, Walter D; Dickinson, Janis L. Cooperative breeding in vertebrates : studies of ecology, evolution, and behavior. Cambridge: Cambridge University Press, 294-317.

Kenyan elephants

1834

1835 Relevant methods from:

1836 The relationship between social behavior and habitat familiarity in African elephants
1837 (*Loxodonta africana*). Noa Pinter-Wollman, Lynne A. Isbell and Lynette A. Hart. Proc. R.
1838 Soc. B (2009) 276, 1009–1014. doi:10.1098/rspb.2008.1538

1839

Translocation and sightings

During September 2005, 150 African elephants were translocated from Shimba Hills
National Reserve on the coast of Kenya (4.08 S to 4.38 S and 39.58 E to 39.38 E) to Tsavo
East National Park (2.08S to 3.78S and 38.18E to 39.38 E), a distance of 160 km. This
translocation was part of the Kenya Wildlife Service (KWS) effort to decrease human–
elephant conflict in the vicinity of Shimba Hills. Twenty elephant groups comprising adult
females, juveniles and calves (average group size 6.8) and 20 independent adult males
were moved over the course of 32 days. The release site differs ecologically from the source
site and is separated from it by dense human population, providing a unique opportunity for
examining the social behaviour of the elephants in a novel environment.

During the translocation, all the elephants were tagged with yellow zip ties on their tails to
distinguish them from the local Tsavo elephant population. Unique white numbers painted on
the translocated elephants' backs, natural ear marks and tusk shapes were used for
individual identification of the translocated elephants (Moss 1996). Elephants' ages were
estimated based on Moss (1996).

The locations, their time and the identities of the translocated and local Tsavo elephants
were recorded in Tsavo East for a year post-translocation using a Geko 201 GPS unit
(Garmin Ltd., USA). Road transects were conducted using a vehicle four to five times a
week, alternating between four routes of similar length (50–70 km) on the existing roads
within Tsavo East National Park. A total of 3371 elephant sightings were recorded, of which
386 and 2985 were the translocated and local elephants, respectively. Of the 150 elephants
translocated, data on 83 were obtained, and are presented here. Because males leave the
social unit in which they were born at the age of 15, and because the social behaviour of
these independent males differs from that of females and their young offspring (Moss &
Poole 1983), such translocated males were excluded from our analyses.

1865

Social association

Elephants were defined as associating with one another if they were sighted within 500 m
from one another within a 2 hour time period, based on McComb et al. (2000, 2003). They
showed that elephants can individually recognize conspecifics' vocalizations over great
distances (1 km). Therefore, the definition of social association used here includes not only
direct interactions but also recognizes the communicative capabilities of elephants to acquire
information about the number and identities (translocated or local) of vocalizing conspecifics

(McComb et al. 2000, 2003). Thus, the definition of social association used here allows for
the acquisition of inadvertent social information about the new environment (Danchin et al.
2004).

1876

References

Danchin, E., Giraldeau, L. A., Valone, T. J. & Wagner, R. H. 2004 Public information: from
nosy neighbors to cultural evolution. Science 305, 487–491. (doi:10.1126/science.1098254)

McComb, K., Moss, C., Sayialel, S. & Baker, L. 2000 Unusually extensive networks of vocal
recognition in African elephants. Anim. Behav. 59, 1103–1109.
(doi:10.1006/anbe.2000.1406)

McComb, K., Reby, D., Baker, L., Moss, C. & Sayialel, S. 2003 Long-distance
communication of acoustic cues to social identity in African elephants. Anim. Behav. 65,
317–329. (doi:10.1006/anbe.2003.2047)

Moss, C. 1996 Getting to know a population. In Studying elephants (ed. K. Kanwanga), pp.
58–74. Nairobi, Kenya: African Wildlife Foundation.

Moss, C. J. & Poole, J. H. 1983 Relationships and social structure of African elephants. In
Primate social relations: an integrated approach (ed. R. A. Hinde), pp. 315–325. Oxford, UK:
Blackwell Scientific Publications.

1891

Kielder voles

Briefly, we monitored a natural population of field voles (*Microtus agrestis*) in Kielder Forest, Northumberland, UK across three studies: 2001–2007, 2008–2010 and 2015–2017. Kielder Forest is a man-made spruce forest and field voles are found in grassy clear-cuts. Trapping was performed across different sites, each a forest clear-cut. At each site, Ugglan small mammal traps (Grahnab, Gnosjo, Sweden) were laid out in a grid and checked regularly in 'primary' and 'secondary' sessions (see below for details). Access to the sites was provided by Forestry England. Newly trapped field voles were injected with a Passive Integrated Transponder (PIT) tag (AVID, UK) for unique identification. At each subsequent capture, we recorded: date, site, trap location and PIT tag ID. This approach allowed us to build up a longitudinal record of the location of tagged voles across multiple sessions. Edges in the social network between individuals (nodes) were defined as unique voles trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night.

Between 2001-2007, trapping was performed from March to November. Primary sessions took place at monthly intervals. At each 'primary' session, voles were trapped at 4 different sites. At each site, 100 Ugglan small mammal traps were laid out in a grid spaced 5 m apart. At each 'primary' session, traps were checked a total of five times ('secondary sessions'). More details for the 2001-2007 study are available in[1].

Between 2008-2010, trapping was performed either from February (2008–2009) or April (2009–2010) to November. Primary sessions took place at monthly intervals. Voles were trapped at a total of 4 different sites – two sites in 2008–2009 and a further two sites in 2009–2010. At each site, 150 Ugglan small mammal traps were laid out in a grid spaced 3–5 m apart. Primary sessions lasted 3 days, and traps were checked a total of five times. More details for the 2008-2010 study are available in[2].

Between 2015-2017, trapping was performed from March to October. Primary sessions took place at approximately two-week intervals. At each primary session, voles were trapped at 4 different sites. During the study, 3 sites were reassigned due to practical constraints, giving a total of 7 different sampling sites. At each site, 150–197 Ugglan small mammal traps were laid out in a grid spaced 3–5 m apart. Primary sessions lasted 3 days, with traps checked each morning and afternoon. More details for the 2015-2017 study are available in[3].

References

1.   Begon, M. et al. Effects of abundance on infection in natural populations: Field voles and cowpox virus. Epidemics 1, 35–46 (2009).

2.   Jackson, J. A. et al. An Immunological Marker of Tolerance to Infection in Wild Rodents. PLOS Biology 12, e1001901 (2014).

3.   Wanelik, K. M. et al. Effects of an IgE receptor polymorphism acting on immunity, susceptibility to infection, and reproduction in a wild rodent. eLife e77666 (2023).

Liverpool wood mice

Live-trapping for this study was carried out in wild wood mouse populations located near Liverpool, UK regularly between May and December for six consecutive years (2009–2014). We sampled 16 trapping grids ranging in size from 2,500 to 10,000 m2, spread across five different woodland sites with 2–3 sites trapped per year. Sites ranged from approximately 2 to 60 km apart. On each grid, trapping stations were placed every 10 m, with two live traps (H.B. Sherman 2 × 2.5 × 6.5 in. folding traps, Tallahassee, FL, USA) at each station baited with grains and bedding material. Full trapping details can be found in (Sweeny, Albery, et al., 2021).

A total of 926 individuals were captured 1,609 times across this 6-year study (max captures per individual = 28, median captures per individual = 4). We used the easting and northing location of the trapping stations where individuals were trapped to estimate density and social network metrics. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night. Density was calculated using individuals' centroids in the sampling year based on annual density kernels and using trapping locations.

References

Sweeny, A. R., Albery, G. F., Venkatesan, S., Fenton, A., & Pedersen, A. B. (2021). Spatiotemporal variation in drivers of parasitism in a wild wood mouse population. Functional Ecology, 35(6), 1277–1287.

Moorea sharks

Owner: Johann Mourier & Serge Planes

Location: Moorea, French Polynesia

The study was conducted at Moorea Island (17°30'S; 149°51'W) in the Society archipelago, French Polynesia. Between 2008 and 2010, a total of seven sites were surveyed on a regular basis along approximately 10 km of the Northern reef of Moorea Island. Among them, six sites were located on the outer reef and were characterized by coral structures from the barrier reef to the drop off (70 m depth) and one site was located inside the lagoon between 2 and 10 m depth within a small channel and characterized by coral patches in a sandy habitat (Mourier et al., 2012; Mourier and Planes 2021). The surveys consisted of 40 min dives (~30 min dedicated to survey) during which individual blacktip reef sharks were identified by photo-identification, using unique, lifelong color-shape of the dorsal fin (Mourier et al., 2012).

Associations between individuals were defined using the "Gambit of the Group", assuming that all individuals observed together are then considered as "associated." An experienced diver conducted a stationary visual census at each site monitored and identifying sharks within a ~100-m radius area (made possible by the high visibility conditions in these tropical waters). As most sharks usually remained together during the time of the dive, we considered the largest number of individuals observed within a 10-min period to be part of a group. We are confident that observed associations represented true grouping structure because groups were spatio-temporally well-defined and sharks were engaged in specific social behavior (e.g., following, parallel swimming or milling; Mourier et al., 2012). To avoid the potential for weak and nonrelevant associations between pairs of individuals with very low number of sightings, we used a restrictive observation threshold to include only individuals observed more than the median number of sightings (median = 14). Thus, all individuals seen less than 15 times were removed from the analyses to ensure that associations were estimated with high accuracy and precision.

Our dataset totaled 180 groups among 105 adult blacktip reef sharks.

References

Mourier, J. & Planes, S. Kinship does not predict the structure of a shark social network. Behavioral Ecology 32, 211-222 (2021).

Mourier, J., Vercelloni, J. & Planes, S. Evidence of social communities in a spatially structured network of a free-ranging shark species. Animal Behaviour 83, 389-401 (2012).

Moray and Tay dolphins

Data are from a long-term individual-based study of bottlenose dolphins on the east coast of Scotland[1]. Boat-based photo-identification (photo-ID) data have been collected since 1989, initially focussing on an area that was subsequently designated as the Moray Firth Special Area of Conservation (92/43/EEC), with occasional surveys further afield[2,3]. During this time the population expanded its distributional range[4,5] and since 2009 photo-ID surveys have also been regularly carried out around Tayside and adjacent waters[5,6]. Photo-ID surveys take place annually from May to September, with occasional surveys at other times of the year. Survey routes are chosen to maximise sighting probability while providing wide coverage of the study areas and all surveys were carried out under NatureScot Animal Scientific Licences. Surveys adhere to standardised protocols[2,5] where photographs were taken of the dorsal fins of as many individual dolphins as possible, graded for quality and using unique markings matched to the Universities of Aberdeen and St Andrews catalogue of known bottlenose dolphins on the Scottish east coast. All dolphins within 100m and engaged in similar activities or travelling in the same direction were considered to be in the same group and associated. Sex was determined using genital photographs or if adults were seen in repeat association with a known calf[7]. Year of birth was estimated from photographs of calves (up to 2 years old) based on their foetal folds, size, colour and behaviour[8].

Sightings data from this population were provided for this study for every bottlenose dolphin group encountered from 1990 to 2021 including the date, location, estimated group size, unique identifier of each dolphin identified, and their sex and year of birth if known. This dataset comprised sightings of 903 individual bottlenose dolphins (males, females, unknown sex and all ages) in 3071 groups and was split into two study areas, Moray (730 dolphins in 2525 groups) and Tay (339 dolphins in 546 groups). Individuals from this population are known to travel across the east coast of Scotland and 166 dolphins were photographed in both areas.

References

1   Cheney, B. J., Arso Civil, M., Hammond, P. S. & Thompson, P. M. Site Condition Monitoring of bottlenose dolphins within the Moray Firth Special Area of Conservation: 2017-2022., (2024).

2   Cheney, B. et al. Long-term trends in the use of a protected area by small cetaceans in relation to changes in population status. Global Ecology and Conservation 2, 118-128, doi:https://doi.org/10.1016/j.gecco.2014.08.010 (2014).

3   Wilson, B., Hammond, P. S. & Thompson, P. M. Estimating size and assessing trends in a coastal bottlenose dolphin population. Ecol Appl 9, 288-300 (1999).

4   Wilson, B., Reid, R. J., Grellier, K., Thompson, P. M. & Hammond, P. S. Considering the temporal when managing the spatial: a population range expansion impacts protected areas-based management for bottlenose dolphins. Anim. Conserv. 7, 331-338, doi:10.1017/s1367943004001581 (2004).

2027   5   Arso Civil, M. et al. Changing distribution of the east coast of Scotland bottlenose
2028   dolphin population and the challenges of area-based management. Aquatic Conservation:
2029   Marine and Freshwater Ecosystems 29, 178-196, doi:10.1002/aqc.3102 (2019).

2030   6   Arso Civil, M. et al. Variations in age- and sex-specific survival rates help explain
2031   population trend in a discrete marine mammal population. Ecology and Evolution 9, 533-544,
2032   doi:doi:10.1002/ece3.4772 (2019).

2033   7   Cheney, B. J., Dale, J., Thompson, P. M. & Quick, N. J. Spy in the sky: a method to
2034   identify pregnant small cetaceans. Remote Sensing in Ecology and Conservation n/a,
2035   doi:https://doi.org/10.1002/rse2.258 (2022).

2036   8   Grellier, K., Hammond, P. S., Wilson, B., Sanders-Reed, C. A. & Thompson, P. M.
2037   Use of photo-identification data to quantify mother-calf association patterns in bottlenose
2038   dolphins. Can. J. Zool.-Rev. Can. Zool. 81, 1421-1427, doi:10.1139/z03-132 (2003).

2039

2040

Mountain Lake mice

Data for the Mountain Lake Mice were collected as part of two 3-year deworming experiments conducted at the Mountain Lake Biological Station, Pembrooke, VA from 2016-2018 (37°22'N, 80°31W). Eight 0.5 ha grids (8 x 8 traps with 10 m spacing) were live-trapped for 2-3 night sessions approximately every 2 weeks using Sherman live traps (H.B. Sherman 2x2.5x6.5 inch folding trap, Tallahassee, FL, USA) from May to August/early September each year. Trapping protocols followed (Sweeny et al. 2020) with each individual receiving a numbered ear tag upon first capture. Movement among grids was minimal with less than 2% of individuals captured on multiple grids. Thus, grids were treated as separate populations for local density and social network calculations. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night. Animal care and use protocols were approved by Princeton University, and the research was supported by a DARPA grant to ALG (68255-LS-DRP).

References

Sweeny, Amy R., Courtney A. Thomason, Edwin A. Carbajal, Christina B. Hansen, Andrea L. Graham, and Amy B. Pedersen. 2020. "Experimental Parasite Community Perturbation Reveals Associations between Sin Nombre Virus and Gastrointestinal Nematodes in a Rodent Reservoir Host." Biology Letters 16 (12): 20200604.

Multimammate mice

Mastomys natalensis, the multimammate mouse, occurs throughout sub-saharan Africa and is an important agricultural pest species as well as host to a number of diseases of human health importance (e.g. plague, Lassa virus and leptospira (Fichet-calvet and Becker-ziaja, 2014; Holt et al., 2006; Mccauley et al., 2015). In Morogoro, Tanzania, where field data for this study was collected, *M. natalensis* undergoes considerable population fluctuations, with densities ranging from 10ha-1 during the breeding season up to 150ha-1 outside the breeding season. Population dynamics have been monitored using a robust design open capture mark recapture on a monthly basis since 1994 and is ongoing (Leirs et al., 2023). For the purposes of this study, data was used for the period of 1994 - 2022. The study design is as follows: 300 single capture Sherman traps were placed on a 300 x 100 m grid and baited with a mixture of peanut butter and cornflour. Trapping occurred for three successive nights; trapped individuals were weighed, sex and sexual condition recorded and individually marked with a unique toe clipping code if it is the first time they were captured. More details about the study site and capture methods can be found in Leirs et. al 2023. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night.

References

Fichet-calvet, E., Becker-ziaja, B., 2014. Lassa Serology in Natural Populations of Rodents 14. https://doi.org/10.1089/vbz.2013.1484

Holt, J., Davis, S., Leirs, H., 2006. A model of Leptospirosis infection in an African rodent to determine risk to humans: Seasonal fluctuations and the impact of rodent control. Acta Trop. 99, 218–225. https://doi.org/10.1016/j.actatropica.2006.08.003

Leirs, H., Kirkpatrick, L., Sluydts, V., Sabuni, C., Borremans, B., Katakweba, A., Massawe, A., Makundi, R., Mulungu, L., Machang'u, R., Mariën, J., 2023. Twenty-nine years of continuous monthly capture-mark-recapture data of multimammate mice (Mastomys natalensis) in Morogoro, Tanzania. Sci. Data 10, 798. https://doi.org/10.1038/s41597-023-02700-3

Mccauley, D.J., Salkeld, D.J., Young, H.S., Makundi, R., Dirzo, R., Eckerlin, R.P., Lambin, E.F., Gaffikin, L., Barry, M., Helgen, K.M., 2015. Effects of Land Use on Plague (Yersinia pestis) Activity in Rodents in Tanzania. Am. J. Trop. Med. Hyg. 92, 776–783. https://doi.org/10.4269/ajtmh.14-0504

Palmyra sharks

We tagged 41 grey reef sharks (*Carcharhinus amblyrhynchos*) that were caught on the forereefs of Palmyra Atoll (5°540 N 162°050 W) in the Central Pacific Ocean. Palmyra is a US Wildlife Refuge and includes large numbers of sharks and other predators (e.g. Bradley et al. 2017). Sharks were caught on hook and line and surgically implanted with Vemco V16 acoustic transmitters (69kHz, semi-randomized delay 60-180 seconds). Transmitters had battery lives of four years and were detected by a network of 55 acoustic receivers (Vemco, VR2W) positioned around the forereef and within the lagoons and back reefs (see Papastamatiou et al. 2018 for full details). Receivers would record the date and time of sharks that swam within range of the receiver (approximately 200-300 m).

We used a gambit of the group approach to generate dynamic social networks, where clusters of sharks co-occurring at receivers through time were identified using Gaussian mixture models (Psorakis et al. 2012, Jacoby et al. 2016). These clusters consisted of individual sharks that visited the same receivers at the same time. Adjacency matrices were produced based on how often dyadic pairs co-occurred within identified clusters. Full model details can be found in Papastamatiou et al. 2020.

References

Bradley D, Conklin E, Papastamatiou YP, McCauley DJ, Pollock K, Pollock A, Kendall BE, Gaines SD, Caselle JE. 2017. Resetting predator baselines in coral reef ecosystems. Sci. Rep. 7:43131

Jacoby DMP, Papastamatiou YP, Freeman R. 2016. Inferring animal social networks and leadership: applications for passive monitoring arrays. J. R. Soc. Interface 13:20160676

Papastamatiou YP, Bodey TW, Friedlander AM, Lowe CG, Bradley D, Weng K, Priestley V, Caselle JE. 2018. Spatial separation without territoriality in shark communities. Oikos. 127:767-779

Papastamatiou YP, Bodey TW, Caselle JE, Bradley D, Freeman R, Friedlander AM, Jacoby DMP. 2020. Multiyear social stability and social information use in reef sharks with diel fission-fusion dynamics. Proc. R. Soc. B. 287: 20201063

Psorakis I, Roberts SJ, Rezek I, Sheldon BC. 2012. Inferring social network structure in ecological systems from spatio-temporal data streams. J. R. Soc. Interface 9: 3055-3066

Potomac dolphins

Field Site and Dolphin Population

The Potomac-Chesapeake Dolphin Project has been conducting annual surveys of wild Tamanend's bottlenose dolphins (*Tursiops erebennus*) in the Potomac River and mid-Chesapeake Bay since July 2015. Approximately 550 km2 in the lower Potomac and mid-Chesapeake Bay are surveyed annually with 90% of survey effort concentrated in a 58 km2 area. Since the project started, over 2000 individuals have been identified with 36% sighted more than one day and 20% returning to the area more than one year. Depending on one's definition, the populations using the area are migratory and inhabit Chesapeake mainstem waters and tributaries in the warm months (April-October) before moving south during the cold months..

Data Collection

All data for this study were collected between 0700 and 1800 between April to October. For the current analyses, survey data annually from 2015 to 2018 were included for a total of 102 surveys with ID data on 1151 individuals (410 were re-sighted). Group sizes ranged from 1-163. The predominant group activity was determined the same way as in Shark Bay. However, group definitions differ slightly due to differences between the two systems. A survey group is defined as a close association or as an aggregation depending on the circumstances. A close association is a discrete group usually comprised of a small number of animals where the 10m chain rule can be used to determine association (as is the case for Shark Bay). An aggregation is a large number of animals that are within a 100m radius and may or may not be delineated into discrete groups as defined by the 10m chain rule. In aggregations, animals are clearly connected to each other with members moving between discrete groups or, in the absence of discrete groups, among each other. Under either circumstance (with or without discrete groups), animals move among each other in such a way that at some point during the survey they have likely been connected by the 10m chain rule.

References

Karniski, C., Patterson, E. M., Krzyszczyk, E., Foroughirad, V., Stanton, M., and Mann, J. 2015. Determining cetacean activity budgets: a comparison of surveys and individual focal follows. Marine Mammal Science. 31:839-852. DOI: 10.1111/mms.12198

Mann, J. 1999. Behavioral Sampling Methods for Cetaceans: A review and critique. Marine Mammal Science, 15(1): 102-122.

McEntee, M.H., Foroughirad, V., Krzyszczyk, E. and Mann, J., 2023. Sex bias in mortality risk changes over the lifespan of bottlenose dolphins. Proceedings of the Royal Society B, 290(2003), p.20230675.

2167 Tsai, Y.J. & Mann, J. 2013. Dispersal, philopatry and the role of fission-fusion dynamics in
2168 bottlenose dolphins. Marine Mammal Science. 29(2) April: 261-279 DOI: 10.1111/j.1748-
2169 7692.2011.00559.x

2170 Prairie dogs

We studied black-tailed prairied dogs on one colony in Grasslands National Park,
Saskatchewan, Canada (49.06N, 107.36W) as a long-term project from 2015 to 2020[1]. Our
goal was to capture all individuals to get a full colony census of the age and sex structure
within the colony as well as population dynamics. We used Tomahawk traps (Tomahawk,
Hazelhurst WI) to capture each individual, and subsequently tagged each pinnae with
alphanumeric ear tags (Monel #1, National Band and Tag Co., Newport, Kentucky) and
painted their dorsal pelage with a unique symbol for future identification. We weighed,
sexed, and aged each prairie dog at first capture.

We conducted our behavioural observations during April (pre-juvenile) and June (post-
juvenile emergence) 2017 within a 3-hectare area located centrally within the colony as all
animals in this area were marked and readily identifiable from a distance with binoculars. We
marked the 3 hectares with a 15 m Cartesian grid system to record the location of individuals
during focal observations. As prairie dogs are a sedentary species, space use and social
structure are not independent of each other and we recorded both social and movement
behaviours. 57 individuals aged 1 year and older were observed across 9 coteries (mean =
6.61 individuals per coterie, range = 2 to 14). We recorded behaviours as we sat > 40 m
from marked individuals during peak activity hours for prairie dogs based on season. We
recorded all affiliative (sniffing, jump-yipping[2], greet-kissing, mutual vigilance, shared
foraging) and agonistic (fighting, chasing, standoffs, territorial defense) interactions between
all individuals in 3-4 hour sessions. We could record all social encounters across all animals
as encounter rates were relatively low. We also conducted focal scans to record all activities
and locations (<1 m) of each individual to establish their home range. We conducted four 20-
min focal scans per individual: two in April and two in June. We recorded an average of 56
locations (range 41-76) for each home range per season. We recorded 380 person-hours of
behavioural observation and 872 interactions between known individuals. We created two
networks from weighted undirected matrices of behavioural data and home ranges from the
focal scan movement data.

2198

2199 References

2200 1.   Kusch, J. M. & Lane, J. E. Determinants of social structure in a northern population of
2201 black-tailed prairie dogs, Cynomys ludovicianus. Animal Behaviour 178, 1–10 (2021).

2202 2.   Hare, J. F., Campbell, K. L. & Senkiw, R. W. Catch the wave: prairie dogs assess
2203 neighbours' awareness using contagious displays. Proceedings of the Royal Society B:
2204 Biological Sciences 281, 20132153 (2014).

2205

2206

Reticulated giraffes

This study was conducted in Ol Pejeta Conservancy (OPC), a 364 km2 semi-arid wildlife reserve located on the equator (0º N, 36º56' E) approximately 220 km north of Nairobi, Kenya. All giraffe at OPC were recognized using individually unique spot patterns along their necks, and classified to age groups as described in (1). At the time of this study, OPC had a population of 212 reticulated giraffe. At the conclusion of the study period, OPC's giraffe population consisted of 160 adults, 20 subadults, 21 juveniles, and 11 neonates. The population exhibited a 50:50 sex ratio.

From January 21 to August 2, 2011, giraffe group composition and membership were recorded for all giraffe groups sighted while driving pre-determined survey routes. Observed giraffe groups were followed off-road until a complete census of the individuals present was accomplished. A group was defined as a set of individuals engaged in the same behavior, or moving in the same direction or toward a common destination, as long as each giraffe was no more than 500 m from at least one other group member. During the study period, a total of 1089 observations of giraffe groups were collected. On average, 30.7 giraffe were observed per day, distributed between four to six groups. Each individual giraffe was observed on average 31.1 times (approximately once per week). A social network was constructed from observed association patterns, using the "gambit of the group" approach.

References

1.   VanderWaal KL, Wang H, McCowan B, Fushing H, Isbell LA. 2014. Multilevel social organization and space use in reticulated giraffe (Giraffa camelopardalis). Behav Ecol 25:17-26.

Rum deer

The study was carried out on an unpredated long-term study population of red deer on the Isle of Rum, Scotland (57°N,6°20′W). The natural history of this matrilineal mammalian system has been studied extensively (Clutton-Brock et al. 1982), and we focussed on females aged 3+ years, as these individuals have the most complete associated census data, and few males live in the study area except during the mating period. Individuals are monitored from birth, providing substantial life history and behavioural data, and >90% of calves are caught and tagged, with tissue samples taken[1].

Census data were collected for the years 1974-2017, totalling 423,070 census observations. Deer were censused by field workers five times a month, for eight months of the year, along one of two alternating routes[1]. Individuals' identities, locations (to the nearest 100M), and group membership were recorded. Grouping events were estimated by seasoned field workers according to a variant of the "chain rule" (Castles et al. 2014), where individuals grazing in a contiguous group within close proximity of each other (each individual under ~10 metres of at least one other individual in the group) were deemed to be associating, with mean 130.4 groups observed per individual across their lifetime (range 6-943). The mortality period falls between Jan-March, when there is least available food, and minimal mortality occurs outside this period. We only used census records in each May-December period, from which we derived annual social network position measures as response variables. We elected to investigate this seasonal period because it stretches from the spring calving period until the beginning of the mortality period, simplifying network construction and avoiding complications arising from mortality events. Our dataset totalled 3356 annual observations among 532 grown females (Figure 1).

We constructed a series of 43 annual social networks using "gambit of the group," where individuals in the same grouping event (as described above) were taken to be associating[3], and using dominance interactions.

References

1.	Clutton-Brock, T. H., Guinness, F. E. & Albon, S. D. Red Deer: Behavior and Ecology of Two Sexes. vol. 15 (University of Chicago Press, Chicago, IL, 1982).

2.	Castles, M. et al. Social networks created with different techniques are not comparable. Animal Behaviour 96, 59–67 (2014).

3.	Franks, D. W., Ruxton, G. D. & James, R. Sampling animal association networks with the gambit of the group. Behavioral Ecology and Sociobiology 64, 493–503 (2010).

Serengeti lions

The research data were collected on the well-studied population of African lions inhabiting a 2700km2 study area within the Serengeti National Park, Tanzania. Multiple rivers and tributaries transect this area, which encompasses both a grassland plains habitat in the southeast, and woodland habitat to the north. There are two main seasons, the wet season spanning November to May, and the dry season spanning June to October. While the woodlands habitat maintains relative stability year-round, the grasslands habitat experiences greater fluctuations in rainfall (Mosser and Packer, 2009). This leads to large seasonal migrations of prey species across the plains, with lower populations densities in the dry season. As such, lion territories shift with the seasons across the landscape, tracking prey species migrations (Packer et al., 2005).

Lionesses exist in egalitarian fission-fusion "prides" composed of related adults and their offspring. In the Serengeti study area these prides can range in size from 2-20 individuals. Within these prides, individuals spend large proportions of time in smaller subgroups, and frequently spend time alone. The composition and size of these subgroups is highly dynamic, fluctuating daily, but female pride-mates usually stay within 5-6 km of each other (within vocal communication range). At maturity (2-3 years of age), approximately 75% of female offspring are recruited into their natal pride, while 25% disperse to form new prides, usually in adjacent territories (Packer, 2023; Pusey and Packer, 1987).

At maturity, all male offspring disperse from their mother's pride in related groups of brothers and cousins. During this nomadic life stage multiple kin cohorts may form alliances, resulting in "coalitions" of up to nine males. Once they reach full body size (~ age 4 years old), these coalitions "take over" residence of a pride of females by outcompeting the existing resident male coalition (Packer, 2023; Pusey and Packer, 1987). Male coalition membership does not change once they are resident within a pride. No evidence of a dominance hierarchy has been recorded between male coalition mates (Packer et al., 2001; Packer, 2023).

The data for this study were collected over a 30-years period from 1984 to 2013. One female per pride was fitted with a GPS collar. Each collared lion was located at least once every two weeks, and individuals within 200m of each other were taken as co-occurring. These co-occurrences were recorded as part of one unique sighting event, with GPS coordinates, date and time stamps. Additionally, opportunistic sightings of individuals within the study area were recorded in the same manner. Identification of individuals with a high level of accuracy was possible using facial markings and whisker spots. For each individual lion within a sighting event, age (estimated from the first sighting as a cub or adult), sex, natal pride (if known), and current pride information was also recorded. We constructed social networks based on whether individuals were observed in the same group on the same date.

References

Mosser, A., and Packer, C. (2009). Group territoriality an dthe benefits of sociality in the African lion, Panthera leo. Animal Behaviour 78, 359-370.

Packer, C., Pusey, A.E., and Eberly, L.E. (2001). Egalitarianism in female African lions.
Science 293, 690–693. https://doi.org/10.1126/science.1062320.

Packer, C., Hilborn, R., Mosser, A., Kissui, B., Borner, M., Hopcraft, G., Wilmshurst, J.,
Mduma, S., and Sinclair, A.R.E. (2005). Ecological Change, Group Territoriality, and
Population Dynamics in Serengeti Lions. Most, 390–393.

Packer, C. (2023). The Lion: Behvaiour, ecology and conservation of an iconic species
(Princeton University Press).

Pusey, A.E., and Packer, C. (1987). The Evolution of Sex-Biased Dispersal in Lions.
Behaviour 101, 275–310.

2319

Shark Bay Dolphin Research Project

Field Site and Dolphin Population: The Shark Bay Dolphin Research Project has been conducting a longitudinal study of bottlenose dolphins (*Tursiops aduncus*) since 1984, monitoring over 1,900 individuals in the eastern gulf of Shark Bay, Australia. The relatively pristine study area extends 600 km2 east of Peron Peninsula and consists of embayment plains (5–13 m), shallow sand flats and seagrass beds (0.5–5 m), and deeper channels (6–13 m). Using boat-based sampling, field observers identify individual dolphins using dorsal fin markings and shape, and also matched photographs to a digital identification catalog. Dolphin ages were estimated based on year of birth, the birth of their first calf (Mann et al. 2000; McEntee et al. 2023), or degree of ventral speckling (Krzyszczyk and Mann 2012). Individuals were considered to be adults if they were 12 or more years of age, or for females, if they had given birth to a calf (Mann et al. 2000). Finally, sex was determined by visual observation of the genital area, genetic analysis, or the presence of a dependent calf. Both sexes remain in their natal area (bisexual philopatry) in this residential population.

Data Collection: All data for this study were collected between 0600 and 1900 during all seasons using survey methods.

A survey is a "snapshot" of a group or individual. When dolphins were sighted, researchers instantaneously estimated initial activity and distance from the research vessel before approaching to within 100 m. Once the research vessel was close enough for observers to identify individuals a survey was initiated on all individuals in the group based on a 10 m "chain rule" (Smolker et al. 1992). For each survey, observers performed a scan of all individuals to assess their behavioral state as one of six categories: foraging, resting, socializing, traveling, other, and unknown. A "predominant group activity" was assigned for the first five minutes of the survey based on the activities of at least 50% of the individuals (Mann 1999) for at least 50% of the time. The final dataset included 7293 group sightings of 910 individuals between 2008 and 2019.

References

Karniski, C., Patterson, E. M., Krzyszczyk, E., Foroughirad, V., Stanton, M., and Mann, J. 2015. Determining cetacean activity budgets: a comparison of surveys and individual focal follows. Marine Mammal Science. 31:839-852. DOI: 10.1111/mms.12198

Mann, J. 1999. Behavioral Sampling Methods for Cetaceans: A review and critique. Marine Mammal Science, 15(1): 102-122.

McEntee, M.H., Foroughirad, V., Krzyszczyk, E. and Mann, J., 2023. Sex bias in mortality risk changes over the lifespan of bottlenose dolphins. Proceedings of the Royal Society B, 290(2003), p.20230675.

Tsai, Y.J. & Mann, J. 2013. Dispersal, philopatry and the role of fission-fusion dynamics in bottlenose dolphins. Marine Mammal Science. 29(2) April: 261-279 DOI: 10.1111/j.1748-7692.2011.00559.x

Silwood wood mice

The field data collection was led by Sarah Knowles, and the data was cleaned and processed by Bryony Allen, Sarah Knowles, and Aura Raulo.

Data were collected over a one-year period (Nov 2014–Dec 2015) from a wild population of rodents in a 2.47 ha mixed woodland plot (Nash's Copse) at Imperial College's Silwood Park campus, UK. Data for this study was collected as part of a longer-term rodent capture-mark-recapture study, where several rodent species were caught (*Apodemus sylvaticus*, *Apodemus sylvaticus*, *Apodemus flavicollis* and *Myodes glareolus*). Trapping was performed every 2-4 weeks, using 122 small folding Sherman traps (5.1 x 6.4 x 16.5cm, H. B Sherman). Traps baited with eight peanuts, a slice of apple and sterile cotton wool for bedding were set at dusk and collected at dawn, with all animals processed, sampled and then released inside the 100m2 grid cell they were captured in. As part of processing, captured individuals were identified to species, sexed, weighed, and aged (juvenile or adult) based on size and pelage characteristics. At first capture, all individuals were injected subcutaneously with a passive integrated transponder tag (PIT-tag) for permanent identification. Ear punches were collected from all mice and stored in ethanol at -20°C to provide genetic material for host genotyping. Live traps were set in an alternating checkerboard design, to ensure even coverage. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night.

Sleepy lizards

The following description is adapted from Payne et al. (2022).

We studied sleepy lizards (*Tiliqua rugosa*), a species of skink native to southern Australia. Sleepy lizards are large (adults are 400-950 g, with snout-vent length 25-35 cm), mainly herbivorous, and can live up to 50 years (Bull, 1995; Bull et al., 2017). Social network studies have shown that sleepy lizards are pair-living with strong male-female pair bonds (Leu et al., 2010, 2011), exhibiting a long-term socially monogamous mating system (Bull et al., 1998; Leu et al., 2015). Sleepy lizards are primarily active during the austral spring, September to December (Bull, 1987). Overnight and during periods of heat stress, they shelter in shaded refugia (i.e., large shrubs, logs, or burrows) (Kerr et al., 2003).

Our field site was an ~ 1.2 km2 area near Bundey Bore Station (33.888240° S, 139.310718° E), South Australia (about 150 km north of Adelaide). The region has a semi-arid Mediterranean climate. The local site is dominated by chenopod shrubs (e.g., Maireana and Atriplex spp.) and patches of black oak trees (Casuarina spp.), with various annual plants growing between and under these shrubs and trees (e.g., the nonnative Ward's weed, Carrichtera annua, a preferred food item for sleepy lizards). In most years, food is more abundant in early spring when conditions are relatively cool and wet, becoming much less abundant later when conditions are hotter and drier.

As part of a long-term monitoring study, we used GPS units (horizontal precision +/- 6 m (Leu et al. 2010a)) to record adult lizard movement during their active season (i.e., the austral spring) from 2008 through 2017, excluding 2012. In 2008 through 2014, GPS units (Leu et al., 2010) took one GPS fix per 10 minutes, while in 2015 through 2017, GPS units took one fix per two minutes. To reduce autocorrelation in the GPS data and ensure consistency across years, following (Michelangeli et al., 2021), we thinned the GPS data from all years to one point per 20 minutes. We removed GPS errors according to fix accuracy (e.g., horizontal dilution greater than three), using an algorithm modified from Bjorneraas (2010) – which identifies errors via displacement, speed, and turning angle – and with manual inspection of GPS tracks for obvious errors.

References

Bjorneraas, K., Van Moorter, B., Rolandsen, C. M., & Herfindal, I. (2010). Screening Global Positioning System Location Data for Errors Using Animal Movement Characteristics. Journal of Wildlife Management, 74(6), 1361–1366. https://doi.org/10.2193/2009-405

Bull, C. M. (1987). A Population Study of the Viviparous Australian Lizard, Trachydosaurus rugosus (Scincidae). Copeia, 1987(3), 749–757.

Bull, C. M. (1995). Population Ecology of the Sleepy Lizard, Tiliqua rugosa, at Mt-Mary, South-Australia. Australian Journal of Ecology, 20(3), 393–402. https://doi.org/10.1111/j.1442-9993.1995.tb00555.x

Bull, C. M., Cooper, S. J. B., & Baghurst, B. C. (1998). Social monogamy and extra-pair
fertilization in an Australian lizard, Tiliqua rugosa. Behavioral Ecology and Sociobiology,
44(1), 63–72. https://doi.org/10.1007/s002650050515

Bull, C. M., Gardner, M. G., Sih, A., Spiegel, O., Godfrey, S. S., & Leu, S. T. (2017). Why Is
Social Behavior Rare in Reptiles? Lessons From Sleepy Lizards. In M. Naguib, J. Podos, L.
W. Simmons, L. Barrett, S. D. Healy, & M. Zuk (Eds.), Advances in the Study of Behavior
(Vol. 49, pp. 1–26). Academic Press. https://doi.org/10.1016/bs.asb.2017.02.001

Kerr, G. D., Bull, C. M., & Burzacott, D. (2003). Refuge sites used by the scincid lizard
Tiliqua rugosa. Austral Ecology, 28(2), 152–160. https://doi.org/DOI 10.1046/j.1442-
9993.2003.01268.x

Leu, S. T., Bashford, J., Kappeler, P. M., & Bull, C. M. (2010). Association networks reveal
social organization in the sleepy lizard. Animal Behaviour, 79(1), 217–225.
https://doi.org/10.1016/j.anbehav.2009.11.002

Leu, S. T., Burzacott, D., Whiting, M. J., & Bull, C. M. (2015). Mate Familiarity Affects Pairing
Behaviour in a Long-Term Monogamous Lizard: Evidence from Detailed Bio-Logging and a
31-Year Field Study. Ethology, 121(8), 760–768. https://doi.org/10.1111/eth.12390

Leu, S. T., Kappeler, P. M., & Bull, C. M. (2011). Pair-living in the Absence of Obligate
Biparental Care in a Lizard: Trading-off Sex and Food? Ethology, 117(9), 758–768.
https://doi.org/10.1111/j.1439-0310.2011.01934.x

Michelangeli, M., Payne, E., Spiegel, O., Sinn, D. L., Leu, S. T., Gardner, M. G., & Sih, A.
(2021). Personality, spatiotemporal ecological variation and resident/explorer movement
syndromes in the sleepy lizard. Journal of Animal Ecology, 91(1), 210–223.
https://doi.org/10.1111/1365-2656.13616

Payne, E., Spiegel, O., Sinn, D. L., Leu, S. T., Gardner, M. G., Godfrey, S. S., Wohlfeil, C., &
Sih, A. (2022). Intrinsic traits, social context, and local environment shape home range size
and fidelity of sleepy lizards. Ecological Monographs, 92(3), e1519.
https://doi.org/10.1002/ecm.1519

2445

Soay sheep

This study was carried out on a long-term study population of Soay sheep (*Ovis aries*) on the St Kilda archipelago. The Soay sheep have lived wild on the archipelago for millennia, and have been monitored since 1985 in our study area in Village Bay on Hirta. The natural history and population dynamics of these sheep have been extensively documented (Clutton-Brock & Pemberton, 2004). Sheep are captured, marked at birth, and followed longitudinally across their lifespan to collect morphological, behavioural, parasitological, and immunological data.

Census data were collected for the years 1988-2018. Sheep are censused 30 times per year (10 each in spring, summer, and autumn). Experienced fieldworkers follow established routes noting the identity, spatial location (to nearest 100m OS grid square), behaviour and group membership of individual sheep. Fieldworkers assign individual sheep to groups using a 'chain rule', where individuals in close proximity are classed as associating (Castles et al., 2014). In total, census data comprises 357,283 total observations of 81,769 groups. Individuals were associated with a mean of 50 unique groups across our total dataset (range 1-464). We used census records to construct annual social networks for each individual sheep, based on group membership in the same observation event.

References

Castles, M., Heinsohn, R., Marshall, H. H., Lee, A. E. G., Cowlishaw, G., & Carter, A. J. (2014). Social networks created with different techniques are not comparable. Animal Behaviour, 96, 59–67.

Clutton-Brock, T. H., & Pemberton, J. M. (2004). Soay Sheep: Dynamics and Selection in an Island Population. Cambridge University Press.

Vulturine guineafowl

The vulturine guineafowl (*Acryllium vulturinum*) is a terrestrial bird species (body size: 60-72 cm) that lives in the arid and semi-arid savanna of East Africa. Vulturine guineafowl are gregarious and typically live in stable groups of 15-65 individuals [1]. Groups are highly cohesive throughout and across the day. Previous work has suggested that groups are non-territorial, establishing highly overlapped home ranges among neighbouring groups [2,3]. Groups frequently roost communally [3] and when the ecological conditions become harsh (dry seasons and droughts), groups can expand their home range areas [2,4] and stable groups merge with preferred groups to form supergroups [1].

The vulturine guineafowl project was established in 2016 and collects data on up to 23 social groups (a total of 1189 birds) around the Mpala Research Centre in central Kenya. These data include fitting birds from each group with a global positioning system (GPS; 15g eObs solar) tag. These tags collect 1 data point per second (1 Hz), 10 consecutive data points every 5 minutes, or 1 data point every 15 minutes at the least, depending on the battery conditions (see [5] for more details). Here we used data from every 15th minute. In addition to the GPS tags, almost all individuals in our population have been fitted with a unique combination of colour bands for individual identification. Each day (morning and evening), a field team census groups in the area, recording the membership of each observed group and its location.

To obtain group size, we used the data from the group compositions for each month (see [6] for details) and combined the GPS data (where groups are) with group membership. In brief, we construct networks of robust co-observations among individuals [7], and then apply a community detection algorithm (walktrap community algorithm, using igraph package [8] in R) to identify groups. From these groups, we could quantify what GPS tags were in the same group and how many birds the group contained, thereby allowing us to estimate local densities.

The contact network was constructed based on the observation that meeting groups tend to mingle completely; each identified group-group contact event assumed that all individuals in one group were meeting all individuals in another group, with a threshold of 20 metres taken to represent a contact event.

References

1.   Papageorgiou, D. et al. The multilevel society of a small-brained bird. Current Biology 29, R1120–R1121 (2019).

2.   Papageorgiou, D., Rozen-rechels, D., Nyaguthii, B. & Farine, D. R. Seasonality impacts collective movements in a wild group-living bird. 1–12 (2021).

3.   Papageorgiou, D., Cherono, W., Gall, G., Nyaguthii, B. & Farine, D. R. Testing the information centre hypothesis in a multilevel society. Journal of Animal Ecology (2024) doi:10.1111/1365-2656.14131.

4.   Klarevas-Irby, J. A., Nyaguthii, B. & Farine, D. R. Moving as a group imposes
constraints on the energetic efficiency of movement. Proceedings B 292, (2025).

5.   He, P. et al. A guide to sampling design for GPS-based studies of animal societies.
Methods Ecol Evol 14, 1887–1905 (2023).

6.   Ogino, M., Strauss, E. D. & Farine, D. R. Challenges of mismatching timescales in
longitudinal studies of collective behaviour. Philosophical Transactions of the Royal Society
B 378, (2023).

7.   Shizuka, D. & Farine, D. R. Measuring the robustness of network community
structure using assortativity. Anim Behav 112, 237–246 (2016).

8.   Csardi, G. & Nepusz, T. The igraph software package for complex network research.
InterJournal Complex Systems Complex Sy, 1695 (2006).

2522

Water dragons

2523

Data were collected as part of an ongoing behavioural study of a population of eastern water
dragons (*Intellagama lesueurii*) inhabiting an urban city park, Roma Street Parklands,
Brisbane, Australia (27°27′46′ S, 153°1′11′ E). Consisting of approximately 336 individuals,
the population inhabits a highly heterogenous, man-made, curated public garden (Strickland
et al, 2014). Behavioural surveys were conducted twice daily (morning 7.30–10.30 and
afternoon 13.00–16.00) approximately three days a week from November 2010 onwards.
Data collection was restricted from September one year until April the following year,
representing the season in which dragons are active and not in brumation.

For each individual encountered, a photo of their left and/or right facial profile, along with
their GPS location was recorded. An individual's sex was assigned based on sexual
dimorphism and dichromatism (Thompson, 1993). The individual's immediate behaviour at
the time of the observation was recorded, which included a spectrum from resting to
agonism (e.g. head bobbing, tail slapping, arm waving, chasing and physical combat).

Profile photographs taken during surveys were used to identify individuals using a previously
established method for this population (Gardiner et al, 2014). This method employed
interactive identification software (I3S Spot, v. 4.0.2) which compared individual facial profile
scale patterns from images taken during behavioural surveys to an existing photo library. We
also took GPS locations of every observation; on each census, individuals with less than
1.85m between them were taken to be in contact.

2543

References

Gardiner, R.Z., Doran, E., Strickland, K., Carpenter-Bundhoo, L. and Frère, C. A face in the
crowd: a non-invasive and cost effective photo-identification methodology to understand the
fine scale movement of eastern water dragons. PloS one 9, 96992 (2014).

Strickland, K., Gardiner, R., Schultz, A.J. and Frere, C.H., 2014. The social life of eastern
water dragons: sex differences, spatial overlap and genetic relatedness. Animal Behaviour
97, 53-61 (2014).

Thompson, M.B. Estimate of the population structure of the eastern water dragon,
Physignathus lesueurii (Reptilia: Agamidae), along riverside habitat. Wildlife Research 20,
613-619 (1993).

2554

2555

Wild crickets

The aim of the project was to monitor a wild population of field crickets (*Gryllus campestris*) living in a meadow in North Spain. This species is annual, and every individual spends most of its life inside or in the vicinity of burrows excavated in the ground. We monitored the natural population for 12 consecutive years. Every year, we found all the burrows in the meadow and marked them with a flag having a unique number. We trapped every individual in the population shortly after they emerged as adults and marked them with a plastic tag with a unique code for each individual. We used up to 140 infra-red video cameras to record 24hrs a day at burrows and an area of about 20cm in diameter around the burrow from the date of first adult emergence to the date when the last adult in the population died. We then watched the video and extracted a number of relevant behaviours for each recorded individual, which were then used to form the social networks. A more detailed description of our methodology can be found in Rodríguez-Muñoz et al. (2019).

References

Rodríguez-Muñoz, R., Boonekamp, J. J., Liu, X. P., Skicko, I., Fisher, D. N., Hopwood, P., & Tregenza, T. (2019). Testing the effect of early-life reproductive effort on age-related decline in a wild insect. Evolution, 73, 317-328. doi: 10.1111/evo.13679

Woodchester badgers

The Woodchester Park badger study is conducted at Woodchester Park, Gloucestershire, UK (51.7°N,2.25°W), and has been ongoing since 1976 [1,2]. For the purposes of this study co-trapping data were used to estimate both population density and social networks (note that co-trapping will only provide a proxy for social associations in this system;[3]). The study site is divided into three zones of approximately equal size, each of which is trapped four times per year from May to January inclusive (no captures occur in the intervening period to avoid catching dependent cubs and their mothers). Box traps constructed of steel mesh are set close to each active main and outlying sett (detected by sett activity surveys in the build-up to trapping) and baited with peanuts for up to 8 days. Traps are then set for two consecutive nights and checked the following mornings. At each active sett, more traps are deployed than expected to be required. On first capture badgers are permanently marked with a unique ID tattoo on the abdomen allowing them to be identified on future captures [4]. Individuals are typically caught (on average) approximately two times per year[2].

Co-trapping networks can then be constructed based on individuals being caught at the same sett on the same day. Because these networks are constructed using co-trapping data only they can include all individuals regardless of their age or sex.

References

1.   McDonald, J. L., Robertson, A. & Silk, M. J. Wildlife disease ecology from the individual to the population: Insights from a long-term study of a naturally infected European badger population. Journal of Animal Ecology 87, 101–112 (2018).

2.   Delahay, R. J. et al. Long-term temporal trends and estimated transmission rates for Mycobacterium bovis infection in an undisturbed high-density badger (Meles meles) population. Epidemiol Infect 141, 1445–1456 (2013).

3.   Silk, M. J., McDonald, R. A., Delahay, R. J., Padfield, D. & Hodgson, D. J. CMRnet: An r package to derive networks of social interactions and movement from mark–recapture data. Methods Ecol Evol 12, 70–75 (2021).

4.   Cheeseman, C. L. & Harris, S. Methods of marking badgers (Meles meles). Journal of Zoology, Lond. 197, 289–292 (1982).

Wytham badgers

The Wytham Badger Project ran 1987 to 2019 and attempted to monitor all badgers resident in the Wytham Woods SSSI, a 4.24 km2 mixed woodland in southern England (51° 46′ N, 1° 20′ W; for further information, see Macdonald & Newman 2022). The population is situated on a hill surrounded by the River Thames on three sides and the A34 motorway on the fourth, mimimizing migration (immigration/emigration rate = 3%: Macdonald and Newman 2002). Over the study period, every active communal burrow system, termed a "sett", was trapped using string-trigger traps for two or (most commonly) three nights, three–four times per year, at regular seasonal intervals. Badgers were transferred to holding cages between 7:00 and 9:00 a.m., transported to a central feldstation, and sedated with 0.2 mL ketamine hydrochloride/kg body weight by intramuscular injection (McLaren et al. 2005). On first capture (typically as a cub or yearling), each badger received a unique numerical inguinal tattoo. The population divided into 23 social groups, established from frequent baitmarking surveys (Buesching et al. 2016), where each social group utilized several setts consisting of 1–10 holes, with on average 5.5 individuals cohabitating in the average sett (range 1 to 28).

In total, the study amassed 11,488 captures of 1823 individuals, to which an enhanced Minimum Number Alive (eMNA; Bright Ross et al. 2002) enumeration procedure was applied. Population size averaged 242 adults ± 15.14 SD, range = 222–263) plus 66 cubs (± 8.1 SD, range = 47–97) from 2005 to 2009. Thereafter, following high, and unexplained, mortality across age classes in 2010, the population settled to a slightly lower but stable phase through to 2015, comprising 195 adults (± 17.06 SD, range = 177–217) and 49 cubs (± 15.47 SD, range = 24–66; Bright Ross et al. 2020). In 2019, population density was 44.55 ± 5.37 badgers km-2, where Wytham Woods has consistently had the highest density of European badgers ever recorded (Macdonald and Newman 2022). Co-trapping networks were then constructed based on individuals being caught at the same sett on the same day.

References

Buesching, C. D., Newman, C., Service, K., Macdonald, D. W., & Riordan, P. (2016). Latrine marking patterns of badgers (Meles meles) with respect to population density and range size. Ecosphere, 7(5), e01328.

Bright Ross, J. G., Newman, C., Buesching, C. D., & Macdonald, D. W. (2022). Preserving identity in capture–mark–recapture studies: increasing the accuracy of minimum number alive (MNA) estimates by incorporating inter-census trapping efficiency variation. Mammalian Biology, 102(3), 567-580.

Macdonald, D. W., & Newman, C. (2002). Population dynamics of badgers (Meles meles) in Oxfordshire, UK: numbers, density and cohort life histories, and a possible role of climate change in population growth. Journal of Zoology, 256(1), 121-138.

Macdonald, D., & Newman, C. (2022). The Badgers of Wytham Woods: a model for
behaviour, ecology, and evolution. Oxford University Press.

McLaren, G. W., Thornton, P. D., Newman, C., Buesching, C. D., Baker, S. E., Mathews, F.,
& Macdonald, D. W. (2005). The use and assessment of ketamine–medetomidine–
butorphanol combinations for field anaesthesia in wild European badgers (Meles
meles). Veterinary anaesthesia and analgesia, 32(6), 367-372.

2653

Wytham tits

The study was conducted in Wytham Woods, Oxford, UK (51°46′ N, 1°20′ W), on a great tit (Parus major) population monitored using standardized protocols since the 1960s [1]. Great tits breed almost exclusively in nest boxes fixed at 1020 GPS-mapped locations across Wytham Woods [2]. Over 98% of breeding great tits occupy a single nest box per year, with breeding spanning April–July and comprising nest building, egg laying, incubation, and offspring rearing. Nestboxes were regularly checked to monitor breeding attempts, identify (or mark) adults (days 6–14 of the nestling phase), and mark nestlings (day 15) with unique British Trust for Ornithology (BTO) metal leg rings, alongside taking standard morphometric measurements [1-3].

In winter (September–March), great tits form highly dynamic, fission-fusion feeding flocks with frequent turnover [3-5]. Since 2007, great tits captured during breeding or winter mist-netting have been fitted with plastic leg rings containing a unique RFID microchip, resulting in the RFID tagging of ~90% of the population [4]. RFID tags enable tracking of individuals at sunflower feeding stations equipped with two RFID antennae (Dorset ID, Aalten, Netherlands) at placed in 65 stratified-grid locations through in winters starting 2011, 2012, and 2013. During these winter seasons, these feeders were open every weekend from End-Nov to End-February (13 weekends), continuously scanning for RFID-tagged birds from pre-dawn to post-dusk.

Winter Population Social Structure

RFID detections generated a spatiotemporal datastream reflecting the bursts of activity as flocks arrived and fed. A machine-learning algorithm (Gaussian mixture model) assigned detections to the most likely flock or 'gathering event', providing a robust and effective method for determining flock co-membership [5]. From the resulting group-by-individual matrices, social networks (association matrices) were constructed using the Simple Ratio Index [5,6].

For this analysis, non-directional weighted networks were built from the yearly (i.e. whole winter season GBI matrix) to measure dyadic association propensity, and the extensive sampling here reduces the typical limitations of the 'gambit of the group' approach [7]. Networks included all RFID-detected individuals linked to a unique BTO ring number. Due to high turnover and movement, kin structure was weak, with only ~1.5% of winter social connections occurring between first-order relatives [7]; therefore, relatedness was not considered a key structuring factor. Nevertheless, this social network dataset is known to be biologically meaningful, and forms part of a broader investigation into great tit social ecology, contributing insights into individual sociality [8,9], social structure and demography [7,10], and links to ecological processes such as information transmission [11,12], foraging [13, 14], breeding settlement and mating behavior [7,14,15].

All work was approved by the University of Oxford, Department of Zoology, Animal Welfare and Ethical Review Board (Approval: APA/1/5/ZOO/NASPA/Sheldon/TitBreedingEcology) and adhered to local animal research guidelines. All birds were caught, tagged, and ringed by licensed BTO ringers.

References

Perrins CM. 1965 Population fluctuations and clutch-size in the great tit, Parus major L. J
Anim Ecol 34, 601–647. (doi:10.2307/2405301)

Wilkin TA, Perrins CM, Sheldon BC. 2007 The use of GIS in estimating spatial variation in
habitat quality: a case study of lay-date in the Great Tit Parus major. Ibis 149, 110–118.
(doi:10.1111/j.1474-919X.2006.00627.x)

Farine DR, Firth JA, Aplin LM, Crates RA, Culina A, Garroway CJ, Hinde CA, Kidd LR,
Milligan ND, Psorakis I, et al. 2015 The role of social and ecological processes in structuring
animal populations: a case study from automated tracking of wild birds. R Soc Open Sci 2,
150057. (doi:10.1098/rsos.150057)

Aplin LM, Farine DR, Morand-Ferron J, Cole EF, Cockburn A, Sheldon BC. 2013 Individual
personalities predict social behaviour in wild networks of great tits (Parus major). Ecol Lett
16, 1365–1372. (doi:10.1111/ele.12181)

Psorakis I, Voelkl B, Garroway CJ, Radersma R, Aplin LM, Crates RA, Culina A, Farine DR,
Firth JA, Hinde CA, et al. 2015 Inferring social structure from temporal data. Behav Ecol
Sociobiol 69, 857–866. (doi:10.1007/s00265-015-1906-0)

Cairns SJ, Schwager SJ. 1987 A comparison of association indices. Anim Behav 35, 1454–
1469. (doi:10.1016/S0003-3472(87)80018-0)

Firth JA, Sheldon BC. 2016 Social carry-over effects underpin trans-seasonally linked
structure in a wild bird population. Ecol Lett 19, 1324–1332. (doi:10.1111/ele.12669)

Aplin LM, Firth JA, Farine DR, Voelkl B, Crates RA, Culina A, Garroway CJ, Hinde CA, Kidd
LR, Psorakis I, et al. 2015 Consistent individual differences in the social phenotypes of wild
great tits, Parus major. Anim Behav 108, 117–127. (doi:10.1016/j.anbehav.2015.07.016)

Firth JA, Voelkl B, Crates RA, Aplin LM, Biro D, Croft DP, Sheldon BC. 2017 Wild birds
respond to flockmate loss by increasing their social network associations to others. Proc. R.
Soc. B Biol. Sci. 284. (doi:10.1098/rspb.2017.0299)

Beck, K. B., Farine, D. R., Firth, J. A., & Sheldon, B. C. (2023). Variation in local population
size predicts social network structure in wild songbirds. Journal of Animal Ecology, 92(12),
2348–2362. https://doi.org/10.1111/1365-2656.14015

Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. 2012 Social networks predict patch
discovery in a wild population of songbirds. Proc Biol Sci 279, 4199–4205.
(doi:10.1098/rspb.2012.1591)

Firth JA, Sheldon BC, Farine DR. 2016 Pathways of information transmission among wild
songbirds follow experimentally imposed changes in social foraging structure. Biol Lett 12,
20160144. (doi:10.1098/rsbl.2016.0144)

McMahon K, Marples NM, Spurgin LG, Rowland HM, Sheldon BC, Firth JA. 2024 Social
network centrality predicts dietary decisions in a wild bird population. iScience 27, 109581.
(doi:10.1016/j.isci.2024.109581)

Firth JA, Voelkl B, Farine DR, Sheldon BC. 2015 Experimental evidence that social
relationships determine individual foraging behavior. Curr Biol 25, 3138–3143.
(doi:10.1016/j.cub.2015.09.075)

Firth JA, Cole EF, Ioannou CC, Quinn JL, Aplin LM, Culina A, McMahon K, Sheldon BC.
2018 Personality shapes pair bonding in a wild bird social system. Nat Ecol Evol, 1696–
1699. (doi:10.1038/s41559-018-0670-8)

2740

2741

Wytham wood mice

This data set is a trapping data set of wild rodents caught in Holly Hill, Wytham Woods 2018-2019. Rodents were trapped approximately fortnightly and new individuals tagged with a PIT-tag for permanent identification. Individuals were then released back to their point of capture. Data were collected over a 1-year period (Nov 2018–Nov 2019) from a wild population of rodents in a 4 ha (200 x 200 meters square) mixed woodland plot. Data for this study was collected as part of a longer-term rodent capture-mark-recapture study, where several rodent species were caught (*Apodemus sylvaticus*, *Apodemus flavicollis* and *Myodes glareolus*). Trapping was performed every 2-4 weeks, using 200 small folding Sherman traps (5.1 x 6.4 x 16.5cm, H. B Sherman). To ensure even trapping coverage, live traps were set with an alternating checkerboard design in every other 10 x 10 meter "Grid cell" of the 200 x 200 meters study area. Traps baited with 6 peanuts, a slice of apple and sterile cotton wool for bedding were set at dusk and collected at dawn, with all animals processed, sampled and then released inside the 100m2 grid cell they were captured in. As part of processing, captured individuals were identified to species, sexed, weighed, and aged (to juvenile or adult) based on size and pelage characteristics. At first capture, all individuals were injected subcutaneously with a passive integrated transponder tag (PIT-tag) for permanent identification. There were a few cases (~2% of captures) where the animal died during trapping and thus was not released. This was most often due to animal being found dead in the trap (which can happen due to epidemics or other poor health especially in the spring) and in a couple of cases due to animal being put down following Schedule 1 due to extreme poor health or injury. Edges in the social network between individuals (nodes) were defined as unique mice trapped nearby (within one adjacent trap distance by Euclidean distance) in the same trapping night.