Density-dependent network structuring within and across wild animal systems

Gregory F Albery*[1,2], Daniel J Becker[3], Josh A Firth[4,5], Matthew Silk[6,7], Amy R Sweeny[8,9], Eric Vander Wal[10], Quinn Webber[11], Bryony Allen[12,13], Simon A Babayan[14], Sahas Barve[15,16], Mike Begon[12], Richard J Birtles[17], Theadora A Block[18,19], Barbara A Block[20], Janette E Bradley[21], Sarah Budischak[22], Cfhristina Buesching[23,24], Sarah J Burthe[25], Aaron B Carlisle[26], Jennifer E Caselle[27], Ciro Cattuto[28], Alexis S Chaine[29], Taylor Chapple[30], Barbara J Cheney[31], Timothy Clutton-Brock[32], Melissa Collier[1], David J Curnick[33], Richard J Delahay[34], Damien R Farine[35,36,37,38], Andy Fenton[12], Francesco Ferretti[39], Helen Fielding[40,41], Vivienne Foroughirad[1,42], Celine Frere[43], Michael G Gardner[44], Eli Geffen[45], Stephanie S Godfrey[46], Andrea L Graham[47], Phil S Hammond[48], Maik Henrich[49], Marco Heurich[49,50,51], Paul Hopwood[7], Amiyaal Ilany[52], Joseph A Jackson[17], Nicola Jackson[43], David Jacoby[53], Ann-Marie Jacoby[54], Miloš Ježek[55], Lucinda Kirkpatrick[56], Alisa Klamm[57], James A Klarevas-Irby[35,36,58,59], Sarah Knowles[4], Lee Koren[52], Ewa Krzyszczyk[60], Jillian M Kusch[61,62], Xavier Lambin[63], Jeffrey E Lane[62], Herwig Leirs[56], Stephan T Leu[64], Bruce E Lyon[18], David W MacDonald[23], Anastasia E Madsen[65,66], Janet Mann[1], Marta Manser[67], Joachim Mariën[56], Apia Massawe[68], Robbie A McDonald[41], Kevin Morelle[55,69], Johann Mourier[70,71,72], Chris Newman[23], Kenneth Nussear[73], Brendah Nyaguthii[37,38,58,74], Mina Ogino[35,36,58], Laura Ozella[75], Yannis Papastamatiou[76], Steve Paterson[77], Eric Payne[78], Amy B Pedersen[9], Josephine M Pemberton[9], Noa Pinter-Wollman[79], Serge Planes[71,72], Aura Raulo[4], Rolando Rodríguez-Muñoz[7], Christopher Sabuni[68], Pratha Sah[1], Robbie J Schallert[80], Ben C Sheldon[4], Daizaburo Shizuka[81], Andrew Sih[78], David L Sinn[82], Vincent Sluydts[56], Orr Spiegel[45], Sandra Telfer[63], Courtney A Thomason[39,83], David M Tickler[84], Tom Tregenza[7], Kimberley VanderWaal[85], Eric L Walters[15], Klara M Wanelik[77,86], Elodie Wielgus[49], Jared Wilson-Aggarwal[7,41], Caroline Wohlfeil[44], Shweta Bansal[1]

*Corresponding author: gfalbery@gmail.com. Affiliations are at the end of the document.

Abstract

High population density should drive individuals to more frequently share space and interact, producing better-connected spatial and social networks. Despite this widely-held assumption, it remains unconfirmed how local density generally drives individuals' positions within wild animal networks. We analysed 34 datasets of simultaneous spatial and social behaviour in >55,000 individual animals, spanning 28 species of fish, reptiles, birds, mammals, and insects. >80% of systems exhibited strongly positive relationships between local density and network centrality, providing broad empirical evidence that local density increases connectedness at the individual level. However, >75% of density-connectedness relationships were nonlinear, and density's importance declined at higher values in >70% of systems, signifying saturating effects. Density's effect was much stronger and less saturating for spatial than social networks, suggesting population density drives individuals to become disproportionately spatially connected rather than socially. These findings reveal fundamental trends underlying societal structuring, with widespread behavioural, ecological, and evolutionary implications.

42   **Keywords**: Behavioural ecology, Spatial ecology, Disease ecology, Epidemiology,
43   Population dynamics, Social network structure, Network analysis, Spatial analysis

44   Introduction

The number of individuals occupying a given space – i.e., population density – is a central
factor governing social systems. At higher densities, individuals are expected to more
frequently share space, associate, and interact, producing more-connected spatial and
social networks and thereby influencing downstream processes such as mating, learning,
and competition. In particular, density-driven increases in network connectedness should
provide more opportunities for parasites to spread between hosts [1–5]. Despite the
fundamental nature of such density-dependent processes, evidence is relatively limited that
individuals inhabiting higher-density areas have more spatial and social connections.
Furthermore, density's effects should differ for asynchronous space sharing (e.g. home
range overlap) *versus* social associations (e.g. den sharing or grouping) or interactions (e.g.
mating or fighting). While several studies have compared animal populations at different
densities to demonstrate variation in social association rates among populations (e.g., [6,7])
or groups (e.g., [8–10]), attempts to identify such density effects *within* continuous
populations of individuals are rarer (but see [11–14]), and their findings have never been
synthesised or compared. We therefore have an incomplete understanding of how density,
as a fundamental ecological parameter, determines socio-spatial dynamics within and
across systems. This inhibits our ability to identify and predict how changes in density – e.g.
through culling, natural mortality, dispersal, or population booms – influence downstream
processes that depend on shared space and social interactions.

The rate at which an individual interacts with conspecifics depends on its spatial and social
behaviour within the context of the surrounding environment and population. Adding more
individuals into the same space should cause them to more frequently spatially overlap and
socially associate or interact (Figure 1). Often, individuals are modelled as randomly moving
and interacting molecules ("mass action" or "mean field"). In this conceptualisation, direct
contact between two molecules is analogous to a social interaction or association; rates of
such interactions are often assumed to increase with density ("density-dependent"; e.g.,
[15]), and/or to be homogenous in space (e.g., [10]). In reality individuals are unlikely to
behave and interact randomly in space, and instead will be influenced by spatially varying
factors including local density [16]. Changes in density may cause individuals to alter their
foraging behaviour [17–19], dispersal [20,21], social preference or avoidance [13,22], mating
behaviour [23], or preferred group size [7]; in some cases, interaction rates may change to
remain constant in the face of changing densities [24]. These and related processes might
produce strong nonlinearities in density-interaction relationships, which can complicate the
predictions of density dependence models of pathogen transmission, for example [2,4,5].
Nevertheless, these nonlinearities are poorly understood and rarely considered.

Several wild animal studies have suggested saturating nonlinear relationships between
density and social association rates [8–10,13]. Such relationships imply that association
rates do not increase passively with density, and that behavioural or demographic processes
likely change as density increases, with the ultimate consequence of slowing association
rates. However, these nonlinearities are difficult to examine between populations or between
species because they introduce a range of confounders and have few replicates along the

density axis [2]. Characterising gradients of density across individuals within a population offers a workaround to this problem, and facilitates an appreciation of the fact that interactions occur between individuals rather than at the population level. Examining between-individual variation is one reason that social network analysis – which allows characterisation and analysis of individual-level social traits, amongst other things – has become so popular in animal ecology in recent years [25–29]. Additionally, recent years have seen a substantial growth in understanding of socio-spatial behaviours, including harmonising the concepts of spatial and social density [2,16,30]. Applying network analyses coupled with this socio-spatial understanding of density could provide an individual-level picture of density's effects on spatial and social connectedness, offering far higher resolution and statistical power and greater ability to detect nonlinearities and between-system differences [2]. By providing new understanding of the correlates and emergent consequences of variation in density, this expansion could help to identify general rules underlying social structuring and network scaling in space.

Critically, different types of interactions or associations should show different relationships with density: for example, the need to compete for food at higher densities could drive a disproportionate increase in aggression [31], but this is unlikely to be true of mating interactions. This rationale is well-understood in disease ecology, as differences in density-contact relationships are thought to drive differences in density dependence of infection – where "contact" is defined as an interaction or association that could spread a pathogen (Figure 1). "Contacts" then form the basis of spatial and social networks used to investigate pathogen transmission dynamics, which should likewise diverge with density just as contacts do. For example, density should drive greater transmission of respiratory pathogens but not sexually transmitted pathogens [1,32]. Establishing these density-contact relationships is integral to understanding disease dynamics and developing control measures [1,33], but we still have a poor understanding of how different interactions (and therefore contact events for different pathogens) are driven by density. This direct/indirect interaction dichotomy is most fundamental to disease ecology [30], but given building interest in the spatial-social interface and relationships between spatial and social networks in behavioural ecology [16], the framework is readily related to other fields. Established density-interaction relationships are diverse and include feral dog bites [11], ant antennations [34] and trophallaxis [24], ungulate group memberships [12,17], rodent co-trapping [8,35], and agamid association patterns [13,14], but no study has yet synthesised how the rates of multiple interaction types relate to density, within or across systems.

Identifying the general rules underlying density dependence requires quantifying density's relationship with proxies of interaction rates at fine scales across a diversity of systems, then identifying the factors determining their slope and shape. To this end, we collate a meta-dataset of over 55,000 individual animals across 34 wildlife systems globally to ask how within-population variation in density determines between-individual interaction rates based on connectedness in spatial and social networks. We fit multiple linear and nonlinear relationships to identify the slope and shape of density effects within each system, and we use meta-analyses to investigate general rules determining their slope and shape across systems. In particular, we focus on comparing space sharing with social interactions and associations as a cross-system case study. Ultimately, we present a *de novo* cross-system analysis of individuals' social and spatial behaviour that traverses fields of behavioural,

population, and disease ecology, which could help to inform general rules governing the
structure of social systems.

Figure 1: Schematic detailing the rationale underlying this study, outlining how population density
drives the formation of spatial and social networks. This depiction uses the Wytham Wood great tits
as an example. Ultimately, we aim to ask whether spatial or social connections generally show a
stronger relationship with density, partly functioning as a proxy for indirect and direct contact events
with the potential to transmit pathogens. This framework moves between concepts of network and
contact formation traversing behavioural ecology, spatial and social network ecology, and disease
ecology.

Figure 2: The phylogenetic (A) and geographic (B) distribution of our 34 examined datasets of spatial and social behaviour, with (C) schematic depicting the methodology for deriving local density values, using the Isle of Rum red deer data as an example. The X and Y axes are bivariate spatial coordinates. The panels within (C) show raw observations of individuals in space that we then average at the individual level to make centroids; we use the centroids to generate annual density distributions, which are then assigned to individuals in the form of local density measures. Animal silhouettes are from phylopic.org; a list of attributions is in the supplement. NB the Potomac dolphins are now defined as Tursiops erebennus*; they are currently incorporated in Panel A as* T. truncatus*, following the Open Tree of Life nomenclature.*

Methods

Data standardisation and behavioural pipeline

Data were manipulated and analysed using R version 4.2.3 [36], and all R code is available at https://github.com/gfalbery/DensityMetaAnalysis. Our 34 datasets each involved at least one continuous uninterrupted spatial distribution of observations in a single population; some datasets comprised multiple such populations. These datasets covered 28 different host species, including sharks, carnivores, cetaceans, ungulates, rodents, elephants, birds,

reptiles, and one insect (Figure 2). In one case (The Firth of Tay and Moray Dolphins) we
used two distinct replicates despite being composed of overlapping groups of individuals,
because of their distinct spatial distributions, which made it difficult to fit a coherent density
distribution.

To standardise the timescale across studies, all systems were analysed as annual replicates
– i.e., social and spatial networks were summarised within each year. Our analyses used 60
system-behaviour replicates, listed in Supplementary Table 1, and totalled 151,507 unique
system-individual-year-behaviour data points.

All spatial coordinates were converted to the scale of kilometres or metres to allow
comparison across systems. To provide an approximation of local density, following prior
methodology [12,37], we took each individual's average location across the year (their
centroid) and created a spatial density kernel using the `adehabitathr` package [38], which
provides a probabilistic distribution of population density across each study system based on
the local frequencies of observed individuals. Each individual was assigned an annual
estimate of local density based on their centroid's location within this spatial density
distribution. We made these density distributions as comparable as possible between
systems by incorporating the density raster using metre squares; however, there were large
differences in density across populations that were difficult to resolve and put on the same
scale (e.g. interactions per individual/km^2 unit of density). Consequently, we scaled and
centred density to have a mean of zero and a standard deviation of one within each
population, which allowed us to focus on differences in relative slope and shape across
systems.

To provide a measure of asynchronous space sharing, we constructed home range overlap
(HRO) networks based on proportional overlap of two individuals' minimum convex polygon
(MCP; i.e., the bounding polygon around all observations of each individual in a given year).
These HRO networks were restricted to only individuals with five or more observations in a
given year to allow us to create convex polygons effectively; 10/34 (29%) systems did not
have sufficient sampling for this analysis. We also repeated our analyses with a series of
higher sampling requirements for observation numbers to ensure that our findings were
robust to this assumption. The MCP approach is relatively low-resolution, and assumes
uniform space use across an individual's home range; however, this approach is less data
intensive – and less sensitive to assumptions – than density kernel-based approaches that
would estimate variation in space use across the home range, allowing us to apply the
models across more systems, more generalisably, and more conservatively.

To provide a measure of social connectedness, we built social networks using various
approaches as defined by the original studies: direct observations of dyadic interactions (e.g.
fighting or mating); gambit of the group (GoG; i.e. membership of the same group) [39]; co-
trapping (i.e. trapped together or in adjacent traps within a given number of trapping
sessions); or direct contact measured by proximity sensors (defined by a certain distance-
based detection threshold). Notably some analyses use indirect interactions – i.e., spatial
overlap – to *approximate* direct interactions, which requires spatiotemporal coincidence,
which we caution against particularly when modelling pathogen transmission [30,40]. While
the two do often correlate, here we are not using HRO to approximate direct interaction

rates, but rather as a measure of indirect interactions (e.g., indicative of transmission of
environmental parasites).

For each social network, we scaled connection strength relative to the number of
observations of each individual in a dyad (i.e., simple ratio index [41]). Our response variable
therefore took the form of strength centrality, scaled to between 0-1 for each dyad, for each
social and spatial network. We focus on comparing density effects on social interactions and
associations with density's effects on space sharing.

210 Density-connectedness models: what forms do density effects
211 take?

We developed a novel workflow to allow us to derive and compare density's effects on
connectedness – and their drivers – in a standardised way across our wild animal systems.
These took three forms: linear models fitted to the whole dataset, nonlinear models fitted to
the whole dataset, and linear models fitted separately to low- and high-density subsets of
each dataset.

Linear models: For each system-behaviour replicate, we first fitted a linear model using the
`lm` function in R, fitting scaled density as an explanatory variable to estimate linear density
effect slopes.

Generalised additive models (GAMs): We fitted GAMs in the `mgcv` package [42] to
identify whether each density effect was better described by a linear or nonlinear
relationship, and to identify the shape of these nonlinear relationships. For each model, we
fitted a default thin plate spline with k=4 knots. This knot number was selected to reduce
overfitting in our models, which formed several fits to the data that were difficult to reconcile
with functional formats. To assess whether nonlinear models fit better than linear models, we
used Akaike Information Criterion (AIC), with a contrast of 2ΔAIC designated to distinguish
between models. See Supplementary Figure 1 for a schematic depicting our analytical
workflow.

Saturation models: To quantify whether density effects were generally saturating (i.e., that
density had steeper relationships with individuals' connectedness at lower density values),
we split the data into two portions: all values below the median density value, and all values
above the median. We then re-ran linear models examining the relationship between density
and strength in each portion. We attempted to investigate nonlinear patterns (especially
saturating effects) across all our systems using a range of other methods (e.g., comparing
specific functional relationships with nonlinear least squares), but found that they were
generally incapable of fitting well to the data in a standardised way across the many datasets
(i.e., non-convergence of nonlinear least squares using semi-automated starting estimates
across systems). As such, this approach represented a tradeoff between tractable,
generalisable model fitting, interpretability, and accurate representation of the relationship's
shape. All else being equal, we posit that investigating the relative slopes of two otherwise-
identical portions of the data is a conservative and informative method of identifying
saturation, which was our main hypothesis for the expected shape of density effects.

Meta-analysis: what factors determine the slope of density-connectedness relationships?

To characterise the typical relative slope of density effects across systems and identify the factors influencing their variation, we fitted hierarchical meta-analytical models using the `metafor` package in R. The response variable was the standardised slope of the linear density effect; because both individual network strength and density were scaled to have mean of zero and standard deviation of one in the linear regression, this is equivalent to the correlation coefficient (r) [43]. We converted all correlation coefficients into Fisher's Z (Z_r) and computed associated sampling variance. We then used an initial model that nested observations within a system-level random effect to account for within- and between-system heterogeneity [44], as 24/34 systems had more than one density effect. We used a random effect for species to account for repeat observations per animal species.

We then added a separate random effect for animal phylogeny [45]. This used a phylogenetic correlation matrix of our 28 animal species derived from the Open Tree of Life via the `rotl` package [46], with the `ape` package used to resolve multichotomies and provide branch lengths [47]. We fitted the model using the `rma.mv()` function with restricted maximum likelihood (REML), weighted by inverse sampling variance, and used variance components to quantify I^2, the contribution of true heterogeneity to the total variance in effect size.

We next fitted models with the same random effects structure that included explanatory variables. To detect whether some animals were more likely to experience density effects, we fitted **Animal group** as a factor with six categories, representing a combination of species' taxonomy and general ecology: aquatic (fish and dolphins), birds, large herbivores (elephants and ungulates), small mammals (rodents and hyraxes), carnivores, and ectotherms (insects and reptiles). We also fitted several explanatory variables indicative of greater statistical power that might increase the strength of density effects: **Geographic area** (km^2, log_{10}-transformed), **Number of years** of study, and **Number of individuals**, all of which we fitted as continuous covariates. Broadly, the animal group model was highly uninformative and competed with the other effects, and we expected that the phylogeny would be more informative, so we report the results of the model without the host group effect fitted.

We ran several different versions of these meta-analyses: first, we fitted meta-analytical models to the **overall linear models** of spatial and social interaction types separately, and then together, to investigate differences between the spatial and social networks in terms of their mean density slope. Next, we fitted duplicated versions of these models, but with the **saturation models**. These models were identical, but each system replicate had two linear estimates: one taken from the first 50% of the data (up to the median), and one to the latter 50%. By fitting a binary fixed effect of "data portion" to the meta-analytical models, this model would tell us whether the slopes were generally higher in the first portion of the data than the last (and therefore showed generally saturating shapes). We were unable to fit meta-analytical models to our GAMMs, as methods capable of meta-analysing nonlinear estimates are not yet well defined.

Figure 3: Meta-analysis revealed drivers of variation in linear density effects on individual network connectedness across systems. A) Our fitted linear model estimates of density effects on network strength. Each point represents the mean estimate from a given system; the error bars denote 95% confidence intervals. Opaque error bars were significant (i.e., do not overlap with 0); transparent ones were not. The estimates are in units of standard deviations for both density and network strength. The colour of the point denotes whether the network being examined was defined using spatial or social connections. B) Meta-analyses revealed that centrality in spatial networks (i.e., home range overlap; red points) had a significantly steeper relationship with density than social networks (blue points). C) When linear models were fitted separately to two portions of the data (values below and above the median), the slopes for the latter portion (pink points) were generally less positive than the former portion (purple points), implying a general saturation shape. In panels B) and C), each coloured point represents a study replicate fitted to the strength estimate; points are sized according to sample size, and jittered slightly on the x axis to reduce overplotting. The large black points represent the mean slope estimated from the meta-analysis, and the error bars represent 95% confidence intervals.

Figure 4: Relationships between density and network connectedness varied substantially across animal systems. Density in individuals per area is on the x axis; network connectedness (strength centrality) is on the y axis. Both values have been standardised to have a mean of zero and a standard deviation of 1 within each system; the axis ticks are in units of 1 standard deviation. Each point represents an individual-year-behaviour replicate; the lines portray the model fit from our GAMs. Red lengths of the smooth=significantly positive; grey=not significantly different from zero; blue=significantly negative. Points are semi-transparent to enhance visibility. Panels are arranged phylogenetically following the tree displayed in Figure 2A; GOG=gambit of the group; HRO=home range overlap. Supplementary Figure 2 shows a version of this plot with variable smoothing parameters to show variation in the possible shape of the curves. Animal silhouettes are from phylopic.org; a set of links and attributions are in the Supplement.

Results and Discussion

We compiled an unprecedented comparative meta-dataset of over ten million observations
of individual animals' spatial and social behaviour, across an expansive range of ecological
systems. We then ran a standardised pipeline to align their spatial and social observations,
identifying strong and predictable relationships between local density and network
connectedness at the individual level.

We observed strong positive relationships between individuals' local population density and
their connectedness in spatial and social networks across a wide range of wild animals: of
our 60 replicates, 48 (80%) were significantly positive when analysed using linear models
(Figure 3A; Supplementary Table 1). Meta-analyses identified a highly significant positive
mean correlation between density and connectedness, both for social networks (Estimate
0.22; 95% CI 0.16, 0.27) and spatial networks (0.45; 0.36, 0.53; Figure 3B). Our study
therefore provides fundamental evidence that high local population density broadly drives
greater connectedness within ecological systems, at the individual level. Slopes were highly
variable across systems for both spatial and social networks (Figure 3A; Q-test of
heterogeneity across systems: $Q_{35} = 5744.83$ and $Q_{23} = 1559.08$, both P<0.0001), indicating
that quantifying these slopes within and between multiple systems and comparing them is
important for understanding animal socio-spatial structure. That is, relationships between
density and individual connectedness differ substantially between populations, and the
biological mechanisms underlying these divergent trends are likely important. As well as
adding resolution and allowing comparisons of density effects across systems, our
methodology facilitated fitting of nonlinear relationships. This approach has only rarely been
applied before, and then at much coarser resolution (see [8,9,11]). As such, this study fills an
important empirical gap by providing insights into the slope and shape of density-
connectedness relationships for a variety of animal groups and their social and spatial
behaviours (Figure 4). Nevertheless, we were able to identify several further general trends
in our data.

Remarkably, density's effect more than doubled in size for spatial compared to social
networks (Figure 3B; r=0.45 *versus* 0.22); there was a difference of 0.28 (CI 0.17, 0.38,
P<0.0001) for this effect when we meta-analysed the two contact types together. This finding
indicates that as density increases, wild animals are more likely to share space with each
other, but that social connections increase at a much slower rate. Similarly, we discovered
that saturating shapes were extremely common: as density increased, its effect on
connectedness decreased, such that 44/60 systems (73%) had a steeper slope at low
density values than at high ones. This effect was strong for social networks (effect on r= -
0.11; CI -0.20, -0.03; P=0.01) and similar in size but marginally insignificant for spatial
networks (-0.14; -0.29, 0.01; P=0.058); due to the greater overall effect for space sharing,
the latter half of the density-spatial connections effect was still higher than the first half of the
density-social connections effect (Figure 3C). In fact, the second half of the social effect
overlapped with zero (Figure 3C; lower CI=-0.0119): that is, considering only the upper half
of density values would not have uncovered a significant effect of density on individuals'
connectedness in social networks. Together, these observations suggest that density-
dependent processes act to limit the increase in social connectedness with density, but
without limiting spatial overlaps to the same extent. Consequently, higher-density areas are
characterised disproportionately by individuals asynchronously sharing space rather than

socially associating, while in lower-density areas individuals are disproportionately more
socially connected proportional to their shared space.

There are many possible social reasons for saturating nonlinearity in density-dependent
network structuring: for example, individuals in higher density areas may begin to avoid each
other, seeking to avoid competition or aggression [31] or exposure to infectious disease [48].
Eastern water dragons (*Intellagama lesueurii*) show greater avoidance at higher densities
[13], supporting this mechanism. Alternatively, in species with high social cognition or stable
bonds, saturation could reflect lower social effort or ability to keep track of social affiliates at
higher densities [49]. In general, individuals likely have a preferred social interaction rate or
group size – a preference that they may increasingly exert at higher densities [7]. It remains
to be seen how this preference varies among individuals, and whether individuals vary in
their preferred social network position given a certain density. Given that individuals vary in
their movement and spatial phenotypes [50–52], and social phenotypes [52–54] in ways that
should manifest for density-dependent behaviours specifically [16], it seems likely that these
slopes could vary between individuals as they do between populations. Future analyses
might fit variable density-connectedness slopes between individuals to identify socio-spatial
syndromes across systems, as has been done previously in single systems including caribou
(*Rangifer tarandus*) [55] and red squirrels (*Tamiasciurus hudsonicus*) [56].

We considered spatial behaviours might explain these trends: for example, density could
create greater competition over resources and therefore reduce energy to roam (and contact
others). Individuals may partition their niches [57], or reduce their territory or home range
sizes [56,58,59], potentially driven by years of plentiful resources supporting higher densities
alongside smaller home ranges which could drive lower association rates. However, our
findings do not seem to support explanations related to small home ranges, because such
explanations should produce an equivalent or stronger reduction in (relative) spatial
connectedness. In contrast, we observed that density drove individuals to become spatially
connected faster than socially, such that the underlying mechanisms likely involve
behaviours and demographic processes that specifically affect social collocation in space
and time. Testing the precise underlying mechanism might require finer-scale behavioural
observations, as described below. Regardless of mechanism, these saturating density-
connectedness relationships strongly support the idea that examining density effects at the
individual level – rather than between populations – is highly informative. For many systems,
"mean field" expectations of homogenous interactions under increasing density likely
produce an inaccurate (i.e., inflated) picture of density's effects.

The fact that spatial networks show stronger and more linear density dependence than
social networks is likely to have important implications for the ecology of wild animal
systems. For example, indirectly transmitted (i.e., environmentally latent) parasites may
exhibit greater density dependence than directly transmitted ones, given that individuals
likely experience disproportionately more indirect contact at higher densities. This
observation contrasts with orthodoxy that contagious parasites are most likely to be density
dependent [60], and supports the value of investigating nonlinear changes in socio-spatial
behaviour and grouping patterns in response to density when considering density
dependence. Saturating density-connectedness functions further have implications for
disease modelling and control. Rather than assuming constant behavioural mixing at higher
densities, epidemiological models could benefit from incorporating density-dependent shifts

in behaviours and demography that influence direct and indirect interaction frequencies, as
previously suggested empirically and by epidemiological theory [15]. These relationships
could influence our targets for culling or vaccination coverage [61]. Given that animals at
high density seem likely to have a relatively shallow relationship between density and
contact rates, reducing population density – for example via culling – might not be effective
at reducing pathogen transmission initially, particularly when considering socially transmitted
pathogens, where contact rates are particularly likely to have saturated (Figure 3C). Similar
problems with culling have already been acknowledged in specific systems – e.g. in canine
rabies [33,62,63] – but our study implies that shallow nonlinear density-contact trends could
be more general than previously thought and could be driven by flexible density-dependent
changes in behaviour and demographic processes.

Operationally, the common nature of saturating density effects will impact researchers' ability
to detect density dependence: that is, density dependence could be harder to observe in
higher-density areas given the shallower slopes we observed. Most of the systems in this
study are relatively long-term studies of known individuals; these populations tend to be in
carefully selected, high-density areas that make it convenient to study the focal animal with
relatively low operational costs. For example, it has previously been noted that the badgers
of Wytham Wood, the red deer of the Isle of Rum, and the Soay sheep of St Kilda are all at
high densities for their respective species [64–66]. As such, we may be inherently
investigating the upper end of density-connectedness relationships in the wild, and it could
be difficult and costly to investigate the effects of low density so widely.

Beyond these general trends, our GAMs revealed that 46/60 density effects on network
connectedness (77%) were significantly nonlinear (ΔAIC>2); these relationships took a wide
variety of shapes, representing a range of nonlinear functions that are hard to generalise
(Figure 4; Supplementary Table 2). Notably, while many GAM smooths were eventually
significantly negative (Figure 4), the vast majority of linear models fitted to the second half of
the data were positive (Figure 3C); this result is likely an artefact of restricted model fitting,
rather than true downturns in connectedness with density. Nonlinearity did not cluster
according to connection type definitions, or according to host group. These observations
were largely corroborated by our meta-analytical models, which found no factors influencing
the slope and shape of density effects (P>0.05). This observation speaks to the complexity
of these relationships within and across systems, while accentuating that simple functional
relationships are often likely to be complicated by contravening ecological factors like habitat
selection [67,68], parasite avoidance [69], and demographic structuring [70]. While we were
unable to identify specific between-system predictors of nonlinearity of density-
connectedness relationships, the finding that most such relationships are strongly nonlinear
is an important consideration for future work.

We acknowledge several limitations of our study, which we nevertheless believe could be
remedied in the future. First, many of our social networks were formed of general
spatiotemporal associations, and relatively few from specific social interactions – particularly
those involving direct physical contact. Our current dataset could therefore benefit from
supplementation with a broader range of direct interactions, particularly involving antagonism
or bonding. For example, datasets concerning aggression or dominance interactions (e.g.
[71]) or grooming alongside spatial behaviour could inform how density dependence affects
the transmission of certain parasites such as mycobacteria [72] or tattoo skin disease [73].

The meta-dataset was also unevenly distributed across animal taxa (Figure 1): there were
no primates or bony fishes and only one invertebrate, while rodents and ungulates were
over-represented. These biases likely emerge through differences in data collection
approaches: for example, although primate social behaviour is often studied with
observations of direct interactions that could augment our data as described above (e.g.
[74]), the spatial data required to build density distributions are rarely collected in these
systems. This is linked to their social structures: our workflow was best suited to studies of
fission-fusion societies or relatively asocial animals, rather than those with wide-ranging
fixed social groups that are more common in primate systems. Finally, given that our data
were observational, we could not account for (or estimate) bidirectional causality between
density and social relationships (point 4 in Figure 1): that is, as well as encountering more
conspecifics in areas of high density, individuals may be drawn to conspecifics, *creating*
areas of high density [16]. To do so might require creating in-depth, high-resolution models
of animal movement and group formation (e.g. [75]), potentially making use of telemetry
approaches and drawing from large-scale open movement repositories like Movebank [76].

Aside from incorporating more specific interaction types, there exist a range of potential
extensions to our analysis. For example, density dependence often varies between age or
sex classes (e.g. [77,78]), and age effects on infection are common and ecologically
important [79,80], as are sex differences [81]. We chose not to analyse how individual
animals' traits alter the shape or slope of density's effects for brevity and simplicity; however,
given that many of the systems nevertheless include these data, future analyses could make
use of this meta-dataset to investigate how density affects connectedness of different
classes of hosts. Further, researchers could investigate other behavioural questions such as
the role of observation biases; the factors influencing the correlation between spatial and
social networks; and the role of environmental drivers and spatial autocorrelation in driving
observed patterns of connectivity [12,16]. Finally, as our analysis approximated density-
contact relationships and not host-parasite interactions specifically, important future work
could investigate whether contact rates (as approximated by network connectedness)
ultimately translate to greater infection risk or parasite burden. Although some previous
investigations have linked density-related metrics to aspects of infection [82,83], density
covaries with a range of other factors including nutrition, cooperation, and competition, all of
which could complicate density-driven increases in exposure [2]. For example, in the case of
ectoparasite transmission, although contact rates in general would likely increase with
density, so too might grooming behaviours that remove parasites; in cases such as these,
density's overall effect on ectoparasite infection may be neutral. In the future, verifying that
within- and between-population variation in density-contact relationships translate to
variation in infection – and whether these trends might be influenced by flexible avoidance
behaviours [37] – will be a vital part of understanding and predicting density-dependent
disease dynamics.

Density is a universal factor underlying the dynamics of animal populations, and its linear
and nonlinear effects on spatial and social network structure are likely to impact myriad
processes in behaviour, ecology, and evolution. Similar to other studies that have reported
general scaling patterns in network analysis [84] and in food web ecology [85], the patterns
we report strongly suggest that animal systems generally become better connected spatially
than socially under increasing density. These might extrapolate to human networks, given
that other scaling patterns in animal networks do [84]. As these patterns seemingly manifest

regardless of animal group and interaction type, they may reflect a generalisable rule
governing the socio-spatial structure of ecological systems. Further refining and
implementing these models could facilitate prediction of network structure in novel systems.

Acknowledgements

GFA and SB were supported by NSF grant number DEB-2211287. GFA acknowledges
support from a College for Life Sciences Fellowship at the Wissenschaftskolleg zu Berlin and
WAI (C-2023-00057). DJB acknowledges support from the Edward Mallinckrodt, Jr.
Foundation and NSF BII 2213854. JAF acknowledges funding from BBSRC
(BB/S009752/1), NERC (NE/S010335/1 and NE/V013483/1), and WAI (C-2023-00057). For
a list of system-specific acknowledgements, please see the supplement.

Code availability

504 All R code is available at https://github.com/gfalbery/DensityMetaAnalysis.

Affiliations

1. Department of Biology, Georgetown University, Washington, DC, USA
2. Wissenschaftskolleg zu Berlin, Berlin, Germany
3. School of Biological Sciences, University of Oklahoma, Norman, OK, USA
4. Department of Biology, University of Oxford, Oxford, UK
5. School of Biology, University of Leeds, Leeds, UK
6. CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
7. Centre for Ecology and Conservation, University of Exeter, Penryn, UK
8. School of Biosciences, University of Sheffield, Sheffield, UK
9. Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, UK
10. Department of Biology, Memorial University of Newfoundland, 45 Arctic Avenue,
St. John's NL
11. University of Guelph, Guelph, Canada
12. Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool,
Liverpool, UK
13. Institute of Zoology, Zoological Society of London, Regents Park, London, UK
14. University of Glasgow, Glasgow, UK
15. Department of Biological Sciences, Old Dominion University, Norfolk, VA, USA
16. Avian Ecology Program, Archbold Biological Station, Venus, FL, USA
17. University of Salford, Salford, UK
18. Department of Ecology and Evolutionary Biology, University of California, Santa Cruz,
CA, USA
19. Canine Companions, 2965 Dutton Ave, Santa Rosa, CA, USA
20. Hopkins Marine Station, Stanford University, Stanford, CA, USA
21. University of Nottingham, Nottingham, UK
22. Pitzer and Scripps Colleges, Claremont, CA, USA
23. Wildlife Conservation Research Unit, The Recanati-Kaplan Centre, Department of
Zoology, University of Oxford, Oxford, UK
24. The University of British Columbia, Irving K. Barber Faculty of Sciences, Okanagan
Department of Biology, Kelowna, British Columbia, Canada

25. UK Centre for Ecology and Hydrology, Bush Estate, Midlothian, UK
26. University of Delaware, Newark, DE, USA
27. Marine Science Institute, UC Santa Barbara, Santa Barbara, CA, USA
28. ISI Foundation, Torino, Italy
29. Evolutionary Ecology Group, Station d'Ecologie Theorique et Experimentale du CNRS, Moulis, France
30. Oregon State University, Corvallis, OR, USA
31. Lighthouse Field Station, School of Biological Sciences, University of Aberdeen, UK
32. Department of Zoology, University of Cambridge, Cambridge, UK
33. Institute of Zoology, Zoological Society of London, London, UK
34. National Wildlife Management Centre, Animal and Plant Health Agency, Sand Hutton, York, UK
35. Department of Collective Behavior, Max Planck Institute for Animal Behavior, 78464 Konstanz, Germany
36. Department of Evolutionary Biology and Environmental Studies, University of Zurich, 8057 Zurich, Switzerland
37. Department of Ornithology, National Museum of Kenya, P.O. Box 40241-001000, Nairobi, Kenya
38. Division of Ecology and Evolution, Research School of Biology, Australian National University, 46 Sullivan Creek Road, Canberra, ACT 2600, Australia
39. Virginia Tech, Blacksburg, VA, USA
40. The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK
41. Environment and Sustainability Institute, University of Exeter, Penryn, UK
42. Department of Marine Biology, Texas AandM University at Galveston, USA
43. School of the Environment, University of Queensland, Brisbane, Queensland, Australia
44. College of Science and Engineering, Flinders University, Adelaide, Australia
45. School of Zoology, Faculty of Life Science, Tel Aviv University, Tel Aviv, Israel
46. Department of Zoology, University of Otago, Otago, New Zealand
47. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA
48. Sea Mammal Research Unit, South Oceans Institute, University of St Andrews, UK
49. Department of Visitor Management and National Park Monitoring, Bavarian Forest National Park, Grafenau, Germany
50. Chair of Wildlife Ecology and Wildlife Management, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany
51. Department of Forestry and Wildlife Management, Faculty of Applied Ecology, Agricultural Sciences and Biotechnology, Campus Evenstad, Inland Norway University of Applied Sciences, Postboks 400 Vestad, 2418 Elverum, Norway
52. Faculty of Life Sciences, Bar Ilan University, Ramat Gan, Israel
53. Lancaster University, Lancaster, UK
54. Division of Marine Science and Conservation, Nicholas School of the Environment, Duke University Marine Laboratory, Beaufort, NC, USA
55. Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech Republic
56. Evolutionary Ecology Group, EVECO, Universiteit Antwerpen, Antwerp, Belgium
57. Department of Nature Conservation and Research, Hainich National Park, Bad Langensalza, Germany
58. Mpala Research Center, P.O Box 555-10400, Nanyuki, 10400, Kenya

59. Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78464
Konstanz, Germany
60. School of Environment and Natural Science, Bangor University, Bangor, Wales
61. Department of Biology, Memorial University of Newfoundland, St John's, NL, Canada
62. Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada
63. University of Aberdeen, Aberdeen, UK
64. School of Animal and Veterinary Sciences, The University of Adelaide, Adelaide,
Australia
65. School of Biological Sciences, University of Nebraska-Lincoln, NE, USA
66. Department of Ecology, Behavior, and Evolution, University of California, San Diego, CA
67. Department of Evolutionary Biology and Environmental Studies, University of Zurich,
Zurich, Switzerland
68. Institute of Pest Management, Sokoine University of Agriculture, Morogoro, Tanzania
69. Max Planck Institute of Animal Behavior, Radolfzell, Germany
70. MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France
71. PSL Research University, Université de Perpignan, Perpignan, France
72. Laboratoire d'Excellence "CORAIL", Papetoai, Moorea, French Polynesia
73. University of Nevada, Reno, NV, USA
74. University of Eldoret, School of Natural Resource Management, Department of Wildlife,
1125-30100 Eldoret, Kenya
75. Department of Veterinary Sciences, University of Turin, Grugliasco (TO), Italy
76. Florida International University, FL, USA
77. University of Liverpool, Liverpool, UK
78. Department of Environmental Science and Policy, University of California Davis, Davis,
CA, USA
79. Department of Ecology and Evolutionary Biology, University of California, Los Angeles,
CA, USA
80. Stanford University, Stanford, CA, USA
81. School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA
82. Department of Wildlife, California State Polytechnic University, Humboldt, Arcata, CA,
USA
83. Texas Tech University, Lubbock, TX, USA
84. University of Western Australia, Crawley WA 6009, Australia
85. Department of Veterinary Population Medicine, University of Minnesota, Minneapolis, US
86. University of Surrey, Guildford, UK

References

1. Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latto J, Sanchez MS,
Smith AB, Swei A. 2005 Should we expect population thresholds for wildlife disease?
Trends Ecol Evol **20**, 511–519. (doi:10.1016/j.tree.2005.07.004)

2. Albery GF. 2022 Density dependence and disease dynamics: moving towards a
predictive framework. *EcoEvoRxiv Preprints* (doi:10.32942/OSF.IO/GAW49)

3. Hopkins SR, Fleming-Davies AE, Belden LK, Wojdak JM. 2020 Systematic review of
modeling assumptions and empirical evidence: does parasite transmission increase

nonlinearly with host density? *Methods Ecol Evol* **11**, 0–2. (doi:10.1111/2041-
210x.13361)

4. Begon M, Bennett M, Bowers RG, French NP, Hazel SM, Turner J. 2002 A
clarification of transmission terms in host-microparasite models: Numbers, densities
and areas. *Epidemiol Infect* **129**, 147–153. (doi:10.1017/S0950268802007148)

5. Ferrari MJ, Perkins SE, Pomeroy LW, Bjrnstad ON. 2011 Pathogens, social networks,
and the paradox of transmission scaling. *Interdiscip Perspect Infect Dis* **2011**.
(doi:10.1155/2011/267049)

6. Webber QMR, Vander Wal E. 2020 Heterogeneity in social network connections is
density-dependent: implications for disease dynamics in a gregarious ungulate. *Behav
Ecol Sociobiol* **74**, 1–13.

7. Beauchamp G. 2011 Functional relationship between group size and population
density in Northwest Atlantic seabirds. *Mar Ecol Prog Ser* **435**, 225–233.
(doi:10.3354/meps09239)

8. Borremans B, Reijniers J, Hens N, Leirs H. 2017 The shape of the contact–density
function matters when modelling parasite transmission in fluctuating populations. *R
Soc Open Sci* **4**. (doi:10.1098/rsos.171308)

9. Davis S, Abbasi B, Shah S, Telfer S, Begon M. 2015 Spatial analyses of wildlife
contact networks. *J R Soc Interface* **12**. (doi:10.1098/rsif.2014.1004)

10. Carson BD, Orians CM, Crone EE. 2023 When does spatial clustering in caterpillars
inuence the relationship between population density and contact rates?
(doi:10.21203/rs.3.rs-2842538/v1)

11. Mancy R *et al.* 2022 Rabies shows how scale of transmission can enable acute
infections to persist at low prevalence. *Science (1979)* **376**, 512–516.

12. Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, Firth
JA. 2021 Multiple spatial behaviours govern social network positions in a wild
ungulate. *Ecol Lett* **24**, 676–686. (doi:10.1101/2020.06.04.135467)

13. Strickland K, Patterson EM, Frère CH. 2018 Eastern water dragons use alternative
social tactics at different local densities. *Behav Ecol Sociobiol* **72**.
(doi:10.1007/s00265-018-2563-x)

14. Strickland K, Frère CH. 2019 Individual Variation in the Social Plasticity of Water
Dragons. *Am Nat* **194**, 194–206. (doi:10.5061/dryad.5j5gh63)

15. Hu H, Nigmatulina K, Eckhoff P. 2013 The scaling of contact rates with population
density for the infectious disease models. *Math Biosci* **244**, 125–134.
(doi:10.1016/j.mbs.2013.04.013)

16. Webber QMR, Albery GF, Farine DR, Pinter-wollman N, Sharma N, Vander Wal E,
Manlove K. 2023 Behavioural ecology at the spatial-social interface. *Biological*
Reviews , 1–40.

17. Albon SD, Staines HJ, Guinness FE, Clutton-Brock TH. 1992 Density-Dependent
Changes in the Spacing Behaviour of Female Kin in Red Deer. *Source: Journal of*
Animal Ecology. **61**.

18. Vas J, Andersen IL. 2015 Density-dependent spacing behaviour and activity budget in
pregnant, domestic goats (Capra hircus). *PLoS One* **10**.
(doi:10.1371/journal.pone.0144583)

19. Connor EF, Cargain MJ. 1994 Density-related foraging behaviour in Closterocerus
tricinctus, a parasitoid of the leaf-mining moth, Cameraria hamadryadella. *Ecol*
Entomol **19**, 327–334. (doi:10.1111/J.1365-2311.1994.TB00249.X)

20. Matthysen E. 2005 Density-dependent dispersal in birds and mammals. *Ecography*
28, 403–416. (doi:10.1111/j.0906-7590.2005.04073.x)

21. Avgar T, Betini GS, Fryxell JM. 2020 Habitat selection patterns are density dependent
under the ideal free distribution. *Journal of Animal Ecology* **89**, 2777–2787.
(doi:10.1111/1365-2656.13352)

22. Vander Wal E, Laforge MP, Mcloughlin PD, Wal E Vander, Laforge MP, Mcloughlin
PD, Vander Wal E, Laforge MP, Mcloughlin PD. 2014 Density dependence in social
behaviour: Home range overlap and density interacts to affect conspecific encounter
rates in a gregarious ungulate. *Behav Ecol Sociobiol* **68**, 383–390.
(doi:10.1007/s00265-013-1652-0)

23. Kokko H, Rankin DJ. 2006 Lonely hearts or sex in the city? Density-dependent effects
in mating systems. In *Philosophical Transactions of the Royal Society B: Biological*
Sciences, pp. 319–334. Royal Society. (doi:10.1098/rstb.2005.1784)

24. Modlmeier AP, Colman E, Hanks EM, Bringenberg R, Bansal S, Hughes DP. 2019
Ant colonies maintain social homeostasis in the face of decreased density. *Elife* **8**, 1–
17. (doi:10.7554/eLife.38473)

25. Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting animal
social network analysis. *Journal of Animal Ecology* **84**, 1144–1163.
(doi:10.1111/1365-2656.12418)

26. Farine DR. 2024 Modelling animal social networks: New solutions and future
directions. *Journal of Animal Ecology* (doi:10.1111/1365-2656.14049)

27. Webber QMR, Vander Wal E. 2019 Trends and perspectives on the use of animal
social network analysis in behavioural ecology: a bibliometric approach. *Anim Behav*
149, 77–87. (doi:10.1016/j.anbehav.2019.01.010)

28.   Pinter-Wollman N *et al.* 2014 The dynamics of animal social networks: Analytical,
conceptual, and theoretical advances. *Behavioral Ecology* **25**, 242–255.
(doi:10.1093/beheco/art047)

29.   Sah P, Mann J, Bansal S. 2018 Disease implications of animal social network
structure: A synthesis across social systems. *Journal of Animal Ecology* **87**, 546–558.
(doi:10.1111/1365-2656.12786)

30.   Albery GF, Kirkpatrick L, Firth JA, Bansal S. 2021 Unifying spatial and social network
analysis in disease ecology. *Journal of Animal Ecology* **90**, 1–17. (doi:10.1111/1365-
2656.13356)

31.   Cubaynes S, Macnulty DR, Stahler DR, Quimby KA, Smith DW, Coulson T. 2014
Density-dependent intraspecific aggression regulates survival in northern Yellowstone
wolves (Canis lupus). *Journal of Animal Ecology* **83**, 1344–1356. (doi:10.1111/1365-
2656.12238)

32.   Lloyd-Smith JO, Getz WM, Westerhoff H V. 2004 Frequency-dependent incidence in
models of sexually transmitted diseases: Portrayal of pair-based transmission and
effects of illness on contact behaviour. *Proceedings of the Royal Society B: Biological*
Sciences **271**, 625–634. (doi:10.1098/rspb.2003.2632)

33.   Morters MK, Restif O, Hampson K, Cleaveland S, Wood JLN, Conlan AJK. 2013
Evidence-based control of canine rabies: A critical review of population density
reduction. *Journal of Animal Ecology* **82**, 6–14. (doi:10.1111/j.1365-
2656.2012.02033.x)

34.   Gordon DM, Paul RE, Thorpe K. 1993 What is the function of encounter patterns in
ant colonies? *Anim Behav* **45**, 1083–1100. (doi:10.1006/ANBE.1993.1134)

35.   Davis S, Abbasi B, Shah S, Telfer S, Begon M, Davis S. 2015 Spatial analyses of
wildlife contact networks. *Journal of the Royal Society, Interface / the Royal Society*
12, 20141004. (doi:10.1098/rsif.2014.1004)

36.   R Development Core Team R. 2011 R: A Language and Environment for Statistical
Computing. *R Foundation for Statistical Computing.* **1**, 409. (doi:10.1007/978-3-540-
74686-7)

37.   Albery GF, Newman C, Bright Ross J, Macdonald DW, Bansal S, Buesching CD.
2020 Negative density-dependent parasitism in a group-living carnivore. *Proceedings*
of the Royal Society B: Biological Sciences **287**, 20202655.
(doi:10.1101/2020.06.15.153726)

38.   Calenge C. 2011 Home range estimation in R: the adehabitatHR package. See
https://cran.r-project.org/web/packages/adehabitatHR/index.html (accessed on 10
March 2020).

39.   Franks DW, Ruxton GD, James R. 2010 Sampling animal association networks with
the gambit of the group. *Behav Ecol Sociobiol* **64**, 493–503. (doi:10.1007/s00265-009-
0865-8)

40. Wanelik KM, Farine DR. 2022 A new method for characterising shared space use networks using animal trapping data. *Behav Ecol Sociobiol* **76**. (doi:10.1007/s00265-022-03222-5)

41. Cairns SJ, Schwager SJ. 1987 A comparison of association indices. *Anim Behav* **35**, 1454–1469. (doi:10.1016/S0003-3472(87)80018-0)

42. Wood SN. 2011 Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J R Stat Soc Series B Stat Methodol* **73**, 3–36. (doi:10.1111/j.1467-9868.2010.00749.x)

43. Koricheva J, Gurevitch J, Mengersen KL. 2013 Handbook of meta-analysis in ecology and evolution. , 498.

44. Konstantopoulos S. 2011 Fixed effects and variance components estimation in three-level meta-analysis. *Res Synth Methods* **2**, 61–76. (doi:10.1002/jrsm.35)

45. Cinar O, Nakagawa S, Viechtbauer W. 2022 Phylogenetic multilevel meta-analysis: A simulation study on the importance of modelling the phylogeny. *Methods Ecol Evol* **13**, 383–395. (doi:10.1111/2041-210X.13760)

46. Michonneau F, Brown JW, Winter DJ. 2016 rotl: an R package to interact with the Open Tree of Life data. *Methods Ecol Evol* **7**, 1476–1481. (doi:10.1111/2041-210X.12593)

47. Paradis E, Schliep K. 2019 ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528. (doi:10.1093/BIOINFORMATICS/BTY633)

48. Gibson AK, Amoroso CR. 2022 Evolution and Ecology of Parasite Avoidance. , 1–21.

49. Colman E, Colizza V, Hanks EM, Hughes DP, Bansal S. 2021 Social fluidity mobilizes contagion in human and animal populations. *Elife* **10**. (doi:10.7554/ELIFE.62177)

50. Michelangeli M, Payne E, Spiegel O, Sinn D, Leu ST, Gardner M, Sih A. 2021 *Personality, spatiotemporal ecological variation, and resident/explorer movement syndromes in the sleepy lizard.* (doi:10.1111/1365-2656.13616)

51. Spiegel O, Leu ST, Bull CM, Sih A. 2017 What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecol Lett* **20**, 3–18. (doi:10.1111/ele.12708)

52. Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014 Individual-level personality influences social foraging and collective behaviour in wild birds. *Proceedings of the Royal Society of London B* **281**, 20141016. (doi:10.1098/rspb.2014.1016)

53. Hunt ER, Mi B, Fernandez C, Wong BM, Pruitt JN, Pinter-Wollman N. 2018 Social interactions shape individual and collective personality in social spiders. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20181366. (doi:10.1098/rspb.2018.1366)

54.   Firth JA, Cole EF, Ioannou CC, Quinn JL, Aplin LM, Culina A, McMahon K, Sheldon
774   BC. 2018 Personality shapes pair bonding in a wild bird social system. *Nat Ecol Evol*
775   **2**, 1696–1699. (doi:10.1038/s41559-018-0670-8)

776   55.   Webber Q, Laforge M, Bonar M, Wal E Vander. 2022 The adaptive value of density-
777   dependent habitat specialization and social network centrality. *Authorea Preprints*
778   (doi:10.22541/AU.165210484.46672609/V1)

779   56.   Webber QMR, Dantzer B, Lane JE, Boutin S, McAdam AG. 2023 Density-dependent
780   plasticity in territoriality revealed using social network analysis. *Journal of Animal*
781   *Ecology* **92**, 207–221. (doi:10.1111/1365-2656.13846)

782   57.   Sheppard CE, Inger R, McDonald RA, Barker S, Jackson AL, Thompson FJ,
783   Vitikainen EIK, Cant MA, Marshall HH. 2018 Intragroup competition predicts individual
784   foraging specialisation in a group-living mammal. *Ecol Lett* **21**, 665–673.
785   (doi:10.1111/ELE.12933)

786   58.   Sanchez JN, Hudgens BR. 2015 Interactions between density, home range
787   behaviors, and contact rates in the Channel Island fox (Urocyon littoralis). *Ecol Evol* **5**,
788   2466–2477. (doi:10.1002/ece3.1533)

789   59.   Wilkin TA, Garant D, Gosler AG, Sheldon BC. 2006 Density effects on life-history
790   traits in a wild population of the great tit Parus major: Analyses of long-term data with
791   GIS techniques. *Journal of Animal Ecology* **75**, 604–615. (doi:10.1111/j.1365-
792   2656.2006.01078.x)

793   60.   Cote IM, Poulin R. 1995 Parasitism and group size in social animals: a meta-analysis.
794   *Behavioral Ecology* **6**, 159–165. (doi:10.1093/beheco/6.2.159)

795   61.   McCallum H, Barlow N, Hone J. 2001 How should pathogen transmission be
796   modelled? *Trends Ecol Evol* **16**, 295–300. (doi:10.1016/S0169-5347(01)02144-9)

797   62.   Rajeev M, Metcalf CJE, Hampson K. 2020 Modeling canine rabies virus transmission
798   dynamics. In *Rabies*, pp. 655–670. Elsevier. (doi:10.1016/b978-0-12-818705-
799   0.00020-0)

800   63.   Townsend SE *et al.* 2013 Designing Programs for Eliminating Canine Rabies from
801   Islands: Bali, Indonesia as a Case Study. *PLoS Negl Trop Dis* **7**.
802   (doi:10.1371/JOURNAL.PNTD.0002372)

803   64.   Macdonald DW, Newman C. 2022 *The Badgers of Wytham Woods: a model for*
804   *behaviour, ecology, and evolution*. Oxford University Press.

805   65.   Clutton-Brock TH, Guinness FE, Albon SD. 1982 *Red Deer: Behavior and Ecology of*
806   *Two Sexes*. Chicago, IL: University of Chicago Press. See
807   https://books.google.co.uk/books/about/Red_Deer.html?id=x4SGuA3t-NoC&pgis=1.

808   66.   Clutton-Brock TH, Pemberton JM. 2004 *Soay sheep : population dynamics and*
809   *selection on St. Kilda*. Cambridge, UK: Cambridge University Press.

67. Tardy O, Massé A, Pelletier F, Mainguy J, Fortin D. 2014 Density-dependent
functional responses in habitat selection by two hosts of the raccoon rabies virus
variant. *Ecosphere* **5**, 1–16. (doi:10.1890/ES14-00197.1)

68. Wilber MQ, Yang A, Boughton R, Manlove KR, Miller RS, Pepin KM, Wittemyer G.
2022 A model for leveraging animal movement to understand spatio-temporal disease
dynamics. *Ecol Lett* , 1–15. (doi:10.1111/ele.13986)

69. Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary Consequences
of   Parasite   Avoidance.   *Trends   Ecol   Evol*   **33**,   619–632.
(doi:10.1016/j.tree.2018.05.001)

70. Lion S, Boots M. 2010 Are parasites '"prudent"' in space? *Ecol Lett* **13**, 1245–1255.
(doi:10.1111/j.1461-0248.2010.01516.x)

71. Strauss ED, Decasien AR, Galindo G, Hobson EA, Shizuka D, Curley JP. 2022
DomArchive: a century of published dominance data. *Philosophical Transactions of*
the Royal Society B **377**. (doi:10.1098/RSTB.2020.0436)

72. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in
wild meerkats. *Proceedings. Biological sciences / The Royal Society* **277**, 633–42.
(doi:10.1098/rspb.2009.1775)

73. Powell SN, Wallen MM, Miketa ML, Krzyszczyk E, Foroughirad V, Bansal S, Mann J.
2021 Sociality and tattoo skin disease among bottlenose dolphins in Shark Bay,
Australia. *Behavioral Ecology* **31**, 459–466. (doi:10.1093/BEHECO/ARZ207)

74. Testard C *et al.* 2020 Rhesus Macaques Build New Social Connections after a Natural
Disaster. *Current Biology* , 1–11. (doi:10.2139/ssrn.3741230)

75. Spiegel O, Leu ST, Sih A, Bull CM. 2016 Socially interacting or indifferent
neighbours☐? Randomization of movement paths to tease apart social preference
and   spatial   constraints.   *Methods   Ecol   Evol* ,   971–979.   (doi:10.1111/2041-
210X.12553)

76. Kays R *et al.* 2021 The Movebank system for studying global animal movement and
demography. *Methods Ecol Evol* **2021**, 1–13. (doi:10.1111/2041-210X.13767)

77. Batsuren E, Zhang X, Song M, Wan X, Li G, Liu J, Huang S, Zhang Z. 2022 Density-
dependent changes of mating system and family structure in Brandt's voles
(*Lasiopodomys brandtii*). *Ecol Evol* , 1–16. (doi:10.1002/ece3.9199)

78. Vander Wal E, van Beest FM, Brook RK. 2013 Density-Dependent Effects on Group
Size Are Sex-Specific in a Gregarious Ungulate. *PLoS One* **8**, e53777.
(doi:10.1371/JOURNAL.PONE.0053777)

79. Ashby B, Bruns E. 2018 The evolution of juvenile susceptibility to infectious disease.
Proceedings of the Royal Society B: Biological Sciences **285**, 20180844.
(doi:10.1098/rspb.2018.0844)

80. Albery GF, Sweeny AR, Webber Q. In press. How behavioural ageing affects
infectious disease.

81. Zuk M. 2009 The sicker sex. *PLoS Pathog* **5**, 1–3. (doi:10.1371/journal.ppat.1000267)

82. Arneberg P, Skorping A, Grenfell B, Read AF. 1998 Host densities as determinants of
abundance in parasite communities. *Proceedings of the Royal Society B: Biological*
Sciences **265**, 1283–1289. (doi:10.1098/rspb.1998.0431)

83. Arneberg P. 2001 An ecological law and its macroecological consequences as
revealed by studies of relationships between host densities and parasite prevalence.
Ecography **24**, 352–358. (doi:10.1111/j.1600-0587.2001.tb00208.x)

84. Rocha LEC, Ryckebusch J, Schoors K, Smith M. 2021 Scaling of social interactions
across animal species. *Sci Rep* **11**, 12584.

85. Lafferty KD, DeLeo G, Briggs CJ, Dobson AP, Gross T, Kuris AM. 2015 A general
consumer-resource population model. *Science (1979)* **349**, 854–857.
(doi:10.1126/SCIENCE.AAA6224/SUPPL_FILE/LAFFERTY.SM.PDF)

861