A distinct isoform of Msp300 (nesprin) organizes the perinuclear microtubule organizing center in adipocytes

Garret M. Morton[1], Maria Pilar Toledo[1], Chunfeng Zheng[1], Yiming Zheng[2#], Timothy L. Megraw[1#*]

Running title: An Msp300 isoform controls the ncMTOC in adipose cells

[1]Department of Biomedical Sciences, Florida State University, Tallahassee, FL, USA

[2]State Key Laboratory of Cellular Stress Biology, School of Life Sciences, Faculty of Medicine and Life Sciences, Xiang'an Hospital of Xiamen University, Xiamen University, Xiamen, China, 361102, and Shenzhen Research Institute of Xiamen University, Shenzhen, China, 518057

[#]Co-corresponding authors

*Correspondence:

timothy.megraw@med.fsu.edu

Abstract

In many cell types, disparate non-centrosomal microtubule-organizing centers (ncMTOCs) replace functional centrosomes and serve the unique needs of the cell types in which they are formed. In Drosophila fat body cells (adipocytes), an ncMTOC is organized on the nuclear surface. This perinuclear ncMTOC is anchored by Msp300, encoded by one of two nesprin-encoding genes in Drosophila. Msp300 and the spectraplakin short stop (shot) are co-dependent for localization to the nuclear envelope to generate the ncMTOC where they recruit the microtubule minus-end stabilizer Patronin (CAMSAP). The *Msp300* gene is complex, encoding at least eleven isoforms. Here we show that two Msp300 isoforms, Msp300-PE and -PG, are required and only one, Msp300-PE, appears sufficient for generation of the ncMTOC. Loss of Msp300-PE and -PG results in severe loss of localization of shot and Patronin, disruption of the MT array, nuclear mispositioning and loss of endosomal trafficking. Furthermore, upon loss of Msp300-PE and -PG, other isoforms are retained at the nuclear surface despite the loss of nuclear positioning and MT organization, indicating that they are not sufficient to generate the ncMTOC. Msp300-PE has an unusual domain structure including a lack of a KASH domain and very few spectrin repeats and appears therefore to have a highly derived function to generate an ncMTOC on the nuclear surface.

Introduction

The <u>l</u>inker of <u>n</u>ucleoskeleton and <u>c</u>ytoskeleton (LINC) complex spans the nuclear envelope and provides a physical connection between the nuclear interior with the cytoplasm and plasma membrane [1, 2]. The essential roles of the LINC complex in cell function are underscored by inherited diseases associated with mutations in LINC complex protein genes [3, 4] including muscular dystrophies [5, 6], cardiomyopathies [7, 8], cerebellar ataxia [9, 10], and arthrogryposis [11]. The LINC complex is integral to cellular processes such as DNA damage response [12, 13], meiotic chromosome movement [14, 15], mechanosensation [1], and nuclear positioning and anchoring [3, 16-19]. Comprising the LINC complex are SUN (<u>Sa</u>d1/<u>UN</u>C-84 homology) and KASH (<u>K</u>larsicht/<u>A</u>NC-1/<u>S</u>yne <u>h</u>omology) proteins. SUN proteins reside at the inner nuclear membrane and contain an N-terminus that extends into the nucleoplasm, where it interacts with lamins of the nucleoskeleton and other nucleoplasm proteins, while the C-terminal SUN domain forms contacts with KASH proteins in the perinuclear space. The KASH proteins reside on the outer nuclear membrane, associate with cytoskeletal components, and are characterized by a C-terminal KASH domain that associates with the SUN domain of the SUN proteins within the nuclear intermembrane space. In the absence of a SUN domain protein interaction in the intermembrane space, the KASH protein does not localize to the outer nuclear membrane [1, 20].

In *Drosophila*, the LINC complex components include two KASH proteins, Klarsicht (Klar) and <u>M</u>uscle-<u>s</u>pecific protein <u>300</u> kDa (Msp300), and one primary SUN protein, Klaroid (Koi), while a second SUN protein, Spag4, is expressed exclusively in the testes. The Drosophila LINC proteins are responsible for the proper positioning of muscle myonuclei [21], similar to their roles in vertebrate muscle [6, 22-24]. Msp300, which was first identified in the muscle [25], promotes myonuclear spacing [26]. These findings implicate nuclear mispositioning in the pathogenesis of diseases such as Emery-Dreifuss muscular dystrophy (EDMD) [26, 27]. As seen with other <u>n</u>uclear <u>e</u>nvelope <u>s</u>pectrin <u>r</u>epeat <u>prote</u>ins (nesprins), Msp300 contains calponin homology (CH) domains, numerous spectrin repeats (SR), and a C-terminal KASH domain. In addition, *Msp300* encodes at least eleven isoforms of various molecular weights expressed from a complex locus (Figure 1). Furthermore, subsets of these isoforms have been shown to exhibit unique functions. In the muscle, isoforms containing the CH domains but lacking the KASH domain were found to contribute to Z-line organization and for optimal muscle function, while isoforms lacking the CH domains but containing the KASH domain were shown to be responsible for maintaining myonuclear positioning [28].

Drosophila fat body cells, which are functionally comparable to mammalian adipocytes and liver cells, are characterized by a nucleus positioned at the center of the cell and MTs organized at the nuclear surface that radiate outwards and connect the nucleus to the plasma membrane [29, 30]. The perinuclear fat body ncMTOC controls critical cellular functions, including microtubule organization, nuclear positioning, and endocytic vesicle trafficking [29]. Two general classes of MTs are organized at the perinuclear ncMTOC, circumferential MTs surround the nucleus, while radial MTs extend from the nucleus into the cytoplasm. The control of MT assembly at the fat body perinuclear ncMTOC has unique molecular features. Assembly of MTs at this ncMTOC requires Patronin, Ninein, and msps (ortholog of ch-TOG), but does not require

γ-tubulin [29]. Patronin (CAMSAP ortholog) is a MT stabilizer involved at many ncMTOCs [31-
37], Ninein is a MT anchor that associates with dynein light intermediate chain [38-40] and with
ensconsin/MAP7 and cooperates with ensconsin in MT assembly [41-43], and msps/ch-TOG is a
MT polymerase that associates with Patronin [29, 44]. Notably, Msp300 is the only LINC
complex protein that establishes the ncMTOC, while Klar and Koi are not required [29],
suggesting a novel and unconventional role for Msp300 at the fat body ncMTOC. Here, we have
further explored the complex biology of Msp300 isoforms and identified one particular and
distinct isoform of Msp300 that is responsible for organization of the perinuclear ncMTOC in the
fat body. By utilizing deletion mutants, transposable element insertion mutations, and RNAi, we
show that one unconventional isoform, Msp300-PE, appears necessary and sufficient for
establishment of the ncMTOC on the nuclear surface.

Results

The *Msp300* locus is complex

The *Msp300* gene is complex, encoding at least 11 isoforms that range in size from 47 to 1,500 kDa (Fig 1A). Based on the curated data on FlyBase, these variants arise from alternative promoters, mRNA splice sites, and transcriptional terminators [45]. The existence of the different isoforms is supported by RNA-seq data of exon-exon junctions and the isolation of partial cDNA clones for all isoforms [46-48]. In addition, tissue-specific expression data of the isoforms indicates that all isoforms are expressed in the fat body [49]. Typically, nesprins consist of many spectrin repeats (SRs), a pair of calponin homology (CH) domains, and a KASH domain [50]. From available expression data, isoforms B, E, and F lack the KASH domain, and isoforms E, F, G, and M lack CH domains. The KASH domain consists of a transmembrane helix and a short C-terminal region that resides in the nuclear intermembrane space [51]. The human giant nesprin isoforms, Nesprin-1 (1.0 MDa) and Nesprin-2 (0.8 MDa), contain 74 and 56 SRs, respectively [52, 53]. Similarly, one of the largest Msp300 isoforms, Msp300-PD (1.41 MDa), contains 52 SRs, as well as 2 CH domains and a KASH domain (Fig 1B-C) [54]. Msp300-PG (1.5 MDa) contains no CH domains, 21 SRs, and a KASH domain. Msp300-PE (1.06 MDa) is an atypical nesprin with no CH domains, only 2 SRs, and no KASH domain (Fig 1D). Instead of a KASH domain, Msp300-PE is predicted to have a transmembrane domain near its C-terminus (Fig 1E), followed by a 35 aa sequence.

Msp300 isoforms E/G/M are expressed more highly in the fat body

We measured the mRNA expression levels of groups of Msp300 isoforms in different tissues, including whole body, brain, flight muscle, ovary and fat body, by reverse transcription quantitative PCR (RT-qPCR). We assessed the expression of three sets of isoform mRNAs: all isoforms, those expressed from the two most 5′ promoters, and the E/G/M group of isoforms (Fig 1A). These data show that *Msp300* is expressed in all tissue types tested (Fig 1F). In the fat body, E/G/M are more highly expressed than the 5′ isoforms. Of the E/G/M group of isoforms in the fat body, isoform G was expressed at the highest level (Fig 1G). In addition, the existence of isoform E is validated by the RT-qPCR expression data.

Knockdown of Msp300 isoforms E and G impair the ncMTOC

To further assess which protein isoforms of *Msp300* are expressed and localized to the nuclear envelope in fat body cells, we used RNAi to knock down expression of isoform groups and stained with a polyclonal Msp300 antibody that targets all isoforms (Fig 1A) to detect remaining isoforms at the nuclear envelope. We generated tissue mosaics of RNAi knockdown using the coinFLP system [55] (Fig 2). With coinFLP, we obtained mosaic tissues where RNAi knockdown cells were generated efficiently at a reliable ratio with control cells, affording an internally-controlled framework to assess RNAi phenotypes. The RNAi knockdown clones are differentiated from control cells by co-expression of nuclear-targeted mCherry (mCherry-nls) in the clones. As an additional control, wild-type (WT) clones were generated in the w^{1118} background (Fig 2A, F).

When all Msp300 isoforms were knocked down (with HMS00368 [56]), the perinuclear

112   Msp300 signal was significantly reduced compared to the control cells, as expected (Fig 2B, F).
113   Consequentially, the nuclei were mispositioned as previously shown for *Msp300* knockdown
114   due to the loss of microtubule organization at the nuclear surface [29]. In contrast, targeting of
115   the 5' isoforms (with GD9539 [57]) did not diminish Msp300 localization to the nuclear surface
116   (Fig 2C, F). Knockdown of isoforms E and G (with GD10406 [57]) results in nuclear

Figure 1. *Msp300* gene isoforms and genetic analysis features. (A) Schematic of the isoforms expressed from *Msp300*. Deletion mutants (red), transposable element insertion mutations (green), and RNAi (orange) are indicated. The isoforms and gene regions affected by these mutations or RNAi elements are depicted. The epitope recognized by the α-Msp300 antibody (blue) is shown. Binding sites of the primers for RT-qPCR are represented by magnified views. (B-E) Schematic representations of Msp300-PD, a 'conventional' Nesprin, and Msp300-PE. Nesprins and most Msp300 isoforms such as PD (B) characteristically contain calponin homology domains, numerous spectrin repeats, and a KASH domain associated with a transmembrane domain (C). The atypical Msp300-PE (D) lacks calponin homology domains and contains only two spectrin repeats. The majority of Msp300-PE is comprised of two types of tandem repeats. Isoform PE also contains a transmembrane domain but lacks a KASH domain (E). (F) Bar graph comparing the mRNA expression levels of isoform groups relative to all isoforms in different tissues from RT-qPCR data from indicated tissues. RNA extractions were performed in triplicate and each RT-qPCR reaction was run in triplicate. Expression levels were normalized to GPDH1 and α-Tubulin reference genes. (G) Expression levels of isoforms E, G, and M relative to the expression level of all three isoforms combined in the fat body. The data are depicted as the mean ± SD. Statistical significance was determined using a two-tailed Student's *t*-test with Welch's correction (*n* = 3). See S4 Data for source data.

mispositioning; however, the Msp300 signal at the nuclear surface was not significantly reduced (Fig 2D, F). The remaining Msp300 signal at the nuclear envelope is likely contributed by the other isoforms not targeted by RNAi. Meanwhile, knockdown of all KASH-containing isoforms (with GD10317 [57]), which includes G, does not significantly reduce perinuclear Msp300 (Fig 2E, F). The persistence of Msp300 signal at nuclei following knockdown of each set of transcripts is consistent with the RT-qPCR data showing that most Msp300 isoforms are expressed in the fat body. Importantly, the loss of nuclear positioning upon knockdown of the E/G isoforms, while retaining Msp300 signal from other isoforms, indicates that Msp300-PE and -PG are uniquely required for nuclear positioning, likely through the regulation of MT organization. The retention of Msp300 signal despite loss of nuclear positioning indicates that other isoforms cannot compensate and lack the function to organize the ncMTOC and that this role is unique to the Msp300-PE,-PG isoforms.

Msp300 isoform E is sufficient for nuclear positioning and microtubule organization, with probable contribution from Isoform G

To determine which isoforms are required to maintain nuclear positioning and MT organization, we examined phenotypes for a variety of mutant lines in addition to the RNAi lines used in Figure 2. We generated mosaic mutant clones using the FLP-FRT recombination system [58], in which control cells express nuclear RFP while mutant cells do not (Fig 3). First, we examined two deletion mutants that impact groups of isoforms: one that deletes the 5′ end of the gene (*Msp300$^{\Delta 5'}$*) [59], disrupting isoforms B, D, H, I, J, K, and L, and a second deletion mutant that deletes a segment encoding the KASH domain (*Msp300$^{\Delta KASH}$*) [60] (see Fig 1A). In both mutants, nuclear positioning and MT organization were unaffected, indicating that the 5′ isoforms and also the KASH domain are not required for generating the ncMTOC (Fig 3A-B, G-H). The lack of requirement for the KASH domain indicates that the LINC complex involved in generating the ncMTOC is atypical.

To probe the isoforms responsible for generating the ncMTOC further, we examined phenotypes for MiMIC transposable element insertion lines within *Msp300* exons that

Figure 2. Multiple Msp300 isoforms localize to the nuclear surface. Immunostaining of fat bodies for Msp300 in *w1118* (A) or following RNAi knockdown by the HMS00368 (B), GD9539 (C), GD10406 (D), or GD10317 (E) RNAi lines. Images shown are representative of the phenotype associated with each line. Mosaic knockdown was achieved using the coinFLP system. Clonal RNAi knockdown cells express nuclear mCherry (red). Nuclei were stained with DAPI and pseudo-colored green, actin (cell membrane) stained red, and Msp300 white. Isoforms targeted by each RNAi line are indicated. Clones characterized by nuclear mispositioning are denoted by a pink arrowhead, while clones with properly positioned nuclei are denoted by a yellow arrow. (F) Quantitation of perinuclear Msp300 localization. Data are represented as the mean ± SD. A one-way ANOVA with Dunnett's T3 test was used (24 ≤ *n* ≤ 30). Source data is found in S4 Data.

conceptually cause *Msp300* loss of function (see Fig 1A). Consistent with the 5′ deletion, the

MI11102 [61] insertion within a coding exon near the 5′ end did not result in any changes to

nuclear positioning or MT organization (Fig 3C, G-H), further supporting that all isoforms

encoded by exons in the 5′ half of the gene are not required for generation of the ncMTOC. In

contrast, disruption of isoforms E, G, and M by MI11557 [61] and MB00410 [62], or just E and G

by MI11862 [61], results in severe nuclear mispositioning and disruption of radial and

Figure 3. Msp300 isoforms E and G are required for nuclear positioning and microtubule organization.
Nuclear positioning and MT organization was assessed by immunostaining fat bodies containing mosaic clones of Δ5' (A), ΔKASH (B), MI11102 (C), MI11557 (D), MB00410 (E), and MI11862 (F) *Msp300* mutations. Representative images are shown. The FLP-FRT technique was used to generate mutant clones, where WT alleles are distinguished by the expression of nuclear RFP. The isoforms affected by each mutant allele are listed (red text). Normal nuclear positioning and MT organization is represented by a yellow arrow, while mispositioned nuclei with disrupted MT organization are indicated with a pink arrowhead. Disruption of isoforms essential to the ncMTOC results in the loss of nuclear centricity (G) and the reduction of circumferential MTs (H). Statistical significance was determined for nuclear centricity and circumferential MT signal using a one-way ANOVA with Dunnett's T3 test and a one sample Student's test with a theoretical mean of 1.000, respectively ($21 \leq n \leq 30$). Bars on the graph represent the mean ± SD of the data. Source data is present in S4 Data.

circumferential MTs (Fig 3D-F, G-H). These data show that these isoforms, or at least the coding regions downstream of these three transposon insertion lines, are necessary for proper function of the ncMTOC. In addition, the results show that disruption of isoforms E and G significantly impairs the fat body ncMTOC and that expression of isoform M is not sufficient to support the ncMTOC. We conclude from these mutant analyses that Msp300 isoforms E and/or G are required to generate the ncMTOC on the fat body nuclear surface.

In addition to targeting sets of isoforms with mutant lines, we also used RNAi to knock down sets of isoforms to complement and corroborate the mutant analysis described above, and also to further discern the isoforms required for the ncMTOC. We used the RNAi lines shown in Figures 1 and 2, and again generated mosaic knockdown with the coinFLP system, but in these experiments we assessed MT organization and nuclear positioning rather than Msp300 expression as done in Figure 2. First, we compared wild-type (w^{1118}) to the internal control cells (Fig 4A, F-G). The HMS00368 RNAi line disrupted all Msp300 isoforms and resulted in severe nuclear mispositioning and MT disruption as shown previously [29] (Fig 4B, F-G). Knockdown of the isoforms expressed from the 5' region of the gene did not affect nuclear positioning or MT organization (Fig 4C, F-G), consistent with the findings from mutants that affect the 5' isoforms as shown in Figure 3. When targeting isoforms E and G with GD10406 RNAi, however, we observed mispositioned nuclei and disorganized MTs (Fig 4D, F-G). This result further shows that isoforms E and/or G are required to organize MTs at the ncMTOC. However, knockdown of isoform G and other isoforms containing the KASH domain using the GD10317 RNAi line resulted in moderate nuclear mispositioning and microtubule disruption (Fig 4E, F-G) indicating that isoform G supports the ncMTOC, but that isoform E is sufficient. We further show that when all isoforms except E and F are knocked down by RNAi (by expressing GD9539 and GD10317 together), the ncMTOC remains intact, whereas knockdown of all isoforms except F (GD9539 + GD10406) resulted in severe disruption of the ncMTOC, indicating that Msp300-PF is not sufficient (Fig S1). Together, the results from the mutants and the RNAi lines show that, while isoform G plays a contributing role in MT organization, isoform E is sufficient to generate the perinuclear ncMTOC. We were unable to test whether isoform G is also sufficient in the absence of E.

Retrograde endocytic vesicle trafficking is dependent on Msp300 isoform E

In addition to nuclear positioning and MT organization, we assessed the effects of

Figure 4. The perinuclear ncMTOC is organized by isoforms E and G. Immunofluorescence microscopy images of WT (w^{1118}) (A), HMS00368 (B), GD9539 (C), GD10406 (D), and GD10317 (E) RNAi lines to assess nuclear positioning and MT organization. Images shown are representative images. Mosaic clones were generated by the coinFLP technique, where knockdown clones are marked with nuclear mCherry (red). RNAi knockdown cells with properly positioned nuclei and organized radial and circumferential MTs are indicated with yellow arrows, and affected nuclei and MTs are indicated with pink arrowheads. Knockdown with GD10317 shows a moderate phenotype characterized by reduced MTs in the absence of nuclear mispositioning (gold arrowhead). Quantitation of nuclear centricity (F) and circumferential MT signal (G). For both analyses, statistics were conducted using a one-way ANOVA with Dunnett's T3 test (n = 30). Bar graphs depict the mean ± SD. S4 Data contains the source data.

Msp300 isoform depletion on endosome trafficking. Prior work showed that Msp300 is required

for the ncMTOC to organize MTs that support retrograde dynein-dependent endocytic vesicle
trafficking [29]. In wild-type cells, Rab5 endosomal vesicles are at the highest density near the
plasma membrane and perinuclearly, and disruption of MTs or dynein results in a loss of Rab5
endosome trafficking to the nuclear periphery [29]. To test the involvement of distinct Msp300
isoforms in endocytic vesicle trafficking, we examined the distribution of the endosomal marker
GFP-Rab5 upon disruption of select isoform(s) by RNAi knockdown in the larval fat body using
SPARC-GAL4. In the w^{1118} control cells and cells expressing $Luciferase^{JF01355}$ (RNAi control), Rab5
vesicles comprise a perinuclear population forming a halo-like pattern (Fig 5A-B, H). Knockdown
of *αTub84B* (positive control) causes impaired retrograde trafficking, as seen by the loss of the
perinuclear Rab5 signal (Fig 5C, H). Similarly, the knockdown of all *Msp300* isoforms using
HMS00368 RNAi results in diminished Rab5 vesicles at the nuclear surface (Fig 5D, H).
Disruption of the 5' isoforms by the GD9539 RNAi line did not impact the Rab5 distribution
pattern, indicative of normal endosome trafficking (Fig 5E, H). In contrast, knockdown of
isoforms E and G with GD10406 RNAi resulted in reduced perinuclear GFP-Rab5 endosome
signal (Fig 5F, H). The knockdown of Isoform G and other KASH-containing isoforms via GD10317
RNAi did not appreciably reduce the perinuclear GFP-Rab5 signal (Fig 5G, H). Together, these
results further show that isoform E is sufficient for ncMTOC function by supporting endosomal
vesicle trafficking in the fat body.
208

Msp300 isoform E is required for the recruitment of the ncMTOC components shot and
Patronin

To further evaluate the role of Msp300 isoforms at the fat body ncMTOC, we examined
their involvement in the recruitment of other ncMTOC components. Msp300 recruits the
spectraplakin short stop (shot) and the microtubule minus-end stabilizer Patronin to the
ncMTOC [29, 30]. In the w^{1118} control, shot localizes to the nuclear surface, along MTs, and on
the plasma membrane (Fig 6A, F). When all *Msp300* isoforms were disrupted, shot no longer
localizes to the nuclear surface [29]. Consistently, we observed a decreased amount of shot at
the nuclear surface when we used HMS00368 RNAi to knock down all isoforms (Fig 6B, F). The
knockdown does not, however, affect the localization of shot at the plasma membrane.
Interestingly, knockdown of the 5' isoforms (with GD9539 RNAi) resulted in an increase in the
perinuclear shot signal (Fig 6C, F). Knockdown of the KASH isoforms that include Isoform G also
showed an increase in the perinuclear localization of shot (Fig 6E, F). Targeting of isoforms E and
G with GD10406, on the other hand, causes a loss of shot from the nuclear surface (Fig 6D, F).
These results indicate that Msp300 isoform E is sufficient for the recruitment of shot to the
ncMTOC, consistent with its role in establishing the perinuclear ncMTOC.

Finally, we investigated the contribution of the isoforms to the recruitment of the
CAMSAP ortholog Patronin to the ncMTOC. Patronin is a MT minus-end stabilizer and a major
regulator of MTs at the fat body perinuclear ncMTOC and complete *Msp300* knockdown results
in the loss of Patronin localization at the nuclear surface [29]. In w^{1118} control and
$Luciferase^{JF01355}$ control RNAi cells expressing Patronin-GFP, Patronin localizes to the nuclear
surface (Fig 7A-B, H), while in $Patronin^{HMS01547}$ RNAi cells perinuclear Patronin is significantly
reduced (Fig 7C, H). Disruption of all isoforms (with HMS00368 RNAi) results in a significant
decrease in the perinuclear localization of Patronin (Fig 7D, H). Knockdown of the 5' isoforms, in

contrast, does not affect the presence of Patronin at the nuclear surface (Fig 7E, H). Patronin

Figure 5. Msp300 isoforms E+G are required for retrograde trafficking of endocytic vesicles. Analysis of retrograde endosome trafficking in w^{1118} (A) cells, *Luciferase*^{JF01355} (B), and *alphaTub84B*^{JF01373} (C), as well as in cells expressing the HMS00368 (D), GD9539 (E), GD10406 (F), and GD10317 (G) RNAi lines. Representative images are shown here. RNAi was expressed throughout the entire fat body tissue with SPARC-GAL4. Images were captured using the same parameters to allow for direct comparison between samples. Insets show a magnified view of a single nucleus (green) and the perinuclear population of Rab5 endosomes (white). Control or unaffected cells exhibit a halo-like pattern of endocytic vesicles surrounding the nucleus (yellow dashed boxes), while knockdown of Msp300 isoforms essential for the MTOC causes loss of this signal (pink boxes). (H) Bar graph depicting perinuclear GFP-Rab5 signal. A one-way ANOVA with Dunnett's T3 test was used for statistical analysis (*n* = 30). The mean ± SD is shown in the graph. Source data is present in S4 Data.

236 perinuclear localization is diminished when Isoforms E and G are knocked down with GD10406

Figure 6. Shot localization to the ncMTOC requires isoform E. Fat bodies were stained for shot in mosaic tissues expressing w^{1118} (A), HMS00368 (B), GD9539 (C), GD10406 (D), or GD10317 (E). Images shown are representative. The coinFLP system was used to express the RNAi, marked with co-expression of nuclear mCherry (red). Recruitment of shot to the nuclear surface by Msp300 results in a perinuclear shot signal (yellow arrows), while the knockdown of necessary Msp300 isoforms causes the loss of this signal (pink arrowheads). (F) Quantitation of perinuclear shot signal. Statistical analysis was performed via a one-way ANOVA and Dunnett's T3 test ($n = 30$). Data are depicted as mean ± SD. S4 Data contains the source data.

RNAi but not when the 3′ KASH-containing isoforms are knocked down (Fig 7F-G, H), consistent with the role for isoform E in MT organization and shot recruitment to the nuclear surface. Together, these data further show that Msp300 isoform E is necessary and may be sufficient for MTOC generation through the recruitment of shot and Patronin.

Msp300-PE has two novel large tandem repeats and a KASH-like C-terminal domain

Most cell types do not generate an MTOC on the nuclear surface [63], and fat body cells appear to utilize a special isoform of Msp300, Msp300-PE to assemble it. To further understand

Figure 7. Patronin localization to the ncMTOC requires isoform E. Patronin localization in w^{1118} (A), *Luciferase^{JF01355}* (RNAi) (B), and *Patronin^{HMS01547}* (RNAi) (C) fat bodies and fat bodies expressing the *Msp300* HMS00368 (D), GD9539 (E), GD10406 (F), or GD10317 (G) RNAi lines. Representative images are shown. RNAi lines were expressed throughout the fat body with SPARC-GAL4. The same confocal microscope parameters were used to capture the images for each sample. Nuclei (DAPI, green) and Patronin-GFP (white). Patronin normally localizes perinuclearly (yellow dashed boxes). Knockdown of Msp300 isoforms essential for the ncMTOC results in loss of Patronin signal at the nuclear surface (pink boxes). (H) Perinuclear Patronin signal in each sample represented in the form of a bar graph. Data is presented as the mean ± SD. Statistics were conducted via a one-way ANOVA with Dunnett's T3 test (n = 30). Source data is present in S4 Data.

what makes Msp300-PE unique, we analyzed its sequence and predicted domain structure with

in silico analysis tools. Msp300-PE contained no other identified domains besides the SRs and

the transmembrane domain based on analysis via the PredictProtein tool [64]. Intriguingly,

Msp300-PE encodes two unique types of novel tandem repeat regions (Fig 8A-D), identified
using the XSTREAM algorithm [65]. The first type, "Type 1," consists of ~20 copies of a 108-
amino acid sequence that is highly similar among the tandem repeats. "Type 2" repeats consist
of 101 amino acids tandemly repeated ~5-7 times in three clusters (labeled 2A-2C in Figure 8)
with a total of 19 repeats. These repeat domains have no previously described function. Using
AlphaFold 3, we analyzed the predicted structures of the repeats [66]. Individually, each cluster
of repeats is predicted to form similar structures despite no sequence similarity, comprised of
beta strands organized in parallel and antiparallel, to form a folded-back sandwich structure
with disordered loops present throughout the protein (Figure 8E-H, and Supplemental Videos 1
and 2). AlphaFold 3 predicts a loose association between the repeats that might induce further
flattening of the structures (Fig 8I). In addition, the predicted structure of the C-terminus of
Msp300-PE is shown. Similar to the KASH domain of other Msp300 isoforms (Fig 8J), the C-
terminal domain (CTD) of Msp300-PE is predicted to contain a membrane-spanning α-helix and
a mostly disordered CTD predicted to reside within the intermembrane space (Fig 8K).

Figure 8. Msp300-PE contains two novel large tandem repeats and a KASH-like C-terminal domain. (A) Repeat 1 (lime green) consists of a 108-amino acid sequence repeated 20 times. (B-D) Repeat 2 (salmon) consists of a different 101-amino acid sequence repeated at least 19 times among 3 clusters. (E) Predicted structure of Repeat 1 and view of Repeat 1 rotated 90° about the *y*-axis. (F) Repeat 2A predicted structure before rotating 90° around the *y*-axis and after. (G) Predicted structure of Repeat 2B and view after rotating 90°C about the *y*-axis. (H) Repeat 2C predicted structure and view of Repeat 2C following 90° rotation around the *y*-axis. (I) Predicted structure of Repeat 1 (lime green) and Repeat 2A-2C (salmon) before and after 90° rotation about the *y*-axis. (J) Predicted structure of the CTD containg the KASH domain (blue), transmembrane domain (green), and linker region (lavender) and view after 90° rotation around the *y*-axis. (K) Predicted structure of the CTD of Msp300-PE with the transmembrane domain (green) and linker region (lavender) shown before 90° rotation around the *y*-axis and after.

Discussion

Here we show that a unique isoform of Msp300, Msp300-PE, is necessary in fat body cells to organize the ncMTOC on the nuclear surface. Msp300-PE and the isoforms expressed from the same transcriptional promotor (Msp300-PG and -PM) are expressed most highly in the fat body compared to the expression of the other isoforms, indicating that these isoforms and the unique protein domains they encode may be responsible for assembly of the ncMTOC on the nuclear surface. Our data also indicate that Msp300-PE is sufficient to generate the ncMTOC on the nuclear surface. Msp300-PE organizes the microtubules through recruitment of shot and Patronin, a major regulator of microtubules at the perinuclear ncMTOC [29] and is consequently required for nuclear positioning and endocytic vesicle trafficking in fat body cells. These findings establish why Msp300, expressed from a complex locus with expression in many cell types, organizes an ncMTOC on the surface of nuclei in a cell-type specific manner. Msp300-PE has a unique domain structure, containing a predicted novel outer nuclear membrane targeting domain and with novel repeat domains which, combined, indicate that Msp300-PE is a novel alternative LINC complex component.

The knockdown of Msp300-PE/PG, or their disruption together with Msp300-PM with transposon insertions, indicates that Msp300-PE and -PG organize the perinuclear MTOC in fat bodies. The loss of Msp300-PG, together with all KASH-containing isoforms, altered nuclear positioning but did not significantly block the function of the MTOC, as MTs and the localization of Rab5-GFP were not significantly affected. Thus, we infer from all the results that Msp300-PE is sufficient to organize the MTOC, while Msp300-PG contributes. It is possible that Msp300-PG is also sufficient, but we did not have the genetic tools to test that in this study. Due to the technical limitations of expressing such large constructs, we were unable to test sufficiency by expressing a transgene of Msp300-PE/PG in a null background; the Msp300-PE coding sequence is almost 29,000 bp in length and Msp300-PG is longer (over 40kbp). Furthermore, while the existence of PE/PG is supported by the available cDNAs corresponding to the exonic regions common to both isoforms, further validation is required to confirm the existence of other isoforms such as PF [48].

In contrast with the majority of the other Msp300 isoforms, and nesprins generally across species, isoform E lacks the characteristic KASH domain, the CH domains, and the high number of spectrin repeats. Also, we show that loss of all KASH-containing isoforms does not affect the ncMTOC in the fat body or the localization of Msp300 to the nuclear surface. However, KASH-less isoforms have been identified for other nesprins in Drosophila and other species including *Drosophila* klar, vertebrate nesprin-1/2, and *C. elegans* ANC-1. A KASH-less isoform of klar was shown to be associated with lipid droplets [67, 68]. In addition, KASH-less splice variants of nesprin-1 and nesprin-2 have been characterized [69-71], and an isoform of nesprin-1 lacking the KASH domain is proposed to be expressed in the embryonic mouse brain [72]. A nesprin-1-giant isoform lacking the KASH domain has been identified in the CNS and is associated with spinocerebellar ataxia [73, 74]. Moreover, In *C. elegans*, ANC-1 regulates nuclear positioning in hypodermis syncytia independently of the KASH domain and the LINC complex [75]. Furthermore, while in *Drosophila* loss of the 5′ isoforms was lethal at the first instar larval stage, deletion of the KASH domain from all isoforms was viable [59]. In Drosophila

muscle, however, Msp300 is required for nuclear positioning in a KASH-dependent manner [21].

Despite the similarities with other KASH-less isoforms, isoform E is particularly unique
due to the presence of the two types of highly conserved ~100 amino acid-long tandem repeats.
Based on the results of a protein BLAST against all insects, the repeats appear to possess
significant homology only among Drosophila species. In addition, the repeats are characterized
by an overall negative charge, as observed in intrinsically disordered regions of proteins [76, 77].
The negatively charged regions may also facilitate binding with other proteins. For example, the
positively charged regions of Patronin, which bind to the negatively charged regions of MTs, may
also interact with the repeats of Msp300. The unique domain structure of Msp300-PE is likely
linked to the mechanism by which this variant nesprin organizes the ncMTOC. Furthermore, the
structural similarity of the C-terminal domain of Msp300-PE with the KASH domain of other
Msp300 isoforms may provide a binding site for a previously unidentified inner nuclear
membrane protein partner of Msp300-PE. Together, Msp300-PE and the predicted inner nuclear
membrane protein may form a novel LINC complex.

Overall, this work shows that Msp300-PE has unique function to generate a perinuclear
ncMTOC. Future work will elucidate the functions of the unique domains and how Msp300-PE is
able to generate an ncMTOC on the surface of nuclei.

Materials and Methods

Genetic data

Information pertaining to the gene sequence and isoforms was obtained from the data available on FlyBase (https://flybase.org/reports/FBgn0261836.htm) [45].

Drosophila stocks

Information for each *Drosophila* stock is contained in Supplementary Table 1. Flies were raised using standard food made of cornmeal, molasses, and yeast and were maintained at 25°C.

Generation and expression of FLP-FRT and coinFLP stocks

The generation of FLP-FRT mutant clones was conducted by crossing y^1 w^* $hsFLP^{22}$; Ubi-RFP-nls neoFRT40A females to males with each *Msp300* allele recombined with neoFRT40A. For each cross, 30-40 females were used.

The generation of coinFLP mosaic RNAi knockdowns was conducted using an adaptation of the coinFLP system [55]. For each cross, 25-35 females of the genotype w^* $hsFLP^1$ coinFLP-GAL4^{attP3};; UAS-mCherry-nls3 were crossed to RNAi line males.

Crosses were established for at least 2-3 days before proceeding with heat shock to induce recombination. FLP-FRT and coinFLP crossed flies were transferred to fresh food vials for 1-2 hours, then to fresh food vials to lay eggs for 2-3 h, and then adults were removed and embryos were aged past the cleavage stage at 29°C for 2 hr. Embryos were then heat-shocked in a 37°C water bath for 30 minutes and then returned to 29°C for coinFLP crosses or 25°C for FLP-FRT crosses. Wandering stage larvae were collected 4-5 days later for fat body dissection.

Sample preparation and immunostaining

Fat bodies from late third-instar wandering larvae were dissected in 1× Dulbecco's phosphate-buffered saline (D-PBS) (GIBCO, Waltham, MA, USA) as previously described [29]. Fat bodies from two larvae were used for each slide. Following dissection, samples were transferred to a poly-L-lysine-coated slide. For Patronin-GFP samples, the fat bodies were fixed in 15 µL of 100% methanol under a siliconized coverslip for 10 min at -20°C. The coverslip was removed and the slides were placed in 1× PBS (137 mM NaCl, 2.7 mM KCl, 8.0 mM Na_2HPO_4, and 1.4 mM KH_2PO_4). All other samples were fixed in 15 µL of 4% paraformaldehyde (PFA) for 8 min before a siliconized coverslip was placed on the slide. The sample was gently flattened and then flash frozen in liquid nitrogen. While still frozen, the coverslip was removed and the slides placed in 1× PBS. For all samples, a ~2 cm diameter hydrophobic ring was drawn around the sample with a Super Pap Pen (Electron Microscopy Sciences, Cat # 71312). Fat bodies were permeabilized, blocked, and stained in 100 µL of staining solution (1× PBS, 1% BSA, 0.1% saponin) for 2 hr at RT or overnight at 4°C using the antibodies and dyes listed in Supplementary Table 2. Slides were extensively washed in 1× PBS following staining. Samples were mounted in 10 µL of mounting medium (80% glycerol, 0.1 M Tris pH 8.8, 0.05% p-phenylenediamine dihydrochloride (Sigma, cat. no. P1519)) and stored at -20°C.

Image acquisition

Fat bodies were imaged with either a Nikon A1 or Nikon AX laser scanning confocal microscope (Nikon, Japan). Specimens were imaged using a 60× oil-immersion objective with a 1.49 numerical aperture (NA). Images were captured with the Nikon NIS-Elements AR software (version 4.6 or 5.42 with Nikon A1 or Nikon AX, respectively) with 0.5 µm between z-sections. All images are presented as maximum intensity projections of the z-stack.

Quantification analysis

Microscopy images were quantified using Fiji [78]. Perinuclear localization was calculated from a single z-slice, while nuclear mispositioning was calculated from a maximum intensity projection. Quantification of perinuclear localization was performed by creating multiple ROIs of the same size throughout the perinuclear region and averaging the total fluorescence intensity of each ROI. The background for each cell was accounted for by subtracting the value of an ROI in the cytoplasm from the value of the perinuclear region. For samples utilizing the coinFLP system, the value for each clone was depicted as ratio of the clone to an adjacent wild-type control cell. For all other samples, the average value of the clones from a slide was divided by the average value of the corresponding w^{1118} control slide. Quantification of nuclear positioning was conducted as previously described [29]. In brief, the coordinates of the cell or nuclear centroid were obtained by defining the boundary of the plasma membrane or nucleus, respectively. Next, the distance between the two coordinates was calculated.

RT-qPCR

For preparation of whole fly and flight muscle RNA, 10 adult flies were used. Ovary RNA was extracted from 10 adult virgin female flies. Brains and fat bodies were dissected from 10 late third-instar wandering larvae. Tissues were washed in 1× D-PBS following dissection. After homogenizing sample via pellet pestle and aspiration with a 20 G needle, RNA was isolated from all samples using TRIzol (Ambion, Austin, TX, USA) following the recommended protocol from the manufacturer. For all extraction steps RNase-free/DNase-free plasticware and solutions were used. Isolated RNA samples were resuspended in UltraPure DNase/RNase-free distilled water and treated with TURBO DNA-*free* kit (Thermo Fisher, Waltham, MA, USA). cDNA was produced using the SuperScript IV (Invitrogen, Waltham, MA, USA) kit with random hexamers primers. Quantitative PCR was performed using the primer pairs shown in Figure 1A along with those for the *αTub84B [79]* and *Gpdh1* reference genes. Primer sequences are listed in Supplementary Table 3. Amplification and quantification were performed using the Brilliant III Ultra-Fast qPCR Master Mix (Agilent, Santa Clara, CA, USA) at a Bio-Rad CFX384 real-time analyzer. Quantification was performed using the $2^{-\Delta\Delta Ct}$ method. The expression levels of the 5′ and E, G, M isoforms were compared to the average expression levels of All isoforms amplified from two separate primer pairs.

Statistics and reproducibility

For microscopy image calculations, three slides, with each containing the fat bodies from two larvae, were imaged. On each slide, $\geqslant 5$ wild-type and $\geqslant 5$ clone cells were included in the quantitation. The use of the mosaic systems in this study provided an internal control, which

reduces variability arising from sample preparation. For non-mosaic tissues, 10 cells from each
slide were quantified. Nuclear positioning and perinuclear localization values are expressed as
the mean ± standard deviation (SD). All samples for RT-qPCR were dissected in triplicate and
each reaction was performed in triplicate. The data were significant based on a *P*-value < 0.05.
For datasets with negative control samples, a one-way ANOVA was performed, while a one
sample *t*-test was used for all other datasets. GraphPad Prism 10 (Dotmatics, Boston, MA, USA)
was used for all analyses.
413
In silico analysis

Analysis of Msp300-PE was performed using the publicly available amino acid sequence
(NCBI Reference Sequence: NP_001188693.1). PredictProtein [64] was used to identify the
residues predicted to comprise the transmembrane domain of Msp300-PE. The locations of the
tandem repeats were determined via XSTREAM [65]. Structure predictions of the repeat regions
and C-termini of PD and PE were performed with AlphaFold 3 [66]. Predicted structures were
visualized using PyMOL (The PyMOL Molecular Graphics System, Version 2.4.0 Schrödinger, LLC).
All analyses were conducted using the default parameters of the tools.

Graphics

All graphics were made using Adobe Illustrator (version 25.3.1). Also, all figures were
composed using Adobe Illustrator.

Acknowledgements

We thank Véronique Morel, Bénédicte Durand, Talila Volk, Michael Welte, and the Bloomington *Drosophila* Stock Center for *Drosophila* stocks; Talila Volk and the Developmental Studies Hybridoma Bank for antibodies; and Yue Julia Wang for assistance with RT-qPCR. We are grateful to Bloomington Drosophila Stock Center (NIH P40OD018537), Vienna Drosophila Resource Center [57], and FlyBase [45] for curating indispensable tools and resources. Thank you to Batory Foods for their generous donation of fly food reagents to support this work.

This work was supported by NIH grant R01GM139971 to T. Megraw, and National Natural Science Foundation of China grant (32370731) and Shenzhen Science and Technology Program grant (JCYJ20230807091308018) to Y. Zheng.

References

1.   Crisp M, Liu Q, Roux K, Rattner JB, Shanahan C, Burke B, et al. Coupling of the nucleus and cytoplasm. The Journal of Cell Biology. 2006;172(1):41-53. doi: 10.1083/jcb.200509124.

2.   Starr DA, Fridolfsson HN. Interactions between nuclei and the cytoskeleton are mediated by SUN-KASH nuclear-envelope bridges. Annu Rev Cell Dev Biol. 2010;26:421-44. doi: 10.1146/annurev-cellbio-100109-104037. PubMed PMID: 20507227; PubMed Central PMCID: PMCPMC4053175.

3.   Horn HF, Brownstein Z, Lenz DR, Shivatzki S, Dror AA, Dagan-Rosenfeld O, et al. The LINC complex is essential for hearing. J Clin Invest. 2013;123(2):740-50. Epub 20130125. doi: 10.1172/JCI66911. PubMed PMID: 23348741; PubMed Central PMCID: PMCPMC3561815.

4.   Mejat A, Misteli T. LINC complexes in health and disease. Nucleus. 2010;1(1):40-52. Epub 2011/02/18. doi: 10.4161/nucl.1.1.10530. PubMed PMID: 21327104; PubMed Central PMCID: PMCPMC3035119.

5.   Storey EC, Fuller HR. Genotype-Phenotype Correlations in Human Diseases Caused by Mutations of LINC Complex-Associated Genes: A Systematic Review and Meta-Summary. Cells. 2022;11(24):4065. Epub 20221215. doi: 10.3390/cells11244065. PubMed PMID: 36552829; PubMed Central PMCID: PMCPMC9777268.

6.   Zhang Q, Bethmann C, Worth NF, Davies JD, Wasner C, Feuer A, et al. Nesprin-1 and -2 are involved in the pathogenesis of Emery Dreifuss muscular dystrophy and are critical for nuclear envelope integrity. Hum Mol Genet. 2007;16(23):2816-33. Epub 20070829. doi: 10.1093/hmg/ddm238. PubMed PMID: 17761684.

7.   Puckelwartz MJ, Kessler EJ, Kim G, Dewitt MM, Zhang Y, Earley JU, et al. Nesprin-1 mutations in human and murine cardiomyopathy. J Mol Cell Cardiol. 2010;48(4):600-8. Epub 20091124. doi: 10.1016/j.yjmcc.2009.11.006. PubMed PMID: 19944109; PubMed Central PMCID: PMCPMC2837775.

8.   Stroud MJ. Linker of nucleoskeleton and cytoskeleton complex proteins in cardiomyopathy. Biophys Rev. 2018;10(4):1033-51. Epub 20180604. doi: 10.1007/s12551-018-0431-6. PubMed PMID: 29869195; PubMed Central PMCID: PMCPMC6082319.

9.   Gros-Louis F, Dupre N, Dion P, Fox MA, Laurent S, Verreault S, et al. Mutations in SYNE1 lead to a newly discovered form of autosomal recessive cerebellar ataxia. Nat Genet. 2007;39(1):80-5. Epub 20061210. doi: 10.1038/ng1927. PubMed PMID: 17159980.

10.   Wang JY, Yu IS, Huang CC, Chen CY, Wang WP, Lin SW, et al. Sun1 deficiency leads to cerebellar ataxia in mice. Dis Model Mech. 2015;8(8):957-67. Epub 20150505. doi: 10.1242/dmm.019240. PubMed PMID: 26035387; PubMed Central PMCID: PMCPMC4527285.

11.   Attali R, Warwar N, Israel A, Gurt I, McNally E, Puckelwartz M, et al. Mutation of SYNE-1, encoding an essential component of the nuclear lamina, is responsible for autosomal recessive arthrogryposis. Human Molecular Genetics. 2009;18(18):3462-9. doi: 10.1093/hmg/ddp290.

12.   Lawrence KS, Tapley EC, Cruz VE, Li Q, Aung K, Hart KC, et al. LINC complexes promote homologous recombination in part through inhibition of nonhomologous end joining. J Cell Biol. 2016;215(6):801-21. Epub 20161212. doi: 10.1083/jcb.201604112. PubMed PMID: 27956467; PubMed Central PMCID: PMCPMC5166498.

13.   Warren DT, Tajsic T, Porter LJ, Minaisah RM, Cobb A, Jacob A, et al. Nesprin-2-dependent ERK1/2 compartmentalisation regulates the DNA damage response in vascular smooth muscle cell ageing. Cell Death Differ. 2015;22(9):1540-50. Epub 20150306. doi: 10.1038/cdd.2015.12. PubMed PMID: 25744025; PubMed Central PMCID: PMCPMC4532777.

14.   Burke B. LINC complexes as regulators of meiosis. Current Opinion in Cell Biology. 2018;52:22-9. doi: 10.1016/j.ceb.2018.01.005.

15.   Koszul R, Kim KP, Prentiss M, Kleckner N, Kameoka S. Meiotic chromosomes move by linkage to
dynamic actin cables with transduction of force through the nuclear envelope. Cell. 2008;133(7):1188-
201. doi: 10.1016/j.cell.2008.04.050. PubMed PMID: 18585353; PubMed Central PMCID:
PMCPMC2601696.
16.   Lee YL, Burke B. LINC complexes and nuclear positioning. Semin Cell Dev Biol. 2018;82:67-76.
Epub 2017/12/02. doi: 10.1016/j.semcdb.2017.11.008. PubMed PMID: 29191370.
17.   Rothballer A, Schwartz TU, Kutay U. LINCing complex functions at the nuclear envelope: what the
molecular architecture of the LINC complex can reveal about its function. Nucleus. 2013;4(1):29-36.
Epub 2013/01/18. doi: 10.4161/nucl.23387. PubMed PMID: 23324460; PubMed Central PMCID:
PMCPmc3585024.
18.   Zhang J, Felder A, Liu Y, Guo LT, Lange S, Dalton ND, et al. Nesprin 1 is critical for nuclear
positioning and anchorage. Hum Mol Genet. 2010;19(2):329-41. Epub 20091028. doi:
10.1093/hmg/ddp499. PubMed PMID: 19864491; PubMed Central PMCID: PMCPMC2796894.
19.   Collins MA, Coon LA, Thomas R, Mandigo TR, Wynn E, Folker ES. Ensconsin-dependent changes
in microtubule organization and LINC complex-dependent changes in nucleus-nucleus interactions result
in quantitatively distinct myonuclear positioning defects. Mol Biol Cell. 2021;32(21):ar27. Epub
20210915. doi: 10.1091/mbc.E21-06-0324. PubMed PMID: 34524872; PubMed Central PMCID:
PMCPMC8693964.
20.   Padmakumar VC, Libotte T, Lu W, Zaim H, Abraham S, Noegel AA, et al. The inner nuclear
membrane protein Sun1 mediates the anchorage of Nesprin-2 to the nuclear envelope. Journal of Cell
Science. 2005/08/01;118(15). doi: 10.1242/jcs.02471.
21.   Elhanany-Tamir H, Yu YV, Shnayder M, Jain A, Welte M, Volk T. Organelle positioning in muscles
requires cooperation between two KASH proteins and microtubules. The Journal of Cell Biology.
2012;198(5):833-46. doi: 10.1083/jcb.201204102.
22.   Grady RM, Starr DA, Ackerman GL, Sanes JR, Han M. Syne proteins anchor muscle nuclei at the
neuromuscular junction. Proc Natl Acad Sci U S A. 2005;102(12):4359-64. Epub 20050304. doi:
10.1073/pnas.0500711102. PubMed PMID: 15749817; PubMed Central PMCID: PMCPMC555524.
23.   Puckelwartz MJ, Kessler E, Zhang Y, Hodzic D, Randles KN, Morris G, et al. Disruption of nesprin-1
produces an Emery Dreifuss muscular dystrophy-like phenotype in mice. Hum Mol Genet.
2009;18(4):607-20. Epub 20081113. doi: 10.1093/hmg/ddn386. PubMed PMID: 19008300; PubMed
Central PMCID: PMCPMC2722216.
24.   Zhang X, Xu R, Zhu B, Yang X, Ding X, Duan S, et al. Syne-1 and Syne-2 play crucial roles in
myonuclear anchorage and motor neuron innervation. Development. 2007;134(5):901-8. Epub
20070131. doi: 10.1242/dev.02783. PubMed PMID: 17267447.
25.   Volk T. A new member of the spectrin superfamily may participate in the formation of embryonic
muscle attachments in Drosophila. Development. 1992;116(3):721-30. doi: 10.1242/dev.116.3.721.
PubMed PMID: 1289062.
26.   Wang S, Reuveny A, Volk T. Nesprin provides elastic properties to muscle nuclei by cooperating
with spectraplakin and EB1. J Cell Biol. 2015;209(4):529-38. doi: 10.1083/jcb.201408098. PubMed PMID:
26008743; PubMed Central PMCID: PMCPMC4442817.
27.   Folker ES, Baylies MK. Nuclear positioning in muscle development and disease. Front Physiol.
2013;4:363. Epub 20131212. doi: 10.3389/fphys.2013.00363. PubMed PMID: 24376424; PubMed
Central PMCID: PMCPMC3859928.
28.   Rey A, Schaeffer L, Durand B, Morel V. Drosophila Nesprin-1 Isoforms Differentially Contribute to
Muscle Function. Cells. 2021;10(11):3061. Epub 20211106. doi: 10.3390/cells10113061. PubMed PMID:
34831284; PubMed Central PMCID: PMCPMC8616381.
29.   Zheng Y, Buchwalter RA, Zheng C, Wight EM, Chen JV, Megraw TL. A perinuclear microtubule-
organizing centre controls nuclear positioning and basement membrane secretion. Nat Cell Biol.

2020;22(3):297-309. Epub 20200217. doi: 10.1038/s41556-020-0470-7. PubMed PMID: 32066907;
PubMed Central PMCID: PMCPMC7161059.
30.   Sun T, Song Y, Dai J, Mao D, Ma M, Ni JQ, et al. Spectraplakin Shot Maintains Perinuclear
Microtubule Organization in Drosophila Polyploid Cells. Dev Cell. 2019;49(5):731-47 e7. Epub 20190418.
doi: 10.1016/j.devcel.2019.03.027. PubMed PMID: 31006649.
31.   Goodwin SS, Vale RD. Patronin regulates the microtubule network by protecting microtubule
minus ends. Cell. 2010;143(2):263-74. Epub 2010/10/16. doi: 10.1016/j.cell.2010.09.022. PubMed PMID:
20946984; PubMed Central PMCID: PMCPMC3008421.
32.   Hendershott MC, Vale RD. Regulation of microtubule minus-end dynamics by CAMSAPs and
Patronin. Proc Natl Acad Sci U S A. 2014;111(16):5860-5. Epub 2014/04/08. doi:
10.1073/pnas.1404133111. PubMed PMID: 24706919; PubMed Central PMCID: PMCPMC4000804.
33.   Jiang K, Hua S, Mohan R, Grigoriev I, Yau KW, Liu Q, et al. Microtubule minus-end stabilization by
polymerization-driven CAMSAP deposition. Dev Cell. 2014;28(3):295-309. Epub 2014/02/04. doi:
10.1016/j.devcel.2014.01.001. PubMed PMID: 24486153.
34.   Akhmanova A, Hoogenraad CC. Microtubule minus-end-targeting proteins. Curr Biol.
2015;25(4):R162-71. doi: 10.1016/j.cub.2014.12.027. PubMed PMID: 25689915.
35.   Tillery MML, Blake-Hedges C, Zheng Y, Buchwalter RA, Megraw TL. Centrosomal and Non-
Centrosomal Microtubule-Organizing Centers (MTOCs) in Drosophila melanogaster. Cells. 2018;7(9):121.
Epub 20180828. doi: 10.3390/cells7090121. PubMed PMID: 30154378; PubMed Central PMCID:
PMCPMC6162459.
36.   Sanchez AD, Feldman JL. Microtubule-organizing centers: from the centrosome to non-
centrosomal sites. Curr Opin Cell Biol. 2017;44:93-101. Epub 2016/09/27. doi:
10.1016/j.ceb.2016.09.003. PubMed PMID: 27666167; PubMed Central PMCID: PMCPMC5362366.
37.   Sallee MD, Feldman JL. Microtubule organization across cell types and states. Curr Biol.
2021;31(10):R506-r11. doi: 10.1016/j.cub.2021.01.042. PubMed PMID: 34033781.
38.   Casenghi M, Barr FA, Nigg EA. Phosphorylation of Nlp by Plk1 negatively regulates its dynein-
dynactin-dependent targeting to the centrosome. J Cell Sci. 2005;118(Pt 21):5101-8. Epub 2005/10/29.
doi: 118/21/5101 [pii]

10.1242/jcs.02622. PubMed PMID: 16254247.
39.   Celestino R, Henen MA, Gama JB, Carvalho C, McCabe M, Barbosa DJ, et al. A transient helix in
the disordered region of dynein light intermediate chain links the motor to structurally diverse adaptors
for cargo transport. PLoS biology. 2019;17(1):e3000100. Epub 2019/01/08. doi:
10.1371/journal.pbio.3000100. PubMed PMID: 30615611; PubMed Central PMCID: PMCPMC6336354.
40.   Redwine WB, DeSantis ME, Hollyer I, Htet ZM, Tran PT, Swanson SK, et al. The human
cytoplasmic dynein interactome reveals novel activators of motility. Elife. 2017;6. Epub 2017/07/19. doi:
10.7554/eLife.28257. PubMed PMID: 28718761; PubMed Central PMCID: PMCPMC5533585.
41.   Zheng Y, Mennella V, Marks S, Wildonger J, Elnagdi E, Agard D, et al. The Seckel syndrome and
centrosomal protein Ninein localizes asymmetrically to stem cell centrosomes but is not required for
normal development, behavior, or DNA damage response in Drosophila. Mol Biol Cell. 2016;27(11):1740-
52. Epub 2016/04/08. doi: 10.1091/mbc.E15-09-0655. PubMed PMID: 27053665; PubMed Central
PMCID: PMCPMC4884065.
42.   Tillery MML, Zheng C, Zheng Y, Megraw TL. Ninein domains required for its localization,
association with partners dynein and ensconsin, and microtubule organization. Mol Biol Cell.
2024;35(9):ar116. Epub 20240718. doi: 10.1091/mbc.E23-06-0245. PubMed PMID: 39024292.
43.   Rosen JN, Azevedo M, Soffar DB, Boyko VP, Brendel MB, Schulman VK, et al. The Drosophila
Ninein homologue Bsg25D cooperates with Ensconsin in myonuclear positioning. J Cell Biol. 2019. Epub
2019/01/11. doi: 10.1083/jcb.201808176. PubMed PMID: 30626718.

44.   Brouhard GJ, Stear JH, Noetzel TL, Al-Bassam J, Kinoshita K, Harrison SC, et al. XMAP215 is a
processive microtubule polymerase. Cell. 2008;132(1):79-88. Epub 2008/01/15. doi: S0092-
8674(07)01547-4 [pii]

10.1016/j.cell.2007.11.043. PubMed PMID: 18191222.
45.   Öztürk-Çolak A, Marygold SJ, Antonazzo G, Attrill H, Goutte-Gattat D, Jenkins VK, et al. FlyBase:
updates to the Drosophila genes and genomes database. Genetics. 2024/05/07;227(1). doi:
10.1093/genetics/iyad211.
46.   Daines B, Wang H, Wang L, Li Y, Han Y, Emmert D, et al. The Drosophila melanogaster
transcriptome by paired-end RNA sequencing. Genome Research. 2011 Feb;21(2). doi:
10.1101/gr.107854.110.
47.   Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, et al. The Developmental
Transcriptome of Drosophila melanogaster. Nature. 2010 Dec 22;471(7339). doi: 10.1038/nature09715.
48.   Rubin GM, Hong L, Brokstein P, Evans-Holm M, Frise E, Stapleton M, et al. A Drosophila
Complementary DNA Resource. Science. 2000-3-24;287(5461). doi: 10.1126/science.287.5461.2222.
49.   Krause SA, Overend G, Dow JAT, Leader DP. FlyAtlas 2 in 2022: enhancements to the Drosophila
melanogaster expression atlas. Nucleic Acids Research. 2021 Oct 29;50(D1). doi: 10.1093/nar/gkab971.
50.   Liem RK. Cytoskeletal Integrators: The Spectrin Superfamily. Cold Spring Harb Perspect Biol.
2016;8(10):a018259. Epub 20161003. doi: 10.1101/cshperspect.a018259. PubMed PMID: 27698030;
PubMed Central PMCID: PMCPMC5046693.
51.   Starr DA, Fischer JA. KASH 'n Karry: the KASH domain family of cargo-specific cytoskeletal
adaptor proteins. Bioessays. 2005;27(11):1136-46. doi: 10.1002/bies.20312. PubMed PMID: 16237665.
52.   Autore F, Pfuhl M, Quan X, Williams A, Roberts RG, Shanahan CM, et al. Large-Scale Modelling of
the Divergent Spectrin Repeats in Nesprins: Giant Modular Proteins. PLOS ONE. 2013;8(5):e63633. doi:
10.1371/journal.pone.0063633.
53.   Simpson JG, Roberts RG. Patterns of evolutionary conservation in the nesprin genes highlight
probable functionally important protein domains and isoforms. Biochem Soc Trans. 2008;36(Pt 6):1359-
67. doi: 10.1042/BST0361359. PubMed PMID: 19021556.
54.   Morel V, Lepicard S, Rey AN, Parmentier ML, Schaeffer L. Drosophila Nesprin-1 controls
glutamate receptor density at neuromuscular junctions. Cell Mol Life Sci. 2014;71(17):3363-79. Epub
20140204. doi: 10.1007/s00018-014-1566-7. PubMed PMID: 24492984.
55.   Bosch JA, Tran NH, Hariharan IK. CoinFLP: a system for efficient mosaic screening and for
visualizing clonal boundaries in *Drosophila*. Development. 2015;142(3):597-606. doi:
10.1242/dev.114603.
56.   Ni J-Q, Zhou R, Czech B, Liu L-P, Holderbaum L, Yang-Zhou D, et al. A genome-scale shRNA
resource for transgenic RNAi in Drosophila. Nature Methods. 2011;8(5):405-7. doi: 10.1038/nmeth.1592.
57.   Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, et al. A genome-wide transgenic
RNAi library for conditional gene inactivation in Drosophila. Nature. 2007;448(7150):151-6. doi:
10.1038/nature05954.
58.   Xu T, Rubin GM. Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development. 1993;117(4):1223-37. doi: 10.1242/dev.117.4.1223.
59.   Technau M, Roth S. The Drosophila KASH domain proteins Msp-300 and Klarsicht and the SUN
domain protein Klaroid have no essential function during oogenesis. Fly (Austin). 2008;2(2):82-91. Epub
20080301. doi: 10.4161/fly.6288. PubMed PMID: 18820478.
60.   Xie X, Fischer JA. On the roles of the Drosophila KASH domain proteins Msp-300 and Klarsicht.
Fly (Austin). 2008;2(2):74-81. Epub 2008/09/30. PubMed PMID: 18820482.

61.   Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, et al. MiMIC: a highly
versatile transposon insertion resource for engineering Drosophila melanogaster genes. Nature
Methods. 2011;8(9):737-43. doi: 10.1038/nmeth.1662.
62.   Bellen HJ, Levis RW, He Y, Carlson JW, Evans-Holm M, Bae E, et al. The <i>Drosophila</i> Gene
Disruption Project: Progress Using Transposons With Distinctive Site Specificities. Genetics.
2011;188(3):731-43. doi: 10.1534/genetics.111.126995.
63.   Tillery MML, Blake-Hedges C, Zheng Y, Buchwalter RA, Megraw TL. Centrosomal and Non-
Centrosomal Microtubule-Organizing Centers (MTOCs) in Drosophila melanogaster. Cells. 2018/09;7(9).
doi: 10.3390/cells7090121.
64.   Bernhofer M, Dallago C, Karl T, Satagopam V, Heinzinger M, Littmann M, et al. PredictProtein -
Predicting Protein Structure and Function for 29 Years. Nucleic Acids Research. 2021/07/02;49(W1). doi:
10.1093/nar/gkab354.
65.   Newman AM, Cooper JB. XSTREAM: a practical algorithm for identification and architecture
modeling of tandem repeats in protein sequences. BMC Bioinformatics. 2007;8(1):382. Epub 20071011.
doi: 10.1186/1471-2105-8-382. PubMed PMID: 17931424; PubMed Central PMCID: PMCPMC2233649.
66.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature 2024 630:8016. 2024-05-08;630(8016). doi:
10.1038/s41586-024-07487-w.
67.   Guo Y, Jangi S, Welte MA. Organelle-specific control of intracellular transport: distinctly targeted
isoforms of the regulator Klar. Mol Biol Cell. 2005;16(3):1406-16. Epub 20050112. doi: 10.1091/mbc.e04-
10-0920. PubMed PMID: 15647372; PubMed Central PMCID: PMCPMC551502.
68.   Welte MA, Gross SP, Postner M, Block SM, Wieschaus EF. Developmental regulation of vesicle
transport in Drosophila embryos: forces and kinetics. Cell. 1998;92(4):547-57. doi: 10.1016/s0092-
8674(00)80947-2. PubMed PMID: 9491895.
69.   Dawe HR, Adams M, Wheway G, Szymanska K, Logan CV, Noegel AA, et al. Nesprin-2 interacts
with meckelin and mediates ciliogenesis via remodelling of the actin cytoskeleton. Journal of Cell
Science. 2009;122(15):2716-26. doi: 10.1242/jcs.043794.
70.   Rajgor D, Mellad JA, Autore F, Zhang Q, Shanahan CM. Multiple novel nesprin-1 and nesprin-2
variants act as versatile tissue-specific intracellular scaffolds. PLoS One. 2012;7(7):e40098. Epub
20120702. doi: 10.1371/journal.pone.0040098. PubMed PMID: 22768332; PubMed Central PMCID:
PMCPMC3388047.
71.   Zhang Q, Ragnauth CD, Skepper JN, Worth NF, Warren DT, Roberts RG, et al. Nesprin-2 is a multi-
isomeric protein that binds lamin and emerin at the nuclear envelope and forms a subcellular network in
skeletal muscle. J Cell Sci. 2005;118(Pt 4):673-87. Epub 20050125. doi: 10.1242/jcs.01642. PubMed
PMID: 15671068.
72.   Zhang X, Lei K, Yuan X, Wu X, Zhuang Y, Xu T, et al. SUN1/2 and Syne/Nesprin-1/2 complexes
connect centrosome to the nucleus during neurogenesis and neuronal migration in mice. Neuron.
2009;64(2):173-87. doi: 10.1016/j.neuron.2009.08.018. PubMed PMID: 19874786; PubMed Central
PMCID: PMCPMC2788510.
73.   Razafsky D, Hodzic D. A variant of Nesprin1 giant devoid of KASH domain underlies the molecular
etiology of autosomal recessive cerebellar ataxia type I. Neurobiol Dis. 2015;78:57-67. Epub 20150402.
doi: 10.1016/j.nbd.2015.03.027. PubMed PMID: 25843669; PubMed Central PMCID: PMCPMC4426048.
74.   Razafsky DS, Ward CL, Kolb T, Hodzic D. Developmental regulation of linkers of the
nucleoskeleton to the cytoskeleton during mouse postnatal retinogenesis. Nucleus. 2013;4(5):399-409.
Epub 20130823. doi: 10.4161/nucl.26244. PubMed PMID: 23974729; PubMed Central PMCID:
PMCPMC3899130.
75.   Hao H, Kalra S, Jameson LE, Guerrero LA, Cain NE, Bolivar J, et al. The Nesprin-1/-2 ortholog ANC-
1 regulates organelle positioning in C. elegans independently from its KASH or actin-binding domains.

Elife. 2021;10:e61069. Epub 20210416. doi: 10.7554/eLife.61069. PubMed PMID: 33860766; PubMed

Central PMCID: PMCPMC8139857.

76.   Uversky VN. Intrinsically Disordered Proteins and Their "Mysterious" (Meta)Physics. Front Phys-

Lausanne. 2019;7:10. doi: 10.3389/fphy.2019.00010. PubMed PMID: WOS:000458061300001.

77.   Wang X, Bigman LS, Greenblatt HM, Yu B, Levy Y, Iwahara J. Negatively charged, intrinsically

disordered regions can accelerate target search by DNA-binding proteins. Nucleic Acids Res.

2023;51(10):4701-12. doi: 10.1093/nar/gkad045. PubMed PMID: 36774964; PubMed Central PMCID:

PMCPMC10250230.

78.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-

source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub 20120628. doi:

10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMCPMC3855844.

79.   Jorgensen V, Chen J, Vander Wende H, Harris DE, McCarthy A, Breznak S, et al. Tunable

Transcriptional Interference at the Endogenous Alcohol Dehydrogenase Gene Locus in Drosophila

melanogaster. G3 (Bethesda). 2020;10(5):1575-83. Epub 20200504. doi: 10.1534/g3.119.400937.

PubMed PMID: 32213532; PubMed Central PMCID: PMCPMC7202008.