TITLE: Preparation of functional metagenomic libraries from low biomass samples using

METa assembly and their application to capture antibiotic resistance genes

Authors: Allman HM[1*], Bernate EP[1*], Franck E[1], Oliaro FJ[2], Hartmann EM[3], Crofts TS[4,5,6]

1. Department of Biomedical Sciences, Florida State University, Tallahassee, FL, USA

2. Conservation Research Department, John G. Shedd Aquarium, Chicago, IL, USA

3. Department of Civil and Environmental Engineering, Northwestern University, Evanston,

IL, USA

4. Department of Animal Sciences, University of Illinois at Urbana-Champaign, Urbana, IL,

USA

5. Carl R. Woese Institute for Genomic Biology, Urbana, IL, USA

6. Division of Nutritional Sciences, University of Illinois at Urbana-Champaign, Urbana, IL

* Co-first authors

Corresponding author: Terence S. Crofts, tcrofts@illinois.edu

ABSTRACT

A significant challenge in the field of microbiology is the functional annotation of

sequence novel genes from microbiomes. The increasing pace of sequencing technology

development has made solving this challenge in a high-throughput manner even more

important. Functional metagenomics offer a sequence-naïve and cultivation-independent

solution. Unfortunately, most methods for constructing functional metagenomic libraries require

large input masses of metagenomic DNA, putting many sample types out of reach. Here, we

show that our functional metagenomic library preparation method, METa assembly, can be used

to prepare useful libraries from much lower input DNA quantities. Standard methods of

functional metagenomic library preparation generally call for 5 µg to 60 µg of input metagenomic

DNA. We demonstrate that the threshold for input DNA mass can be lowered at least to 30.5 ng,

a three-log decrease from prior art. We prepared functional metagenomic libraries using

between 30.5 ng and 100 ng of metagenomic DNA, and found that, despite their limited input

mass, they were sufficient to link MFS transporters lacking substrate-specific annotations to

tetracycline resistance and to capture a gene encoding a sequence novel GNAT family

acetyltransferase that represents a new streptothricin acetyltransferase, *satB*. Our preparation

of functional metagenomic libraries from aquatic samples and a human stool swab demonstrate

that METa assembly can be used to prepare functional metagenomic libraries from microbiomes

that were previously incompatible with this approach.

Figure 1. Functional metagenomics and METa assembly. A) The general steps in the preparation and application of a functional metagenomic library consist of: 1) Extraction of metagenomic DNA from a source microbiome. 2) Fragmentation of metagenomic DNA to preferred size range. 3) Packaging of inserts into expression vectors. 4) Transformation of host cells with vector library. 5) Screen or selection of functional metagenomic library for a phenotype of interest. 6) Collection of screened or selected metagenomic fragments. 7) Sequencing of selected inserts. 8) Open reading frame calling and annotation to identify potential genes underlying phenotypes of interest. B) Steps 2) and 3) above are modified in Mosaic ends tagmentation (METa) assembly. Fragmentation is achieved using Tn5 transposase tagmentation with mosaic end sequence oligos. Tagmented DNA is gap filled by polymerase and directly cloned (without amplification) into an expression vector with matching mosaic end sequences defining the cloning site using assembly cloning.

INTRODUCTION

Microbiomes, assemblages of bacteria, fungi, viruses and other microorganisms, are

host to some of the highest levels of genetic diversity on the planet. Due to the difficulty in

culturing all but a select low percentage of organisms in microbiomes (often estimated to be

~0.1% to 1%) current methods for studying microbiomes rely on DNA extracted from these

communities, metagenomic DNA[1–4]. High-throughput sequencing of metagenomic DNA provides

unrivaled insights into the taxonomic structure of a microbial community and can shed

significant light on its functional capacity as well[3]. However, even with the sequencing

revolution, the genetic diversity of microbiomes is still beyond our ability to accurately annotate

novel microbial genes. Complementing homology and structure-based methods for assigning

function to novel genes is the high-throughput, sequence- and cultivation-naïve method of

functional metagenomics. In this method, functional metagenomic libraries, sometimes referred

to as shotgun cloning libraries, are prepared by fragmenting metagenomic DNA and cloning

those fragments *en masse* into an expression microorganism. Once a library is prepared, any

genes encoded within captured metagenomic DNA fragments are in theory able to be

expressed by the host cell to give a phenotype that can be selected or screened for[4–7] (**Figure**

1A). Functional metagenomic libraries can be divided into two categories based on the average

size of their captured metagenomic DNA fragments, with small insert libraries generally

capturing fragments between 2 kb and 10 kb in length and large insert libraries (*i.e,* cosmid,

fosmid, and bacterial artificial chromosomes [BAC]) capturing fragments between 15 kb and

100s of kb in length[7].

While functional metagenomics is a powerful method for linking novel genes to functions

without requiring the growth of non-model organisms in the lab, it has its own shortcomings.

These issues include the potential for gene toxicity, limited expression of foreign genes in library

hosts such as *E. coli*, limited coverage of target metagenomes, and requirements for large

quantities of metagenomic DNA for preparation of the functional metagenomic library itself. We

recently reported the development of a new method for preparing small insert functional

metagenomic libraries that we call METa assembly[8]. This functional metagenomic library

preparation method relies on tagmentation to fragment input metagenomic DNA and an

assembly-based cloning step to replace blunt ligation (**Figure 1B**). We previously demonstrated

that this method circumvents one shortcoming of functional metagenomics, limited

metagenomic coverage, by showing that METa assembly libraries are more efficient at capturing

DNA and can produce approximately 80-fold larger libraries per input DNA mass. We also

showed that METa assembly could be used to prepare functional metagenomic libraries

encoding tens of gigabase pairs (Gb) of captured DNA using sub-µg inputs of metagenomic

DNA, far lower than the standard 5 µg to 20 µg input[8,9].

Here we build on our earlier results by testing if METa assembly can be used to prepare

functional metagenomic libraries with even lower quantities of input DNA. We found that we

could consistently prepare functional metagenomic libraries starting with 50 ng or less input

DNA or even DNA extracted from a swab dipped into a 10 mg/ml human stool slurry. These

libraries were large enough that, when selected for resistance to the antibiotics tetracycline and

nourseothricin, they yielded genes predicted to encode novel proteins. These findings suggest

that even low biomass microbiomes and samples can be studied by functional metagenomics

using METa assembly.

METHODS

Strains and materials

Routine cultivation of *E. coli* was performed at 37°C with shaking at 250 rpm for aeration

in the case of liquid cultures. Cultivation for cloning and starter cultures was performed using

lysogeny broth (Miller) (LB) with the addition of kanamycin at 50 µg/ml to maintain plasmids.

Antimicrobial susceptibility testing was performed using Cation-Adjusted Mueller-Hinton media

(MH) (Teknova, 101320-364). *E. coli* clones used in this study are of the DH10B lineage and

carry plasmids derived from pZE21 unless otherwise noted[8,10,11]. *E. coli* clones were maintained

at -80°C in 15% glycerol in LB. Transposase enzymes and mosaic ends oligos were previously

prepared and stored at -20°C[8,12]. Standard chemicals used in this study were of high purity for

molecular biology and microbiology. Kanamycin sulfate (VWR Life Science, 75856-68),

nourseothricin (Research Products International, N51200-1.0), tetracycline (Aldon Corp,

TT0070-5GR), chloramphenicol (Thomas Scientific, 216035-25G), ciprofloxacin hydrochloride

(Corning, 61-277-RG), azithromycin dihydrate (TCI, A2076), and florfenicol (Fisher Scientific,

F08115G) were stored at 4°C as powders. Working solutions were stored at -20°C as filtered

aqueous solutions at 50 mg/ml concentration (kanamycin and nourseothricin), 5 mg/ml in

ethanol (tetracycline), 50 mg/ml in ethanol (chloramphenicol), 10 mg/ml in alkaline water

(ciprofloxacin), or 50 mg/ml in DMSO (azithromycin and florfenicol).

Preparation of 50 ng test functional metagenomic libraries

102   To determine if agarose extraction could be used to purify and size-select DNA following

low mass tagmentation reactions, we used a DNA ladder (ThermoFisher Scientific 1 kb plus,

10787018) to mimic fragmented DNA. Mock tagmentation reactions were made with a final DNA

concentration of 50 ng/µl by combining 1 µl of diluted DNA ladder with 3 µl of autoclaved milliQ

water and 1 µl of 5X Taps tagmentation buffer (50 mM TAPS pH 8.5, 25 mM $MgCl_2$, 50% v/v

DMF in water)[12]. We added 0.25 µl of 1% sodium dodecyl sulfate (SDS) to the 5 µl mock

reactions to simulate reaction quenching followed by 1 µl of 6x DNA dye (15% Ficoll 400 w/v,

0.25% Orange G w/v in water) before loading on to a 0.7% agarose gel in TAE buffer containing

SybrSafe dye (ThermoFisher Scientific, S33102). DNA bands were visualized over blue light

and a clean razor was used to cut out gel portions containing bands greater than 2 kb in length.

DNA was extracted from the agarose gel fragments using a DNA extraction kit (New England

Biolabs, T1020L) following manufacturer instructions with modification. Each agarose fragment

was weighed and incubated with four volumes of gel dissolving buffer for at least 10 minutes at

55°C. The dissolved agarose was then applied to a spin column, spun through, then re-applied

a second time. The standard protocol was followed until the elution step where each column

was eluted twice sequentially with 6 µl of 55°C autoclaved milliQ water. DNA concentration was

measured using the Quant-It DNA quantification system (Invitrogen).

To test the preparation of functional metagenomic libraries using 50 ng of input

metagenomic DNA, we used a sample of previously extracted metagenomic DNA from a

Canada goose fecal pellet that was stored at -20°C[8]. Briefly, metagenomic DNA for that sample

was extracted using a DNeasy PowerSoil kit (Qiagen, catalog no. 12888-100) with modifications

including incubation of fecal material in Qiagen buffer CD1 at 65°C for 10 minutes and 95°C for

10 minutes[8,13]. For library preparation, we followed our previously published META assembly

protocol[8]. Briefly, triplicate 5 µl tagmentation reactions containing 50 ng of metagenomic DNA,

1x TAPS-DMF buffer, and 25 ng of in-house transposase[12] loaded with mosaic end sequence

oligos were incubated at 55°C for 7 minutes followed by quenching by addition of SDS to reach

0.05% final concentration and incubation for 5 more minutes at 55°C. To each reaction, 2 µl of

6x loading dye was added and fragmented metagenomic DNA in the range of 2 kb to 9 kb was

extracted by gel excision as described above. The 12 µl of size-selected metagenomic DNA

fragments were gap-filled by the addition of 12 µl of 2x Q5 DNA polymerase (New England

Biolabs, M0492S) (pre-incubated at 98°C for 30 seconds) and held at 72°C for 15 minutes

before purification by PCR and DNA cleanup kit (New England Biolabs, T1030S) according to

manufacturer instructions with the following modifications: DNA dissolved in two volumes of

binding buffer was applied to the spin column and the flow-through was applied a second time.

At elution, we used two sequential applications of 6 µl of 55°C autoclaved milliQ water.

Metagenomic DNA inserts were cloned into our plasmid pZE21-ME using 2x NEBuilder HiFi

assembly mix (New England Biolabs, E2621L) according to manufacturer instructions

(assuming an insert DNA mass of 15 ng and an average fragment size of 2 kb). The triplicate

libraries were purified by PCR and DNA cleanup kit and the full 12 µl of purified library DNA was

introduced into 25 µl of *E. coli* DH10B electrocompetent cells (New England Biolabs, C3020K)

by electroporation at 1.8 kV in 0.1 mm cuvettes. Actual voltage reached and the electroporation

time constant were recorded for all electroporation reactions to detect failed or low efficiency

transformations. All three libraries were rescued post-transformation by inoculation into 1 ml of

SOC outgrowth medium (New England Biolabs, B9020S) pre-warmed to 37°C and incubated

with aeration at 37°C for 1 hour.

Following recovery, 100 µl of 100-fold, 10,000-fold, and 1,000,000-fold diluted cultures

were plated on to LB agar with kanamycin and incubated overnight at 37°C. The colonies were

counted to estimate the number of unique clones resulting from each transformation. The

remaining cells (~1 ml) were inoculated into 50 ml of LB with kanamycin and incubated at room

temperature overnight with aeration to amplify the functional metagenomic library. Once cultures

reached a 1 cm pathlength optical density at 600 nm (OD_{600}) of between 0.6 and 1.0, cells were

collected by centrifugation at 4,000 rcf for 7 minutes at 4°C. Pellets were resuspended in 10 ml

of LB containing kanamycin and 15% glycerol and stored at -80°C as 1 ml aliquots. The

triplicate functional metagenomic library sizes were determined using the Dantas lab calculator

(http://dantaslab.wustl.edu/LibSizeCalc/) considering unique cell count, determined above,

average metagenomic insert size, and the proportion of clones containing an empty plasmid.

This proportion and the average insert size were determined by performing colony PCR as

previously described on 9 clones from each replicate library[8].

Preparation of an aquatic functional metagenomic library from a Shedd Aquarium

microbiome

Preparation of a functional metagenomic library from aquatic metagenomic DNA began

with a previously extracted metagenomic DNA sample. The initial sample was collected from the

Shedd Aquarium in Chicago in the Underwater Beauty exhibit in May, 2019 and metagenomic

DNA was extracted[14]. A functional metagenomic library was prepared from 50 ng of DNA as

described above and evaluated by colony counting and colony PCR (n=14 colonies) to

determine library size. After preparation of the functional metagenomic library, quantification of

the input DNA by fluorescent dye (Quant-It) revealed that only 30.5 ng of metagenomic DNA

had been used as input for this library and this corrected DNA mass was used in determining

library preparation efficiency.

Functional metagenomic selection for antibiotic resistance from the aquarium functional

metagenomic library

The aquarium functional metagenomic library was selected for resistance to the

antibiotics tetracycline, ciprofloxacin, and florfenicol. MH agar plates containing 50 µg/ml

kanamycin were prepared to contain each antibiotic at the following concentrations previously

established to be inhibitory to *E. coli*[15]: 8 µg/ml tetracycline, 0.5 µg/ml ciprofloxacin, and 8 µg/ml

florfenicol. An aliquot of the functional metagenomic library was thawed from -80°C storage and

a volume of suspension calculated to contain approximately 10-fold more total cells than the

unique cell count was plated on each antibiotic and incubated overnight at 37°C. A parallel

series of agar plates were plated with a similar cell count of *E. coli* cells containing only empty

vectors as a control.

Following incubation, 15 colonies each from the tetracycline and florfenicol selections

were collected and re-streaked onto LB agar plates containing kanamycin and either

tetracycline or florfenicol to obtain pure cultures of resistant clones. Plasmids were extracted

from each clone by miniprep kit (New England Biolabs, T1010L) and whole plasmid sequencing

was performed by Plasmidsaurus using Oxford Nanopore Technology with custom analysis and

annotation. The metagenomic inserts were identified by the presence of flanking mosaic end

sequences (5'- agatgtgtataagagacag-3') and clustered on the Clustal Omega webserver[16] to de-

replicate metagenomic fragments that were captured in the selection more than once. The

metagenomic inserts targeted for further study (TET1, TET3, TET13) were analyzed by the

MetaGeneMark program[17] to identify potential open reading frames which were subjected to

analysis by Basic Local Alignment Search Tool (BLAST)[18,19] against the NCBI non-redundant

database and The Comprehensive Antibiotic Resistance Database (CARD)[20,21]. The annotated

metagenomic DNA fragments were made into schematics using ApE plasmid editor[22] and

Microsoft PowerPoint.

Preparation of a functional metagenomic library from a fecal swab

An aliquot of the ZymoBIOMICS fecal reference standard (Zymo Research, D6323) was

used as a source for a fecal microbiome swab functional metagenomic library. An aliquot of

fecal standard was thawed from -80°C storage and a sterile swab was dipped into the tube and

gently wrung out against the tube's side as it was removed. Metagenomic DNA was extracted

from the swab using a PowerSoil Pro kit (Qiagen, 47014). The swab was agitated in 800 µl of

C1 solution to release material and DNA extraction was performed following manufacturer

instructions with the eluted DNA reduced in volume using a vacuum concentrator. A functional

metagenomic library was prepared by METa assembly as above, with the tagmentation reaction

scaled to match the 100.8 ng of metagenomic DNA available from the extraction. Library size

was evaluated by colony count and colony PCR as above (n=14 colonies).

Functional metagenomic selection for nourseothricin resistance from the stool functional

metagenomic library

In our previous nourseothricin resistance selection[8], we used agar plates containing 64

µg/ml nourseothricin but this concentration of drug is likely well beyond the *E. coli* minimal

inhibitory concentration (MIC). To determine conditions appropriate for a functional

metagenomic selection for nourseothricin resistance genes, we performed an agar dilution

assay on MH agar with drug concentrations ranging from 0.5 µg/ml to 2048 µg/ml in two-fold

jumps[23]. Each agar plate was inoculated with 100 µl of a 0.5 McFarland unit suspension of *E.*

coli DH10B harboring an empty pZE21-ME plasmid. Plates were incubated at 37°C for roughly

24 hours prior to being observed for growth to determine an appropriate concentration for

functional metagenomic selections.

A functional metagenomic selection for nourseothricin resistance genes was carried out

as above with the stool swab functional metagenomic library plated to carry an estimated ten-

fold excess of cells over unique clones. The functional metagenomic library was plated on MH

agar with 16 µg/ml nourseothricin and 50 µg/ml kanamycin. In parallel, a suspension of a control

E. coli DH10B clone with empty vector was also plated as above at a similar predicted titer. The

plates were incubated at 37°C overnight. A single colony on the nourseothricin plate was picked

and re-streaked to obtain a pure culture and processed by miniprep kit to obtain a plasmid for

sequencing. The plasmid sequence was analyzed as before. The predicted acetyltransferase

open reading was synthesized into a pZE21-derivative plasmid[10,11] by Integrated DNA

Technologies (IDT) and an aliquot of electrocompetent *E. coli* DH10B was transformed with the

plasmid for further resistance study.

Phylogenetic analyses

For phylogenetic analysis of the putative tetracycline efflux pumps (TET1, TET3, and

TET13) we downloaded CARD amino acid sequences[20,21] for the ontology term "major facilitator

superfamily (MFS) antibiotic efflux pump" and added the functionally-selected predicted efflux

pump amino acid sequences to the FASTA file. For analysis of the predicted nourseothricin

resistance protein, we downloaded CARD protein sequences for the ontology term "acylation of

antibiotic conferring resistance", which includes the streptothricin, aminoglycoside, apramycin,

virginiamycin, capreomycin, and chloramphenicol acetyltransferase sequences. We also

downloaded the amino acid sequence for the NAT nourseothricin acetyltransferase[24].

Phylogenetic trees were prepared by first aligning the protein sequences using the

Clustal Omega web server with default settings[16]. Next a maximum likelihood phylogenetic tree

was prepared from the alignment using IQ-TREE[25,26] with the "Find best and apply" substitution

model, standard 100 replicates bootstrap and otherwise default settings. The resulting

consensus trees were visualized using the Interactive Tree of Life (iTOL) display and annotation

tool[27].

The streptothricin acetyltransferase multiple protein alignment was prepared using

Clustal Omega as above using only the STAT, NAT, SAT-2, SAT-3, SAT-4, SatA, and SatB amino

acid sequences. This also provided a percent identity matrix for the streptothricin

acetyltransferase proteins. The alignment was edited and visualized in Jalview[28] and amino acid

residue numbering was set according to the *Bacillus anthracis* SatA crystal structure

sequence[29].

Microbroth dilution antimicrobial susceptibility testing

The antimicrobial resistance of *E. coli* clones expressing metagenomic DNA fragments

(TET1, TET3, TET13, and NTC1), the *satB* open reading frame, or the *stat* gene (pGDP1 stat

was a gift from Gerard Wright (Addgene plasmid # 112886 ; http://n2t.net/addgene:112886;

RRID:Addgene_112886)) were quantified by microbroth dilution assay[23,30]. Briefly, the

corresponding clones and control *E. coli* carrying an empty plasmid were streaked for single

colonies on LB agar with kanamycin. Approximately 10 colonies for each clone were suspended

in 2 ml of MH media with kanamycin and brought to an OD_{600} of between 0.1 and 0.15 to

correspond to a 0.5 McFarland standard density, then diluted 100-fold in MH broth with

kanamycin. We prepared a 96-well plate (Costar, 3370) to contain 50 µl of MH broth with 50

µg/ml kanamycin and 2-fold increasing concentrations of tetracycline, chloramphenicol,

azithromycin, or nourseothricin at twice the final target concentration. We added 50 µl of the

diluted bacterial suspensions to this plate, with each clone assayed in quadruplicate. The 96-

well plates were sealed with a Breathe-Easy membrane (MilliporeSigma, Z380059) and

incubated at 37°C with shaking for 20 to 24 hours. Culture density was recorded as OD_{600} with a

1 cm pathlength correction using an Epoch 2 plate reader (BioTek) and concentration-response

curves were fitted to four parameter Hill equations in GraphPad Prism 10.2.3 (GraphPad

Software, La Jolla, CA, USA) to determine 50% inhibitory concentration (IC_{50}) values.

Differences between IC_{50} values for control *E. coli* and the tested clones were evaluated for

significance in GraphPad Prism using lognormal Brown-Forsythe ANOVA tests with multiple

comparison correction using the Dunnett T3 method.

SatB structural prediction

Structural modelling of the SatB protein was performed using AlphaFold2 through the

ColabFold v1.5.5 Jupyter notebook[31,32]. The predicted amino acid sequence was entered as

query sequence and default parameters were used to generate the model. The model was

superimposed against the SatA crystal structure (3PP9)[29,33,34] using the Mol* 3D viewer[35] hosted

by the Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB)[36].

Mol* 3D calculated the root mean squared deviation (RMSD) of the resulting superposition of

SatB on SatA.

Sequence availability

Sequences for the TET1, TET3, TET13, and NTC1 metagenomic DNA fragments and the

proposed *satB* gene can be found under GenBank accession numbers PV822507, PV822508,

PV822509, PV8225110, and PV822511.

RESULTS

METa assembly can reproducibly prepare Gb sized libraries with 50 ng of input

metagenomic DNA

Our published META assembly protocol tagments metagenomic DNA at a concentration

of 10 ng/µl, meaning a 50 ng DNA tagmentation reaction would have a 5 µl volume where

accurate pipetting starts to become difficult. To test the practicality of extracting useful DNA

fragments from this size reaction, we diluted 50 ng of a DNA ladder in tagmentation buffer in

triplicate to a concentration of 10 ng/µl to make mock 5 µl tagmentation reactions. The triplicate

mock fragmented samples were loaded on to an agarose gel for excision of bands ranging from

approximately 2 kb to 10 kb (**Supplemental figure 1A**). This resulted in recovered DNA

quantities (acting as stand-ins for fragmented metagenomic DNA) averaging 12.7 ng (± 2.7

standard deviation). Given that the total mass of the DNA bands 2 kb and higher was 16 ng, this

represented a 79% yield and suggested that size-selection of fragmented DNA from a 50 ng

tagmentation reaction could be feasible (**Supplemental figure 1B**).

Next, we tested this explicitly by preparing a 50 ng metagenomic DNA functional

metagenomic library in triplicate. Using previously extracted goose fecal metagenomic DNA as

Table 1. Summary statistics of functional metagenomic libraries

Library	Input DNA	DNA mass (ng)	Insert size (kb)	Proportion	Library size (Gb)	Efficiency (Gb/µg)	Selections[1]
50 ng goose 1	Goose gut microbiome	50	2.48	1.00	0.89	17.8	n/a
50 ng goose 2[2]	Goose gut microbiome	50	2.62	1.00	0.10	2.09	n/a
50 ng goose 3	Goose gut microbiome	50	3.13	1.00	2.46	49.3	n/a
Aquatic	Shedd aquarium	30.5	2.77	0.88	0.72	23.6	TET, CIP, FFC
Fecal	Human fecal reference	100.8	2.70	0.90	1.48	14.7	NTC

[1]no selection (n/a), tetracycline (TET), ciprofloxacin (CIP), florfenicol (FFC), nourseothricin (NTC)
[2]Replicate sparked during electroporation

input[8], we performed three tagmentation reactions in parallel and used gel excision to size-

select for DNA fragments between 2 kb and 10 kb (**Supplemental figure 2A**). These fragments

were used to prepare three functional metagenomic libraries by METa assembly. During the

electroporation step, replicates 1 and 3 reached full voltage (1.8 kV) and had time constants of

4.6 and 4.8 seconds respectively. Replicate 2 reported a voltage of 1.65 kV and a 1.0 second

time constant, consistent with the sample 'sparking'. Replicate #1 had an average insert size of

2.48 kb, no empty vectors, and an average library size of 0.89 Gb with a range of 0.69 Gb to 1.1

Gb. For replicate #2 these values were 2.62 kb, no empty vectors, and 0.1 Gb with a range of

0.08 Gb to 0.13 Gb. Replicate #3 values were 3.13 kb, no empty vectors, and 2.46 Gb with a

range of 1.61 Gb to 3.27 Gb. Averaging across the three libraries we found an overall average

insert size of 2.75 kb with no empty vectors (**Supplemental figure 2B**), an average library size

of 1.15 Gb, and a library preparation efficiency of 23 Gb/µg of input DNA. Notably, these

averages include replicate #2 with a dramatically smaller library size (100 Mb compared to 890

Mb and 2.5 Gb) (**Table 1**). If replicate #2 is excluded, the average library size for the 50 ng

functional metagenomic libraries increases to 1.68 Gb with an efficiency of 33.6 Gb/µg.

METa assembly preparation of a functional metagenomic library from a low biomass

aquarium microbiome

Two areas where input mass is limiting for classic functional metagenomic library

preparation are 1) when the microbiome of interest is low in biomass (*e.g.,* aquatic sources as

opposed to soil or fecal samples) and 2) when the microbiome of interest is not readily available

for re-sampling. We decided to test low-input mass METa assembly on a microbiome that fulfills

both limitations. In a previous project, metagenomic DNA was extracted from the Underwater

Beauty exhibit at the Shedd Aquarium in Chicago, Illinois following filtration capture of planktonic

cells[14]. We used a sample of previously extracted metagenomic DNA to prepare a functional

metagenomic library. Tagmentation was performed on 30.5 ng of metagenomic DNA and

fragments from 2 kb to 10 kb were size selected by gel excision. Following transformation,

random colonies from the functional metagenomic library were used as template for PCR to

amplify captured metagenomic inserts to determine library statistics. We found that this aquatic

functional metagenomic library captured 720 Mb of metagenomic DNA (range of 0.42 Gb to 1.17

Gb) with an average insert size of 2.77 kb and with 88% of plasmids containing an insert. The

library preparation efficiency was calculated to be 23.6 Gb/µg (**Table 1**).

Tetracycline resistance in an aquarium functional metagenomic library

To evaluate the usefulness of this functional metagenomic library, we used it to

investigate antimicrobial resistance against three antibiotics that are commonly used in

aquaculture: tetracycline, florfenicol, and ciprofloxacin[37,38]. The selection on ciprofloxacin did not

yield any resistant colonies but colonies were recoverable from both the tetracycline and

florfenicol plates. As an initial investigation, fifteen colonies from only the tetracycline selection

were collected and submitted for whole plasmid sequencing. We dereplicated repeated inserts,

which resulted from plating a ten-fold excess of total cells over unique cells, resulting in three *E.*

coli clones with unique metagenomic DNA fragments we termed TET1, TET3, and TET13. The

metagenomic DNA captured in these cones had top nucleotide BLAST hits to *Lentibacter*

algarum strain SH36 (95% coverage, 82.09% identity), *Legionella pneumophila* strain NY23

(30% coverage, 66.87% identity), and *Saccharophagus degradans* strain FZY0027 (32%

coverage, 70.26% identity), respectively.

Figure 2. Tetracycline selected metagenomic DNA fragments from an aquarium microbiome. A) Gene schematics of the tetracycline-selected TET1, TET3, and TET13 metagenomic DNA fragments with predicted efflux pumps highlighted (TET1 teal, TET3 red, TET13 blue). Other predicted open reading frames are shown as empty arrows and plasmid backbone markers are highlighted as follows: Promoter (green), mosaic end sequences (orange), terminator (red). B) Maximum likelihood consensus phylogenetic tree of MFS efflux pumps with predicted efflux pump amino acid sequences from TET1 (teal), TET3 (blue), and TET13 (red) fragments highlighted.

We annotated the metagenomic DNA fragments captured in these three clones

(**Supplemental figure 3**) and found them to each contain predicted MFS family efflux pumps

with low full-length amino acid identity (<30%) to their closest match in The Comprehensive

Antibiotic Resistance Database (CARD)[20,21] (**Figure 2A**). When placed on a phylogenetic tree

alongside known antibiotic resistance efflux pumps, TET1 and TET13 clustered with each other,

a bicyclomycin efflux pump (BCR-1), and efflux pumps associated chloramphenicol resistance

(PexA, MdfA, and Cml

family pumps) (**Figure 2B**).

TET3 instead clustered

with efflux pumps with a

variety of targets including

tetracycline (TetV, TAP),

macrolides (MefD), and

multiple antimicrobials

(Hp1181, EfmA,) (**Figure**

2B).

The resistance of all

three clones to tetracycline,

chloramphenicol, and

azithromycin were

measured by microbroth

dilution. All three clones

showed increased

resistance to tetracycline,

with TET1 and TET3

notably showing an ability

to grow in the presence of

Figure 3. Efflux pump containing metagenomic DNA fragments from an aquarium confer tetracycline resistance. Microbroth dilution assay curves and calculated 50% inhibitory concentration (IC_{50}) values for *E. coli* clones carrying the indicated metagenomic DNA fragments (TET1, TET3, or TET13). A) and B) tetracycline, C) and D) chloramphenicol, or E) and F) azithromycin. (**** $p<0.0001$, ** $p<0.005$, * $p<0.05$, n=4 for all).

8 µg/ml tetracycline, the breakpoint for resistance determination in *Enterobacterales* like *E. coli*

(**Figure 3A**). All three *E. coli* clones gained a statistically significant increase in their tetracycline

50% inhibitory concentration (IC_{50}) value (**Figure 3B**). Two of the clones showed a statistically

significant change in chloramphenicol resistance (TET1 being more susceptible and TET13

being more resistant) (**Figure 3C and 3D**). However, the limited change in absolute terms (less

that a two-fold change) suggests they are not biologically significant. None of the three clones

showed a significant change in azithromycin resistance compared to the control clone (**Figure**

3E and 3F).

META assembly preparation of a functional metagenomic library from a human stool

swab

Another area where the ability to prepare a functional metagenomic library using minimal

DNA resources could be beneficial is within clinical settings. For example, fecal swabs provide

access to the medically important gut microbiome, but extraction of metagenomic DNA from this

sample type falls far below what classic functional metagenomic library preparation methods call

for[15,39]. To test the possibility of medical swabs being used as input for META assembly, we

turned to the ZymoBIOMICS fecal reference with TruMatrix technology fecal standard. Each

aliquot of this standard is expected to contain 10 mg (wet weight) of human fecal material in 1

ml of DNA/RNA shield preservation buffer. We sampled this microbiome by dipping a sterile

swab into a thawed tube of the material followed by DNA extraction from the swab, resulting in

collection of 100.8 ng of DNA. We prepared a functional metagenomic library by META

assembly and evaluated it as above and found 90% of colonies contained an insert, each with

an average size of 2.7 kb, for a total expected library size of 1.48 Gb (range of 0.62 Gb to 2.32

Gb) and a library preparation efficiency of 14.7 Gb/μg (**Table 1**).

Nourseothricin resistance in a human stool functional metagenomic library

We performed a functional metagenomic selection for nourseothricin resistance in the

human gut microbiome using the Zymo fecal library. In our previous work we performed

Figure 4. Predicted novel streptothricin acetyltransferase from the human gut microbiome. A) Gene schematics of the NTC1 metagenomic DNA fragment with the predicted acetyltransferase gene highlighted ('*satB*', blue). Predicted genes captured on the same DNA fragment are shown as empty arrows. Plasmid backbone elements are: Promoter (green), mosaic end sequence (orange), terminator (red). B) Maximum likelihood consensus phylogenetic tree of the predicted SatB protein (blue) in the context of known streptothricin acetyltransferases (SATs, NAT, and STAT) and aminoglycoside, chloramphenicol, virginiamycin, apramycin, and capreomycin acetyltransferases.

nourseothricin selections using a concentration of 64 µg/ml[8] but here we found that 16 µg/ml is

more than sufficient. This selection resulted in the capture of a single nourseothricin resistant

colony which we collected and extracted a plasmid from to sequence the resistance-conferring

metagenomic DNA fragment. A parallel experiment in which an equal titer of control *E. coli* cells

containing an empty plasmid was plated on agar containing 16 µg/ml nourseothricin resulted in

the growth of no colonies.

Following whole plasmid sequencing of the resistant colony, we found that the NTC1

metagenomic DNA fragment showed limited identity and coverage with organisms in the NCBI

database, with the top nucleotide BLAST hit being *Clostridia* bacterium i40-0019-1A8 (20%

coverage, 76.08% identity). The Zymo fecal reference has been shotgun sequenced by other

groups and we confirmed the presence of short reads in these sequencing runs (*i.e.,*

SRX26261613[40]) with >98% nucleotide identity to the NTC1 metagenomic DNA fragment,

confirming its origin in the Zymo stool material.

The metagenomic DNA fragment was predicted to contain three open reading frames

including a potential GCN5-Related N-Acetyltransferase (GNAT) (**Figure 4A**), the enzyme

family to which antibiotic-inactivating acetyltransferase enzymes belong. We prepared a

phylogenetic tree with the predicted streptothricin acetyltranferase, which we termed SatB,

alongside the six other streptothricin acetyltransferase family members (STAT, NAT, SatA, SAT-

2, SAT-3, and SAT-4) and aminoglycoside, virginiamycin, capreomycin, and chloramphenicol

acetyltransferases from the CARD database (**Figure 4B**). This tree places SatB in the same

clade as the other streptothricin acetyltransferases but at a distance comparable to other distinct

members of the family.

To explore this further we investigated the ability of the NTC1 metagenomic DNA

fragment and of the

predicted *satB* open

reading frame to confer

nourseothricin resistance in

E. coli. As a positive control

we also included an *E. coli*

clone expressing the *stat*

gene[41]. All three clones

showed an ability to grow

at higher nourseothricin

concentrations than a

Figure 5. Streptothricin resistance is conferred by the NTC1 metagenomic DNA fragment and *satB*. A) Results of a microbroth dilution assay of *E. coli* clones grown in the presence of variable nourseothricin concentrations (NTC1, *E. coli* carrying the NTC1 metagenomic DNA fragment; *stat*, *E. coli* expressing the *stat* streptothricin acetyltransferase; *satB*, *E. coli* expressing the predicted *satB* open reading frame from the NTC1 fragment). B) IC_{50} values calculated from dose-response curves. (**** $p<0.0001$, *** $p<0.0005$, ** $p<0.005$, n=4 for all).

control *E. coli* clone with an empty plasmid (**Figure 5A**). Expression of the NTC1 metagenomic

DNA fragment was sufficient to increase the nourseothricin resistance of *E. coli* by

approximately 64-fold and expression of the *stat* gene or the putative *satB* gene provided even

greater resistance,

allowing *E. coli* to

grow at

nourseothricin

concentrations at

least 1,000-fold

greater than when

carrying an empty

plasmid (**Figure**

5B).

Given the

position of the

predicted SatB

enzyme on the

antimicrobial

acetyltransferase

phylogenetic tree

(**Figure 4B**) and the

ability of the *satB*

gene to confer

Figure 6. SatB comparison to SatA and other streptothricin acetyltransferases. A) Multiple sequence alignment of the six named streptothricin acetyltransferase enzymes and the proposed SatB enzyme. Conserved residues are highlighted in color, and residues predicted to be important in strepothricin binding are indicated with blue stars (positions 136, 145, 149, 154, and 164) and the predicted catalytic glutamate is highlighted by a red star (position 137) (numbering system corresponds to SatA). B) Heat map of percent amino acid identities between each of the acetyltransferase enzymes, darker colors indicate higher percent identity. C) AlphaFold2 model of the predicted SatB enzyme (blue) superimposed on the SatA crystal structure (red) (PDB 3PP9) (RMSD 2.16 Å).

streptothricin resistance (**Figure 5**) we next considered our argument that SatB represents a

new member of the streptothricin acetyltransferase family. A multiple sequence alignment of

SatB and other streptothricin acetyltransferases showed that SatB has identical amino acid

residues at predicted key positions for catalysis (Glu137) and streptothricin binding (Leu136,

Ala145, Tyr149, Phe154, and Tyr164, numbering based on SatA crystal structure) (**Figure 6A**).

A percent amino acid identity matrix of streptothricin acetyltransferases and a single

aminoglycoside acetyltransferase indicates that the SatB sequence does not share high levels

of identity with any of the other proteins, its highest identity being 37% compared to SAT-4

(**Figure 6B**) (the full-length Needleman-Wunsch identity of SatB with SAT-4 is only 33%). SatB

is as unique compared to the other acetyltransferases (average percent identity of 29%) as

each acetyltransferase protein is to the rest of the enzymes in the matrix (average percent

identity of 29%).

Finally, we used AlphaFold2[31,32] to model the three-dimensional structure of SatB and

compared it to the solved X-ray crystal structure of SatA from *Bacillus anthracis* (3PP9)[29,33].

When superimposed, the two models show similar secondary structure and have a global root

mean squared error (RMSD) of 2.16 Å (**Figure 6C**). In this predicted structure, SatB has an N-

terminal region with an alpha helix not found in SatA (**Figure 6C**) which is also evidence in the

multiple sequence alignment (**Figure 6A**).

DISCUSSION

Here we report that the META assembly method for preparing functional metagenomic

libraries can be extended to preparing libraries from low biomass and otherwise precious

microbiome samples with limited available metagenomic DNA. As an illustration of this potential,

we prepared modestly sized functional metagenomic libraries from metagenomic DNA extracted

from an aquarium and from a swab of a human stool sample. We further demonstrated the utility

of the resulting functional metagenomic libraries by applying them to identify likely tetracycline

efflux pumps and a novel streptothricin acetyltransferase.

A key result of our experiments is that using META assembly to prepare functional

metagenomic libraries allows us move the amplification-free minimum metagenomic DNA input

to approximately two orders of magnitude lower than previous protocols[9,15]. An advantage of our

tagmentation-based fragmentation approach over physical fragmentation (*e.g., via* sonication) is

its ready scalability to lower volumes and therefore lower input DNA masses. However, we

suspected that DNA losses during purification and fragment size-selection by gel excision would

make sub-200 ng libraries impractical. Instead, we show that, across our experiments, 50 ng or

lower input metagenomic DNA masses can be used to prepare functional metagenomic libraries

encoding useful amounts of metagenomic DNA (**Table 1**). We propose, and demonstrate, that

this opens new sample types to analysis by functional metagenomic selections or screens.

These include samples that are intrinsically of low biomass, either due to difficulty in collection

or their rarity (*e.g.,* aquatic metagenomes, but also historically banked microbiomes or samples

from remote regions). These also include microbiomes sampled using medical swabs. While our

example stool functional metagenomic library used Zymo fecal reference material, the mass of

the metagenomic DNA extracted from the swab and used to prepare a functional metagenomic

library was less than what has been reported in extractions from other fecal swabs (*i.e.,* ~100 ng

vs ~370 ng[39]). Functional metagenomic selections have been proposed to be useful for medical

applications[42], and our results support the feasibility of this.

Prior studies that have prepared functional metagenomic libraries from limited

metagenomic DNA samples have relied on multiple displacement amplification (MDA) reactions

to amplify DNA for input into cosmid large insert functional metagenomic libraries. This

approach allowed researchers to convert nanogram or even picogram quantities of

metagenomic DNA into the microgram quantities needed for library preparation[43,44]. However,

MDA reactions, as well as other amplification-based approaches to increasing DNA mass, come

with multiple issues, including selection-based bias (preferential amplification of a subset of

templates in the input DNA mixture due to GC content or DNA conformation), drift bias

(preferential amplification of a template by random chance), mutation (due to polymerase error

present even in high-fidelity enzymes), and risk of amplification of contaminating DNA templates

(such as those found in 'kitomes')[45,46]. The risk of these biases may be unavoidable for groups

preparing functional metagenomic libraries from picogram DNA quantities, but our results show

that METa assembly can likely replace MDA for functional metagenomic libraries prepared from

nanogram DNA masses.

Our previous smallest METa assembly input, 200 ng, resulted in a functional

metagenomic library encoding approximately 13.5 Gb with a preparation efficiency of ~68

Gb/µg[8]. Here, our 50 ng, 30.5 ng, and 100.8 ng libraries averaged an efficiency of between 20.5

Gb/µg and 24.1 Gb/µg, depending on the inclusion or exclusion of a library that sparked during

transformation, suggesting that lower input DNA masses may come with decreased library

preparation efficiency (**Table 1**). It is possible that this is due to DNA losses during the multiple

purification steps involved in library preparation. Nevertheless, these levels of efficiency still

perform more than an order of magnitude better than the average for blunt ligation-based

techniques (1 Gb/µg)[8].

While the functional metagenomic libraries prepared here are smaller than our previous

high-mass input libraries, their usefulness is demonstrated by their capture of genes for

tetracycline efflux pumps and streptothricin resistance (**Figures 2 and 4**). The three predicted

tetracycline efflux pumps show homology to previously identified MFS efflux pumps, with the

predicted amino acid sequences having near complete coverage and between 65% and almost

100% identity to hits in the NCBI database. However, they have only low identity to CARD hits,

with full length alignments showing only 28% to 29% identity. It is common for proteins identified

through functional metagenomic selections for antibiotic resistance to have high identity to

proteins in the NCBI database, but for those proteins to not previously have been identified as

being antibiotic resistance enzymes[47]. Tetracycline efflux pumps are found within many

environments[48–51] but the difficulty in predicting the substrate specificity of MFS transporters

from sequence alone makes the identification of potential resistance genes from environmental

microbiomes difficult. This is illustrated by the contrast between the apparent clustering of the

TET1 and TET13 MFS pumps with amphenicol efflux and TET3 with mixed macrolide,

tetracycline, and other antimicrobial efflux (**Figure 2B**) and their respective lack of biologically

relevant resistance to chloramphenicol (**Figure 3CD**) or azithromycin (**Figure 3EF**). Our work

highlights the important role that functional metagenomic selections can play in filling this type of

knowledge gap.

Our previous work[8,52] and those of others[41,53] suggests that streptothricin resistance is

common across soil microbiomes and we chose to explore if this is the case for the human gut

microbiome as well. The ZymoBIOMICS fecal standard, the source of out stool functional

metagenomic library input DNA, has already been sequenced and annotated for known

antibiotic resistance genes by the Zymo corporation, but the data associated with this product

does not list any known streptothricin resistance genes. Our ability to capture a sequence-novel

(**Figure 6**) streptothricin acetyltransferase (termed SatB) from even the modestly sized stool

functional metagenomic library suggests that streptothricin resistance may be a feature of

animal-associated microbiomes as well. Expression of the *satB* gene in *E. coli* conferred a very

high level of nourseothricin resistance (**Figure 5**) and we are planning on testing what effect the

extended N-terminal chain of SatB (**Figure 6A and C**) might have on its enzymatic activity and

specificity. The presence of this resistance element in the human gut microbiome has important

implications for the proposed future use of streptothricin antibiotics in human medicine[54,55] and

suggests that analogs that are not substrates for acetyltransferase enzymes should be explored

before clinical adoption of this antimicrobial class[52].

568   While we see the opening of amplification-free low biomass functional metagenomic

libraries as a significant advancement, limitations to functional metagenomics remain. First,

functional metagenomic libraries largely rely on *E. coli* cells as the host organism for the

functional metagenomic library and therefore is subject to potential limits on what metagenomic

genes can be successfully expressed by this host[56]. In theory, this can be surmounted using

alternative hosts[57–59], and the added efficiency of META assembly may be important when using

strains with lower transformation efficiency than commercial *E. coli* strains. Second, there are

potentially diminishing returns to preparing smaller functional metagenomic libraries, although

the libraries prepared here were shown to still be useful by their capture of new antibiotic

resistance genes even without covering more than a fraction of the genetic diversity found in

many microbiomes. Tagmentation reactions are regularly performed on metagenomic DNA

samples with 5 ng or less mass[12,60,61] suggesting that the limit for functional metagenomic

libraries could be pushed even further.

581   In conclusion, while high-throughput DNA sequencing has helped circumvent the great

plate count anomaly and has revolutionized microbiome research, methods for accurately

annotating novel genes have not managed to keep pace. Machine-learning tools will likely be

able to help with this issue but will also require large amounts of high-quality sequence-function

correlation data. Functional metagenomics is well placed to supply this data and, in the

meantime, can make significant strides in discovering novel functions from known and unknown

genes. The fulfillment of this potential by functional metagenomics can be significantly aided by

the ability to prepare the libraries more efficiently and using less input DNA, a precondition that

we demonstrate is reached here.

Acknowledgements

CRediT contributions: Conceptualization – TSC; Data curation –TSC; Formal analysis – HMA, EPB, EF, TSC; Funding acquisition – EMH, TSC; Investigation – HMA, EPB, EF, TSC; Project administration – TSC Resources – FJO EMH, TSC; Supervision – EMH, TSC; Visualization – HMA, EPB, EF, TSC; Writing - original draft – HMA, EPB, TSC; Writing - reviewing and editing – HMA, EPB, FJO, EMH, TSC. See supplemental figure 5 for graphical representation of contributions. We would like to thank Chrissy Cabay, William Van Bonn, and James Clark for their help in acquiring the Shedd Aquarium sample and Jinglin Hu for the metagenomic DNA. This work was supported in part by a grant from the Florida State University Council for Research and Creativity.

References

1. Staley, James T; Konopka, A., and Staley Allan, J.T.K. (1985). Microorganisms in Aquatic and Terrestrial Habitats. Annual review of microbiology *39*, 321–346.

2. Boolchandani, M., D'Souza, A.W., and Dantas, G. (2019). Sequencing-based methods and resources to study antimicrobial resistance. Nature Reviews Genetics *20*, 356–370. https://doi.org/10.1038/s41576-019-0108-4.

3. Dick, G. (2018). Genomic Approaches in Earth and Environmental Sciences (John Wiley & Sons, Ltd) https://doi.org/10.1002/9781118708231.

4. Handelsman, J., Rondon, M.R., Brady, S.F., Clardy, J., and Goodman, R.M. (1998). Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. Chemistry & biology *5*, R245-9. https://doi.org/10.1016/S1074-5521(98)90108-9.

5. Crofts, T.S., Gasparrini, A.J., and Dantas, G. (2017). Next-generation approaches to understand and combat the antibiotic resistome. Nature reviews. Microbiology *15*, 422–434. https://doi.org/10.1038/nrmicro.2017.28.

6. Lam, K.N., Cheng, J., Engel, K., Neufeld, J.D., and Charles, T.C. (2015). Current and future resources for functional metagenomics. Front. Microbiol. *6*. https://doi.org/10.3389/fmicb.2015.01196.

7. Taupp, M., Mewis, K., and Hallam, S.J. (2011). The art and design of functional metagenomic
screens. Current opinion in biotechnology *22*, 465–472.
https://doi.org/10.1016/j.copbio.2011.02.010.

8. Crofts, T.S., McFarland, A.G., and Hartmann, E.M. (2021). Mosaic Ends Tagmentation
(META) Assembly for Highly Efficient Construction of Functional Metagenomic Libraries.
mSystems *6*, e0052421. https://doi.org/10.1128/mSystems.00524-21.

9. Mahmud, B., Boolchandani, M., Patel, S., and Dantas, G. (2023). Functional Metagenomics
to Study Antibiotic Resistance. Methods Mol Biol *2601*, 379–401. https://doi.org/10.1007/978-
1-0716-2855-3_21.

10.   Lutz, R., and Bujard, H. (1997). Independent and tight regulation of transcriptional units
in escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic
Acids Research *25*, 1203–1210. https://doi.org/10.1093/nar/25.6.1203.

11.   Crofts, T.S., Sontha, P., King, A.O., Wang, B., Biddy, B.A., Zanolli, N., Gaumnitz, J., and
Dantas, G. (2019). Discovery and Characterization of a Nitroreductase Capable of Conferring
Bacterial Resistance to Chloramphenicol. Cell chemical biology *26*, 559-570.e6.
https://doi.org/10.1016/j.chembiol.2019.01.007.

12.   Picelli, S., Björklund, A.K., Reinius, B., Sagasser, S., Winberg, G., and Sandberg, R.
(2014). Tn5 transposase and tagmentation procedures for massively scaled sequencing
projects. Genome research *24*, 2033–2040. https://doi.org/10.1101/gr.177881.114.

13.   Cao, J., Hu, Y., Liu, F., Wang, Y., Bi, Y., Lv, N., Li, J., Zhu, B., and Gao, G.F. (2020).
Metagenomic analysis reveals the microbiome and resistome in migratory birds. Microbiome
8, 26. https://doi.org/10.1186/s40168-019-0781-8.

14.   Hu, J., Hellgeth, N., Cabay, C., Clark, J., Oliaro, F.J., Van Bonn, W., and Hartmann, E.M.
(2022). Towards understanding microbial degradation of chloroquine in large saltwater
systems. Science of the Total Environment *807*, 150532.
https://doi.org/10.1016/j.scitotenv.2021.150532.

15.   Boolchandani, M., Patel, S., and Dantas, G. (2017). Antibiotics P. Sass, ed. (Springer
New York) https://doi.org/10.1007/978-1-4939-6634-9.

16.   Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O., Madhusoodanan,
N., Kolesnikov, A., and Lopez, R. (2022). Search and sequence analysis tools services from
EMBL-EBI in 2022. Nucleic Acids Research *50*, W276–W279.
https://doi.org/10.1093/nar/gkac240.

17.   Zhu, W., Lomsadze, A., and Borodovsky, M. (2010). Ab initio gene identification in
metagenomic sequences. Nucleic acids research *38*, e132.
https://doi.org/10.1093/nar/gkq275.

18.   Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman,
D.J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search
programs. Nucleic Acids Research *25*, 3389–3402. https://doi.org/10.1093/nar/25.17.3389.

19.   Schaffer, A.A. (2001). Improving the accuracy of PSI-BLAST protein database searches
659   with composition-based statistics and other refinements. Nucleic Acids Research *29*, 2994–
660   3005. https://doi.org/10.1093/nar/29.14.2994.

661   20.   Alcock, B.P., Huynh, W., Chalil, R., Smith, K.W., Raphenya, A.R., Wlodarski, M.A.,
662   Edalatmand, A., Petkau, A., Syed, S.A., Tsang, K.K., et al. (2023). CARD 2023: expanded
663   curation, support for machine learning, and resistome prediction at the Comprehensive
664   Antibiotic Resistance Database. Nucleic Acids Research *51*, D690–D699.
665   https://doi.org/10.1093/nar/gkac920.

666   21.   Alcock, B.P., Raphenya, A.R., Lau, T.T.Y., Tsang, K.K., Bouchard, M., Edalatmand, A.,
667   Huynh, W., Nguyen, A.L.V., Cheng, A.A., Liu, S., et al. (2020). CARD 2020: Antibiotic
668   resistome surveillance with the comprehensive antibiotic resistance database. Nucleic Acids
669   Research *48*, D517–D525. https://doi.org/10.1093/nar/gkz935.

670   22.   Davis, M.W., and Jorgensen, E.M. (2022). ApE, A Plasmid Editor: A Freely Available
671   DNA Manipulation and Visualization Program. Frontiers in Bioinformatics *2*, 1–15.
672   https://doi.org/10.3389/fbinf.2022.818619.

673   23.   Wiegand, I., Hilpert, K., and Hancock, R.E.W. (2008). Agar and broth dilution methods to
674   determine the minimal inhibitory concentration (MIC) of antimicrobial substances. Nature
675   Protocols *3*, 163–175. https://doi.org/10.1038/nprot.2007.521.

676   24.   Krügel, H., Fiedler, G., Smith, C., and Baumberg, S. (1993). Sequence and
677   transcriptional analysis of the nourseothricin acetyltransferase-encoding gene nat1 from
678   Streptomyces noursei. Gene *127*, 127–131. https://doi.org/10.1016/0378-1119(93)90627-F.

679   25.   Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., and Minh, B.Q. (2016). W-IQ-TREE: a
680   fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res *44*, W232–
681   W235. https://doi.org/10.1093/nar/gkw256.

682   26.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., Von
683   Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for
684   Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution *37*, 1530–1534.
685   https://doi.org/10.1093/molbev/msaa015.

686   27.   Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the
687   phylogenetic tree display and annotation tool. Nucleic Acids Research, gkae268.
688   https://doi.org/10.1093/nar/gkae268.

689   28.   Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and Barton, G.J. (2009).
690   Jalview Version 2—a multiple sequence alignment editor and analysis workbench.
691   Bioinformatics *25*, 1189–1191. https://doi.org/10.1093/bioinformatics/btp033.

692   29.   Burckhardt, R.M., and Escalante-Semerena, J.C. (2019). Insights into the Function of
693   the *N*-Acetyltransferase SatA That Detoxifies Streptothricin in *Bacillus subtilis* and *Bacillus
694   anthracis*. Appl Environ Microbiol *85*, e03029-18. https://doi.org/10.1128/AEM.03029-18.

695   30.   Committee, E., Testing, S., Microbiology, C., and Escmid, I.D. (2003). Determination of
696   minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. Clinical
697   Microbiology and Infection *9*, ix–xv. https://doi.org/10.1046/j.1469-0691.2003.00790.x.

698   31.   Kim, G., Lee, S., Levy Karin, E., Kim, H., Moriwaki, Y., Ovchinnikov, S., Steinegger, M.,
699   and Mirdita, M. (2025). Easy and accurate protein structure prediction using ColabFold. Nat
700   Protoc 20, 620–642. https://doi.org/10.1038/s41596-024-01060-5.

701   32.   Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M.
702   (2022). ColabFold: making protein folding accessible to all. Nat Methods 19, 679–682.
703   https://doi.org/10.1038/s41592-022-01488-1.

704   33.   Halavaty, A.S., Wawrzak, Z., Onopriyenko, O., Edwards, A., Savchenko, A., Peterson,
705   S., Anderson, W.F., and Center for Structural Genomics of Infectious Diseases (CSGID)
706   (2011). 1.6 Angstrom resolution crystal structure of putative streptothricin acetyltransferase
707   from Bacillus anthracis str. Ames in complex with acetyl coenzyme A: 3pp9.
708   https://doi.org/10.2210/pdb3pp9/pdb https://doi.org/10.2210/pdb3pp9/pdb.

709   34.   Burckhardt, R.M., and Escalante-Semerena, J.C. (2017). In Bacillus subtilis, the SatA
710   (formerly YyaR) acetyltransferase detoxifies streptothricin via lysine acetylation. Applied and
711   Environmental Microbiology 83, 1–11. https://doi.org/10.1128/AEM.01590-17.

712   35.   Sehnal, D., Bittrich, S., Deshpande, M., Svobodová, R., Berka, K., Bazgier, V., Velankar,
713   S., Burley, S.K., Koča, J., and Rose, A.S. (2021). Mol* Viewer: modern web app for 3D
714   visualization and analysis of large biomolecular structures. Nucleic Acids Research 49,
715   W431–W437. https://doi.org/10.1093/nar/gkab314.

716   36.   Berman, H.M. (2000). The Protein Data Bank. Nucleic Acids Research 28, 235–242.
717   https://doi.org/10.1093/nar/28.1.235.

718   37.   Schar, D., Klein, E.Y., Laxminarayan, R., Gilbert, M., and Van Boeckel, T.P. (2020).
719   Global trends in antimicrobial use in aquaculture. Sci Rep 10, 21878.
720   https://doi.org/10.1038/s41598-020-78849-3.

721   38.   Bondad-Reantaso, M.G., MacKinnon, B., Karunasagar, I., Fridman, S., Alday-Sanz, V.,
722   Brun, E., Le Groumellec, M., Li, A., Surachetpong, W., Karunasagar, I., et al. (2023). Review
723   of alternatives to antibiotic use in aquaculture. Reviews in Aquaculture 15, 1421–1451.
724   https://doi.org/10.1111/raq.12786.

725   39.   Biehl, L.M., Garzetti, D., Farowski, F., Ring, D., Koeppel, M.B., Rohde, H., Schafhausen,
726   P., Stecher, B., and Vehreschild, M.J.G.T.G.T. (2019). Usability of rectal swabs for
727   microbiome sampling in a cohort study of hematological and oncological patients. PLOS ONE
728   14, e0215428. https://doi.org/10.1371/journal.pone.0215428.

729   40.   Yang, W., Luyten, Y., Reister, E., Mangelson, H., Sisson, Z., Auch, B., Liachko, I.,
730   Roberts, R.J., and Ettwiller, L. (2025). Proxi-RIMS-seq2 applied to native microbiomes
731   uncovers hundreds of known and novel m5C methyltransferase specificities. Nucleic Acids
732   Research 53, gkaf226. https://doi.org/10.1093/nar/gkaf226.

733   41.   Cox, G., Sieron, A., King, A.M., De Pascale, G., Pawlowski, A.C., Koteva, K., and Wright,
734   G.D. (2017). A Common Platform for Antibiotic Dereplication and Adjuvant Discovery. Cell
735   chemical biology 24, 98–109. https://doi.org/10.1016/j.chembiol.2016.11.011.

42.   van der Helm, E., Imamovic, L., Hashim Ellabaan, M.M., van Schaik, W., Koza, A., and
Sommer, M.O.A. (2017). Rapid resistome mapping using nanopore sequencing. Nucleic
acids research *45*, e61. https://doi.org/10.1093/nar/gkw1328.

43.   Neufeld, J.D., Chen, Y., Dumont, M.G., and Murrell, J.C. (2008). Marine methylotrophs
revealed by stable-isotope probing, multiple displacement amplification and metagenomics.
Environmental Microbiology *10*, 1526–1535. https://doi.org/10.1111/j.1462-
2920.2008.01568.x.

44.   Verastegui, Y., Cheng, J., Engel, K., Kolczynski, D., Mortimer, S., Lavigne, J., Montalibet,
J., Romantsov, T., Hall, M., McConkey, B.J., et al. (2014). Multisubstrate Isotope Labeling and
Metagenomic Analysis of Active Soil Bacterial Communities. mBio *5*, e01157-14.
https://doi.org/10.1128/mBio.01157-14.

45.   Sabina, J., and Leamon, J.H. (2015). Bias in Whole Genome Amplification: Causes and
Considerations. In Whole Genome Amplification Methods in Molecular Biology., T. Kroneis,
ed. (Springer New York), pp. 15–41. https://doi.org/10.1007/978-1-4939-2990-0_2.

46.   Salter, S.J., Cox, M.J., Turek, E.M., Calus, S.T., Cookson, W.O., Moffatt, M.F., Turner, P.,
Parkhill, J., Loman, N.J., and Walker, A.W. (2014). Reagent and laboratory contamination can
critically impact sequence-based microbiome analyses. BMC Biol *12*, 87.
https://doi.org/10.1186/s12915-014-0087-z.

47.   Sommer, M.O.A., Dantas, G., and Church, G.M. (2009). Functional characterization of
the antibiotic resistance reservoir in the human microflora. Science (New York, N.Y.) *325*,
1128–1131. https://doi.org/10.1126/science.1176950.

48.   Hartmann, E.M., Hickey, R., Hsu, T., Betancourt Román, C.M., Chen, J., Schwager, R.,
Kline, J., Brown, G.Z., Halden, R.U., Huttenhower, C., et al. (2016). Antimicrobial Chemicals
Are Associated with Elevated Antibiotic Resistance Genes in the Indoor Dust Microbiome.
Environmental science & technology *50*, 9807–9815.
https://doi.org/10.1021/acs.est.6b00262.

49.   Riesenfeld, C.S., Goodman, R.M., and Handelsman, J. (2004). Uncultured soil bacteria
are a reservoir of new antibiotic resistance genes. Environmental Microbiology *6*, 981–989.
https://doi.org/10.1111/j.1462-2920.2004.00664.x.

50.   Leclercq, S.O., Wang, C., Zhu, Y., Wu, H., Du, X., Liu, Z., and Feng, J. (2016). Diversity
of the Tetracycline Mobilome within a Chinese Pig Manure Sample. Applied and
environmental microbiology *82*, 6454–6462. https://doi.org/10.1128/AEM.01754-16.

51.   Shuai, W., Itzhari, D., Ronen, Z., and Hartmann, E.M. (2023). Mitigation of antimicrobial
resistance genes in greywater treated at household level. Science of The Total Environment
890, 164136. https://doi.org/10.1016/j.scitotenv.2023.164136.

52.   Franck, E., and Crofts, T.S. (2024). History of the streptothricin antibiotics and evidence
for the neglect of the streptothricin resistome. npj Antimicrob Resist *2*, 3.
https://doi.org/10.1038/s44259-023-00020-5.

53.   Baltz, R.H. (2007). Antimicrobials from actinomycetes: Back to the future. Microbe *2*,
125–131.

54.   Morgan, C.E., Kang, Y.-S., Green, A.B., Smith, K.P., Dowgiallo, M.G., Miller, B.C.,
Chiaraviglio, L., Truelson, K.A., Zulauf, K.E., Rodriguez, S., et al. (2023). Streptothricin F is a
bactericidal antibiotic effective against highly drug-resistant gram-negative bacteria that
interacts with the 30S subunit of the 70S ribosome. PLoS Biol *21*, e3002091.
https://doi.org/10.1371/journal.pbio.3002091.

55.   Dowgiallo, M.G., Miller, B.C., Kassu, M., Smith, K.P., Fetigan, A.D., Guo, J.J., Kirby, J.E.,
and Manetsch, R. (2022). The convergent total synthesis and antibacterial profile of the
natural product streptothricin F. Chemical Science *13*, 3447–3453.
https://doi.org/10.1039/d1sc06445b.

56.   Gabor, E.M., Alkema, W.B.L., and Janssen, D.B. (2004). Quantifying the accessibility of
the metagenome by random expression cloning techniques. Environmental Microbiology *6*,
879–886. https://doi.org/10.1111/j.1462-2920.2004.00640.x.

57.   Van Schaik, J., Li, Z., Cheadle, J., and Crook, N. (2023). Engineering the Maize Root
Microbiome: A Rapid MoClo Toolkit and Identification of Potential Bacterial Chassis for
Studying Plant–Microbe Interactions. ACS Synth. Biol. *12*, 3030–3040.
https://doi.org/10.1021/acssynbio.3c00371.

58.   Apjok, G., Számel, M., Christodoulou, C., Seregi, V., Vásárhelyi, B.M., Stirling, T.,
Eszenyi, B., Sári, T., Vidovics, F., Nagrand, E., et al. (2023). Characterization of antibiotic
resistomes by reprogrammed bacteriophage-enabled functional metagenomics in clinical
strains. Nat Microbiol *8*, 410–423. https://doi.org/10.1038/s41564-023-01320-2.

59.   Lam, K.N., Martens, E.C., and Charles, T.C. (2018). Developing a Bacteroides System
for Function-Based Screening of DNA from the Human Gut Microbiome. mSystems *3*, 1–13.
https://doi.org/10.1128/mSystems.00195-17.

60.   Baym, M., Kryazhimskiy, S., Lieberman, T.D., Chung, H., Desai, M.M., and Kishony, R.K.
(2015). Inexpensive multiplexed library preparation for megabase-sized genomes. PLoS
ONE *10*, 1–15. https://doi.org/10.1371/journal.pone.0128036.

61.   Rinke, C., Low, S., Woodcroft, B.J., Raina, J., Skarshewski, A., Le, X.H., Butler, M.K.,
Stocker, R., Seymour, J., Tyson, G.W., et al. (2016). Validation of picogram- and femtogram-
input DNA libraries for microscale metagenomics. PeerJ *4*, e2486.
https://doi.org/10.7717/peerj.2486.

806