Title: Longitudinal trajectories of brain development from infancy to school age and their relationship to literacy development

Authors: Ted K. Turesky[1], Elizabeth Escalante[1], Megan Loh[1], Nadine Gaab[1,2]

Affiliations
[1]Harvard Graduate School of Education, Cambridge, MA
[2]Harvard Medical School, Boston, MA

Correspondence
Ted K. Turesky
50 Church Street
Cambridge, MA 02139
Theodore_Turesky@gse.harvard.edu

Preprint URL: https://doi.org/10.1101/2024.06.29.601366

Classifications: Biological Sciences/Neuroscience

Keywords: brain, development, infant, literacy, longitudinal, MRI

Competing interests: The authors do not have any competing interests to report.

Abstract

Reading is one of the most complex skills that we utilize daily, and it involves the early development and interaction of various lower-level subskills, including phonological processing and oral language. These subskills recruit brain structures, which begin to develop long before the skill manifests and exhibit rapid development during infancy. However, how longitudinal trajectories of early brain development in these structures support long-term acquisition of literacy subskills and subsequent reading is unclear. Children underwent structural and diffusion MRI scanning at multiple timepoints between infancy and second grade and were tested for literacy subskills in preschool and decoding and word reading in early elementary school. We developed and implemented a reproducible pipeline to generate longitudinal trajectories of early brain development to examine associations between these trajectories and literacy (sub)skills. Furthermore, we examined whether familial risk of reading difficulty and children's home literacy environments, two common literacy-related covariates, influenced those trajectories. Results showed that individual differences in curve features (e.g., intercepts and slopes) for longitudinal trajectories of volumetric, surface-based, and white matter organization measures were linked directly to phonological processing and indirectly to first-grade decoding and word reading skills via phonological processing. Altogether, these findings suggest that the brain bases of phonological processing, previously identified as the strongest behavioral predictor of reading and decoding skills, may already begin to develop by birth but undergo further refinement between infancy and preschool. The present study underscores the importance of considering academic skill acquisition from the very beginning of life.

Significance Statement

Reading is crucial for academic, vocational, and health outcomes, but acquiring proficient reading skills is a protracted developmental process involving lower-level subskills and brain structures that undergo rapid development starting in infancy. We examined how longitudinal trajectories of early brain development support long-term acquisition of reading using a reproducible pipeline we developed specifically for infant-to-school-age longitudinal MRI data. Findings suggest that the brain bases of reading-related skills begin to develop by birth but continue building between infancy and preschool. This study emphasizes the importance of considering academic skill acquisition as a dynamic process preceding the emergence of the skill, and it offers a roadmap for future studies to examine relationships between early brain development and academic skill acquisition.

1. Introduction

"…the best way to determine how a child learns is to follow them closely while they are learning." (1, 2).

Reading acquisition is a multifactorial, developmental process that begins long before the skill manifests. Behaviorally, the acquisition of reading necessitates the acquisition and complex interplay of lower-level literacy subskills. In turn, these subskills represent waypoint products of brain development that began in utero. Therefore, understanding how reading skill emerges requires examining its behavioral subskills and the developmental trajectories of the brain areas subserving it starting from the very beginnings of life.

Literacy development represents a model process through which to examine academic skill acquisition, because literacy develops hierarchically, with lower-level "subskills" interacting and driving the emergence of higher-level academic skills (e.g., 3). For instance, phonological processing, which refers to the ability to detect, understand, and manipulate speech sounds (4–7), is the most consistent predictor of subsequent decoding and word reading (4, 8–10). Meanwhile, various studies have shown that both word reading and oral language skills, which encompass abilities supporting listening comprehension (11; e.g., vocabulary and syntactic knowledge), are important subskills for reading comprehension (12). However, these subskills themselves have protracted development and the brain structures that support them begin developing long before they manifest, with the most rapid development transpiring perinatally (13, 14).

Several cross-sectional and prospective studies have been conducted linking performance of these key literacy subskills to brain architecture. For instance, cross-sectional studies in preschoolers and kindergarteners have shown that phonological processing is associated with brain structure and function in left temporoparietal, occipitotemporal, and inferior frontal regions and tracts (15–20). This set of regions has also been linked with preschool and kindergarten oral language skills in some studies (21–23), but other studies have not shown associations (17, 24). Looking to early development, infant brain function and white matter organization have been shown to relate prospectively to (pre)school-age literacy outcomes (25–29), including phonological processing and oral language skills (30–32). Taken together, these studies suggest that certain brain areas may be important for academic skill *performance* at a particular time during development or as 'neural scaffolding' that supports subsequent language development (33). However, recent work in school-age children using large-scale, multi-site cross-sectional datasets showed little evidence for stable associations between individual differences in white matter organization and reading performance. Rather, using a separate

longitudinal dataset, investigators found associations between slopes of white matter growth and
reading gains, suggesting a dynamic interplay between brain development and learning to read
(34). Moreover, the brain itself is a dynamic system that undergoes its most rapid development
during infancy and early childhood (13, 14), making cross-sectional and prospective designs
suboptimal to capture individual differences in early skill *acquisition* (2) As such, individual
longitudinal trajectories, which capture heterogenous rates of brain and skill development
between infancy and second grade, are needed to examine the acquisition of (sub)skills important
for literacy.

Therefore, the overall goal of the current study is to examine the relationship between
longitudinal trajectories of early brain development and acquisition of reading-related subskills,
which we undertook in four objectives. The first objective was to generate individual longitudinal
trajectories of early brain development spanning infancy to school age. Prior longitudinal studies
have examined trajectories of early brain development but only up to the second year of life (e.g.,
35), while others have examined developmental trajectories of brain structure spanning infancy
to school age but only by harmonizing cross-sectional and longitudinal datasets acquired and
processed with varying methods (e.g., 36). However, to our knowledge, no purely longitudinal
studies have mapped the development of brain structure, including white matter organization,
from infancy to school age, using a consistent processing pipeline and one that is appropriate for
early development. Indeed, methodological challenges associated with infant MRI data have
restricted the number and breadth of longitudinal studies in the early developmental period. This
is not only because acquiring and processing infant MRI data requires specialized procedures
and tools to generate accurate brain estimates (37–42), but also because the use of different
procedures and tools for different developmental stages can introduce bias in developmental
analyses. However, the alternative—to use identical procedures and tools for all developmental
stages—would likely cause estimates to vary in accuracy across ages and could introduce
spurious effects and/or obscure true effects in developmental comparisons (43). Some infant-
specific tools offer age-specific adjustments (e.g., different sets of templates) within the first two
years of life, and the small number of studies that have examined longitudinal trajectories of brain
development in the first two years have opted for a balanced approach, with age-appropriate
adjustments to limited processing steps in a way that does not require fundamentally different
techniques for different ages (e.g., additional sequences or different smoothing kernels; (35, 44).
However, methods to accommodate a wider early developmental range are lacking.

In the current study, we extend this work by developing reproducible pipelines for
generating longitudinal trajectories of volumetric, surface-based, and white matter organization

brain measures from infancy to school age. To do this, we leveraged both infant-specific and
standard MRI procedures and tools. We then compared among several candidate linear and
nonlinear mixed effects models, varying according to function (e.g., linear, logarithmic) and
random parameters (e.g., intercepts alone versus intercepts and slopes), to identify the model
with the most parsimonious fit (45). In general, prior longitudinal studies have reported rapid
growth at birth that tapers with age (35, 46–48), except for cortical thickness, which peaks
between ages one and two years (49), suggesting logarithmic functions might provide a better fit
compared with other functions for most measures.

Our second objective was to examine the association of individual differences in early
brain development to long-term literacy development. As such, we extracted curve features (e.g.,
intercepts and slopes) from individual longitudinal trajectories for brain areas and tracts previously
shown to relate to reading-related skill development in longitudinal studies in older children (34,
50–59) or prospective studies in infants (30–32). We then tested these curve features for
correlations with preschool/early kindergarten phonological processing, a key literacy subskill.
Based on converging evidence from prior neuroimaging (30–32) and genetics studies (60, 61),
we hypothesized that phonological processing skill would relate to curve intercepts (i.e., brain
estimates at birth) and slopes (i.e., rate of brain development). If the foundations of literacy
development are largely present at birth and stable across early development, then we expect
most of these brain-behavior associations to be with curve intercepts, whereas if the brain
development subserving literacy subskills is more protracted or dynamic, as suggested by Roy
and colleagues (34), then we expect brain-behavior associations to be with curve slopes.

Our third objective was to examine the roles of risk factors related to literacy skills in
shaping longitudinal trajectories of early brain development. Reading difficulty is heritable, as 40-
60% of children with a familial risk (e.g., first-degree relative with a history) of reading difficulty
(FHD+) themselves develop reading difficulty (62, 63) Brain imaging studies of FHD+
preschoolers show reduced gray matter volume, activation, and fractional anisotropy in left
occipito-temporal and temporo-parietal regions (16–18) and tracts (57) compared with FHD-
preschoolers; these reductions overlap with those observed in children with reading difficulty (20,
64–70), suggesting that the phenotypes characteristic of reading difficulty manifest in some
children before the start of formal reading instruction. Concordantly, similar alterations have been
observed earlier in development, where FHD+ infants exhibited lower fractional anisotropy in left
arcuate fasciculus compared with FHD- infants (71), distinguishable patterns of functional
connectivity in left fusiform gyrus as shown by a support vector machine classifier (72), and
alterations in neural responses to basic speech sounds as measured by event-related potentials

(73–75). Possibly underlying this heritability, work in genetics, including recent large-scale

genome-wide studies, has shown that reading-related skills (60) and reading difficulty (76, 77) are

associated with variation in genes involved in early developmental processes, including

neurogenesis and axon guidance (78).

Another risk factor repeatedly shown to affect children's literacy skills is the home literacy

environment, which includes caregiver–child shared reading and reading-related resources, in

preschool literacy skills (79–84) and in infancy/toddlerhood (83, 85–90). Recent work has also

identified links between the home literacy environment and brain architecture in

preschoolers/kindergarteners (21, 24, 91–94) and infants (95, 96). Overall, these studies offer

strong evidence that genetic and environmental factors related to the development of reading-

related skills are likely to affect longitudinal trajectories of brain development starting perinatally.

Therefore, we hypothesized that the FHD status and home literacy environment would

significantly contribute, as covariates, to longitudinal trajectories of brain development in left

hemisphere temporo-parietal, occipito-temporal, and inferior frontal regions and tracts.

Our fourth objective was to broaden the scope of our examination of early brain-literacy

associations to other literacy-related subskills and subsequent reading skills. Reading acquisition

is a hierarchical process in which multiple distinct but interacting subskills (including but not limited

to phonological processing) converge to effect higher-order skills. First, we tested the specificity

of associations with phonological processing by examining another subskill important for reading

comprehension, oral language skill. Second, we tested whether phonological processing

mediated relationships between curve features of early brain development and subsequent

measures of decoding and word reading skill. Taken altogether, findings from this study will inform

our understanding of how early brain development contributes to later reading-related skill

acquisition.

2. Methods

2.1. Participants

Children examined in this study were obtained from two longitudinal cohorts: the New England dataset and the Calgary open-source dataset (https://osf.io/axz5r/). Only children with high-quality (i.e., post-QC, please see below) MRI data from at least two timepoints between birth and early adolescence were included. Due to the multi-modal nature of the study, varying numbers of longitudinal datasets were used for structural (n = 98 with 276 observations) versus diffusion analyses (n = 128 with 396 observations). For the New England cohort, family history (i.e., first-degree relative with a history) of reading difficulty was also examined and present in 40 of 80 children. For children ages ≤ 24 months, demographic information was provided by caregivers. Please see Table 1 for overall demographic details, Supplementary Table 1 for demographic details by modality, and Supplementary Methods for participant information, including race demographics, unique to each cohort. This study was approved by the Institutional Review Boards of Boston Children's Hospital (IRB-P00023182), Harvard University (IRB21-0916), and the University of Calgary Conjoint Health Research Ethics Board (REB13-0020). Participants' parents gave informed consent, and children gave verbal assent if over 25 months old. Data examined here partially overlap with brain images analyzed in previous studies examining infant brain development (30–32, 36, 48, 54, 71, 72, 96–102).

Table 1. Participant demographics		
General information	Numbers participants | observations	137 | 441
	Number observations per participant	3 ± 1
	Age at literacy-related subskill and cognitive testing (months)	63 ± 5.3
	Age at decoding/word reading testing (months)	82 ± 6.4
Covariates	Biological sex (F/M)	73/64
	Maternal education (years)	17 ± 2.1
	Cohort ([New England]/Calgary)	80/57
	Family history of reading difficulty (+/-)	40/40
	Home literacy environment (a.u.)	0.037 ± 0.41
Literacy-related subskills	Phonological processing standard score	106 ± 14
	Oral language standard score	113 ± 13
Decoding/word reading	Word attack standard score	112 ± 14
	Word identification standard score	110 ± 17
Cognitive abilities	Nonverbal general cognitive ability	106 ± 13

2.2. Environmental variables

Socioeconomic status was measured with maternal education, consistent with prior brain imaging

studies on socioeconomic status (103–108). Maternal education measures were collected during

each timepoint, although inter-time-point variability was low, and these were averaged to generate

one socioeconomic status measure across the developmental window. Rather than using ordinal

coding, years of education as a continuous measure ranging from 12 to 20 years were used.

For the New England cohort, parents also completed at each time point questionnaires

relating to children's home literacy environments home literacy environment, which includes the

extent of parent-child shared reading and access to reading-related resources (109). Responses

were indicated using ordinal scales ranging from 1 to 6. As responses were non-normally

distributed ($p < 0.05$ according to the Shapiro-Wilk normality test; Supplementary Figure 1),

except for "Time read to per week" for all timepoints and "Frequency with which family members

share rhymes or jokes with the child" for one timepoint, they were normalized and then averaged

at each timepoint according to procedure used previously (30). As with maternal education, home

literacy environment estimates, which exhibited low inter-timepoint variability, were averaged to

generate one estimate per individual across the developmental window. These overall home

literacy environment estimates, which were normally distributed (Shapiro-Wilk $W = 0.99$, $p > 0.05$;

Supplementary Figure 2), were used in later statistical analyses.

2.3. Literacy and cognitive measures

Two literacy subskills were administered to children prior to the beginning of formal reading

instruction: phonological processing and oral language. These constructs were selected as

representative subskills supporting literacy development (3, 4, 8–10, 12, 110); however, they

constitute a small subset of literacy-related measures collected for this cohort. No outliers were

detected using the *isoutlier* function in MATLAB, which sets an outlier threshold at three scaled

median absolute deviations from the median.

Phonological processing was measured in both New England and Calgary cohorts. For

the New England cohort, the *phonological processing* composite was estimated from three

subtests from the WJ-IV Tests of Cognitive Abilities: word access, word fluency, and substitution

(111). Word access measures phonetic coding by asking children to identify words containing

certain sounds. Word fluency measures speed of lexical access by asking children to name as

many words as possible beginning with a certain sound in one minute. Substitution measures

children's ability to produce a new word by replacing one sound from a provided word with another

sound. Children in the Calgary cohort were administered the phonological processing subtest of

the NEPSY-II, which measures phonemic awareness (112). Unlike the New England cohort, each
child from the Calgary cohort completed this subtest multiple times. To harmonize phonological
processing scores across the two datasets, we used NEPSY-II scores from when the child was
closest in age to the average age at which the New England cohort completed the WJ-IV (64
months) and not earlier than 50 months of age.

The composite oral language, which was only measured in the New England cohort, was
estimated from two subtests from the WJ-IV Tests of Oral Language: picture vocabulary and oral
comprehension (113). Picture vocabulary measures lexical knowledge by asking children to
specify a picture corresponding to a given word or naming an object. Oral comprehension
measures oral listening, vocabulary, and reasoning by asking children to identify missing words
from short passages. All assessments were administered and double-scored by testers trained
by a clinical psychologist and then raw scores were converted to standard scores.

In addition, we measured decoding and word reading with two untimed subtests from the
Woodcock Reading Mastery Tests III (114): word attack and word identification. For word attack,
children were presented with pseudowords that they needed to *decode* using phonological
abilities. For word identification, children were presented with individual real words that they
needed to *read*. Word attack and word identification subtests were administered at the beginning
of formal reading instruction.

Lastly, we measured children's nonverbal general cognitive ability at preschool/early
kindergarten-age using the Matrix Reasoning subtest of the Kaufman Brief Intelligence Test: 2nd
Edition (KBIT-2, (115)). Herein, children were asked to identify the piece missing from a matrix of
visual images.

All raw estimates for each of the WJ-IV, NEPSY-II, and WRMT subtests were non-
normally distributed ($p < 0.05$ according to the Shapiro-Wilk normality test), except for
preschool/early kindergarten picture vocabulary and late kindergarten/grade 1 word attack
(Supplementary Figure 3). All standardized (composite) estimates used in subsequent analyses
were normally distributed ($p > 0.05$; Supplementary Figure 4).

2.4. MRI data acquisition and processing
All data were acquired on a 3.0 T scanner with a 32-channel head coil. Please see Supplementary
Methods for cohort-specific acquisition parameters.

The rapid brain growth transpiring immediately after birth posed serious challenges for structural
MRI processing because standard methods that are optimal for older children are suboptimal for

infants and vice versa (43). To circumvent bias associated with choosing a single pipeline for

multiple developmental stages, we implemented pipelines that are age-appropriate but not

fundamentally different for each developmental stage.

2.4.1. Structural MRI processing and quality control

Raw magnetization-prepared rapid gradient-echo (MPRAGE) images were visually inspected for

artifacts by a trained rater and scored as "fail," "check," or "pass" (116). Only images scored as

"check" or "pass" underwent image processing. The standard, unmodified FreeSurfer v7.3

(https://surfer.nmr.mgh.harvard.edu/) "recon" pipeline was used for brains > 50 months.

Processing procedures for brains < 50 months were similar in concept to those described in (117).

Namely, Infant FreeSurfer (for brains ≤ 24 months) or standard FreeSurfer (for brains 25 to 50

months) was used to extract the brain from the skull, correct for intensity inhomogeneity, and

segment MPRAGE images by tissue class (gray matter, white matter, cerebrospinal fluid) and

subcortical brain region (40); https://surfer.nmr.mgh.harvard.edu/fswiki/infantFS). To improve

tissue classification accuracy, MPRAGE images were submitted in parallel to iBEATv2.0 Docker

1.0.0 (118–120); https://github.com/iBEAT-V2/iBEAT-V2.0-Docker), has been validated for birth

to age six years (118). Resulting segmentations from each software package were then

hybridized using in-house MATLAB code that combined the cortical pial and white matter

boundaries labelled by iBEATv2.0 with the subcortical parcellations of Infant FreeSurfer (brains ≤

24 months) or standard Freesurfer (brains 25 to 50 months), effectively relabeling cortical gray

and white matter in the FreeSurfer-style segmentation; for details, please see the Supplementary

Methods. Structural processing was finalized by submitting these FreeSurfer-style hybrid

segmentations to a modified version of the standard FreeSurfer v7.3 "recon" pipeline, which, for

brains ≤ 24 months, incorporated elements from Infant FreeSurfer. For a schematic of the

structural processing pipeline, please see Supplementary Figure 5.

Resulting white and pial surfaces were visually inspected by two (brains > 50 months) or

three (brains ≤ 24 months) trained raters on a 3-point scale (0, 1, and 2) and datasets with average

ratings > 1.5 were retained for subsequent analyses. Finally, parcellations were visualized to

ensure accuracy of anatomical labels. For reproducibility purposes, we did not perform manual

editing to correct tissue mislabeling in the remaining images; however, structural measures from

edited and unedited pediatric brain images processed with FreeSurfer have been shown to be

highly correlated (121), including in children (122). Measures of gray and white matter volume,

surface area, cortical thickness, and mean curvature, generated in the final FreeSurfer steps,

were extracted from 8 a priori left hemisphere regions delineated with the Desikan-Killiany atlas—

banks of the superior temporal sulcus, fusiform gyrus, inferior parietal lobule, middle temporal

gyrus, pars opercularis, pars triangularis, superior temporal gyrus, and supramarginal gyrus—

based on their reported involvement in reading-related subskills (15–23, 30–32, 54, 60; for

reviews, please see 123, Table 3 and 124).

2.4.2. Diffusion MRI processing and quality control

Preprocessed and tractography for all diffusion-weighted image (DWI) data, regardless of age,

were performed with MRtrix3 based on the pipeline established for the Developing Human

Connectome Project (41, 125). DWI data were first denoised using Marchenko–Pastur principal

component analysis (126–128) and then corrected for susceptibility distortions, eddy currents,

motion, and intensity inhomogeneity using FSL's topup and eddy (with slice-to-volume correction)

functions (129–133), and Advanced Normalization Tools (ANTs) N4 bias correction tool (134).

Subsequently, three tissue response functions for spherical deconvolution were estimated using

the Dhollander algorithm, a 0.1 fractional anisotropy threshold, and eight maximum harmonic

degrees (135). Fiber orientation densities (FODs) were computed with multi-shell, multi-tissue

constrained spherical deconvolution (136, 137) and then normalized using multi-tissue informed

log-domain intensity normalization.

Two million streamlines were tracked from the normalized FOD maps using the

Anatomically Constrained Tractography (ACT) technique. Importantly, ACT has been shown to

greatly improve tractography, but it relies on accurate tissue segmentations, which are typically

challenging to generate with infant brain data. Using the hybrid segmentations generated for

brains ≤ 50 months (please see above) circumvented this challenge. Thus, FreeSurfer-style

hybrid segmentations for brains ≤ 50 months and standard FreeSurfer segmentations for brains

> 50 months were registered to the preprocessed DWI images using ANTs and then converted

to five-tissue-type images for ACT. The remaining whole-brain tractography parameters included

seeding at the gray/white matter boundary and tracking with the iFOD1 probabilistic algorithm;

step size, minimum and maximum length, and maximum step angle were set to default (138, 139).

Resulting whole-brain tractography was then submitted to the open-source instantiation

of Automated Fiber Quantification (AFQ; (140) for waypoint- and probabilistic-atlas-based fiber

tract segmentation. The standard pyAFQ pipeline was used for brains > 24 months (141), whereas

pyBabyAFQ was used for brains ≤ 24 months (142). Please see Supplementary Methods for a

summary of differences between the two AFQ instantiations. Next, tracts were resampled to 100

equidistant nodes, and diffusion properties (fractional anisotropy; mean diffusivity) were quantified

for each node for the left hemisphere tracts of interest—arcuate fasciculus, superior longitudinal

fasciculus, and inferior longitudinal fasciculus. Mean fractional anisotropy and mean diffusivity

values for each tract were used to examine model fits (please see section 2.5.1), whereas a more

fine-grained, node-based approach was taken for brain-behavior analyses (please see section

2.5.2). For the latter, to better align tract cores across participants, five nodes on either end were

removed, reducing the total number of nodes per tract to 90. Finally, tracts of interest were visually

inspected by two trained raters on a 3-point scale (0, 1, and 2) and datasets with average ratings

≥ 1 were retained for subsequent analyses. For a schematic of the diffusion processing pipeline,

please see Supplementary Figure 6.

2.5. Statistical analyses

An overview of statistical operations, all of which were performed in RStudio version 4.2.2, can

be found in Supplementary Figure 7.

2.5.1. Longitudinal trajectory estimation

Prior to modeling, estimates of brain structure and white matter organization underwent a final

quality control procedure to remove estimates for brain areas/tracts of interest if, for brains > 25

months, they were preceded or followed by ≥ 10% annual change (positive or negative) in any

measure and for brains ≤ 24 months, inter-observation changes were negative (or positive for

mean diffusivity). To mitigate data loss, we next identified which observation—earlier or later—

was more likely to be inaccurate, using an outlier detection procedure (described in the

Supplementary Methods). All remaining participants after these procedures had multiple

observations.

To generate longitudinal trajectories (i.e., growth curves), we next submitted cleaned,

longitudinal structural and white matter organization estimates to linear mixed effects models

using linear, logarithmic, and quadratic functions from the R 'lme4' package. Intercepts and slopes

were modeled as fixed and random effects, and covariates for biological sex, socioeconomic

status, and cohort (New England or Calgary) were entered as fixed effects. Vijayakumar and

colleagues recommends comparing model fits quantitatively (45); consequently, we computed

Bayesian Information Criterion (BIC) metrics and the model (i.e., function and number of random

terms) that provided the best fit (i.e., lowest BIC value) was selected for subsequent analyses.

Correlations between intercepts and slopes were tested to ensure associations between random

terms were minimized. Most curve features were normally distributed according to the Shapiro-

Wilk normality test ($p > 0.05$) and histograms depicting their distributions are provided in

Supplementary Figures 9-15.

2.5.2. Associations between longitudinal trajectories of early brain development and phonological
processing

Next, individual-level curve features (i.e., intercepts and slopes) were extracted and tested for
correlations (Pearson) with literacy subskills. The extracted curve features represent individual
variation in intercepts or slopes beyond what would be predicted based on the covariates (e.g.,
biological sex); consequently, we did not include these same covariates in the tests of
correlations. To determine whether curve features of volumetric and surface-based measures
were associated with phonological processing in the full sample after accounting for multiple brain
regions (8 tests), a significance threshold was set to $p_{FDR} < 0.05$ and applied to measures (e.g.,
gray matter volume) separately. As diffusion analyses were performed node-wise (90 nodes), we
corrected for multiple comparisons at $p_{FWE} < 0.05$ using a permutation-based, threshold-free
cluster enhancement method (143), implemented in the permuco package in R (144); corrections
were performed separately for each tract. These methods are similar to those used previously
(24, 96, 108). When significant, individual growth curves were separated into three groups
according to scores on their behavioral assessment with low < 85, 85 ≤ average ≤ 115, and high
> 115, averaged by group, and then plotted.

2.5.3. Sensitivity analyses

We conducted four sensitivity analyses to test the reliability of our results. First, we performed a
replication analysis on gray/white matter volume, surface area, and mean diffusivity with nonlinear
mixed effects models using asymptotic functions from the R 'nlme' package (145), similar to that
described in Alex and colleagues (36); https://github.com/knickmeyer-lab/ORIGINs_ICV-and-
Subcortical-volume-development-in-early-childhood). Intercepts and asymptotes were modeled
as fixed and random effects; rate constants were modeled as fixed effects. Second, instead of
testing correlations between curve features and outcomes, we entered outcomes as main and
interaction terms in linear mixed effects models. Third, adhering to recommendations to report
both raw and TIV-corrected results (45), we recomputed brain-behavior associations for
volumetric and surface-based measures using semipartial correlations (Pearson) with the random
terms from longitudinal modeling with TIV as covariates of no interest. Fourth, we submitted
volumetric and surface-based brain-behavior associations to semipartial correlations (Pearson)
with average (across timepoint) Euler numbers, which quantifies topological defects (146). For
additional details on these sensitivity analyses and Euler quantification, please see the
Supplementary Methods.

2.5.4. Specificity analyses

To determine whether brain-behavior effects were specific to certain brain measures, regions,

curve features, and behavioral outcomes, we performed two specificity analyses. First, we

generated whole-brain maps depicting variability in associations with phonological processing

according to brain measures (e.g., gray matter volume), region, and curve feature. Second, limited

to the New England children, we tested correlations between curve features of brain development

and oral language skill, another reading subskill, and nonverbal general cognitive abilities. For

volumetric and surface-based measures, FDR correction accounted for brain regions and

behavioral measures (24 tests). For diffusion, cluster-level FWE correction accounted for nodes

(90 tests).

2.5.5. Literacy-related factors as fixed effects in models of early brain development

We determined whether literacy-related covariates (i.e., home literacy environment and FHD

status) contributed to the fit of the growth curve in two ways. First, we examined each covariate

as fixed effects in models of brain development for measures and regions whose curve features

related to phonological processing (please see brain-behavior associations in 2.5.2). For brain

structure, a significance threshold of $p_{FDR} < 0.05$ was applied to correct for the multiple brain

development models that fit this criterion. Diffusion analyses, which were performed node-wise

for each tract separately (90 tests), were corrected for multiple comparisons using a lenient cluster

size threshold of 5 nodes, as threshold-free cluster enhancement method was not available for

the statistical tests applied to covariates. Second, we compared brain development models with

and without each covariate using BIC metrics. For diffusion, BIC metrics were obtained using tract

averages (rather than separately on individual nodes). Correction for multiple comparisons was

again set to $p_{FDR} < 0.05$ and applied as described above for brain structure, this time also including

statistics for average diffusion measures.

2.5.6. Indirect effects between longitudinal trajectories of early brain development and word

reading outcomes via literacy subskills

As with previous work (147), indirect effects were tested when literacy subskills were related to

curve features of brain development and decoding and word reading outcomes (after FDR

correction). Formal indirect effects modeling was conducted using the Mediation package in R

with 20,000 bootstrapped samples, and effects were determined significant when the 95%

confidence interval for the average causal mediation effect did not include 0. All mediation models

included covariates for maternal education, home literacy environment, and family history of

reading difficulty. As in the sensitivity analyses, models were run twice, once without controlling

for TIV and once including TIV curve features. Consistent with prior literature (36, 147, 148), no

additional correction for multiple comparisons was applied.

2.6. Data availability

All code used to process and analyze has been made openly available at

https://github.com/TeddyTuresky/Longitudinal-Trajectories-Early-Brain-Development-Language.

The Calgary dataset is also freely available at https://osf.io/axz5r/. All Boston/Cambridge data

used in this study will be made publicly available at upon acceptance of the manuscript.

3. Results

3.1. *Longitudinal trajectories of brain structure and white matter organization in left hemisphere literacy-related brain regions and tracts*

Our first objective was to generate accurate longitudinal trajectories of early brain development in regions and tracts reportedly involved in reading-related subskills (15–23, 30–32, 54, 60; for reviews, please see 123, Table 3 and 124). Accordingly, growth curves for regional estimates of gray and white matter volume, surface area, cortical thickness, and mean curvature and for tract estimates of fractional anisotropy and mean diffusivity were generated from 137 children (F/M = 73/64) with 441 observations. Every child had at least two observations between birth and school age with 77 having at least one observation in infancy (Figure 1; please see Supplementary Figure 16 for age distributions by modality). Linear mixed effects models included individual-level intercepts and slopes as well as covariates for biological sex, socioeconomic status, and cohort. All structural and diffusion measures for all regions and tracts were best fit with logarithmic functions according to Bayesian Information Criterion (Supplementary Table 2), whereby rates of development were steeper perinatally and then slowed across the first ten years following birth (Figure 2; Supplementary Figures 17-22).

Figure 1. Age distribution of longitudinal dataset from infancy to early adolescence. All children had structural and/or diffusion MRI data from at least two observations (dots). Blue, New England cohort; orange, Calgary cohort.

Figure 2. Average longitudinal trajectories from infancy to early adolescence by measure. Raw
estimates for each brain measure and region were submitted to linear mixed effects models using
a logarithmic function. Individual growth curves predicted by this model were averaged to show
the overall longitudinal trajectory of the sample for each volumetric/surface-based (blue lines) and
diffusion (green lines) measure. Absolute brain estimates were then converted to percent change
values to visualize all brain measures along a single axis. For growth curves for separate brain
regions and tracts, please see Supplementary Figures 17-22.

3.2. Associations between growth curve features and phonological processing

Our second objective was to examine whether brain development affects literacy development.

Consequently, curve features (i.e., intercepts and slopes) were extracted from individual-level

growth curves and tested for correlations (Pearson) with preschool/early kindergarten

phonological processing scores, a subskill critical for literacy (3, 8).

For volumetric and surface-based measures, curve features of gray/white matter volume

and surface area in several left hemisphere regions exhibited associations with phonological

processing. Specifically, greater phonological processing was associated with (a) greater

intercepts of gray matter volume in the left banks of the superior temporal sulcus, (b) greater

slopes of white matter volume in left occipitotemporal, temporoparietal, and inferior frontal

regions, and (c) greater surface area intercepts and slopes in inferior parietal lobule, pars

triangularis, and superior temporal gyrus (Supplementary Figure 23). Next, individual growth

curves corresponding to these measures and regions were separated into three groups according

to phonological processing scores with low < 85, 85 ≤ average ≤ 115, and high > 115, averaged

by group, and then plotted to visualize variation in longitudinal trajectories by phonological

processing score. Most notably, children with lower phonological processing scores tended to

have less gray matter volume in left banks of the superior temporal sulcus at birth but maintained
similar rates of development compared with higher scoring children. At the same time, they had
lower rates of white matter volume growth in left occipitotemporal, temporoparietal, and inferior
frontal regions (Figure 3).

Figure 3. Longitudinal trajectories of brain structure from infancy to early adolescence according
to phonological processing skill in preschool/early kindergarten. Graphs depict average
trajectories for children with low (< 85), average (85 – 115), and high (> 115) standardized
phonological processing scores for measures and regions whose (A) intercepts, (B) slopes, or (C)
both intercepts and slopes correlated with phonological processing (p_{FDR} < 0.05). Correlation
statistics are reported adjacent to their corresponding plots; intercept and slope statistics for
surface area averaged here for visualization purposes but reported separately in Supplementary
Figure 23. As a group, children with low phonological processing in preschool/early kindergarten
tended to have attenuated longitudinal trajectories, either because they began with lower
estimates, as with less gray matter volume in the left banks of the superior temporal gyrus at birth
(upper left graph), or because they had slower rates of development, as with white matter volume
in other left hemisphere brain regions.

Curve features of white matter organization in the left arcuate fasciculus also correlated

with phonological processing outcomes, as greater phonological processing skill was associated

with greater slopes of mean diffusivity (Supplementary Figure 24). When distinguishing

longitudinal trajectories according to literacy subskills (i.e., low < 85, 85 ≤ average ≤ 115, and high

> 115), as with volumetric and surface-based measures above, children with low phonological

processing scores tended to have higher mean diffusivity at birth but greater rates of mean

diffusivity development (i.e., more negative slopes) compared with higher scoring children (Figure

4). No significant associations were observed for the superior or inferior longitudinal fasciculus.

arcuate fasciculus

r = 0.23
p = 1.9x10^{-2}
nodes 6-25

Phonological processing skill: Low   Average   High

Figure 4. Longitudinal trajectories of mean diffusivity from infancy to early adolescence according
to phonological processing skill in preschool/early kindergarten. Graph depicts average
trajectories for children with low (< 85), average (85 – 115), and high (> 115) standardized
phonological processing scores for the left arcuate fasciculus nodes whose slopes correlated with
phonological processing (p_{FWE} < 0.05). Children with low phonological processing tended to
exhibit faster development (i.e., more negative slope) in anterior arcuate fasciculus.

3.3. *Sensitivity Analyses*

We performed four sensitivity analyses to test the robustness of the observed brain-behavior

relationships. For the first sensitivity analysis, we re-fit our volumetric, surface area, and mean

diffusivity estimates with nonlinear mixed effects models using asymptotic functions

(Supplementary Figures 25-28). This alternative model to characterize longitudinal trajectories

was especially important for mean diffusivity findings, where intercepts and slopes from the main

analysis were correlated for many nodes in each tract examined.

We observed that greater phonological processing was still associated with greater

intercepts of gray matter volume development in the left banks of the superior temporal sulcus

and of surface area in left inferior parietal lobule, pars triangularis, and superior temporal gyrus

(Supplementary Table 3A, C). Interestingly, when trajectories were again split according to low (<

85), average (85 – 115), and high (> 115) standardized phonological processing scores, intercepts

were saliently divided, more so than when using linear models (Supplementary Figure 3A,C).

Whereas our main analyses allowed us to examine relationships between brain development and

literacy subskills directly by using a random slopes term, the nonlinear model used in this

sensitivity analysis replaces the random slopes term with a random asymptote term. As such,

relationships between brain development and language outcomes may only be inferred where

brain measures and regions exhibit asymptote-behavior correlations without corresponding

intercept-behavior correlations. Models for middle and superior temporal cortices and

supramarginal gyrus white matter volume did not converge. However, other brain regions whose

white matter volume and surface area slopes correlated positively with phonological processing

in the main analysis also exhibited significant associations between asymptotes and phonological

processing. Effect sizes in were generally smaller for white matter volume and larger for surface

area compared with the main analysis (Supplementary Tables 3B, C; Supplementary Figure

3B,C). Regarding white matter organization, greater phonological processing was associated with

lower mean diffusivity intercepts in left arcuate fasciculus (Supplementary Tables 3D).

Our second sensitivity analysis involved adding phonological processing main and age x

phonological processing interaction terms as covariate analogues to brain-behavior correlations

with intercepts and slopes, respectively. All results for gray and white matter volume and mean

diffusivity present in the main analysis persisted in this sensitivity analysis; however, surface area

effects did not (Supplementary Table 4).

For the third sensitivity analysis, we recomputed brain-behavior associations for

volumetric and surface-based measures using semipartial correlations, controlling for curve

features of the longitudinal trajectory for total intracranial volume (TIV). Relative to the results of

the main analysis, effect sizes were reduced when including curve features of TIV. However,

associations with phonological processing remained significant for intercepts of the banks of the

superior temporal sulcus gray matter and inferior parietal lobule, pars triangularis, and superior

temporal gyrus surface area. Slopes of white matter volume development in inferior parietal lobule

and pars triangularis also remained significant (Supplementary Table 5).

In the fourth sensitivity analysis, we again recomputed brain-behavior associations, this

time controlling for Euler numbers as a reproducible alternative to manual quality control (149).

On average, effect sizes showed no drop relative to the main analysis (r_{avg} = 0.35). All brain-

behavior associations significant in the main analysis remained significant with the inclusion of

Euler numbers, except for the association between phonological processing and slopes of
superior temporal gyrus surface area (Supplementary Table 6).

3.4. *Specificity analyses*

We also performed analyses to determine whether brain-behavior associations were limited to
specific brain measures, regions, and curve features or contingent upon specific behavioral
measures. Whole-brain analyses showed higher brain-behavior effects for white matter volume
slopes compared (numerically) to other measures, especially in left inferior parietal lobule;
however, effects did not appear to be specific to the left hemisphere (Supplementary Figure 32).
In a subset of data, we also tested whether associations were specific to phonological processing,
to other reading subskills, or to cognitive measures in general. Brain-behavior effects with
phonological processing persisted in this subset but similar effects were not observed for oral
language skills or nonverbal general cognitive ability (Supplementary Table 7).

3.5. *Contributions of literacy-related factors to longitudinal trajectories of brain structure and white*
matter organization

Next, we sought to examine whether the observed brain-behavior associations are driven by two
common literacy-related factors: family history of reading difficulty and the home literacy
environment. When modeled as fixed effects, neither constituted significant contributors to the
longitudinal trajectories of early brain development that predicted phonological processing
(Supplementary Table 8). Furthermore, BIC estimates for models with versus without literacy-
related covariates were not significantly different for any brain measures or regions
(Supplementary Table 9).

3.6. *Indirect effects between longitudinal trajectories of early brain development and word*
reading outcomes via phonological processing

Finally, we examined whether literacy subskills mediated the relationship between brain
development curve features (e.g., intercepts and slopes) and decoding and word reading
outcomes. As a prerequisite for mediation, the mediator (i.e., phonological processing) must be
associated with both the predictor (i.e., brain estimate) and outcome (i.e., word reading). We
limited the potential mediator to phonological processing, which related to both decoding ($r = 0.48$;
$p < 0.005$) and word reading ($r = 0.53$; $p < 0.001$), as measured by the Woodcock Reading Mastery
Tests III word attack and word identification subtests (114), because brain-behavior associations
with oral language were few and inconsistent in sensitivity analyses, and we limited the potential

predictors to brain measures and regions/tracts surviving FDR correction in the main analysis.
Indirect effects were reported when the 95% confidence intervals (CI), based on 20,000
bootstrapped samples, for the average causal mediation effect did not include 0 (please see
Methods).

Phonological processing mediated the relationship between brain and decoding and
between brain and word reading for the following measures and regions: intercepts of gray matter
volume development in the left banks of the superior temporal sulcus (Figure 5A). Phonological
processing skill also mediated associations between decoding/word reading and slopes of white
matter volume development and intercepts and slopes of surface area development in left
temporo-parietal and inferior frontal regions (Figure 5B, C; Supplementary Tables 10, 11). Indirect
effect sizes nominally attenuated when controlling for TIV curve features for decoding (average
estimate with TIV = 1.6 x 10^{-2}, average estimate without TIV = 2.0 x 10^{-2}) and word reading
(average estimate with TIV = 2.4 x 10^{-2}, average estimate without TIV = 3.1 x 10^{-2}; Supplementary
Tables 12, 13). Lastly, phonological processing skill mediated associations between slopes of
mean diffusivity development in left arcuate fasciculus and decoding and word reading (Figure
5D).

Figure 5. Phonological processing skill mediates the relationship between early brain
development and decoding and word reading. Indirect effects (filled arrows) were found for
(A) intercepts of gray matter volume in the left banks of the superior temporal sulcus; (B)
slopes of white matter volume and (C) intercepts and slopes of surface area in left temporo-
parietal and inferior frontal regions; and (D) slopes of mean diffusivity in left arcuate fasciculus
(nodes 6-25, green). Note: indirect effects are depicted for surface area slopes only; surface
area intercept effects are reported in Supplementary Figures 10, 11.

4. Discussion

The brain regions and tracts that eventually support decoding and word reading begin to develop long before the skills themselves emerge. Here, we examined the relationship between longitudinal trajectories of early brain development and acquisition of reading-related (sub)skills in four objectives. First, we generated longitudinal trajectories of early brain structure, including white matter organization, from infancy to school age in regions and tracts previously linked to literacy development (30–32, 34, 50–58) using a novel processing and analysis pipeline appropriate for the early developmental period. Findings showed that longitudinal trajectories were best modeled using logarithmic, compared with linear and quadratic, functions. Second, we examined associations between curve features of longitudinal trajectories and a key literacy subskill: phonological processing. Results showed that curve intercepts (i.e., birth brain estimates) of gray matter volume and surface area and curve slopes (i.e., early brain development) of white matter volume, surface area, and mean diffusivity predicted phonological processing measured in preschool/early kindergarten. While effects were robust in hypothesized left temporo-parietal, occipito-temporal, and inferior frontal regions and tracts, specificity analyses suggest that these brain-behavior associations are not limited to these regions. The predominance and magnitude of slope-outcome associations in comparison with intercept-outcome associations suggests a less stable, more dynamic relationship between brain and literacy development (34). Third, we examined whether familial risk of reading difficulty and home literacy environment, two common literacy-related covariates, influenced those trajectories and found that they did not. Fourth, we expanded the scope of our inquiry to long-term literacy development, showing that phonological processing mediated associations between early brain development and decoding and word reading skills between late kindergarten and second grade. Overall, these findings suggest that the neural foundations for the subsequent development of phonological processing may be partially present at birth but are still forming in the years between birth and preschool and eventually support the development of decoding and word reading skills.

Trajectories of early brain development have been generated from longitudinal studies up to the first two years of life (e.g., 35) and from combined cross-sectional and longitudinal datasets spanning infancy to school age (e.g., 36). However, methodological challenges associated with infant MRI and longitudinal designs in general have created a relative vacuum of longitudinal MRI studies spanning infancy to school age (43). The current study fills this gap as a purely longitudinal examination of early structural brain development in which longitudinal trajectories are generated with a pipeline designed for the early developmental period. Overall, longitudinal trajectories of gray and white matter volume, surface area, fractional anisotropy, and mean diffusivity exhibited

rapid postnatal growth that slowed with age, which largely comports with previous early

developmental studies using cross-sectional (150–154) and combined longitudinal and cross-

sectional designs (13, 36, 46, 48, 155, 156). As shown previously (49, 154), the curves for cortical

thickness and mean curvature were generally less steep compared with other brain measures,

both initially following birth and into childhood. Nonetheless, empirical comparisons with linear

and quadratic functions showed that logarithmic models were more parsimonious for all measures

examined.

After generating longitudinal trajectories, we examined how this brain development

influenced literacy development by relating curve features of individual trajectories to reading-

related subskills measured in preschool/early kindergarten as well as decoding and word reading

skills measured between kindergarten and second grade. The finding of direct links to

phonological processing and indirect links to decoding and word reading from curve intercepts of

gray matter volume (i.e., gray matter volume at birth) specifically in the left banks of the superior

temporal sulcus is compelling in the context of a recent large-scale genomics study showing

common genetic influences on surface area in this region and reading-related skills (60), including

from a gene involved in neurogenesis and axon formation (78). Combined with observed

intercept-outcome associations for surface area in other left hemisphere regions and prior work

showing an association between reading difficulty and left temporo-parietal sulcal patterns

determined in utero (157), these findings offer convergent evidence for a mechanistic pathway

through which genetic factors shape the foundations of reading development via prenatal left

temporo-parietal brain development. While this intercept-outcome effect remained robust through

all four sensitivity analyses, this interpretation should be viewed with caution, as specificity

analyses suggest intercept-outcome effects may not be limited to the left banks of the superior

temporal sulcus.

The intercept-outcome findings also support the hypothesis that school-age literacy skill

builds on an early-developing foundation (33) or 'neural scaffold' (30–32). However, the paucity

of intercept-outcome associations in contrast to the multitude and magnitude of slope-outcome

associations suggests that this neural scaffold develops substantially over the first several years

of life. While the mechanisms driving these divergent associations with outcomes will ultimately

require further investigation, it is conceivable that they reflect links to distinct early precursors of

phonological skills. For instance, the intercept-outcome associations may reflect stable relations

with foundational perception skills that develop earlier (e.g., prosody, differentiating phonemes)

and remain necessary for the development of more advanced phonological processing skills,

whereas slope-outcome associations reflect dynamic associations with more advanced

phonological processing skills such as syllable or phoneme deletion or substitution. Overall, the

presence of slope-outcome associations offers insights into the early development of a neural

scaffold for literacy and in doing so, underscores the importance of examining individual

longitudinal trajectories (2), as opposed to cross-sectional and prospective associations.

It was also interesting that most of the slope-outcome findings were with white matter

properties, and nearly all of these remained significant in all sensitivity analyses, with some effects

becoming even more robust (e.g., please see inferior parietal white matter in second sensitivity

analysis). While no prior study has examined the relationship between long-term literacy subskills

and longitudinal trajectories of white matter development beginning in infancy (or early

development of surface-based measures), there is a small corpus of literature that has linked

reading-related skill performance to short-term, school-age developmental changes in left

temporo-parietal white matter volume (158) and organization, most consistently in the left arcuate

fasciculus (34, 56, 57, 158). This is especially compelling when considering that the slope-

outcome relationship we observed with white matter organization was specific to the left arcuate

fasciculus and to phonological processing, which is highly predictive of subsequent word reading

and decoding performance (4, 8–10). Work from genomics might offer further explanation for the

dominance of white matter associations, as genes involved in axon guidance (159), axon

formation (78), and oligodendrocyte maturation (160) have also been linked to reading-related

skills (60) and reading disability (76, 77). Although white matter volume and organization are not

strongly associated and thought to be sensitive to different properties (161), their reliance on

myelination, the primary function of oligodendrocytes, and other axonal properties suggests that

they may represent another mechanism through which genetic factors may shape early brain

development.

Interestingly, specificity analyses showed direct brain-behavior associations with

phonological processing but not with oral language skills or nonverbal general cognitive abilities.

Prior brain imaging studies in preschoolers/kindergarteners comport with this finding as

phonological processing seems to exhibit more consistent associations with brain architecture

(15–20), compared with oral language skills (21–23); c.f., (17, 24). While both subskills are

considered critical for higher-level reading-related skills, they are thought to function on two

distinguishable developmental pathways, whereby phonological processing is more involved in

recognizing and decoding printed words and oral language skills are more essential for the

development of reading comprehension skills (please see Scarborough's Reading Rope; (8). It

should be noted that this framework also fits with our mediation findings, which show indirect

relations between early brain development and decoding and (printed) word reading via

phonological processing. Furthermore, whereas phonological processing is limited in its scope,

as specifically one's ability to recognize and manipulate sounds in a word (4–7), oral language

was measured in the current study as a composite of vocabulary and oral comprehension

subtests, with the latter requiring children to engage with semantic and syntactic cues. As different

brain regions may be more specialized for certain component language skills (e.g., pars

opercularis may be more involved in syntactic processing while pars triangularis and middle

temporal gyrus may be more involved in semantics; (162, 163)), aggregating distinct oral

language skills may have obscured brain-behavior associations. Conversely, it is also conceivable

that oral language acquisition requires additional, moderating factors that were not modeled in

the present study (e.g., social interactions (164) or conversational turn-taking (21)). Nevertheless,

the necessity of phonological processing for literacy development and its relation to early brain

development reported here underscore the importance of examining reading-related subskills at

the very beginnings of life.

Turning to literacy-related covariates, familial history of reading difficulty (FHD) was

hypothesized to influence longitudinal trajectories of early brain development, based on previous

findings showing FHD-related alterations in brain architecture in infancy (71, 72) and preschool

(16, 18, 57). Consequently, the observation that FHD status did not influence longitudinal

trajectories in any regions or tracts examined was initially unexpected. However, it is important to

recognize that a child who has an older sibling or parent with a reading difficulty (FHD+) does not

necessarily have a genetic susceptibility for reading difficulty (165), nor will they necessarily

develop reading challenges given the multifactorial nature of reading difficulties. Rather, FHD

status should be considered one of multiple risk factors that can contribute to long-term literacy

development (166, 167), either through intergenerational transmission of genes or environment

(168). Consistent with this, roughly half of FHD+ children develop typical reading skills (62, 63),

and those who do develop typical reading skills have, as a group, been shown to recruit right- and

inter-hemispheric compensatory pathways (169, 170), a pattern similar to children with reading

difficulty who subsequently show improvements (171). Although the specificity analysis examining

brain-behavior effects in non-a-priori brain regions did not point to literacy-related effects solely in

the left hemisphere, right hemisphere regions and tracts were not thoroughly assayed in the

current study. Therefore, it may behoove future studies with larger sample sizes to examine the

development of right hemisphere regions and tracts in the context of FHD status and to distinguish

FHD+ children who develop typical reading skills from FHD+ children who do not.

The home literacy environment also did not significantly influence longitudinal trajectories

in the regions and tracts examined, despite reported links to brain structure and function in infants

(95, 96) and preschoolers/kindergarteners (21, 24, 91–94). However, with few exceptions (e.g.,

common associations to arcuate fasciculus fractional anisotropy in infants (96) and kindergarten

(24)), specific brain measures and regions/tracts linked to home literacy environment variables

identified at the infant time point were not also identified at the preschool/kindergarten time point,

suggesting that brain-home literacy environment relations may vary across this developmental

window. Future longitudinal studies with more frequent sampling of observations will be needed

to test whether this explanation is accurate. Also, as examination of home literacy was limited to

longitudinal trajectories for brain measures and regions/tracts that were associated with

phonological processing, it is likely that measures and regions/tracts related to the home literacy

environment went untested in the current study.

This study had five main limitations. First, consistent with prior work (36), the models we

fit to early brain development generated smooth longitudinal trajectories. In actuality, it is unlikely

that early brain development transpires as predictably, especially during sensitive and critical

periods (172) or specific learning milestones (2). Although participants in the current study were

sampled over three times on average, which is preferred for modeling growth curves (173) and

more than most longitudinal imaging studies (174), future studies would benefit from increased

sampling, particularly around learning milestones germane to literacy development. Second, the

sample size in the current study was relatively small when compared with the sample sizes used

in multi-site, combined cross-sectional and longitudinal studies (e.g., 13, 36). While sensitivity

analyses for the most part demonstrated the robustness of the results, future studies with larger

sample sizes will be needed to confirm the findings presented here. Third, cortical thickness,

mean curvature, fractional anisotropy, and mean diffusivity exhibited high correlations between

random parameters (i.e., intercept-slope correlations) when modeled with logarithmic functions,

which spurred concerns over the accuracy of the estimated random parameters. For the current

findings, inaccurate random parameters could have generated false negatives for the former three

measures and false positives for mean diffusivity. Consequently, for the former three parameters,

even though they were consistently better fit with logarithmic functions, we re-analyzed intercept-

outcome and slope-outcome associations using random parameters from models with quadratic

functions, which had considerably lower intercept-slope correlations. Despite this, results

remained non-significant. Meanwhile, the first sensitivity analysis addressed concerns for mean

diffusivity by showing that nonlinear mixed effects models using asymptotic functions decoupled

intercepts and slopes while maintaining the significant results observed in the main analysis. It is

also important to note that this limitation does not apply to findings for gray and white matter

volume or surface area. Fourth, we examined somewhat narrowly literacy-related factors that

could contribute to early brain development by only including FHD status and home literacy
environment, and it is likely that factors not included also contribute to the longitudinal trajectories
examined (e.g., teaching quality, educational opportunities, executive functioning skills (166,
175). Fifth, our longitudinal analysis pipeline identifies and removes brain estimates preceding or
following developmental changes that are too steep to occur neuroanatomically and more likely
to emerge from region or tract mislabeling, despite our quality control efforts. Consequently, it is
possible that estimates without steep developmental changes also suffer mislabeling but remain
undetected. Overall, interpretations of findings should be considered in the context of these
limitations.

In conclusion, this study examined associations between longitudinal trajectories of early
brain development beginning in infancy and long-term reading acquisition, specifically literacy
subskills. Longitudinal trajectories were generated using a novel, reproducible pipeline we
designed specifically for examining early brain development and included familial risk of a reading
difficulty and environmental covariates. Findings indicate that preschool/early kindergarten
phonological processing, one of the strongest predictors of subsequent word reading
development, relates to gray matter volume and surface area at birth and development of white
matter volume, surface area, and mean diffusivity across early development. These results offer
further evidence for a neural scaffold for literacy development, which is present at birth and
continues forming across the first several years of life. The present study also provides a roadmap
for future longitudinal studies to examine the relationship between early brain development and
acquisition of other academic skills. Understanding when the foundations for reading emerge can
deliver important insights into the development of instructional approaches and preventative, and
intervention strategies.

857
858
859
860

Acknowledgements
This work was funded by NIH–NICHD R01 HD065762 (2016-2021), the William Hearst Fund
(Harvard University), and the Harvard Catalyst/NIH (5UL1RR025758) to N.G. We would like to
thank all participating families for their long-term dedication to this study. We are grateful for all
additional members of the research team who contributed to data acquisition and quality control,
including Victoria Hue, Ja'Kala Barber, Emily Hu, Olivia Baldi, Brooke Jordan, Jade Dunstan,
Clarisa Carruthers, Carolyn King, Doroteja Rubez, Kathryn Garrisi, Ally Lee, Omeed Moini, Anaïs
Colin, Zumin Chen, Colby Weiss, and Olivia Oh. We also thank Rebecca Petersen for her input
on psychometric assessments, Jennifer Zuk for work on infant scanning procedures, Borjan
Gagoski and Ross Mair for efforts on scanning sequences, Luke Miratrix and Fernando Aguate
for guidance on (non)linear mixed effects models, and Banu Ahtam for feedback.
872

References

1. R. Siegler, Children's learning. *American Psychologist* **6**, 769–778 (2005).
2. M. Bonte, S. Brem, Unraveling individual differences in learning potential: A dynamic framework for the case of reading development. *Dev Cogn Neurosci* [Preprint] (2024).
3. Y.-S. Kim, Toward Integrative Reading Science: The Direct and Indirect Effects Model of Reading. *J Learn Disabil* **53**, 469–491 (2020).
4. R. K. Wagner, J. K. Torgesen, The nature of phonological processing and its causal role in the acquisition of reading skills. *Psychol Bull* **101**, 192–212 (1987).
5. R. Wagner, *et al.*, "The Nature of Prereaders' Phonological Processing Abilities" (1987).
6. U. Goswami, A neural oscillations perspective on phonological development and phonological processing in developmental dyslexia. *Lang Linguist Compass* **13** (2019).
7. U. Goswami, "Phonological development across different languages" in *Routledge International Handbook of English, Language & Literacy Teaching*, (2010), pp. 98–109.
8. H. S. Scarborough, "Connecting early language and literacy to later reading (dis)abilities: Evidence, theory, and practice" in *Handbook of Early Literacy Research*, S. Neuman, D. Dickinson, Eds. (Guilford Press, 2001), pp. 97–110.
9. K. Moll, A. Loff, M. J. Snowling, Cognitive Endophenotypes of Dyslexia. *Scientific Studies of Reading* **17**, 385–397 (2013).
10. M. Melby-Lervåg, S.-A. H. Lyster, C. Hulme, Phonological Skills and Their Role in Learning to Read: A Meta-Analytic Review. *Psychol Bull* **138**, 322–352 (2012).
11. P. Kendeou, P. van den Broek, M. J. White, J. S. Lynch, Predicting Reading Comprehension in Early Elementary School: The Independent Contributions of Oral Language and Decoding Skills. *J Educ Psychol* **101**, 765–778 (2009).
12. Y. S. G. Kim, Why the Simple View of Reading Is Not Simplistic: Unpacking Component Skills of Reading Using a Direct and Indirect Effect Model of Reading (DIER). *Scientific Studies of Reading* **21**, 310–333 (2017).
13. R. A. I. Bethlehem, *et al.*, Brain charts for the human lifespan. *Nature* **604**, 525–533 (2022).
14. J. Gilmore, R. C. Knickmeyer, W. Gao, Imaging structural and functional brain development in early childhood. *Nat Rev Neurosci* **19**, 123–137 (2018).
15. F. Hoeft, *et al.*, Prediction of Children's Reading Skills Using Behavioral, Functional, and Structural Neuroimaging Measures. *Behavioral Neuroscience* **121**, 602–613 (2007).
16. N. M. Raschle, M. Chang, N. Gaab, Structural brain alterations associated with dyslexia predate reading onset. *Neuroimage* **57**, 742–749 (2011).
17. N. Raschle, P. L. Stering, S. N. Meissner, N. Gaab, Altered neuronal response during rapid auditory processing and its relation to phonological processing in prereading children at familial risk for dyslexia. *Cerebral Cortex* **24**, 2489–2501 (2014).
18. N. Raschle, J. Zuk, N. Gaab, Functional characteristics of developmental dyslexia in left-hemispheric posterior brain regions predate reading onset. *Proceedings of the National Academy of Sciences* **109**, 2156–2161 (2012).
19. Z. M. Saygin, *et al.*, Tracking the Roots of Reading Ability: White Matter Volume and Integrity Correlate with Phonological Awareness in Prereading and Early-Reading Kindergarten Children. *Journal of Neuroscience* **33**, 13251–13258 (2013).
20. M. Vandermosten, B. Boets, J. Wouters, P. Ghesquière, A qualitative and quantitative review of diffusion tensor imaging studies in reading and dyslexia. *Neurosci Biobehav Rev* **36**, 1532–1552 (2012).
21. R. Romeo, *et al.*, Language Exposure Relates to Structural Neural Connectivity in Childhood. *J Neurosci* **38**, 7870–7877 (2018).
22. R. Romeo, *et al.*, Beyond the 30-million-word gap: Children's conversational exposure is associated with language-related brain function. *Psychol Sci* **29**, 700–710 (2018).

23. R. A. Marks, *et al.*, Spoken language proficiency predicts print-speech convergence in
beginning readers. *Neuroimage* **201** (2019).
24. K. Davison, *et al.*, Examining the relationship between shared book reading at home,
white matter organization in kindergarten, and subsequent language and reading abilities:
a longitudinal investigation. *J Cogn Neurosci* **35**, 259–275 (2022).
25. H. Lyytinen, *et al.*, The development of children at familial risk for dyslexia: Birth to early
school age. *Ann Dyslexia* **54**, 184–220 (2004).
26. T. K. Guttorm, P. H. T. Leppänen, J. A. Hämäläinen, K. M. Eklund, H. J. Lyytinen,
Newborn event-related potentials predict poorer pre-reading skills in children at risk for
dyslexia. *J Learn Disabil* **43**, 391–401 (2010).
27. T. K. Guttorm, A. Poikkeus, K. M. Eklund, P. Lyytinen, H. Lyytinen, Brain event-related
potentials (ERPs) measured at birth predict later language development in children with
and without familial risk for dyslexia. *Cortex* **41**, 291–303 (2005).
28. T. L. Van Zuijen, A. Plakas, B. A. M. Maassen, N. M. Maurits, A. Van der Leij, Infant
ERPs separate children at risk of dyslexia who become good readers from those who
become poor readers. *Dev Sci* **16**, 554–563 (2013).
29. P. H. T. Leppänen, *et al.*, Infant brain responses associated with reading-related skills
before school and at school age. *Neurophysiologie Clinique* **42**, 35–41 (2012).
30. X. Yu, *et al.*, Functional Connectivity in Infancy and Toddlerhood Predicts Long-Term
Language and Preliteracy Outcomes. *Cerebral Cortex* **32**, 725–736 (2021).
31. J. Zuk, *et al.*, White matter in infancy is prospectively associated with language outcomes
in kindergarten. *Dev Cogn Neurosci* **50** (2021).
32. X. Tang, *et al.*, Longitudinal associations between language network characteristics in the
infant brain and school-age reading abilities are mediated by early-developing
phonological skills. *Dev Cogn Neurosci* **68** (2024).
33. R. Cusack, M. Ranzato, C. J. Charvet, Helpless infants are learning a foundation model.
Trends Cogn Sci (2024). https://doi.org/10.1016/j.tics.2024.05.001.
34. E. Roy, *et al.*, White matter and literacy: A dynamic system in flux. *Dev Cogn Neurosci* **65**
(2024).
35. J. H. Gilmore, *et al.*, Longitudinal development of cortical and subcortical gray matter
from birth to 2 years. *Cerebral Cortex* **22**, 2478–2485 (2012).
36. A. M. Alex, *et al.*, A global multicohort study to map subcortical brain development and
cognition in infancy and early childhood. *Nat Neurosci* **27**, 176–186 (2024).
37. M. Prastawa, J. H. Gilmore, W. Lin, G. Gerig, Automatic segmentation of MR images of
the developing newborn brain. *Med Image Anal* **9**, 457–466 (2005).
38. N. Raschle, *et al.*, Pediatric neuroimaging in early childhood and infancy: Challenges and
practical guidelines. *Ann N Y Acad Sci* **1252**, 43–50 (2012).
39. A. Makropoulos, S. J. Counsell, D. Rueckert, A review on automatic fetal and neonatal
brain MRI segmentation. *Neuroimage* **170**, 231–248 (2018).
40. L. Zöllei, J. E. Iglesias, Y. Ou, P. E. Grant, B. Fischl, Infant FreeSurfer: An automated
segmentation and surface extraction pipeline for T1-weighted neuroimaging data of
infants 0–2 years. *Neuroimage* **218**, 116946 (2020).
41. M. Pietsch, *et al.*, A framework for multi-component analysis of diffusion MRI data over
the neonatal period. *Neuroimage* **186**, 321–337 (2019).
42. M. Grotheer, *et al.*, White matter myelination during early infancy is linked to spatial
gradients and myelin content at birth. *Nat Commun* **13**, 1–12 (2022).
43. T. K. Turesky, J. Vanderauwera, N. Gaab, Imaging the rapidly developing brain: Current
challenges for MRI studies in the first five years of life. *Dev Cogn Neurosci* **47**, 100893
(2021).
44. W. Gao, *et al.*, Functional network development during the first year: Relative sequence
and socioeconomic correlations. *Cerebral Cortex* **25**, 2919–2928 (2015).

45. N. Vijayakumar, K. L. Mills, A. Alexander-Bloch, C. K. Tamnes, S. Whittle, Structural brain
development: A review of methodological approaches and best practices. *Dev Cogn*
Neurosci **33**, 129–148 (2018).
46. R. C. Knickmeyer, *et al.*, A Structural MRI Study of Human Brain Development from Birth
to 2 Years. *Journal of Neuroscience* **28**, 12176–12182 (2008).
47. G. Li, *et al.*, Mapping region-specific longitudinal cortical surface expansion from birth to 2
years of age. *Cerebral Cortex* **23**, 2724–2733 (2013).
48. J. Reynolds, M. N. Grohs, D. Dewey, C. Lebel, Global and regional white matter
development in early childhood. *Neuroimage* **196**, 49–58 (2019).
49. F. Wang, *et al.*, Developmental topography of cortical thickness during infancy. *Proc Natl*
Acad Sci U S A **116**, 15855–15860 (2019).
50. M. Ben-Shachar, R. F. Dougherty, G. K. Deutsch, B. A. Wandell, "The Development of
Cortical Sensitivity to Visual Word Forms" (2011).
51. S. V. Di Pietro, I. I. Karipidis, G. Pleisch, S. Brem, Neurodevelopmental trajectories of
letter and speech sound processing from preschool to the end of elementary school. *Dev*
Cogn Neurosci **61** (2023).
52. S. Brem, *et al.*, Brain sensitivity to print emerges when children learn letter-speech sound
correspondences. *Proc Natl Acad Sci U S A* **107**, 7939–7944 (2010).
53. X. Yu, *et al.*, Emergence of the neural network underlying phonological processing from
the prereading to the emergent reading stage: A longitudinal study. *Hum Brain Mapp* **39**,
2047–2063 (2018).
54. J. Reynolds, X. Long, M. N. Grohs, D. Dewey, C. Lebel, Structural and functional
asymmetry of the language network emerge in early childhood. *Dev Cogn Neurosci* **39**
(2019).
55. G. Dehaene-Lambertz, K. Monzalvo, S. Dehaene, The emergence of the visual word
form: Longitudinal evolution of category-specific ventral visual areas during reading
acquisition. *PLoS Biol* **16** (2018).
56. J. Yeatman, R. F. Dougherty, M. Ben-Shachar, B. A. Wandell, Development of white
matter and reading skills. *Proc Natl Acad Sci U S A* **109**, E3045–E3053 PNAS (2012).
57. Y. Wang, *et al.*, Development of tract-specific white matter pathways during early reading
development in at-risk children and typical controls. *Cerebral Cortex* **27**, 2469–2485
(2017).
58. O. Ozernov-Palchik, D. Sury, T. K. Turesky, X. Yu, N. Gaab, Longitudinal changes in
brain activation underlying reading fluency. *Hum Brain Mapp* **44**, 18–34 (2023).
59. E. Huber, A. Mezer, J. D. Yeatman, Neurobiological underpinnings of rapid white matter
plasticity during intensive reading instruction. *Neuroimage* **243** (2021).
60. E. Eising, *et al.*, Genome-wide analyses of individual differences in quantitatively
assessed reading- and language-related skills in up to 34,000 people. *Proc Natl Acad Sci*
U S A **119** (2022).
61. A. Carrión-Castillo, P. M. Paz-Alonso, M. Carreiras, Brain structure, phenotypic and
genetic correlates of reading performance. *Nat Hum Behav* **7**, 1120–1134 (2023).
62. G. P. Vogler, J. C. DeFries, S. N. Decker, Family history as an indicator of risk for reading
disability. *J Learn Disabil* **18**, 419–421 (1985).
63. M. J. Snowling, M. Melby-Lervåg, Oral language deficits in familial dyslexia: A meta-
analysis and review. *Psychol Bull* **142**, 498–545 (2016).
64. F. Hoeft, *et al.*, Functional and morphometric brain dissociation between dyslexia and
reading ability. *Proceedings of the National Academy of Sciences* **104**, 4234–4239
(2007).
65. F. Richlan, M. Kronbichler, H. Wimmer, Meta-analyzing brain dysfunctions in dyslexic
children and adults. *Neuroimage* **56**, 1735–42 (2011).

66.   F. Richlan, M. Kronbichler, H. Wimmer, Structural abnormalities in the dyslexic brain: A
1025   meta-analysis of voxel-based morphometry studies. *Hum Brain Mapp* **34**, 3055–3065
1026   (2013).

1027   67.   F. Richlan, M. Kronbichler, H. Wimmer, Functional abnormalities in the dyslexic brain: a
1028   quantitative meta-analysis of neuroimaging studies. *Hum Brain Mapp* **30**, 3299–308
1029   (2009).

1030   68.   C. Steinbrink, *et al.*, The contribution of white and gray matter differences to
1031   developmental dyslexia: Insights from DTI and VBM at 3.0 T. *Neuropsychologia* **46**,
1032   3170–3178 (2008).

1033   69.   K. M. Hasan, *et al.*, Diffusion tensor quantification and cognitive correlates of the
1034   macrostructure and microstructure of the corpus callosum in typically developing and
1035   dyslexic children. *NMR Biomed* **25**, 1263–1270 (2012).

1036   70.   E. Temple, *et al.*, Disrupted neural responses to phonological and orthographic
1037   processing in dyslexic children: an fMRI study. *Neuroreport* **12**, 299–307 (2001).

1038   71.   N. Langer, *et al.*, White Matter Alterations in Infants at Risk for Developmental Dyslexia.
1039   *Cereb Cortex* **27**, 1027–1036 (2017).

1040   72.   X. Yu, *et al.*, Patterns of Neural Functional Connectivity in Infants at Familial Risk of
1041   Developmental Dyslexia. *JAMA Netw Open* **5**, E2236102 (2022).

1042   73.   T. K. Guttorm, P. H. T. Leppänen, A. Tolvanen, H. Lyytinen, Event-related potentials in
1043   newborns with and without familial risk for dyslexia: Principal component analysis reveals
1044   differences between the groups. *J Neural Transm* **110**, 1059–1074 (2003).

1045   74.   T. K. Guttorm, P. H. T. Leppänen, U. Richardson, H. Lyytinen, Event-Related Potentials
1046   and Consonant Differentiation in Newborns with Familial Risk for Dyslexia. *J Learn
1047   Disabil* **34**, 534–544 (2001).

1048   75.   M. van Herten, *et al.*, Differences in AERP responses and atypical hemispheric
1049   specialization in 17-month-old children at risk of dyslexia. *Brain Res* **1201**, 100–105
1050   (2008).

1051   76.   C. Doust, *et al.*, Discovery of 42 genome-wide significant loci associated with dyslexia.
1052   *Nat Genet* **54**, 1621–1629 (2022).

1053   77.   K. M. Price, *et al.*, Hypothesis-driven genome-wide association studies provide novel
1054   insights into genetics of reading disabilities. *Transl Psychiatry* **12** (2022).

1055   78.   M. Watabe-Uchida, K. A. John, J. A. Janas, S. E. Newey, L. Van Aelst, The Rac Activator
1056   DOCK7 Regulates Neuronal Polarity through Local Phosphorylation of Stathmin/Op18.
1057   *Neuron* **51**, 727–739 (2006).

1058   79.   A. C. Payne Grover, W. L. Andrea Angel, "The Role of Home Literacy Environment in the
1059   Development of Language Ability in Preschool Children from Low-income Families"
1060   (1994).

1061   80.   A. G. Bus, M. H. van IJzendoorn, A. D. Pellegrini, Joint book reading makes for success
1062   in learning to read: A meta-analysis on intergenerational transmission of literacy. *Rev
1063   Educ Res* **65**, 1–21 (1995).

1064   81.   S. R. Burgess, S. A. Hecht, C. J. Lonigan, Relations of the home literacy environment
1065   (HLE) to the development of reading-related abilities: A one-year longitudinal study. *Read
1066   Res Q* **37**, 408–426 (2002).

1067   82.   M. A. Foster, R. Lambert, M. Abbott-Shim, F. McCarty, S. Franze, A model of home
1068   learning environment and social risk factors in relation to children's emergent literacy and
1069   social outcomes. *Early Child Res Q* **20**, 13–36 (2005).

1070   83.   S. A. Schmitt, A. M. Simpson, M. Friend, A longitudinal assessment of the home literacy
1071   environment and early language. *Infant Child Dev* **20**, 409–431 (2011).

1072   84.   L. G. Hamilton, M. E. Hayiou-Thomas, C. Hulme, M. J. Snowling, The Home Literacy
1073   Environment as a Predictor of the Early Literacy Development of Children at Family-Risk
1074   of Dyslexia. *Scientific Studies of Reading* **20**, 401–419 (2016).

85. J. Karrass, J. M. Braungart-Rieker, Effects of shared parent-infant book reading on early language acquisition. *J Appl Dev Psychol* **26**, 133–148 (2005).

86. E. Hoff, The Specificity of Environmental Influence: Socioeconomic Status Affects Early Vocabulary Development Via Maternal Speech. *Child Dev* **74**, 1368–1378 (2003).

87. A. Muhinyi, M. L. Rowe, Shared reading with preverbal infants and later language development. *J Appl Dev Psychol* **64** (2019).

88. M.-L. Laakso, A.-M. Poikkeus, P. Lyytinen, Shared reading interaction in families with and without genetic risk for dyslexia: implications for toddlers' language development. *Infant Child Dev* **8**, 179–195 (1999).

89. J. L. Malin, N. J. Cabrera, M. L. Rowe, Low-income minority mothers' and fathers' reading and children's interest: Longitudinal contributions to children's receptive vocabulary skills. *Early Child Res Q* **29**, 425–432 (2014).

90. Ö. E. Demir-Lira, L. R. Applebaum, S. Goldin-Meadow, S. C. Levine, Parents' early book reading to children: Relation to children's later language and literacy outcomes controlling for other parent language input. *Dev Sci* **22** (2019).

91. S. J. Powers, Y. Wang, S. D. Beach, G. D. Sideridis, N. Gaab, Examining the relationship between home literacy environment and neural correlates of phonological processing in beginning readers with and without a familial risk for dyslexia: an fMRI study. *Ann Dyslexia* **66**, 337–360 (2016).

92. J. S. Hutton, J. Dudley, T. Horowitz-Kraus, T. DeWitt, S. K. Holland, Associations between home literacy environment, brain white matter integrity and cognitive abilities in preschool-age children. *Acta Paediatr* **109**, 1376–1386 (2020).

93. J. S. Hutton, T. Horowitz-Kraus, A. L. Mendelsohn, T. DeWitt, S. K. Holland, Home reading environment and brain activation in preschool children listening to stories. *Pediatrics* **136**, 466–478 (2015).

94. J. S. Hutton, *et al.*, Shared Reading Quality and Brain Activation during Story Listening in Preschool-Age Children. *Journal of Pediatrics* **191**, 204-211.e1 (2017).

95. L. S. King, M. C. Camacho, D. F. Montez, K. L. Humphreys, I. H. Gotlib, Naturalistic language input is associated with resting-state functional connectivity in infancy. *Journal of Neuroscience* **41**, 424–434 (2021).

96. T. K. Turesky, *et al.*, Home language and literacy environment and its relationship to socioeconomic status and white matter structure in infancy. *Brain Struct Funct* **227**, 2633–2645 (2022).

97. J. Zuk, J. Vanderauwera, T. Turesky, X. Yu, N. Gaab, Neurobiological predispositions for musicality: White matter in infancy predicts school-age music aptitude. *Dev Sci* **26** (2023).

98. C. Lebel, *et al.*, Prepartum and Postpartum Maternal Depressive Symptoms Are Related to Children's Brain Structure in Preschool. *Biol Psychiatry* **80**, 859–868 (2016).

99. C. Thieba, *et al.*, Factors Associated With Successful MRI Scanning in Unsedated Young Children. *Front Pediatr* **6**, 1–5 (2018).

100. D. Paniukov, R. M. Lebel, G. Giesbrecht, C. Lebel, Cerebral blood flow increases across early childhood. *Neuroimage* **204** (2020).

101. M. Walton, D. Dewey, C. Lebel, Brain white matter structure and language ability in preschool-aged children. *Brain Lang* **176**, 19–25 (2018).

102. J. Reynolds, X. Long, D. Paniukov, M. Bagshawe, C. Lebel, Calgary Preschool magnetic resonance imaging (MRI) dataset. *Data Brief* **29** (2020).

103. L. M. Betancourt, *et al.*, Effect of socioeconomic status (SES) disparity on neural development in female African-American infants at age 1 month. *Dev Sci* **19**, 947–956 (2016).

104. N. H. Brito, W. P. Fifer, M. M. Myers, A. J. Elliott, K. G. Noble, Associations among family socioeconomic status, EEG power at birth, and cognitive skills during infancy. *Dev Cogn Neurosci* **19**, 144–151 (2016).

105. G. M. Lawson, J. T. Duda, B. B. Avants, J. Wu, M. J. Farah, Associations between
children's socioeconomic status and prefrontal cortical thickness. *Dev Sci* **16**, 641–652
(2013).
106. E. C. Merz, N. Tottenham, K. G. Noble, Socioeconomic Status, Amygdala Volume, and
Internalizing Symptoms in Children and Adolescents. *Journal of Clinical Child and*
Adolescent Psychology **47**, 312–323 (2018).
107. K. G. Noble, *et al.*, Family income, parental education and brain structure in children and
adolescents. *Nat Neurosci* **18**, 773–778 (2015).
108. O. Ozernov-Palchik, *et al.*, The relationship between socioeconomic status and white
matter microstructure in pre-reading children: A longitudinal investigation. *Hum Brain*
Mapp **40**, 741–754 (2018).
109. M. K. Denney, J. P. English, M. M. Gerber, J. Leafstedt, M. L. Ruz, Family and home
literacy practices: mediating factors for preliterate English learners at risk in *Annual*
Meeting of the American Educational Research Associations, (2001), pp. 1–17.
110. Y. S. G. Kim, Executive Functions and Morphological Awareness Explain the Shared
Variance between Word Reading and Listening Comprehension. *Scientific Studies of*
Reading **27**, 451–474 (2023).
111. F. A. Schrank, K. S. McGrew, N. Mather, *Woodcock-Johnson IV Tests of Cognitive*
Abilities (Riverside, 2014).
112. M. Korkman, U. Kirk, S. Kemp, *NEPSY–Second Edition (NEPSY-II)*, 2nd Ed. (The
Psychological Corporation, 2007).
113. F. Schrank, N. Mather, K. McGrew, *Woodcock-Johnson IV Tests of Oral Language*
(Riverside Publishing Co, 2014).
114. R. W. Woodcock, *Woodcock Reading Mastery Tests (WRMT-III)* (NCS Pearson, 2011).
115. A. S. Kaufman, N. L. Kaufman, Kaufman Brief Intelligence Test, Second Edition.
PsycTESTS Dataset [Preprint] (2004).
116. L. L. Backhausen, *et al.*, Quality control of structural MRI images applied using
FreeSurfer-a hands-on workflow to rate motion artifacts. *Front Neurosci* **10** (2016).
117. V. S. Natu, *et al.*, Infants' cortex undergoes microstructural growth coupled with
myelination during development. *Commun Biol* **4** (2021).
118. L. Wang, *et al.*, iBEAT V2.0: a multisite-applicable, deep learning-based pipeline for infant
cerebral cortical surface reconstruction. *Nat Protoc* **18**, 1488–1509 (2023).
119. G. Li, *et al.*, Measuring the dynamic longitudinal cortex development in infants by
reconstruction of temporally consistent cortical surfaces. *Neuroimage* **90**, 266–279
(2014).
120. G. Li, *et al.*, Computational neuroanatomy of baby brains: A review. *Neuroimage*
[Preprint] (2019).
121. C. S. McCarthy, *et al.*, A comparison of FreeSurfer-generated data with and without
manual intervention. *Front Neurosci* **9** (2015).
122. E. P. Pulli, *et al.*, Feasibility of FreeSurfer Processing for T1-Weighted Brain Images of 5-
Year-Olds: Semiautomated Protocol of FinnBrain Neuroimaging Lab. *Front Neurosci* **16**
(2022).
123. B. Ahtam, *et al.*, Morphological Features of Language Regions in Individuals with
Tuberous Sclerosis Complex. *J Autism Dev Disord* **54**, 3155–3175 (2024).
124. K. Chyl, G. Fraga-González, S. Brem, K. Jednoróg, Brain dynamics of (a)typical reading
development—a review of longitudinal studies. *NPJ Sci Learn* [Preprint] (2021).
125. M. Bastiani, *et al.*, Automated processing pipeline for neonatal diffusion MRI in the
developing Human Connectome Project. *Neuroimage* **185**, 750–763 (2019).
126. L. Cordero-Grande, D. Christiaens, J. Hutter, A. N. Price, J. V. Hajnal, Complex diffusion-
weighted image estimation via matrix recovery under general noise models. *Neuroimage*
200, 391–404 (2019).

127. J. Veraart, *et al.*, Denoising of diffusion MRI using random matrix theory. *Neuroimage*
142, 394–406 (2016).
128. J. Veraart, E. Fieremans, D. S. Novikov, Diffusion MRI noise mapping using random
matrix theory. *Magn Reson Med* **76**, 1582–1593 (2016).
129. J. L. R. Andersson, S. Skare, J. Ashburner, How to correct susceptibility distortions in
spin-echo echo-planar images: Application to diffusion tensor imaging. *Neuroimage* **20**,
870–888 (2003).
130. J. L. R. Andersson, *et al.*, Towards a comprehensive framework for movement and
distortion correction of diffusion MR images: Within volume movement. *Neuroimage* **152**,
450–466 (2017).
131. J. L. R. Andersson, S. N. Sotiropoulos, An integrated approach to correction for off-
resonance effects and subject movement in diffusion MR imaging. *Neuroimage* **125**,
1063–1078 (2016).
132. S. Skare, R. Bammer, Jacobian weighting of distortion corrected EPI data in *Proceedings*
of the International Society for Magnetic Resonance in Medicine, (2010).
133. S. M. Smith, *et al.*, Advances in functional and structural MR image analysis and
implementation as FSL. *Neuroimage* **23**, 208–219 (2004).
134. N. J. Tustison, *et al.*, N4ITK: Improved N3 bias correction. *IEEE Trans Med Imaging* **29**,
1310–1320 (2010).
135. T. Dhollander, R. Mito, D. Raffelt, A. Connelly, Improved white matter response function
estimation for 3-tissue constrained spherical deconvolution in *Proc Intl Soc Mag Reson*
Med, (2019).
136. B. Jeurissen, J. D. Tournier, T. Dhollander, A. Connelly, J. Sijbers, Multi-tissue
constrained spherical deconvolution for improved analysis of multi-shell diffusion MRI
data. *Neuroimage* **103**, 411–426 (2014).
137. J. D. Tournier, F. Calamante, D. G. Gadian, A. Connelly, Direct estimation of the fiber
orientation density function from diffusion-weighted MRI data using spherical
deconvolution. *Neuroimage* **23**, 1176–1185 (2004).
138. J. D. Tournier, F. Calamante, A. Connelly, MRtrix: Diffusion tractography in crossing fiber
regions. *Int J Imaging Syst Technol* **22**, 53–66 (2012).
139. R. E. Smith, J. D. Tournier, F. Calamante, A. Connelly, Anatomically-constrained
tractography: Improved diffusion MRI streamlines tractography through effective use of
anatomical information. *Neuroimage* **62**, 1924–1938 (2012).
140. J. Yeatman, R. Dougherty, N. Myall, B. Wandell, H. Feldman, Tract profiles of white
matter properties: automating fiber-tract quantification. *PLoS One* **7**, e49790 (2012).
141. J. Kruper, *et al.*, Evaluating the Reliability of Human Brain White Matter Tractometry.
Aperture Neuro **1**, 1–25 (2021).
142. M. Grotheer, *et al.*, Human white matter myelinates faster in utero than ex utero. (2023).
https://doi.org/10.1073/pnas.
143. S. M. Smith, T. E. Nichols, Threshold-free cluster enhancement: Addressing problems of
smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* **44**,
83–98 (2009).
144. J. Frossard, O. Renaud, Permutation tests for regression, anova, and comparison of
signals: The permuco package. *J Stat Softw* **99**, 1–32 (2021).
145. J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar, R Core Team, nlme: Linear and Nonlinear
Mixed Effects Models. *R Package Version 3* [Preprint] (2017). Available at: https://bugs.r-
project.org.
146. A. F. G. Rosen, *et al.*, Quantitative assessment of structural image quality. *Neuroimage*
169, 407–418 (2018).

147. T. Turesky, *et al.*, Brain Morphometry and Diminished Physical Growth in Bangladeshi
Children Growing up in Extreme Poverty: a Longitudinal Study. *Dev Cogn Neurosci* **52**,
101029 (2021).
148. T. K. Turesky, *et al.*, Brain morphometry and chronic inflammation in Bangladeshi
children growing up in extreme poverty. *Imaging Neuroscience* **2**, 1–16 (2024).
149. J. Monereo-Sánchez, *et al.*, Quality control strategies for brain MRI segmentation and
parcellation: Practical approaches and recommendations - insights from the Maastricht
study. *Neuroimage* **237** (2021).
150. J. Matsuzawa, M. Matsui, T. Konishi, Age-related Volumetric Changes of Brain Gray and
White Matter in Healthy Infants and Children. *Cerebral Cortex* **11**, 335–342 (2001).
151. S. Groeschel, B. Vollmer, M. D. King, A. Connelly, Developmental changes in cerebral
grey and white matter volume from infancy to adulthood. *International Journal of
Developmental Neuroscience* **28**, 481–489 (2010).
152. S. C. L. Deoni, D. C. Dean, J. O. Muircheartaigh, H. Dirks, B. A. Jerskey, Investigating
white matter development in infancy and early childhood using myelin water faction and
relaxation time mapping. **63**, 1038–1053 (2012).
153. S. C. L. Deoni, *et al.*, Breastfeeding and early white matter development: A cross-
sectional study. *Neuroimage* **82**, 77–86 (2013).
154. J. Remer, *et al.*, Quantifying cortical development in typically developing toddlers and
young children, 1–6 years of age. *Neuroimage* **153**, 246–261 (2017).
155. C. E. Sanchez, J. E. Richards, C. R. Almli, Neurodevelopmental MRI brain templates for
children from 2 weeks to 4 years of age. *Dev Psychobiol* **54**, 77–91 (2011).
156. X. Geng, *et al.*, NeuroImage Quantitative tract-based white matter development from birth
to age 2 years. *Neuroimage* **61**, 542–557 (2012).
157. K. Im, N. Raschle, S. A. Smith, P. Ellen Grant, N. Gaab, Atypical Sulcal Pattern in
Children with Developmental Dyslexia and At-Risk Kindergarteners. *Cerebral Cortex* **26**,
1138–1148 (2016).
158. C. A. Myers, *et al.*, White Matter Morphometric Changes Uniquely Predict Children's
Reading Acquisition. *Psychol Sci* **25**, 1870–1883 (2014).
159. K. Hannula-Jouppi, *et al.*, The axon guidance receptor gene ROBO1 is a candidate gene
for developmental dyslexia. *PLoS Genet* **1**, 0467–0474 (2005).
160. L. Reiche, *et al.*, C21orf91 Regulates Oligodendroglial Precursor Cell Fate—A Switch in
the Glial Lineage? *Front Cell Neurosci* **15** (2021).
161. A. M. Fjell, *et al.*, The relationship between diffusion tensor imaging and volumetry as
measures of white matter properties. *Neuroimage* **42**, 1654–1668 (2008).
162. A. D. Friederici, The cortical language circuit: From auditory perception to sentence
comprehension. *Trends Cogn Sci* [Preprint] (2012).
163. J. Wang, M. L. Rice, J. R. Booth, Syntactic and semantic specialization and integration in
5-to 6-year-old children during auditory sentence processing. *J Cogn Neurosci* **32**, 36–49
(2019).
164. P. K. Kuhl, Early language acquisition: Cracking the speech code. *Nat Rev Neurosci*
[Preprint] (2004).
165. X. Yu, J. Zuk, N. Gaab, What Factors Facilitate Resilience in Developmental Dyslexia?
Examining Protective and Compensatory Mechanisms Across the Neurodevelopmental
Trajectory. *Child Dev Perspect* **12**, 240–246 (2018).
166. O. Ozernov-Palchik, X. Yu, Y. Wang, N. Gaab, Lessons to be learned: How a
comprehensive neurobiological framework of atypical reading development can inform
educational practice. *Curr Opin Behav Sci* **10**, 45–58 (2016).
167. H. Catts, Y. Petscher, Early identification of dyslexia: Current advancements and future
directions. *Perspect Lang Lit* **44**, 33–36 (2018).

168.   S. A. Hart, C. Little, E. van Bergen, Nurture might be nature: cautionary tales and
1277   proposed solutions. *NPJ Sci Learn* **6**, 1–12 (2021).
1278   169.   J. Zuk, *et al.*, Multifactorial pathways facilitate resilience among kindergarteners at risk for
1279   dyslexia: A longitudinal behavioral and neuroimaging study. *Dev Sci* **24**, 1–18 (2021).
1280   170.   X. Yu, *et al.*, Putative protective neural mechanisms in prereaders with a family history of
1281   dyslexia who subsequently develop typical reading skills. *Hum Brain Mapp* **41**, 2827–
1282   2845 (2020).
1283   171.   F. Hoeft, *et al.*, Neural systems predicting long-term outcome in dyslexia. *Proceedings of
1284   the National Academy of Sciences* **108**, 361–366 (2011).
1285   172.   C. A. Nelson, L. J. Gabard-Durnam, Early Adversity and Critical Periods :
1286   Neurodevelopmental Consequences of Violating the Expectable Environment. *Trends
1287   Neurosci* **43**, 133–143 (2020).
1288   173.   P. J. Curran, K. Obeidat, D. Losardo, Twelve frequently asked questions about growth
1289   curve modeling. *Journal of Cognition and Development* **11**, 121–136 (2010).
1290   174.   E. H. Telzer, *et al.*, Methodological considerations for developmental longitudinal fMRI
1291   research. *Dev Cogn Neurosci* [Preprint] (2018).
1292   175.   E. Roy, *et al.*, Differences in educational opportunity predict white matter development.
1293   *Dev Cogn Neurosci* **67** (2024).
1294

Supplementary Methods

1. Participants

New England cohort

Children participated in a longitudinal investigation of brain and literacy development from infancy to school age (NIH–NICHD R01 HD065762). Families of these children were recruited from the New England Area using the Research Participant Registry provided by the Division of Developmental Medicine at Boston Children's Hospital, and with flyers and ads disseminated in local schools and newspapers, at community events, and on social media. Children were enrolled as infants with the expectation that participation would continue at subsequent developmental stages across the first decade of life. Children were excluded from the study if at any timepoint they were diagnosed with neurological, sensory, or motor disorders or had contraindications for MRI evaluation (e.g., metal implants). All children included were from English-speaking families and were born at gestational week 37 or later. Race demographics, whose reporting is encouraged to improve equitability in neuroscience (1), are as follows: 70% White/Caucasian, 6% Black/African American, 5% Asian, 16% Multiracial, and 5% Hispanic. Neuroimaging and behavioral testing were conducted at Boston Children's Hospital prior to 2021 and at the Center for Brain Science at Harvard University from 2021 onward. At Boston Children's Hospital, children's anatomical MRI scans (please see parameters below) did not show any potentially malignant brain features, as reviewed by a pediatric neuroradiologist. This study was approved by the Institutional Review Boards of Boston Children's Hospital (IRB-P00023182) and Harvard University (IRB21-0916). Informed written consent was provided by each participating infant's parent(s) and children gave verbal assent for participation after 50 months of age.

Calgary cohort

Children in the Calgary, Alberta area were recruited from the ongoing study on pregnancy outcomes and nutrition (2). Children in this study were predominantly (roughly 90%) from Caucasian families, but also included Asian/Pacific Islander, African, Filipino, Latino/Hispanic, and Multiracial children. Children were born at gestational week 36 or later and did not have diagnosed genetic, neurological, or neurodevelopmental disorders. The study was approved by the University of Calgary Conjoint Health Research Ethics Board (REB13-0020). Parent and/or guardian consent and child assent were acquired for all participants. For additional information, please see the open-source repository (https://osf.io/axz5r/) and previous publications (3–9).

2. MRI data acquisition

New England cohort

All data were acquired on a 3.0 T Siemens scanner with a 32-channel head coil. Please note, sequence parameters varied to optimize acquisition for the increasing head size and neuroanatomy of the participants across the developmental window. Structural T1-weighted magnetization-prepared rapid gradient-echo (MPRAGE) scans were acquired with the following parameters: TR = 2270-2520 ms, TE = 1.66-1.73 ms, field of view = 192-224 mm, 1 mm^3 voxels, 144-176 sagittal slices. Diffusion echo planar images were acquired using the following parameters: TR = 3800-8320 ms, TE = 88-89 ms, flip angle = 90°, field of view = 180-256 mm, voxel size = 2 x 2 x 2 mm^3, 62-78 slices, 30 b = 1000 s/mm^2 gradient directions, 10-11 b = 0 s/mm^2 non-diffusion-weighted volumes. Diffusion data were acquired with slice-acceleration (SMS/MB) factor = 2 and one reverse phase encoding (i.e., posterior-to-anterior) volume. Please note, sequence parameters varied to optimize acquisition for the increasing head size and neuroanatomy of the participants across the developmental window.

Calgary cohort

All data were acquired on a 3.0 T General Electric scanner with a 32-channel head coil. Structural T1-weighted scans were acquired with the following parameters: TR = 8.23 ms, TE = 3.76 ms, field of view = 230 mm, 0.45 x 0.45 x 0.9 mm^3 voxels, 210 slices. Diffusion echo planar images were acquired using the following parameters: TR = 6750 ms, TE = 79 ms, flip angle = 90°, field of view = 200 mm, voxel size = 0.78 x 0.78 x 2.2 mm^3, 50-55 slices, 30 b = 750 s/mm^2 gradient directions, 5 b = 0 s/mm^2 non-diffusion-weighted volumes. Diffusion acquisition did not use a slice-acceleration factor and did not include a reverse phase encoding (i.e., posterior-to-anterior) volume.

3. Algorithm for combining FreeSurfer and iBEATv2.0 segmentations

Segmentations from each software package were then hybridized using in-house MATLAB code that combined the cortical pial and white matter boundaries labeled by iBEATv2.0 with the FreeSurfer subcortical parcellations (i.e., effectively relabeling cortical gray and white matter in the FreeSurfer segmentation using iBEATv2.0 tissue-class demarcations). Herein, iBEATv2.0 3-class tissue segmentations were first resampled to the space of the aseg file. iBEATv2.0 gray matter and cerebrospinal fluid voxels that overlapped with the aseg gray matter and cerebrospinal fluid labels, respectively, inherited the latter's labels. iBEATv2.0 white matter voxels overlapping with aseg white *or gray* matter received the aseg white matter label; as iBEATv2.0 does not

distinguish hemispheres in its labels and FreeSurfer generally overestimates gray matter in this age group, gray matter information was also used here for relabeling. iBEATv2.0 voxels unlabeled by these procedures were then submitted to a nearest neighbor search to identify aseg labels at minimum Euclidean distance. As FreeSurfer generally performed better compared with iBEATv2.0 on subcortical areas, all subcortical (non-cerebellum areas) were relabeled with subcortical FreeSurfer labels. Finally, hybridized segmentation files were converted to FreeSurfer-style white matter files and submitted to a modified version of the standard FreeSurfer v7.3 "recon" pipeline.

4. Preparation of Euler numbers

Similar to Bethlehem and colleagues 2022, for further quality control, we also extracted Euler numbers (10), which quantify topological defects in FreeSurfer's cortical reconstruction and have been shown to correlate with visual ratings and mark artifactual images (11). Euler numbers from each hemisphere were averaged across observations for each participant (average: (11); sum: (12)) and used in sensitivity analyses.

5. Implementations of pyAFQ and pyBabyAFQ

Whole-brain tractography was then submitted for fiber tract segmentation to the open-source instantiation of Automated Fiber Quantification (AFQ; (13, 14). Herein, waypoint regions-of-interest (ROIs) and a probabilistic fiber atlas were mapped from a standard template to the individual brain. Fibers were delineated into separate tracts according to the waypoint ROIs through which they passed and the tract they were most likely to belong based on the probability atlas. Additional fibers were removed (i.e., outlier detection) if they deviated sufficiently from the core of the tract. While all diffusion data underwent the above processes, parameters differed by age group. For brains > 25 months, the standard pyAFQ pipeline was used, in which ROIs and the probability fiber atlas were mapped from an adult MNI template and fibers were removed if they were more than five standard deviations from the core of the tract (14). In contrast, for brains ≤ 24 months, we used pyBabyAFQ (15), an implementation in the pyAFQ suite designed to accommodate the smaller neuroanatomy in infants. Accordingly, ROIs and the probabilistic fiber atlas are mapped from an infant template (16), the ROIs are smaller compared to those used by standard pyAFQ, an additional ROI was used for tracts with acute curves, and the fiber outlier detection threshold was reduced to four standard deviations from the tract core. Importantly, varying the parameters used to segment the tracts by age group preserves the accuracy with

which tracts are segmented and reduces age-related bias that would emerge if using standard (i.e., suboptimal) parameters for younger children.

6. Quality control procedures for longitudinal trajectory estimation

Prior to modeling, estimates of brain structure and white matter organization, excepting cortical thickness and mean curvature, underwent a final quality control procedure to remove neuroanatomically implausible observations by setting annual change thresholds. For instance, it was unlikely that gray matter volume in any particular brain area changed by more than 10% per year in children over 25 months. Accordingly, for brains > 25 months, observations for brain regions and tracts of interest (please see above) were flagged if they were preceded or followed by ≥ 10% annual change (positive or negative). To improve model convergence in the first sensitivity analysis (with nonlinear mixed effects models), the threshold for mean diffusivity was lowered from 10% to 5%. To mitigate data loss, we next identified which timepoint—the earlier timepoint or later timepoint—was more likely to be inaccurate, using an outlier detection procedure. Herein, we generated average (across participants) estimates for each brain area/tract and each brain measure by timepoint (roughly, grade level) to which to compare the observations flagged in the previous step. Out of the pair of flagged observations, the one farther from the average estimate corresponding to its timepoint was discarded and the other flagged observation was preserved. This process was done iteratively to handle cases in which children over 50 months had MRI observations from more than two timepoints. A similar quality control procedure was used for brains ≤ 24 months, but it needed to account for the rapid brain growth already thoroughly reported (17, 18). Consequently, observations were only flagged when inter-timepoint changes were negative for gray/white matter volume, surface area, or fractional anisotropy, or positive for mean diffusivity. No inter-timepoint threshold was used for cortical thickness or mean curvature for brains ≤ 24 months. All remaining participants after these additional quality control procedures had multiple observations (i.e., longitudinal datasets), including one from ≤ 24 months.

7. Sensitivity analyses

To test the reliability of our results, we performed a replication analysis on gray/white matter volume, and surface area with nonlinear mixed effects models using asymptotic functions from the R 'nlme' package (19), similar to that described in Alex and colleagues (18); https://github.com/knickmeyer-lab/ORIGINs_ICV-and-Subcortical-volume-development-in-early-childhood). Intercepts and asymptotes were modeled as fixed and random effects; rate constants were modeled as fixed effects. The reason that this model was not used in the main analysis is

that it does not use a random slopes term, and a key aspect of the current study is to examine the relation between brain growth and literacy subskills. However, it should be noted that the nonlinear model does provide an indirect examination of the relation between growth and literacy subskills; e.g., if there is no association between the intercept and the literacy subskill, but there is an association between the asymptote and the literacy subskill, then it could be inferred that there is an association between brain growth and the literacy subskill. Although both linear and nonlinear models have different requirements (e.g., linear models require relationships between predictors and outcomes to be linear), the functional forms used on the current dataset in both cases fit the data closely (Supplementary Figure 8), suggesting random parameters were similar or proportional across models.

For the main analysis, we opted to model brain development prior to testing brain-behavior associations because this comports with the temporal order theorized, that brain development effects subsequent behavioral skills. Also, our sample size is larger for longitudinal brain data alone compared with longitudinal brain data plus reading-related outcomes; therefore, longitudinal models of brain development would be improved if not including outcomes in the model. However, in practice, contributions of phonological processing main and age x phonological interaction terms should be analogous to associations between phonological processing and curve intercepts and slopes, respectively. Consequently, we also examined phonological processing main and age x phonological processing interaction terms included in the linear mixed effects models.

In addition, we did not initially control for total intracranial volume (TIV) when modeling longitudinal trajectories of brain structure, consistent with other work examining developmental trajectories of brain structure beginning in infancy (12, 18, 20–23). Further, recent work has shown that TIV correction may be problematic, reducing brain-behavior predictive accuracies for gray matter volume and surface area or potentially generating spurious predictions for cortical thickness (24). Given these concerns and recommendations to report both raw and TIV-corrected results (25), we thought that an appropriate use of TIV would be to recompute brain-behavior associations for volumetric and surface-based measures using semipartial correlations (Pearson) with the random terms from longitudinal modeling with TIV as covariates of no interest. We report results of the linear mixed model run on brain-behavior relations with TIV.

Lastly, for volumetric and surface-based measures, visual ratings of cortical surfaces were used to identify sub-optimal datasets (visual ratings of tract reconstructions were for measures of white matter organization). However, Euler numbers, which quantify the number the topological defects in FreeSurfer's cortical surface reconstruction (10), have been shown to consistently correlate with quality ratings (11) and to serve as reproducible alternatives to manual quality

control, including manual editing (26). Comparable to Rosen and colleagues, visual ratings and Euler numbers were highly correlated after controlling for age and biological sex (r = 0.44, p = 2.3 x 10^{-28}). Therefore, to account for residual variance due to segmentation quality in a data-driven, reproducible manner, we submitted volumetric and surface-based brain-behavior associations to semipartial correlations (Pearson) with average (across timepoint) Euler numbers.

8. References for the Supplement

1.   E. K. Webb, J. A. Etter, J. A. Kwasa, Addressing racial and phenotypic bias in human neuroscience methods. *Nat Neurosci* **25**, 410–414 (2022).
2.   B. J. Kaplan, *et al.*, The Alberta Pregnancy Outcomes and Nutrition (APrON) cohort study: Rationale and methods. *Matern Child Nutr* **10**, 44–60 (2014).
3.   D. Paniukov, R. M. Lebel, G. Giesbrecht, C. Lebel, Cerebral blood flow increases across early childhood. *Neuroimage* **204** (2020).
4.   M. Walton, D. Dewey, C. Lebel, Brain white matter structure and language ability in preschool-aged children. *Brain Lang* **176**, 19–25 (2018).
5.   J. Reynolds, M. N. Grohs, D. Dewey, C. Lebel, Global and regional white matter development in early childhood. *Neuroimage* **196**, 49–58 (2019).
6.   J. Reynolds, X. Long, D. Paniukov, M. Bagshawe, C. Lebel, Calgary Preschool magnetic resonance imaging (MRI) dataset. *Data Brief* **29** (2020).
7.   J. Reynolds, X. Long, M. N. Grohs, D. Dewey, C. Lebel, Structural and functional asymmetry of the language network emerge in early childhood. *Dev Cogn Neurosci* **39** (2019).
8.   C. Lebel, *et al.*, Prepartum and Postpartum Maternal Depressive Symptoms Are Related to Children's Brain Structure in Preschool. *Biol Psychiatry* **80**, 859–868 (2016).
9.   C. Thieba, *et al.*, Factors Associated With Successful MRI Scanning in Unsedated Young Children. *Front Pediatr* **6**, 1–5 (2018).
10.   A. M. Dale, B. Fischl, M. I. Sereno, "Cortical Surface-Based Analysis I. Segmentation and Surface Reconstruction" (1999).
11.   A. F. G. Rosen, *et al.*, Quantitative assessment of structural image quality. *Neuroimage* **169**, 407–418 (2018).
12.   R. A. I. Bethlehem, *et al.*, Brain charts for the human lifespan. *Nature* **604**, 525–533 (2022).
13.   J. Yeatman, R. Dougherty, N. Myall, B. Wandell, H. Feldman, Tract profiles of white matter properties: automating fiber-tract quantification. *PLoS One* **7**, e49790 (2012).
14.   J. Kruper, *et al.*, Evaluating the Reliability of Human Brain White Matter Tractometry. *Aperture Neuro* **1**, 1–25 (2021).
15.   M. Grotheer, *et al.*, Human white matter myelinates faster in utero than ex utero. (2023). https://doi.org/10.1073/pnas.
16.   F. Shi, *et al.*, Infant brain atlases from neonates to 1- and 2-year-olds. *PLoS One* **6** (2011).
17.   J. Gilmore, R. C. Knickmeyer, W. Gao, Imaging structural and functional brain development in early childhood. *Nat Rev Neurosci* **19**, 123–137 (2018).
18.   A. M. Alex, *et al.*, A global multicohort study to map subcortical brain development and cognition in infancy and early childhood. *Nat Neurosci* **27**, 176–186 (2024).
19.   J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar, R Core Team, nlme: Linear and Nonlinear Mixed Effects Models. *R Package Version 3* [Preprint] (2017). Available at: https://bugs.r-project.org.

20. S. Groeschel, B. Vollmer, M. D. King, A. Connelly, Developmental changes in cerebral grey and white matter volume from infancy to adulthood. *International Journal of Developmental Neuroscience* **28**, 481–489 (2010).

21. J. Matsuzawa, M. Matsui, T. Konishi, Age-related Volumetric Changes of Brain Gray and White Matter in Healthy Infants and Children. *Cerebral Cortex* **11**, 335–342 (2001).

22. S. C. L. Deoni, D. C. Dean, J. O. Muircheartaigh, H. Dirks, B. A. Jerskey, Investigating white matter development in infancy and early childhood using myelin water faction and relaxation time mapping. **63**, 1038–1053 (2012).

23. J. Remer, *et al.*, Quantifying cortical development in typically developing toddlers and young children, 1–6 years of age. *Neuroimage* **153**, 246–261 (2017).

24. E. Dhamala, *et al.*, Proportional intracranial volume correction differentially biases behavioral predictions across neuroanatomical features, sexes, and development. *Neuroimage* **260** (2022).

25. N. Vijayakumar, K. L. Mills, A. Alexander-Bloch, C. K. Tamnes, S. Whittle, Structural brain development: A review of methodological approaches and best practices. *Dev Cogn Neurosci* **33**, 129–148 (2018).

26. J. Monereo-Sánchez, *et al.*, Quality control strategies for brain MRI segmentation and parcellation: Practical approaches and recommendations - insights from the Maastricht study. *Neuroimage* **237** (2021).

Supplemental Tables

Supplementary Table 1. Participant demographics		Structure	Diffusion
General information	N. participants	98	128
	N. observations	276	396
	N. observations per participant	3 ± 1	3 ± 1
	Age at literacy-related subskill testing (months)	63 ± 5.3	63 ± 5.1
	Age at decoding/word reading testing (months)	82 ± 6.4	82 ± 6.4
Covariates	Biological sex (F/M)	46/52	67/61
	Maternal education (years)	17 ± 2.1	17 ± 2.1
	Cohort ([New England]/Calgary)	59/39	77/51
	Family history of reading difficulty (+/-)	30/29	39/38
	Home literacy environment (a.u.)	0.073 ± 0.40	0.041 ± 0.42
Literacy-related subskills	Phonological processing standard score	108 ± 13	106 ± 14
	Oral language standard score	115 ± 13	113 ± 13
Decoding/word reading	Word attack standard score	115 ± 13	112 ± 14
	Word identification standard score	113 ± 17	109 ± 17
Cognitive abilities	Nonverbal general cognitive ability	109 ± 12	106 ± 13

Supplementary Table 2. Comparison of Linear Mixed Effects Models for Gray Matter Volume							
Brain Region/Tract	Random Intercept			Random Intercept and Slope			
A. Gray Matter Volume	Logarithmic	Linear	Quadratic	Logarithmic	Linear	Quadratic	p
bankssts	3264	3332	3292	**3242**	3330	3283	7.97E-14
fusiform	4143	4266	4160	**4116**	4270	4131	1.72E-12
inferiorparietal	4459	4581	4478	**4456**	4589	4482	1.37E-07
middletemporal	4265	4419	4303	**4233**	4412	4279	3.59E-18
parsopercularis	3913	4016	3955	**3866**	4016	3922	2.09E-22
parstriangularis	3797	3943	3812	**3777**	3946	3797	2.77E-10
superiortemporal	4354	4490	4400	**4335**	4492	4387	6.55E-17
supramarginal	4201	4325	4235	**4178**	4326	4207	3.43E-15
B. White Matter Volume							
bankssts	3311	3356	3334	**3253**	3326	3292	1.09E-20
fusiform	3765	3799	3772	**3706**	3777	3733	2.60E-17
inferiorparietal	4030	4081	4046	**3968**	4047	4006	6.51E-20
middletemporal	3735	3759	3734	**3664**	3718	3686	4.74E-18
parsopercularis	3639	3685	3653	**3552**	3636	3586	5.60E-25
parstriangularis	3396	3422	3399	**3285**	3360	3311	8.61E-28
superiortemporal	4031	4044	4039	**3949**	3977	3966	1.30E-22
supramarginal	4171	4218	4188	**4068**	4168	4103	3.70E-29
C. Surface Area							
bankssts	2923	2970	2943	**2900**	2963	2931	4.09E-12
fusiform	3523	3641	3554	**3503**	3644	3547	6.85E-14
inferiorparietal	3807	3902	3831	**3793**	3903	3829	6.81E-11
middletemporal	3517	3595	3552	**3511**	3595	3552	1.10E-11
parsopercularis	3189	3267	3225	**3150**	3254	3207	3.07E-19
parstriangularis	3110	3202	3143	**3072**	3194	3124	2.80E-18
superiortemporal	3555	3639	3591	**3536**	3632	3580	7.96E-15
supramarginal	3645	3724	3676	**3615**	3714	3650	3.24E-16
D. Cortical Thickness							
bankssts	-235	-226	-225	**-258**	-233	-235	4.62E-10
fusiform	-392	-372	-389	**-425**	-394	-409	9.38E-11
inferiorparietal	-385	-375	-382	**-403**	-386	-395	2.34E-07
middletemporal	-268	-217	-260	**-290**	-228	-288	2.87E-09
parsopercularis	-335	-321	-322	**-361**	-332	-341	2.85E-11

parstriangularis	-298	-277	-298	**-324**	-287	-314	1.30E-08
superiortemporal	-336	-283	-319	**-348**	-286	-330	4.38E-09
supramarginal	-334	-315	-330	**-339**	-314	-333	1.33E-04
E. Mean Curvature							
bankssts	-1611	-1594	-1615	**-1649**	-1614	-1654	2.21E-11
fusiform	-1448	-1432	-1447	**-1472**	-1439	-1471	4.21E-08
inferiorparietal	-1498	-1484	-1494	**-1554**	-1518	-1543	6.66E-16
middletemporal	-1435	-1423	-1435	**-1447**	-1426	-1448	1.72E-07
parsopercularis	-1505	-1495	-1502	**-1536**	-1513	-1526	2.98E-10
parstriangularis	-1482	-1470	-1479	**-1502**	-1478	-1492	9.68E-08
superiortemporal	-1525	-1513	-1525	**-1533**	-1515	-1535	5.17E-07
supramarginal	-1441	-1425	-1444	**-1469**	-1440	-1475	2.17E-10
F. Fractional Anisotropy							
Arcuate fasc.	**-1740**	-1564	-1685	-1733	-1557	-1685	NA
Sup. long. fasc.	**-1749**	-1530	-1700	-1742	-1525	-1713	NA
Inf. long. fasc.	-1667	-1526	-1607	**-1662**	-1523	-1611	NA
G. Mean Diffusivity							
Arcuate fasc.	-6706	-6501	-6593	**-6750**	-6525	-6647	2.44E-37
Sup. long. fasc.	-6950	-6711	-6816	**-7001**	-6741	-6868	2.14E-43
Inf. long. fasc.	-6384	-6214	-6294	**-6410**	-6232	-6321	2.76E-28
Bold indicates lowest BIC value; I, intercept; S, slope							

Supplementary Table 3. Brain-behavior associations with nonlinear mixed effects model						
Brain Region/Tract	Intercept			Asymptote		
A. Gray Matter Volume	r	p	pFDR	r	p	pFDR
bankssts	0.32	8.8E-03	3.4E-02	0.28	2.0E-02	5.3E-02
fusiform	0.02	8.6E-01	8.6E-01	0.08	4.8E-01	4.8E-01
inferiorparietal	0.28	1.3E-02	3.4E-02	0.28	1.3E-02	5.1E-02
middletemporal	0.20	8.2E-02	1.1E-01	0.20	8.2E-02	1.1E-01
parsopercularis	0.17	1.4E-01	1.6E-01	0.22	5.9E-02	1.1E-01
parstriangularis	0.29	9.4E-03	3.4E-02	0.32	5.1E-03	4.0E-02
superiortemporal	0.22	5.0E-02	8.0E-02	0.19	9.6E-02	1.1E-01
supramarginal	0.28	1.7E-02	3.4E-02	0.20	9.5E-02	1.1E-01
B. White Matter Volume						
bankssts	0.26	2.6E-02	6.6E-02	0.26	2.7E-02	2.7E-02
fusiform	0.11	3.4E-01	3.4E-01	0.30	8.0E-03	1.3E-02
inferiorparietal	0.30	1.0E-02	5.0E-02	0.43	1.1E-04	5.4E-04
middletemporal	NA	NA	NA	NA	NA	NA
parsopercularis	0.12	3.0E-01	3.4E-01	0.29	1.1E-02	1.4E-02
parstriangularis	0.23	4.8E-02	8.0E-02	0.40	4.8E-04	1.2E-03
superiortemporal	NA	NA	NA	NA	NA	NA
supramarginal	NA	NA	NA	NA	NA	NA
C. Surface Area						
bankssts	0.30	1.4E-02	2.7E-02	0.29	1.5E-02	2.9E-02
fusiform	0.14	2.4E-01	2.4E-01	0.20	8.4E-02	9.6E-02
inferiorparietal	0.35	1.8E-03	7.2E-03	0.39	5.4E-04	2.3E-03
middletemporal	0.22	6.2E-02	8.3E-02	0.23	5.0E-02	8.1E-02
parsopercularis	0.21	7.8E-02	8.9E-02	0.21	7.2E-02	9.6E-02
parstriangularis	0.39	5.8E-04	4.6E-03	0.39	5.8E-04	2.3E-03
superiortemporal	0.30	1.1E-02	2.7E-02	0.30	1.0E-02	2.7E-02
supramarginal	0.26	3.0E-02	4.9E-02	0.20	9.7E-02	9.7E-02
D. Mean Diffusivity*						
Arcuate fasc.	-0.22	3.3E-02	NA	0.03	6.5E-01	NA
*averaged across $p_{FWE} < 0.05$ significant nodes						

Supplementary Table 4. Contributions of outcomes to linear mixed effects model						
Brain Region/Tract	Main Effect			Interaction		
A. Gray Matter Volume	Effect size	p	pFDR	Effect size	p	pFDR
bankssts	1.8E+01	5.5E-04	4.4E-03	-1.1E+00	5.2E-01	9.0E-01
fusiform	1.1E+01	5.2E-01	5.2E-01	-7.4E-01	8.7E-01	9.0E-01
inferiorparietal	4.0E+01	7.3E-02	1.4E-01	3.1E+00	5.8E-01	9.0E-01
middletemporal	2.9E+01	6.6E-02	1.4E-01	-6.7E-01	9.0E-01	9.0E-01
parsopercularis	1.7E+01	1.4E-01	1.9E-01	-7.1E-01	8.2E-01	9.0E-01
parstriangularis	8.3E+00	2.1E-01	2.4E-01	2.4E+00	1.8E-01	9.0E-01
superiortemporal	3.4E+01	8.7E-02	1.4E-01	-3.2E+00	6.2E-01	9.0E-01
supramarginal	4.5E+01	1.4E-02	5.5E-02	-2.3E+00	7.0E-01	9.0E-01
B. White Matter Volume						
bankssts	3.3E+00	3.3E-01	3.7E-01	1.9E+00	1.8E-01	1.8E-01
fusiform	-1.0E+01	3.5E-01	3.7E-01	7.3E+00	1.5E-02	1.7E-02
inferiorparietal	-2.9E+01	2.7E-02	1.3E-01	2.0E+01	7.3E-07	5.8E-06
middletemporal	-1.8E+01	9.2E-02	1.7E-01	9.9E+00	1.7E-03	3.5E-03
parsopercularis	-9.8E+00	5.0E-02	1.3E-01	5.0E+00	5.4E-03	7.2E-03
parstriangularis	-7.2E+00	1.1E-01	1.7E-01	5.5E+00	1.6E-04	6.2E-04
superiortemporal	-2.3E+01	3.3E-02	1.3E-01	1.3E+01	2.4E-04	6.3E-04
supramarginal	-1.1E+01	3.7E-01	3.7E-01	1.2E+01	4.9E-03	7.2E-03
C. Surface Area						
bankssts	4.9E+00	7.8E-03	6.2E-02	-2.1E-02	9.7E-01	9.7E-01
fusiform	1.4E+00	7.3E-01	7.3E-01	1.0E+00	3.5E-01	5.6E-01
inferiorparietal	9.4E+00	2.1E-01	4.2E-01	3.2E+00	1.1E-01	4.3E-01
middletemporal	3.0E+00	5.5E-01	6.2E-01	1.4E+00	2.9E-01	5.6E-01
parsopercularis	1.6E+00	4.5E-01	6.0E-01	4.8E-01	5.3E-01	7.0E-01
parstriangularis	1.9E+00	2.0E-01	4.2E-01	1.1E+00	2.6E-02	2.1E-01
superiortemporal	4.5E+00	3.5E-01	5.6E-01	1.8E+00	2.2E-01	5.6E-01
supramarginal	1.4E+01	1.9E-02	7.6E-02	-7.2E-01	7.1E-01	8.1E-01
D. Mean Diffusivity*						
Arcuate fasc.	-2.4E-06	9.2E-03	NA	6.0E-07	6.9E-03	NA
*averaged across p_{FWE} < 0.05 significant nodes						

Supplementary Table 5. Brain-behavior associations controlling for TIV				
Brain Region	Measure-Curve Feature	r	p	pFDR
bankssts	GMV-intercept	0.37	2.1E-03	1.7E-02
fusiform	WMV-slope	0.21	6.8E-02	1.1E-01
inferiorparietal	WMV-slope	0.41	2.7E-04	2.2E-03
middletemporal	WMV-slope	0.23	5.0E-02	1.0E-01
parsopercularis	WMV-slope	0.19	1.1E-01	1.4E-01
parstriangularis	WMV-slope	0.35	3.0E-03	1.2E-02
superiortemporal	WMV-slope	0.23	4.6E-02	1.0E-01
supramarginal	WMV-slope	0.15	1.9E-01	2.2E-01
inferiorparietal	SA-intercept	0.31	6.7E-03	2.7E-02
parstriangularis	SA-intercept	0.35	2.4E-03	1.9E-02
superiortemporal	SA-intercept	0.26	2.9E-02	7.5E-02
inferiorparietal	SA-slope	0.20	9.2E-02	3.7E-01
parstriangularis	SA-slope	0.24	4.3E-02	3.4E-01
superiortemporal	SA-slope	0.01	9.6E-01	9.9E-01

Supplementary Table 6. Brain-behavior associations controlling for Euler numbers				
Brain Region	Measure-Curve Feature	r	p	pFDR
bankssts	GMV-intercept	0.38	1.3E-03	1.1E-02
fusiform	WMV-slope	0.33	3.7E-03	4.9E-03
inferiorparietal	WMV-slope	0.50	5.4E-06	4.3E-05
middletemporal	WMV-slope	0.36	1.6E-03	3.1E-03
parsopercularis	WMV-slope	0.30	8.9E-03	1.0E-02
parstriangularis	WMV-slope	0.46	5.1E-05	2.1E-04
superiortemporal	WMV-slope	0.38	6.1E-04	1.6E-03
supramarginal	WMV-slope	0.33	3.4E-03	4.9E-03
inferiorparietal	SA-intercept	0.32	5.3E-03	2.1E-02
parstriangularis	SA-intercept	0.36	1.7E-03	1.4E-02
superiortemporal	SA-intercept	0.28	1.6E-02	4.2E-02
inferiorparietal	SA-slope	0.33	3.1E-03	1.3E-02
parstriangularis	SA-slope	0.37	1.2E-03	9.7E-03
superiortemporal	SA-slope	0.27	1.9E-02	5.0E-02

Supplementary Table 7. Associations between brain measures and literacy-related and cognitive (sub)skills							
		Phonological processing		Oral language		Nonverbal general cognitive ability	
Brain Region/Tract	Measure-Curve Feature	r	p	r	p	r	p
bankssts	GMV-intercept	0.38	1.5E-03	0.17	2.2E-01	0.09	5.4E-01
fusiform	WMV-slope	0.32	4.8E-03	0.13	3.4E-01	0.18	1.9E-01
inferiorparietal	WMV-slope	0.48	1.7E-05	0.28	4.4E-02	0.09	5.3E-01
middletemporal	WMV-slope	0.36	1.7E-03	0.20	1.7E-01	0.14	3.0E-01
parsopercularis	WMV-slope	0.30	8.8E-03	0.16	2.5E-01	0.25	7.3E-02
parstriangularis	WMV-slope	0.43	1.3E-04	0.15	2.8E-01	0.21	1.3E-01
superiortemporal	WMV-slope	0.36	1.4E-03	0.14	3.3E-01	0.13	3.6E-01
supramarginal	WMV-slope	0.29	9.7E-03	0.09	5.2E-01	0.24	8.3E-02
inferiorparietal	SA-intercept	0.33	3.8E-03	0.00	1.0E+00	0.07	6.0E-01
parstriangularis	SA-intercept	0.35	2.2E-03	0.25	7.7E-02	0.26	5.7E-02
superiortemporal	SA-intercept	0.28	1.7E-02	0.19	1.8E-01	0.30	2.7E-02
inferiorparietal	SA-slope	0.35	2.0E-03	0.22	1.2E-01	0.00	1.0E+00
parstriangularis	SA-slope	0.36	1.6E-03	0.30	3.2E-02	0.24	8.6E-02
superiortemporal	SA-slope	0.27	2.0E-02	0.16	2.7E-01	0.17	2.1E-01
Arcuate fasc.	MD-slope	0.24	1.5E-02	NA	NA	NA	NA
*averaged across p_{FWE} < 0.05 significant nodes; NA indicates no nodes were significant after correction							

Supplementary Table 8. Contributions of literacy-related covariates to growth curves					
		FHD		HLE	
Brain Region/Tract	Measure	Effect size	p	Effect size	p
bankssts	GMV	6.8E+01	5.9E-01	2.2E+02	1.7E-01
fusiform	WMV	1.0E+02	5.1E-01	1.4E+02	4.8E-01
inferiorparietal	WMV	-1.1E+02	6.7E-01	3.9E+02	2.5E-01
middletemporal	WMV	-8.4E+01	6.5E-01	5.5E+02	2.1E-02
parsopercularis	WMV	7.0E+01	4.2E-01	9.6E+00	9.3E-01
parstriangularis	WMV	1.9E+02	2.5E-02	-3.0E+01	7.9E-01
superiortemporal	WMV	3.4E+01	8.6E-01	1.8E+02	4.8E-01
supramarginal	WMV	-2.7E+02	3.2E-01	3.5E+02	3.2E-01
inferiorparietal	SA	1.0E+01	9.5E-01	2.2E+02	2.8E-01
parstriangularis	SA	8.2E+01	3.1E-02	5.3E+00	9.2E-01
superiortemporal	SA	6.5E+01	5.1E-01	1.0E+02	4.2E-01
Arcuate fasc.	MD	NA	NA	NA	NA
FHD, family history of reading difficulty					
HLE, home literacy environment					
NA indicates no nodes were significant after FWE correction					

Supplementary Table 9. Comparison of Linear Mixed Effects Models with versus without Literacy-Related Covariates				
Brain Region/Tract	Measure	without covariate	with covariate	p
A. Family history of reading difficulty				
bankssts	GMV	**2328.2**	2332.5	3.7E-01
fusiform	WMV	**2505.3**	2509.9	5.1E-01
inferiorparietal	WMV	**2670.1**	2675.0	6.7E-01
middletemporal	WMV	**2496.7**	2501.6	6.6E-01
parsopercularis	WMV	**2311.7**	2315.9	3.4E-01
parstriangularis	WMV	**2274.7**	2275.0	2.8E-02
superiortemporal	WMV	**2530.3**	2535.3	8.6E-01
supramarginal	WMV	**2647.6**	2651.3	2.4E-01
inferiorparietal	SA	**2451.3**	2456.4	9.7E-01
parstriangularis	SA	**2012.9**	2013.6	3.6E-02
superiortemporal	SA	**2354.4**	2359.0	5.0E-01
Arcuate fasc.	MD	**-3358.7**	-3354.7	2.6E-01
B. Home literacy Environment				
bankssts	GMV	**2328.2**	2331.4	1.7E-01
fusiform	WMV	**2505.3**	2509.9	4.9E-01
inferiorparietal	WMV	**2670.1**	2673.9	2.5E-01
middletemporal	WMV	2496.7	**2496.4**	2.1E-02
parsopercularis	WMV	**2311.7**	2316.8	9.3E-01
parstriangularis	WMV	**2274.7**	2279.7	7.9E-01
superiortemporal	WMV	**2530.3**	2534.8	4.8E-01
supramarginal	WMV	**2647.6**	2651.8	3.2E-01
inferiorparietal	SA	**2451.3**	2455.3	2.9E-01
parstriangularis	SA	**2012.9**	2018.0	9.2E-01
superiortemporal	SA	**2354.4**	2358.8	4.2E-01
Arcuate fasc.	MD	**-3358.7**	-3353.4	1.0E+00
Bold indicates lower BIC value				

Supplementary Table 10. Indirect effects between brain and decoding via phonological processing

Brain Region/Tract	Measure-Curve Feature	Effect size	Lower CI	Upper CI	p
bankssts	GMV-intercept	1.7E-02	3.4E-03	3.8E-02	1.1E-02
fusiform	WMV-slope	6.8E-03	-1.0E-03	1.9E-02	9.4E-02
inferiorparietal	WMV-slope	9.1E-03	5.5E-04	2.2E-02	3.2E-02
middletemporal	WMV-slope	9.4E-03	6.1E-04	2.3E-02	3.4E-02
parsopercularis	WMV-slope	2.0E-02	3.0E-03	4.1E-02	1.6E-02
parstriangularis	WMV-slope	2.2E-02	2.8E-03	4.9E-02	1.6E-02
superiortemporal	WMV-slope	8.3E-03	7.8E-04	2.2E-02	2.0E-02
supramarginal	WMV-slope	5.0E-03	-1.3E-03	1.3E-02	1.2E-01
inferiorparietal	SA-intercept	8.5E-03	-9.7E-04	1.5E-02	8.8E-02
parstriangularis	SA-intercept	4.6E-02	2.3E-03	1.0E-01	3.6E-02
superiortemporal	SA-intercept	1.6E-02	-1.9E-03	3.6E-02	8.3E-02
inferiorparietal	SA-slope	2.3E-02	2.0E-03	5.9E-02	2.4E-02
parstriangularis	SA-slope	7.4E-02	3.7E-03	1.8E-01	3.4E-02
superiortemporal	SA-slope	1.8E-02	-3.8E-03	4.6E-02	1.1E-01
Arcuate fasc.*	MD-slope	2.6E+05	6.8E+04	6.2E+05	1.4E-02
*averaged across p_{FWE} < 0.05 significant nodes					

Supplementary Table 11. Indirect effects between brain and word reading via phonological processing					
Brain Region/Tract	Measure-Curve Feature	Effect size	Lower CI	Upper CI	p
bankssts	GMV-intercept	2.6E-02	9.2E-03	5.4E-02	4.0E-04
fusiform	WMV-slope	9.7E-03	-1.3E-03	2.5E-02	8.3E-02
inferiorparietal	WMV-slope	1.4E-02	2.7E-03	3.2E-02	7.0E-03
middletemporal	WMV-slope	1.4E-02	1.6E-03	3.3E-02	2.6E-02
parsopercularis	WMV-slope	2.9E-02	6.2E-03	5.8E-02	1.1E-02
parstriangularis	WMV-slope	3.4E-02	6.9E-03	7.2E-02	1.3E-02
superiortemporal	WMV-slope	1.3E-02	3.2E-03	3.2E-02	4.2E-03
supramarginal	WMV-slope	7.1E-03	-1.6E-03	1.8E-02	1.0E-01
inferiorparietal	SA-intercept	1.2E-02	3.8E-04	2.1E-02	4.5E-02
parstriangularis	SA-intercept	6.6E-02	3.3E-03	1.4E-01	4.1E-02
superiortemporal	SA-intercept	2.6E-02	3.1E-03	4.9E-02	2.6E-02
inferiorparietal	SA-slope	3.7E-02	7.1E-03	8.6E-02	1.3E-02
parstriangularis	SA-slope	1.1E-01	5.3E-03	2.5E-01	3.9E-02
superiortemporal	SA-slope	3.1E-02	-3.0E-04	6.6E-02	5.2E-02
Arcuate fasc.*	MD-slope	3.1E+05	7.0E+04	6.8E+05	2.0E-02
*averaged across p_{FWE} < 0.05 significant nodes					

Supplementary Table 12. Indirect effects between brain and decoding via phonological processing controlling for TIV					
Brain Region	Measure-Curve Feature	Effect size	Lower CI	Upper CI	p
bankssts	GMV-intercept	1.7E-02	3.2E-03	3.8E-02	1.0E-02
fusiform	WMV-slope	3.5E-03	-4.6E-03	1.6E-02	3.7E-01
inferiorparietal	WMV-slope	8.5E-03	-1.5E-05	2.5E-02	5.1E-02
middletemporal	WMV-slope	7.1E-03	-3.1E-03	2.3E-02	1.6E-01
parsopercularis	WMV-slope	1.4E-02	-5.3E-04	3.7E-02	6.5E-02
parstriangularis	WMV-slope	1.7E-02	-1.7E-03	4.6E-02	8.5E-02
superiortemporal	WMV-slope	7.7E-03	-5.2E-04	2.2E-02	7.9E-02
supramarginal	WMV-slope	2.2E-03	-5.9E-03	1.1E-02	5.6E-01
inferiorparietal	SA-intercept	8.4E-03	-1.2E-03	1.5E-02	1.0E-01
parstriangularis	SA-intercept	4.5E-02	1.6E-03	1.1E-01	3.9E-02
superiortemporal	SA-intercept	1.6E-02	-3.4E-03	3.7E-02	1.0E-01
inferiorparietal	SA-slope	1.8E-02	-1.4E-03	5.9E-02	7.8E-02
parstriangularis	SA-slope	5.2E-02	-1.2E-02	1.7E-01	1.4E-01
superiortemporal	SA-slope	8.3E-03	-3.1E-02	4.3E-02	6.7E-01

Supplementary Table 13. Indirect effects between brain and word reading via phonological processing controlling for TIV					
Brain Region	Measure-Curve Feature	Effect size	Lower CI	Upper CI	p
bankssts	GMV-intercept	2.6E-02	8.4E-03	5.3E-02	1.7E-03
fusiform	WMV-slope	5.0E-03	-6.6E-03	2.0E-02	3.8E-01
inferiorparietal	WMV-slope	1.3E-02	1.4E-03	3.3E-02	2.0E-02
middletemporal	WMV-slope	1.0E-02	-4.9E-03	3.2E-02	1.5E-01
parsopercularis	WMV-slope	2.1E-02	-8.0E-04	4.9E-02	6.0E-02
parstriangularis	WMV-slope	2.6E-02	-3.8E-03	6.3E-02	9.3E-02
superiortemporal	WMV-slope	1.2E-02	-2.4E-04	3.2E-02	5.6E-02
supramarginal	WMV-slope	2.9E-03	-8.7E-03	1.4E-02	5.7E-01
inferiorparietal	SA-intercept	1.2E-02	-6.5E-05	2.1E-02	5.1E-02
parstriangularis	SA-intercept	6.6E-02	2.5E-03	1.4E-01	4.2E-02
superiortemporal	SA-intercept	2.6E-02	5.2E-04	5.1E-02	4.7E-02
inferiorparietal	SA-slope	2.8E-02	-1.3E-03	8.5E-02	6.1E-02
parstriangularis	SA-slope	7.2E-02	-2.6E-02	2.2E-01	1.7E-01
superiortemporal	SA-slope	1.4E-02	-4.3E-02	6.4E-02	6.0E-01

Supplementary Figures

Supplementary Figure 1. Distribution of home literacy environment variables. Histograms depict the distribution of home literacy environment questionnaire responses across five developmental timepoints. New England data only, as these data were not collected in the Calgary dataset.

Supplementary Figure 2. Distribution of home literacy environment scores. Histogram depicts the distribution of home literacy environment scores normalized from the responses shown in Supplementary Figure 1. New England data only, as these data were not collected in the Calgary dataset.

Supplementary Figure 3. Distribution of raw literacy-related outcomes. Histograms depict the distributions of New England data for (A) subtests constituting phonological processing and oral language composite scores and KBIT-2, and (B) word ID and word attack assessments. (C) Histogram for raw phonological processing scores from the Calgary dataset. WJ-IV, Woodcock-Johnson edition IV; KBIT-2, Kaufman Brief Intelligence Test edition 2; WRMT, Woodcock Reading Mastery Tests; NEPSY-II, Neuropsychological Assessment.

Supplementary Figure 4. Distribution of standardized literacy-related outcomes. Histograms depict the distributions of for (A) literacy-related subskills, (B) decoding/word reading assessments and (C) nonverbal general cognitive abilities. Phonological processing scores sum WJ-IV estimates from the New England dataset and NEPSY-II estimates from the Calgary dataset. WJ-IV, Woodcock-Johnson edition IV; NEPSY-II, Neuropsychological Assessment; WRMT, Woodcock Reading Mastery Tests; KBIT-2, Kaufman Brief Intelligence Test edition 2.

Supplementary Figure 5. Infant Structural Processing Pipeline Overview. Raw images without visual artifacts were processed using a combination of iBEATv2.0 Docker, Infant FreeSurfer, FreeSurfer, and in-house scripts. Cortical surfaces were visually inspected for tissue classification accuracy (please see Methods section for details).

Diffusion Images + reverse PE → **MRtrix3 Preprocessing and FODs Generation**

Tissue Segmentation → **MRtrix Anatomically Constrained Tractography** → **py(Baby)AFQ**

Arcuate Fasciculus

Supplementary Figure 6. Diffusion Processing Pipeline Overview. Diffusion data were denoised and then corrected for susceptibility distortions, eddy currents, head motion, and intensity inhomogeneity using MRtrix with FSL and ANTs implementations. Fiber orientation densities (FODs) were generated using constrained spherical deconvolution. Fibers were tracked with Anatomically Constrained Tractography, which leveraged the tissue segmentations from the structural processing pipeline (Supplementary Figure 1): the hybrid segmentation for brains < 50 months and the standard FreeSurfer segmentation for brains > 50 months. Fibers were segmented into tracts using pyBabyAFQ for brains ≤ 24 months and pyAFQ for brains > 50 months (the left arcuate fasciculus is depicted as an example). PE, phase encoding.

Supplementary Figure 7. Statistical Analysis. (A) Brain estimates for regions and tracts of interest, (B) cleaning changes that are not neurophysiologically plausible. Remaining observations are then bridged (pink segment). Please see Methods for determining whether to remove the estimate preceding or following effects models, here using logarithmic functions, to generate (D) individual growth curves. (E) Associations between features (i.e., random intercepts and slopes) regions/tracts with significant brain-behavior associations were divided into low (< 85, dotted line), average (85 – 115, solid line), and high (> 115, dashed line) groups within groups, and plotted.

Supplementary Figure 9. Distribution of curve features of gray matter volume. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 10. Distribution of curve features of white matter volume. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 11. Distribution of curve features of surface area. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 12. Distribution of curve features of cortical thickness. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 13. Distribution of curve features of mean curvature. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 14. Distribution of curve features of fractional anisotropy. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 15. Distribution of curve features of mean diffusivity. Top row depicts histograms for curve intercepts. Bottom row depicts histograms for curve slopes. Brain estimates were scaled for visualization purposes, but skewness was not altered.

Supplementary Figure 16. Age distributions by modality of longitudinal datasets from infancy to early adolescence. All children had (A) structural and/or (B) diffusion MRI data from at least two observations (dots). Blue, New England cohort; orange, Calgary cohort.

Supplementary Figure 17. Longitudinal trajectories of gray matter volume between infancy and early adolescence. Raw estimates of gray matter volume (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue line).

Supplementary Figure 18. Longitudinal trajectories of white matter volume between infancy and early adolescence. Raw estimates of white matter volume (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue line).

Supplementary Figure 19. Longitudinal trajectories of surface area between infancy and early adolescence. Raw estimates of surface area (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue line).

Supplementary Figure 20. Longitudinal trajectories of cortical thickness between infancy and early adolescence. Raw estimates of cortical thickness (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue line).

Supplementary Figure 21. Longitudinal trajectories of mean curvature between infancy and early adolescence. Raw estimates of mean curvature (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue line).

Supplementary Figure 22. Longitudinal trajectories of fractional anisotropy and mean diffusivity between infancy and early adolescence. Raw diffusion estimates (gray spaghetti plot backdrop) were submitted to linear mixed effects models using a logarithmic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (green line).

Supplementary Figure 23. Statically significant associations between growth curve features of brain volume and surface area and preschool/early kindergarten phonological processing. (A) Gray matter volume in the left banks of the superior temporal sulcus exhibited an association between growth curve intercepts and phonological processing, whereas (C) all associations with white matter volume were between growth curve slopes and phonological processing. Both surface area intercepts (B) and slopes (D) exhibited associations with phonological phonological processing. All associations pass $p_{FDR} < 0.05$. WJ-IV, Woodcock-Johnson IV.

Supplementary Figure 24. Statistically significant associations between growth curve features of white matter organization and preschool/early kindergarten phonological processing skill. Mean diffusivity in the left arcuate fasciculus exhibited associations between growth curve slopes and phonological processing. Association is cluster-level corrected at $p_{FWE} < 0.05$. Note: removal of outlier (orange circle) did not abolish cluster-level significance. WJ-IV, Woodcock-Johnson IV.

Supplementary Figure 25. Longitudinal trajectories of gray matter volume between infancy and early adolescence generated for sensitivity analyses. Raw estimates of gray matter volume (gray spaghetti plot backdrop) were submitted to nonlinear mixed effects models using an asymptotic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue lines).

Supplementary Figure 26. Longitudinal trajectories of white matter volume between infancy and early adolescence generated for sensitivity analyses. Raw estimates of white matter volume (gray spaghetti plot backdrop) were submitted to nonlinear mixed effects models using an asymptotic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue lines).

Supplementary Figure 27. Longitudinal trajectories of surface area between infancy and early adolescence generated for sensitivity analyses. Raw estimates of surface area (gray spaghetti plot backdrop) were submitted to nonlinear mixed effects models using an asymptotic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (blue lines).

Supplementary Figure 28. Longitudinal trajectories of mean diffusivity between infancy and early adolescence generated for sensitivity analyses. Raw estimates of mean diffusivity (gray spaghetti plot backdrop) were submitted to a nonlinear mixed effects model using an asymptotic function. Individual growth curves predicted by this model were averaged to show the overall longitudinal trajectory of the sample (green lines).

Supplementary Figure 29. Longitudinal trajectories of brain structure from infancy to early adolescence according to phonological processing skill in preschool/early kindergarten for sensitivity analyses. Graphs depict average trajectories for children with low (< 85), average (85 – 115), and high (> 115) standardized phonological processing scores for measures and regions whose (A) intercepts, (B) slopes, or (C) both intercepts and asymptotes correlated with phonological processing (p_{FDR} < 0.05). Correlation statistics are reported adjacent to their corresponding plots; intercept and asymptote statistics for surface area averaged here for visualization purposes but reported separately in Supplemental Table 3.

Supplementary Figure 30. Statistically significant associations between growth curve features of white matter organization and preschool/early kindergarten phonological processing for sensitivity analyses. Mean diffusivity in the left arcuate fasciculus exhibited a negative association between growth curve intercepts and phonological processing. Association is cluster-level corrected at $p_{FWE} < 0.05$. WJ-IV, Woodcock-Johnson IV.

arcuate fasciculus

Phonological processing skill: Low   Average   High

Supplementary Figure 31. Longitudinal trajectories of mean diffusivity from infancy to early adolescence according to phonological processing skill in preschool/early kindergarten for sensitivity analyses. Graph depicts average trajectories for children with low (< 85), average (85 – 115), and high (> 115) standardized phonological processing scores for the left arcuate fasciculus nodes whose intercepts correlated negatively with phonological processing (p_{FWE} cluster-level < 0.05).

Supplementary Figure 32. Maps of brain-behavior associations. Brain maps show associations between curve features (intercepts and slopes) and preschool/early kindergarten phonological processing across brain regions in the Desikan-Killiany atlas for gray matter volume, white matter volume, and surface area. Note: no model convergence for superior frontal volume and no white matter volume measures were available for anterior cingulate cortex.