1   Hierarchical population genetic structure and signatures of adaptation in

2   *Lates calcarifer*

4   Matthew A. Campbell[1*] and Joy A. Becker[1]

6   [1] The University of Sydney, School of Life and Environmental Sciences, 380 Werombi Road,

7   Camden New South Wales 2570, Australia

9   [*] Corresponding Author

10   Matthew A. Campbell, m.campbell@sydney.edu.au; https://orcid.org/0000-0002-5826-0329

12   Joy A. Becker, https://orcid.org/0000-0003-4749-4004

Abstract

Context

Lates calcarifer known as barramundi perch, is widespread in the tropical Indo-Pacific region and is an important taxon in aquaculture, recreational and commercial fisheries. With mitochondrial barcoding study of barramundi perch, three clear mitochondrial clusters have emerged.

Aims

To test for congruent signal between mitochondrial and nuclear genomes and to test for population genetic structure. We further estimated the evolutionary significance of genetic clusters in terms of signatures of adaptation in the nuclear genome.

Methods

We conducted analyses of mitochondrial sequence data and whole-genome sequencing data from across the range of barramundi perch (n=60). We applied phylogenetic, phylogenomic and population genetic methods.

Key Results

We find congruent signal for three lineages of barramundi perch in both the mitochondrial and nuclear genomes. Substantial genetic differentiation is present between and within all three lineages of barramundi perch. Calculation of divergence times indicates a separation of ~ 7 million years ago, and another separation dating to ~1 million years ago.

Conclusions

Three lineages of barramundi perch are apparent, with candidate adaptive loci identified between and within each lineage, indicating both discreteness and evolutionary significance.

Implications

This study highlights the need for comprehensive sampling and integrative study of genotypes and phenotypes across the range of barramundi perch.

Additional Key Words

1.0 Introduction

Lates calcarifer (Bloch 1790) is a valuable commercial taxon with numerous common names given its wide distribution and importance in historic and contemporary subsistence activities. Having been depicted in rock art and Dreamtime stories for millennia, 'barramundi' (Australian Aboriginal word meaning 'large, scaled silver fish') and also known as 'namamkorl' in the Kunwinjku dialect from the peoples in Kakadu and Western Arnhem Land (Northern Territory), hold deep cultural significance in Australian Aboriginal heritage https://pad.katalyst.com.au/plant_or_animals/barramundi (Cluney, 2004; Edwards-Vandenhoek, 2015; Shamsi et al., 2020). The common name 'barramundi' is widely used in Australian English while 'Asian sea bass' is also used in English and scientific works. We use the common name is barramundi perch (BP) in this manuscript.

Barramundi perch is distributed from the Persian Gulf eastwards to Taiwan and Australia, and into Oceania (**Figure 1**). Aquaculture beginning in Thailand in the 1970's has increased greatly and BP is widely propagated across its range with some aquaculture outside its native range. Barramundi perch is also an important taxon for conservation aquaculture with annual restocking in lakes and impoundments across northern Australia (e.g. Manton Dam, Northern Territory; Lake Kununurra, Western Australia) to support recreational fisheries and tourism in remote communities. The BP is a protandrous hemaphrodite, catadromous, and tolerant of a wide range of salinities (Moore, 1979; Moore and Reynold, 1982). The sequential hermaphrodism of the taxon requires the importation of young males into aquaculture facilities as older fish become females, a potential source of hybridization and mixing of stocks of different origin. Other qualities BP such as high fecundity, ease of culture and rapid growth (i.e. harvestable size of 3 kg in less than two years) encourage its production (Rimmer, 2009).

The widespread culture of this species has led to genetic studies to understand population genetic structure of BP. Mitochondrial barcoding revealed a deep divergence between Australian and Myanmar fish (K2P distance ~ 9.50%) (Ward et al., 2008). Broader sampling geographically of mitochondrial lineages has substantiated this divergence, showing a distinctive clade of Indian subcontinent fish including the Bay of Bengal (IND, additional locations the Andaman Islands and Myanmar) separate from Southeast Asia (SEA including Singapore, Malaysia, Thailand and

Indonesia) and Northern Australia (Vij et al., 2014). Vij et al. (2014) did include the sequences

from Ward et al. (2008) but did not include New Guinea, a region known to have biogeographic

similarities to Northern Australia (Wallace, 1859). Studies utilizing more complete genomic and

more complete geographic sampling are very limited. Restriction site-Associated DNA

sequencing (RADseq) was applied to population-level samples of BP (n = 131) from six

localities (Thailand, Malaysia, Indonesia, Western Australia, Eastern Australia and Papua New

Guinea) found a high degree of differentiation of SEA fish from Australia + New Guinea

(AUS+NG). Pairwise F_{st} within SEA was 0.005-0.009 but 0.36-0.443 between SEA and

AUS+NG without evidence of contemporary gene flow (Wang et al., 2016). Some investigation

of population genomic structure was applied to a range-wide data set at a high level as part of a

genome assembly paper but did not investigate concordance between mtDNA and genomic DNA

or population genetic metrics (Vij et al., 2016). No study to date, has analyzed both

mitochondrial data and genomic data from the same individuals to evaluate concordance between

data sources and resolve possible discordance between mitochondrial and genomic signals.

While there is evidence of genetic differentiation of BP, the evolutionary significance of these

differences are not substantiated and there remains a lack of recognition of distinctive species

that has serious implications in management of BP. For example, there is potential range overlap

of the IND lineage with the SEA lineage that has not been evaluated, and the forced

hybridization of these lineages during propagation is possible. They may exist as separate stocks

in a mixed stock fishery or form a natural hybrid zone. There is also the movement of broodstock

that may mix fish of different genetic lineages throughout the range of BP. While the recognition

of species is indeed important, so is the recognition of units below the species level that

contribute to diversity such as ecotypes or life history variants. The ecological effects of

intraspecific variation can meet our exceed species effect e.g. (Des Roches et al., 2018).

Conservation of both species-level and intraspecific variation can serve to promote ecological,

cultural and economic functions of organisms, resilience and genetic variation for future adaptive

potential e.g. (Layton et al., 2021; Lehnert et al., 2023; Schindler et al., 2010). Identification of

both specific and intraspecific variation can be considered both through isolation and adaptation

(Waples, 1991). We follow with an approach to identify units of BP through both isolation, that

is discreteness of higher-level groups, and testing for evolutionary significant differences

between genetic groupings (i.e. adaptive genetic loci) (Lehnert et al., 2023).

106 2.0 Methods

2.1 Whole-Genome Sequence Data Set

Genome-wide sequence data from BP was downloaded from the NCBI Sequence Read Archive

under the BioProject accession PRJNA311498. The 61 samples in PRJNA311498 covered a

broad distribution, from the Indian subcontinent, parts of Southeast Asia, Northern Australia and

Papua New Guinea. Two additional sequences of BP were obtained from Bangladesh

(PRJNA1021005) and for phylogenetic analyses an outgroup obtained (*L. japonicus,* n =1,

PRJDB13763). The paired-end data was aligned to the BP reference genome GCF001640805.2

(TLL_Latcal_v3) with the Burrows-Wheeler aligner (BWA) with the mem algorithm (Li and

Durbin, 2010, 2009). Alignments were sorted, properly paired reads kept and PCR duplicates

removed with SAMTools (Danecek et al., 2021; Li et al., 2009). Total read counts, filtered read

counts and depth of coverage were calculated from alignments with SAMTools. Those samples

with coverage $\leq$ 3X were excluded from further analyses.

2.2 Mitochondrial Phylogeny

A mitochondrial alignment from the WGS data set was created by creating a Binary Alignment

Map (BAM) file for each individual with more than 3X coverage only including reads aligned to

the reference mitochondrial genome (NC_007439.1, samtools view). These BAM files were

converted to consensus FASTQ files with the bam2cns subprogram of proovread (Hackl et al.,

2014) then to FASTA files turning all bases with quality scores below 20 to N with seqtk (seq -a

-q20 -n N) (https://github.com/lh3/seqtk). The mitochondrial Cytochrome Oxidase I (mtCOI)

gene was extracted from the complete mitochondrial FASTA-formatted sequences with

SAMtools (faidx NC_007439.1:6329-7879). For rooting, the corresponding region of the *L.*

japonicus mitochondrial genome was extracted from the NCBI reference sequence

(NC_034339.1) and sequences were aligned with MAFFT (Katoh et al., 2002; Katoh and Toh,

2008). A Maximum-Likelihood (ML) phylogeny of sequences was generated with IQTREE

specifying a Hasegawa-Kishino-Yano model of nucleotide substitution with gamma-distributed

rate variation (-m HKY+G) and support for nodes evaluated with rapid bootstrapping (-bb

10000) (Hoang et al., 2017; Nguyen et al., 2014). The tree was imported into R and visualized

with the ggtree package (Yu et al., 2017) collapsing nodes to polytomies receiving less than 75%

Bootstrap Support (BS) and rooting by the longest branch.

2.3 Genome-Wide Phylogenetic and Network Analyses

We generated a genome-wide set of called SNPs with Analysis of Next Generation Sequencing

Data (ANGSD) from the 24 main linkage groups in the BP assembly (Korneliussen et al., 2014).

Quality of called SNPs was enforced by ensuring that a SNP was present in at least 90% of

individuals, had a minimum Minor Allele Frequency (MAF) of 0.05, a minimum mapping

quality of 20, minimum base quality of 20, a SNP p – val of 1.00 x 10^{-6} and a posterior cutoff of

0.90. A SAMTools model was specified and a PLINK-formatted file created as the output. We

converted the PLINK-formatted file to a VCF-formatted file with PLINK (Purcell et al., 2007).

Linked SNPS were removed with BCFTools +prune (-l 0.20 -w 10000).

The multispecies coalescent was applied to model gene-tree and species-tree discordance

resulting from Incomplete Lineage Sorting (ILS) e.g. (Maddison, 1997) on a set of called SNPs

that included the outgroup *L. japonicus*. A species-tree was generated using SVDQuartets in

PAUP* version 4.0a (Chifman and Kubatko, 2015, 2014; Swofford, 2003). Samples were

pooled by sampling region for the species tree. A random set of quartets (evalQuartets=random),

partially ambiguous SNPs were distributed over all compatible site patterns (ambigs=distribute),

and 1,000 bootstrap replicates (bootstrap=standard nreps=1000) were options specified.

For ML phylogenetic analysis and network analysis we called SNPS with BP samples and a

phylogeny of SNPs was inferred with IQTREE by identifying a model of nucleotide evolution

with ModelFinder Plus and indicating the ascertainment bias correction should be applied (-m

MFP+ASC) (Kalyaanamoorthy et al., 2017; Nguyen et al., 2011). IQTREE was used to further

filter SNPs for suitability for ascertainment bias correction. Support for nodes was evaluated

with 10,000 rapid boostraps (Hoang et al., 2017). As concatenated phylogenetic analyses do not

explicitly model hybridization or ILS, we created an implicit network to identify potential

hybridization or ILS in our data set with SplitsTree (Huson and Bryant, 2005). The same

alignment provided to IQTREE was analyzed with SplitsTree.

2.4 Population Genetic Structure and Statistics

Total signal in the data set was assessed by the generation of covariance matrix with ANGSD

and a PC analysis. The covariance matrix was generated by with a single read sampling approach

due to a wide range of individual coverage (minimum 4.73, maximum 34.81, mean 9.38 and

median 6.41) with the options -doIBs 1 and -doCov 1 in ANGSD. The 24 main linkage groups of

the reference assembly were sampled for this analysis. A variable site was required to be present

in >90% of individuals with a minimum MAF of 0.05. A minimum mapping quality of 10 was

specified with a minimum base quality of 20 and a GATK genotype likelihood model was used

(-GL 2). The resulting covariance matrix was imported into R for PC analysis.

To calculate genetic differentiation in terms of F_{st} was calculated by generating Site Frequency

Spectra (SFS) for each sampling region with ANGSD and the realSFS subprogram, splitting

Indonesia into two locations based on metadata available with the sequences ("K" n=5 and and

"SJ" n=5). Vietnam (n=2), Indonesia "SU" (n=1) and Bangladesh (n=2) were excluded due to

small sample sizes. Two-dimensional SFS were calculated between each possible pair of

sampling regions and F_{st} calculated (see code repository accompanying this paper). Nucleotide

diversity (π) and Tajima's D (*D*) estimates were calculated for each chromosome with realSFS

using the saf2theta option followed by the thetaStat subprogram of ANGSD and the do_stat

option.

Admixture analysis was conducted with NGSadmix (Skotte et al., 2013) on genotype likelihoods

generated with ANGSD as for the PC analysis using a single read sampling approach. The

resulting BEAGLE-formatted file was used with NGSadmix to produce an admixture plot of *K* =

2-5 genetic clusters.

2.5 Tests for Adaptative Genetic Variation

Genotype calls produced for phylogenetic network analysis after pruning were imported into R

for analysis with PCAdapt (Luu et al., 2017; Privé et al., 2020). The *pcadapt* function was then

used to generate a scree plot and identify an objective number of PCs with the first 20 PCs

examined. From the resulting object from the *pcadapt* function, outlier loci were identified by

filtering for a Benjamini and Hochberg (1995) and Bonferroni corrected p – values with

significance values less than 0.10. We repeated this analysis for individuals in each main genetic

lineage by filtering for variants with a minimum minor allele frequency of 0.05 and examining

the first 5 PCs.

2.6 Divergence Time Estimation

Existing efforts have covered a broad geographic and lineage sampling of *L. calcarifer* with

barcoding (mitochondrial *cox1*) (Vij et al., 2014; Ward et al., 2008). The effort of this study was

then to calculate divergence times of the three lineages (IND, SEA and AUS+NG). We

downloaded barcode sequences from the Barcode of Life Database and Genbank for Latidae.

With BLAST (blastn) we confirmed the major lineages of *Lates calcarifer* in barcode data

(Altschul et al., 1997) then selected representative sequences from these three lineages to

conduct a divergence time analysis (**Table 1**). We also searched for representation across Latidae

for broad taxonomic representation of this family including sequences from *L. japonicus*, *L.*

microlepis and *L. niloticus*. We included representatives of the two main mitochondrial lineages

present in *Psammoperca* corresponding to *P. waigiensis* and *P. datnioides*. Centropomidae has

been confirmed to be the sister lineage of Latidae in molecular phylogenetic study e.g.

(Campbell et al., 2013; Chen et al., 2007; Li et al., 2011) and inclusion allows dating the Time to

Most Recent Common Ancestor (TMRCA) of Centropomidae and Latidae to be 48.6 million

years ago (mya) (Otero, 2004). We also included sequences from the outgroup lineages

Channidae and Nandidae as an additional calibration point dating the TMRCA of those taxa to

be 48 mya (Santini et al., 2009).

Sequences were aligned with MAFFT (Katoh et al., 2002; Katoh and Toh, 2008), in frame

coding was confirmed by translation in the NCBI ORF finder and a NEXUS file made for a

time-calibrated phylogenetic analysis in MrBayes (Ronquist et al., 2012). The third codon

position was excluded due to the age of the data set and expected saturation occurring in

mitochondrial sequences from carangimorph taxa at these timescales e.g. (Campbell et al., 2014).

A nucleotide substitution model of General Time Reversible (GTR) with a proportion of

invariant sites (I) and gamma-distributed rate variation (Γ) was specified. Constraints were

applied to Centropomidae + Latidae to date with an exponential distribution used,

offsetexp(48.60, 63.18), and the outgroup, offsetxp(48.00,62.40). Initially, three runs and three

chains were specified with a run length of 1 million generations sampled ever 1,000 generations

and a 25% burnin applied. Evaluation of parameters and convergence with Tracer (Rambaut et

al., 2018) was assessed, and run length increased until sufficient ESS for all parameters (>200)

was reached. Tree summarization was done in MrBayes with the sumt command and an all

compatible consensus tree made.

3.0 Results

3.1 Whole-Genome Sequence Data Set

From the original 63 *L. calcarifer* sequences, 60 had coverage over 3X and were included in

additional analyses (**Table 1**). Sample sizes for regions were 13 for putative members of the

IND lineage, 30 for putative members of the SEA lineage, and 17 for putative members of the

AUS+NG lineage from ten sampling regions (**Figure 1**). The *L. japonicus* sequence aligned to

the *L. calcarifer* reference genome with high success with 110X coverage and was down-

sampled to 20X coverage with SAMtools to be more consistent with genome coverage measures

of *L. calcarifer* individuals.

3.2 Mitochondrial Phylogeny

The 61 samples (60 *L. calcarifer* and one *L. japonicus*) produced an alignment of 1,576 base

pairs in length with 71 distinct patterns and 128 parsimony-informative sites. In the alignment,

1,441 sites were invariant. The ML phylogeny indicates a clear division of *L. calcarifer* IND

individuals with two Cambodian-origin fish from all other fish, BS=85% (**Figure 2**). Support

for monophyly of *L. calcarifer* SEA + *L. calcarifer* AUS+NG is maximal, and support for a

clade of *L. calcarifer* AUS+NG is moderate (BS=80%). Individuals sourced from the putative *L.*

calcarifer SEA clade do not form clear geographic clusters. The Assemble Species by Automatic

Partition algorithm indicated the strongest support for two species, with a threshold distance of

4.10% (results provided in the Data Supplement). This division separates the *L. calcarifer* IND

mitochondrial lineage from *L. calarifer* SEA + *L. calcarifer* AUS+NG.

3.3 Genome-Wide Phylogenetic and Network Analyses

For analysis with SVDQuartets to generate a rooted species-tree, 8,529 variants that all have

unique alignment patterns were analyzed and is presented as **Figure 3**. The SNP alignment

consisted of 6,456 parsimony informative and 2,073 singleton sites. Three lineages are apparent

in the species-tree with maximal support for monophyly corresponding to the *L. calcarifer* IND,

L. calcarifer SEA and *L. calcarifer* AUS+NG. The SEA and AUS+NG lineages are highly

supported (BS=100%) as sister lineages. Within the IND lineage, the India Eastern Coast and

Bangladesh sampling locations are indicated to be sister lineages (BS=100%). The AUS+NG

lineage individuals from the Northern Territory and Papua New Guinea are indicated to be most

closely related (BS=100%). The first division within the SEA lineage is between Indonesia-SU +

Indonesia-K and receives maximal support for monophyly and branching patterns. The

Philippines sampling location separates with the next successive branching within the SEA

lineage (BS=100%). Relationships among Cambodia, Indonesia-SJ, Thailand and Vietnam

sampling locations are not clearly resolved.

For ML phylogenetic analysis and phylogenetic network analysis an alignment of 60 BP

individuals with 7,661 SNPs having 7,661 distinct unique alignment patterns and 5,571

parsimony informative sites was created. The alignment contains 2,090 singleton sites. Support

for monophyly of *L. calcarifer* IND is maximal, as well as geographic division with *L. calcarifer*

IND into Eastern and Western Indian subcontinent (**Figure 4A**). *L. calcarifer* SEA as a grouping

receives low support (BS = 76%) with concordance between sampling locations and placement

not complete. Indonesia-K and the Philippines sampling locations are highly supported to be

monophyletic, but Cambodia, Thailand and Indonesia-SJ are not. Support for a *L. calcarifer*

AUS+NG is maximal, with maximal support for division between Queensland and Northern

Territory + Papua New Guinea fish. Monophyly of samples from the Northern Territory is

poorly supported (BS = 68%).

The phylogenetic network resolves an IND lineage containing two sub-lineages (India Eastern

Coast + Bangladesh, India Western Coast) with separation of all three sampling locations

(**Figure 4B**). The SEA lineage is present, with reticulations with the IND lineage occurring with

a subgroup of 10 individuals, disrupting monophyly of some sampling locations. The Indonesia-

K and Philippines locations are monophyletic, but the Cambodia, Thailand and Indonesia-SJ

sampling locations are not. The AUS+NG grouping is composed of three subclades Queensland,

Northern Territory and Papua New Guinea. In the network, Northern Territory and Papua New

Guinea fish are most-closely related to each other.

3.4 Population Genetic Structure and Statistics

The PC analysis included 7,321,852 variable sites in the genome and indicated a separation into

three larger genetic clusters corresponding to IND, SEA and AUS+NG individuals (**Figure 5**).

The first PC separated AUS+NG, SEA and IND individuals (PC 1 12.57% of variation) with the

second PC showing variation within SEA and IND clusters (PC2 4.71%). Clear separation of

Indian East Coast + Bangladesh and Indian West Coast individuals is present on both PC 2 and

PC 3 (PC 3 3.51%).

Genetic differentiation (F_{st}) was lowest within main genetic lineages, such as Thailand and

Cambodia (F_{st} = 0.06) within the *L. calcarifer* SEA lineage and between the Indian East Coast

and Indian West Coast sampling locations (F_{st} = 0.11) within the *L. calcarifer* IND lineage

(**Figure 6A**). The highest levels of genetic differentiation are found between main genetic

lineages, between AUS+NG and SEA F_{st} = 0.15-0.35, between AUS+NG and IND F_{st} = 0.22-

0.33, between SEA and IND F_{st} = 0.12-0.26). Nucleotide diversity is the lowest in the AUS+NG

lineage (π = 20,067-22,775) and highest in the IND lineage (π = 207,299-214,485) (**Figure**

6B). Tajima's D was strongly negative (< -1) in the Indonesia-SJ, Thailand, India Eastern Coast

and India Western Coast sampling locations (**Figure 6C**). The AUS+NG sampling locations had

D values between -0.58 and -0.77.

Admixture analysis of 7,321,852 genotype likelihoods from 60 BP individuals for K = 2-5 is

presented as **Figure 7**. At K = 2, the Bangladesh, Indian Eastern Coast and Indian Western

Coast have high ancestry coefficients (Q) > 0.99 for a single genetic cluster and the AUS+NG

lineage and Indonesia-K sampling location have high ancestry coefficients for the other. The

remaining SEA sampling locations have a mixture of contributions from both genetic clusters.

At K = 3 the AUS+NG individuals form a clearly genetic cluster with Q > 0.99 for ancestry

coefficients, the SEA sampling locations are heterogenous mixture proportions from sampling

locations largely composed of the third genetic cluster, with the Indonesia-K and Indonesia-SU

sampling locations exhibiting a near uniform ~0.27% admixture proportion with the genetic

cluster typical of AUS+NG sampling locations. Higher values of K lead to separating of genetic

units within the SEA lineage that may or may not correspond to sampling locations.

3.5 Tests for Adaptation

The variation within the pruned genotype calls of 22,088 SNPs from the n=60 WGS sequences

with >3X coverage supported an optimal $K = 3$ PCs (**Supplemental Figure S1**). These PCs

separated the three main lineages of *L. calcarifer*, with the third PC separating the Indonesia-K

sampling location and to a lesser degree the Indonesia-SU individual from other SEA. Benjamini

and Hochberg (1995) adjusted p – values (BH) identified 134 candidate adaptive SNPs and

Bonferroni corrected p – values (BF) indicates 34 candidate adaptive SNPs across the three PC

axes.

Filtering to IND individuals (n=13) resulted in 12,075 SNPs with a minimum MAF of 0.05, with

the first PC separating the Bangladesh, India Eastern Coast and India Western Coast with 738

BH and 379 BF candidate adaptive SNPs identified on this axis (**Supplemental Figure S2**). The

second PC axis separated India Eastern Coast with 285 BH and 59 BF candidate adaptive SNPS.

From the SEA lineage, 30 individuals with 17,556 SNPs with a minimum MAF of 0.05 were

analyzed. The first three PCs were considered, separating Indonesia-K and Indonesia-SU

sampling locations on PC 1, the Philippines along PC 2 and some Thailand and Cambodia

individuals along PC 3 with 190 BH and 29 BF candidate adaptive SNPs identified in total

(**Supplemental Figure S3**). The data set of AUS+NG individuals (n=17) for loci with a

minimum MAF of 0.05 resulted in 14,642 SNPs. We found strong support for $K = 2$ PCs, the

first PC separating Queensland from the Northern Territory and Papua New Guinea

(**Supplemental Figure S4**). The second PC separated Papua New Guinea from the other

locations with 234 BH candidate adaptive SNPs and 57 BF candidate adaptive SNPs across the

two PC axes.

3.6 Divergence Time Estimation

We compiled a barcode alignment of 13 sequences with 660 characters resulting in 440

characters (399 invariant) when including the first and second codon positions only. Sufficient

effective sample sizes (ESS >200, range of 712-3,002) was reached with 20 million generations

and two runs of three chains each as evaluated after a 25% burnin. The resulting consensus tree

presented in **Figure 8.** The median divergence time of *L. calcarifer* IND from the other two

lineages is estimated to be 6.76 mya with a 95% Highest Posterior Density (HPD) of 1.31-14.24

mya. The divergence of *L. calcarifer* AUS+PNG from *L. calcarifer* SEA is estimated to be 1.02

mya with a 95% HPD range of 0.00-5.03 mya. The relationships of *L. calcarifer* lineages are

highly supported, with the placement of *L. calcarifer* having a Posterior Probability (PP) of 0.99,

and the sister relationship between *L. calcarifer* AUS+PNG and *L. calcarifer* SEA having a PP

of 0.97. The NEXUS file for use with MrBayes and the consensus tree are provided in Data

Supplement of this paper.

361 **4.0 Discussion**

We demonstrate concordance of nuclear and mitochondrial genomes of three main genetic

lineages of BP with substantial differences quantified in terms of genome-wide estimates of F_{st},

candidate adaptive loci and deep divergence times. We observed F_{st} values up to 0.36 and in

terms of divergence times, we estimated 6.76 million years for the divergence of IND from other

BP and a divergence time estimate of 1.02 million years for AUS+NG from SEA. The previously

published estimate based on RADseq data for divergence times by Wang et al. (2016) is

restricted to AUS+NG from the SEA lineage and is an approximate 644,100-878,900 years

before present. Both of these estimates, the estimate presented here from fossil-calibrated

phylogeny and Wang et al. (2016), support a divergence during the glacial cycling and changes

in sea level and conditions during the Quaternary. The three main lineages of BP exhibit

discreteness (genetic isolation) and evolutionary significance (i.e. adaptive differentiation).

Within each of the three main lineages, there was additional evidence for discreteness and

evolutionary significance. The IND sampling locations were divided phylogenetically (**Figures 3**

& 4), and in PC analysis (**Supplemental Figure S2**) with candidate adaptive SNPs identified.

Previously, with the IND lineage, substructure was not clear with mitochondrial data and we can

find a clear division within IND (Bangladesh, India Eastern Coast and India Western Coast) with

genomic data. The SEA lineage exhibited notable separation of the Indonesia-K and Indonesia-

SU sampling locations, the Philippines sampling location and some Thailand fish (**Figure 4,**

Supplemental Figure S3). Clear differentiation of all Philippines and Thailand fish may be

impacted by hybridization (**Figure 7**). The RADseq data set of Wang et al. (2016) did not show

clear evidence of genetic structure with three sampling locations within SEA, but broader

sampling with microsatellites did appear to support genetic structuring (Yue et al., 2009). The

AUS+NG sampling exhibited a clear division between Queensland and other sampling locations

(**Figures 3 & 4**) and all three sampling locations could be clearly differentiated on two PC axes

with candidate adaptive loci apparent (**Supplemental Figure S4**). Within the AUS+NG lineage

there are several studies demonstrating genetic differentiation within that lineage e.g.

(Chenoweth et al., 1998; Loughnan et al., 2019; Shaklee and Salini, 1985; Wang et al., 2016)

with 16 discrete genetic populations identified in Australia only (Keenan, 1994).

4.2 What is Lates calcarifer*?*

Our fundamental understanding of nature is based at the species level and the lack of recognition

of species is a major knowledge shortfall in conservation and management (Brown and

Lomolino, 1998). Molecular genetic techniques have continued to identify candidate species of

fishes which may or may not have diagnostic morphological characters typical of species

descriptions e.g. (Campbell et al., 2023; Moyle and Campbell, 2022). Instead of being a

historical aspect of biology, the need for continued taxonomic work is very important e.g. (Delić

et al., 2017; Godfray et al., 2004). Taxonomic work often involves the detailed anatomical

investigation of organisms that can reveal phenotypic differences related not only to speciation,

but also life history variation. What remains outstanding is if BP can be accurately represented

within Linnean taxonomy or if an alternative approach is better suited to the biological realities

of BP.

The most recent taxonomic effort relevant to BP led to the description of *L. lakdiva* Pethiyagoda

& Gill 2012 from western Sri Lanka and *L. usiwara* Pethiyagoda & Gill 2012 from eastern

Myanmar (Pethiyagoda and Gill, 2012). *Lates lakdiva* from Sri Lanka apparently cannot be

propagated in aquaculture using methods used for BP elsewhere indicative of divergences from

those species. The specimens of *L. usiwara* were the same as featured in the barcoding study of

Ward et al. (2008) highlighting deep mitochondrial divergences, but in a wider context the barcode

sequences of *L. usiwara* are identical to samples from the Indian subcontinent (Vij et al., 2014).

However, in comparison to *L. calcarifer*, a differential diagnosis was possible and the distribution of *L. usiwara* is perhaps indicative of a recently derived species or ecotype. The anatomical study undertaken by Vij et al. (2014) along with barcoding did not clearly separate all these newly described species of *Lates* from *L. calcarifer* noting small sample sizes used in the description of *L. usiwara* and *L. lakdiva* and conclude that there is substantial mismatch between anatomical descriptions of new species of *Lates* and candidate species of *Lates* apparent genetically. Barcode sequences, it should be noted, will not reflect proportion of ancestry as do genomic data and ascertaining the ancestry proportions of fish used in taxonomic studies is critical going forward due human translocations of *Lates*. Furthermore, it is possible for a high degree of genome-wide differentiation or substantial reproductive isolation be present that would not be apparent in barcode data. Pethiyagoda and Gill (2012) themselves observe that due to the large body size and sequential hermaphodism of *L. calcarifer sensu lato*, they were restricted to examining many juveniles for their study.

The three genetic lineages of BP remain unclear in distribution and have unclear anatomical differentiation (Vij et al., 2014). Portions of the range, such as Taiwan, remain unsampled for genomic data and genetic structuring is apparent across all three lineages (**Figures 3, 4, 5 and 6, Supplemental Figures S1-S4**). Differentiation between and within the main lineages can be clarified with additional sampling and representation of nominal taxa in genomic data sets (i.e. *L. usiwara* and *L. lakdiva*). Given the strong genetic differentiation and deep inferred time scales (**Figure 8**), one possible resolution is to recognize at least two species-level taxa that correspond to the IND lineage and a combined SEA and AUS+NG lineage. At a potential species-level, the taxonomy of *L. calcarifer* is complex and the recognition of types and their origins is also challenging, reviewed by Pethiyagoda and Gill (2012). The type locality of *L. calcarifer* (Bloch 1790) may be Tamil Nadu, India (10.4°N, 79.3°E) (Fricke et al., 2023). Thus, the *L. calcarifer* IND lineage may be represented by this name. There are numerous available names that may be applicable to *L. calcarifer* SEA, but this requires substantial time and expertise to resolve. There is, however, an available name, *L. cavifrons* (Alleyne & Macleay 1877) that could be applied to *L. calcarifer* AUS+NG. The collection locality is either the Torres Strait or the coast of New Guinea with a lost holotype with the drawing providing the current synonymy (Pethiyagoda and Gill, 2012).

Without formal taxonomic recognition, the three main genetic lineages of BP can be recognized for management purposes such as mixed stock fisheries. Vij et al. (2014) note that the assumption of single species of BP has undoubtedly influenced previous studies and microsatellite data from Thailand showing the highest allelic richness may reflect genotyping of two divergent genetic lineages along the west coast of Thailand (Yue et al., 2009). If different species of BP are recognized, then questions of biology can be researched such as the causes of speciation and what the mechanisms for reproductive isolation of IND may be. Development of an anatomical key usable in the field that can separate these species e.g. (Wang et al., In Revision) would be of great practical utility and could be used to resolve the distributions and overlap of IND and SEA lineages. A high-throughput sequencing assay that can discriminate the main lineages could provide a rapid assessment of genetic lineage, hybridization levels and possible provenance in studies of food adulteration.

4.2 Admixture

Overall, analyses that can quantify hybridization in this study are challenging because of the high innate diversity of the SEA lineage and limited sample sizes, notably that of the IND lineage (n = 13). The sequences were also obtained opportunistically from the NCBI Sequence Read Archive and not collected by the authors. With these cautions in mind, we find evidence of admixture between major genetic lineages within our data set as evidenced by the placement of Cambodia-origin samples, with a largely SEA genomic background within the mitochondrial IND lineage (**Figures 2, 4 & 5**). When examining all 60 individuals at $K = 3$, the two Cambodia individuals with IND mtDNA have ancestry coefficients (Q) of 0.29 and 0.43 with the IND genetic cluster, however, there is such a degree of variation in the SEA group with respect to admixture proportions and inconsistency across sampling locations that these measurements should not be accepted at face value. The Cambodia group has two individuals with Q = 0.00 for the IND genetic cluster and one with 0.17 for example when examining all individuals. Separate analysis of Cambodia and *L. calcarifer* IND individuals only show $Q = 0.23$ and 0.33 for the two Cambodia individuals with IND mtDNA and no evidence for nuclear genomic admixture of other Cambodian individuals with IND (**Figure 7**).

Nucleotide diversity in the SEA Indonesia-SJ location is very high (**Figure 6B)** and no doubt

influences the reduced F_{st} of this lineage to other sampling regions (**Figure 6A**). Combined with

the large negative *D* value which arises from admixture e.g. (Hahn et al., 2013) this sampling

location may contain admixed individuals. A similar pattern of high nucleotide diversity, low-F_{st},

and strongly negative *D* is apparent with the Thailand sampling location (**Figure 6**). Both the

Indonesia-SJ and Thailand sampling locations show substantial contributions of the IND genetic

cluster in their ancestry (**Figure 7**), but it is not uniformly distributed across sampling locations.

If the distribution of ancestry is uniform across sampling locations, it would suggest an older age

of hybridization. The variable admixture proportions in sampling locations suggest recent

hybridization events, e.g. Cambodia at $K = 4$. Furthermore, the admixture analyses suggest that

Thailand at $K = 4$ is represented by more than one distinct genetic unit from within the SEA

lineage with additional contributions from the IND lineage.

Clearly, samples of known provenance, the use of informative loci that are diagnostic to

lineages, and boosting of sampling locations and sample sizes focused across the possible

intergrade zone IND and SEA would help resolve these ambiguities. Testing for levels of

admixture with higher sample sizes e.g. NewHybrids (Anderson and Thompson, 2002) could

resolve if the observed admixture is recent (F1s, F2s) or more advanced backcrosses, providing

evidence for disentangling artificial and natural hybridization events. Furthermore, the use of

archive and museum samples to provide genetic baselines prior to use of BP as an aquaculture

species can be used to create a genetic baseline to compare modern genetics against to identify

potential translocations and adaptation to changing environments e.g. (Turbek et al., 2023)

4.3 Conclusion

We corroborate the findings of mitochondrial barcoding identifying three divergent lineages of

BP through the analysis of genomic data. The analysis of genome-wide nuclear variants also

provides evidence of hierarchical divergences, that is within the IND, SEA and AUS+NG

lineages there is genetic structuring. The SEA lineage demonstrates a complex history with a

combination of natural genetic differentiation and what is likely human mediated admixture

events. Phenotypic and life history variation within BP remain poorly characterized and are other

aspects of variation that can substantially contribute to ecosystem productivity and commercial

applications. Research indicates that within Australia for instance, there is strong local adaptation
related to temperature (Newton et al., 2013) that could be applied to various goals. The wide and
varied use of BP in commercial, recreation and subsistence fisheries as well as aquaculture
creates differing priorities across user groups. Further exploration of BP variation can support
conservation aquaculture and food production as well as the preservation of native BP stocks.
510

Data Availability
512 Genomic sequence data are available from the NCBI Sequence Read Archive under BioProject
513 accessions PRJNA311498, PRJNA1021005 and PRJDB13763. Alignments for phylogenetic
514 analyses include the control block for SVDQuartets are included in the Data Supplement of this
515 paper. Sequence data examined in this study for time-calibrated phylogenetic analysis are
516 publicly available with mitochondrial sequence accessions reported in Table 1.
517

Declaration of Funding
519 Support for MC was received from the School of Life and Environmental Sciences at the
520 University of Sydney and the Charles Gilbert Heydon Travelling Fellowship.
521

Conflicts of Interest
523 The authors declare no conflicts of interest.
524

Lay Summary
526 Barramundi Perch, aka Barramundi, are a widespread fish widely propagated and highly sought
527 after for food and recreation. We find that the Barramundi contains three main genetic lineages
528 with numerous subdivisions, and that these divisions are linked to adaptive changes and may be
529 considered evolutionarily significant. These findings indicate diversity within Barramundi that
530 may be useful for both conservation and exploitation.
531

Figures

Figure 1 Distribution of *Lates calcarifer sensu lato* from AquaMaps (Kaschner et al., 2015) with probability of occurrence (Overall Probability) colored from yellow (low) to red (high). The type locality of *L. calcarifer* is indicated by a purple square. Sampling of main genetic lineages included in this paper are indicated by shape, AUS+NG circles, SEA triangles and IND diamonds. Locations of WGS are approximate as specific coordinates are not available. Cambodia is represented by four separate collection locations (shown as one).

Figure 2 Maximum-Likelihood phylogeny of mitochondrial Cytochrome Oxidase I from 60
barramundi perch individuals derived from whole-genome sequence data. The tree is rooted by
L. japonicus. Node shape indicates the putative lineage of origin and collection information is
provided as tip labels. Nodes with < 75% bootstrap support are collapsed to polytomies with
nodes ≥ 75% bootstrap support but < 90% indicated as grey squares and nodes ≥ 90% bootstrap
support as black rectangles.

547 **Figure 3** Species-tree from SVDQuartets of Barramundi Perch sampling locations in this study

548 based on 8,529 variants rooted by *Lates japonicus*. Values at nodes are bootstrap support values .

Figure 4 (A) Maximum-Likelihood phylogeny of genome-wide SNPs of 60 barramundi perch individuals. Nodes with < 75%

bootstrap support are collapsed to polytomies with nodes ≥ 75% bootstrap support but < 90% indicated as grey squares and nodes ≥

90% bootstrap support as black rectangles. **(B)** Implicit phylogenetic network of genome-wide SNPs from 60 barramundi perch

individuals. Node shape indicates the putative lineage of origin and color indicates collection location.

555
556

557   **Figure 5** Principal Component (PC) analysis of the whole-genome sequence data set of 60

558   barramundi perch individuals utilizing 7,321,852 genotype likelihoods. The first PC is on the x –

559   axis in both figures with the second PC on y – axis in (A) and third PC in (B).

Figure 6 Population genetic statistics of barramundi perch in the WGS data set. **(A)** Pairwise F_{st} values between regions. **(B)** Nucleotide diversity of sampling regions in the WGS data set. **(C)** Tajima's D of sampling regions in the WGS data set. Indonesia is represented by two sampling regions. Indonesia-SU, Vietnam and Bangladesh were excluded due to low sample sizes ($n \leq 2$).

Figure 7 Genome-wide admixture analysis of the WGS data set (n=60) from 7,321,852 genotype
likelihoods for $K = 2$ - 5 genetic clusters. Individuals are plotted on the x – axis and ancestry
coefficient (Q) on the y – axis.

Figure 8 Time-calibrated phylogeny of main mitochondrial lineages of barramundi perch. Node

age is from the posterior medians with 95% Highest Posterior Density (HPD) indicated by

horizontal bars.

Supplemental Figure S1 PCAdapt analyses for identification of adaptive genetic loci. **(A)**

Proportion of explained variance for each Principal Component (PC), supporting $K = 3$ PCs. **(B)**

Plot of PC 1 & PC 2, IND, SEA and AUS+NG. **(C)** Plot of PC 1 & PC 3, PC 3 separates

Indonesia-K and Indonesia-SU sampling locations from all other samples.

Supplemental Figure S2 PCAdapt analyses for identification of adaptive genetic loci from the

IND genetic lineage. **(A)** Proportion of explained variance for each Principal Component (PC).

(B) Plot of PC 1 & PC 2.

590
591
Supplemental Figure S3 PCAdapt analyses for identification of adaptive genetic loci within the SEA lineage. **(A)** Proportion of

explained variance for each Principal Component (PC), supporting at least $K = 3$ PCs. **(B)** Plot of PC 1 & PC 2, **(C)** Plot of PC 1 & PC

3.

Supplemental Figure S4 PCAdapt analyses for identification of adaptive genetic loci within the

AUS+NG lineage. **(A)** Proportion of explained variance for each Principal Component (PC). **(B)**

Plot of PC 1 & PC 2, separating Queensland, Northern Territory and Papua New Guinea

sampling locations.

Tables

Table 1 Sample origins by region of whole-genome sequence data set for *L. calcarifer sensu lato*. The sample size and average coverage is presented for each sampling region.

Region	Additional Geographic Information	Source	Sample Size	Average Coverage
Northern Territory, Australia	Darwin Harbor	Wild-Caught	6	5.80
Queensland, Australia	Hinchinbrook Channel	Wild-Caught	6	5.34
Papua New Guinea	Fly River	Wild-Caught	5	5.83
Indonesia	K - Kalimantan	Unknown	5	5.73
Indonesia	SJ – South of Jakarta	Wild-Caught	5	17.8
Indonesia	SU - Sulawesi	Unknown	1	6.13
Philippines		Hatchery Broodstock	5	6.32
Vietnam		Hatchery Broodstock	2	6.21
Cambodia	Three locations	Wild-Caught	5	5.90
Thailand	East Coast	Wild-Caught	7	15.60
Bangladesh	Bakkhali River	Unknown	2	10.04
India Eastern Coast	Tamil Nadu	Wild-Caught	7	8.50
India Western Coast	Mumbai	Wild-Caught	4	20.22

607   **Table 2** Taxa and sequences included in time-calibrated phylogeny of *Lates calcarifer sensu*

608   *lato*. *Lates calcarifer* is represented by three lineages, Australia + New Guinea (AUS+NG),

609   Southeast Asia (SEA) and Indian subcontinent (IND) as described in the text. The family is

610   given, with the taxon, corresponding BOLD or NCBI accession and the geographic origin of the

611   sample if known.

Family	Taxon	Accession	Origin
Latidae	*Lates calcarifer* AUS+NG	EU189379.1	Western Australia
	Lates calcarifer SEA	KJ573921.1	Indonesia
	Lates calcarifer IND	KJ573895.1	Andaman and Nicobar Islands
	Lates japonicus	LC269832.1	Japan
	Lates microlepis	ISZA082-21	Democratic Republic of the Congo
	Lates niloticus	LT853908.1	Not Specified
	Hypopterus macropterus	LC269834.1	Australia
	Psammoperca waigiensis	OQ387793.1	Philippines, Luzon
	Psammoperca datnioides	NC035737.1	Australia, Darwin
Centropomidae	*Centropomus ensiferus*	MW183492.1	Brazil, Para, Braganca
	Centropomus undecimalis	JQ841102.1	Belize, Belize City
Channidae	*Channa maculata*	KP112195.1	China, Hubei, Wuhan
Nandidae	*Nanus nandus*	KX245097.1	India

Bibliography

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25, 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Anderson, E.C., Thompson, E.A., 2002. A model-based method for identifying species hybrids using multilocus genetic data. Genetics 160, 1217–1229. https://doi.org/10.1093/genetics/160.3.1217

Brown, J.H., Lomolino, M.V., 1998. Biogeography. Sinauer Press, Sunderland, MA.

Campbell, M.A., Badger, M.E., Buckmaster, N., Starostka, A.B., Hawks, T., Finger, A.J., 2023. Molecular phylogenetic and population genetic relationships of a putative species of sucker (*Catostomus* sp.) from Surprise Valley in the Great Basin, USA. Transactions of the American Fisheries Society 152, 273–286. https://doi.org/10.1002/tafs.10407

Campbell, M.A., Chen, W.-J., López, J.A., 2013. Are flatfishes (Pleuronectiformes) monophyletic? Mol. Phylogenet. Evol. 69, 664–673. https://doi.org/10.1016/j.ympev.2013.07.011

Campbell, M.A., López, J.A., Satoh, T.P., Chen, W.-J., Miya, M., 2014. Mitochondrial genomic investigation of flatfish monophyly. Gene 551, 176–182. https://doi.org/10.1016/j.gene.2014.08.053

Chen, W.-J., Ruiz-Carus, R., Ortí, G., 2007. Relationships among four genera of mojarras (Teleostei: Perciformes: Gerreidae) from the western Atlantic and their tentative placement among percomorph fishes. J. Fish. Biol. 70, 202–218. https://doi.org/10.1111/j.1095-8649.2007.01395.x

Chenoweth, S.F., Hughes, J.M., Keenan, C.P., Lavery, S., 1998. Concordance between dispersal and mitochondrial gene flow: isolation by distance in a tropical teleost, Lates calcarifer (Australian barramundi). Heredity 80, 187–197. https://doi.org/10.1046/j.1365-2540.1998.00292.x

Chifman, J., Kubatko, L., 2015. Identifiability of the unrooted species tree topology under the coalescent model with time-reversible substitution processes, site-specific rate variation, and invariable sites. Journal of Theoretical Biology 374, 35–47. https://doi.org/10.1016/j.jtbi.2015.03.006

Chifman, J., Kubatko, L., 2014. Quartet inference from SNP data under the coalescent model. Bioinformatics 30, 3317–3324. https://doi.org/10.1093/bioinformatics/btu530

Cluney, B.J., 2004. The Northern Territory wild-catch barramundi *Lates calcarifer* fishery: a comparative investigation of the social, cultural, economic and environmental values of key stakeholders. (Master's). CDU.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools. GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008

Delić, T., Trontelj, P., Rendoš, M., Fišer, C., 2017. The importance of naming cryptic species and the conservation of endemic subterranean amphipods. Scientific Reports 7, 3391. https://doi.org/10.1038/s41598-017-02938-z

Des Roches, S., Post, D.M., Turley, N.E., Bailey, J.K., Hendry, A.P., Kinnison, M.T., Schweitzer, J.A., Palkovacs, E.P., 2018. The ecological importance of intraspecific variation. Nature Ecology & Evolution 2, 57–64. https://doi.org/10.1038/s41559-017-0402-5

Edwards-Vandenhoek, S., 2015. This my country, I'm painting here: the role of social practice
art and participatory design in building resilience and place-making in the Warmun
Aboriginal Community, Kimberley, Western Australia., in: Australia Council of
University Art and Design Schools Conference Proceedings. Adelaide.

Fricke, R., Eschmeyer, W.N., van der Laan, R. (Eds.), 2023. Eschmeyer's Catalog of Fishes:
Genera, Species, References.

Godfray, H.C.J., Knapp, S., Mace, G.M., 2004. The role of taxonomy in species conservation.
Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences
359, 711–719. https://doi.org/10.1098/rstb.2003.1454

Hackl, T., Hedrich, R., Schultz, J., Förster, F., 2014. proovread : large-scale high-accuracy
PacBio correction through iterative short read consensus. Bioinformatics 30, 3004–3011.
https://doi.org/10.1093/bioinformatics/btu392

Hahn, C., Bachmann, L., Chevreux, B., 2013. Reconstructing mitochondrial genomes directly
from genomic next-generation sequencing reads—a baiting and iterative mapping
approach. Nucleic Acids Research. https://doi.org/10.1093/nar/gkt371

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S., 2017. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution 35,
518–522. https://doi.org/10.1093/molbev/msx281

Huson, D.H., Bryant, D., 2005. Application of phylogenetic networks in evolutionary studies.
Molecular Biology and Evolution 23, 254–267. https://doi.org/10.1093/molbev/msj030

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods
14, 587–589. https://doi.org/10.1038/nmeth.4285

Kaschner, K., Kesner-Reyes, K., Garilao, C., Rius-Barile, J., Rees, T., Froes, R., 2015.
AquaMaps: Predicted range maps for aquatic species. [WWW Document]. AquaMaps.
URL www.aquamaps.org (accessed 8.8.16).

Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30, 3059–3066.
https://doi.org/10.1093/nar/gkf436

Katoh, K., Toh, H., 2008. Recent developments in the MAFFT multiple sequence alignment
program. Brief. Bioinform. 9, 286–298. https://doi.org/10.1093/bib/bbn013

Keenan, C., 1994. Recent evolution of population structure in Australian barramundi, Lates
calcarifer (Bloch): An example of isolation by distance in one dimension. Mar.
Freshwater Res. 45, 1123–1148.

Korneliussen, T.S., Albrechtsen, A., Nielsen, R., 2014. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinform 15, 356. https://doi.org/10.1186/s12859-014-0356-4

Layton, K.K.S., Snelgrove, P.V.R., Dempson, J.B., Kess, T., Lehnert, S.J., Bentzen, P., Duffy,
S.J., Messmer, A.M., Stanley, R.R.E., DiBacco, C., Salisbury, S.J., Ruzzante, D.E.,
Nugent, C.M., Ferguson, M.M., Leong, J.S., Koop, B.F., Bradbury, I.R., 2021. Genomic
evidence of past and future climate-linked loss in a migratory Arctic fish. Nature Climate
Change 11, 158–165. https://doi.org/10.1038/s41558-020-00959-7

Lehnert, S.J., Bradbury, I.R., Wringe, B.F., Van Wyngaarden, M., Bentzen, P., 2023.
Multifaceted framework for defining conservation units: An example from Atlantic
salmon (Salmo salar) in Canada. Evolutionary Applications n/a.
https://doi.org/10.1111/eva.13587

Li, C., Ricardo, B.R., Smith, L.W., Ortí, G., 2011. Monophyly and interrelationships of Snook
706   and Barramundi (Centropomidae sensu Greenwood) and five new markers for fish
707   phylogenetics. Mol. Phylogenet. Evol. 60, 463–471.
708   https://doi.org/10.1016/j.ympev.2011.05.004
709   Li, H., Durbin, R., 2010. Fast and accurate long-read alignment with Burrows-Wheeler
710   transform. Bioinformatics 26, 589–595. https://doi.org/10.1093/bioinformatics/btp698
711   Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler
712   transform. Bioinformatics 25, 1754–1760. https://doi.org/10.1093/bioinformatics/btp324
713   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
714   Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The sequence
715   alignment/map format and SAMtools. Bioinformatics 25, 2078–2079.
716   https://doi.org/10.1093/bioinformatics/btp352
717   Loughnan, S.R., Smith-Keune, C., Beheregaray, L.B., Robinson, N.A., Jerry, D.R., 2019.
718   Population genetic structure of barramundi (*Lates calcarifer*) across the natural
719   distribution range in Australia informs fishery management and aquaculture practices.
720   Mar. Freshwater Res. 70, 1533–1542.
721   Luu, K., Bazin, E., Blum, M.G.B., 2017. pcadapt: an R package to perform genome scans for
722   selection based on principal component analysis. Molecular Ecology Resources 17, 67–
723   77. https://doi.org/10.1111/1755-0998.12592
724   Maddison, W.P., 1997. Gene trees in species trees. Systematic Biology 46, 523–536.
725   https://doi.org/10.1093/sysbio/46.3.523
726   Moore, R., 1979. Natural sex inversion in the Giant Perch (*Lates calcarifer*). Mar. Freshwater
727   Res. 30, 803–813.
728   Moore, R., Reynold, L., 1982. Migration patterns of barramundi, *Lates calcarifer* (Bloch), in
729   Papua New Guinea. Mar. Freshwater Res. 33, 671–682.
730   Moyle, P.B., Campbell, M.A., 2022. Cryptic species of freshwater sculpin (Cottidae: *Cottus*) in
731   California. Zootaxa 5154, 501–527. https://doi.org/10.11646/zootaxa.5154.5.1
732   Newton, J.R., Zenger, K.R., Jerry, D.R., 2013. Next-generation transcriptome profiling reveals
733   insights into genetic factors contributing to growth differences and temperature
734   adaptation in Australian populations of barramundi (Lates calcarifer). Marine Genomics
735   11, 45–52. https://doi.org/10.1016/j.margen.2013.07.002
736   Nguyen, H.-V., Legras, J.-L., Neuvéglise, C., Gaillardin, C., 2011. Deciphering the hybridisation
737   history leading to the lager lineage based on the mosaic genomes of *Saccharomyces*
738   *bayanus* strains NBRC1948 and CBS380T. PLoS ONE 6, e25821.
739   https://doi.org/10.1371/journal.pone.0025821
740   Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2014. IQ-TREE: A Fast and
741   Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
742   Molecular Biology and Evolution 32, 268–274. https://doi.org/10.1093/molbev/msu300
743   Otero, O., 2004. Anatomy, systematics and phylogeny of both Recent and fossil latid fishes
744   (Teleostei, Perciformes, Latidae). Zool. J. Linn. Soc-Lond. 141, 81–133.
745   https://doi.org/10.1111/j.1096-3642.2004.00111.x
746   Pethiyagoda, R., Gill, A.C., 2012. Description of two new species of sea bass (Teleostei: Latidae:
747   *Lates*) from Myanmar and Sri Lanka. Zootaxa 3314.
748   https://doi.org/10.11646/zootaxa.3314.1.1

Privé, F., Luu, K., Vilhjálmsson, B.J., Blum, M.G.B., 2020. Performing Highly Efficient
Genome Scans for Local Adaptation with R Package pcadapt Version 4. Molecular
Biology and Evolution 37, 2153–2154. https://doi.org/10.1093/molbev/msaa053
Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J.,
Sklar, P., de Bakker, P.I.W., Daly, M.J., Sham, P.C., 2007. PLINK: A Tool Set for
Whole-Genome Association and Population-Based Linkage Analyses. The American
Journal of Human Genetics 81, 559–575. https://doi.org/10.1086/519795
Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A., 2018. Posterior
summarization in Bayesian phylogenetics using Tracer 1.7. Systematic Biology 67, 901–
904. https://doi.org/10.1093/sysbio/syy032
Rimmer, M.A., 2009. *Lates calcarifer*, in: Crespi, V., New, M. (Eds.), Cultured Aquatic Species
Fact Sheets. FAO.
Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: Efficient Bayesian
phylogenetic inference and model choice across a large model space. Syst. Biol. 61, 539–
542. https://doi.org/10.1093/sysbio/sys029
Santini, F., Harmon, L., Carnevale, G., Alfaro, M., 2009. Did genome duplication drive the
origin of teleosts? A comparative study of diversification in ray-finned fishes. BMC Evol.
Bio. 9, 194. https://doi.org/10.1186/1471-2148-9-194
Schindler, D.E., Hilborn, R., Chasco, B., Boatright, C.P., Quinn, T.P., Rogers, L.A., Webster,
M.S., 2010. Population diversity and the portfolio effect in an exploited species. Nature
465, 609–612. https://doi.org/10.1038/nature09060
Shaklee, J., Salini, J., 1985. Genetic variation and population subdivision in Australian
barramundi, Lates calcarifer (Bloch). Mar. Freshwater Res. 36, 203–218.
Shamsi, S.M., Williams, M., Mansourian, Y., 2020. An introduction to Aboriginal fishing
cultures and legacies in seafood sustainability. Sustainability 12.
https://doi.org/10.3390/su12229724.
Skotte, L., Korneliussen, T.S., Albrechtsen, A., 2013. Estimating individual admixture
proportions from Next Generation Sequencing data. Genetics 195, 693.
https://doi.org/10.1534/genetics.113.154138
Swofford, D.L., 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).
Turbek, S.P., Bossu, C., Rayne, C., Gruppi, C., Kus, B.E., Whitfield, M., Smith, T.B., Paxton,
E.H., Bay, R.A., Ruegg, K.C., 2023. Historical DNA reveals climate adaptation in an
endangered songbird. Nature Climate Change 13, 735–741.
https://doi.org/10.1038/s41558-023-01696-3
Vij, S., Kuhl, H., Kuznetsova, I.S., Komissarov, A., Yurchenko, A.A., Van Heusden, P., Singh,
S., Thevasagayam, N.M., Prakki, S.R.S., Purushothaman, K., Saju, J.M., Jiang, J.,
Mbandi, S.K., Jonas, M., Hin Yan Tong, A., Mwangi, S., Lau, D., Ngoh, S.Y., Liew,
W.C., Shen, X., Hon, L.S., Drake, J.P., Boitano, M., Hall, R., Chin, C.-S., Lachumanan,
R., Korlach, J., Trifonov, V., Kabilov, M., Tupikin, A., Green, D., Moxon, S., Garvin, T.,
Sedlazeck, F.J., Vurture, G.W., Gopalapillai, G., Kumar Katneni, V., Noble, T.H., Scaria,
V., Sivasubbu, S., Jerry, D.R., O'Brien, S.J., Schatz, M.C., Dalmay, T., Turner, S.W.,
Lok, S., Christoffels, A., Orbán, L., 2016. Chromosomal-level assembly of the Asian
Seabass genome using long sequence reads and multi-layered scaffolding. PLOS
Genetics 12, e1005954. https://doi.org/10.1371/journal.pgen.1005954

Vij, S., Purushothaman, K., Gopikrishna, G., Lau, D., Saju, J.M., Shamsudheen, K.V., Kumar, K.V., Basheer, V.S., Gopalakrishnan, A., Hossain, M.S., Sivasubbu, S., Scaria, V., Jena, J.K., Ponniah, A.G., Orbán, L., 2014. Barcoding of Asian seabass across its geographic range provides evidence for its bifurcation into two distinct species. Frontiers in Marine Science 1.

Wallace, A.R., 1859. The geographical distribution of birds. Ibis 1, 449–454.

Wang, D., Wong, M.-K., Lee, M.-Y., Campbell, M.A., Chen, W.-J., In Revision. Systematics of the superfamily Trichiuroidea (Teleostei: Scombriformes): undescribed species and a new family of commercially important demersal fishes in tropical and temperate seas. Zoological Journal of the Linnean Society.

Wang, L., Wan, Z.Y., Lim, H.S., Yue, G.H., 2016. Genetic variability, local selection and demographic history: genomic evidence of evolving towards allopatric speciation in Asian seabass. Molecular Ecology 25, 3605–3621. https://doi.org/10.1111/mec.13714

Waples, R.S., 1991. Definition of "species" under the Endangered Species Act: application to Pacific salmon. NOAA technical memorandum NMFS F/NWC ; 194.

Ward, R.D., Holmes, B.H., Yearsley, G.K., 2008. DNA barcoding reveals a likely second species of Asian sea bass (barramundi) (*Lates calcarifer*). J. Fish Biol 72, 458–463. https://doi.org/10.1111/j.1095-8649.2007.01703.x

Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y., 2017. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology and Evolution 8, 28–36. https://doi.org/10.1111/2041-210X.12628

Yue, G.H., Zhu, Z.Y., Lo, L.C., Wang, C.M., Lin, G., Feng, F., Pang, H.Y., Li, J., Gong, P., Liu, H.M., Tan, J., Chou, R., Lim, H., Orban, L., 2009. Genetic variation and population structure of Asian seabass (Lates calcarifer) in the Asia-Pacific region. Aquaculture 293, 22–28. https://doi.org/10.1016/j.aquaculture.2009.03.053