1   **PTEN Regulates Myofibroblast Activation in Valvular Interstitials Cells based on**

2   **Subcellular Localization**

3   Dilara Batan[1,2,*], Georgios Tseropoulos[1,2,*], Carrie Bishop[1], Bruce E. Kirkpatrick[1,2,3], Kaustav
4   Bera[1,2], Alex Khang[1,2], Mary Weiser-Evans[4,5,6], Kristi S. Anseth[1,2]
5
[1] Department of Chemical and Biological Engineering, University of Colorado, Boulder,
Colorado 80303, United States
2 The BioFrontiers Institute, University of Colorado, Boulder, Colorado 80303, United States
[3] Medical Scientist Training Program, School of Medicine, University of Colorado, Aurora,
Colorado 80045, United States
[4] Department of Medicine, Division of Renal Diseases and Hypertension, University of Colorado,
Anschutz Medical Campus, 12700 East 19th Avenue, C281, Research Complex 2, Room 7101,
Aurora, Colorado 80045, USA.
[5] Center for Fibrosis Research and Translation, School of Medicine, University of Colorado,
Anschutz Medical Campus, Aurora, Colorado 80045, USA.
[6] Department of Medicine, Cardiovascular Pulmonary Research Program, University of Colorado,
Anschutz Medical Campus, Aurora, Colorado 80045, USA.
18
19   *Denotes equal contribution
20
21
22

23   **Abstract**
24

Aortic valve stenosis (AVS) is characterized by altered mechanics of the valve leaflets, which

disrupts blood flow through the aorta and can cause left ventricle hypotrophy. These changes in

the valve tissue result in activation of resident valvular interstitial cells (VICs) into

myofibroblasts, which have increased levels of αSMA in their stress fibers. The persistence of

VIC myofibroblast activation is a hallmark of AVS. In recent years, the tumor suppressor gene

phosphatase and tensin homolog (PTEN) has emerged as an important player in the regulation of

fibrosis in various tissues (e.g., lung, skin), which motivated us to investigate PTEN as a

potential protective factor against matrix-induced myofibroblast activation in VICs. In aortic

valve samples from humans, we found high levels of PTEN in healthy tissue and low levels of

PTEN in diseased tissue. Then, using pharmacological inducers to treat VIC cultures, we

observed PTEN overexpression prevented stiffness-induced myofibroblast activation, whereas

genetic and pharmacological inhibition of PTEN further activated myofibroblasts. We also

observed increased nuclear PTEN localization in VICs cultured on stiff matrices, and nuclear

PTEN also correlated with smaller nuclei, altered expression of histones and a quiescent

fibroblast phenotype. Together, these results suggest that PTEN not only suppresses VIC

activation, but functions to promote quiescence, and could serve as a potential pharmacological

target for the treatment of AVS.

<u>Introduction</u>

Aortic Valve Stenosis (AVS) is a progressive cardiovascular disease that directly affects

the aortic valve leaflet matrix composition and mechanical properties, leading to disrupted blood

flow through the aorta[1] and eventual complications associated with left ventricular pressure-

overload.[2] When aortic valve leaflets thicken and stiffen as AVS progresses, significant

deficiencies in cardiac output result, which can lead to heart failure and death. AVS affects

around 2% of the population older than 65 with higher occurrences in males compared to

females.[3] Symptomatic AVS patients suffer high mortality rates if the condition is left untreated.[4]

Currently, there are no known effective pharmacological treatments for AVS, and surgical or

transcatheter valve replacement remains the standard of care for patients with severe disease.

However, many patients are ineligible for surgery, as valve replacement has many risks

including valve leaks, bleeding, and vessel damage.[1,5] As a result, a deeper molecular and

mechanistic understanding of AVS is necessary if one hopes to identify alternative

pharmaceutical treatment strategies.

Valvular interstitial cells (VICs) are the main population of cells residing in aortic valve

leaflets, and these fibroblast cells are responsible for valve maintenance and repair following

injury. Because of their key role in valve homeostasis, VICs are highly sensitive to

physicochemical changes in the matrix microenvironment.[6–8] Specifically, VICs transition from

quiescent fibroblasts to activated myofibroblasts, characterized by formation of alpha-smooth

muscle actin (αSMA) stress fibers, increased proliferation, and elevated extracellular matrix

(ECM) deposition.[9] During fibrosis, the myofibroblast phenotype persists over time, leading to a

pathogenic accumulation of ECM , stiffening of the local microenvironment, and creating a

positive feedback loop that can accelerate disease progression.[10] Investigating

mechanotransduction pathways capable of reversing this process and restoring the quiescent

fibroblast phenotype continues to be a promising avenue for potential treatments.[11–15]

Hydrogels provide highly tunable microenvironments that can be adjusted to recapitulate

the stiffness of the valve matrix at different stages of fibrosis progression and enable researchers

to study VIC mechanosensing in a controlled manner.[16] In this work, the crosslinking density of

poly(ethylene glycol) (PEG) hydrogels was tailored to control the matrix mechanical

properties,[17] while the fibronectin-derived adhesive peptide, RDGS, was used to promote VIC-

matrix interactions.[18] When cultured on soft PEG hydrogels (~1-5 kPa), VICs maintained a

quiescent fibroblast phenotype, but transitioned to an activated myofibroblast phenotype on

stiffer PEG matrices (E'~13-30kPa).[15,19–22] Thus, these biomaterial systems provided a reliable *in*

vitro platform for VIC cultures in healthy and disease scenarios.

As valvular tissue stiffens during AVS progression, signaling through mechanosensitive

pathways can reveal essential signaling hubs governing cell phenotype and potentially serve as

druggable targets. Importantly, elevated levels of phospho-AKT (p-AKT), which indicates

increased activation of phosphoinositide 3-kinase (PI3K)/AKT pathway, have been observed in

the fibroblast to myofibroblast transition (FMT).[23] In this regard, the pathway has been presented

as a master regulator of idiopathic pulmonary fibrosis.[23] Similarly, we have previously

demonstrated that PI3K/AKT signaling is increased on stiff substrates and is necessary for

substrate stiffness induced myofibroblast activation in VICs.[11,24] These findings suggest that

targeting the PI3K/AKT pathway may be a promising approach for treating fibrotic diseases.

Phosphatase and tensin homolog (PTEN), which is a negative regulator of the PI3K/AKT

pathway, has been proposed as a potential target for therapeutic intervention in fibrotic

disease.[25–27] For example, Lu *et al.* demonstrated that overexpression of PTEN in the mouse

model superPTEN protects against angiotensin II induced vascular fibrosis.[26] Further, the same

group later observed that small molecule drugs, such a 5-azacytidine and itraconazole, increase

PTEN levels and are able to ameliorate effects of increased signaling through the PI3K/AKT

pathway.[26,28] Complementary to the overexpression studies, Parapuram *et al*. demonstrated that

loss of PTEN leads to overproduction of collagen and is enough to induce fibrosis in lung and

skin tissues.[27,29] Similar studies were conducted in mammary gland tissue by Jones *et al*. and

further confirmed that PTEN regulates extracellular matrix deposition and collagen organization,

steps that are essential in fibrosis development.[30] Collectively, these findings suggest that

targeting the PI3K/AKT through PTEN may hold promise for developing novel therapies for

fibrotic diseases. However, no previous work has been conducted specifically investigating the

effects of PTEN in the context of AVS. Motivated by this, we aimed to explore the role of PTEN

in VIC myofibroblast activation and its potential as a therapeutic target for aortic valve disease.

Results herein reveal a major role of PTEN in regulating VIC fibroblast and myofibroblast

phenotypes. Initially, we observe a loss of PTEN protein in diseased human valve leaflet

sections compared to healthy ones. To test the specific role of mechanotransduction on PTEN

expression and VIC phenotype, we tailored the modulus/stiffness of PEG hydrogels to direct

VICs towards either a fibroblast (E' ~3 kPa) or a myofibroblast (E' ~13.4 kPa) majority

population. Next, we demonstrate that pharmacologically increased PTEN levels promote a more

quiescent phenotype for VICs cultured on stiff hydrogels, while chemical and genetic inhibition

of PTEN increased myofibroblast activation on soft hydrogels. Interestingly, the minority

fibroblast population on stiff matrices had higher nuclear localization of PTEN compared to the

majority myofibroblast population. In addition, the population of cells with high PTEN nuclear

localization correlated with smaller nuclei and VIC quiescence. The regulation of VIC activation

by PTEN was further verified by shRNA knockdown, as well as overexpression (nuclear

localization and exclusion signal tagged) experiments *in vitro*. Our results led us to hypothesize

that nuclear PTEN may play an additional role, beyond cytoplasmic regulation of PI3K/AKT,

maintaining VIC quiescence by regulating histone expression on stiff matrices. Overall, in this

study we show that PTEN is a major regulator of VIC fibroblast to myofibroblast transition and

can serve as a future target for pharmaceutical interventions.

<u>Materials and Methods</u>

119
<u>Hydrogel precursor synthesis</u>

Poly(ethylene glycol) photodegradable diacrylate PEGdiPDA (MW approximately 3400 Da) was

synthesized and characterized according to a previous method.[31] >90% acrylate functionalization

was verified by [1]H NMR. The adhesive peptide, acrylamide diethylene glycol-diethylene glycol-

glycine-arginine-glycine-aspartic acid-serine-glycine (Ac-OOGRGDSG), was synthesized via 9-

fluorenylmethyloxycarbonyl (Fmoc)/1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]

pyridinium 3-oxid hexafluorophosphate (HATU) coupling strategy on Rink Amide

methylbenzhydryl amine (MBHA) resin (0.500 mmol) using a Protein Technologies Tribute

automated solid-phase peptide synthesis machine and characterized using electrospray ionization

(ESI) mass spectroscopy.[31]

Formation of hydrogels

The preparation of photo-softening PEGdiPDA hydrogels was adapted from previously

described protocols.[31] PEGdiPDA was co-polymerized with poly(ethylene glycol) monoacrylate

(PEGA; $M_n \sim 400$ Da; Monomer-Polymer and Dajac Laboratories) and acryl-OOGRGDSG in

PBS via redox-initiated free radical polymerization. Gel solutions were prepared with 7.0 wt%

PEGdiPDA, 6.8 wt% PEGA, 5 mM acryl-OOGRGDSG, 0.2 M ammonium persulfate (APS) and

0.1 M tetramethylethylenediamine (TEMED). Gels were formed on acrylated cover glass after

the addition of APS/TEMED, with diameters of 12, 15, or 25 mm and a thickness of 100 µm.

Gels were allowed to cure for 6 min and washed and stored in PBS before cell seeding. Soft

hydrogels were prepared by irradiating the initial hydrogels with UV light ($\lambda = 365$ nm; $I_0 = 10$

mW/cm^2) for 6 minutes.

Swollen rheological characterization

Optically thin PEGdiPDA hydrogels (thickness = 200 µm) were made between two sigmacoted

glass slides separated with a spacer using the formulations above and allowed to swell in PBS

overnight. A rheometer (TA Instruments DH-R3) with a parallel plate accessory was used to

measure the shear modulus of the PEGdiPDA hydrogel before and after photo-degradation. The

equilibrium storage modulus (G′) was measured using a dynamic time sweep ($\gamma = 1\%$; $\omega = 1$

rad/s), as determined to be in the linear regime for this material. For *in situ* degradation, a quartz

plate was used, and the rheometer was equipped with a 365 nm light source ($\lambda = 365$ nm; $I_0 = 10$

mW/cm^2; Omnicure 1000, LumenDynamics). After network formation, the hydrogel was

exposed to 365 nm light for 6 minutes, and the change in G' was monitored using the same

dynamic time sweep parameters. The same dynamic time sweep measurements were used to

obtain final G' as a function of light exposure time. G' was measured and converted to Young's

modulus (E) using a Poisson's ratio of 0.5.[32]

Valvular Interstitial Cell (VIC) isolation and culture

Primary VICs were isolated from aortic valve leaflets from fresh male porcine hearts (Hormel) as

previously described.[14,20,21,24]. Briefly, aortic valve leaflets were excised from the hearts and

rinsed in Earle's Balanced Salt Solution (EBSS, Life Technologies) supplemented with 50 U/mL

penicillin and 50 µg/mL streptomycin (Thermo Fisher), and 1 µg/mL amphotericin B (Thermo

Fisher). Leaflets were incubated in a 250 U/mL type II collagenase (Worthington) solution in

EBSS for 30 min under 5% CO2 at 37˚C with gentle shaking. Valvular endothelial cells were

removed by vortexing leaflets at maximum speed for 30 seconds. Leaflets were transferred to

fresh collagenase solution and incubated for 60 min under 5% CO2 at 37˚C with gentle shaking.

Leaflets were vortexed again for 2 minutes to dislodge valvular interstitial cells (VICs), and the

entire solution was passed through a 100 µm cell strainer. The strained cell solution was

centrifuged at 0.2g x 20 minutes. Pelleted cells were resuspended in VIC expansion media

compromising of M199 media (Life Technologies) 15% fetal bovine serum (FBS, Life

Technologies), 50 U/mL penicillin, 50 µg/mL streptomycin (Thermo Fisher), and 1µg/mL

amphotericin B (Thermo Fisher). Cells were cultured in T-75 flasks for expansion until 80%

confluency was obtained. Cells were harvested using trypsin (Life Technologies) to be frozen in

50% fetal bovine serum (FBS) (Life Technologies), 45% M199 media (Life Technologies), and

5% DMSO (Sigma-Aldrich).

For experimental use, cells were thawed and cultured in T-75 flasks in 15% FBS

supplemented M199 media until 80% confluent. Cells were counted after harvesting using an

automated hemocytometer (Countess). A seeding density of 15,000 cells/cm^2 was used for VICs

on hydrogels, with growth media consisting of M199 media supplemented with 1% FBS,

penicillin-streptomycin, and fungizone. 12 mm hydrogels were used for immunostaining

experiments; 25 mm hydrogels were used for real-time quantitative polymerase chain reaction

(RT-qPCR) to increase the measurable RNA content.

Hydrogel Immunostaining

VICs were fixed in 4% paraformaldehyde (PFA, Electron Microscopy Sciences) in PBS for 20

minutes and rinsed twice with PBS. Cells were permeabilized with 0.1% TritonX-100 (Fisher

Scientific) in PBS for 20 minutes followed by 1 hour of blocking in 5% bovine serum albumin

(BSA, Sigma-Aldrich). Primary antibodies were diluted in 5% BSA and incubated at 4˚C

overnight at given concentrations: α-smooth muscle actin (αSMA, mouse, Cat. No. ab7817,

Abcam) at 1:300 dilution, PTEN (rabbit, Cat. No. D8H1X, Cell Signaling) at 1:300 dilution.

Primary antibody was washed off the cells twice with PBST (0.5% Tween-20 in PBS), and once

in PBS. Samples were incubated in secondary antibody solution of 1:200 dilution goat anti-

mouse Alexa 488 (Life Technologies), 1:500 dilution of goat anti-rabbit Alexa 648 (Life

Technologies), 1:5000 dilution of HCS Cell Mask (Life Technologies), and 1:1000 dilution of

DAPI (Life Technologies) for 1 hour before rinsing twice with PBST (0.005% Tween-20 in

PBS) and once in PBS. For imaging, hydrogels were inverted onto a glass-bottom 24-well plate

(Cellvis) and visualized using an Operetta high-content confocal microscope using a 20x air

objective (Perkin Elmer). Immunofluorescence intensity in immunostained cells was quantified

using Harmony software (Perkin Elmer). PTEN nuclear localization was quantified by dividing

PTEN average nuclear signal by PTEN average cytoplasmic signal.

Isolation of mRNA and Real-Time Quantitative Polymerase Chain Reaction (RT-qPCR)

RNA was isolated according to manufacturer's instructions using an RNeasy Micro Kit

(Qiagen). RNA concentration was quantified using ND-1000 NanoDrop Spectrophotometer

(Thermo Fischer). cDNA was produced using iScript Reverse Transcription Supermix kit (Bio-

Rad) on a Mastercycler (Eppendorf). Relative mRNA levels were measured with SYBR Green

reagents on an iCycler (Bio-Rad). Results were normalized to the housekeeping gene RPL30

(Ribosomal protein L30). Custom primers are presented in **Table S1**.

207
PTEN lentiviral knockdown

Two shRNA sequences were purchased from Millipore Sigma and cloned into the pLKO.1

lentiviral vector. Virus was produced utilizing second generation packaging plasmids PSPAX2

and pnd2.g as described previously.[33] Briefly, HEK 293T cells were transfected with the

packaging plasmids and shRNA for PTEN and two viral batches were collected after 24 and 48

hours. P1 VICs were transduced with the viral constructs for 3 hours, and VICs that incorporated

the virus were selected with puromycin (1ug/ml) for 3 days. The genetically modified VICs were

allowed to proliferate on TCPS until 80% confluence and subsequently passaged on soft or stiff

hydrogels (E' ~3 kPa, E' ~13.4 kPa, respectively). The potency of shRNA PTEN knockdown

was validated though immunohistochemistry for PTEN staining and qPCR testing for PTEN

mRNA, and the most potent knockdown sequence was selected for experiments reported herein.

Transfection with NES and NLS plasmid constructs

VICs were seeded on stiff hydrogels at a density of 6000 cells / cm^2 and incubated overnight.

Transfection media was prepared the next day using lipofectamine 3000 at a dilution of 0.5ul in

50ul of Opti-MEM reduced serum media per well and vortexed. The plasmids (Addgene PTEN

NES Plasmid# 10932, Addgene PTEN NLS Plasmid# 10933) were amplified using the Qiagen^{TN}

mini prep kit and diluted in a concentration of 1ug/ul. The control condition consisted of

lipofectamine and Opti-MEM only. The growth media was removed from wells and transfection

media was added. The transfection plate was incubated for 1 hour and then washed three times

with Opti-MEM. Subsequently, growth media was added and the plate was incubated for 3 days,

after which the plate was fixed and prepared for immunohistochemical characterization.

Heart valve tissue histology

Samples of aortic healthy and diseased (AVS) valve leaflets were acquired from deceased or

heart transplant patients at the University of Colorado Anschutz Medical Campus **(Table S2)**.

Tissues were fixed in 4% paraformaldehyde for 10 minutes and embedded in Tissue Tek O.C.T.

for subsequent cryosectioning. The sections were washed 3 times with PBS and permeabilized

with 0.1% TritonX for 1 hour and blocked with 5% goat serum, 5% BSA, and 0.01% TritonX for

1 hour. The sections were treated with the following primary antibodies (αSMA, mouse, Cat.

No. ab7817, Abcam at 1:300 dilution, PTEN rabbit, Cat. No. D8H1X, Cell Signaling at 1:300

dilution) sequentially over 2 days with 2 overnight incubations with the primary antibody in

blocking buffer (dilution 1:200) and 1 hour incubation with secondary antibody (1:400). The

tissues were imaged on an Nikon A1R confocal microscope and analyzed using ImageJ.

Statistical Analysis
244

Data are presented as mean ± standard deviation with a minimum of three biological and four

technical replicates for all studies unless otherwise stated. Outliers were removed using Robust

Outlier (ROUT) method (set at 1%). Significance was defined as *P<0.05, **P<0.01,

***P<0.005, ****P<0.001 using a t-test for comparison of two samples and one-way analysis of

variance (ANOVA) for comparison of three or more samples. Statistics were performed in

GraphPad Prism.

Results

PTEN levels are lower in diseased human valve leaflets compared to healthy valves

As previous studies have demonstrated that loss of PTEN is involved in fibrosis development,

we wanted to test whether PTEN levels were changed with AVS development. To do this, we

immunostained human valve leaflet tissue sections from healthy and diseased patients for DAPI,

αSMA, and PTEN and performed histological analysis targeted towards the levels of αSMA and

PTEN. In healthy tissue, we observed high levels of PTEN and low levels of αSMA (**Figure 1**

A-C). In diseased tissue, significantly higher levels of αSMA and minimal levels of PTEN were

observed, indicating an inverse correlation, and implicating a protective role for PTEN against

fibrosis development in valve leaflets (**Figure 1 A**). These results motivated us to investigate the

role of PTEN in VIC activation *in vitro,* building on previous work that utilizes soft and stiff

hydrogels to recapitulate VIC myofibroblast differentiation and differential expression of αSMA,

dependent on substrate stiffness.

265
266
Figure 1: PTEN and αSMA levels inversely correlate in healthy and AVS human aortic
valve tissue. A. Representative histological images of AVS and healthy human aortic valve
tissue (male) stained for DAPI (blue), PTEN (magenta) and αSMA (green) levels in valvular
interstitial cells. Scale bar = 50 µm **B.** Quantification of average αSMA intensity (normalized to
DAPI per mm^2) for healthy and diseased tissue within the marked area in A. (normalized to
DAPI per mm^2, n=5 fields of view per tissue, t-test, ***p<0.001). C. Quantification of PTEN
intensity (normalized per mm^2) for healthy and diseased tissue within the marked area. (n=5
fields of view per tissue, t-test, ***p<0.001)

281 *PTEN levels inversely correlate with αSMA levels in VIC hydrogel cultures*
282

Motivated by the results from human valve samples, we wanted to test the effect of substrate

stiffness on PTEN expression levels. To do this, we initially cultured VICs on soft and stiff

hydrogels to identify conditions which promote a majority fibroblast phenotype on soft

hydrogels and a majority myofibroblast phenotype on stiff hydrogels. We synthesized PEG-

acrylate hydrogels supplemented with a fibronectin-derived peptide sequence, arginine-glycine-

aspartic acid-serine (RGDS) to allow for cell adhesion and facilitation of cell-matrix interactions.

We formulated soft (Young's modulus (E')□=□~2.8□kPa) and stiff hydrogels (Young's

modulus (E')□=□~13.4□kPa) to match the modulus of healthy and diseased valve leaflet tissue

to mimic the local microenvironmental VIC mechanosensing in AVS **(Supplementary Figure**

1). Consistent with previous data, we observed a quiescent fibroblast VIC phenotype on soft

hydrogels and an activated myofibroblast phenotype on stiff gels, as identified by the presence

and intensity of αSMA **(Figure 2 A,B)**. In addition to the αSMA protein produced, the VIC

morphology cultured on soft hydrogels was significantly smaller in cell and nuclear area

compared to VICs cultured on stiff hydrogels (

Supplementary Figure B-C). Next, to determine if PTEN plays a protective role against VIC

myofibroblast activation, we examined protein levels in VICs cultured on the same soft and stiff

hydrogels. We observed that VICs displayed significantly higher PTEN levels when cultured on

soft hydrogels compared to stiff hydrogels, inversely correlating with αSMA levels and

myofibroblast activation **(Figure C,D)**.

302
303
304
Figure 2: **PTEN levels inversely correlate with myofibroblast activation in VICs A.**
Representative images of VICs cultured on soft and stiff hydrogels for 3 days, αSMA (green)
and cytoplasm (red). Scale bar = 100 µm **B.** Quantification of αSMA (green) intensity for A.
(n>750 cells from 4 hydrogels, unpaired t test, ****p<0.001). **C.** Representative images of
PTEN (gray) levels in VICs cultured on soft and stiff hydrogels for 3 days. **D.** Quantification of
PTEN intensity for C. (n>440 cells from 3 hydrogels, unpaired t test, ****p<0.001). **E.**
Quantification of αSMA (green) intensity for F. **F.** Representative images of VICs cultured on
soft and stiff hydrogels for 3 days treated with DMSO vehicle or 5 µM AKT inhibitor MK-2206,
αSMA (green) and cytoplasm(red). Scale bar = 100 µm (n>750 cells from 4 hydrogels,
unpaired t test, ****p<0.001). **G.** Representative images of VICs cultured on soft and stiff
hydrogels for 3 days, AKT (red) and cytoplasm (yellow). Scale bar = 100 µm **H.** Quantification
of AKT (red) intensity for G. (n>920 cells from 3 hydrogels, unpaired t test). **I.** Representative
images of VICs cultured on soft and stiff hydrogels for 3 days, pAKT (green) and cytoplasm
(yellow). Scale bar = 100 µm **J.** Quantification of pAKT (green) intensity for I. (n>1000 cells
from 3 hydrogels, unpaired t test, ****p<0.001).

Since PTEN is a known inhibitor of the PI3K/AKT pathway, we sought to verify its potential

role in influencing stiffness-induced PI3K/AKT activity in VICs myofibroblast cultured on soft

and stiff hydrogels. Specifically, we used a highly specific and widely used small molecule AKT

inhibitor, MK-2206, and introduced it to activated VICs cultured on stiff hydrogels.[34–36] We

found that treatment with 5µM MK-2206 significantly reduced VIC αSMA expression when

cultured on stiff hydrogels compared to untreated controls, and promoted a quiescent fibroblast

population (**Figure 2 E,F, <u>Supplementary Figure</u>**). Further, we observed similar levels of AKT

in VICs cultured on soft and stiff hydrogels; however, pAKT levels were significantly higher in

VICs cultured on stiff hydrogels (**<u>Figure</u> G-J**). This observation supports the notion that

PI3K/AKT is a key mechanoregulator of VIC myofibroblast activation in stiff

microenvironments.[24]

PTEN activators decrease VIC activation in stiff hydrogel microenvironments

Next, we wanted to investigate if increasing PTEN levels in VICs cultured on stiff

hydrogels would play a protective role against myofibroblast activation. To test this, we used two

small molecule PTEN activators that have been shown to increase cellular PTEN levels.[28,37]

Initially, we treated VICs cultured on TCPS with varying doses of the PTEN activators, 5-

azacytidine or itraconazole, and compared these to the DMSO vehicle control. We tested PTEN

mRNA levels using a concentration range (100 nm, 1 µM, and 10 µM) of 5-azacytidine and

itraconazole and observed that PTEN levels were significantly increased with 10 µM treatment

of both 5-azacytidine (**Figure A**) and itraconazole (**Supplementary Figure**).

cytoplasm αSMA

344
<u>Figure 3:</u> PTEN activators decrease myofibroblast activation in VICs A. Relative mRNA
expression levels of PTEN in VICs cultured on TCPS with treatment of vehicle or increasing
concentration of the PTEN activator, 5-azacytidine, n=4 wells, means ± SD shown, one-way
ANOVA test, *p < .05 ,**p < .01. **B.** Representative images of VICs cultured on soft and stiff
hydrogels for 3 days in the presence of vehicle or PTEN activator 5-azacytidine, αSMA (green)
and cytoplasm (red). Scale bar = 100 µm. **C.** Quantification of αSMA intensity for B (n>600
cells from 4 hydrogels, one-way ANOVA test, ****p<0.001). **D-G.** Relative mRNA expression
levels of fibrotic genes in VICs cultured on soft or stiff hydrogels treated with vehicle or PTEN
activator 5-azacytidine for 3 days, n=4 hydrogels, one-way ANOVA test, *p < .05 ,**p < .01,
p < .001,*p<0.001.
355

After verifying that PTEN levels could be elevated using small molecule activators, we

treated VICs cultured on stiff hydrogels with either vehicle or 5-azacytidine and itraconazole

with previously determined concentrations **(Supplementary Figure A-D)**. We observed that

VICs treated with PTEN activators displayed significantly lower levels of αSMA compared to

the vehicle treated stiff hydrogel culture controls at levels resembling the soft hydrogel negative

controls **(Figure B,C, Supplementary Figure A,B)**. In addition to having lower levels of

αSMA, VICs treated with PTEN activators also displayed a significantly smaller cell size and

nuclear area when compared to DMSO treated controls **(Supplementary Figure A,B)**.

To more broadly assess the role of PTEN in myofibroblast activation, VICs were cultured

in the presence of PTEN activators or DMSO vehicle, and qPCR was performed to determine the

relative expression of myofibroblast associated genes, namely alpha-Smooth Muscle Actin

(ACTA2), Connective Tissue Growth Factor (CTGF), Serpin Peptidase Inhibitor clade E

member 1 (SERPINE1), and Collagen type 1A1 (COL1A1). The expression of all four genes

was significantly decreased in VICs cultured on stiff hydrogels treated with PTEN activators

compared to the vehicle controls at levels resembling the soft hydrogel negative control

conditions **(Figure   D-G, Supplementary Figure   C,D)**. These results suggest that PTEN

influences multiple markers of VIC myofibroblasts and that the effects are observable after 72

hours of treatment.

Chemical and genetic PTEN inhibition increases VIC myofibroblast activation in soft hydrogel

microenvironments

To further validate the role of PTEN in myofibroblast activation, we next performed a

series of studies inhibiting PTEN activity by introducing chemical and genetic perturbations.

Initially we tested the effects of a potent PTEN phosphatase inhibitor bpV(HOpic) on VIC

myofibroblast activation, as measured by αSMA expression.[38] Quiescent VICs cultured on soft

hydrogels treated with 100 nM of bpV(HOpic) displayed significantly higher levels of αSMA

protein compared to cells treated with the DMSO control **(Figure 1 A,B)**. In addition to αSMA

protein content, ACTA2 gene expression levels were also elevated with PTEN inhibitor

treatment **(Figure 1 C)**. These results suggest that inhibition of PTEN phosphatase activity alone

can promote myofibroblast activation in VICs. However, interestingly, we observed non-

significant changes in mRNA levels of COL1A1 and SERPINE 1, suggesting more than just

phosphatase activity of PTEN is important to induce these transcriptional changes (

Supplementary Figure).

In addition to chemical PTEN inhibition, we utilized lentiviral shRNA knockdowns of

PTEN in transduced early passage VICs. PTEN protein levels in the knockdown VICs were

significantly reduced compared to the scrambled shRNA controls on both soft and stiff

hydrogels. **(Figure 1 D,F, Supplementary Figure A,C)**. We cultured scrambled or PTEN

knockdown VICs on both soft and stiff hydrogels. On soft hydrogels, consistent with our

previous data, we observed enhanced PTEN expression and low levels of VIC myofibroblast

activation (**Figure 1 D**), but PTEN knockdown on soft hydrogels resulted in increased levels of

αSMA **(Figure 1 E,F)**. On stiff hydrogels, we observed lowered levels of PTEN with higher

levels of αSMA (**Supplementary Figure A-C**). However, the effect of PTEN on VICs cultured

on stiff hydrogels was much lower compared to soft hydrogel conditions, as VICs are already

activated on stiff hydrogels, but the effects observed were statistically significant

(Supplementary Figure B,C). After confirming that PTEN protein levels were lower with our

shRNA constructs, we next quantified mRNA expression levels of known fibrotic genes **(Figure**

1 G). Compared to scrambled shRNA controls, PTEN knockdown VICs displayed higher levels

of ACTA2, and COL1A1 expression. Collectively, these results further validate PTEN's role as a

regulator of the fibroblast-to-myofibroblast transition in VICs, using hydrogels as a mechano-

regulator to introduce quiescence or activating matrix signals.

406
407
<u>Figure 1:</u> PTEN inhibitors increase myofibroblast activation in VICs. A. Representative
images of VICs cultured on soft and stiff hydrogels for 3 days in the presence of DMSO vehicle
or 100nM of PTEN inhibitor bpV(Hopic), αSMA (green) and cytoplasm(red). Scale bar = 100
µm. **B.** Quantification of αSMA intensity for A. (n>600 cells from 4 hydrogels, one-way
ANOVA test, ****$p<0.001$). **C.** Relative mRNA expression levels of ACTA2 in VICs cultured
on soft hydrogels with treatment of DMSO vehicle or 100nM of PTEN inhibitor bpV(Hopic),
and stiff hydrogels, n=4 wells, means ± SD shown. **D.** Representative images of scrambled and
PTEN knockdown VICs cultured on stiff hydrogels for 3 days, αSMA (green) and
cytoplasm(red). Scale bar = 100 µm. **E.** Quantification of αSMA fluorescence intensity (n>30
cells, t-test ,****$p<0.001$) **F.** Quantification of PTEN fluorescence intensity (n>30 cells, t-test,
****$p<0.001$). **G.** Relative mRNA expression levels of various fibrotic genes in VICs treated
with the scrambled or PTEN on TCPS, n=1 vector.
420

421
422 *High nuclear/cytoplasmic ratio of PTEN correlates with low αSMA levels in VIC on stiff*
423 *hydrogels*
424
In addition to the canonical role of PTEN in dephosphorylating PIP3 at the cell membrane,

additional roles of PTEN and its nuclear localization have been recently identified, namely that

PTEN is able to act as a transcription factor and alter global chromatin architecture and gene

expression levels.[39] Further, we have previously shown that matrix stiffness can influence the

persistence of VIC myofibroblast activation through epigenetic remodeling, so we next sought to

test whether nuclear PTEN is involved in this process. Specifically, we used quantitative image

analysis to investigate the subcellular localization of PTEN in VICs cultured on both soft and

stiff hydrogels.

Initially, we quantified αSMA levels of VIC cultures on soft and stiff hydrogels and separated

the population into two classes: fibroblasts and myofibroblasts based on the intensity of αSMA

signal. We observed that the quiescent VIC fibroblasts had a significantly higher nuclear to

     cytoplasmic PTEN ratio compared to myofibroblasts cultured on stiff hydrogels, where PTEN

remained largely cytoplasmic (

A,B). We observed no statistically significant differences in the nuclear/cytoplasmic PTEN ratio

between fibroblasts and myofibroblasts in VICs cultured on soft hydrogels, where VICs are

quiescent. (

C-D). Together, these results suggest that PTEN may serve an additional role in the nucleus,

protecting against stiffness-induced matrix activation of myofibroblasts.

Figure 5: High nuclear/cytoplasmic ratio of PTEN correlates with low αSMA levels on stiff
hydrogels. A. Single cell images of VICs cultured on stiff hydrogels for 3 days αSMA (green)
and PTEN (red). Scale bar = 10 µm. B. Quantification of PTEN nuclear to cytoplasmic ratio in
fibroblast and myofibroblast populations on stiff hydrogels (n>250 cells from 4 hydrogels, one-
way ANOVA test, ****p<0.001). C. Single cell images of VICs cultured on soft hydrogels for 3
days αSMA (green) and PTEN (red). Scale bar = 10 µm. D. Quantification of PTEN nuclear to
cytoplasmic ratio in fibroblast and myofibroblast populations on soft hydrogels (n>250 cells
from 4 hydrogels, one-way ANOVA test, ****p<0.001). E. Quantification of nuclear area of
cells with high and low nuclear to cytoplasmic PTEN ratio on soft hydrogels (n>250 cells from 4
hydrogels).

To verify the role of PTEN localization in myofibroblast activation, we ectopically

expressed PTEN in VICs with two distinct nucleotide sequences that signal either nuclear

(nuclear localization signal-NLS) or cytoplasmic (nuclear exclusion signal-NES) localization.

Both plasmid constructs contain a hemagglutinin (HA) tag to verify transfection efficiency. VICs

were plated on stiff 2D hydrogels and transfected with either NES PTEN or NLS PTEN plasmids

with ~4-7% efficiency **(Figure 6 D)**. After 3 days, we observed increased PTEN

nuclear/cytoplasmic ratio in the NLS PTEN condition compared to NES PTEN and untransfected

control, indicating that the transfected plasmid constructs localized PTEN as expected **(Figure 6**

A, B). More importantly, αSMA was significantly downregulated among the cells transfected

with NLS PTEN, while αSMA levels were slightly upregulated in the NES PTEN compared to

untransfected control **(Figure 6 A, C)**, indicating that nuclear PTEN plays a protective role

against myofibroblast activation on stiff hydrogels.

As nuclear PTEN has been shown to play a role in nuclear dynamics and epigenetic

modifications, we wanted to test if there was a correlation between high nuclear PTEN and

nuclear area in VICs.[39] For VICs cultured on stiff hydrogels, we separated the cell population

into two cohorts based on their nuclear to cytoplasmic PTEN ratio. VICs with a high

nuclear/cytoplasmic PTEN ratio (nuclear/ cytoplasmic PTEN $\geq$ than 1.5) had significantly

smaller nuclei compared to VICs with low nuclear/cytoplasmic PTEN ratio (nuclear/

cytoplasmic PTEN < than 1.5) **(Figure 6 E)**. This observation lead us to a more in depth

investigation of the role of PTEN in chromatin dynamics.

Figure 6: Ectopic expression of NLS PTEN downregulates αSMA on stiff hydrogels. A.
Single cell images of VICs cultured on stiff hydrogels for 3 days αSMA (green), PTEN (red) and
anti-HA tag (white). Scale bar = 10 µm. **B.** Quantification of αSMA among untransfected VICs
and VICs with ectopic expression of NES and NLS PTEN on stiff hydrogels. (n=40 cells from 6
hydrogels, one-way ANOVA test, ****p<0.001) **C.** Quantification of the ratio of nuclear to
cytoplasmic PTEN among untransfected, NES and NLS VICs (n=40 cells from 6 hydrogels, one-
way ANOVA test, ****p<0.001). **D.** Quantification of the % transfection efficiency between
NES and NLS PTEN. (n=40 cells from 6 hydrogels, one-way ANOVA test, ****p<0.001) **E.**
Quantification of nuclear area of cells with high and low nuclear to cytoplasmic PTEN ratio on
stiff hydrogels (n>250 cells from 4 hydrogels).

489
490

Since the nuclear/cytoplasmic ratio of PTEN affected nuclear size in addition to αSMA

levels, we subsequently studied the effect of the localization of PTEN on VIC epigenetic

modifications. PTEN has been implicated in the regulation of histone activity in the context of

fibrosis and heart failure, thus, we were interested in the correlation between the protective role

of PTEN against αSMA upregulation and the levels of the tri-methylation of histone 3 on lysine-

9 (H3K9me3) on VICs cultured for 3 days on soft and stiff hydrogels transfected with NES

PTEN

<u>Figure 7</u>: Nuclear localization of PTEN upregulates H3K9me3 on stiff hydrogels and
protects against myofibroblast activation. A. Single cell immunocytochemical images of
untransfected VICs and VICs with ectopic expression of NES and NLS PTEN cultured on stiff
hydrogels for 3 days stained for αSMA (green), PTEN (red) and H3K9me3 (white). Scale bar =
10 µm. **B.** Quantification of αSMA intensity among untransfected VICs and VICs with ectopic
expression of NES and NLS PTEN on stiff hydrogels. (n=50 cells from 4 hydrogels, one-way
ANOVA test, ****p<0.001) **C.** Quantification of H3K9me3 nuclear intensity. (n=50 cells from 4
hydrogels, one-way ANOVA test, ****p<0.001) **D.** Quantification of the ratio of nuclear to
cytoplasmic PTEN among untransfected, NES and NLS VICs (n=50 cells from 4 hydrogels, one-
way ANOVA test, ****p<0.001).[40–42]

509
or NLS PTEN. Consistent with our previous results, the high nuclear/cytoplasmic ratio of PTEN,

induced by NLS PTEN transfection, downregulated αSMA on stiff hydrogels but also

upregulated H3K9me3, increasing its nuclear levels. In the NES PTEN as well as the

untransfected control conditions, H3K9me3 levels remained significantly lower **(Figure 7A-D)**.

Interestingly, on soft hydrogels, where VICs display low αSMA levels, the increase in

nuclear/cytoplasmic PTEN ratio, similarly, resulted in increased H3K9me3 levels

(Supplementary Figure 9A-D). This, indicated that the PTEN-H3K9me3 upregulation

modulates αSMA levels, but their interactions are independent of VIC myofibroblast activation.

519
Figure 8: PTEN levels correlate with H3K8me3 levels in healthy human heart valve tissue
while inversely correlating with αSMA on diseased tissue. A. Representative histological
images of AVS and healthy human heart valve tissue (male) stained for DAPI (blue), αSMA
(green), PTEN (red) and H3K9me3 (white) levels in valvular interstitial cells. Scale bar = 40 µm
B. Quantification of average αSMA intensity (normalized to DAPI per mm^2) for healthy and
diseased tissue. (normalized to DAPI per mm^2, n=5 fields of view per tissue, 2 biological
replicates, t-test, ***p<0.001). **C.** Quantification of average H3K9me3 intensity (normalized to
DAPI per mm^2) for healthy and diseased tissue. (normalized to DAPI per mm^2, n=5 fields of
view per tissue, 2 biological replicates, t-test, ***p<0.001). **D.** Colocalization analysis of
H3K9me3 and DAPI and αSMA. Plot of the classical Pearson's coefficient. (n=5 fields of view
per tissue, 2 biological replicates)

Finally, we were interested in the translation of our *in vitro* results of the PTEN-

H3K9me3 interactions on human healthy and fibrotic heart valve tissue. Interestingly,

histological analysis positively correlated the levels of PTEN with the levels of H3K9me3 in

healthy tissue and inversely correlated with αSMA levels in fibrotic tissue (**Figure 8 A-C**).

Colocalization analysis of H3K9me3 with DAPI, confirmed its nuclear localization albeit

H3K9me3 only partially localizing with PTEN in human tissues (**Figure 8 D**), suggesting an

important but complex role of PTEN in the regulation of αSMA in human heart valve tissue.

Discussion

Results presented herein suggest that PTEN regulates myofibroblast activation in VICs,

which is consistent with the emerging notion that PTEN plays a central protective role against

fibrosis development in numerous tissues and organ systems.[25–27,29,36] as well as myogenic

differentiation of smooth vascular smooth muscle cells[62]. In line with this, we observed

decreased levels of PTEN in diseased human valve leaflet tissues compared to healthy ones. The

results linking PTEN with aortic valve stenosis complement what has been shown in tissue

samples from patients with pulmonary fibrosis, where an inverse correlation of the expression of

PTEN and αSMA was observed.[43] Further, loss of PTEN protein and downregulation in its

phosphatase activity have both been linked to development of pulmonary fibrosis.[44–48] To further

investigate the role of PTEN, we exploited biomaterial matrices to study the cross-talk between

mechanosensing and matrix signaling, in the absence of other confounding factors, that can

influence the VIC fibroblast-to-myofibroblast transition. We focused on elucidating how

regulation of PTEN activity alters VIC activation/quiescence in both soft (healthy) and stiff

(fibrotic-like) hydrogel microenvironments. Our findings suggest that PTEN has multiple modes

of action depending on PTEN localization and matrix stiffness (**Figure 9**). We observe total

PTEN expression levels decrease with matrix stiffness, inverse to the effects of stiffness on VIC

myofibroblast activation. However, the spatial distribution of PTEN matters, as high nuclear to

cytoplasmic ratio of PTEN on stiff hydrogels correlates with the quiescent fibroblast phenotype,

as observed in

and may protect against myofibroblast activation when VICs are grown in stiff, fibrotic-

like microenvironments.

566
Figure 9: Schematic of PTEN activity in VICs on soft and stiff hydrogels. A. VICs cultured
on soft hydrogels are quiescent due to the high PTEN activity in both the cellular cytosol and
nucleus. Cytosolic PTEN acts as an antagonist of PI3K/AKT, while nuclear PTEN is involved in
the regulation of histone expression. **B.** VICs differentiate towards myofibroblasts on stiff
hydrogels due to the downregulation of PTEN, resulting in increased levels of phosphor-
PI3K/AKT and elevated αSMA expression levels. However, when PTEN is ectopically
expressed in the nucleus, αSMA levels are downregulated through increased expression of
H3K9me3, maintaining VIC quiescence.

Further, we show that PTEN directly influences myofibroblast activation in VICs. PTEN

is a known inhibitor of the PI3K/AKT pathway, and results from our lab reported by Wang *et al.*

have shown that the phosphorylation of PI3K/AKT pathway is central to matrix stiffness induced

myofibroblast activation in VICs.[24]   We validate the involvement of AKT in myofibroblast

activation by pharmacological inhibition of phospho-AKT through MK-2206, which resulted in

decreased αSMA levels. This led us to examine the relative role of PTEN, an AKT antagonist,

abrogating the effects of mechanosensing. Namely, PTEN activators decrease, while PTEN

inhibitors increase myofibroblast activation, respectively. Furthermore, in addition to being

regulated by matrix stiffness, PTEN levels are also epigenetically regulated. We have

demonstrated increased mRNA expression of PTEN by 5-azacytidine and itraconazole treatment

(**Figure** , **Supplementary Figure**). Epigenetic silencing of PTEN has been proposed by

methylation of the CpG islands within the promoter, which is involved in several cancers.[49–53] As

such, methyltransferase inhibitors 5-azacytidine and decitabine have been previously identified

as PTEN activators by blocking methylation of the promoter and increasing gene expression.[28,54]

Interestingly, global methylation levels are reported to increase in AVS, which suggests that

DNA methylation inhibitors might also be used to reverse this process.[55]

While relative cell PTEN levels are strong indicators of myofibroblast activation in this

study and others, subcellular localization of PTEN has also been shown to play important roles in

influencing cell phenotype.[43,56–58] We observe that high nuclear localization of PTEN correlates

with the quiescent fibroblast phenotype and smaller nuclei for VICs on stiff gels, but no

correlations were observed with VICs on soft hydrogels. This suggests high nuclear localization

of PTEN may protect against myofibroblast activation. Although loss of PTEN has been

associated with fibrosis, subcellular localization of PTEN to the nucleus has not been previously

studied in relation to fibrotic disease. To investigate the role of PTEN nuclear localization in

myofibroblast activation, we ectopically expressed NES and NLS PTEN in VICs grown on stiff

substrates, where nuclear PTEN significantly downregulated αSMA expression. A high level of

nuclear PTEN, regulating epigenetic modifications, might suggest resistance to the transient

myofibroblast state and continued protection against activation.

Complementing this observation, recent work from our lab linked histone regulation and

chromatin condensation with myofibroblast activation.[19,59] Specifically, Walker *et al.* used

ATAC-seq to show that transient myofibroblasts have a more open chromatin structure, which

subsequently condenses again to the stabilized phenotypic condition.[59,19] PTEN has been shown

to interact with chromatin by directly binding to histone H1 and controlling epigenetic

modifications[60] , while nuclear PTEN levels correlate with H3K9me3 intensity, a marker for

chromatin condensation, and depletion of H3K9me3 is observed in PTEN knockout cells.[56]

Further, our results are consistent with PTEN levels *in vivo* that correlate with H3K9me3 levels

and potentially a protective effect against αSMA upregulation and fibrosis. Mechanistically,

PTEN may reinforce the H3K9me3 suppression of the PPARγ gene, decreasing the expression of

fibrotic markers such as αSMA and COL1.[61] Taken together, we propose that nuclear PTEN

promotes the VIC quiescent fibroblast phenotype by indirectly controlling αSMA expression at

an epigenetic level. However, the mechanism that leads to the heterogeneity of nuclear PTEN

observed in the VIC population remains a topic for future investigation.

In conclusion, we have shown that PTEN and its subcellular localization play an

important role in regulating myofibroblast activation in VICs. Increased global levels of PTEN

expression, either by low matrix stiffness or by pharmacological treatment, protect against

myofibroblast activation. High levels of nuclear PTEN correlate with a H3K9me2 expression

levels and the quiescent fibroblast phenotype, even on stiff hydrogel matrices. Importantly, this

correlation is recapitulated in human healthy and fibrotic heart valve tissues, where we have

observed mutually exclusive expression of αSMA and PTEN. Thus, we demonstrated that PTEN

may be a valuable pharmacological target for non-invasive therapeutics for valve fibrosis and

lead to further understanding of the dynamics of AVS progression.

Acknowledgments: We thank Dr. Robert Weiss for valuable discussions and feedback. We

acknowledge National Institutes of Health R01 HL142935 and R01 HL132353 awarded to KSA.

We acknowledge funding from the National Institutes of Health T32 HL007822 awarded to

A.K., Helen Hay Whitney Foundation award number F1339 to K.B., American Heart

Association 20PRE35200068 awarded to D.B.

Author contributions: D.B., G.T., and K.S.A. designed and performed all the experiments. B.K.

and A.K. performed image analysis, mathematical analysis, and data 32 analysis. D.B., B.E.K.,

G.T., K.B., M.W.E. and K.S.A. contributed to writing and editing the paper.

Competing interests: The authors report no competing interests.

639
640
641
642 **Supplementary Material**
643

644
645
Supplementary Figure 1: **VICs display increased cellular and nuclear area on stiff**
hydrogels. A. Young's modulus for synthesized soft and stiff swollen hydrogels by shear
rheology. (n = 8, unpaired t test, ****p < .001). **B.** Quantification of cell size of VICs cultured
on soft or stiff hydrogels for 3 days (n>740 cells from 4 hydrogels, unpaired t test,
****p<0.001). **C.** Quantification of nuclear area of VICs cultured on soft or stiff hydrogels for 3
days (n>740 cells from 4 hydrogels, unpaired t test, ****p<0.001).
652

653
654

Supplementary Figure 2: **VICs display decreased myofibroblast activation with AKT**
inhibition A. Representative images of VICs cultured on soft and stiff hydrogels for 3 days,
αSMA (green) and cytoplasm (yellow). Scale bar = 100 μm **B.** Quantification of αSMA (green)
intensity for A. (n>1000 cells from 4 hydrogels, one-way ANOVA test, ****p<0.001).

661
662 **<u>Supplementary Figure 3</u>: Itraconazole treatment increases PTEN gene expression.** Relative
663 mRNA expression levels of PTEN in VICs cultured on TCPS with treatment of vehicle or
664 increasing concentration of PTEN activator itraconazole, n=4 wells, means ± SD shown, one-
665 way ANOVA test, **p < .01.

666
667
<u>Supplementary Figure 4</u>: PTEN activators decrease myofibroblast activation in VICs A.
Representative images of VICs cultured on TCPS for 3 days with indicated concentrations of
PTEN activators 5-azacytidine, αSMA (green) and nuclei(DAPI). Scale bar = 100 µm. **B.**
Quantification of myofibroblast activation for A, n=4 wells, one-way ANOVA test. **C.**
Representative images of VICs cultured on TCPS for 3 days with indicated concentrations of
PTEN activators itraconazole, αSMA (green) and nuclei(DAPI). Scale bar = 100 µm. **D.**
Quantification of myofibroblast activation for C, n=4 wells, one-way ANOVA test,
****p<0.001.

677
678

cytoplasm αSMA

Supplementary Figure 5: PTEN activators decrease myofibroblast activation in VICs A.
Representative images of VICs cultured on soft and stiff hydrogels for 3 days in the presence of
DMSO vehicle or10 µM itraconazole (PTEN activator), αSMA (green) and cytoplasm(red).
Scale bar = 100 µm. **B.** Quantification of αSMA intensity for A. (n>600 cells from 4 hydrogels,
one-way ANOVA test, *p < .05, ****p<0.001). **C-F.** Relative mRNA expression levels of
various fibrotic genes in VICs cultured on soft or stiff hydrogels treated with DMSO vehicle or
PTEN activator itraconazole (10µM), n=4 hydrogels, one-way ANOVA test, *p < .05 ,**p < .01,
p < .001,*p<0.001.

702
703

704
705 **Supplementary Figure 6: PTEN activators decrease cellular and nuclear area of VICs**
706 **cultured on stiff hydrogels. A.** Quantification of cell size of VICs cultured on soft or stiff
707 hydrogels for 3 days (n>600 cells from 4 hydrogels, unpaired t test, ****p<0.001). **B.**
708 Quantification of cell size of VICs cultured on soft or stiff hydrogels for 3 days (n>600 cells
709 from 4 hydrogels, unpaired t test, ****p<0.001).

710
711 **Supplementary Figure 7: PTEN phosphatase inhibition does not increase mRNA levels of**
712 **fibrotic genes A,B.** Relative mRNA expression levels of various fibrotic genes in VICs cultured
713 on soft or stiff hydrogels treated with DMSO vehicle or PTEN inhibitor bpV(Hopic) (100 nM),
714 n=3 hydrogels, one-way ANOVA test, *p < .05 ,**p < .01.
715

716
717 **Supplementary Figure 8:** **PTEN knockdown increases αSMA levels in VICs cultured on**
718 **stiff hydrogels A.** Representative images of scrambled and PTEN knockdown VICs cultured on
719 stiff hydrogels for 3 days, αSMA (green) and cytoplasm(red). Scale bar = 100 µm. **B.**
720 Quantification of αSMA fluorescence intensity (n>30 cells, t-test ,****p<0.001) **C.**
721 Quantification of PTEN fluorescence intensity (n>30 cells, t-test, ****p<0.001).
722

Supplementary Figure 9: Nuclear localization of PTEN upregulates H3K9me3 on soft hydrogels. A. Single cell immunocytochemical images of VICs cultured on soft hydrogels for 3 days αSMA (green), PTEN (red) and H3K9me3 (white). Scale bar = 10 µm. **B.** Quantification of αSMA intensity among untransfected VICs and VICs with ectopic expression of NES and NLS PTEN on soft hydrogels. (n=50 cells from 4 hydrogels, one-way ANOVA test, ****$p<0.001$) **C.** Quantification of H3K9me3 nuclear intensity. (n=50 cells from 4 hydrogels, one-way ANOVA test, ****$p<0.001$) **D.** Quantification of the ratio of nuclear to cytoplasmic PTEN among untransfected, NES and NLS VICs (n=50 cells from 4 hydrogels, one-way ANOVA test, ****$p<0.001$).

Table S1: **Custom Primers used for RT-qPCR**

GENE:	FW sequence 5′ to 3′	RV sequence 5′ to 3′
PTEN	GGCGGAACTTGCAATCCTCA	AACTTGTCTTCCCGTCGTGT
L30	GCTGGGGTACAAGCAGACTC	AGATTTCCTCAAGGCTGGGC
COL1A1	GGGCAAGACAGTGATTGAATACA	GGATGGAGGGAGTTTACAGGAA
CTGF	CTGGTCCAGACCACAGAGTGG	GCAGAAAGCGTTGTCATTGG
ACTA2	GCAAACAGGAATACGATGAAGCC	AACACATAGGTAACGAGTCAGAGC
SERPINE 1	ACGGCCATTATCGTCTCAGC	AAAAGGACCGTTCCTGTGGG

738
739
740   **Table S2:** **Patient Information of the tissue used**

Condition	Age	Sex	Race (Ethnicity)	Diagnosis	LVEF (%)
Healthy	45	M	american indian	non-ischemic	35
Stenotic	67	M	W	ischemic	29.5

References

1. Otto CM, Prendergast B. Aortic-Valve Stenosis — From Patients at Risk to Severe Valve Obstruction. *N Engl J Med*. 2014;371:744–756.

2. Yutzey KE, Demer LL, Body SC, Huggins GS, Towler DA, Giachelli CM, Hofmann-Bowman MA, Mortlock DP, Rogers MB, Sadeghi MM, Aikawa E. Calcific aortic valve disease: A consensus summary from the alliance of investigators on calcific aortic valve disease. *Arterioscler Thromb Vasc Biol*. 2014;34:2387–2393.

3. Schlotter F, Halu A, Goto S, Blaser MC, Body SC, Lee LH, Higashi H, DeLaughter DM, Hutcheson JD, Vyas P, Pham T, Rogers MA, Sharma A, Seidman CE, Loscalzo J, Seidman JG, Aikawa M, Singh SA, Aikawa E. Spatiotemporal Multi-Omics Mapping Generates a Molecular Atlas of the Aortic Valve and Reveals Networks Driving Disease. *Circulation* [Internet]. 2018;138:377–393. Available from: https://www.ahajournals.org/doi/10.1161/CIRCULATIONAHA.117.032291

4. Goody PR, Hosen MR, Christmann D, Niepmann ST, Zietzer A, Adam M, Bönner F, Zimmer S, Nickenig G, Jansen F. Aortic valve stenosis: From basic mechanisms to novel therapeutic targets. *Arterioscler Thromb Vasc Biol*. 2020;885–900.

5. Liang F, Wang X, Wang Q, Yan P, Yao L. Predilatation and paravalvular leakage risk in TAVR. *Lancet* [Internet]. 2020;396:600. Available from: http://dx.doi.org/10.1016/S0140-6736(20)30624-3

6. Liu AC, Joag VR, Gotlieb AI. The emerging role of valve interstitial cell phenotypes in regulating heart valve pathobiology. *Am J Pathol* [Internet]. 2007;171:1407–1418. Available from: http://dx.doi.org/10.2353/ajpath.2007.070251

7. Kendall RT, Feghali-Bostwick CA. Fibroblasts in fibrosis: Novel roles and mediators. *Front Pharmacol*. 2014;5 MAY:1–13.

8. Humphrey JD, Dufresne ER, Schwartz MA. Mechanotransduction and extracellular matrix homeostasis. *Nat Rev Mol Cell Biol* [Internet]. 2014;15:802–812. Available from: http://dx.doi.org/10.1038/nrm3896

9. Hinz B. Myofibroblasts. *Exp Eye Res* [Internet]. 2016;142:56–70. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26192991

10. Duscher D, Maan ZN, Wong VW, Rennert RC, Januszyk M, Rodrigues M, Hu M, Whitmore AJ, Whittam AJ, Longaker MT, Gurtner GC. Mechanotransduction and fibrosis. *J Biomech* [Internet]. 2014;47:1997–2005. Available from: http://www.nature.com/articles/ncb2756

11. Wang H, Leinwand LA, Anseth KS. Cardiac valve cells and their microenvironment-insights from in vitro studies. Nat. Rev. Cardiol. 2014;11:715–727.

12. Ceccato TL, Starbuck RB, Hall JK, Walker CJ, Brown TE, Killgore JP, Anseth KS, Leinwand LA. Defining the Cardiac Fibroblast Secretome in a Fibrotic Microenvironment. *J Am Heart Assoc*. 2020;9:e017025.

13. Kuehlmann B, Bonham CA, Zucal I, Prantl L, Gurtner GC. Mechanotransduction in Wound Healing and Fibrosis. *J Clin Med*. 2020;9:1423.

14. Ma H, Macdougall LJ, GonzalezRodriguez A, Schroeder ME, Batan D, Weiss RM, Anseth KS. Calcium Signaling Regulates Valvular Interstitial Cell Alignment and Myofibroblast Activation in Fast-Relaxing Boronate Hydrogels. *Macromol Biosci*. 2020;20:1–14.

15. Batan D, Peters DK, Schroeder ME, Aguado BA, Young MW, Weiss RM, Anseth KS.

Hydrogel cultures reveal Transient Receptor Potential Vanilloid 4 regulation of
myofibroblast activation and proliferation in valvular interstitial cells. *FASEB J* [Internet].
2022 [cited 2023 Jan 5];36:1–16. Available from:
https://onlinelibrary.wiley.com/doi/full/10.1096/fj.202101863R

16. Caliari SR, Burdick JA. A practical guide to hydrogels for cell culture. *Nat Methods 2016*
135 [Internet]. 2016 [cited 2023 Jan 10];13:405–414. Available from:
https://www.nature.com/articles/nmeth.3839

17. Tibbitt MW, Anseth KS. Hydrogels as extracellular matrix mimics for 3D cell culture.
Biotechnol Bioeng [Internet]. 2009 [cited 2023 Jan 10];103:655–663. Available from:
https://onlinelibrary-wiley-com.colorado.idm.oclc.org/doi/full/10.1002/bit.22361

18. Mann BK, Gobin AS, Tsai AT, Schmedlen RH, West JL. Smooth muscle cell growth in
photopolymerized hydrogels with cell adhesive and proteolytically degradable domains:
Synthetic ECM analogs for tissue engineering. *Biomaterials*. 2001;22:3045–3051.

19. Walker CJ, Batan D, Bishop CT, Ramirez D, Aguado BA, Schroeder ME, Crocini C,
Schwisow J, Moulton K, Macdougall L, Weiss RM, Allen MA, Dowell R, Leinwand LA,
Anseth KS, Leslie Leinwand CA. Extracellular matrix stiffness controls cardiac valve
myofibroblast activation through epigenetic remodeling. *Bioeng Transl Med*. 2022;7.

20. Grim JC, Aguado BA, Vogt BJ, Batan D, Andrichik CL, Schroeder ME, Gonzalez-
Rodriguez A, Max Yavitt F, Weiss RM, Anseth KS. Secreted factors from
proinflammatory macrophages promote an osteoblast-like phenotype in valvular
interstitial cells. *Arterioscler Thromb Vasc Biol*. 2020;E296–E308.

21. Aguado BA, Walker CJ, Grim JC, Schroeder ME, Batan D, Vogt BJ, Rodriguez AG,
Schwisow JA, Moulton KS, Weiss RM, Heistad DD, Leinwand LA, Anseth KS. Genes
That Escape X Chromosome Inactivation Modulate Sex Differences in Valve
Myofibroblasts. *Circulation*. 2022;145:513–530.

22. Kloxin AM, Benton JA, Anseth KS. In situ elasticity modulation with dynamic substrates
to direct cell phenotype. *Biomaterials* [Internet]. 2010 [cited 2023 Jan 10];31:1. Available
from: /pmc/articles/PMC2791128/

23. Wang J, Hu K, Cai X, Yang B, He Q, Wang J, Weng Q. Targeting PI3K/AKT signaling
for treatment of idiopathic pulmonary fibrosis. *Acta Pharm Sin B* [Internet]. 2022 [cited
2023 Feb 2];12:18–32. Available from:
www.elsevier.com/locate/apsbwww.sciencedirect.com

24. Wang H, Tibbitt MW, Langer SJ, Leinwand LA, Anseth KS. Hydrogels preserve native
phenotypes of valvular fibroblasts through an elasticity-regulated PI3K/AKT pathway.
Proc Natl Acad Sci U S A [Internet]. 2013;110:19336–19341. Available from:
http://www.pnas.org/content/110/48/19336.short%7B%5C%25%7D5Cnhttp://www.ncbi.
nlm.nih.gov/pubmed/24218588

25. Cai B, Yang L, Jung Y Do, Zhang Y, Liu X, Zhao P, Li J. PTEN: An Emerging Potential
Target for Therapeutic Intervention in Respiratory Diseases. 2022 [cited 2022 Dec
8];Available from: https://doi.org/10.1155/2022/4512503

26. Lu S, Strand KA, Mutryn MF, Tucker RM, Jolly AJ, Furgeson SB, Moulton KS,
Nemenoff RA, Weiser-Evans MCM. PTEN (phosphatase and tensin homolog) protects
against ang II (angiotensin II)-induced pathological vascular fibrosis and remodeling -
Brief report. *Arterioscler Thromb Vasc Biol*. 2020;40:394–403.

27. Parapuram SK, Thompson K, Tsang M, Hutchenreuther J, Bekking C, Liu S, Leask A.
Loss of PTEN expression by mouse fibroblasts results in lung fibrosis through a CCN2-

dependent mechanism. *Matrix Biol*. 2015;43:35–41.

28.   Strand KA, Lu S, Mutryn MF, Li L, Zhou Q, Enyart BT, Jolly AJ, Dubner AM, Moulton
KS, Nemenoff RA, Koch KA, Labarbera D V., Weiser-Evans MCM. High throughput
screen identifies the DNMT1 (DNA Methyltransferase-1) inhibitor, 5-azacytidine, as a
potent inducer of PTEN (Phosphatase and Tensin Homolog): Central Role for PTEN in 5-
azacytidine protection against pathological vascular remodeling. *Arterioscler Thromb*
Vasc Biol. 2020;40:1854–1869.

29.   Parapuram SK, Shi-Wen X, Elliott C, Welch ID, Jones H, Baron M, Denton CP, Abraham
DJ, Leask A. Loss of PTEN Expression by Dermal Fibroblasts Causes Skin Fibrosis. *J*
Invest Dermatol. 2011;131:1996–2003.

30.   Jones CE, Sharick JT, Colbert SE, Shukla VC, Zent JM, Ostrowski MC, Ghadiali SN,
Sizemore ST, Leight JL. Pten regulates collagen fibrillogenesis by fibroblasts through
SPARC. *PLoS One* [Internet]. 2021;16:1–21. Available from:
http://dx.doi.org/10.1371/journal.pone.0245653

31.   Kloxin AM, Kasko AM, Salinas CN, Anseth KS. Photodegradable hydrogels for dynamic
tuning of physical and chemical properties. *Science (80-)*. 2009;324:59–63.

32.   Yang C, Tibbitt MW, Basta L, Anseth KS. Mechanical memory and dosing influence stem
cell fate. *Nat Mater*. 2014;13:645–652.

33.   Mehrotra P, Ikhapoh I, Lei P, Tseropoulos G, Zhang Y, Wang J, Liu S, Bronner ME,
Andreadis ST. Wnt/BMP Mediated Metabolic Reprogramming Preserves Multipotency of
Neural Crest-Like Stem Cells. *Stem Cells* [Internet]. 2023 [cited 2023 May 5];41:287–
305. Available from: https://academic.oup.com/stmcls/article/41/3/287/6976057

34.   Malkomes P, Lunger I, Luetticke A, Oppermann E, Haetscher N, Serve H, Holzer K,
Bechstein WO, Rieger MA. Selective AKT Inhibition by MK-2206 Represses Colorectal
Cancer-Initiating Stem Cells. *Ann Surg Oncol* [Internet]. 2016 [cited 2023 May
14];23:2849–2857. Available from: https://pubmed.ncbi.nlm.nih.gov/27059026/

35.   Liu R, Liu D, Trink E, Bojdani E, Ning G, Xing M. The Akt-specific inhibitor MK2206
selectively inhibits thyroid cancer cells harboring mutations that can activate the PI3K/Akt
pathway. *J Clin Endocrinol Metab* [Internet]. 2011 [cited 2023 May 14];96. Available
from: https://pubmed.ncbi.nlm.nih.gov/21289267/

36.   Qiu T, Tian Y, Gao Y, Ma M, Li H, Liu X, Wu H, Zhang Y, Ding H, Cao M, Zhang J, Dai
J, Chen J, Cai H. PTEN loss regulates alveolar epithelial cell senescence in pulmonary
fibrosis depending on Akt activation. *Aging (Albany NY)* [Internet]. 2019 [cited 2023 Apr
19];11:7492. Available from: /pmc/articles/PMC6781970/

37.   Bollong MJ, Yang B, Vergani N, Beyer BA, Chin EN, Zambaldo C, Wang D, Chatterjee
AK, Lairson LL, Schultz PG. Small molecule-mediated inhibition of myofibroblast
transdifferentiation for the treatment of fibrosis. *Proc Natl Acad Sci U S A*.
2017;114:4679–4684.

38.   Schmid AC, Byrne RD, Vilar R, Woscholski R. Bisperoxovanadium compounds are
potent PTEN inhibitors. *FEBS Lett*. 2004;566:35–38.

39.   Yang J, Yin Y. PTEN in Chromatin Remodeling. *Cold Spring Harb Perspect Med*
[Internet]. 2020 [cited 2023 Apr 18];10. Available from: /pmc/articles/PMC6996446/

40.   Benitez JA, Ma J, D'Antonio M, Boyer A, Camargo MF, Zanca C, Kelly S, Khodadadi-
Jamayran A, Jameson NM, Andersen M, Miletic H, Saberi S, Frazer KA, Cavenee WK,
Furnari FB. PTEN regulates glioblastoma oncogenesis through chromatin-associated
complexes of DAXX and histone H3.3. *Nat Commun 2017 81* [Internet]. 2017 [cited 2024

May 16];8:1–15. Available from: https://www.nature.com/articles/ncomms15223

41. Shen WC, Lai YC, Li LH, Liao K, Lai HC, Kao SY, Wang J, Chuong CM, Hung SC.
Methylation and PTEN activation in dental pulp mesenchymal stem cells promotes
osteogenesis and reduces oncogenesis. *Nat Commun 2019 101* [Internet]. 2019 [cited 2024
May 16];10:1–13. Available from: https://www.nature.com/articles/s41467-019-10197-x

42. Yan F, Chen Z, Cui W. H3K9me2 regulation of BDNF expression via G9a partakes in the
progression of heart failure. *BMC Cardiovasc Disord* [Internet]. 2022 [cited 2024 May
16];22:1–13. Available from: https://link.springer.com/articles/10.1186/s12872-022-
02621-w

43. White ES, Atrasz RG, Hu B, Phan SH, Stambolic V, Mak TW, Hogaboam CM, Flaherty
KR, Martinez FJ, Kontos CD, Toews GB. Negative regulation of myofibroblast
differentiation by PTEN (phosphatase and tensin homolog deleted on chromosome 10).
Am J Respir Crit Care Med. 2006;173:112–121.

44. Geng J, Huang X, Li Y, Xu X, Li S, Jiang D, Liu Z, Dai H. Phosphatase and tensin
homolog deleted on chromosome 10 contributes to phenotype transformation of
fibroblasts in idiopathic pulmonary fibrosis via multiple pathways. *Exp Biol Med*
[Internet]. 2016 [cited 2023 Apr 27];241:157–165. Available from:
https://journals.sagepub.com/doi/10.1177/1535370215600100

45. Xie B, Zheng G, Li H, Yao X, Hong R, Li R, Yue W, Chen Y. Effects of the tumor
suppressor PTEN on the pathogenesis of idiopathic pulmonary fibrosis in Chinese
patients. *Mol Med Rep* [Internet]. 2016 [cited 2023 Apr 27];13:2715–2723. Available
from: http://www.spandidos-publications.com/10.3892/mmr.2016.4852/abstract

46. Nho RS, Hergert P. IPF Fibroblasts Are Desensitized to Type I Collagen Matrix-Induced
Cell Death by Suppressing Low Autophagy via Aberrant Akt/mTOR Kinases. *PLoS One*
[Internet]. 2014 [cited 2023 Apr 27];9:e94616. Available from:
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0094616

47. Im J, Hergert P, Nho RS. Reduced FoxO3a expression causes low autophagy in idiopathic
pulmonary fibrosis fibroblasts on collagen matrices. *Am J Physiol - Lung Cell Mol Physiol*
[Internet]. 2015 [cited 2023 Apr 27];309:L552–L561. Available from:
https://journals.physiology.org/doi/10.1152/ajplung.00079.2015

48. Mao P, Li J, Huang Y, Wu S, Pang X, He W, Liu X, Slutsky AS, Zhang H, Li Y.
MicroRNA-19b mediates lung epithelial-mesenchymal transition via phosphatidylinositol-
3,4,5-trisphosphate 3-phosphatase in response to mechanical stretch. *Am J Respir Cell Mol*
Biol [Internet]. 2017 [cited 2023 Apr 27];56:11–19. Available from: http://www.tm4.org/

49. Brito MB, Goulielmaki E, Papakonstanti EA. Focus on PTEN regulation. *Front Oncol*.
2015;5:166.

50. Kang YH, Lee HS, Kim WH. Promoter Methylation and Silencing of PTEN in Gastric
Carcinoma. *Lab Investig 2002 823* [Internet]. 2002 [cited 2023 Apr 24];82:285–291.
Available from: https://www.nature.com/articles/3780422

51. García JM, Silva J, Peña C, Garcia V, Rodriguez R, Cruz MA, Cantos B, Provencio M,
España P, Bonilla F. Promoter methylation of the PTEN gene is a common molecular
change in breast cancer. *Genes, Chromosom Cancer* [Internet]. 2004 [cited 2023 Apr
24];41:117–124. Available from: https://onlinelibrary-wiley-
com.colorado.idm.oclc.org/doi/full/10.1002/gcc.20062

52. Goel A, Arnold CN, Niedzwiecki D, Carethers JM, Dowell JM, Wasserman L, Compton
C, Mayer RJ, Bertagnolli MM, Boland CR. Frequent Inactivation of PTEN by Promoter

Hypermethylation in Microsatellite Instability-High Sporadic Colorectal Cancers. *Cancer Res* [Internet]. 2004 [cited 2023 Apr 24];64:3014–3021. Available from: https://aacrjournals.org/cancerres/article/64/9/3014/517889/Frequent-Inactivation-of-PTEN-by-Promoter

53. Mirmohammadsadegh A, Marini A, Nambiar S, Hassan M, Tannapfel A, Ruzicka T, Hengge UR. Epigenetic Silencing of the PTEN Gene in Melanoma. *Cancer Res* [Internet]. 2006 [cited 2023 Apr 24];66:6546–6552. Available from: https://aacrjournals.org/cancerres/article/66/13/6546/525928/Epigenetic-Silencing-of-the-PTEN-Gene-in-Melanoma

54. Xing XQ, Li B, Xu SL, Zhang CF, Liu J, Deng YS, Yang J. 5-Aza-2'-deoxycytidine, a DNA methylation inhibitor, attenuates hypoxic pulmonary hypertension via demethylation of the PTEN promoter. *Eur J Pharmacol* [Internet]. 2019 [cited 2023 Apr 24];855:227–234. Available from: https://pubmed.ncbi.nlm.nih.gov/31085236/

55. Nwachukwu N, Hagler M, Kafa R, Roos C, Miller J. Evidence for altered DNA methylation as a major regulator of gene expression in calcific aortic valve disease (671.15). *FASEB J*. 2014;28.

56. Gong L, Govan JM, Evans EB, Dai H, Wang E, Lee SW, Lin HK, Lazar AJ, Mills GB, Lin SY. Nuclear PTEN tumor-suppressor functions through maintaining heterochromatin structure. *Cell Cycle* [Internet]. 2015 [cited 2023 Feb 16];14:2323–2332. Available from: https://www.tandfonline.com/doi/abs/10.1080/15384101.2015.1044174

57. Horita H, Wysoczynski CL, Walker LA, Moulton KS, Li M, Ostriker A, Tucker R, McKinsey TA, Churchill MEA, Nemenoff RA, Weiser-Evans MCM. Nuclear PTEN functions as an essential regulator of SRF-dependent transcription to control smooth muscle differentiation. *Nat Commun* [Internet]. 2016 [cited 2023 Apr 19];7. Available from: /pmc/articles/PMC5411712/

58. Liu J-L, Sheng X, Hortobagyi ZK, Mao Z, Gallick GE, Yung WKA. Nuclear PTEN-Mediated Growth Suppression Is Independent of Akt Down-Regulation. *Mol Cell Biol* [Internet]. 2005 [cited 2023 May 11];25:6211. Available from: /pmc/articles/PMC1168816/

59. Walker CJ, Crocini C, Ramirez D, Killaars AR, Grim JC, Aguado BA, Clark K, Allen MA, Dowell RD, Leinwand LA, Anseth KS. Nuclear mechanosensing drives chromatin remodelling in persistently activated fibroblasts. *Nat Biomed Eng*. 2021;5:1485–1499.

60. Chen ZH, Zhu M, Yang J, Liang H, He J, He S, Wang P, Kang X, McNutt MA, Yin Y, Shen WH. PTEN Interacts with Histone H1 and Controls Chromatin Condensation. *Cell Rep* [Internet]. 2014 [cited 2023 Apr 25];8:2003. Available from: /pmc/articles/PMC4201947/

61. Jiang Y, Wang S, Zhao Y, Lin C, Zhong F, Jin L, He F, Wang H. Histone H3K9 demethylase JMJD1A modulates hepatic stellate cells activation and liver fibrosis by epigenetically regulating peroxisome proliferator-activated receptor γ. *FASEB J*. 2015;29:1830–1841.

62. Horita H, Wysoczynski CL, Walker LA, Moulton KS, Li M, Ostriker A, Tucker R, McKinsey TA, Churchill MEA, Nemenoff RA, Weiser-Evans MCM. Nuclear PTEN fucntions as an essential regularor of SRF-dependent transcription to control smooth muscle differentiation. *Nat Commun*. 2016;7.1;10830