Large-scale genome analyses provide insights into Hymenoptera evolution

Chun He[1], Yi Yang[1], Xianxin Zhao[1], Junjie Li[1], Yuting Cai[1], Lijia Peng[1], Yuanyuan Liu[1],

Shijiao Xiong[2], Yang Mei[1], Zhichao Yan[3], Jiale Wang[4], Shan Xiao[5], Ziwen Teng[6], Xueke

Gao[7], Hui Xue[7], Qi Fang[1], Gongyin Ye[1], Xinhai Ye[8,*]

[1]State Key Laboratory of Rice Biology & Ministry of Agricultural and Rural Affairs Key

Laboratory of Molecular Biology of Crop Pathogens and Insects, Institute of Insect

Sciences, Zhejiang University, Hangzhou, China

[2]Xianghu Laboratory, Hangzhou, China

[3]Department of Entomology, Nanjing Agricultural University, Nanjing, China

[4]Institute of Quality Standard and Monitoring Technology for Agro-Products of

Guangdong Academy of Agricultural Sciences, Guangzhou, China

[5]Ningbo Academy of Agricultural Sciences, Ningbo, China

[6]College of Plant Health & Medicine, Qingdao Agricultural University, Qingdao, China

[7]State Key Laboratory of Cotton Bio-breeding and Integrated Utilization, Institute of

Cotton Research, Chinese Academy of Agricultural Sciences, Anyang, China.

[8]College of Advanced Agriculture Sciences, Zhejiang A&F University, Hangzhou, China

*Correspondence: yexinhai@zafu.edu.cn (X.H. Ye)

Abstract

The order Hymenoptera includes a large number of species with diverse lifestyles and is

known for its significant contributions to natural ecosystems. To better understand the

evolution of this diverse order, we performed large-scale comparative genomics on 131

species from 13 superfamilies, covering most representative groups. We used these

genomes to reveal an overall pattern of genomic change in terms of gene content and

evolutionary rate throughout hymenopteran history. We identified genes that possibly

contributed to the evolution of several key innovations, such as parasitoidism, wasp-

waist, sting, and secondary phytophagy. We also discovered the distinct genomic

trajectories between the clade containing major parasitoid wasps (Parasitoida) and

stinging species (Aculeata) since their divergence, which are involved in many aspects of

genomic change, such as rapidly evolving gene families, gene gain and loss, and

metabolic pathway evolution. In addition, we explored the genomic features

accompanying the three independent evolution of secondary phytophagy. Our work

provides insights for understanding genome evolution and the genomic basis of

diversification in Hymenoptera.

Introduction

Hymenoptera is a diverse order of insects, which includes more than 153,000 described

species, such as sawflies, wasps, ants, and bees[1–3]. These hymenopterans play a pivotal

role in natural ecosystems, serving a variety of characters (e.g., herbivores, parasitoids,

predators, and pollinators)[4,5]. The remarkable diversity of species and the apparent

changes in lifestyles have prompted interest in the evolution of Hymenoptera. The current

phylogenetic analyses based on various datasets have largely established the

phylogenetic framework of Hymenoptera and identified several key innovative events in

their evolution[3,4,6,7]. In brief, the origin of the Hymenoptera is estimated to be

approximately 281 million years ago (Mya) in the Permian period[3]. It is likely that the

ancestral hymenopterans are plant-feeders, and this lifestyle is still retained in the

majority of sawflies, which represent the early branches of the Hymenoptera tree[3,4]. The

emergence of parasitoidism occurred in the last common ancestor of Orussoidea and

wasp-waisted wasps (also known as Apocrita)[3,4]. This transition from phytophagy to

parasitoidism is regarded as a key innovation in the evolution of Hymenoptera and may

have been a significant driver of species diversification, as approximately 70% of all

described hymenopterans are parasitoids[3,4,8]. In Apocrita, species have evolved a highly

constricted "waist" between the first and second abdominal segments, and this wasp-

waist allows for flexibility in female ovipositors, which may be an important adaptation for

parasitoid lifestyles[4,9]. Another morphological innovation is the acquisition of the

venomous stinger (the modified female ovipositor from an egg-laying apparatus to a

stinging apparatus) in Aculeata[3,4]. And some stinging species in the Aculeata (e.g., ants

and certain groups of bees and wasps) have also evolved into eusociality independently[3].

Furthermore, it is important to note that following the evolution of the basal hymenopteran

diet from phytophagy to parasitoidism (equivalent to carnivory), certain gall wasps, fig

wasps, and pollen-collecting bees have independently evolved again to phytophagy (i.e.,

secondary phytophagy)[4]. A recent phylogenetic study has indicated that the evolution of

secondary phytophagy had a significant influence on the diversification rate in

Hymenoptera[4]. Overall, the evolution of Hymenoptera has been accompanied by several

key innovations and evolutionary changes of lifestyles; however, the genetic mechanisms

underlying these evolutionary processes remain largely unknown.

The sequencing of numerous hymenopteran genomes has opened new avenues for a

deeper understanding of Hymenoptera evolution, yet a comprehensive, large-scale

comparative genomic analysis of Hymenoptera is lacking. To fill this gap, we performed a

large-scale comparative genomics study containing 131 species, which represent major

lineages of Hymenoptera. Using the genomic data, we systematically investigated: (1)

the general patterns of genome evolution in terms of gene content and evolutionary rate

throughout hymenopteran history; (2) genomic changes occurred at the evolutionary

branches associated with key innovations; (3) differences in genome evolution between

two major clades in the Hymenoptera phylogeny, the Parasitoida and Aculeata; and (4)

genomic features related to three independently evolutionary events of secondary

phytophagy. Our findings not only offer a comprehensive picture of the genome evolution

of hymenopteran insects but also highlight several genomic changes potentially related to

evolutionary innovations, shedding light on the genomic basis of adaptation and

diversification within this remarkable order.

Results and Discussion

Hymenoptera phylogenomics

To explore the evolutionary history of Hymenoptera, we obtained high-quality genomes

for 131 hymenopteran insects from 13 superfamilies (29 families), covering many

representative lineages with unique lifestyles such as sawflies, wasps, ants, and bees

(Supplementary Table 1). These genomes were all publicly available, and 78 of them

were sequenced using long-read sequencing technologies, showing the overall high

contiguity (mean Contig N50 4.19 Mb). Furthermore, BUSCO (Benchmarking Universal

Single-Copy Orthologs) assessments indicated that these genomes have a high level of

gene completeness (highest 99.80%, lowest 85.10%, mean 97.26%, s.d. = 0.03)

(Supplementary Table 1). In order to minimize the impact of potential gene annotation

issues on subsequent analysis, we then employed an alignment-based approach to

identify potential broken and chimeric genes[10], correcting about 3.6% of total genes

(Supplementary Table 2). After gene corrections, orthology inference produced 33,405

orthogroups (OGs, also referred to as gene families), with 25.1% of them present in over

70% of hymenopterans (Fig. 1, Supplementary Fig. 1, and Supplementary Table 3). Of

the total number of genes analyzed, 78,789 (4.77%) were identified as singletons,

meaning that no homolog was predicted in any genome.

We next reconstructed the phylogeny of hymenopterans based on protein sequences

from 1002 single-copy genes (Fig. 1). Only nine (6.98%) nodes in this tree fail to receive

100% bootstrap support (Supplementary Fig. 2). Our species tree was mostly consistent

with previous hymenopteran phylogenies using transcriptomic and mitochondrial data[3,7].

We showed that: (i) phytophagous sawflies from Tenthredinoidea and Cephoidea are

placed in the early branches of Hymenoptera; (ii) Parasitoid Orussoidea is a sister to all

wasp-waisted wasps, also known as Apocrita; and (iii) in Apocrita, parasitoid wasps from

several superfamilies such as Ichneumonoidea, Chalcidoidea, Cynipoidea, and

Platygastroidea form one clade (i.e., Parasitoida), and stinging species of Aculeata form

its sister clade (Fig. 1). This phylogenetic relationship within Apocrita differs from the

results of studies using ultraconserved elements[4,6]. These studies support that

Ichneumonoidea is a sister to the clade containing all other species of Apocrita. There

may be several reasons for this conflict, such as the type of data used and the samples

analyzed. Further studies should expand the sampling to establish a more robust tree for

Hymenoptera. In our subsequent analyses, we used the phylogenetic tree constructed

from the genomic data, and the divergence times were estimated using 13 calibration

points based on a previous study[3] (Fig. 1 and Supplementary Fig. 2).

Genomic changes throughout Hymenoptera history

We next investigated the changes in three different genomic features of protein-coding genes, including gene family evolution, protein domain rearrangement, and the evolutionary rate of protein-coding genes, to explore the overall dynamics of Hymenoptera genome evolution. First, we inferred the history of gene family evolution and identified 63,907 gene family expansions and 106,004 contractions (Supplementary Figs. 3–4). This pattern suggests a generally reductive mode of gene family evolution in Hymenoptera, which is also supported by a small-scale comparative genomic study of this order[11]. The pincer wasp *Gonatopus flavifemur* (Chrysidoidea, Dryinidae) is notable among hymenopterans for having the largest number of gene family expansion (736) and contraction (2564) events (Supplementary Fig. 3). We also estimated rates for gene family size evolution and found the average global rate of gene gain and loss of gene families in Hymenoptera was estimated to be 58.29 gene gain and loss per million years (My). The rates of most hymenopteran branches are relatively comparable, but we also noticed several notable accelerations of the gene gain and loss rates in different lineages. For example, the clade comprising *Nasonia*, *Trichomalopsis*, and *Pteromalus* in Pteromalidae exhibited an average gene gain and loss rate of 605.74 per My. Furthermore, a noticeable acceleration in gene gain and loss rates can be observed in the branches of leafcutter ants (Formicidae), as previously noted in studies[12,13] (Supplementary Figs. 5–6 and Supplementary Table 4).

We identified 230 gene families that exhibited significant expansions or contractions during Hymenoptera evolution, and GO (Gene Ontology) enrichment analysis showed that they were enriched for the steroid metabolic process, the maltose metabolic process, and sensory perception of smell (FDR-adjusted $P < 0.05$; Supplementary Tables 5–6 and Supplementary Fig. 7). Among these, several rapidly evolved digestion-related gene families, such as Maltase, Esterase, and Trypsin, may be associated with adaptation to different diets in Hymenoptera (Supplementary Table 5). The rapid evolution of the Fatty acyl-CoA reductase family involved in the biosynthesis of insect pheromones may be driving the evolution of pheromone diversity[14–16]. The significant expansions and

contractions of Odorant receptor families may be related to the rapid adaptations to

detect the different odors from the living environments (such as pheromones and odors

related to host plants or host insects) of diverse hymenopterans. We also found that the

size of the cuticular protein family changed rapidly during the Hymenoptera evolution

(Supplementary Fig. 8), which might be associated with the evolutionary diversity of the

cuticular exoskeleton and morphology[17,18]. Detailed analysis further indicated that the

size variation of the cuticular protein family mainly occurred in the CPR-RR-1, CRP-RR-

2, and CPAP1 subfamilies (Supplementary Table 7 and Supplementary Fig. 8).

We also investigated the evolution of protein domain families based on the Pfam

annotations. Similarly, we identified 105 Pfam domain families with significantly

accelerated rates, including OS-D (Insect pheromone-binding family, A10/OS-D,

PF03392.12), FA_desaturase (Fatty acid desaturase, PF00487.23), Chitin_bind_4 (Insect

cuticle protein, PF00379.22), 7tm_2 (7 transmembrane receptor, PF00002.23), and

Lipase (Lipase, PF00151.18), consistent with the significantly changed gene families

(Supplementary Table 8). Furthermore, we reconstructed the rearrangements of protein

domains in Hymenoptera evolution (see Methods). In total, 33,943 domain arrangement

changes were observed, and 41.8% of them were formed by a fusion of two ancestral

domains (Supplementary Table 9). We noticed the higher rates of domain

rearrangements in several lineages, such as *G. flavifemur*, *Temnothorax longispinosus*,

and *Diadromus collaris*, which may contribute to the evolutionary innovations of these

species (Supplementary Table 9 and Supplementary Fig. 9).

We next explored the changes in the evolutionary rate of the protein-coding genes in

Hymenoptera. We first estimated the rates of amino acid substitutions across the

Hymenoptera phylogeny to reveal the overall pattern of protein sequence evolution (Fig.

1). Our analysis estimated an average amino acid substitution rate of 0.0025

substitutions per site per My with a standard deviation of 0.0055 (Supplementary Table

10). We found that the overall accelerated protein evolution events were dispersed

across the multiple lineages of Hymenoptera, such as some clades in Anthophila

(Apoidea) and Braconidae (Ichneumonoidea). These evolutionary rate shifts may be

related to phenotypic transitions. For example, a comparative genomic study has

suggested that the accelerated evolution of a number of proteins is linked to the

independent miniaturization across parasitoid wasps[19]. Then, we examined the

evolutionary rate of each orthologous protein-coding gene by measuring both the

normalized amino acid substitution rates and the selective constraint (ratio of non-

synonymous to synonymous substitutions, d_N/d_S). Functional analysis showed that genes

associated with gene expression evolved most rapidly, followed by genes related to

immunity, metabolism, and sensory perception (Fig. 2B). The rapid evolution of genes in

these functional categories may play a role in phenotypic innovation and adaptive

evolution in Hymenoptera. In contrast, genes with slower evolutionary rates were mainly

enriched in some housekeeping processes, such as cellular component organization and

cell development (Supplementary Fig. 10). In addition, we also conducted whole genome

alignment analysis (Cactus alignment) to investigate the evolutionary constraints and

acceleration at the whole genome level (see Methods). We chose the bumblebee

Bombus terrestris (Apidae) as a reference because the availability of ATAC-Seq data is

helpful to infer potential regulatory elements[20]. In the 27-way alignment, 59.5%

(233.69/392.96 Mb) of the *B. terrestris* genome was aligned to at least one species, and

21.5% and 8.5% of the genome were identified as constrained and accelerated regions,

respectively, by calculating the phyloP score (Supplementary Fig. 11A). We then

identified the top 5% most accelerated and most conserved genes, measured by the

average phyloP score of the coding sequence of each gene (Supplementary Table 11).

GO enrichment analyses indicated that the most constrained genes were enriched in

structure morphogenesis and development, while the most accelerated genes were

involved in sensory perception and steroid metabolic processes, consistent with our

previous findings in Fig. 2B (Supplementary Figs. 11B–C and Supplementary Tables 12–

13). We also identified 481,924 highly conserved elements using phastCons, and 27,602

(5.7%) of them were conserved across all species in our 27-way alignment

(Supplementary Fig. 12A). Among them, 151 non-coding regions were overlapped with

the accessible chromatin regions supported by ATAC-Seq in the reference species

B. terrestris, suggesting their potential functions in gene regulation. We further found that the genes located near these conserved non-coding elements were functionally related to nervous system development and cell morphogenesis (FDR-adjusted *P* < 0.05; Supplementary Fig. 12B and Supplementary Table 14).

Genomic characteristics accompanying key innovations in Hymenoptera evolution

We next examined several key nodes in the evolutionary history of Hymenoptera, investigating the genomic changes that occurred at these nodes with the aim of correlating the association of genomic changes and phenotypic innovations. Here, we focused on four key evolutionary positions: the origin of parasitoidism (node A in Fig. 2A), the emergence of the wasp-waist (node B), the stem of Parasitoida that is prior to the radiation of the majority of parasitoid wasps (node C), and the emergence of the stinger (node D). The genomic features related to secondary phytophagy in several independent lineages will be discussed in a subsequent section. We did not investigate the genome evolution of eusocial hymenopterans due to the significant contributions of some recent works in this area[21–26].

We first focused on the last common ancestor of the parasitoid wood wasp *Orussus abietinus* (Orussidae) and other wasp-waisted species (Apocrita), which represents the position of parasitoidism origin in the Hymenoptera tree. Our analysis identified 115 contracted and no expanded gene families at this node (Fig. 2A and Supplementary Table 15). Furthermore, we also found 137 missing gene families (that is, gene families present in over 70% of outgroup species and are lost in the test group) and only five novel gene families (also known as novel core genes, which are retained in over 70% of species in the test group) (Supplementary Tables 16–17). This pattern of gene family evolution suggests that the reductive evolution of gene families may have played an important role in the origin of parasitoidism. GO enrichment analysis showed that these contracted and missing families were significantly enriched for functions associated with detection of stimulus, cholesterol transport and metabolism, lipid metabolism, and xenobiotic metabolic processes (FDR-adjusted *P* < 0.05; Supplementary Tables 18–19).

The changes in these gene families in the ancestral parasitoid wasp could be associated

with the shift in lifestyle and diet from a phytophagous sawfly to an insectivorous

parasitoid. For instance, the genes related to detection of stimulus (e.g., Gustatory and

odorant receptors, Glutamate receptors, and Transient receptor potential channels) may

contribute to early parasitoid wasps locating potential hosts in novel environments (i.e., in

dead wood for the most parasitoids in Orussidae[11]). The contracted gene families with

cholesterol-transporting and metabolizing functions may be associated with alterations in

the nutritional composition of the foods consumed after the acquisition of parasitoidism.

In addition, we identified 96 genes that show evidence of positive selection in the

parasitoid ancestor (aBSREL, FDR-adjusted $P < 0.05$; Supplementary Table 20). These

genes were associated with many developmental processes, including muscle tissue

development, eye morphogenesis, respiratory system development, and larval salivary

gland morphogenesis (Supplementary Table 21). These genomic changes may be

related to the dramatic changes in morphological characteristics that are considered

adaptations to a parasitoid lifestyle, such as simplified larval morphology lacking eyes

and legs in Orussidae[11,27].

In the last common ancestor of wasp-waisted species (Apocrita), we also identified more

contracted gene families (30) than expanded gene families (7) (Fig. 2A and

Supplementary Table 22). The contraction of Cuticle protein genes that have roles in

cuticle development at the Apocrita stem may contribute to the emergence of wasp-waist.

We also discovered 119 positively selected genes with diverse functions such as

endocytosis and regulation of enzyme activity (aBSREL, FDR-adjusted $P < 0.05$;

Supplementary Table 23). At the stem of stingers, the Aculeata, 434 contracted gene

families were found, but only 12 expanded gene families were found (Supplementary

Table 24). The expanded genes included the Elongation of very long chain fatty acid

protein genes that are involved in female pheromone biosynthesis[28,29]. The contracted

genes with functions related to cuticle development (e.g., Cuticle protein genes and

Laccase genes) were also identified. Moreover, our analysis revealed 175 positively

selected genes at the stem of the Aculeata (Supplementary Table 25). These included the

genes that regulate the highly conserved pathways for development, such as Wnt

signaling pathway[30] (e.g., *FZD2*, *LRP6*, and *MLLT3*) and Notch signaling pathway[31] (e.g.,

Ift172, *NCSTN*, and *POFUT1*). Changes in these genes may have contributed to the

unique morphological features of stinging hymenopterans.

The Parasitoida includes a large number of parasitoid wasp species, and the stem of the

Parasitoida clade showed 235 contracted and 15 expanded gene families (Fig. 2A and

Supplementary Table 26). We found the expansion of Venom allergen genes and Venom

metalloproteinase genes, which may contribute to venom functions to enhance the

probability of parasitic success. Additionally, we inferred that the expanded salivary

secreted peptide family may also play a role in feeding on hosts and regulation of host

immunity, based on current knowledge of the functions of the salivary proteins in

parasitoid wasps[32–34]. This result suggests that the expansion of gene families involved in

parasitoid-host interaction occurred prior to the radiation of parasitoid wasps. Taken

together, our findings provide a number of candidate genes that may be linked to the

evolution of key innovations in Hymenoptera.

Divergent trajectories of genomic change between Parasitoida and Aculeata

The Parasitoida and Aculeata clades diverged from the ancestral wasp-waisted wasp at

around 243 Mya, representing two major groups of hymenopterans with clear differences

in lifestyle, morphology, and diversity[3] (Fig. 1). The Parasitoida clade includes primarily

parasitoid wasps that lack a stinger, and the Aculeata clade comprises stinging species

with diverse life habits, including stinging parasitoids, social wasps, ants, and bees[3,8]. We

next sought to explore and compare the genomic changes within these two clades, which

may provide insights into the independent evolution of two major groups of

hymenopterans.

Depending on the results of gene family evolution, we determined the distribution of the

rapidly evolving events for each gene family on the branches in the Parasitoida and

Aculeata clades. This thus allowed us to explore whether there are gene families with a

clear distributional preference for rapid evolutionary events in the two clades. A total of 91

gene families were identified that exhibited rapid evolutionary events, with a significantly

higher prevalence observed in the Parasitoida compared to Aculeata (FDR-adjusted $P <$

0.05, Chi-square test, odds ratio > 1; Fig. 3A, Supplementary Fig. 13, and Supplementary

Table 27). These included a number of Histone genes (i.e., H2A, H2B, H3, and H4) that

are involved in the structure of chromatin[35,36], suggesting the potential distinct chromatin

structures in parasitoid wasps. The enrichment of rapidly evolving Trypsin, Glucose

dehydrogenase, and UDP-glucosyltransferase families in the Parasitoida may be

associated with the evolutionary adaptations of digestion and detoxification metabolism,

enabling the parasitoid wasps to adapt to diverse arthropod hosts. The rapid evolution of

the Peptidoglycan recognition protein family in the Parasitoida may contribute to the

antimicrobial defenses of parasitoid wasps[37,38]. Furthermore, we observed significant

changes in the SET and MYND domain-containing protein family, an epigenetic

regulator[39], in the Parasitoida, which suggests that epigenetic regulation may play an

important role in the evolution of parasitoid wasps. By contrast, only 12 gene families

were identified as enriched rapidly evolving families in the Aculeata, including the THAP

domain-containing protein, RCC1 and BTB domain-containing protein, Odorant receptor,

and Gustatory receptor (FDR-adjusted $P <$ 0.05, Chi-square test, odds ratio > 1;

Supplementary Table 28 and Supplementary Fig. 14). In addition, we estimated and

compared the rates of gene gain and loss for each gene family between the Parasitoida

and Aculeata clades to determine the gene families with significantly different gene gain

and loss rates within the two clades. This analysis may capture additional gene families

that have undergone significant evolutionary changes between the Parasitoida and

Aculeata, which may have been overlooked in previous analyses due to the limited

number of evolutionary branches where such changes occurred. In this analysis, we

identified 1604 gene families with significantly higher rates of gene gain and loss in the

Parasitoida compared to Aculeata (FDR-adjusted $P <$ 0.05, One-tailed Mann-Whitney U

test; Fig. 3B and Supplementary Table 29). GO analysis of the gene families with high

turnover rates in the Parasitoida showed enrichment for functional categories such as

polysaccharide digestion, intracellular sterol transport, and prostaglandin secretion (FDR-

adjusted $P < 0.05$; Fig. 3C and Supplementary Table 30). Additionally, we noticed some

genes related to venom functions were included in this gene set, suggesting venom may

play a key role in the evolution and diversity of parasitoid wasps in Parasitoida (Fig. 3B;

discussed in detail in the next section). Similarly, we found a small number of gene

families (26) with faster gene gain and loss rates in the Aculeata compared to Parasitoida

(FDR-adjusted $P < 0.05$, One-tailed Mann-Whitney U test; Supplementary Table 31).

We next compared the selective constraint (d_N/d_S) of single-copy OGs between the

Parasitoida and Aculeata. The result showed none of the OGs had significantly different

d_N/d_S between the two clades (Two-tailed Mann-Whitney U test; Supplementary Table

32). We hypothesized that this result may be attributed to the presence of significant

variability in the d_N/d_S ratio of the branches within the two clades. In order to investigate

the degree of variation in d_N/d_S between the Parasitoida and Aculeata, we calculated the

coefficient of variation (CoV) of the d_N/d_S in the two clades (Supplementary Table 33).

Our findings revealed that the d_N/d_S variation of the Parasitoida was significantly larger

than that of the Aculeata ($P = 7.92e{-}84$, One-tailed Mann-Whitney U test; Supplementary

Fig. 15). And 73.8% of the analyzed OG exhibited a greater CoV of the Parasitoida than

that observed in the Aculeata (Supplementary Table 33). This result suggests that the

Parasitoida may be subject to more variable selection pressures compared to the

Aculeata. This is likely to be related to the diverse lifestyles and hosts of the parasitoid

wasps in Parasitoida. We also identified 13 OGs with obviously higher CoV of the d_N/d_S in

the Parasitoida (i.e., the OGs with the top 10% CoV in Parasitoida and the bottom 10%

CoV in Aculeata), including exocytosis-related STXBP5, calcium-binding protein

Calmodulin, and cytoskeleton organization-related DIAPH, suggesting that they may

have evolved under different selective pressures and may play key roles in Parasitoida

evolution (Supplementary Table 34).

We further explored the variation in metabolic pathways between the Parasitoida and

Aculeata, which may reflect differences in the life history of these two groups. To

compare the pathway repertoire variation between the two clades, we first performed EC

(Enzyme Commission number) annotations for all 131 hymenopterans (Supplementary

Table 35) and constructed ancestral KEGG (Kyoto Encyclopedia of Genes and

Genomes) pathways based on eight phytophagous sawflies (Tenthredinoidea) located on

the early branches of the Hymenoptera tree (see Methods). Then, we mapped the

present/absent information of each EC of the species from the Parasitoida and Aculeata

to the ancestral pathways. This allows us to study the pathway-level changes of

Parasitoida and Aculeata after their divergence. We examined the pathway coverage

(PC, i.e., the fraction of ECs in an ancestral pathway that were annotated in a species) of

each pathway in each species (Supplementary Table 36). Overall, our results showed

significantly lower PC for Parasitoida compared to Aculeata (P = 2.68e-46, One-tailed

Mann-Whitney U test; Supplementary Fig. 16), suggesting that the specialized

parasitoidism may have simplified the pathways of parasitoid wasps, and similar

phenomena have also been demonstrated in other parasitic species, such as parasitic

worms[40], dicyemids[41], protists[42], and plants[43]. Our detailed analysis further revealed that

37 pathways from 14 KEGG superpathways exhibited significantly lower PC in

Parasitoida when compared to Aculeata, which are involved in lipid metabolism,

metabolism of cofactors and vitamins, carbohydrate metabolism, etc. (FDR-adjusted P <

0.05, One-tailed Mann-Whitney U test; Fig. 3D, Supplementary Fig. 17, and

Supplementary Table 37). Conversely, five pathways showed lower PC for Aculeata, and

these metabolic processes included purine metabolism, pentose phosphate pathway, and

glyoxylate and dicarboxylate metabolism (FDR-adjusted P < 0.05, One-tailed Mann-

Whitney U test; Fig. 3D, Supplementary Fig. 17, and Supplementary Table 38). Among

these PC variations, we inferred the lower PC of arachidonic acid metabolism (ko00590)

and folate biosynthesis (ko00790) in Parasitoida caused by the pervasive absence of

carbonyl reductase (K00079, EC. 1.1.1.184/189/197). In addition, we found that the loss

of the same enzyme may have different effects on these two pathways. Specifically, the

absence of carbonyl reductase in the arachidonic acid metabolic pathway has disrupted

the pathway from prostaglandin E2 to prostaglandin F2alpha. The widespread absence

of this pathway in parasitoid wasps suggests that they may have other ways of acquiring

prostaglandin F2alpha, probably from their diet. However, the absence of carbonyl

reductase in the folate biosynthetic pathway did not result in pathway disruption due to

the redundancy in this pathway. This may represent a streamlined evolution of the

pathway in parasitoid wasps. We next compared the CoV of the PC between the two

clades and found six superpathways exhibited significantly higher variation in

Parasitoida, also covering the pathways we discussed above (FDR-adjusted P < 0.05,

One-tailed Mann-Whitney U test; Supplementary Fig. 18). This finding suggests that

these pathways are more variable in parasitoid wasp species, and it is likely that these

variations occurred several times independently in Parasitoida evolution. This may

suggest that the rapid changes in these pathways are related to the adaptation of

parasitoid wasps to diverse parasitoid lifestyles, such as adaptation to different hosts and

different parasitic strategies.

Massively duplicated genes and gene losses in Parasitoida

The preceding section of this study has demonstrated that parasitoid wasps exhibit high

rates of gene gain and loss, which likely results in significant changes in the number of

members of certain gene families. Here we first explored the gene families with

significantly larger gene numbers in Parasitoida than those in other taxa, which were

likely to be caused by the massive gene duplications that occurred in Parasitoida. In total,

we identified 95 gene families with a significantly larger number of genes in Parasitoida

species compared to other hymenopterans (FDR-adjusted P < 0.05, One-tailed Mann-

Whitney U test; Supplementary Table 39 and Supplementary Fig. 19). Functional

annotation of these families was diverse, but we observed they were frequently related to

chemoreception (such as Odorant receptor, Gustatory receptor, and General odorant-

binding protein), detoxification (such as Multidrug resistance-associated protein and

UDP-glycosyltransferase), and protein degradation (such as Proteases and Protease

inhibitors). Among these gene families, we also found 41 families with no annotation

information, designated as Putative uncharacterized proteins. In contrast, we only found

nine enlarged gene families for Aculeata, with three families belonging to the Odorant

receptor (Supplementary Fig. 20 and Supplementary Table 40). The findings suggest that

large-scale gene duplication was more prevalent in Parasitoida, and the identified gene

families that exhibited massive expansions may have played roles in the adaptive

evolution of parasitoid wasps and promoted the diversity of these species.

Among the gene families identified with large-scale gene duplications in Parasitoida, we

found several families whose members are often known as venom components, the

important weapons of parasitoid wasps to overcome host immune systems or manipulate

host development[44,45]. For instance, our analysis indicated that the Venom

metalloproteinase family has undergone an extensive expansion, specifically in the

Parasitoida clades, which may have contributed to the recruitment of the family member

as a venom component in parasitoid wasps through neofunctionalization (Fig. 4A and

Supplementary Fig. 21). Indeed, Venom metalloproteinase has been reported as venom

protein in at least 19 parasitoid wasps from 5 families of 3 superfamilies in Parasitoida[46–

62]. And some studies have demonstrated that these venom proteins play distinct roles

(such as regulating host development[57] and inhibiting host immunity[63]) in different

parasitoid-host systems, suggesting the duplications of this family in parasitoid wasps

may also be related to the diversity of venom functions. Additionally, the duplication of

Venom metalloproteinase genes may also help diversify nonvenomous functions. In

Anastatus fulloi (Chalcidoidea, Eupelmidae), our gene expression profiling showed

diverse expression patterns of Venom metalloproteinase genes, suggesting that they

have functions other than venom functions (Fig. 4B). For example, nine of them exhibited

specifically higher expression in the larval stage, suggesting their possible roles in

digestion or parasitoid-host interaction as salivary proteins (Fig. 4B and Supplementary

Table 41). Similarly, the expanded Venom allergen family in Parasitoida may also be

related to the venom evolution of parasitoid wasps (Fig. 4C and Supplementary Fig. 22).

And we found 15 parasitoid species have used the members of this family as their

venom[47,53,54,58–61,64–69]. Furthermore, the gene family also exhibited diverse gene

expression patterns, which suggests that the gene may have multiple functions (Fig. 4D

and Supplementary Table 42). Other enlarged gene families in Parasitoida that have

evidence related to venom components include Neprilysin[70,71], Serpin[53,72,73] and

Trypsin[62]. Overall, our findings indicate that the extensive duplication of certain genes in

parasitoid wasps has played a role in the evolution of venom diversity. This may enhance

the adaptability of parasitoid wasps to different hosts, thereby contributing to the species

diversity in Parasitoida.

We also identified 199 gene families that were completely missing in a higher proportion

of Parasitoida than in other taxa (i.e., the gene family is lost in more than 20% of

parasitoids but in no more than 2% of other taxa) (Supplementary Table 43). Several

losses in these gene families may be associated with the parasitoid lifestyle of species in

Parasitoida. For example, we hypothesized that the loss of some gene

families/subfamilies associated with detoxification metabolism (such as UDP-

glycosyltransferase, Cytochrome P450, and Nephrin) may be related to the reduction of

toxic substances contained in the hosts of some parasitoids. Eleven gene families with

GO terms related to embryo development exhibited a higher loss proportion in parasitoid

wasps (Supplementary Table 43). This suggests that the early developmental processes

of parasitoid wasps may have been altered compared to other hymenopterans, possibly

in response to adaptations to parasitic life. In addition, our analysis also recovered our

previous finding that Vitellogenin and its receptor (Vitellogenin receptor) have been

repeatedly lost in many parasitoid wasps[74]. The loss of these genes may result in a

reduction or absence of yolk proteins in parasitoid wasp eggs, which is consistent with

some phenotypic observations[75–77]. It is proposed that this may be related to the fact that

parasitoid wasp eggs are able to obtain nutrients from the host.

Genomic features of independent secondary phytophagy

During the evolution of Hymenoptera, a shift from phytophagy to parasitoidism occurred

at the ancestral node of Orussidae and Apocrita, after which some hymenopterans from

different superfamilies re-evolved into phytophagy (i.e., secondary phytophagy). And a

recent evolutionary study has proposed that secondary phytophagy in Hymenoptera

plays a key role in their species diversity[4]. However, the genomic basis for the

hymenopterans that evolved back to phytophagy is still unknown. Moreover, it is worth

exploring whether there are any similarities in the genomic features underlying these

independent transitions (i.e., convergent or parallel evolution). In the present study, our

Hymenoptera phylogeny captured three independent transitions to secondary

phytophagy, locating at the terminal branch leading to the gall wasp *Belonocnema kinseyi*

(Cynipoidea, Cynipidae) (node E in Fig. 2A), the ancestral branch of two fig wasps

Ceratosolen solmsi and *Eupristina verticillata* (Chalcidoidea, Agaonidae) (node F), and

the ancestral branch of pollen-feeding bees (Apoidea, Anthophila) (node G; Fig. 1). First,

we examined the genomic changes, including gene family expansions/contractions and

genes with positive selection on these target branches. In general, the genomic changes

of these three branches showed a high degree of branch specificity, both in terms of the

number of changes and the composition of genes or gene families (Fig. 5A and

Supplementary Tables 44–46). In addition, our analysis did not reveal any instances of

gene family expansion, contraction, or positive selection of genes that were common to

all three branches (Fig. 5A). This result suggests that the genomic changes behind the

three independent transitions to phytophagy in Hymenoptera may be distinct. This may

be attributed to differences in the types of plant-derived foods consumed (e.g., plant galls

for the gall wasp and pollen for bees). Nevertheless, we did identify some genomic

changes that were shared by two branches or specific to one branch, which may be

associated with secondary phytophagy (Fig. 2A). For example, we observed that the

Maltase family has expanded on the branches related to the secondary phytophagy of

both the gall wasp and fig wasps, which probably play a role in enhancing the

carbohydrate metabolism of these species (Supplementary Table 44). This is also

supported by the expansion of Glucose dehydrogenase family in the gall wasp *B. kinseyi*.

We also noticed the family contractions (e.g., Venom metalloproteinase and Venom

allergen) related to venom functions in the gall wasp and fig wasps (Supplementary Table

45). This may be indicative of the degradation of the venom components originally used

to attack insect hosts following their transformation to phytophagy. In bees, a large

expansion of Odorant receptors was observed, probably to improve the ability to

recognize and distinguish volatile substances in plants so as to effectively recognize and

select food. Additionally, we found the Peritrophin family expanded, which is regarded as

the most important physical immune barrier of the gut in arthropods[78], possibly as an

adaptation from sarcophagy to pollen feeding. In contrast, the shrinkage of the Uricase

gene could be an evolutionary response to the lower metabolic capacity for purine

degradation[79]. In *B. kinseyi*, the expansion of Glutathione S-transferase could also be

required for the detoxification of phytochemicals[80,81] (Fig. 2A). Meanwhile, positive

selection analysis resulted in 97, 73, and 148 positively selected genes in fig wasps,

bees, and the gall wasp, respectively, of which 272 genes were exclusive (Fig. 5A and

Supplementary Table 46). We further discovered that 67.28% of OGs to which exclusively

positively selected genes belong were clustered into 19 functional clusters composed of

OGs with genes evolving under positive selection exclusively in three transition branches,

probably indicating the functional convergence signal (Fig. 5B and Supplementary Table

47). However, no more than 30% of expanded and contracted OGs could be clustered

(Supplementary Figs. 23–24 and Supplementary Tables 48–49). Overall, the genomic

signatures of dietary changes can be summarized into three functional categories:

chemoreceptor, digestion, and detoxification.

Plants exploit the glucosinolate-myrosinase system to produce toxic hydrolysis products

to defend themselves from herbivores[82,83]. In *Athalia rosae* and some aphids,

glucosinolate can be rapidly sequestered into the hemolymph, where it is hydrolyzed by

myrosinases[11,84–86]. Thus, a high copy number of the myrosinase gene may characterize

phytophagous species[11]. However, in our study, the copy number of myrosinase was

indeed high in phytophagous sawflies, but there was no expansion in secondary

phytophagous species (Supplementary Fig. 25). Therefore, we speculate that secondary

phytophagous species may employ strategies of detoxification metabolism that are

different from those of phytophagous species.

Conclusion

In this study, our evolutionary and comparative genomic analyses revealed many

genomic changes (such as gene family evolution, protein domain evolution, evolutionary

rate, and selection) in the evolutionary history of Hymenoptera. We correlated some

genomic changes with specific innovations, including the evolution of parasitoidism,

wasp-waist, sting, and secondary phytophagy. In comparing the two major hymenopteran

clades, Parasitoida and Aculeata, we uncovered distinct genomic features indicative of

their independent evolutionary trajectories, including variations in pathway coverage and

gene family dynamics, which may reflect adaptation of their divergent lifestyles (highly

parasitoidism vs. complex free-living habits, which are not limited to parasitoidism).

Furthermore, we investigated the genomic basis for the hymenopterans that evolved

back to phytophagy in three independent lineages. Overall, the results of our study

provide insights into the evolution and diversification of Hymenoptera.

Methods

Data preparation

We collected publicly available genomic and annotation data for 131 Hymenoptera

species from 13 superfamilies and 29 families. To avoid redundancy, we selected a

representative protein isoform (i.e., the isoform with the longest amino acid sequence) for

each gene in each species. Assembly quality of each genome was assessed with

QUAST 5.2.0[87], and completeness was assessed with BUSCO 5.5.0[88] (-m prot -l

insecta_odb10). Finally, the average BUSCO score reached 97.26%. For more detailed

information about species and public data repositories, see Supplementary Table 1. To

reduce the impact of potential gene annotation errors, we corrected potential chimeric

genes and broken genes. We first used Broccoli v1.2[89] to identify OGs from all protein

sequences of 131 species. Genes assigned to two or more OGs by Broccoli v1.2 were

considered chimeric genes. Those that belonged to the same OG, were sequenced on

the same strand on the same chromosome, showing low sequence similarity (less than

30% calculated using MAFFT v7.310[90]), and matching different portions of the longest

homolog protein in *Nasonia vitripennis* were considered broken genes. Detailed

correction procedures for chimeric and broken genes are referenced in a previous

study[10].

Phylogenetic analysis

We identified 33,405 OGs from all protein sequences of 131 species with OrthoFinder

2.5.4[91]. The protein sequences of 1002 single-copy genes were aligned with MAFFT

v7.310[90] and concatenated into supergenes. Then, these sequences were used to

construct a maximum-likelihood phylogenetic tree using IQ-TREE 2.0.3[92] with 1000

ultrafast bootstraps. The best-fitting model estimated by ModelFinder[93] was JTT+F+R10.

Species divergence times were estimated with r8s1.81[94] using 13 time points based on

previous studies[3]. The time points were as follows: Hymenoptera: 281 Mya,

Tenthredinoidea: 106–176 Mya, Orussoidea + Apocrita: 211–289 Mya, Apocrita: 203–276

Mya, Ichneumonoidea: 151–218 Mya, Chalcidoidea: 105–159 Mya, Cynipoidea: 69–125

Mya, Aculeata: 160–224 Mya, Apoidea: 128–182 Mya, Vespoidea: 114–180 Mya,

Formicidae: 65–127 Mya, Apidae: 68–99 Mya, and Braconidae: 116–177 Mya.

Gene family evolution

We used CAFE5[95] to determine gene family expansion and contraction for each branch,

using the result of OrthoFinder and a species-level phylogeny with divergence times as

inputs. Gene families with a P value less than 0.05 were considered to have significant

expansions and contractions. In order to estimate the differences in the distribution of

rapid evolutionary events between Parasitoida and Aculeata, we used Chi-square tests,

and if FDR-adjusted $P < 0.05$ and odds ratio > 1, the gene family was considered to have

a significantly different occurrence rate of rapid evolutionary events between Parasitoida

and Aculeata. The gene gain and loss rate for each clade was calculated as the sum of

gene gains and losses for all OGs divided by the divergence time. For each OG, the

gene gain and loss rates of different branches were calculated as the count change of

OG between it and its parent clade divided by the corresponding divergence time.

Cuticular protein family analysis

BLASTP analyses were performed against protein sequences of all species using

cuticular protein sequences from hymenopteran species (i.e., *Apis mellifera*, *Microplitis*

mediator, *P. puparum*, and *N. vitripennis*) as probes (-evalue 1e-5). Meanwhile, the

genomes of all species were searched by TBLASTN with the above probes. The domains

of proteins aligned with BLASTP and TBLASTN were predicted using hidden Markov

model searches with HMMER v3.4 (https://github.com/EddyRivasLab/hmmer) against

Pfam database (http://pfam.xfam.org/). For the cuticular proteins analogous to the

peritrophins (CPAP) family, the CBM_14 domain (PF01607) was required, of which the

CPAP1 subfamily had one and CPAP3 had three. For the cuticular proteins with Rebers

and Riddiford (R&R) consensus (CPRs), the Chitin_bind_4 domain (PF00379) was

required, and they were divided into RR-1 and RR-2 subfamilies according to the best

hit. The CPF family was identified based on the Cuticle_3 domain (PF11018) and the

Tweedle (TWDL) family based on the DUF243 domain (PF03103).

Selection pressure analysis

We selected 27 representative species of the major groups of Hymenoptera and

identified 2060 one-to-one orthologue protein-coding genes using OrthoFinder 2.5.4[91]; for

detailed species, see Supplementary Table 50. HyPhy[96] package was used to detect

gene selection signals at key points in Hymenoptera evolution. aBSREL[97] was used for

detecting positively selected genes (FDR-adjusted $P < 0.05$, Bonferroni-Holm correction).

To explore the selection pressure of genes in each clade, we calculated d_N/d_S using

PAML 4.9[98] in the free-ratio model, assuming that the d_N/d_S values of all branches in the

phylogenetic tree are not equal. Mann-Whitney U tests were performed to identify OGs

with a significant difference in d_N/d_S between Parasitoida and Aculeata. To further test

d_N/d_S variation in Parasitoida and Aculeata, we used the CoV, calculated by dividing the

standard deviation by the mean of d_N/d_S in each OG in each group, and compared them

using a Mann-Whitney U test.

Protein domain analysis

We utilized HMMER 3.3.2 to annotate the protein domains in all species and calculated

the number of domain families classified based on PfamA accession. CAFE5[95] was used

to identify significantly accelerated domain families in the same way that gene families

were. The domain rearrangements were reconstructed using DomRates[99] with PfamScan

(https://github.com/aziele/pfam_scan) results as inputs.

GO annotation and enrichment analysis

We annotated proteins from all species using eggNOG-mapper v2[100] against the

eggNOG5 database. Based on the GO annotation results, we performed GO enrichment

analysis of the candidate gene set using GOATOOLS[101] with the go-basic.obo database

(http://geneontology.org/). The result of the *P* values for each GO term was corrected for

an FDR < 0.05 (Benjamini-Hochberg multi-test correction).

Molecular evolution analysis on functional categories

The package constellatoR (https://github.com/MetazoaPhylogenomicsLab/constellatoR)

was used to assign GO annotations to each OG and cluster GOs with similar functions

using the results of eggNOG-mapper v2 and OrthoFinder as inputs, thus classifying OGs

into different functional categories. For each OG in each category, the normalized

evolutionary rate was estimated by averaging the ratio of amino acid substitution rates of

interspecific homologous proteins to inter-species amino acid substitution rates. PAML

4.9[98] was used to calculate the d_N/d_S ratio for each OG using its respective phylogenetic

tree.

Whole genome alignment and constraint analysis

We used a 27-way whole genome alignment generated by the reference-free

aligner Cactus[102] and PHAST v1.6[103] programs (https://github.com/CshlSiepelLab/phast).

We utilized msa_view to retrieve 4-fold degenerate sites based on *B. terrestris* coding

gene annotation, and we fitted a phylogenetic model with phyloFit from PHAST v1.6. To

quantify the constraint of individual nucleotides, we utilized phyloP (part of the PHAST

v1.6 package) to compute the per-base constraint based on the resulting model.

Columns with a positive score were considered constrained bases, and those with a

negative score were accelerated-evolving bases. Similarly, we used phastCons[104] (part of

the PHAST v1.6 package) to find conserved elements. We excluded conserved elements

less than 20 bp in length. To further investigate the chromatin accessibility of conserved

no-coding elements, we downloaded raw ATAC-Seq data for the egg, larva, pupa, and

adult stages of *B. terrestris*[20] from NCBI (accessions in Supplementary Table 51).

Trimmomatic 0.33[105] was used to filter raw reads, removing low-quality reads and

adaptor sequences. Clean reads were mapped to the *B. terrestris* reference genome with

Bowtie2[106]. Then, we used Picard 3.1.1 (https://github.com/broadinstitute/picard) to

eliminate PCR duplicates and SAMtools 1.6[107] to delete mitochondrial reads. MACS2[108]

was utilized to identify peaks, and the irreproducible discovery rate (IDR) analysis was

performed to assess the reproducibility by comparing the consistency of two biological

replicates at the same developmental stage. Peaks with IDR <= 0.05 were reproducible

across two replicates and were kept for joint analysis with identified conserved no-coding

elements.

Metabolic pathway evolution

We used eggNOG-mapper v2[100] to perform EC annotations for all 131 hymenopterans,

and then ancestral pathways were constructed according to the intersections of

annotated ECs in eight sawflies (Tenthredinoidea) and the enzymes of each pathway in

KEGG. Pathway coverage was measured as the number of ECs in the pathway

annotated divided by the total number of ECs in the constructed ancestral pathway. We

considered pathways for which KEGG had a reference pathway for *A. rosae*, and the

reconstructed pathway contained at least five ECs. Coverage of each pathway between

the Parasitoida and Aculeata species was compared by a Mann-Whitney *U* test, and the

FDR-adjusted *P* of less than 0.05 was considered significant higher or lower. The CoVs

of pathway coverage in Parasitoida and Aculeata were calculated as the standard

deviation divided by the mean of pathway coverage in these two clades, respectively.

Functional convergence analysis

To investigate the level of functional convergence of interested OGs in three secondary

phytophagy transitional ancestral branches, the R package constellatoR was used to

calculate the similarity based on the semantic similarity matrix between OGs and cluster

them with an exemplar OG. We then used the plotConstellation function to visualize the

cluster shared by the interested OGs of the three transition nodes.

Transcriptome analysis

Raw transcriptome sequencing data for the second instar larva, third instar larva, fourth instar larva, female pupa, female adult, and venom gland of *A. fulloi* were downloaded from NCBI (accessions in Supplementary Table 52). Fastp 0.22.0[109] was used for filtration and quality control, and Salmon 0.14.1[110] was used to calculate the expression level of transcripts (measured as transcripts per million, TPM).

Acknowledgments

This work was supported by the Program of the National Natural Science Foundation of China (NSFC) (Grant No. 32202376 to XHY), the Key Program of NSFC (Grant No. 32330085 to GYY), the Young Elite Scientists Sponsorship Program by China Association for Science and Technology (Grant No. 2022QNRC001 to XHY), the Program of NSFC (Grant No. 32302408 to YY, No. 32302337 to XXZ, No. 32302444 to JLW), and the Scientific Research Startup Fund Project of Zhejiang A&F University (Grant No. 2023LFR119 to XHY). We are grateful to all colleagues who have contributed to the sequencing of Hymenoptera genomes.

Author contributions

Project conception: X.H.Y. Manuscript preparation: C.H., Y.Y., X.X.Z., X.H.Y., and G.Y.Y. Data collection and preparation: C.H., Y.Y., L.J.P., Y.Y.L., S.X., Y.M., X.K.G., and H.X. Phylogenetic analysis: C.H. and Y.Y. Comparative genomic analysis: C.H., X.H.Y., Y.Y., X.X.Z., J.J.L., and Y.T.C. Data interpretation: X.H.Y., C.H., Y.Y., S.J.X., J.L.W., Z.C.Y., Z.W.T., Q.F., and G.Y.Y. All authors read and approved the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Grimaldi, D. & Engel, M. S. *Evolution of the Insects*. (Cambridge University Press, 2005).

2.   Huber, J. T. Biodiversity of Hymenoptera. in *Insect Biodiversity: Science and Society*
(eds. Foottit, R. G. & Adler, P. H.) 303–323 (Wiley, 2009).

3.   Peters, R. S. et al. Evolutionary history of the Hymenoptera. *Curr. Biol.* **27**, 1013–
1018 (2017).

4.   Blaimer, B. B. et al. Key innovations and the diversification of Hymenoptera. *Nat.*
Commun. **14**, 1212 (2023).

5.   Sharkey, M. J. Phylogeny and classification of Hymenoptera. *Zootaxa* **1668**, 521–
548 (2007).

6.   Branstetter, M. G. et al. Phylogenomic insights into the evolution of stinging wasps
and the origins of ants and bees. *Curr. Biol.* **27**, 1019–1025 (2017).

7.   Tang, P. et al. Mitochondrial phylogenomics of the Hymenoptera. *Mol. Phylogenet.*
Evol. **131**, 8–18 (2019).

8.   Polaszek, A. & Vilhemsen, L. Biodiversity of hymenopteran parasitoids. *Curr. Opin.*
Insect Sci. **56**, 101026 (2023).

9.   Vilhelmsen, L., Mikó, I. & Krogmann, L. Beyond the wasp-waist: structural diversity
and phylogenetic significance of the mesosoma in apocritan wasps (Insecta:
Hymenoptera). *Zool. J. Linn. Soc.* **159**, 22–194 (2010).

10.   Mantica, F. et al. Evolution of tissue-specific expression of ancestral genes across
vertebrates and insects. *Nat. Ecol. Evol.* **8**, 1140–1153 (2024).

11.   Oeyen, J. P. et al. Sawfly genomes reveal evolutionary acquisitions that fostered the
mega-radiation of parasitoid and eusocial Hymenoptera. *Genome Biol. Evol.* **12**, 1099–
1188 (2020).

12.   Wissler, L., Gadau, J., Simola, D. F., Helmkampf, M. & Bornberg-Bauer, E.
Mechanisms and dynamics of orphan gene emergence in insect genomes. *Genome Biol.*
Evol. **5**, 439–455 (2013).

13.   Thomas, G. W. C. et al. Gene content evolution in the arthropods. *Genome Biol.* **21**,
15 (2020).

14.   Dou, X. Y., Zhang, A. J. & Jurenka, R. Functional identification of fatty acyl
reductases in female pheromone gland and tarsi of the corn earworm, *Helicoverpa zea*.
Insect Biochem. Mol. Biol. **116**, 103260 (2020).

15.   Carot-Sans, G., Muñoz, L., Piulachs, M. D., Guerrero, A. & Rosell, G. Identification
and characterization of a fatty acyl reductase from a *Spodoptera littoralis* female gland
involved in pheromone biosynthesis. *Insect Mol. Biol.* **24**, 82–92 (2015).

16.   Teerawanichpan, P., Robertson, A. J. & Qiu, X. A fatty acyl-CoA reductase highly
expressed in the head of honey bee (*Apis mellifera*) involves biosynthesis of a wide
range of aliphatic fatty alcohols. *Insect Biochem. Mol. Biol.* **40**, 641–649 (2010).

17.   Xiong, G. et al. Body shape and coloration of silkworm larvae are influenced by a
novel cuticular protein. *Genetics* **207**, 1053–1066 (2017).

18.   Pan, P. L. et al. A comprehensive omics analysis and functional survey of cuticular
proteins in the brown planthopper. *Proc. Natl. Acad. Sci. USA.* **115**, 5175–5180 (2018).

19.   Xu, H. X. et al. Comparative genomics sheds light on the convergent evolution of
miniaturized wasps. *Mol. Biol. Evol.* **38**, 5539–5554 (2021).

20.   Zhao, X. M., Su, L., Xu, W. L., Schaack, S. & Sun, C. Genome-wide identification of
accessible chromatin regions in bumblebee by ATAC-seq. *Sci. Data* **7**, 367 (2020).

21. Kapheim, K. M. et al. Genomic signatures of evolutionary transitions from solitary to
group living. *Science* **348**, 1139–1143 (2015).

22. Woodard, S. H. et al. Genes involved in convergent evolution of eusociality in bees.
Proc. Natl. Acad. Sci. USA **108**, 7472–7477 (2011).

23. Lacy, K. D. & Kronauer, D. J. C. Evolution: how sweat bees gained and lost
eusociality. *Curr. Biol.* **33**, R770–R773 (2023).

24. Mikhailova, A. A., Rinke, S. & Harrison, M. C. Genomic signatures of eusocial
evolution in insects. *Curr. Opin. Insect Sci.* **61**, 101136 (2024).

25. Jones, B. M. et al. Convergent and complementary selection shaped gains and
losses of eusociality in sweat bees. *Nat. Ecol. Evol.* **7**, 557–569 (2023).

26. Wyatt, C. D. R. et al. Social complexity, life-history and lineage influence the
molecular basis of castes in vespid wasps. *Nat. Commun.* **14**, 1046 (2023).

27. Vilhelmsen, L. Larval anatomy of Orussidae (Hymenoptera). *J. Hymenopt. Res.* **12**,
346–354 (2003).

28. Chen, Y. Y. et al. Female semiochemicals stimulate male courtship but dampen
female sexual receptivity. *Proc. Natl. Acad. Sci. USA* **120**, e2311166120.

29. Chertemps, T. et al. A female-biased expressed elongase involved in long-chain
hydrocarbon biosynthesis and courtship behavior in *Drosophila melanogaster. Proc. Natl.*
Acad. Sci. USA **104**, 4273–4278 (2007).

30. Rim, E. Y., Clevers, H. & Nusse, R. The Wnt pathway: from signaling mechanisms to
synthetic modulators. *Annu. Rev. Biochem.* **91**, 571–598 (2022).

31. Zhou, B. H. et al. Notch signaling pathway: architecture, disease, and therapeutics.
Signal Transduct. Target. Ther. **7**, 95 (2022).

32. Richards, E. H. Salivary secretions from the ectoparasitic wasp, *Eulophus*
pennicornis contain hydrolases, and kill host hemocytes by apoptosis. *Arch. Insect*
Biochem. Physiol. **79**, 61–74 (2012).

33. Shi, J. M. et al. The larval saliva of an endoparasitic wasp, *Pteromalus puparum*,
suppresses host immunity. *J. Insect Physiol.* **141**, 104425 (2022).

34. Pang, L. et al. Larval secretions of parasitoid wasps are new effectors that impair
host immune defences. *Crop Health* **1**, 11 (2023).

35. Peterson, C. L. & Laniel, M.-A. Histones and histone modifications. *Curr. Biol.* **14**,
R546–R551 (2004).

36. Martire, S. & Banaszynski, L. A. The roles of histone variants in fine-tuning chromatin
organization and function. *Nat. Rev. Mol. Cell Biol.* **21**, 522–541 (2020).

37. Kurata, S. Peptidoglycan recognition proteins in *Drosophila* immunity. *Dev. Comp.*
Immunol. **42**, (2014).

38. Mei, X. H. et al. Peptidoglycan recognition protein 6 (PGRP6) from Asian corn borer,
Ostrinia furnacalis (Guenée) serve as a pattern recognition receptor in innate immune
response. *Arch. Insect Biochem. Physiol.* **111**, e21955 (2022).

39. Xiao, D. Y. et al. The roles of SMYD4 in epigenetic regulation of cardiac development
in zebrafish. *PLoS Genet.* **14**, (2018).

40. International Helminth Genomes Consortium. Comparative genomics of the major
parasitic worms. *Nat. Genet.* **51**, 163–174 (2019).

41. Lu, T. M., Kanda, M., Furuya, H. & Satoh, N. Dicyemid mesozoans: a unique

parasitic lifestyle and a reduced genome. *Genome Biol. Evol.* **11**, 2232–2243 (2019).

42.   Ralton, J. E., Sernee, M. F. & McConville, M. J. Evolution and function of

carbohydrate reserve biosynthesis in parasitic protists. *Trends Parasitol.* **37**, 988–1001

(2021).

43.   Chen, X. L. et al. *Balanophora* genomes display massively convergent evolution with

other extreme holoparasites and provide novel insights into parasite–host interactions.

Nat. Plants **9**, 1627–1642 (2023).

44.   Asgari, S. & Rivers, D. B. Venom proteins from endoparasitoid wasps and their role

in host-parasite interactions. *Annu. Rev. Entomol.* **56**, 313–335 (2011).

45.   Moreau, S. & Asgari, S. Venom proteins from parasitoid wasps and their biological

functions. *Toxins (Basel)* **7**, 2385–2412 (2015).

46.   Teng, Z. W. et al. Protein discovery: combined transcriptomic and proteomic

analyses of venom from the endoparasitoid *Cotesia chilonis* (Hymenoptera: Braconidae).

Toxins (Basel) **9**, 135 (2017).

47.   Zhao, W. et al. Comparative transcriptome analysis of venom glands from *Cotesia*

vestalis and *Diadromus collaris*, two endoparasitoids of the host *Plutella xylostella. Sci.*

Rep. **7**, (2017).

48.   Goecks, J. et al. Integrative approach reveals composition of endoparasitoid wasp

venoms. *PLoS One* **8**, e64125 (2013).

49.   Burke, G. R. & Strand, M. R. Systematic analysis of a wasp parasitism arsenal. *Mol.*

Ecol. **23**, 890–901 (2014).

50.   Werren, J. H. et al. Functional and evolutionary insights from the genomes of three

parasitoid *Nasonia* species. *Science* **327**, 343–348 (2010).

51.   Dennis, A. B. et al. Functional insights from the GC-poor genomes of two aphid

parasitoids, *Aphidius ervi* and *Lysiphlebus fabarum. BMC Genom.* **21**, 376 (2020).

52.   Formesyn, E. M., Heyninck, K. & de Graaf, D. C. The role of serine- and

metalloproteases in *Nasonia vitripennis* venom in cell death related processes towards a

Spodoptera frugiperda Sf21 cell line. *J. Insect Physiol.* **59**, 795–803 (2013).

53.   Ye, X. H. et al. Comprehensive isoform-level analysis reveals the contribution of

alternative isoforms to venom evolution and repertoire diversity. *Genome Res.* **33**, 1554-

1567 (2023).

54.   Yang, L. et al. Identification and comparative analysis of venom proteins in a pupal

ectoparasitoid, *Pachycrepoideus vindemmiae. Front. Physiol.* **11**, 9 (2020).

55.   Lin, Z. et al. Insights into the venom protein components of *Microplitis mediator*, an

endoparasitoid wasp. *Insect Biochem. Mol. Biol.* **105**, 33–42 (2019).

56.   Parkinson, N., Conyers, C. & Smith, I. A venom protein from the endoparasitoid

wasp *Pimpla hypochondriaca* is similar to snake venom reprolysin-type metalloproteases.

J. Invertebr. Pathol. **79**, 129–131 (2002).

57.   Price, D. R. G. et al. A venom metalloproteinase from the parasitic wasp *Eulophus*

pennicornis is toxic towards its host, tomato moth (*Lacanobia oleracae*). *Insect Mol. Biol.*

18, 195–202 (2009).

58.   Dorémus, T. et al. Venom gland extract is not required for successful parasitism in

the polydnavirus-associated endoparasitoid *Hyposoter didymator* (Hym. Ichneumonidae)

despite the presence of numerous novel and conserved venom proteins. *Insect Biochem.*

Mol. Biol. **43**, 292–307 (2013).

59. Alvarado, G. et al. Bioinformatic analysis suggests potential mechanisms underlying parasitoid venom evolution and function. *Genomics* **112**, 1096–1104 (2020).

60. Vincent, B. et al. The venom composition of the parasitic wasp *Chelonus inanitus* resolved by combined expressed sequence tags analysis and proteomic approach. *BMC Genom.* **11**, 693 (2010).

61. Gatti, J.-L. et al. Proteo-trancriptomic analyses reveal a large expansion of metalloprotease-like proteins in atypical venom vesicles of the wasp *Meteorus pulchricornis* (Braconidae). *Toxins (Basel)* **13**, 502 (2021).

62. Yu, K. L. et al. Multi-omic identification of venom proteins collected from artificial hosts of a parasitoid wasp. *Toxins (Basel)* **15**, 377 (2023).

63. Lin, Z. et al. A metalloprotease homolog venom protein from a parasitoid wasp suppresses the Toll pathway in host hemocytes. *Front. Immunol.* **9**, 2301 (2018).

64. Yang, Y. et al. Genome of the pincer wasp *Gonatopus flavifemur* reveals unique venom evolution and a dual adaptation to parasitism and predation. *BMC Biol.* **19**, 145 (2021).

65. Colinet, D. et al. Extensive inter- and intraspecific venom variation in closely related parasites targeting the same host: the case of *Leptopilina* parasitoids of *Drosophila*. *Insect Biochem. Mol. Biol.* **43**, 601–611 (2013).

66. Crawford, A. M. et al. The constituents of *Microctonus sp.* parasitoid venoms. *Insect Mol. Biol.* **17**, 313–324 (2008).

67. Zhu, J. Y. Deciphering the main venom components of the ectoparasitic ant-like bethylid wasp, *Scleroderma guani*. *Toxicon* **113**, 32–40 (2016).

68. Liu, N. Y. et al. Venomics reveals novel ion transport peptide-likes (ITPLs) from the parasitoid wasp *Tetrastichus brontispae*. *Toxicon* **141**, 88–93 (2018).

69. Becchimanzi, A. et al. Venomics of the ectoparasitoid wasp *Bracon nigricans*. *BMC Genom.* **21**, 34 (2020).

70. Ye, X. H. et al. Genomic signatures associated with maintenance of genome stability and venom turnover in two parasitoid wasps. *Nat. Commun.* **13**, 6417 (2022).

71. Colinet, D. et al. Identification of the main venom protein components of *Aphidius ervi*, a parasitoid wasp of the aphid model *Acyrthosiphon pisum*. *BMC Genom.* **15**, 342 (2014).

72. Zhou, L. Z. et al. Two venom serpins from the parasitoid wasp *Microplitis mediator* inhibit the host prophenoloxidase activation and antimicrobial peptide synthesis. *Insect Biochem. Mol. Biol.* **152**, 103895 (2023).

73. Yan, Z. C. et al. A serpin gene from a parasitoid wasp disrupts host immunity and exhibits adaptive alternative splicing. *PLoS Pathog.* **19**, e1011649 (2023).

74. Zhao, X. X. et al. Genomic signatures associated with the evolutionary loss of egg yolk in parasitoid wasps. Preprint at https://doi.org/10.1101/2023.12.30.573744 (2024).

75. Ralec, A. Egg contents in relation to host-feeding in some parasitic hymenoptera. *Entomophaga* **40**, 87–93 (1995).

76. Grbic, M. & Strand, M. R. Shifts in the life history of parasitic wasps correlate with pronounced alterations in early development. *Proc. Natl. Acad. Sci. USA* **95**, 1097–1101 (1998).

77. Donnell, D. M. Vitellogenin of the parasitoid wasp, *Encarsia formosa* (Hymenoptera: Aphelinidae): gene organization and differential use by members of the genus. *Insect Biochem. Mol. Biol.* **34**, 951–961 (2004).

78. Hegedus, D., Erlandson, M., Gillott, C. & Toprak, U. New insights into peritrophic matrix synthesis, architecture, and function. *Annu. Rev. Entomol.* **54**, 285–302 (2009).

79. Zhu, L. F. et al. Adaptive evolution to a high purine and fat diet of carnivorans revealed by gut microbiomes and host genomes. *Environ. Microbiol.* **20**, 1711–1722 (2018).

80. Gloss, A. D. et al. Evolution in an ancient detoxification pathway is coupled with a transition to herbivory in the Drosophilidae. *Mol. Biol. Evol.* **31**, 2441 (2014).

81. Gloss, A. D. et al. Evolution of herbivory remodels a *Drosophila* genome. Preprint at https://doi.org/10.1101/767160 (2019).

82. Winde, I. & Wittstock, U. Insect herbivore counteradaptations to the plant glucosinolate–myrosinase system. *Phytochemistry* **72**, 1566–1575 (2011).

83. Lv, Q. Q., Li, X. F., Fan, B. F., Zhu, C. & Chen, Z. X. The cellular and subcellular organization of the glucosinolate–myrosinase system against herbivores and pathogens. *Int. J. Mol. Sci.* **23**, 1577 (2022).

84. Abdalsamee, M. K., Giampà, M., Niehaus, K. & Müller, C. Rapid incorporation of glucosinolates as a strategy used by a herbivore to prevent activation by myrosinases. *Insect Biochem. Mol. Biol.* **52**, 115–123 (2014).

85. Müller, C. Interactions between glucosinolate- and myrosinase-containing plants and the sawfly *Athalia rosae*. *Phytochem. Rev.* **8**, 121–134 (2009).

86. Bridges, M. et al. Spatial organization of the glucosinolate-myrosinase system in brassica specialist aphids is similar to that of the host plant. *Proc. Biol. Sci.* **269**, 187–191 (2002).

87. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

88. Manni, M., Berkeley, M. R., Seppey, M. & Zdobnov, E. M. BUSCO: assessing genomic data quality and beyond. *Curr. Protoc.* **1**, e323 (2021).

89. Derelle, R., Philippe, H. & Colbourne, J. K. Broccoli: combining phylogenetic and network analyses for orthology assignment. *Mol. Biol. Evol.* **37**, 3389–3396 (2020).

90. Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

91. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

92. Minh, B. Q. et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

93. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., von Haeseler, A. & Jermin, L. S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589 (2017).

94. Sanderson, M. J. r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. *Bioinformatics* **19**, 301–302 (2003).

95. Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models variation in

evolutionary rates among gene families. *Bioinformatics* **36**, 5516–5518 (2021).

96. Pond, S. L. K., Frost, S. D. W. & Muse, S. V. HyPhy: hypothesis testing using

phylogenies. *Bioinformatics* **21**, 676–679 (2005).

97. Smith, M. D. et al. Less is more: an adaptive branch-site random effects model for

efficient detection of episodic diversifying selection. *Mol. Biol. Evol.* **32**, 1342–1353

(2015).

98. Yang, Z. H. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.*

24, 1586–1591 (2007).

99. Dohmen, E., Klasberg, S., Bornberg-Bauer, E., Perrey, S. & Kemena, C. The

modular nature of protein evolution: domain rearrangement rates across eukaryotic life.

BMC Evol. Biol. **20**, 30 (2020).

100. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: functional annotation, orthology assignments, and domain

prediction at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

101. Klopfenstein, D. V. et al. GOATOOLS: a python library for gene ontology analyses.

Sci. Rep. **8**, 10872 (2018).

102. Armstrong, J. et al. Progressive Cactus is a multiple-genome aligner for the

thousand-genome era. *Nature* **587**, 246–251 (2020).

103. Hubisz, M. J., Pollard, K. S. & Siepel, A. PHAST and RPHAST: phylogenetic

analysis with space/time models. *Brief. Bioinform.* **12**, 41–51 (2011).

104. Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and

yeast genomes. *Genome Res.* **15**, 1034–1050 (2005).

105. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

106. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*

Methods **9**, 357–359 (2012).

107. Li, H. et al. The sequence alignment/map format and SAMtools. *Bioinformatics* **25**,

2078–2079 (2009).

108. Gaspar, J. M. Improved peak-calling with MACS2. Preprint at

https://doi.org/10.1101/496521 (2018).

109. Chen, S. F., Zhou, Y. Q., Chen, Y. R. & Gu, J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

110.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides

fast and bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419

(2017).

Figures

Fig. 1 | Phylogenetic analysis and genomic comparisons of Hymenoptera. The

maximum-likelihood phylogenetic tree on the left was constructed with 1000 ultrafast

bootstraps under a JTT+F+R10 model using protein sequences from 1002 single-copy

genes. Almost all the nodes received 100% bootstrap support, except for nine. The

branch color corresponds to the rate of substitutions per site per 100 million years along

that particular branch. The divergence times among species were estimated using 13

calibration points. Several key innovations, such as parasitoidism, wasp-waist, and

stinger, are marked on the phylogeny, and three independent transitions to secondary

phytophagy are indicated by red arrows. On the right side, the figure shows i) genome

assembly sizes of hymenopteran species; ii) BUSCO profiles for these genomes; and iii)

orthology profiles. In the orthology analysis, the protein-coding genes of each species

were subdivided to represent different types of orthology clusters.

Fig. 2 | Gene family evolution and rapidly evolving genes in Hymenoptera. A) Gene

family expansion and contraction at the nodes of key innovations in Hymenoptera

evolution. In each box, the first row shows the number of gene families that expanded or

contracted, respectively. Below are the corresponding representative gene families, and

the size changes of each gene family are also indicated. **B**) Rapidly evolving OGs and

functional categories in terms of evolutionary rate and selection constraint (d_N/d_S).

Functional categories (GO Biological Process terms) are sorted by normalized

evolutionary rates, and the top 25 GO terms are shown. The normalized evolutionary rate

of each OG is measured as the mean of the amino acid substitution rates of each OG

normalized by the genome-wide amino acid substitution rates. OGs were further grouped

according to their functional similarity. d_N/d_S of each OG refers to the ratio of the non-

synonymous substitution rate to the synonymous substitution rate, measured by the one-

ratio model in codeml.

1005

Fig. 3 | Different evolutionary genome dynamics between Parasitoida and Aculeata.

A) The distribution of the rapidly evolving events for gene families on the branches of the

Parasitoida and Aculeata clades. Three gene families, Histone H3, Trypsin, and UDP-

glucosyltransferase, are shown here, which have a significantly higher occurrence rate of

rapid evolutionary events in Parasitoida than in Aculeata (FDR-adjusted P < 0.05, Chi-

square test, odds ratio >1). **B**) Gene families that have a higher rate of gene gain and

loss in Parasitoida than in Aculeata. The gene gain and loss rate is measured as the sum

of gene gains and losses per million years. Asterisks indicate significant shifts between

Parasitoida and Aculeata (One-tailed Mann-Whitney U test; *≤05; **≤0.01; ***≤0.001,

****≤0.0001). **C**) GO enrichment analysis of the gene families with a faster rate of gene

gain and loss in Parasitoida than in Aculeata. The gray box shows the GO terms for the

10 biological processes with the lowest adjusted P value (FDR-adjusted P < 0.05,

Benjamini-Hochberg multi-test correction). The size of the blue circles represents the

number of genes in the corresponding GO. **D**) KEGG pathway coverage comparison

between Parasitoida and Aculeata. The expanded lines from the point represent the first

and third quartiles of pathway coverage in Parasitoida and Aculeata, respectively.

1022

Fig. 4 | Expansions of Venom allergen and Venom metalloprotease gene families in

Parasitoida. A) The size distribution of the Venom metalloprotease gene families in

Hymenoptera. The number on the bar indicates the maximum gene copy count across

the species for that family. **B**) Expression pattern of the Venom metalloprotease genes in

Anastatus fulloi at five developmental stages and the venom gland. Expression is

measured using z-score values of transcripts per million (TPM) by row. L2, second instar

larva; L3, third instar larva; L4, fourth instar larva; YP, pupal stage; AD, female adult; VG,

venom gland. Only genes with a TPM greater than 0 in at least one period are shown. **C**)

The size distribution of the Venom allergen gene families in Hymenoptera. **D**) Expression

patterns of the Venom allergen genes in *A. fulloi* at five developmental stages and the

venom gland.

A

B

1034

Fig. 5 | Genomic features of three independent origins of secondary phytophagy.

A) UpSet charts showing the number of shared and unique expanded, contracted, and

positively selected gene families among three clades of independent transitions to

secondary phytophagy: Cynipidae, Agaonidae, and Anthophila, with intersection size

representing the number of shared OGs in each category. No expanded or contracted

OGs, or positively selected genes, were shared by all three groups. **B**) Functional

convergence of exclusively positively selected genes in the three branches of secondary

phytophagy origins. Each dot represents an OG to which the exclusively positively

selected gene belongs. The centroid OG of each polygon is labeled with potential

functions. Of the OGs with exclusively positively selected genes, 67.28% were clustered

into 19 functional clusters, indicating functional convergence signals.