Intra- and interspecific variations in flight performance of oak-associated Agrilinae (Coleoptera: Buprestidae) using computerised flight mills

Le Souchu Elodie[1,3], Sallé Aurélien[1], Bankhead-Dronnet Stéphanie[1], Laparie Mathieu[2]*, Sauvard Daniel[2]*

[1] University of Orléans, Laboratory of Physiology, Ecology and Environment (P2E), INRAE USC 1328, Orléans, France
[2] INRAE, URZF (Forest Zoology Research Unit), Orléans, France
* These authors contributed equally to the work.

[3] Corresponding author: Le Souchu Elodie
Correspondence: elodie.lesouchu2@gmail.com
Present address: Swedish University of Agricultural Sciences (SLU), Department of Ecology - Inst för Ekologi, Box 7044, 750 07 Uppsala, Sweden

ABSTRACT

Several Agrilinae species (Coleoptera: Buprestidae) are secondary pests of broadleaf forests, and some of them are also major invasive pests. These thermophilous borers are expected to be favoured by climate change and the global deterioration of forest health, and ultimately expand their range and damage. Flight behaviour and performance of these insects are poorly known despite their critical role in dispersal inside and outside native ranges and their relevance for management purposes. This study aimed to assess intra- and interspecific variability in active flight of several Agrilinae species and effects of sex and mass on this variability to contribute to filling this gap. We assessed the flight performance of eleven species associated with oaks (nine *Agrilus*, one *Coraebus* and one *Meliboeus*) plus one *Agrilus* species associated with the herbaceous layer. Computer-monitored flight mills were used to measure flight parameters, including periods, durations, distances and velocity in 250 beetles. Overall, flight capacities were rather homogeneous among species, with a dominance of poor flyers and only *Coraebus undatus* showed outstanding performance. Beetles generally performed several short flight bouts within one trial, and only a few individuals sustained long flight. The maximal total distance covered across multiple assays until death ranged from 170 to 16 097 m depending on the species, with a median between 35 and 966 m (excluding individuals that never flew). On top of this interspecific variability, flight distances also varied greatly among individuals, but were not influenced by sex. Preflight body mass had mixed effects depending on the species, presumably related to different dispersal patterns. In our experimental conditions, most species had limited average dispersal capacities over multiple flight trials. Overall, dispersal over long distances and colonisation events probably depend mainly on a small proportion of individuals which largely exceeded the median performance.
Keywords: *Agrilus*, secondary pest, range expansion, tethered flight, dispersal behaviour, locomotor activity

                                          Introduction

The jewel beetle subfamily Agrilinae (Coleoptera: Buprestidae) includes numerous
species, especially in the genus *Agrilus* which gathers more than 3 300 described species
worldwide (Jendek & Grebennikov, 2023). Some of these thermophilous and heliophilous
insects (Reed et al., 2018) are known to be phytosanitary threats on a wide diversity of
woody and some non-woody plants (Jendek & Poláková, 2014) and can have important
economic and ecological impacts in their native area. Larvae develop in the roots, stems,
trunks or branches of their host plant. Within trees, larval development could typically last
several years. Imaginal moult takes place in spring, within host tissues. After emergence, they
usually move towards the foliage to perform a maturation feed and mate. Then females
search for a suitable host to lay their eggs (Bílý, 2002).
As opportunistic pests, Agrilinae are associated with forest declines and diebacks (Le
Souchu et al., 2024). In temperate forests, wood-borers such as *Agrilus biguttatus* (Fabricius)
and *Agrilus viridis* (Linnaeus) are aggravating agents of large scale oak and beech forest
declines (Sallé et al., 2014; Brown et al., 2015; Brück-Dyckhoff et al., 2019; Macháčová et al.,
2022). A related Agrilinae species, *Coraebus undatus* (Fabricius) has also been associated
with oak decline (Du Merle & Attié, 1992; Sallé et al., 2014). *Agrilus* species are also known
for their high invasive potential, supported by several examples of human-mediated
invasions (e.g., Wu et al., 2017; Hızal & Arslangündoğdu, 2018; Bozorov et al., 2019). Indeed,
their long and cryptic pre-imaginal life in the wood facilitates unnoticed transportations and
introductions (Chamorro et al., 2015; Ruzzier et al., 2023). One of the prominent invasive
pests worldwide is the Emerald Ash Borer (EAB), *Agrilus planipennis* Fairmaire. It is
responsible for most of the worldwide economic damage attributed to Buprestidae (Renault
et al., 2022). This species has invaded North America from Asia (Herms & McCullough, 2014)
and arrived in Russia in 2003, now threatening the western part of Europe (Orlova-
Bienkowskaja et al., 2020). Other *Agrilus* species have been identified as potential future
invaders, such as *Agrilus anxius* (Evans et al., 2020) or *Agrilus fleischeri* Obenberger
(Volkovitsh et al., 2020), including species belonging to the European assemblage of oak-
associated borers. Current forest decline phenomena are beneficial to these borers (Le
Souchu et al., 2024), together with climate change which is suspected to assist their
northward expansion (Pederson & Jørum, 2009; Brown et al., 2015). Within this context,
evaluating flight capacities of intercepted or introduced species is of paramount importance
to evaluate their dispersal capacities and implement efficient monitoring and control
strategies.
Active flight is a central function in the life strategies of most insects (Dickinson et al.,
1999; Dudley, 2002). This mode of locomotion is involved in the search for a mate, trophic
resources or favourable environmental conditions, as well as in dispersal and colonisation of
new geographic areas (Dudley, 2002). The distance covered during the lifetime of an
individual can vary considerably among genera, species, and even individuals. It can be
influenced by numerous factors, ranging from intrinsic features (physiological traits,
reserves, morphological characteristics and behaviours, mated status, *etc.*), to external
(a)biotic factors (temperature, crowding, food availability, *etc.*) and their spatiotemporal
variations, immediate or not (Minter et al., 2018). The study of flight capacities and their
variations at multiple focal scales can provide valuable insights to better understand the
ecology of species and, in particular, help to estimate their dispersal potential (see e.g.,
Robinet et al., 2012). Yet, in the case of *Agrilus* spp., despite their economic and ecological
damage and their predisposition to be accidentally introduced into new environments, this

information remains scarce in the literature. Their flight capacities have rarely been tested,
let alone in European species. Therefore, the numerous works on *A. planipennis* are used to
estimate the expected dispersal capacities of potential invasive *Agrilus* spp*.* or introduced
ones (e.g., Baranchikov et al., 2019a; b) and highlight their high flight capacities which may
facilitate further dispersal (Taylor et al., 2010; USDA-APHIS, 2011). While male *Agrilus* may
be more prone to staying close to their birth host to look for a mate (Rodriguez-Soana et al.,
2007), gravid females are able to disperse several kilometres away (Taylor et al., 2010).
However, it is often considered that most females will disperse only tens of meters if the
resource distribution allows it (Mercader et al., 2009).
Several methods exist to estimate insect flight capacities. Laboratory approaches are
usually favoured (Minter et al., 2018; Naranjo, 2019) due to the short lifespan of insects and
the difficulty to track them in the field visually or using mark-recapture protocols; in addition,
most insects, including *Agrilus* spp., are too tiny to carry even the smallest tags. These
approaches use either free flight protocols (or free flight chamber, Taylor et al., 2010; e.g.,
wind tunnel, Crall et al., 2017) or tethered techniques (e.g., flight mill, Sauvard et al., 2018).
The latter are usually preferred, especially flight mills, which are rather inexpensive, produce
a wealth of data, have no limit on flying distance, and are easy to replicate under controlled
conditions. As insects tethered on a flight mill never reach any target, flight mills provide
information on flight capacities rather than actual flight dynamics and behaviours *in natura*.
Their advantages are therefore offset by the fact that they create artificial conditions, with
multiple potential biases. However, their easy replication allows comparing individuals and
species in the exact same conditions, and many authors showed that tethered flight results
are consistent with general knowledge on the studied insect performances, or even actual
field flight capacity when it is available (Robinet et al., 2012, on processionary moth; Sauvard
et al., 2018, on yellow-legged hornet).
In this study, we aimed at determining flight capacities of several *Agrilus* species, and
specifically their intra- and interspecific variations. The focus on *Agrilus* behaviour and
biology may help establish management strategies and improve detection tools. We
measured flight potential of oak-associated species using wild individuals tethered to
computer-monitored flight mills. An herbaceous species, *Agrilus hyperici* (Creutzer), similar
in size to most of the oak-associated species we sampled, was also added to the pool of
species. This species was intentionally introduced in North America from France in 1953 as
a biological control agent to target *Hypericum perforatum* L. and its dispersal have been
monitored (Jendek & Grebennikov, 2009; Gov. of British Columbia, 2018). This addition
allowed to (1) confirm that the performance of *A. hyperici* deals with the knowledge on its
dispersion *in natura*, and (2) compare flight performance of forest species to that of an
herbaceous-associated species. First, we expected noticeable flight differences among
species, as previously observed within multiple coleopteran families (Nitidulidae, Van Dam
et al., 2000; Cetoniinae, Dubois et al., 2009). Second, we hypothesised high inter-individual
variations in flight measurements, especially a strong sexual dimorphism with females
putatively flying greater distances than males, as is the case for mated females of
A. planipennis (Taylor et al., 2010).

                                Material and methods

Sampling and rearing design

We conducted three consecutive years of experiments, from 2021 to 2023, in France. Live
Agrilus were caught using two methods. First, we used green multi-funnel traps (ChemTica
Internacional, San Jose, Costa Rica) with twelve fluon-coated funnels. These traps are
commonly used to sample Agrilinae (Rassati et al., 2019; Le Souchu et al., 2024). From June
23 to August 06 in 2021 and from May 09 to July 27 in 2022, respectively six and eight traps
were set at the University of Orléans (Longitude: 47.85 N, Latitude: 1.94 E; WGS 84, DDD),
and four and six traps at the INRAE site of Ardon (47.83 N, 1.91 E). These two sites are
separated by 2.6 km. Selected forest patches in these areas were dominated by *Quercus robur*
(L.) and *Quercus petraea* (Matt.) Liebl. A fresh oak leaf was placed in each collector to provide
food and support to trapped insects, and no preservative liquid was used in order to keep
them alive. Traps were hung among the lower branches in the canopy (i.e., approximately 15
m above the ground), to capture adults after their emergence, during their search for a mate
or a suitable host. No lure was added to the traps and the collectors were emptied every
morning during weekdays. Additionally, we captured *A. hyperici* on *H. perforatum* and oak
borers on oak logs using a sweep net. In 2022, *A. hyperici* were sampled in three locations, at
Saint-Denis-en-Val (47.89 N, 1.98 E) on June 26 and 28, at the University of Orléans on July
04, and at the INRAE site of Ardon on July 06. In 2023, we similarly captured *A. hyperici* in
Vouzon (47.64 N, 2.05 E) on June 16. The same day, several specimens of oak borers were
captured in a log deposit nearby (47.65 N, 2.01 E). All combined, the oak-associated species
caught were: *Agrilus angustulus* (Illiger)*, Agrilus graminis* Kiesenwetter*, Agrilus hastulifer*
(Ratzeburg)*, Agrilus laticornis* (Illiger)*, Agrilus sulcicollis* Lacordaire*, C. undatus, Meliboeus*
fulgidicollis (P.H. Lucas); and species specialised in others hosts (indicated between
parentheses) were: *Agrilus graecus* Obenberger (*Viscum* spp.)*, Agrilus olivicolor*
Kiesenwetter (*Carpinus* spp.)*, A. viridis* (*Fagus* spp.) and *A. hyperici* (*H. perforatum*). All
individuals are listed in the supplementary table S1.
Live Agrilinae collected were maintained in the INRAE site of Ardon, in a climatic chamber
(Galli GTEST-CL-0700) at a 12:8-hour cycle of 25:20 °C $\pm$ 0.2 °C with 2-hour gradients
between both phases, 16:8-hour L:D photoperiod using three LED spots (Lédis, 3.5 W, 300
lumens, 4500 K), and 60 % $\pm$ 5 % relative humidity (warm phase and photophase were
centred on solar midday, 2 pm local time). Each insect was put in an individual plastic vial
capped with wet cotton to allow ventilation and provide a water source. Since species were
determined only after death, all vials but those containing *A. hyperici* (which were known in
advance due to the different sampling method) were supplied with a fresh oak leaf renewed
every two days, whereas vials containing *A. hyperici* were supplied with fresh *H. perforatum*
leaves*.

Flight mill design

Rotational flight mills were designed to estimate beetle flight capacities. Each flight mill
(Figure 1) was built on a wooden square base (19 × 19 cm), with an upwards wooden stand
in the centre and a wooden bracket on one corner, extending up to the centre. A magnet
(Supermagnete, https://www.supermagnete.de/, S-05-05-N) was placed under the bracket
and another one on the stand, both vertically aligned to hold an entomological pin (size 2).
To minimise friction, the pinhead had no contact with the lower magnet, and only the point
of the pin was in contact with the upper magnet. The pin supported a horizontal 16-cm arm,

composed of two identical 1-mm-diameter carbon rods connected by a 2-mm-diameter 2-cm aluminium tube; a hole was drilled in the middle of the tube, through which the pin was inserted and glued. The arm carried a small magnet (Supermagnete, S-03-02-N) glued on the lower side, near one end of the tube. The insect was placed at the end of the arm opposite to this magnet and drove the arm while flying. For heavy species (*C. undatus*), a counterweight made of reusable adhesive paste was also placed on the arm end opposite to the insect.

A complete lap corresponded to a travel distance of 0.5 metre. Arm moves were tracked using two Hall effect sensors (DiodesZetex AH337-PG-B) able to detect magnetic fields. They were positioned on the stand just below the arm, approximately 60° from each other around the pin, and detected every small magnet transit over them. Each Hall sensor was connected to a single board computer (Raspberry Pi), to which signals were transmitted. A total of 26 flight mills were built, and managed using 4 Raspberry Pi (1 Raspberry Pi B+, managing 2 flight mills; 1 Raspberry Pi 2 B, 1 Raspberry Pi 3 B and 1 Raspberry Pi 3 B+, each managing 8 flight mills).

An in-house Crystal program running on these computers managed the signals. The program initialised each flight test with flight user-set metadata (mill identifier, insect identifier and gender if known in advance), and start time, creating two dedicated CSV files (one per Hall sensor), hereafter referred to as log files. Then, each associated signal detection was time-stamped and recorded in the appropriate file. At the end of the test, end time was also recorded and the log files were closed.

Flight mills were installed in climate chambers (three PHCBi MIR-254-PE, with six flight mills each, and two Galli GTEST-TM-0700, with four flight mills each; 25 °C $\pm$ 0.5 °C, 50 % $\pm$ 10 % relative humidity). Each chamber shelf carried a pair of flight mills, which were lit by two LED spots (same as in the rearing chamber). In order to stimulate flight, fresh oak branches with leaves (or *H. perforatum* branches when *A. hyperici* was tested) were placed on each shelf, between the flight mills.

Figure 1. Flight mill used for this study, with a close-up view of the sensor and axis part. The electronics of the system are not shown.

Experimental design

As Agrilinae are diurnal and thermophilous insects (Reed et al., 2018), we tested them for
flight capacities during daytime and at 25 °C $\pm$ 0.5 °C. After its sampling, each beetle was
maintained one day in rearing conditions (initial rest). The first flight test occurred the next
working day; it lasted eight hours. The beetle was then tested again on the flight mill each
working day until its death, but these flight tests lasted only four hours. However, test
frequency was lowered to a test per week after the tenth one. Test time was chosen so that
insects were extracted from the rearing chamber at least an hour after the beginning of the
warm phase of their rearing thermal cycle; first tests started at approximately 10 am, and
next tests at about 2 pm. In 2021, due to initial low availability of flight mills, only the first
test was done with some beetles; as two of them remained alive three weeks after, they had
then been tested (during 4 hours) each working day until their death.
Beetles were weighed on a microbalance (Mettler Toledo AG204, d = 0.1 mg) before each
flight test. They were then placed in a 3 °C walk-in chamber to slow them down and ease
glueing a strip of paper narrower than their width on the dorsal side of their pronotum using
cyanoacrylate glue, the most common glue used to tether insects (Naranjo, 2019). The length
and width of strips were adjusted to the dimensions of insects. This operation in cold
conditions lasted from 15 to 30 minutes, then insects were returned to ambient temperature
and immediately attached to flight mills with reusable adhesive paste between the mill's arm
and the strip of paper. This setup allowed rapidly installing and removing tested insects at
the beginning and the end of experiments. At the end of flight tests, paper strips were
delicately removed from insects, as well as any visible residue of glue, and insects were
weighed again before being returned to their rearing vial and chamber. However, no clear
variation of the fresh mass was observed during a flight mill trial or during beetle life and
only the initial preflight mass was used.

Data analysis

In-house *Ruby* scripts were developed to analyse the log files generated by the *Crystal*
program used to monitor flight mills. They gathered data from the two log files generated by
a flight mill during a test, separated flight phases (identified using unequal alternance of
signals from the two Hall sensors) from noise and oscillations, and recorded their timestamp,
duration and length in the raw dataset used in further analyses. The flight phases were
hereafter referred to as flight bouts.
We focused our study on several measures of flight capacities, traditionally used to assess
flight performance, which included total flight distance (m), mean flight bout distance (m),
total flight duration (s), mean flight bout duration (s), number of flight bouts and latency to
first flight bout (s). The latency to first flight bout was used as a measure of the propensity to
fly. However, several measures were correlated and we chose to not include the mean flight
bout duration and the total flight duration in analyses. The selection method with the
correlation tests are available as supplementary material (Document S2). We also explored
the relationship between the fresh body mass before the first flight mill trial, hereafter
referred to as preflight mass, with the performance. For statistical reasons, species with
fewer than six individuals in total were removed from all per-species analyses or figures, and
only included in approaches where all species were pooled together. Some flight tests
experienced technical problems, such as insect escape or electronic issues, and were
excluded from analyses. As indicated above, test frequency changed after the tenth test, and,
in 2021, two beetles completed their tests, except the first one, after a long rest period; the

corresponding data, hereafter referred to as unusual trial frequency data, was also often excluded from analyses.

Standard linear model (LM) was used to assess linear regression. Kruskal-Wallis tests were used to assess the difference in flight capacities between groups. The Benjamini-Hochberg method permitted to correct p-values for False Discovery Rate. Kruskal-Wallis tests were followed by a post-hoc Dunn test with the Holm adjustment method.

LMERs were used to evaluate the effect of sex and preflight body mass on response variables associated with flight performance of the three most abundant species (*A. hyperici, A. laticornis* and *A. olivicolor*). Every response variable was log-transformed to improve the models, and Q-Q plots were used to review them. Since data from all three years were analysed together, the year of experiment (i.e., 2020, 2021, 2022) was included as a random effect. For each response variable, the null model was compared to the model with preflight mass and sex as explanatory variables, based on their respective AICc (Akaike Information Criterion scores) with a correction for small sample sizes.

When it was needed to compare data from the first eight-hour trial with the subsequent four-hour trials, we truncated the former one to four hours. This resulted in only limited data discarded, as on average 83 % of total distance of eight-hour trials is flown during the first four hours. Unusual trial frequency data was discarded from analyses.

All analyses and figures were realised using *R* 4.2 and *R* 4.3 (R Core Team, 2022, 2023).

Results

General beetle characteristics

A total of 250 beetles were tested during the three-year experiment, with 102 individuals in 2021, 125 in 2022, and 23 in 2023 (Table 1). Twelve species were collected, among which three dominated the samples: *A. laticornis*, *A. hyperici* and *A. olivicolor*. Sex ratio was found to be relatively balanced in the three most numerous species, and often biased towards fewer males than females in rarer species, but their small sample sizes warrants caution on this result.

The overall lifespan after capture was low and highly variable (Figure S3). The median was only 4 days, with a quarter of beetles living more than 6 days, 5 % (15 beetles) more than 13 days and 1 % (4 beetles) between 38 and 56 days.

Preflight mass differed significantly among species (Kruskal-Wallis, $\chi^2(7) = 67.4$, $p < 0.001$, n = 208). According to Dunn's test, *C. undatus*, *A. hastulifer* and *A. graminis* were significantly heavier than the other species tested (Figure 2). *Coraebus undatus* was clearly out of category in terms of size and mass.

Species	2021 Females		2021 Males		2022 Females		2022 Males		2023 Females		2023 Males		Total	
Agrilus angustulus	1	(0)			1	(1)	2	(2)	5	(5)	2	(2)	11	(10)
Agrilus graecus *			1	(0)									1	(0)
Agrilus graminis	2	(2)	3	(3)	3	(3)	1	(1)					9	(9)
Agrilus hastulifer	7	(6)	3	(3)	2	(2)							12	(11)
Agrilus hyperici					14	(14)	13	(11)	8	(8)	7	(7)	42	(40)
Agrilus laticornis	29	(25)	21	(19)	31	(26)	41	(35)	1	(1)			123	(106)
Agrilus obscuricollis *	2	(2)	1	(1)	2	(2)							5	(5)
Agrilus olivicolor	8	(6)	8	(6)	5	(5)	1	(1)					22	(18)
Agrilus sulcicollis *			1	(1)	1	(1)	2	(2)					4	(4)
Agrilus viridis *	3	(3)	2	(1)									5	(4)
Coraebus undatus	4	(4)			2	(2)	1	(1)					7	(7)
Meliboeus fulgidicollis	6	(5)			1	(0)	2	(2)					9	(7)

295
296 **Table 2.** Median, mean and maximum total distance flown by the insects during their lifespan. First trial and Sum correspond to the data of the first eight-hour trial and the sum of all trials respectively.

| Species | Median total flight distance (m) | | | | Mean total flight distance (m) | | | | Highest total flight distance (m) | |
| | With non-flying individuals | | Without non-flying individuals | | With non-flying individuals | | Without non-flying individuals | | | |
	First trial	Sum	First trial	Sum	First trial	Sum	First trial	Sum	First trial	Sum
All with *Coraebus undatus*	40	54	95	119	200 ± 28	356 ± 74	290 ± 38	487 ± 100	3 028	16 097
All without *Coraebus undatus*	35	53	87	115	167 ± 22	274 ± 38	245 ± 31	379 ± 51	2 246	4 636
Agrilus angustulus	70	71	71	92	186 ± 100	205 ± 92	207 ± 109	225 ± 99	1 048	1 048
Agrilus graecus	738	738	738	738	738	738	738	738	738	738
Agrilus graminis	74	149	87	342	98 ± 39	663 ± 500	147 ± 47	995 ± 730	350	4 636
Agrilus hastulifer	143	182	212	212	245 ± 90	486 ± 189	300 ± 101	531 ± 202	910	2 135
Agrilus hyperici	25	56	316	460	341 ± 91	671 ± 152	546 ± 130	916 ± 189	2 246	2 989
Agrilus laticornis	35	53	67	95	121 ± 21	144 ± 21	169 ± 27	195 ± 26	1 132	1 132
Agrilus obscuricollis	0	0	144	144	58 ± 41	58 ± 41	144 ± 64	144 ± 64	208	208
Agrilus olivicolor	1	1	27	35	26 ± 11	28 ± 10	53 ± 19	52 ± 15	170	170
Agrilus sulcicollis	68	68	68	68	141 ± 96	141 ± 96	141 ± 96	141 ± 96	426	426
Agrilus viridis	364	782	428	894	367 ± 153	771 ± 277	489 ± 130	964 ± 256	739	1 644
Coraebus undatus	966	966	1 070	966	1 205 ± 410	3172 ± 2 168	1 405 ± 424	3172 ± 2 168	3 028	16 097
Meliboeus fulgidicollis	0	0	61	61	72 ± 60	79 ± 69	168 ± 131	211 ± 174	429	559

General flight characteristics

First analyses were performed on first (eight-hour) trial data. All species and sexes
combined, 222 individuals (123 females, 99 males) have been analysed, taking into account
the removal of individuals for which some issue occurred during the trial. Main results are
displayed in figure 2.

Figure 2. Values for a selection of response variables measured during the first (eight-
hour) trial, per sex and species. Females: left (green), males: right (purple).

Regardless of the species, flight distance flown during the first trial was highly variable:
70 insects (31 %) did not fly, 44 % (98 beetles) flew between 1 m and 211 m, 25 %
(56 beetles) flew more than 211 m, 10 % (23 beetles) more than 596 m, 5 % (12 beetles)
more than 948 m, and 1 % (3 beetles) more than 2 138 m (Figure 3).The longest total
distance flown within an eight-hour trial was 3 028 m for a female *C. undatus*, 2 246 m for a
female *A. hyperici*, and 429 m for a male *M. fulgidicollis.* Remarkably, some individuals of
C. undatus (two females) and *A. hyperici* (one female, four males) flew distances long enough
to stand out from the right tail of the distributions, even after their first trial, and may be
considered extreme (Figure 3). The flight speed of *Coraebus* and other species showed lower
variation and was respectively 0.40 m/s $\pm$ 0.03 m/s and 0.26 m/s $\pm$ 0.13 m/s.
Flight distance flown by the tested insects over their lifespan (i.e., including all the trials)
varied also greatly. The median flight distance (Table 2) for *Agrilus* and *Meliboeus* was 53 m
(115 m excluding the 28 % of non-flying insects); a quarter of the beetles flew more than
243 m (401 m), 10 % more than 730 m (1 039 m), and 1 % (3 beetles) flew between 2 890 m
and 4 636 m (2 916 m and 4 636 m). Distances were longer for *C. undatus*: median distance
of 967 m with one exceptional beetle above 16 km. Flown distance was globally linked with
the number of trials (LM, p < 0.001, R^2 = 0.42, *C. undatus* excluded); no significant variation
with species or sex had been observed.

Figure 3. Distribution of total distance flown per trial depending on sex, all 12 species
pooled together ($n_{females}$ = 123, n_{males} = 99). Darker bars represent the first trial (eight
hours) while lighter bars represent all consecutive trials (four hours), cumulated on
top of the first trial. Individual flights $\geq$ 1300 m are marked with an arrow indicating
the corresponding species, with the trial number and individual identifier in brackets.

Regardless of their species, most tested individuals performed several short flight bouts
within a single eight-hour trial (Figure 2). Flight bout number was highly variable: 25 or less
for half of trials with flight activity, but more than 58 for a quarter (39 trials), and 16 trials
(10 %) overreaching this number with more than 150 flight bouts over eight hours. Three
trials even exceeded 300 flight bouts: two by females *A. hyperici* (407 and 379 flight bouts)
and one by a female *A. laticornis* (308 flight bouts). The distance flown during a flight bout
was also highly variable (median 2 m, 10% above 9.5 m, up to 653 m).

Interspecific variation in flight performance

Among the four flight performance variables measured during the first (eight-hour) trial, two of them varied significantly among species (sex pooled): total flight distance (Kruskal-Wallis, $\chi^2(7) = 20.4$, $p = 0.005$, n = 208) and mean distance flown per flight bout (Kruskal-Wallis, $\chi^2(7) = 23$, $p = 0.002$, n = 208). However, Dunn's tests highlighted only a few differences among groups: between *A. olivicolor* and *C. undatus* for the total flight distance, between *A. hastulifer* and *A. laticornis* for the mean distance flown per flight bouts. Latency to first flight bout ($\chi^2(7) = 8.46$, $p = 0.29$, n = 208) and number of flight bouts ($\chi^2(7) = 13.7$, $p = 0.06$, n = 208) were not influenced by species.

Factors of intraspecific variation in flight performance

Intraspecific differences in the three most abundant species (*A. hyperici, A. laticornis* and *A. olivicolor*) demonstrated that endogenous factors influenced some of their flight capacities. Preflight mass positively affected total flight distance, as well as the number of flight bouts in *A. hyperici* and *A. laticornis*, but not in *A. olivicolor* (Table 3). Response variables showed no sex bias, as either fixed effects had no significant influence, or null models were better. However, marginal R^2 were weak and other factors likely have impacted flight performance. Only *A. hyperici* showed standing out marginal R^2 with number of moves ($R^2 = 0.25$) and total flight length ($R^2 = 0.28$) while every other R^2 did not exceed 0.13.

Table 3. LMER tests results for the effects of sex and preflight mass on a selection of response variables during the first (eight-hour) trial, with year of experiment as random effect. Response variables in each Linear Mixed-Effects Models were log-transformed. The reference level for the categorial factor 'sex' is female. ΔAICc = AICc (full model) – AICc (null model). The full model was considered to better explain the response variable than the null model when ΔAICc was lower than -2, and the opposite when ΔAICc was higher than -2. Conditional R^2 corresponds to the variance of both fixed and random effects while marginal R^2 considers only the variance of the fixed effect.

Species	Variables	ΔAICc	Sex (female)			Mass			Cond. R2	Marg. R2
			Estimate	SE	t value	Estimate	SE	t value		
A. hyperici	Total flight distance	-22.51	1.09	0.85	1.28	1 025.36 ***	264.91	3.87		0.28
	Number of flight bouts	-19.98	0.45	0.61	0.73	679.24 ***	189.78	3.58		0.25
	Mean flight bout distance	8.62	-	-	-	-	-	-	-	-
	Latency to first flight bout	-11.33	0.85	1.07	0.79	-308.19	369.73	-0.83	0.09	0.07
A. laticornis	Total flight distance	-17.81	0.81	0.44	1.85	461.98 **	150.70	3.07	0.14	0.09
	Number of flight bouts	-15.85	0.54	0.34	1.56	351.75 **	116.63	3.02	0.11	0.09
	Mean flight bout distance	-3.27	0.05	0.15	0.35	5.67	51.74	0.11	0.12	0.001
	Latency to first flight bout	-8.95	-0.30	0.61	-0.48	13.63	203.92	0.07		0.004
A. olivicolor	Total flight distance	-10.82	1.58	0.97	1.63	456.42	346.36	1.31	0.45	0.12
	Number of flight bouts	-9.36	1.13	0.67	1.71	294.03	238.59	1.23	0.42	0.13
	Mean flight bout distance	7.55	-	-	-	-	-	-	-	-
	Latency to first flight bout	1.67	-	-	-	-	-	-	-	-

Consecutive flight performance and patterns

In four-hour trials, the distribution of flight performance was similar to that of the first (eight-hour) one (Figure 3), but flight performance varied over consecutive trials. Data on *C. undatus* were excluded from these analyses, due to their specificities discussed below. Overall, raw flight performance decreased over consecutive trials (Figure 4 left, circle symbols; Kruskal-Wallis, $\chi^2(9) = 46.2$, $p < 0.001$, $n = 501$). Differences were mainly observed between the first trial and others. Extreme long distance flights ($> 1\,250$ m, arbitrary threshold) only took place during the four first trials (and mainly in the first one). Flight performance increased in the late trials, when insects benefited from longer rest periods, but remained before first trials one (Figure 4 left, plus symbols). However, the global distribution showed that beetles flew a greater distance on their first trial, with no variation between subsequent trials (Figure 4 right; Kruskal-Wallis, $\chi^2(9) = 140$, $p < 0.001$, $n = 501$).

Figure 4. Changes in flight performance during consecutive trials. **Left:** Distribution of distance flown during four-hour trials across consecutive trials. **Right:** Distribution of relative flight performance across consecutive trials; for each beetle and trial, relative flight performance was computed as the flight distance during the trial divided by the mean flight distance during all trials. For both plots, data of the eight-hour trial was truncated to the first four hours. Unusual trial frequency data and data from *C. undatus* were discarded from analyses; however, late trial data are displayed on the left panel (plus symbols). Different letters above the data indicate a significant difference according to Dunn's test.

Variable flight patterns over time were observed among species, with *A. hyperici, A. graminis,* and *C. undatus* standing out from the others because a fraction of their individuals remained alive and covered long distances after the first or even subsequent trials (Figures 5 and S4). No such pattern was observed in other species, either due to a generally low flight activity, or comparatively lower lifespan in rearing conditions, reducing the number of consecutive trials per individual. Regardless of species, 35 % of individuals were dead after the first trial, 70 % after the second and 86 % after the third. Most individuals (93 %) were dead after four trials. The vast majority of individuals that survived across multiple trials were unable to sustain long flight distances over time (mainly feeding the "[0]" category in figures 5 and S4), such as *A. laticornis* and *A. hastulifer.* However, some individuals diverged from this trend. A female *C. undatus* and a male *A. hyperici* flew long distances even after their first flight mill trial, as well as a female *A. hyperici* flying more than 800 m during its fifth trial (although not indicated with an arrow in figure 3; Figures 5 and

S4). Remarkably, the extreme *C. undatus* female flew several kilometres in five consecutive trials, cumulating over 16 km of active flight after its capture.

Figure 5. Individual flight trajectories across four consecutive trials in the three most numerous species (*A. laticornis*, *A. hyperici* and *A. olivicolor*), with the first trial spanning eight hours and the following four hours. Total flight distance is discretized in categories of increasing range to simplify individual trajectories. The NA category corresponds to missing data due to a technical issue during the trial (typically, a temporary escape of an individual).

                                    Discussion

Flight performance and behaviour

This study allowed a first analysis of flight characteristics of several European Agrilinae,
including flight performance and behaviour. It revealed several common patterns and some
specificities among species. A clear observation was the high variation in most flight
performance parameters among individuals. Prior history (including age and flight history)
is a likely factor for this variation, but is unknown, albeit probably only part of the picture,
since flight activity is typically known as an intrinsically variable trait among individuals and
high inter-individual variability is common in flight mill experiments (Taylor et al., 2010;
Sauvard et al., 2018). Flight performance decreased over the course of the trials, but changed
little over a few ones; remarkably, the best flyer in *C. undatus* was able to fly long distances
during several consecutive trials.

At the species level, the assessment of flight performance of several Agrilinae revealed
rather homogeneous flight capacities among species, under the experimental conditions of
tethered flights. Only the few *C. undatus* individuals presented substantial differences. Even
if their number was too low to exhibit many significant results, these beetles actually
performed some flights which largely exceeded the dispersal distance of any other species.
However, this species is also much bigger than others. The huge intraspecific variation and
the relatively low sample size, moreover unbalanced between species, had probably impeded
identifying specific flight characteristics. *Coraebus undatus* excepted, the low variation in
beetle size among the sampled species and the low flight activity of most tested individuals
might have also reduced our ability to discriminate the results among species. Data available
on flight capacities of other *Agrilus* species in similar experiment indicates better
performance: median of 750 m for *A. planipennis* (Taylor et al., 2010) and mean between 200
to 1 800 m for *Agrilus auroguttatus* (Schaeffer), depending on conditions (Lopez et al., 2014).
These values are rather similar to *C. undatus* performance, but, as this latter species, *A.*
planipennis and *A. auroguttatus* are also bigger than the *Agrilus* and *Meliboeus* species we
tested. From a more general perspective, Agrilinae performances are lower than those of
some bigger beetles such as *Monochamus galloprovincialis* (Olivier) (2 km per day; David et
al., 2014) and *Rhynchophorus ferrugineus* (Olivier) (2.6 km per day; Ávalos et al., 2014), and
far from those of some long-flight insects such as hornets (median about 20 km per day;
Sauvard et al., 2018).

At the trial level, most flying individuals showed no sustained flight, and instead
completed multiple short flight bouts. This flying pattern could correspond to "routine flight"
(Van Dyck & Baguette, 2005), typically associated with foraging or the search for a sexual
partner or shelter. In contrast, these authors described longer flights as "dispersal flights",
presumably allowing insects to escape from bad conditions or predators, and were almost
never observed during our experiment. However, opposing the two flight types may not
always be relevant. In fact, our study showed that *Agrilus* species could spread by performing
a series of short flights ultimately resulting in relatively long total distances covered, instead
of single sustained flights (e.g., Ávalos et al., 2014).

Globally, tested beetles tended to be poor dispersers in the experimental setting, with
most of them not taking off or flying over very short ranges (generally less than 200 m). Only
C. undatus seemed to fly significantly longer distances (median of 1 000 m). This observation
may suggest resident strategies, i.e., individuals that tend to stay in the range of their
emergence area. One third of beetles did not fly at all. This is congruent with studies on other

beetle species, which found a similar percentage of non-flyers, such as for *Ips sexdentatus*
Boern (Jactel, 1993) and *R. ferrugineus* (Ávalos et al., 2014). These authors hypothesised that
such a high proportion of non-flying insects might be due to the tethered of the beetles and
to the flight mill itself, or be a consequence of the individual incapacity to fly. As some non-
flying beetles in a trial also demonstrated being able to fly during a later trial, not flying was
probably due to a combination of behavioural variations, propensity to take-off, and
constraints of the flight mills, but not solely the latter. Nevertheless, as flight performance
distribution had a very long tail, some insects largely exceeded the median performance of
their conspecifics and could be considered good dispersers. As such, they may perform better
at reaching new distant habitats if enough patches of viable hosts are available on their path
(Lopez et al., 2014), which is normally the case for oak-associated species in European
temperate areas. Within *Agrilus* species, an *A. hyperici* appeared to be the most active
disperser; interestingly, this species was the only one that was not associated with forest
habitats in our sample, and their relatively high performance could be related to its more
scattered herbaceous habitat. *A. hyperici* was introduced in western Canada during the last
century and, even though no accurate data is available, it seems to have largely spread in this
region since; however, this took several decades despite active human assistance (Gov. of
British Columbia, 2018). This observation *in natura* is consistent with modest flight
capacities in this species. Alongside their flight capacities, it is possible that small Agrilinae
species occasionally take advantage of winds to maximize dispersal , as observed in some
bark beetles up to several tens of kilometres (Nilssen, 1984). Moreover, the actual realized
dispersal in Agrilinae could result from the combination of short-distance flight and human-
assisted transportation (i.e., with wood material), sometimes over long distances, as
observed in *A. planipennis* (Muirhead et al., 2006), *Agrilus bilineatus* (Weber) (Baranchikov
et al., 2019b) and related species (e.g., Wu et al., 2017; Ruzzier et al., 2023).

No sexual dimorphism in flight performance

We expected greater flight performance in females than males, as already demonstrated
with mated females of *A. planipennis* (Taylor et al., 2010). Yet, sex did not impact any flight
variables in the overall sample. Those results are concordant, however, with observations on
A. auroguttatus whose flight capacities were not affected by sex either (Lopez et al., 2014).
Lack of sexual dimorphism in flight performance seems then frequent in Agrilinae species.
Both sexes are suspected to participate in dispersal in the tested species. Males are known to
use flight to actively search for females, while mated females typically disperse to find a
suitable host for egg laying (Chamorro et al., 2015), and both behaviours may contribute to
species dispersal. However, it is rather considered that females are the ones which are able
to disperse far away from their host when necessary while males stay close to their natal or
maturation area (Rodriguez-Soana et al., 2007; Mercader et al., 2009; Taylor et al., 2010).

Importance of body mass on flight performance

Body mass is known to be positively correlated with flight performance in many insect
species, and is generally considered to reflect the available fuel load (Minter et al., 2018).
Such a relation was observed in the majority of the species we tested, including the best flyer
(*C. undatus*) and the two most numerous species (*A. laticornis* and *A. hyperici*). Relation to
body mass seemed to differ between these groups; in *C. undatus*, heavier individuals tend to
fly immediately and to perform longer bouts; in the two *Agrilus*, heavier individuals tend, on
the contrary, to increase the number of bouts. In contrast to these species, performance of

other ones, including the third most numerous one (*A. olivicolor*), was not impacted by preflight mass, which corroborates what was observed in some other *Agrilus* species, i.e., *A. auroguttatus* (Lopez et al., 2014) and *A. planipennis* (Taylor et al., 2010). This difference could be related to the relative importance of reserves *vs.* dead weight (mainly structure) in mass variation, but also to insect strategy (noticeably, they could mainly fly when reserves are high, or, on the contrary, when they are low, to forage in order to increase them). It should be noted, however, that even when preflight mass was related to flight performance, it explains only a small part of its variation. Other factors should be considered, and the effect of some of them, diet, mated status and age, has been demonstrated in other *Agrilus* species (Taylor et al., 2010; Lopez et al., 2014). Future studies should include these factors to further investigate Agrilinae flight performance. Nevertheless, our results on species of different body mass (*Agrilus* and *Meliboeus* species *vs. C. undatus*) and previous literature on *A. planipennis* and *A. auroguttatus* suggest that the positive correlation between body mass and flight performance stands among Agrilinae species.

Limitations of the study and interpretation caveats

The sampling design based on interception traps allowed capturing multiple species associated with oak forests, and not only oak borers. However, it prevented controlling for additional factors pertaining to the prior experience of individuals (flight, diet, mated status) and inherently biased captures towards dispersing individuals. This may cause either (1) undersampling of the most resident individuals, even more so than those identified in this study, or (2) sampling of individuals that already spent energy on dispersal before the experiment, or a combination of both. Together with the limited number of captured individuals for some species and the impossibility to control some factors, this warrants using a different sampling strategy in further studies focusing on only oak-borers. We used dry collectors in green multi-funnel traps which are a relevant method for Buprestidae (Allison & Redak, 2017). However, our sampling design likely let a fraction of individuals escape even if the collector was emptied every workday, possibly those showing the highest locomotor activity. Indeed, similar traps but equipped with wet collectors were installed for another study nearby one of the sampling sites, and captured a more abundant and richer community of beetles (A. Sallé, pers. comm.). Emergence traps can yield very good results (e.g., Taylor et al., 2010; Lopez et al., 2014 for borers), but would have restricted the sample to borers of the equipped hosts only, instead of targeting all circulating Buprestidae in oak forests.

The intrinsic dispersal potential of a flying insect species is generally determined mainly by the overall flight capacity of its individuals throughout their life, more specifically that of its best flyers. In the Agrilinae species studied here (except perhaps *C. undatus*), this capacity appeared low, with about 25% of insects not flying in the experimental conditions they were subjected to, and actively flying individuals rarely overreaching one kilometre. As flight performance appeared to be rather maintained over consecutive trials, flight capacity was highly dependent on lifespan. However, the beetles proved to be difficult to maintain alive in rearing conditions, with a median lifespan of four days across all species. As a comparison, *A. planipennis* was reported to live between 2 and 17 weeks under laboratory conditions, with an average of 7 weeks for males and 9 weeks for females (Chamorro et al., 2015 and references within), albeit figures indicated a considerable variability in lifespan in this species as well. Given our observations of limited leaf damage and faeces accumulation in the rearing vials, we suspect insects failed to feed properly in the conditions they were offered,

for unknown reasons. Our method was nevertheless the same one used successfully by other
studies of flight performance of *Agrilus* spp. (Taylor et al., 2010; Lopez et al., 2014). Combined
effects of starvation and daily flight trials in flight mills likely impacted survival severely in
the present experiment, ultimately reducing lifespan and underestimating the overall flight
capacity of these species. Moreover, lifespan values are to be taken with caution since the age
of individuals at the time of their capture remains unknown.

Flight mills under controlled conditions are a proven method to assess the flight potential
of insects (e.g., Sauvard et al., 2018 and references therein). Since the method can easily be
standardized under controlled conditions, flight mills are especially valuable to compare
species or investigate the influence of factors on flight activity. However, tethered flight
performance cannot be considered a direct measurement of free flight or natural flight (Ribak
et al., 2017; Sauvard et al., 2018; Naranjo, 2019), and cautious interpretation is advised. Flight
mills create artificial conditions and could both hinder take-off (inertia) or promote flight
(support making the insects weightless, inertia of the rotating arm). The absence of a moving
natural landscape in the vision of the specimens is likely to influence their locomotor
behaviour, and the influence of sensed angular velocity or weightlessness on tethered insects'
propensity to fly is unknown. Tethered flight experiments have already been suspected of
under- or overestimating flight abilities due to the lack of tarsal contact and the absence of a
perch for taking off or landing (Minter et al., 2018) or because of the device itself (Taylor et
al., 2010). Yet, they remain a highly relevant and standard method to compare conspecifics
or sympatric species in a given set of conditions to approach their motivation to disperse
under those settings, or even physiological expenditures in flight dispersal when their flight
propensity is high. In other respects, flight mills likely limited the flight speed in our
experiment, as was found in another flight mill study on *A. planipennis* with a three-fold
difference between free and tethered flight speeds (Taylor et al., 2010).

Conclusions and perspectives

This work provided a first comparative approach on the flight behaviour and performance
in several Agrilinae species associated with oak forests. It showed several common traits
among the focal species, namely (1) the considerable inter-individual variation, (2) the low
average flight performance compared to other insect species using similar experimental
designs, and (3) the relative homogeneity of this pattern among most of the species
investigated, despite the probable existence of different flight behaviour among species as
seen with long sustained flights in *C. undatus*. This provides insights into the poorly
understood dispersal ecology of oak-associated Agrilinae, suggesting a generally low average
dispersal propensity and the importance of scarce events carried by a few extreme
individuals in shaping the ultimate colonization and spread patterns at the population and
species levels.

While the present work aimed at investigating flight in Agrilinae using a multispecific
approach that, to our knowledge, has rarely been attempted with flight mill experiments, an
oak-borer species considered a key aggravating agent of oak decline in Europe is missing in
our sampling: *A. biguttatus*. This beetle is currently expanding its range with climate change
(Pederson & Jørum, 2009; Brown et al., 2015) and genetic molecular markers have
highlighted population differentiation at the European scale (Le Souchu et al. submitted). Yet,
little is known about the dispersal ability of this species, let alone its variation among
populations. Flight mills may prove a relevant method to better understand the genetic
structure and movements of populations in this species, especially as it is considered a major

threat outside its current range (e.g., in North America, USDA-APHIS, 2011). As the biggest
European *Agrilus* species, at a size similar to that of *A. planipennis* (Volkovitsh et al., 2019)
in which flight ability is described in most details among *Agrilus* species, comparing both
species would be insightful to better understand the spread potential of *A. biguttatus.*

Appendices

597 The supplementary material is available in the open repository Data Recherche Gouv at
598 https://doi.org/10.57745/TY9KLG.

Acknowledgements

600 We are thankful for the technical assistance offered by Maxime Stanislawek, Clément
601 Beaumont and Mathieu Plateau.

Funding

603 This work was supported by the Région Centre-Val de Loire Project no. 2018-00124136
604 (CANOPEE) coordinated by A. Sallé.

Conflict of interest disclosure

606 The authors declare no conflicts of interest and they comply with the PCI rule of having
607 no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

The data and code that support the findings of this study are openly available in three
repositories. The main one, at https://doi.org/10.57745/TY9KLG, includes the material
specific to the present study: supplementary data, videos of insects flying in the rotational
flight mills, raw log files from the flight mills, associated processed data and final *R* scripts.
Git-versioned *R* scripts are also available at
https://forgemia.inra.fr/mathieu.laparie/agrilusflight. The source of the *Crystal* program
used to monitor flight mills and produce log files is available at
https://doi.org/10.57745/DIWSWF. The *Ruby* pipelines used to process log files into flight
data are available at https://doi.org/10.57745/YLMCNU.

References

Allison JD, Redak RA (2017) The Impact of Trap Type and Design Features on Survey and
Detection of Bark and Woodboring Beetles and Their Associates: A Review and Meta-
Analysis. *Annual Review of Entomology*, **62**, 127–146. https://doi.org/10.1146/annurev-
ento-010715-023516
Ávalos JA, Martí-Campoy A, Soto A (2014) Study of the flying ability of *Rhynchophorus*
ferrugineus (Coleoptera: Dryophthoridae) adults using a computer-monitored flight mill.
Bulletin of Entomological Research, **104**, 462–470.
https://doi.org/10.1017/S0007485314000121

Baranchikov Y, Björklund N, Bonifacio LF, Matthews-Berry S, Petrice T, Sallé A, Wang X-Y, Suffert M, Picard C (2019a) Pest risk Analysis for *Agrilus fleischeri* (Coleoptera: Buprestidae). https://doi.org/10.13140/RG.2.2.33002.75204

Baranchikov Y, Björklund N, Bonifacio LF, Matthews-Berry S, Petrice T, Sallé A, Wang X-Y, Suffert M, Picard C (2019b) Pest risk analysis for *Agrilus bilineatus* (Coleoptera: Buprestidae), two-lined chestnut borer. https://doi.org/10.13140/RG.2.2.19580.97926

Bílý S (2002) Summary of the bionomy of the Buprestid beetles of Central Europe (Coleoptera: Buprestidae). *Acta Entomological Musei Nationalis Pragae*, **10**, 104.

Bozorov TA, Luo Z, Li X, Zhang D (2019) *Agrilus mali* Matsumara (Coleoptera: Buprestidae), a new invasive pest of wild apple in western China: DNA barcoding and life cycle. *Ecology and Evolution*, **9**, 1160–1172. https://doi.org/10.1002/ece3.4804

Brown N, Inward DJG, Jeger M, Denman S (2015) A review of *Agrilus biguttatus* in UK forests and its relationship with acute oak decline. *Forestry*, **88**, 53–63. https://doi.org/10.1093/forestry/cpu039

Brück-Dyckhoff C, Petercord R, Schopf R (2019) Vitality loss of European beech (*Fagus sylvatica* L.) and infestation by the European beech splendour beetle (*Agrilus viridis* L., Buprestidae, Coleoptera). *Forest Ecology and Management*, **432**, 150–156. https://doi.org/10.1016/j.foreco.2018.09.001

Chamorro ML, Jendek E, Haack R, Petrice T, Woodley N, Konstantinov A, Volkovitsh M, Yang X-K, Grebennikov V, Lingafelter S (2015) *Illustrated Guide to the emerald ash borer* Agrilus planipennis *Fairmaire and Related Species (Coleoptera, Buprestidae)*. Pensoft Publishers, Sofia, Bulgaria.

Crall JD, Chang JJ, Oppenheimer RL, Combes SA (2017) Foraging in an unsteady world: bumblebee flight performance in field-realistic turbulence. *Interface Focus*, **7**, 20160086. https://doi.org/10.1098/rsfs.2016.0086

David G, Giffard B, Piou D, Jactel H (2014) Dispersal capacity of *Monochamus galloprovincialis*, the European vector of the pine wood nematode, on flight mills. *Journal of Applied Entomology*, **138**, 566–576. https://doi.org/10.1111/jen.12110

Dickinson MH, Lehmann F-O, Sane SP (1999) Wing Rotation and the Aerodynamic Basis of Insect Flight. *Science*, **284**, 1954–1960. https://doi.org/10.1126/science.284.5422.1954

Du Merle P, Attié M (1992) Coraebus undatus (Coleoptera: Buprestidae) sur chêne liège dans le Sud-Est de la France: estimation des dégâts, relations entre ceux-ci et certains facteurs du milieu. *Annales des sciences forestières*, **49**, 571–588.

Dubois GF, Vernon P, Brustel H (2009) A flight mill for large beetles such as Osmoderma eremita (Coleoptera: Cetoniidae). In: *Buse J, Alexander KNA, Ranius T, Assmann T, Saproxylic Beetles: Their Role and Diversity in European Woodland and Tree Habitats-Proceedings of the 5th Symposium and Workshop. Pensoft Publishers, Sofia*, pp. 219–224.

Dudley R (2002) *The biomechanics of insect flight: form, function, evolution*. Princeton Univ.y Press, Princeton, N.J.

Evans HF, Williams D, Hoch G, Loomans A, Marzano M (2020) Developing a European Toolbox to manage potential invasion by emerald ash borer (*Agrilus planipennis*) and bronze birch borer (*Agrilus anxius*), important pests of ash and birch. *Forestry: An International Journal of Forest Research*, **93**, 187–196. https://doi.org/10.1093/forestry/cpz074

Gov. of British Columbia (2018) *Agrilus hyperici (Creutzer)*. Canada.

Herms DA, McCullough DG (2014) Emerald Ash Borer Invasion of North America: History, Biology, Ecology, Impacts, and Management. *Annual Review of Entomology*, **59**, 13–30. https://doi.org/10.1146/annurev-ento-011613-162051

Hızal E, Arslangündoğdu Z (2018) The First Record of Two-Lined Chestnut Borer *Agrilus bilineatus* (Weber, 1801) (Coleoptera: Buprestidae) from Europe. *Entomological News*, **127**, 333–335. https://doi.org/10.3157/021.127.0404

Jactel H (1993) Individual variability of the flight potential of *Ips sexdentatus* Boern. (Coleoptera: Scolytidae) in relation to day of emergence, sex, size, and lipid content. *The Canadian Entomologist*, **125**, 919–930. https://doi.org/10.4039/Ent125919-5

Jendek E, Grebennikov V (2009) *Agrilus sulcicollis* (Coleoptera: Buprestidae), a new alien species in North America. *The Canadian Entomologist*, **141**, 236–245. https://doi.org/10.4039/n09-021

Jendek E, Grebennikov VV (2023) Summary of native geographic distribution of all 3,341 species of the most speciose animal genus Agrilus (Coleoptera: Buprestidae). *Journal of Insect Biodiversity*, **39**. https://doi.org/10.12976/jib/2023.39.2.1

Jendek E, Poláková J (2014) *Host Plants of World* Agrilus *(Coleoptera, Buprestidae)*. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-319-08410-7

Le Souchu E, Bouget C, Sallé A (2024) Environmental drivers of local and temporal variations in the community of oak-associated borers (Coleoptera: Buprestidae). *European Journal of Forest Research*. https://doi.org/10.1007/s10342-023-01644-y

Lopez VM, McClanahan MN, Graham L, Hoddle MS (2014) Assessing the Flight Capabilities of the Goldspotted Oak Borer (Coleoptera: Buprestidae) With Computerized Flight Mills. *Journal of Economic Entomology*, **107**, 1127–1135. https://doi.org/10.1603/EC13525

Macháčová M, Nakládal O, Samek M, Baťa D, Zumr V, Pešková V (2022) Oak Decline Caused by Biotic and Abiotic Factors in Central Europe: A Case Study from the Czech Republic. *Forests*, **13**, 1223. https://doi.org/10.3390/f13081223

Mercader RJ, Siegert NW, Liebhold AM, McCullough DG (2009) Dispersal of the emerald ash borer, *Agrilus planipennis*, in newly-colonized sites. *Agricultural and Forest Entomology*, **11**, 421–424. https://doi.org/10.1111/j.1461-9563.2009.00451.x

Minter M, Pearson A, Lim KS, Wilson K, Chapman JW, Jones CM (2018) The tethered flight technique as a tool for studying life-history strategies associated with migration in insects: Tethered flight for studying insect movement. *Ecological Entomology*, **43**, 397–411. https://doi.org/10.1111/een.12521

Muirhead JR, Leung B, Van Overdijk C, Kelly DW, Nandakumar K, Marchant KR, MacIsaac HJ (2006) Modelling local and long-distance dispersal of invasive emerald ash borer *Agrilus planipennis* (Coleoptera) in North America. *Diversity and Distributions*, **12**, 71–79. https://doi.org/10.1111/j.1366-9516.2006.00218.x

Naranjo SE (2019) Assessing Insect Flight Behavior in the Laboratory: A Primer on Flight Mill Methodology and What Can Be Learned. *Annals of the Entomological Society of America*, **112**, 182–199. https://doi.org/10.1093/aesa/say041

Nilssen A (1984) Long-range aerial dispersal of bark beetles and bark weevils (Coleoptera, Scolytidae and Curculionidae) in northern Finland. In: *Annales Entomologici Fennici*, pp. 37–42. Entomological Society of Finland.

Orlova-Bienkowskaja MJ, Drogvalenko AN, Zabaluev IA, Sazhnev AS, Peregudova EY, Mazurov SG, Komarov EV, Struchaev VV, Martynov VV, Nikulina TV, Bieńkowski AO (2020) Current range of *Agrilus planipennis* Fairmaire, an alien pest of ash trees, in European Russia and Ukraine. *Annals of Forest Science*, **77**, 29. https://doi.org/10.1007/s13595-020-0930-z

Pederson H, Jørum P (2009) The jewel beetle *Agrilus biguttatus* (Fabricius, 1777) found in Denmark (Coleoptera, Buprestidae). *Entomologiske Meddelelser*, **77**, 19–26.

R Core Team (2022) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

R Core Team (2023) *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.
Rassati D, Marini L, Marchioro M, Rapuzzi P, Magnani G, Poloni R, Di Giovanni F, Mayo P,
Sweeney J (2019) Developing trapping protocols for wood-boring beetles associated with
broadleaf trees. *Journal of Pest Science*, **92**, 267–279. https://doi.org/10.1007/s10340-
018-0984-y
Reed K, Denman S, Leather SR, Forster J, Inward DJG (2018) The lifecycle of *Agrilus*
biguttatus: the role of temperature in its development and distribution, and implications
for Acute Oak Decline. *Agricultural and Forest Entomology*, **20**, 334–346.
https://doi.org/10.1111/afe.12266
Renault D, Angulo E, Cuthbert RN, Haubrock PJ, Capinha C, Bang A, Kramer AM, Courchamp F
(2022) The magnitude, diversity, and distribution of the economic costs of invasive
terrestrial invertebrates worldwide. *Science of The Total Environment*, **835**, 155391.
https://doi.org/10.1016/j.scitotenv.2022.155391
Ribak G, Barkan S, Soroker V (2017) The aerodynamics of flight in an insect flight-mill (R
Gurka, Ed,). *PLOS ONE*, **12**, e0186441. https://doi.org/10.1371/journal.pone.0186441
Robinet C, Imbert C-E, Rousselet J, Sauvard D, Garcia J, Goussard F, Roques A (2012) Human-
mediated long-distance jumps of the pine processionary moth in Europe. *Biological*
Invasions, **14**, 1557–1569. https://doi.org/10.1007/s10530-011-9979-9
Rodriguez-Soana CR, Miller JR, Poland TM, Kuhn TM, Otis GW, Turk T, Ward DL (2007)
Behaviors of Adult *Agrilus planipennis* (Coleoptera: Buprestidae). *The Great Lakes*
Entomologist, **40**. https://doi.org/10.22543/0090-0222.2168
Ruzzier E, Haack RA, Curletti G, Roques A, Volkovitsh MG, Battisti A (2023) Jewels on the go:
exotic buprestids around the world (Coleoptera, Buprestidae). *NeoBiota*, **84**, 107–135.
https://doi.org/10.3897/neobiota.84.90829
Sallé A, Nageleisen L-M, Lieutier F (2014) Bark and wood boring insects involved in oak
declines in Europe: Current knowledge and future prospects in a context of climate
change. *Forest Ecology and Management*, **328**, 79–93.
https://doi.org/10.1016/j.foreco.2014.05.027
Sauvard D, Imbault V, Darrouzet É (2018) Flight capacities of yellow-legged hornet (Vespa
velutina nigrithorax, Hymenoptera: Vespidae) workers from an invasive population in
Europe (W Blenau, Ed,). *PLOS ONE*, **13**. https://doi.org/10.1371/journal.pone.0198597
Taylor RAJ, Bauer LS, Poland TM, Windell KN (2010) Flight Performance of *Agrilus*
planipennis (Coleoptera: Buprestidae) on a Flight Mill and in Free Flight. *Journal of Insect*
Behavior, **23**, 128–148. https://doi.org/10.1007/s10905-010-9202-3
USDA-APHIS (2011) New pest response guidelines: exotic wood-boring and bark beetles.
Van Dam WA, Williams RN, Taylor RAJ (2000) Flight performance of some nitidulid beetles
(Coleoptera) using a computer-monitored flight mill. *J. Agric. Urban Entomol.*, **17**, 143–
151.
Van Dyck H, Baguette M (2005) Dispersal behaviour in fragmented landscapes: Routine or
special movements? *Basic and Applied Ecology*, **6**, 535–545.
https://doi.org/10.1016/j.baae.2005.03.005
Volkovitsh MG, Kovalev AV, Orlova-Bienkowskaja MJ (2020) Current Distribution and
Diagnostic Features of Two Potentially Invasive Asian Buprestid Species: *Agrilus mali*
Matsumura and *A. fleischeri* Obenberger (Coleoptera: Buprestidae). *Insects*, **11**, 493.
https://doi.org/10.3390/insects11080493
Volkovitsh MG, Orlova-Bienkowskaja MJ, Kovalev AV, Bieńkowski AO (2019) An illustrated
guide to distinguish emerald ash borer (*Agrilus planipennis*) from its congeners in Europe.

Forestry: An   International   Journal   of   Forest   Research,   cpz024.
https://doi.org/10.1093/forestry/cpz024
Wu Y, Trepanowski NF, Molongoski JJ, Reagel PF, Lingafelter SW, Nadel H, Myers SW, Ray AM
(2017) Identification of wood-boring beetles (Cerambycidae and Buprestidae)
intercepted in trade-associated solid wood packaging material using DNA barcoding and
morphology. *Scientific Reports*, **7**, 40316. https://doi.org/10.1038/srep40316