Intra- and interspecific variations in flight performance of oak-associated Agrilinae (Coleoptera: Buprestidae) using computerised flight mills

Le Souchu Elodie[1,3], Sallé Aurélien[1], Bankhead-Dronnet Stéphanie[1], Laparie Mathieu*[2], Sauvard Daniel*[2]

[1] University of Orléans, Laboratory of Physiology, Ecology and Environment (P2E), INRAE USC 1328 - Orléans, France
[2] INRAE, URZF (Forest Zoology Research Unit) - Orléans, France
* These authors contributed equally to the work.

[3] Corresponding author: Le Souchu Elodie
Correspondence: elodie.lesouchu2@gmail.com
Present address: Swedish University of Agricultural Sciences (SLU), Department of Ecology - Inst för Ekologi, Box 7044, 750 07 Uppsala, Sweden

ABSTRACT

Several Agrilinae species (Coleoptera: Buprestidae) are secondary pests of broadleaf forests, and some of them are also major invasive pests. These thermophilous borers are expected to be favoured by climate change and the global deterioration of forest health, and ultimately expand their range and damage. Flight behaviour and performance of these insects are poorly known despite their critical role in dispersal inside and outside native ranges and their relevance for management purposes. This study aimed to assess intra- and interspecific variability in active flight of several Agrilinae species and effects of sex and mass on this variability to contribute to filling this gap. We assessed the flight performance of eleven species associated with oaks (nine *Agrilus*, one *Coraebus* and one *Meliboeus*) plus one *Agrilus* species associated with the herbaceous layer. Computer-monitored flight mills were used to measure flight parameters, including periods, durations, distances and velocity in 250 beetles. Overall, flight capacities were rather homogeneous among species, with a dominance of poor flyers and only *Coraebus undatus* showed outstanding performance. Beetles generally performed several short flight bouts within one trial, and only a few individuals sustained long flight. The maximal total distance covered across multiple assays until death ranged from 170 to 16 097 m depending on the species, with a median between 35 and 966 m (excluding individuals that never flew). On top of this interspecific variability, flight distances also varied greatly among individuals, but were not influenced by sex. Preflight body mass had mixed effects depending on the species, presumably related to different dispersal patterns. In our experimental conditions, most species had limited average dispersal capacities over multiple flight trials. Overall, dispersal over long distances and colonisation events probably depend mainly on a small proportion of individuals which largely exceeded the median performance.

Keywords: *Agrilus*, secondary pest, range expansion, tethered flight, dispersal behavioural, locomotor activity

 Introduction

The jewel beetle subfamily Agrilinae (Coleoptera: Buprestidae) includes numerous
species, especially in the genus *Agrilus* which gathers more than 3 300 described species
worldwide (Jendek & Grebennikov, 2023). Some of these thermophilous and heliophilous
insects (Reed et al., 2018) are known to be phytosanitary threats on a wide diversity of
woody and some non-woody plants (Jendek & Poláková, 2014) and can have important
economic and ecological impacts in their native area. Larvae develop in the roots, stems,
trunks or branches of their host plant. Within trees, larval development could typically last
several years. Imaginal moult takes place in spring, within host tissues. After emergence, they
usually move towards the foliage to perform a maturation feed and mate. Then females
search for a suitable host to lay their eggs (Bílý, 2002).
As opportunistic pests, Agrilinae are associated with forest declines and diebacks (Le
Souchu et al., 2024). In temperate forests, wood-borers such as *Agrilus biguttatus* (Fabricius)
and *Agrilus viridis* (Linnaeus) are aggravating agents of large scale oak and beech forest
declines (Sallé et al., 2014; Brown et al., 2015; Brück-Dyckhoff et al., 2019; Macháčová et al.,
2022). A related Agrilinae species, *Coraebus undatus* (Fabricius) has also been associated
with oak decline (Du Merle & Attié, 1992; Sallé et al., 2014). *Agrilus* species are also known
for their high invasive potential, supported by several examples of human-mediated
invasions (*e.g.,* Wu et al., 2017; Hızal & Arslangündoğdu, 2018; Bozorov et al., 2019). Indeed,
their long and cryptic pre-imaginal life in the wood facilitates unnoticed transportations and
introductions (Chamorro et al., 2015; Ruzzier et al., 2023). One of the prominent invasive
pests worldwide is the Emerald Ash Borer (EAB), *Agrilus planipennis* Fairmaire. It is
responsible for most of the worldwide economic damage attributed to Buprestidae (Renault
et al., 2022). This species has invaded North America from Asia (Herms & McCullough, 2014)
and arrived in Russia in 2003, now threatening the western part of Europe (Orlova-
Bienkowskaja et al., 2020). Other *Agrilus* species have been identified as potential future
invaders, such as *Agrilus anxius* (Evans et al., 2020) or *Agrilus fleischeri* Obenberger
(Volkovitsh et al., 2020), and including species belonging to the European assemblage of oak-
associated borers. Current forest decline phenomena are beneficial to these borers (Le
Souchu et al., 2024), together with climate change which is suspected to assist their
northward expansion (Pederson & Jørum, 2009; Brown et al., 2015). Within this context,
evaluating flight capacities of intercepted or introduced species is of paramount importance
to evaluate their dispersal capacities and implement efficient monitoring and control
strategies.
Active flight is a central function in the life strategies of most insects (Dickinson et al.,
1999; Dudley, 2002). This mode of locomotion is involved in the search for a mate, trophic
resources or favourable environmental conditions, as well as in dispersal and colonisation of
new geographic areas (Dudley, 2002). The distance covered during the lifetime of an
individual can vary considerably among genera, species, and even individuals. It can be
influenced by numerous factors, ranging from intrinsic features (physiological traits,
reserves, morphological characteristics and behaviours, mated status, *etc.*), to external
(a)biotic factors (temperature, crowding, food availability, *etc.*) and their spatiotemporal
variations, immediate or not (Minter et al., 2018). The study of flight capacities and their
variations at multiple focal scales can provide valuable insights to better understand the
ecology of species and, in particular, help to estimate their dispersal potential (see *e.g.,*
Robinet et al., 2012). Yet, despite the economic and ecological damage caused by *Agrilus* spp.
and their predisposition to be accidentally introduced into new environments, this

information remains scarce in the literature. Their flight capacities have rarely been tested,
let alone in European species. Therefore, the numerous works on *A. planipennis* are used to
estimate the expected dispersal capacities of potential invasive *Agrilus* spp. or introduced
ones (*e.g.,* Baranchikov et al., 2019a; b) and highlight their high flight capacities which may
facilitate further dispersal (Taylor et al., 2010; USDA-APHIS, 2011). While males *Agrilus* are
expected to stay within the range of their host, to search for a mate (Rodriguez-Soana et al.,
2007), gravid females are able to disperse several kilometres far from their host (Taylor et
al., 2010). However, it is considered that females will mostly disperse less than 100 m if viable
host plants are available nearby (Mercader et al., 2009).

In this study, we aimed at determining flight capacities of several *Agrilus* species, and
specifically their intra- and interspecific variations. The focus on *Agrilus* behaviour and
biology may help establish management strategies and improve detection tools. We
measured flight potential of oak-associated species using wild individuals tethered to
computer-monitored flight mills. An herbaceous species, *Agrilus hyperici* (Creutzer), similar
in size to most of the oak-associated species we sampled, was also added to the pool of
species. This species was intentionally introduced in North America from France in 1953 as
a biological control agent to target *Hypericum perforatum* L. and its dispersal have been
monitored (Jendek & Grebennikov, 2009; Gov. of British Columbia, 2018). This addition
allowed to (1) confirm that the performance of *A. hyperici* deals with the knowledge on its
dispersion in natura, and (2) compare flight performance of forest species to that of an
herbaceous-associated species. First, we expected noticeable flight differences among
species, as previously observed within multiple coleopteran families (Nitidulidae, Van Dam
et al., 2000; Cetonidae, Dubois et al., 2009). Second, we hypothesised high inter-individual
variations in flight measurements, especially a strong sexual dimorphism with females
putatively flying greater distances than males, as is the case for mated females of
A. planipennis (Taylor et al., 2010).

Material and methods

Sampling and rearing design

We conducted three consecutive years of experiments, from 2021 to 2023, in France. Live
Agrilus were caught using two methods. First, we used green multi-funnel traps (ChemTica
Internacional, San Jose, Costa Rica) with twelve fluon-coated funnels. These traps are
commonly used to sample Agrilinae (Rassati et al., 2019; Le Souchu et al., 2024). From
June 23 to August 06 in 2021 and from May 09 to July 27 in 2022, respectively six and eight
traps were set at the University of Orléans (Longitude: 47.85 N, Latitude: 1.94 E; WGS 84,
DDD), and four and six traps at the INRAE site of Ardon (47.83 N, 1.91 E). These two sites are
separated by 2.6 km. Selected forest patches in these areas were dominated by *Quercus robur*
(L.) and *Quercus petraea* (Matt.) Liebl. A fresh oak leaf was placed in each collector to provide
food and support to trapped insects, and no preservative liquid was used in order to keep
them alive. Traps were hung among the lower branches in the canopy (*i.e.,* approximately
15 m above the ground), to capture adults after their emergence, during their search for a
mate or a suitable host. No lure was added to the traps and the collectors were emptied every
morning during weekdays. Additionally, we captured *A. hyperici* on *H. perforatum* and oak
borers on oak logs using a sweep net. In 2022, *A. hyperici* were sampled in three locations, at
Saint-Denis-en-Val (47.89 N, 1.98 E) on June 26 and 28, at the University of Orléans on
July 04, and at the INRAE site of Ardon on July 06. In 2023, we similarly captured *A. hyperici*

in Vouzon (47.64 N, 2.05 E) on June 16. The same day, several specimens of oak borers were captured in a log deposit nearby (47.65 N, 2.01 E). All combined, the oak-associated species caught were: *Agrilus angustulus* (Illiger), *Agrilus graminis* Kiesenwetter, *Agrilus hastulifer* (Ratzeburg), *Agrilus laticornis* (Illiger), *A. sulcicollis*, *C. undatus*, *Meliboeus fulgidicollis* (P.H. Lucas); and species specialised in others hosts (indicated between parentheses) were: *Agrilus graecus* Obenberger (*Viscum* spp.), *Agrilus olivicolor* Kiesenwetter (*Carpinus* spp.), *A. viridis*. (*Fagus* spp.) and *A. hyperici* (*H. perforatum*). All individuals are listed in the Supplementary Table S1.

Live Agrilinae collected were maintained in the INRAE site of Ardon, in a climatic chamber (Galli GTEST-CL-0700) at a 12:8-hour cycle of 25:20 °C $\pm$ 0.2 °C with 2-hour gradients between both phases, 16:8-hour L:D photoperiod using three LED spots (Lédis, 3.5 W, 300 lumens, 4500 K), and 60 % $\pm$ 5 % relative humidity (warm phase and photophase were centred on solar midday, 2 pm local time). Each insect was put in an individual plastic vial capped with wet cotton to allow ventilation and provide a water source. Since species were determined only after death, all vials but those containing *A. hyperici* (which were known in advance due to the different sampling method) were supplied with a fresh oak leaf renewed every two days, whereas vials containing *A. hyperici* were supplied with fresh *H. perforatum* leaves.

Flight mill design

Rotational flight mills were designed to estimate beetle flight capacities. Each flight mill (Figure 1) was built on a wooden square base (19 × 19 cm), with an upwards wooden stand in the centre and a wooden bracket on one corner, extending up to the centre. A magnet (Supermagnete, https://www.supermagnete.de/, S-05-05-N) was placed under the bracket and another one on the stand, both vertically aligned to hold an entomological pin (size 2). To minimise friction, the pinhead had no contact with the lower magnet, and only the point of the pin was in contact with the upper magnet. The pin supported an horizontal 16-cm arm, composed of two identical 1-mm-diameter carbon rods connected by a 2-mm-diameter 2-cm aluminium tube; a hole was drilled in the middle of the tube, through which the pin was inserted and glued. The arm carried a small magnet (Supermagnete, S-03-02-N) glued on the lower side, near one end of the tube. The insect was placed at the end of the arm opposite to this magnet and drove the arm while flying. For heavy species (*C. undatus*), a counterweight made of reusable adhesive paste was also placed on the arm end opposite to the insect.

One lap of the arm corresponded to a travel distance of 0.5 metre. Arm moves were tracked using two Hall effect sensors (DiodesZetex AH337-PG-B) able to detect magnetic fields. They were positioned on the stand just below the arm, approximately 60° from each other around the pin, and detected every small magnet transit over them. Each Hall sensor was connected to a single board computer (Raspberry Pi), to which signals were transmitted. A total of 26 flight mills were built, and managed using 4 Raspberry Pi (1 Raspberry Pi B+, managing 2 flight mills; 1 Raspberry Pi 2 B, 1 Raspberry Pi 3 B and 1 Raspberry Pi 3 B+, each managing 8 flight mills).

An in-house *Crystal* program running on these computers managed the signals. The program initialised each flight test with flight user-set metadata (mill identifier, insect identifier and gender (if known in advance)), and start time, creating two dedicated CSV files (one per Hall sensor), hereafter referred to as log files. Then, each associated signal detection was time-stamped and recorded in the appropriate file. At the end of the test, end time was also recorded and the log files were closed.

Flight mills were installed in climate chambers (three PHCBi MIR-254-PE, with six flight
mills each, and two Galli GTEST-TM-0700, with four flight mills each; 25 °C ± 0.5 °C, 50 % ±
10 % relative humidity). Each chamber shelf carried a pair of flight mills, which were lit by
two LED spots (same as in the rearing chamber). In order to stimulate flight, fresh oak
branches with leaves (or *H. perforatum* branches when *A. hyperici* was tested) were placed
on each shelf, between the flight mills.

Figure 1. Flight mill used for this study, with a close-up view of the sensor and axis part.
192   The electronics of the system are not shown.

Experimental design

As Agrilinae are diurnal and thermophilous insects (Reed et al., 2018), we tested them for
flight capacities during daytime and at 25 °C ± 0.5 °C. After its sampling, each beetle was
maintained one day in rearing conditions (initial rest). The first flight test occurred the next
working day; it lasted eight hours. The beetle was then tested again on the flight mill each
working day until its death, but these flight tests lasted only four hours. However, test
frequency was lowered to a test per week after the tenth one. Test time was chosen so that
insects were extracted from the rearing chamber at least an hour after the beginning of the
warm phase of their rearing thermal cycle; first tests started at approximately 10 am, and
next tests at about 2 pm. In 2021, due to initial low availability of flight mills, only the first
test was done with some beetles; as two of them remained alive three weeks after, they had
then been tested (during 4 hours) each working day until their death.
Beetles were weighed on a microbalance (Mettler Toledo AG204, d = 0.1 mg) before each
flight test. They were then placed in a 3 °C walk-in chamber to slow them down and ease
gluing a strip of paper narrower than their width on the dorsal side of their pronotum using
cyanoacrylate glue, the most common glue used to tether insects (Naranjo, 2019). The length
and width of strips were adjusted to the dimensions of insects. This operation in cold
conditions lasted from 15 to 30 minutes, then insects were returned to ambient temperature
and immediately attached to flight mills with reusable adhesive paste between the mill's arm
and the strip of paper. This setup allowed rapidly installing and removing tested insects at

the beginning and the end of experiments. At the end of flight tests, paper strips were
delicately removed from insects, as well as any visible residue of glue, and insects were
weighed again before being returned to their rearing vial and chamber.

Data analysis

In-house *Ruby* scripts were developed to analyse the log files generated by the *Crystal*
program used to monitor flight mills. They gathered data from the two log files generated by
a flight mill during a test, separated flight phases (identified using unequal alternance of
signals from the two Hall sensors) from noise and oscillations, and recorded their timestamp,
duration and length in the raw dataset used in further analyses. The flight phases were
hereafter referred to as flight bouts.
We focused our study on several measures of flight capacities, traditionally used to assess
flight performance, which included total flight distance (m), mean flight bout distance (m),
total flight duration (s), mean flight bout duration (s), number of flight bouts and latency to
first flight bout (s). The latency to first flight bout was used as a measure of the propensity to
fly. We also explored the relationship between the fresh body mass before the first flight mill
trial, hereafter referred to as preflight mass, with the performance. For statistical reasons,
species with fewer than six individuals in total were removed from all per-species analyses
or figures, and only included in approaches where all species were pooled together. Some
flight tests experienced technical problems, such as insect escape or electronic issues, and
were excluded from analyses. As indicated above, test frequency changed after the tenth test,
and, in 2021, two beetles completed their tests (except the first one) after a long rest period;
the corresponding data, hereafter referred to as unusual trial frequency data, was also often
excluded from analyses.
Standard linear model (LM) was used to assess linear regression. Kruskal-Wallis tests
were used to assess the difference in flight capacities between groups. The Benjamini-
Hochberg method permitted to correct p-values for False Discovery Rate. Kruskal-Wallis
tests were followed by a post-hoc Dunn test with the Holm adjustment method.
LMERs were used to evaluate the effect of sex and preflight body mass on response
variables associated with flight performance of the three most abundant species (*A. hyperici*,
A. laticornis and *A. olivicolor*). Every response variable was log-transformed to improve the
models, and Q-Q plots were used to review them. Since data from all three years were
analysed together, the year of experiment (*i.e.,* 2020, 2021, 2022) was included as a random
effect. For each response variable, the null model was compared to the model with preflight
mass and sex as explanatory variables, based on their respective AICc (Akaike Information
Criterion scores) with a correction for small sample sizes.
All analyses and figures were realised using *R* 4.2 and R 4.3 (R Core Team, 2022, 2023).

Results

General results

A total of 250 beetles were tested during the three-year experiment, with 102 individuals
in 2021, 125 in 2022, and 23 in 2023 (Table 1). Twelve species were collected, among which
three dominated the samples: *A. laticornis*, *A. hyperici* and *A. olivicolor*. Sex ratio was found
to be relatively balanced in the three most numerous species, and often biased towards fewer
males than females in rarer species, but their small sample sizes warrants caution on this
result. No clear variation of the fresh mass was observed during a flight mill trial or during

beetle life, likely due to the rather low precision of our balance regarding to the mass of the
beetles. Consequently, only the initial preflight mass was used.

Table 1. Sample sizes for each species and sex over the three-year experiment. Values
outside brackets indicate the number of individuals for each species and sex collected
and tested in flight mills, and values within brackets indicate the number of those
insects that could be incorporated in analyses and figures for the first trial. The four
rarest species (indicated by an asterisk) were discarded from all analyses or figures
that included a species factor.

Species	2021 Females		2021 Males		2022 Females		2022 Males		2023 Females		2023 Males		Total	
Agrilus angustulus	1	(0)			1	(1)	2	(2)	5	(5)	2	(2)	11	(10)
*Agrilus graecus ***			1	(0)									1	(0)
Agrilus graminis	2	(2)	3	(3)	3	(3)	1	(1)					9	(9)
Agrilus hastulifer	7	(6)	3	(3)	2	(2)							12	(11)
Agrilus hyperici					14	(14)	13	(11)	8	(8)	7	(7)	42	(40)
Agrilus laticornis	29	(25)	21	(19)	31	(26)	41	(35)	1	(1)			123	(106)
*Agrilus obscuricollis ***	2	(2)	1	(1)	2	(2)							5	(5)
Agrilus olivicolor	8	(6)	8	(6)	5	(5)	1	(1)					22	(18)
*Agrilus sulcicollis ***			1	(1)	1	(1)	2	(2)					4	(4)
*Agrilus viridis ***	3	(3)	2	(1)									5	(4)
Coraebus undatus	4	(4)			2	(2)	1	(1)					7	(7)
Meliboeus fulgidicollis	6	(5)			1	(0)	2	(2)					9	(7)

The overall lifespan after capture was low and highly variable (Supplementary Figure S1).
The median was only 4 days, with a quarter of beetles living more than 6 days, 5 % (15
beetles) more than 13 days and 1 % (4 beetles) between 38 and 56 days. Beetles that
survived the longest despite repeated trials had a noticeably high feeding activity. Lifespan
values are to be taken with caution since the age of individuals at the time of their capture
remains unknown.
Flight distance distribution also varied greatly over the lifespan of the tested insects. The
median flight distance (Table 2) for *Agrilus* and *Meliboeus* was 53 m (115 m excluding the
28 % of non-flying insects); a quarter of the beetles flew more than 243 m (401 m), 10 %
more than 730 m (1 039 m), and 1 % (3 beetles) flew between 2 890 m and 4 636 m (2 916 m
and 4 636 m). Distances were longer in the seven *C. undatus*: median distance 967 m with
one exceptional beetle above 16 km. Flown distance was globally linked with the number of
trials (LM, $p < 0.001$, $R^2 = 0.42$, *C. undatus* excluded); no significant variation with species
or sex had been observed.
280

281 **Table 2**. Median, mean and maximum total distance flown by the insects during their lifespan. First trial and Sum correspond to the data of the
282 first eight-hour trial and the sum of all trials respectively.

| Species | Median total flight distance (m) | | | | Mean total flight distance (m) | | | | Highest total flight distance (m) | |
| | With non-flying individuals | | Without non-flying individuals | | With non-flying individuals | | Without non-flying individuals | | | |
	First trial	Sum	First trial	Sum	First trial	Sum	First trial	Sum	First trial	Sum
All with *C. undatus*	40	54	95	119	200 ± 28	356 ± 74	290 ± 38	487 ± 100	3 028	16 097
All without *C. undatus*	35	53	87	115	167 ± 22	274 ± 38	245 ± 31	379 ± 51	2 246	4 636
A. angustulus	70	71	71	92	186 ± 100	205 ± 92	207 ± 109	225 ± 99	1 048	1 048
A. graecus	738	738	738	738	738	738	738	738	738	738
A. graminis	74	149	87	342	98 ± 39	663 ± 500	147 ± 47	995 ± 730	350	4 636
A. hastulifer	143	182	212	212	245 ± 90	486 ± 189	300 ± 101	531 ± 202	910	2 135
A. hyperici	25	56	316	460	341 ± 91	671 ± 152	546 ± 130	916 ± 189	2 246	2 989
A. laticornis	35	53	67	95	121 ± 21	144 ± 21	169 ± 27	195 ± 26	1 132	1 132
A. obscuricollis	0	0	144	144	58 ± 41	58 ± 41	144 ± 64	144 ± 64	208	208
A. olivicolor	1	1	27	35	26 ± 11	28 ± 10	53 ± 19	52 ± 15	170	170
A. sulcicollis	68	68	68	68	141 ± 96	141 ± 96	141 ± 96	141 ± 96	426	426
A. viridis	364	782	428	894	367 ± 153	771 ± 277	489 ± 130	964 ± 256	739	1 644
C. undatus	966	966	1 070	966	1 205 ± 410	3172 ± 2 168	1 405 ± 424	3172 ± 2 168	3 028	16 097
M. fulgidicollis	0	0	61	61	72 ± 60	79 ± 69	168 ± 131	211 ± 174	429	559

284 General flight characteristics among species

285   Comparisons among species were done on first (eight-hour) trial data, which allowed to
286 include as many insects as possible without having to manage repeated measurements. All

Figure 2. Values for a selection of response variables measured during the first trial
(eight hour), per sex and species. Females: left (green), males: right (purple).

species and sexes combined, 222 individuals (123 females, 99 males) have been analysed,
taking into account the removal of individuals for which some issue occurred during the trial.
Main results are displayed in figure 2. Preflight mass differed significantly among species
(Kruskal-Wallis, $\chi^2(7) = 67.4$, p < 0.001, n = 208). According to Dunn's test, *C. undatus*,
A. hastulifer and *A. graminis* were significantly heavier than the other species tested
(Figure 2). Even if *C. undatus* was not significantly heavier than *A. hastulifer* and *A. graminis*,
presumably because small sample size and high inter-individual variation, this species was
clearly out of category in terms of size and mass.
Regardless of the species, flight distance distribution during the first trial was highly
variable: 70 insects (31 %) did not fly, 44 % (98 beetles) flew between 1 m and 211 m, 25 %
(56 beetles) flew more than 211 m, 10 % (23 beetles) more than 596 m, 5 % (12 beetles)
more than 948 m, and 1 % (3 beetles) more than 2 138 m (Figure 3). The longest total
distance flown within an eight-hour trial was 3 028 m for a female *C. undatus*, 2 246 m for a
female *A. hyperici*, and 429 m for a male *M. fulgidicollis*. Remarkably, some individuals of
C. undatus (two females) and *A. hyperici* (one female, four males) flew distances long enough
to stand out from the right tail of the distributions, even after their first trial, and may be
considered extreme (Figure 3). Despite the exceptional performances of a female *C. undatus*,
no significant difference has been observed between sexes. The flight speed of *Coraebus* and
other species showed low variation and was respectively 0.40 m/s $\pm$ 0.03 m/s and
0.26 m/s $\pm$ 0.13 m/s.
Regardless of their species, most tested individuals performed several short flight bouts
within a single eight-hour trial (Figure 2). Flight bout number was highly variable: 25 or less
for half of trials with flight activity, but more than 58 for a quarter (39 trials), and 16 trials
(10 %) overreaching this number with more than 150 flight bouts over eight hours. Three
trials even exceeded 300 flight bouts: two by females *A. hyperici* (407 and 379 flight bouts)
and one by a female *A. laticornis* (308 flight bouts). Flight bout duration was also highly
variable (median 12 s, 10 % above 36 s, up to 1 406 s), so was the corresponding flown
distance (median 2 m, 10 % above 9.5 m, up to 653 m).

Correlations among variables

Considering the large body size of *C. undatus* compared to other species, global (*i.e.,* non-
species-specific) correlations among variables were analysed with and without *C. undatus* in
the pool of species (Figure 4 and Supplementary Figure S3). In both cases, strong correlations
were found between distance variables (total flight distance, mean flight bout distance) and
their duration counterparts (total flight duration, mean flight bout duration). Considering
these strong correlations, duration variables were discarded from further analyses and
emphasis was put on distance variables instead.
When *C. undatus* was included in the multidimensional clustering of variables, preflight
mass clustered tightly with total flight distance, while the association became looser when
excluding *C. undatus* from the pool. In addition, when *C. undatus* was excluded, preflight mass
correlated with mean flight bout distance. The number of flight bouts was correlated with
total flight duration or distance only when *C. undatus* was excluded, suggesting that other
species performed multiple flights while *C. undatus* showed sustained flight bouts. With or
without *C. undatus*, the latency to the first flight bout was inversely correlated with total flight
distance.

Figure 3. Distribution of total distance flown per trial depending on sex, all 12 species pooled together ($n_{females} = 123$, $n_{males} = 99$). Darker bars represent the first trial (eight hour) while lighter bars represent all consecutive trials (four hours), cumulated on top of the first trial. Individual flights $\geq$ 1300 m are marked with an arrow indicating the corresponding species, with the trial number and individual identifier in brackets.

Figure 4. Correlation network among a selection of response variables measured for the first (eight-hour) trials, with all 12 species pooled together (left) and all but *C. undatus* (right). Highly correlated variables appear closer to each other, the proximity of points being determined by multidimensional clustering using a Principal Coordinates Analysis with *stats::cmdscale((1 - abs(correlations)), k = 2)* in *R* (see corr package; Kuhn et al., 2022) and are joined by stronger paths. Path colour is determined by the correlation coefficient and sign, and correlations whose absolute value is below 0.2 are not shown.

Given that species may show different flight strategies and are not equally represented in the sample, correlations with preflight mass were further analysed within the eight most numerous species only (Figure 5). In all species but *A. olivicolor*, higher preflight mass negatively impacted the latency to the first flight bout, in other words it increased the

propensity to take off, or showed no relationship. This effect did not appear clearly in the multivariate analysis (Figure 4 and Suplementary Figure S3), but could have been concealed by near-zero correlation coefficients in the two most abundant species, *A. laticornis* and *A. hyperici*. Mixed effects with mean flight bout distance, total flight distance and number of flight bouts were found. Noticeable positive correlations were found for preflight mass with the former two variables in *C. undatus* only, and with the latter two variables in *M. fulgidicollis*, *A. hyperici*, *A. laticornis* and *A. hastulifer*. Some particularly high correlations coefficients found in *M. fulgidicollis* are, however, to be considered cautiously considering the low sample size and scarce active flights recorded in this species (four in total, see Figure 8), suggesting that such high coefficients resulted from a fraction of the individuals and could correspond to a biassed sample.

Figure 5. Correlation coefficients between preflight mass and a selection of response variables during the first (eight-hour) trial, sorted by increasing value, in the eight most abundant species in the samples.

Flight performance

Two flight performance variables measured during the first (eight-hour) trial varied significantly among species (sex pooled): total flight distance (Kruskal-Wallis, $\chi^2(7) = 20.4$, $p = 0.005$, $n = 208$) and mean distance flown per flight bout (Kruskal-Wallis, $\chi^2(7) = 23$, $p = 0.002$, $n = 208$). However, Dunn tests highlighted only few differences among groups: between *A. olivicolor* and *C. undatus* for total flight distance, between *A. hastulifer* and *A. laticornis* for mean distance flown per flight bouts. Move latency ($\chi^2(7) = 8.46$, $p = 0.29$, $n = 208$) and number of flight bouts ($\chi^2(7) = 13.7$, $p = 0.06$, $n = 208$) were not influenced by species.

Intraspecific differences in the three most abundant species (*A. hyperici*, *A. laticornis* and *A. olivicolor*) demonstrated that endogenous factors influenced some of their flight capacities. Preflight mass positively affected total flight distance, as well as the number of flight bouts in *A. hyperici* and *A. laticornis*, but not in *A. olivicolor* (Table 3). Response variables showed no sex bias, as either fixed effects had no significant influence, or null models were better. However, marginal R^2 were weak and other factors likely have impacted flight performance. Only *A. hyperici* showed standing out marginal R^2 with number of moves ($R^2 = 0.23$) and total flight length ($R^2 = 0.27$) while every other R^2 did not exceed 0.13.

Table 3. LMER tests results for the effects of sex and preflight mass on a selection of response variables during the first (eight-hour) trial, with year of experiment as random effect. Response variables in each Linear Mixed-Effects Models were log-transformed. The reference level for the categorial factor 'sex' is female. ΔAICc = AICc (full model) – AICc (null model). The full model was considered to better explain the response variable than the null model when ΔAICc was lower than -2, and the opposite when ΔAICc was higher than -2. Conditional (Cond.) R^2 corresponds to the variance of both fixed and random effects while marginal (Marg.) R^2 considers only the variance of the fixed effect. *** : $p < 0.001$, ** : $p < 0.01$, * : $p < 0.05$

Species	Variables	ΔAICc	Sex (female) Estimate	SE	t value	Mass Estimate	SE	t value	Cond. R2	Marg. R2
A. hyperici	Total flight distance	-22.51	1.09	0.85	1.28	1 025.36 ***	264.91	3.87		0.28
	Number of flight bouts	-19.98	0.45	0.61	0.73	679.24 ***	189.78	3.58		0.25
	Mean flight bout distance	8.62	-	-	-	-	-	-	-	-
	Latency to first flight bout	-11.33	0.85	1.07	0.79	-308.19	369.73	-0.83	0.09	0.07
A. laticornis	Total flight distance	-17.81	0.81	0.44	1.85	461.98 **	150.70	3.07	0.14	0.09
	Number of flight bouts	-15.85	0.54	0.34	1.56	351.75 **	116.63	3.02	0.11	0.09
	Mean flight bout distance	-3.27	0.05	0.15	0.35	5.67	51.74	0.11	0.12	0.001
	Latency to first flight bout	-8.95	-0.30	0.61	-0.48	13.63	203.92	0.07		0.004
A. olivicolor	Total flight distance	-10.82	1.58	0.97	1.63	456.42	346.36	1.31	0.45	0.12
	Number of flight bouts	-9.36	1.13	0.67	1.71	294.03	238.59	1.23	0.42	0.13
	Mean flight bout distance	7.55	-	-	-	-	-	-	-	-
	Latency to first flight bout	1.67	-	-	-	-	-	-	-	-

Consecutive flight patterns

Flight performance varied over consecutive trials. To compare data from the first eight-hour trial with the subsequent four-hour trials, we truncated the former one to four hours. This resulted in only limited data discarded, as on average 83 % of total distance is flown during the first four hours. Unusual trial frequency data was discarded from analyses. Data on *C. undatus* were also excluded from these analyses, due to their specificities discussed below. Overall, raw flight performance decreased over consecutive trials (Figure 6 left, circle symbols; Kruskal-Wallis, $\chi^2(9) = 46.2$, $p < 0.001$, $n = 501$). Differences were mainly observed between the first trial and others. Extreme long distance flights ($> 1\,250$ m, but this threshold is arbitrary) only took place during the four first trials (and mainly in the first one). Flight performance increased in the late trials, when insects benefited from longer rest periods, but remained before first trials one (Figure 6 left, + symbols). Individual patterns over time were analysed using flight distance standardised by mean performance of each beetle. For each individual, the standardised flight distance showed large variations between successive trials (Supplementary Figure S2). However, the global distribution showed that beetles flew a greater distance on their first trial, with no variation between subsequent trials (Figure 6 right; Kruskal-Wallis, $\chi^2(9) = 140$, $p < 0.001$, $n = 501$).

Figure 6. Evolution of flight performance during consecutive trials. Left: Distribution of
distance flown during four-hour trials across consecutive trials. Right: Distribution of
relative flight performance across consecutive trials; for each beetle and trial, relative
flight performance was computed as the flight distance during the trial divided by the
mean flight distance during all trials. For both plots, data of the eight-hour trial was
truncated to the first four hours. Unusual trial frequency data and data from *C. undatus*
were discarded from analyses; however, late trial data are displayed on the left panel
(+ symbols). Different letters above the data indicate a significant difference according
to Dunn's test.

Variable flight patterns over time were observed among species, with *A. hyperici*,
A. graminis, and *C. undatus* standing out from the others because a fraction of their
individuals covered long distances after the first or even subsequent trials (Figure 7;
Figure 8). No such pattern was observed in another species, either due to a generally low
flight activity, or comparatively lower lifespan in rearing conditions, reducing the number of
consecutive trials per individual. Regardless of species, 35 % of individuals were dead after
the first trial, 70 % after the second and 86 % after the third. Most individuals (93 %) were
dead after four trials, and the remaining (7 %) were composed of individuals of the three
above-mentioned species (12 *A. hyperici*, 1 *A. graminis* and 1 *C. undatus*) as well as of 2
A. hastulifer. The vast majority of individuals that survived across multiple trials were unable
to sustain long flight distances over time (mainly feeding the "[0]" category in figures 7
and 8), such as *A. laticornis* and *A. hastulifer*. However, some individuals diverged from this
trend. A female *C. undatus* and a male *A. hyperici* flew long distances even after their first
flight mill trial, despite an experiment duration reduced to four hours in next trials (Figure 3;
Figure 7), as well as a female *A. hyperici* flying more than 800 m during its fifth trial (although
not indicated with an arrow in Figure 3; Figure 8). Remarkably, the extreme *C. undatus*
female flew several kilometres in five consecutive trials, cumulating over 16 km of active
flight after its capture (Figure 3). Finally, none of the individuals of the other species has
reached the fourth trial.

Figure 7. Individual flight trajectories across four consecutive trials in the three most numerous species (*A. laticornis*, *A. hyperici* and *A. olivicolor*), with the first trial spanning eight hours and the following four hours. Total flight distance is discretized in categories of increasing width to simplify individual trajectories. The NA category corresponds to missing data due to a technical issue during the trial (typically, a temporary escape of an individual).

Figure 8. Individual flight trajectories across four consecutive trials in the less numerous species with the first trial spanning eight hours and the following four hours. Total flight distance is discretized in categories of increasing width to simplify individual trajectories.

                                            Discussion

This study allowed a first analysis of flight characteristics of several European Agrilinae,
including flight performance and behaviour. It revealed several common patterns and some
specificities among species. A clear observation was the high variation in most flight
performance parameters among individuals. Prior history (including age and flight history)
is a likely factor for this variation, but is unknown, albeit probably only part of the picture,
since flight activity is typically known as an intrinsically variable trait among individuals in
other studies as well (Sauvard et al., 2018). Flight performance did not greatly vary during
consecutive trials, and remarkably, the best flyer in *C. undatus* was able to fly long distances
during several consecutive trials. High inter-individual variability is common in flight mill
experiments (Taylor et al., 2010; Sauvard et al., 2018). Such locomotor strategies with low
average dispersal rate but high inter-individual variability could help species to primarily
exploit local resources, easily accessible, but also to survive local disasters thanks to few
dispersers.
At the species level, the assessment of flight performance of several Agrilinae revealed
rather homogeneous flight capacities among species, under the experimental conditions of
tethered flights. Only the few *C. undatus* individuals presented substantial differences. Even
if their number was too low to exhibit many significant results, these beetles actually
performed some flights which largely exceeded the dispersal distance of any other species.
However, this species is also much bigger than others. The huge intraspecific variation and
the relatively low sample size, moreover unbalanced between species, had probably impeded
identifying specific flight characteristics. *C. undatus* excepted, the low variation in beetle size
among the sampled species and the low flight activity of most tested individuals might have
also reduced our ability to discriminate the results among species. Data available on flight
capacities of other *Agrilus* species in similar experiment indicates better performance:
median of 750 m for *A. planipennis* (Taylor et al., 2010) and mean between 200 to 1 800 m
for *Agrilus auroguttatus* (Schaeffer), depending on conditions (Lopez et al., 2014). These
values are rather similar to *C. undatus* performance, but, as this latter species, *A. planipennis*
and *A. auroguttatus* are also bigger than the *Agrilus* and *Meliboeus* species we tested. From
a more general perspective, Agrilinae performances are lower than those of some bigger
beetles such as *Monochamus galloprovincialis* (Olivier) (2 km per day; David et al., 2014)
and *Rhynchophorus ferrugineus* (Olivier) (2.6 km per day; Ávalos et al., 2014), and far from
those of some long-flight insects such as hornets (median about 20 km per day; Sauvard et
al., 2018).
At the trial level, most flying individuals showed no sustained flight, and instead
completed multiple short flight bouts. This flying pattern could correspond to "routine flight"
(Van Dyck & Baguette, 2005), typically associated with foraging or the search for a sexual
partner or shelter. In contrast, these authors described longer flights as "dispersal flights",
presumably allowing insects to escape from bad conditions or predators, and were almost
never observed during our experiment. However, opposing the two flight types may not be
relevant. The high correlation between the number of flight bouts and the total distance flown
by individuals highlighted that, instead of covering long distances at once, *Agrilus* species
could spread by performing a series of short flights (*e.g.,* Ávalos et al., 2014).
Globally, over the course of a day, tested beetles tended to be poor dispersers in the
experimental setting, with most of them not taking off or flying over very short ranges
(generally less than 200 m). Only *C. undatus* seemed to fly significantly longer distances
(median of 1 000 m). This observation may suggest resident strategies, *i.e.,* individuals that

tend to stay in the range of their emergence area. One third of beetles did not fly at all. This
is congruent with studies on other beetle species, which found a similar percentage of non-
flyers, such as for *Ips sexdentatus* Boern (Jactel, 1993) and *R. ferrugineus* (Ávalos et al.,
2014). These authors hypothesised that such a high proportion of non-flying insects might
be due to the tethered of the beetles and to the flight mill itself, or be a consequence of the
individual incapacity to fly. As some non-flying beetles in a trial also demonstrated being able
to fly during a later trial, not flying was probably due to a combination of behavioural
variations, propensity to take-off, and constraints of the flight mills, but not solely the latter.
Nevertheless, as flight performance distribution had a very long tail, some insects largely
exceeded the median performance of their conspecifics and could be considered good
dispersers. As such, they may perform better at reaching new distant habitats if enough
patches of viable hosts are available on their path (Lopez et al., 2014), which is normally the
case for oak-associated species in European temperate areas. Within *Agrilus* species, an
A. hyperici appeared to be the most active disperser; interestingly, this species was the only
one that was not associated with forest habitats in our sample, and their relatively high
performance could be related to its more scattered herbaceous habitat.
We expected a greater flight performance in females than males, as already demonstrated
with mated females of *A. planipennis* (Taylor et al., 2010). Yet, sex did not impact any flight
variables in the overall sample. Those results are concordant, however, with observations on
A. auroguttatus whose flight capacities were not affected by sex either (Lopez et al., 2014).
Lack of sexual dimorphism in flight performance seems then frequent in Agrilinae species.
Both sexes are then suspected to participate in dispersal in the tested species. Males are
known to use flight to actively search for females, while mated females typically disperse to
find a suitable host for egg laying (Chamorro et al., 2015), and both behaviours may
contribute to species dispersal. However, it is rather considered that females are the ones
which are able to disperse far away from their host when necessary while males stay close to
their natal or maturation area (Rodriguez-Soana et al., 2007; Mercader et al., 2009; Taylor et
al., 2010).
Insect body mass is also known to influence flight performance of insects. There is often a
positive correlation between this performance and the body mass, which is generally
considered to reflect the available fuel load (Minter et al., 2018). Such a relation was observed
in the majority of the species we tested, including the best flyer (*C. undatus*) and the two most
numerous species (*A. laticornis* and *A. hyperici*). Relation to body mass seemed to differ
between these groups; in *C. undatus*, heavier individuals tend to fly immediately and to
perform longer bouts; in the two *Agrilus*, heavier individuals tend, on the contrary, to
increase the number of bouts. In contrast to these species, performance of other ones,
including the third most numerous one (*A. olivicolor*), was not impacted by preflight mass,
which corroborates what was observed in some other *Agrilus* species, *i.e., A. auroguttatus*
(Lopez et al., 2014) and *A. planipennis* (Taylor et al., 2010). This difference could be related
to the relative importance of reserves vs. dead weight (mainly structure) in mass variation,
but also to insect strategy (noticeably, they could mainly fly when reserves are high, or, on
the contrary, when they are low, to forage in order to increase them). It should be noted,
however, that even when preflight mass was related to flight performance, it explains only a
small part of its variation. Other factors should be considered, and the effect of some of them,
diet, mated status and age, has been demonstrated in other *Agrilus* species (Taylor et al.,
2010; Lopez et al., 2014). Future studies should include these factors to further investigate
Agrilinae flight performance.

Flight mills under controlled conditions are a proven method to assess the flight potential
of insects. They are especially valuable to compare species or investigate the influence of
factors on flight activity. However, tethered flight performance recorded with this design
cannot be considered a direct measurement of free flight or natural flight (Ribak et al., 2017;
Sauvard et al., 2018; Naranjo, 2019), and must be taken with caution. Flight mill creates
artificial conditions and could both hinder (inertia) or promote (support) flight. The absence
of a moving natural landscape in the vision of the specimens is likely to influence their
locomotor behaviour, and the influence of sensed angular velocity or weightlessness on
tethered insects' propensity to fly is unknown. Tethered flight experiments have already
been suspected of under- or overestimating flight abilities due to the lack of tarsal contact
and the absence of a perch for taking off or landing (Minter et al., 2018) or because of the
device itself (Taylor et al., 2010). Yet, they remain a highly relevant method to compare
conspecifics or sympatric species in a given set of conditions and approach their motivation
to disperse under those conditions, or even physiological expenditures in flight dispersal
when their flight propensity is high. In other respects, flight mill likely limited the flight speed
in our experiment, as was found in another flight mill study on *A. planipennis* with a three-
fold difference between free and tethered flight speed (Taylor et al., 2010).

The sampling design based on interception traps allowed capturing multiple species
associated with oak forests, and not only oak borers. However, it prevented controlling for
additional factors pertaining to the prior experience of individuals (flight, diet, mated status)
and inherently biassed captures towards dispersing individuals. This may cause either (1)
undersampling of the most resident individuals, even more so than those tested in this study,
or (2) sampling of individuals that already spent energy on dispersal before the experiment.
Together with the limited number of captured individuals for some species and the
impossibility to control some factors, this warrants using a different sampling strategy in
further studies focusing on only oak-borers. We used dry collectors in green multi-funnel
traps which are a relevant method for Buprestidae (Allison & Redak, 2017). However, our
sampling design likely let a fraction of individuals escape even if the collector was emptied
every workday, possibly those showing the highest locomotor activity. Indeed, similar traps
but equipped with wet collectors were installed for another study nearby one of the sampling
sites, and captured a more abundant and richer community of beetles (A. Sallé, pers. comm.).
Emergence traps can yield very good results (*e.g.,* Taylor et al., 2010; Lopez et al., 2014 for
borers), but would have restricted the sample to borers of the equipped hosts only, instead
of targeting all circulating Buprestidae in oak forests.

An oak-borer species considered a key aggravating agent of oak decline in Europe is
missing in our sampling: *A. biguttatus*. This beetle is currently expanding its range with
climate change (Pederson & Jørum, 2009; Brown et al., 2015). and genetic molecular markers
highlighted population differentiation at the European scale (Le Souchu et al. submitted). Yet,
little is known about the dispersal ability of this species, let alone its variation among
populations. Flight mills may prove a relevant method to better understand the genetic
structure and movements of populations in this species, especially as it is considered a major
threat outside its current range (see for instance North America where its potential arrival is
closely monitored due to the danger its invasion would pose, USDA-APHIS, 2011). As the
biggest European *Agrilus* species, at a size similar to that of *A. planipennis* (Volkovitsh et al.,
2019) in which flight ability is described in most details among *Agrilus* species, comparing
both species would be insightful to better understand the spread potential of *A. biguttatus*.

Intrinsic dispersal potential of a flying insect species is generally mainly determined by
the overall flight capacity of its individuals during their life, more specifically that of its best

flyers. In Agrilinae (except perhaps *C. undatus*), this capacity was generally low in our experiment conditions (a quarter of non-flying insects, only a few percent of individuals above a kilometre). As flight performance did not greatly vary during their life, this capacity was highly dependent on lifespan. However, during this experiment, beetles proved to be difficult to maintain alive in rearing conditions, with a median lifespan of only four days. As a comparison, *A. planipennis* was reported to live between 2 and 17 weeks under laboratory conditions, with an average of 7 weeks for males and 9 weeks for females (Chamorro et al., 2015 and references within). Moreover, European Agrilinae lifespan is generally estimated at several weeks in natural conditions. Given our observations of limited leaf damage and faeces accumulation in the rearing vials, we suspect a failure to feed properly in the conditions they were offered, for unknown reasons. Combined effects of starvation and daily trials in flight mills likely impacted survival severely in the present experiment and dramatically reduced lifespan. The overall flight capacity we observed was then certainly underestimated. Even taking that into account, our study suggests that the overall flight capacity of small *Agrilus* and *Meliboeus* species is modest, with only a few percent of beetles flying some kilometres. Corresponding field observations are rare. *A. hyperici* was introduced in western Canada during the last century; even if data is imprecise, it seems to have largely spread in this region, but that took several decades, despite human help (Gov. of British Columbia, 2018). This is consistent with modest flight capacities of this species. Alongside their flight capacities, it is possible that small Agrilinae species occasionally disperse in natura using wind, as observed in some bark beetles, sometimes for several tens of kilometres (Nilssen, 1984). Moreover, Agrilinae actual dispersal could result from association of short-distance flight capacity and human-assisted transportation (noticeably with wood material), sometimes over long distances, as observed in *A. planipennis* (Muirhead et al., 2006), *Agrilus bilineatus* (Weber) (Baranchikov et al., 2019b) and related species (*e.g.,* Wu et al., 2017; Ruzzier et al., 2023).

Conclusion

This work provided a first comparative approach on several Agrilinae flight behaviour and performance. Despite experimental limitations, it showed several common traits among species, noticeably the huge intraspecific variations of flight performance, its low value, and its relative homogeneity among species. It also suggests the existence of several flight behavioural patterns among species.

Appendices

The supplementary material is available in the open repository Data Recherche Gouv at https://doi.org/10.57745/TY9KLG.

Acknowledgements

We are thankful for the technical assistance offered by Maxime Stanislawek, Clément Beaumont and Mathieu Plateau.

Funding

This work was supported by the Région Centre-Val de Loire Project no. 2018-00124136 (CANOPEE) coordinated by A. Sallé.

Conflict of interest disclosure

The authors declare no conflicts of interest and they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

The data and code that support the findings of this study are openly available in three repositories. The main one, at https://doi.org/10.57745/TY9KLG, includes the material specific to the present study: supplementary data, videos of insects flying in the rotational flight mills, raw log files from the flight mills, associated processed data and final R scripts. Git-versionned R scripts are also available at https://forgemia.inra.fr/mathieu.laparie/agrilusflight. The source of the Crystal program used to monitor flight mills and produce log files is available at https://doi.org/10.57745/DIWSWF. The Ruby pipelines used to process log files into flight data are available at https://doi.org/10.57745/YLMCNU.

References

Allison JD, Redak RA (2017) The Impact of Trap Type and Design Features on Survey and Detection of Bark and Woodboring Beetles and Their Associates: A Review and Meta-Analysis. *Annual Review of Entomology*, **62**, 127–146. https://doi.org/10.1146/annurev-ento-010715-023516

Ávalos JA, Martí-Campoy A, Soto A (2014) Study of the flying ability of *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) adults using a computer-monitored flight mill. *Bulletin of Entomological Research*, **104**, 462–470. https://doi.org/10.1017/S0007485314000121

Baranchikov Y, Björklund N, Bonifacio LF, Matthews-Berry S, Petrice T, Sallé A, Wang X-Y, Suffert M, Picard C (2019a) Pest risk Analysis for *Agrilus fleischeri* (Coleoptera: Buprestidae). https://doi.org/10.13140/RG.2.2.33002.75204

Baranchikov Y, Björklund N, Bonifacio LF, Matthews-Berry S, Petrice T, Sallé A, Wang X-Y, Suffert M, Picard C (2019b) Pest risk analysis for *Agrilus bilineatus* (Coleoptera: Buprestidae), two-lined chestnut borer. https://doi.org/10.13140/RG.2.2.19580.97926

Bílý S (2002) Summary of the bionomy of the Buprestid beetles of Central Europe (Coleoptera: Buprestidae). *Acta Entomological Musei Nationalis Pragae*, **10**, 104.

Bozorov TA, Luo Z, Li X, Zhang D (2019) *Agrilus mali* Matsumara (Coleoptera: Buprestidae), a new invasive pest of wild apple in western China: DNA barcoding and life cycle. *Ecology and Evolution*, **9**, 1160–1172. https://doi.org/10.1002/ece3.4804

Brown N, Inward DJG, Jeger M, Denman S (2015) A review of *Agrilus biguttatus* in UK forests and its relationship with acute oak decline. *Forestry*, **88**, 53–63. https://doi.org/10.1093/forestry/cpu039

Brück-Dyckhoff C, Petercord R, Schopf R (2019) Vitality loss of European beech (*Fagus sylvatica* L.) and infestation by the European beech splendour beetle (*Agrilus viridis* L., Buprestidae, Coleoptera). *Forest Ecology and Management*, **432**, 150–156. https://doi.org/10.1016/j.foreco.2018.09.001

Chamorro ML, Jendek E, Haack R, Petrice T, Woodley N, Konstantinov A, Volkovitsh M, Yang X-K, Grebennikov V, Lingafelter S (2015) *Illustrated Guide to the emerald ash borer*

Agrilus planipennis *Fairmaire and Related Species (Coleoptera, Buprestidae)*. Pensoft Publishers, Sofia, Bulgaria.

David G, Giffard B, Piou D, Jactel H (2014) Dispersal capacity of *Monochamus galloprovincialis*, the European vector of the pine wood nematode, on flight mills. *Journal of Applied Entomology*, **138**, 566–576. https://doi.org/10.1111/jen.12110

Dickinson MH, Lehmann F-O, Sane SP (1999) Wing Rotation and the Aerodynamic Basis of Insect Flight. *Science*, **284**, 1954–1960. https://doi.org/10.1126/science.284.5422.1954

Du Merle P, Attié M (1992) Coraebus undatus (Coleoptera: Buprestidae) sur chêne liège dans le Sud-Est de la France: estimation des dégâts, relations entre ceux-ci et certains facteurs du milieu. *Annales des sciences forestières*, **49**, 571–588.

Dubois GF, Vernon P, Brustel H (2009) A flight mill for large beetles such as Osmoderma eremita (Coleoptera: Cetoniidae). In: *Buse J, Alexander KNA, Ranius T, Assmann T, Saproxylic Beetles: Their Role and Diversity in European Woodland and Tree Habitats-Proceedings of the 5th Symposium and Workshop. Pensoft Publishers, Sofia*, pp. 219–224.

Dudley R (2002) *The biomechanics of insect flight: form, function, evolution*. Princeton Univ.y Press, Princeton, N.J.

Evans HF, Williams D, Hoch G, Loomans A, Marzano M (2020) Developing a European Toolbox to manage potential invasion by emerald ash borer (*Agrilus planipennis*) and bronze birch borer (*Agrilus anxius*), important pests of ash and birch. *Forestry: An International Journal of Forest Research*, **93**, 187–196. https://doi.org/10.1093/forestry/cpz074

Gov. of British Columbia (2018) *Agrilus hyperici (Creutzer)*. Canada.

Herms DA, McCullough DG (2014) Emerald Ash Borer Invasion of North America: History, Biology, Ecology, Impacts, and Management. *Annual Review of Entomology*, **59**, 13–30. https://doi.org/10.1146/annurev-ento-011613-162051

Hızal E, Arslangündoğdu Z (2018) The First Record of Two-Lined Chestnut Borer *Agrilus bilineatus* (Weber, 1801) (Coleoptera: Buprestidae) from Europe. *Entomological News*, **127**, 333–335. https://doi.org/10.3157/021.127.0404

Jactel H (1993) Individual variability of the flight potential of *Ips sexdentatus* Boern. (Coleoptera: Scolytidae) in relation to day of emergence, sex, size, and lipid content. *The Canadian Entomologist*, **125**, 919–930. https://doi.org/10.4039/Ent125919-5

Jendek E, Grebennikov V (2009) *Agrilus sulcicollis* (Coleoptera: Buprestidae), a new alien species in North America. *The Canadian Entomologist*, **141**, 236–245. https://doi.org/10.4039/n09-021

Jendek E, Grebennikov VV (2023) Summary of native geographic distribution of all 3,341 species of the most speciose animal genus Agrilus (Coleoptera: Buprestidae). *Journal of Insect Biodiversity*, **39**. https://doi.org/10.12976/jib/2023.39.2.1

Jendek E, Poláková J (2014) *Host Plants of World* Agrilus *(Coleoptera, Buprestidae)*. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-319-08410-7

Kuhn M, Jackson S, Cimentada J (2022) *corrr: Correlations in R*.

Le Souchu E, Bouget C, Sallé A (2024) Environmental drivers of local and temporal variations in the community of oak-associated borers (Coleoptera: Buprestidae). *European Journal of Forest Research*. https://doi.org/10.1007/s10342-023-01644-y

Lopez VM, McClanahan MN, Graham L, Hoddle MS (2014) Assessing the Flight Capabilities of the Goldspotted Oak Borer (Coleoptera: Buprestidae) With Computerized Flight Mills. *Journal of Economic Entomology*, **107**, 1127–1135. https://doi.org/10.1603/EC13525

Macháčová M, Nakládal O, Samek M, Baťa D, Zumr V, Pešková V (2022) Oak Decline Caused
by Biotic and Abiotic Factors in Central Europe: A Case Study from the Czech Republic.
Forests, **13**, 1223. https://doi.org/10.3390/f13081223
Mercader RJ, Siegert NW, Liebhold AM, McCullough DG (2009) Dispersal of the emerald ash
borer, *Agrilus planipennis*, in newly-colonized sites. *Agricultural and Forest Entomology*,
11, 421–424. https://doi.org/10.1111/j.1461-9563.2009.00451.x
Minter M, Pearson A, Lim KS, Wilson K, Chapman JW, Jones CM (2018) The tethered flight
technique as a tool for studying life-history strategies associated with migration in insects:
Tethered flight for studying insect movement. *Ecological Entomology*, **43**, 397–411.
https://doi.org/10.1111/een.12521
Muirhead JR, Leung B, Van Overdijk C, Kelly DW, Nandakumar K, Marchant KR, MacIsaac HJ
(2006) Modelling local and long-distance dispersal of invasive emerald ash borer *Agrilus*
planipennis (Coleoptera) in North America. *Diversity and Distributions*, **12**, 71–79.
https://doi.org/10.1111/j.1366-9516.2006.00218.x
Naranjo SE (2019) Assessing Insect Flight Behavior in the Laboratory: A Primer on Flight Mill
Methodology and What Can Be Learned. *Annals of the Entomological Society of America*,
112, 182–199. https://doi.org/10.1093/aesa/say041
Nilssen A (1984) Long-range aerial dispersal of bark beetles and bark weevils (Coleoptera,
Scolytidae and Curculionidae) in northern Finland. In: *Annales Entomologici Fennici* , pp.
37–42. Entomological Society of Finland.
Orlova-Bienkowskaja MJ, Drogvalenko AN, Zabaluev IA, Sazhnev AS, Peregudova EY, Mazurov
SG, Komarov EV, Struchaev VV, Martynov VV, Nikulina TV, Bieńkowski AO (2020) Current
range of *Agrilus planipennis* Fairmaire, an alien pest of ash trees, in European Russia and
Ukraine. *Annals of Forest Science*, **77**, 29. https://doi.org/10.1007/s13595-020-0930-z
Pederson H, Jørum P (2009) The jewel beetle *Agrilus biguttatus* (Fabricius, 1777) found in
Denmark (Coleoptera, Buprestidae). *Entomologiske Meddelelser*, **77**, 19–26.
R Core Team (2022) *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.
R Core Team (2023) *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.
Rassati D, Marini L, Marchioro M, Rapuzzi P, Magnani G, Poloni R, Di Giovanni F, Mayo P,
Sweeney J (2019) Developing trapping protocols for wood-boring beetles associated with
broadleaf trees. *Journal of Pest Science*, **92**, 267–279. https://doi.org/10.1007/s10340-
018-0984-y
Reed K, Denman S, Leather SR, Forster J, Inward DJG (2018) The lifecycle of *Agrilus*
biguttatus : the role of temperature in its development and distribution, and implications
for Acute Oak Decline. *Agricultural and Forest Entomology*, **20**, 334–346.
https://doi.org/10.1111/afe.12266
Renault D, Angulo E, Cuthbert RN, Haubrock PJ, Capinha C, Bang A, Kramer AM, Courchamp F
(2022) The magnitude, diversity, and distribution of the economic costs of invasive
terrestrial invertebrates worldwide. *Science of The Total Environment*, **835**, 155391.
https://doi.org/10.1016/j.scitotenv.2022.155391
Ribak G, Barkan S, Soroker V (2017) The aerodynamics of flight in an insect flight-mill (R
Gurka, Ed,). *PLOS ONE*, **12**, e0186441. https://doi.org/10.1371/journal.pone.0186441
Robinet C, Imbert C-E, Rousselet J, Sauvard D, Garcia J, Goussard F, Roques A (2012) Human-
mediated long-distance jumps of the pine processionary moth in Europe. *Biological*
Invasions, **14**, 1557–1569. https://doi.org/10.1007/s10530-011-9979-9

Rodriguez-Soana CR, Miller JR, Poland TM, Kuhn TM, Otis GW, Turk T, Ward DL (2007)
Behaviors of Adult *Agrilus planipennis* (Coleoptera: Buprestidae). *The Great Lakes*
Entomologist, **40**. https://doi.org/10.22543/0090-0222.2168

Ruzzier E, Haack RA, Curletti G, Roques A, Volkovitsh MG, Battisti A (2023) Jewels on the go:
exotic buprestids around the world (Coleoptera, Buprestidae). *NeoBiota*, **84**, 107–135.
https://doi.org/10.3897/neobiota.84.90829

Sallé A, Nageleisen L-M, Lieutier F (2014) Bark and wood boring insects involved in oak
declines in Europe: Current knowledge and future prospects in a context of climate
change. *Forest Ecology and Management*, **328**, 79–93.
https://doi.org/10.1016/j.foreco.2014.05.027

Sauvard D, Imbault V, Darrouzet É (2018) Flight capacities of yellow-legged hornet (Vespa
velutina nigrithorax, Hymenoptera: Vespidae) workers from an invasive population in
Europe (W Blenau, Ed,). *PLOS ONE*, **13**. https://doi.org/10.1371/journal.pone.0198597

Taylor RAJ, Bauer LS, Poland TM, Windell KN (2010) Flight Performance of *Agrilus*
planipennis (Coleoptera: Buprestidae) on a Flight Mill and in Free Flight. *Journal of Insect*
Behavior, **23**, 128–148. https://doi.org/10.1007/s10905-010-9202-3

USDA-APHIS (2011) New pest response guidelines: exotic wood-boring and bark beetles.

Van Dam WA, Williams RN, Taylor RAJ (2000) Flight performance of some nitidulid beetles
(Coleoptera) using a computer-monitored flight mill. *J. Agric. Urban Entomol.*, **17**, 143–
151.

Van Dyck H, Baguette M (2005) Dispersal behaviour in fragmented landscapes: Routine or
special movements? *Basic and Applied Ecology*, **6**, 535–545.
https://doi.org/10.1016/j.baae.2005.03.005

Volkovitsh MG, Kovalev AV, Orlova-Bienkowskaja MJ (2020) Current Distribution and
Diagnostic Features of Two Potentially Invasive Asian Buprestid Species: *Agrilus mali*
Matsumura and *A. fleischeri* Obenberger (Coleoptera: Buprestidae). *Insects*, **11**, 493.
https://doi.org/10.3390/insects11080493

Volkovitsh MG, Orlova-Bienkowskaja MJ, Kovalev AV, Bieńkowski AO (2019) An illustrated
guide to distinguish emerald ash borer (*Agrilus planipennis*) from its congeners in Europe.
Forestry: An International Journal of Forest Research, cpz024.
https://doi.org/10.1093/forestry/cpz024

Wu Y, Trepanowski NF, Molongoski JJ, Reagel PF, Lingafelter SW, Nadel H, Myers SW, Ray AM
(2017) Identification of wood-boring beetles (Cerambycidae and Buprestidae)
intercepted in trade-associated solid wood packaging material using DNA barcoding and
morphology. *Scientific Reports*, **7**, 40316. https://doi.org/10.1038/srep40316

Upper magnet in contact with the pin tip

Detection magnet

Hall effect sensors

Lower magnet not in contact with pinhead

Arm

Adhesive paste to hold the paper strip glued on the insect

Singleboard computer