Restoration of Type 17 immune signaling is not sufficient for protection during

influenza-associated pulmonary aspergillosis

Aijaz Ahmad[1], Ravineel Bhan Singh[1], Kara Nickolich[1], Matthew Pilewski[1], Caden Ngeow[1], Kwame Frempong-Manso[1], Keven Robinson[1]*

[1]Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, Department of Medicine, University of Pittsburgh, Pittsburgh, PA 15213, USA

Aijaz Ahmad: https://orcid.org/0000-0003-2845-0727

Keven Robinson: https://orcid.org/0000-0001-7558-4002

Running title: Type 17 immunity in influenza-associated pulmonary aspergillosis

***Corresponding Author**

Tel: +1 412 648 3098

robinsonkm@upmc.edu; kmr82@pitt.edu

ABSTRACT

Influenza-associated pulmonary aspergillosis (IAPA) is a severe complication of influenza

infection that occurs in critically ill patients and results in higher mortality compared to

influenza infection alone. Interleukin-17 (IL-17) and the Type 17 immune signaling pathway

cytokine family are recognized for their pivotal role in fostering protective immunity against

various pathogens. In this study, we investigate the role of IL-17 and Type 17 immune

signaling components during IAPA. Wild-type mice were challenged with influenza A H1N1

(Flu) and then exposed to *Aspergillus fumigatus* ATCC42202 resting conidia on day 6 post-

influenza infection, followed by the quantification of cytokines and chemokines at 48 hours

post-fungal infection. Gene and protein expression levels revealed that IL-17 and Type 17

immune cytokines and antimicrobial peptides are downregulated during IAPA compared to

mice singularly infected solely with *A. fumigatus*. Restoration of Type 17 immunity was not

sufficient to provide protection against the increased fungal burden observed during IAPA.

These findings contrast those observed during post-influenza bacterial super-infection, in

which restoration of Type 17 immune signaling protects against exacerbation seen during

super-infection. Our study highlights the need for future studies to understand the immune

mechanisms that increase susceptibility to fungal infection.

Keywords: Interleukin-17; IAPA; antimicrobial peptides; *Aspergillus fumigatus*

Importance

IAPA significantly elevates the risk of mortality in patients with severe influenza. Type-17

immunity is critical to host defense during fungal infections and, therefore, vital to

understand its role during IAPA. The observations in this study reveal that Type 17 immunity

is impaired during IAPA, potentially increasing susceptibility to secondary infection with

Aspergillus fumigatus. However, restoration of IL-17 signaling alone is not sufficient to reduce fungal burden in our murine IAPA model. These observations differ from those observed in post-influenza bacterial super-infections, suggesting that the mechanisms underlying viral-fungal super-infection are different than those that underly viral-bacterial super-infection. By elucidating the complex interactions between the host immune system, influenza, and *A. fumigatus*, these findings are vital for developing strategies to enhance immune responses and improve survival rates during IAPA.

INTRODUCTION

Influenza-associated pulmonary aspergillosis (IAPA) is a severe complication of influenza infection that occurs in critically ill patients and results in higher mortality compared to influenza infection alone [1]. The pathology of IAPA manifests through invasive growth of *Aspergillus fumigatus* within the lungs during influenza infection [2]. Influenza virus damages the respiratory epithelium, compromising the barrier function and allowing opportunistic fungi such as *A. fumigatus* to invade and colonize the lung tissue. The dysregulated immune state of the host, caused by the viral infection, provides an optimal environment for fungal growth and dissemination.

The interleukin-17 (IL-17) cytokine family is a pivotal component in mediating protective immunity against various pathogens [3]. IL-17, primarily produced by a subset of T helper cells known as Th17 cells, has been implicated in the immune response against extracellular pathogens. IL-17 exerts its immunomodulatory effects by promoting the synthesis of pro-inflammatory molecules, including cytokines, chemokines, and antimicrobial peptides. These molecules collectively facilitate the recruitment and activation of neutrophils and other immune effectors to sites of infection, thereby augmenting the host defense against invading pathogens [4].

Despite the well-established role of IL-17 in immunity against various pathogens, its specific

involvement in the context of IAPA remains understudied. Therefore, understanding the

immune response in this context is vital for improving therapeutic strategies and patient

outcomes. Our current study investigates the role of IL-17 and IL-17-related cytokines in

IAPA, employing a murine model to delineate their impact on disease pathogenesis and

immune responses. Our observation sheds light on the complex host immune system that

occurs during IAPA, aiming to uncover novel therapeutic avenues and enhance patient

outcomes in managing IAPA.

MATERIALS AND METHODS

Animals: Six to eight-week-old male C57BL/6 mice were purchased from Taconic Farms

(Germantown, NY). The mice were kept in a pathogen-free environment and co-housed in the

same facility before the commencement of the studies. All animal studies were performed

according to the protocol for the care and use of animals sanctioned by the University of

Pittsburgh Institutional Animal Care and Use Committee. All the studies used age- and sex-

matched mice.

Pathogens and superinfection model: Influenza A/PR/8/34 H1N1 was propagated in

chicken eggs as previously described [5] or by using Madin-Darby canine kidney (MDCK)

cells. The cells were maintained in DMEM with 10% FBS (Bio-Techne, Minneapolis, MN),

penicillin (100 U/ml), streptomycin (100 ug/ml) (Invitrogen, Waltham, MA). The cells were

washed with PBS, and infected 0.001 MOI of influenza virus A/Puerto Rico/8/1934 (H1N1)

in DMEM with 0.2% bovine serum albumin (Invitrogen, Waltham, MA), and 2 µg/ml of L-

tosylamido-2-phenyl ethyl chloromethyl ketone (TPCK) (Sigma-Aldrich, MO). The virus

containing supernatant was harvested after 72 hours and the viral titer was determined by

standard plaque assay. Mice were infected with 100 PFU of influenza A/PR/8/34 H1N1 (in 50

μl sterile PBS) from a frozen stock or control PBS by oropharyngeal aspiration. Infected mice

were incubated for 6 days, at which time mice received 2.5×10^7 conidia of *A. fumigatus*

ATCC42202 inoculum or PBS control. At 48 hours post-fungal infection, all the mice were

euthanized to harvest lungs for further studies.

111
Lung inflammation analysis: After harvesting, mouse lungs were lavaged with 1 ml of

sterile PBS to perform inflammatory cell differential counts. The upper lobe of the right lung

was homogenized in sterile PBS for counting fungal colonies and cytokine analysis,

conducted either with Lincoplex (Millipore, MO, USA) or ELISA assays (R&D Systems,

MN, USA), following the manufacturer's guidelines. The middle and lower lobes of the right

lung were snap-frozen and then homogenized under liquid nitrogen for RNA extraction using

the RNA isolation Kit (Agilent Technologies, TX, USA). The RNA analysis was carried out

via standard RT-qPCR employing Bio-Rad SSO advanced Universal Probes Supermix (CA,

USA). Gene expression analysis was performed from two replicate samples. It was calculated

using the formula $\Delta Cq=2^{Cq \text{ target gene-Cq reference gene}}$, where the quantitation cycle (Cq) was the

average Cq value of the target gene minus the *HPRT* reference gene's mean.

123
Flow cytometry: Flow cytometry analysis was conducted on the whole left lung. After

harvesting, the left lung underwent collagenase digestion, following a previously described

protocol [6]. The resulting single-cell preparations were *in vitro* stimulated with PMA (50

ng/ml) and ionomycin (750 ng/ml) for four h at 37 °C. Subsequently, cells were stained with

antibodies, fixed and permeabilized, and stained with fluorescent-conjugated antibodies (BD

Biosciences). The analysis used a Cytek Aurora™ CS System (Cytek® Biosciences

Bethesda, MD, USA).

131
132 **RESULTS**

133 **Type 17 cytokines are inhibited during IAPA**

134 We have previously published a murine model of IAPA that demonstrates increased morbidity

135 in mice co-infected with influenza and *A. fumigatus* [7]. Type 17 immunity plays a critical

136 role in host defense against *A. fumigatus* and other fungal pathogens [8-12]. IL-17 and other

137 Type 17 immune cytokines also play a critical role in the development of bacterial super-

138 infection during influenza [13-14]. Therefore, we hypothesized that the Type 17 immune

139 response would play an essential role during IAPA. To model this hypothesis, C57BL/6J male

140 mice were challenged with a sublethal dose of influenza A PR/8/34 H1N1 (100 PFU) for 6

141 days, followed by 2.5×10^7 □*A. fumigatus* (ATCC strain 42202) conidia, and after 48 hours,

142 fungal burden was assessed. Mice super-infected with influenza and *A. fumigatus* had

143 decreased expression of IL-17, IL-22, IL-23, and IL-1β compared to those infected with *A.*

144 *fumigatus* alone (**Fig 1A**). This downregulation could potentially contribute to increased

145 morbidity, mortality, and fungal burden compared to singular infection with *A. fumigatus*

146 alone. In addition to the gene expression changes, protein expression analysis also showed a

147 decreased production of IL-17, IL-22, IL-23, and IL-1β proteins in super-infected mice (**Fig**

148 **1B**).

149

150 **Decreased IL-17-producing γδ T cells during IAPA**

151 Flow cytometry analysis was conducted to quantify IL-17-producing T cells in our model. We

152 measured IL17$^+$ CD4$^+$ T and γδ T cells in our model due to their well-established role as

153 primary producers of IL-17. We observed a notable decrease in the total number of IL-17

154 producing γδ T cells in super-infected mice compared to singular *A. fumigatus* infection (**Fig**

155 **1C**). Comparatively, we observed no difference in the abundance of IL17$^+$ CD4$^+$ T cells

156 between the different groups (**Fig 1C**). When interpreting these results, it is important to

157 consider the timing of mouse harvesting post- *A. fumigatus* infection. Specifically, the

analysis was conducted at a 48-hour time point post-infection. At this early stage, the immune

response is still evolving, with distinct subsets of immune cells playing varying roles. In

particular, it is noted that there were more innate γδ T cells present at this time point, with T

cells potentially infiltrating at later stages of infection.

Reduced expression of IL-17/IL-22 associated antimicrobial peptides in IAPA

As IL-17 and IL-22 are distinct lineages of Type 17 cells [15], we studied their effector

function by examining gene expression levels of IL-17- and IL-22- associated antimicrobial

peptides in our model. The expression levels of *Reg3β*, *Reg3γ, and Lcn2* were significantly

reduced in super-infected mice compared to those infected solely with *A. fumigatus* (**Fig 1D**).

This reduction of IL-17- and IL-22- associated antimicrobial peptides may play a role in

fungal clearance during IAPA.

Restoration of Type 17 immunity does not enhance fungal clearance during IAPA

With the observation that Type 17 cytokines and antimicrobial peptides were significantly

downregulated during IAPA compared to *A. fumigatus* alone, we hypothesized that their

restoration could provide protection during IAPA. To test this hypothesis, we overexpressed

IL-17 proposing that it would rescue Type-17 immunity and enhance fungal clearance.

Overexpression of IL-17 significantly upregulated the levels of IL-17a mRNA (**Fig 2B**) and

downstream inflammatory mediators TNF and CXCL1 during IAPA (**Fig 2C**) [16,17];

however, fungal burden remained unchanged **(Fig 2A)**. Interestingly, the total number of

inflammatory cells in the bronchoalveolar lavage fluid and the total numbers of neutrophils

and macrophages measured by cytospin differential also remained unchanged with

upregulation of Type 17 immunity (**Fig S1**). Additionally, we administered murine

recombinant IL-17 protein during IAPA and fungal burden was unchanged (**Fig S2**). Next, we

restored components of the Type 17 immune signaling pathway that are both upstream and

downstream of IL-17 to determine effects on fungal clearance. Since both IL-1β and IL-23

are known to induce IL-17 production from γδ T cell [18], and we observed decreased

production during IAPA (**Fig 1A-B**), we administered IL-1β and IL-23+IL-1β together for

synergistic effects during IAPA. There was a significant increase of IL-17 mRNA in the mice

that received exogenous IL-1β and IL-23/IL-1β (**Fig 2E**); however, fungal burden remained

unchanged (**Fig 2D**). Additionally, IL-1β and IL-23+IL-1β did not alter IL-22 mRNA

expression (**Fig 2E**). Super-infected mice were administered either with Reg3β or Reg3γ to

explore potential therapeutic interventions. Although both these peptide levels have been

reduced in super-infected mice, neither changed fungal clearance during IAPA (**Fig 2F**).

Additionally, we observed no difference in the total cell count of immune cells in

bronchoalveolar lavage fluid (**Fig S3**). Collectively, these results indicate that restoration of

IL-17 signaling is not sufficient to restore fungal clearance during IAPA. These findings

suggest the involvement of mechanisms beyond Type 17 inhibition causing decreased fungal

clearance during IAPA and highlight the complexity of IAPA pathogenesis.

DISCUSSION

Although viral-fungal co-infections are associated with high mortality, limited data exists

regarding pathophysiology and lung immunology. Influenza and *A. fumigatus* have

individually been studied due to their clinical significance; however, the synergy and

complexities that arise when these pathogens co-exist within the same host remain poorly

understood. Invasive pulmonary aspergillosis was classically considered a disease of

immunocompromised patients; however, recent clinical observations have reported invasive

pulmonary aspergillosis following influenza infection (IAPA) in immunocompetent patients

[19]. The increased risk of developing aspergillosis in patients with preceding influenza can

be partially attributed to viral-induced epithelial disruption, the first line of host defense

against fungal infections. However, recent evidence suggests that second (phagocytosis and

the killing of *Aspergillus* conidia by phagocytes) and third lines (extracellular mechanisms,

mediated by neutrophils, to kill the *Aspergillus*) of the antifungal host responses are also

impaired in patients with IAPA [1,20]. IAPA has also been documented to provoke a severe

inflammatory response, resulting in a cytokine storm within lung tissue [21]. IAPA has been

described for decades and has been increasingly recognized since the 2009 H1N1 influenza

pandemic. Notably, *Aspergillus* species are also implicated in causing super-infection during

SARS-CoV-2 (COVID-19) infections [22,23], underscoring the importance of studying viral-

fungal super-infections. Additionally, studies suggest that the hyperinflammatory responses

driven by systemic cytokines in COVID-19 patients also contribute to CAPA [24].

The Type 17 cytokine family is recognized for its pivotal role in fostering protective

immunity against a spectrum of pathogens [25]. Previous research has substantiated the

involvement of IL-17 in the context of viral-bacterial superinfections [13]. The IL-17

pathway has also promoted *Aspergillus* clearance within pulmonary tissues [8-10].

Mechanistically, the protective role for IL-17 is mediated by the recruitment of neutrophils

through chemokine signaling and the production of antimicrobial peptides (AMP) production

[26]. This study aims to delineate the specific roles of IL-17 and Type 17 immune-associated

cytokines and chemokines in IAPA.

Our results demonstrate that preceding influenza infection impairs Type17 immunity

during IAPA. Both gene expression and protein quantification of key Type 17 immune

cytokines, particularly IL-17 and IL-22, are decreased during IAPA compared to singular

infection. IL-17, a pro-inflammatory cytokine, plays an essential role in fungal infections by

recruiting neutrophils and other immune cells to the site of infection and by inducing the

production of antimicrobial peptides [27]. Additionally, IL-17 synergistically collaborates

with IL-22 for a robust immune response against fungal infection [28]. The synergistic action

of IL-17 and IL-22 is crucial for a robust immune response, and their inhibition can lead to a

compromised ability to control fungal infections. Decreased IL-17 production was also seen

by Lee et al., using a similar murine model of IAPA; however, we also show a reduction in

other Type 17-immune associated cytokines and antimicrobial peptides [29]. IL-1β, a pro-

inflammatory cytokine critical to Type 17 immunity, is reduced in super-infection compared

to singular *A. fumigatus* challenge. IL-1β has also been reported to induce neutrophils and

macrophages recruitment to lungs during microbial invasion [7] and stimulates endothelial

adhesion molecules, different cytokines and chemokines, and the Th17 adaptive immune

response [30]. Our results align with the previous findings of downregulation of *IL1B* that

was observed in humans during IAPA [20]. Furthermore, in our murine model of IAPA, the

impaired production of IL-17 and IL-22 is also associated with reduced levels of

antimicrobial peptides, weakening the host's defense mechanisms. These findings highlight

that the downregulation of Type 17 immunity during IAPA compromises the host's defense

mechanisms and thereby increasing susceptibility to *Aspergillus* infections.

Despite the observed reduction in Type 17 immunity during IAPA, restoration of

various components of the IL-17 signaling pathway, both upstream and downstream of IL-17,

did not lead to improved fungal clearance. In contrast to prior studies that showed restoration

of Type 17 immune pathway components rescued bacterial clearance during post-influenza

bacterial super-infection [13,14,31], the current study indicates that restoration of IL-17

signaling alone is not sufficient to reduce fungal burden in a murine IAPA model. IL-17 may

play a protective role but not a restorative role during IAPA. Notably, the immune response is

dynamic during infection and augmentation of Type 17 signaling at other time points during

IAPA may produce different results. Interestingly, downregulation of IL-1β in our mouse

model is consistent with the findings in human patients with IAPA versus influenza alone

[20]. This consistency between our findings in mice and human patients strengthens the

validity of our results and suggests that the observed decrease in cytokines is a robust effect

that is relevant across species. Importantly, these studies show that immune regulation during

post-influenza fungal super-infection and post-influenza bacterial super-infection are not

driven by the same mechanisms. It underscores the need for additional studies to understand

the immune mechanisms that increase susceptibility to fungal infection during influenza and

how delineation of the specific cell types and immune pathways that are necessary for fungal

host defense during viral infection may lead to future therapeutics.

ETHICS APPROVAL

The University of Pittsburgh Institutional Animal Care and Use Committee approved the

study under Protocol No. 21063690.

ACKNOWLEDGMENTS

The authors appreciate the National Institute of Allergy and Infectious Diseases Grant R01

(Award No. R01AI153337 to KMR) and the National Heart, Lung and Blood Institute Grant

K08 (Award No. K08HL133445 Award No to KMR).

DATA AVAILABILITY

All the associated data is provided with this manuscript or in supplementary files.

DECLARATION OF INTEREST

The authors have no conflict of interest with any organization or entity with a financial

interest in the subject matters or materials discussed in this manuscript.

SUPPLEMENTAL MATERIAL

Supplemental Figures S1-S3.

REFERENCES

1. Schauwvlieghe AFAD, Rijnders BJA, Philips N, Verwijs R, Vanderbeke L, Van

Tienen C, Lagrou K, Verweij PE, Van de Veerdonk FL, Gommers D, Spronk P,

Bergmans DCJJ, Hoedemaekers A, Andrinopoulou ER, van den Berg CHSB,

Juffermans NP, Hodiamont CJ, Vonk AG, Depuydt P, Boelens J, Wauters J, Dutch-

Belgian Mycosis study group. 2018. Invasive aspergillosis in patients admitted to the

intensive care unit with severe influenza: A retrospective cohort study. Lancet Respir

Med 6(10): 782–792. https://doi.org/10.1016/S2213-2600(18)30274-1.

2. van de Veerdonk FL, Kolwijck E, Lestrade PPA, Hodiamont CJ, Rijnders BJA, van

Paassen J, Haas P-J, Oliveira Dos Santos C, Kampinga GA, Bergmans DCJJ, van Dijk

K, de Haan AFJ, van Dissel J, van der Hoeven HG, Verweij PE, Dutch Mycoses Study

Group. 2017. Influenza-associated aspergillosis in critically ill patients. Am J Respir

Crit Care Med https://doi.org/10.1164/rccm.201612-2540LE.

3. Kuwabara T, Ishikawa F, Kondo M, Kakiuchi T. 2017. The role of IL-17 and related

cytokines in inflammatory autoimmune diseases. Mediators Inflamm 2017:3908061.

https://doi.org/10.1155/2017/3908061

4. Korn T, Bettelli E, Oukka M, Kuchroo VK. 2009. IL-17 and Th17 Cells. Annu Rev

Immunol 27:485–517. https://doi.org/10.1146/annurev.immunol.021908.132710

5. Robinson KM, Ramanan K, Clay ME, McHugh KJ, Pilewski MJ, Nickolich KL,

Corey C, Shiva S, Wang J, Muzumdar R, Alcorn JF. 2018. The inflammasome

potentiates influenza/*Staphylococcus aureus* superinfection in mice. JCI Insight 3.

https://doi.org/10.1172/jci.insight.97470

6.   Robinson KM, Ramanan K, Clay ME, McHugh KJ, Rich HE, Alcorn JF. 2018. Novel

protective mechanism for interleukin-33 at the mucosal barrier during influenza-

associated   bacterial   superinfection.   Mucosal   Immunol   11:199–208.

https://doi.org/10.1038/mi.2017.32

7.   Tobin JM, Nickolich KL, Ramanan K, Pilewski MJ, Lamens KD, Alcorn JF,

Robinson KM. 2020. Influenza suppresses neutrophil recruitment to the lung and

exacerbates secondary invasive pulmonary aspergillosis. J Immunol 205:480–488.

https://doi.org/10.4049/jimmunol.2000067

8.   Werner JL, Gessner MA, Lilly LM, Nelson MP, Metz AE, Horn D, Dunaway CW,

Deshane J, Chaplin DD, Weaver CT, Brown GD, Steele C. 2011. Neutrophils produce

interleukin 17A (IL-17A) in a dectin-1- and IL-23-dependent manner during invasive

fungal infection. Infect Immun 79:3966–3977. https://doi.org/10.1128/iai.05493-11

9.   Taylor PR, Leal SM, Sun Y, Pearlman E. 2014. *Aspergillus* and *Fusarium* corneal

infections are regulated by Th17 cells and IL-17-producing neutrophils. J Immunol

192:3319–3327. https://doi.org/10.4049/jimmunol.1302235

10.   Gessner MA, Werner JL, Lilly LM, Nelson MP, Metz AE, Dunaway CW, Chan YR,

Ouyang W, Brown GD, Weaver CT, Steele C. 2012. Dectin-1-dependent interleukin-

22 contributes to early innate lung defense against *Aspergillus fumigatus*. Infect

Immun 80:410–417. https://doi.org/10.1128/iai.05939-11

11.   Sparber F, LeibundGut-Landmann S. 2019. Interleukin-17 in antifungal immunity.

Pathogens 8. https://doi.org/10.3390%2Fpathogens8020054

12.   Malacco NLS de O, Rachid MA, Gurgel IL da S, Moura TR, Sucupira PHF, de Sousa

LP, de Souza D da G, Russo R de C, Teixeira MM, Soriani FM. 2018. Eosinophil-

associated innate IL-17 response promotes *Aspergillus fumigatus* lung pathology.

Front Cell Infect Microbiol 8:453. https://doi.org/10.3389/fcimb.2018.00453

13.   Kudva A, Scheller EV, Robinson KM, Crowe CR, Choi SM, Slight SR, Khader SA,

Dubin PJ, Enelow RI, Kolls JK, Alcorn JF. 2011. Influenza A inhibits Th17-mediated

host defense against bacterial pneumonia in mice. J Immunol 186:1666–1674.

https://doi.org/10.4049/jimmunol.1002194

14.   Robinson KM, Choi SM, McHugh KJ, Mandalapu S, Enelow RI, Kolls JK, Alcorn

JF. 2013. Influenza A exacerbates *Staphylococcus aureus* pneumonia by attenuating

IL-1β   production   in   mice.   J   Immunol   191:5153–5159.

https://doi.org/10.4049/jimmunol.1301237

15.   Aujla SJ, Chan YR, Zheng M, Fei M, Askew DJ, Pociask DA, Reinhart TA,

McAllister F, Edeal J, Gaus K, Husain S, Kreindler JL, Dubin PJ, Pilewski JM,

Myerburg MM, Mason CA, Iwakura Y, Kolls JK. 2008. IL-22 mediates mucosal host

defense   against   Gram-negative   bacterial   pneumonia.   Nat   Med   14:275–281.

https://doi.org/10.1038/nm1710

16.   Wang CQF, Akalu YT, Suarez-Farinas M, Gonzalez J, Mitsui H, Lowes MA, Orlow

SJ, Manga P, Krueger JG. 2013. IL-17 and TNF synergistically modulate cytokine

expression while suppressing melanogenesis: potential relevance to psoriasis. J Invest

Dermatol 133:2741–2752. https://doi.org/10.1038%2Fjid.2013.237

17.   Sharma AK, Mulloy DP, Le LT, Laubach VE. 2014. NADPH oxidase mediates

synergistic effects of IL-17 and TNF-α on CXCL1 expression by epithelial cells after

lung ischemia-reperfusion. Am J Physiol Lung Cell Mol Physiol 306:L69-79.

https://doi.org/10.1152/ajplung.00205.2013

18.   Sutton CE, Lalor SJ, Sweeney CM, Brereton CF, Lavelle EC, Mills KHG. 2009.

Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells,

amplifying Th17 responses and autoimmunity. Immunity 31:331–341.

https://doi.org/10.1016/j.immuni.2009.08.001

19.   Shah MM, Hsiao EI, Kirsch CM, Gohil A, Narasimhan S, Stevens DA. 2018. Invasive

359   pulmonary aspergillosis and influenza co-infection in immunocompetent hosts: case

360   reports and review of the literature. Diagn Microbiol Infect Dis 91:147–152.

361   https://doi.org/10.1016/j.diagmicrobio.2018.01.014.

362   20.   Feys S, Gonçalves SM, Khan M, Choi S, Boeckx B, Chatelain D, Cunha C, Debaveye

363   Y, Hermans G, Hertoghs M, Humblet-Baron S, Jacobs C, Lagrou K, Marcelis L,

364   Maizel J, Meersseman P, Nyga R, Seldeslachts L, Starick MR, Thevissen K,

365   Vandenbriele C, Vanderbeke L, Vande Velde G, Van Regenmortel N, Vanstapel A,

366   Vanmassenhove S, Wilmer A, Van De Veerdonk FL, De Hertogh G, Mombaerts P,

367   Lambrechts D, Carvalho A, Van Weyenbergh J, Wauters J. 2022. Lung epithelial and

368   myeloid innate immunity in influenza-associated or COVID-19-associated pulmonary

369   aspergillosis: an observational study. Lancet Respir Med 10:1147–1159.

370   https://doi.org/10.1016/S2213-2600(22)00259-4

371   21.   Gonçalves SM, Pereira I, Feys S, Cunha C, Chamilos G, Hoenigl M, Wauters J, van

372   de Veerdonk FL, Carvalho A. 2024. Integrating genetic and immune factors to

373   uncover pathogenetic mechanisms of viral-associated pulmonary aspergillosis. MBio

374   15:e0198223. https://doi.org/10.1128/mbio.01982-23.

375   22.   Zhu X, Ge Y, Wu T, Zhao K, Chen Y, Wu B, Zhu F, Zhu B, Cui L. 2020. Co-

376   infection with respiratory pathogens among COVID-2019 cases. Virus Res

377   285:198005. https://doi.org/10.1016/j.virusres.2020.198005

378   23.   van Arkel AL, Rijpstra TA, Belderbos HN, Van Wijngaarden P, Verweij PE,

379   Bentvelsen RG. 2020. COVID-19–associated pulmonary aspergillosis. Am J Respir

380   Crit Care Med 202(1):132-5. https://doi.org/10.1164/rccm.202004-1038LE

381   24.   Arastehfar A, Carvalho A, van de Veerdonk FL, Jenks JD, Koehler P, Krause R,

382   Cornely OA, S Perlin D, Lass-Flörl C, Hoenigl M. 2020. COVID-19 Associated

Pulmonary Aspergillosis (CAPA)-From Immunology to Treatment. J Fungi (Basel) 6.

https://doi.org/10.3390/jof6020091.

25.   Huangfu L, Li R, Huang Y, Wang S. 2023. The IL-17 family in diseases: from bench

to bedside. Signal Transduct Target Ther 8:402. https://doi.org/10.1038/s41392-023-

01620-3.

26.   Mills KHG. 2023. IL-17 and IL-17-producing cells in protection versus pathology.

Nat Rev Immunol 23:38–54. https://doi.org/10.1038/s41577-022-00746-9.

27.   Puerta-Arias JD, Mejía SP, González Á. 2020. The Role of the Interleukin-17 Axis

and Neutrophils in the Pathogenesis of Endemic and Systemic Mycoses. Front Cell

Infect Microbiol 10:595301. https://doi.org/10.3389/fcimb.2020.595301.

28.   Sawada Y, Setoyama A, Sakuragi Y, Saito-Sasaki N, Yoshioka H, Nakamura M.

2021. The Role of IL-17-Producing Cells in Cutaneous Fungal Infections. Int J Mol

Sci 22. https://doi.org/10.3390/ijms22115794.

29.   Lee CK, Oliveira LVN, Akalin A, Specht CA, Lourenco D, Gomez CL, Ramirez-

Ortiz ZG, Wang JP, Levitz SM. 2023. Dysregulated pulmonary inflammatory

responses exacerbate the outcome of secondary aspergillosis following influenza.

MBio 14:e0163323. https://doi.org/10.1128/mbio.01633-23.

30.   Sahoo M, Ceballos-Olvera I, del Barrio L, Re F. 2011. Role of the inflammasome, IL-

1β, and IL-18 in bacterial infections. ScientificWorldJournal 11:2037–2050.

https://doi.org/10.1100/2011/212680.

31.   Robinson KM, Choi SM, McHugh KJ, Mandalapu S, Enelow RI, Kolls JK, Alcorn

JF. 2013. Influenza A exacerbates Staphylococcus aureus pneumonia by attenuating

IL-1β   production   in   mice.   J   Immunol   191:5153–5159.

https://doi.org/10.4049/jimmunol.1301237.

FIGURE LEGENDS

Figure 1: Type 17 immune pathway is downregulated during influenza-associated pulmonary aspergillosis (IAPA). WT mice were infected with influenza A H1N1 PR/8/34 on day 0 and subsequently challenged with 2.5×10^7 *aspergillus fumigatus* (AF) at 6 dpii. Lung samples were collected at 48 h post-AF challenge. (**A**) IL-17 and related pro-inflammatory cytokine gene expression quantified by RT-qPCR, (**B**) Protein levels of IL-17-related cytokines quantified using Lincoplex assay or ELISA, (**C**) Flow cytometry analysis of IL-17-producing cells, CD4$^+$ IL-17$^+$ T cells and γδ T cells, (**D**) IL-17/IL-22-associated antimicrobial peptide gene expression quantified by RT-qPCR. Data were compiled from two independent experiments and are presented as means ± SEM, with statistical significance marked as *$p < 0.05$, ** $p < 0.005$, and *** $p < 0.0005$ by unpaired student's T-test.

Figure 2: Restoration of Type 17 immune signaling is not sufficient to provide protection during IAPA. WT mice were infected with influenza A H1N1 PR/8/34 on day 0 and subsequently challenged with 2.5×10^7 *aspergillus fumigatus* (AF) at 6 dpii. Lung samples were collected at 48 h post-AF challenge. (**A-C**) Mice were administered IL-17 expressing adenovirus (IL-17) or control adenovirus during IAPA and (**A**) fungal burden measured by CFU, (**B**) IL-17a gene expression quantified by RT-qPCR, and (**C**) TNF⍺ and CXCL1 cytokines measured by ELISA. (**D-E**) Mice were administered IL-1β or IL-1β+IL-23 murine recombinant protein during IAPA and (**D**) fungal burden measured by CFU, (**E**) IL-17a and IL-22 gene expression quantified by RT-qPCR. (**F**) Mice were administered Reg3β or Reg3γ murine recombinant proteins during IAPA and fungal burden measured by CFU. Data were compiled from two independent experiments and are presented as means ± SEM, with statistical significance marked as *$p < 0.05$, ** $p < 0.005$, and **** $p < 0.0001$ by unpaired student's T-test.

Figure 1: **Type 17 immune pathway is downregulated during influenza-associated pulmonary aspergillosis (IAPA).** WT mice were infected with influenza A H1N1 PR/8/34 on day 0 and subsequently challenged with 2.5×10^7 *aspergillus fumigatus* (AF) at 6 dpii. Lung samples were collected at 48 h post-AF challenge. (**A**) IL-17 and related pro-inflammatory cytokine gene expression quantified by RT-qPCR, (**B**) Protein levels of IL-17-related cytokines quantified using Lincoplex assay or ELISA, (**C**) Flow cytometry analysis of IL-17-producing cells, CD4$^+$ IL-17$^+$ T cells and γδ T cells, (**D**) IL-17/IL-22-associated antimicrobial peptide gene expression quantified by RT-qPCR. Data were compiled from two independent experiments and are presented as means ± SEM, with statistical significance marked as *$p < 0.05$, ** $p < 0.005$, and *** $p < 0.0005$ by unpaired student's T-test.

Figure 2: Restoration of Type 17 immune signaling is not sufficient to provide protection during IAPA. WT mice were infected with influenza A H1N1 PR/8/34 on day 0 and subsequently challenged with 2.5×10^7 *aspergillus fumigatus* (AF) at 6 dpii. Lung samples were collected at 48 h post-AF challenge. (**A-C**) Mice were administered IL-17 expressing adenovirus (IL-17) or control adenovirus during IAPA and (**A**) fungal burden measured by CFU, (**B**) IL-17a gene expression quantified by RT-qPCR, and (**C**) TNF□ and CXCL1 cytokines measured by ELISA. (**D-E**) Mice were administered IL-1β or IL-1β+IL-23 murine recombinant protein during IAPA and (**D**) fungal burden measured by CFU, (**E**) IL-17a and IL-22 gene expression quantified by RT-qPCR. (**F**) Mice were administered Reg3β or Reg3γ murine recombinant proteins during IAPA and fungal burden measured by CFU. Data were compiled from two independent experiments and are presented as means ± SEM, with statistical significance marked as *$p < 0.05$, ** $p < 0.005$, and **** $p < 0.0001$ by unpaired student's T-test.