High Throughput Repurposing Screen Reveals Compounds with Activity Against

***Toxoplasma gondii* Bradyzoites**

Taher Uddin[1], Jing Xia[1], Yong Fu[1], Case W. McNamara[2], Arnab K. Chatterjee[2], L. David

Sibley[1*]

[1]Department of Molecular Microbiology, Washington University School of Medicine, St. Louis,

MO, USA.

[2]Calibr at Scripps Research, La Jolla, California, USA.

*Correspondence: Sibley@wustl.edu

KEYWORDS: High throughput screening, toxoplasmosis, chronic infection

ABSTRACT

Toxoplasma gondii causes widespread chronic infections that are not cured by current treatments

due to inability to affect semi-dormant bradyzoite stages within tissue cysts. To identify

compounds to eliminate chronic infection, we developed a HTS using a recently characterized

strain of *T. gondii* that undergoes efficient conversion to bradyzoites in intro. Stage-specific

expression of luciferase was used to selectively monitor growth inhibition of bradyzoites by the

Library of Pharmacological Active Compounds, consisting of 1280 drug-like compounds. We

identified 44 compounds with >50% inhibitory effects against bradyzoites, including several new

highly potent compounds several of which have precedent for antimicrobial activity. Subsequent

characterization of the compound Sanguinarine sulfate revealed potent and rapid killing against

in vitro produced bradyzoites and bradyzoites harvested from chronically infected mice. These

findings provide a platform for expanded screening and identify promising compounds for

further preclinical development against *T. gondii* bradyzoites responsible for chronic infection.

Toxoplasma gondii is a widespread parasite of animals that causes zoonotic infections in humans

[1]. Serological studies suggest that ~ 1/3 of humans are chronically infected with *T. gondii*

worldwide, although prevalence rates vary widely by geographic region [2]. Human infections are

caused by ingestion of undercooked meat, harboring tissue cysts, or ingestion of oocysts shed by

infected cats [3]. The acute stage of infection is predominated by fast growing tachyzoites that

disseminate widely, including to sites of immune privilege like the brain, followed by

differentiation into cysts harboring semi-dormant bradyzoites, which divide slowly and

asynchronously [4]. Although most infections are only mildly symptomatic and controlled by the

immune system, they persist chronically and hence predispose individuals to subsequent

reactivation if they become immunocompromised [5]. The current standard of care treatment does

not eradicate chronic infection, hence infected individuals remain at risk of reactivation for life.

Long known as a cause of congenital infection [6], recent studies highlight toxoplasmosis as a

cause of ocular disease due to newly acquired infections in otherwise healthy adults [7]. Frequent

and severe outbreaks of ocular toxoplasmosis have been described in South America [8] and India

[9, 10], and may occur in other localities. Notably, infection of healthy individuals in South America

often leads to severe, recurrent ocular toxoplasmosis [11], with an estimated disease burden of 30

million individuals requiring treatment annually in Brazil alone [12]. Additionally, prior

toxoplasma infection has been linked to increased cognitive decline in Alzheimer's Disease,

although not all studies have conformed this association, likely due to the complexity of risk

factors involved [13].

Current therapies for treatment of toxoplasmosis rely on inhibition of the folate pathway in

the parasite [14]. The standard of care therapy (i.e. sulfadiazine and pyrimethamine) inhibits the

rapidly growing tachyzoite stage but has minimal activity on bradyzoites within tissue cysts and

consequently does not eliminate chronic infection [15]. Unfortunately, there are also significant

adverse effects of this treatment regimen due to intolerance or allergic reactions [16, 17] and

contraindication during the first two trimesters of pregnancy [18]. Although drug resistance is not

frequently encountered in treatment of toxoplasmosis, some isolates are naturally resistant to

sulfonamides due to natural variants in dihydropteroate synthetase, or other molecular

mechanisms, thus complicating treatment in some cases [17]. Clindamycin and several macrolide

antibiotics have also been shown to inhibit growth in vitro and in animal models [19, 20] and such

antibiotics have been used to treat toxoplasmosis in humans [21]. However, these compounds are

not specific to the parasite and disrupt the normal endogenous microbiota leading to possible

emergence of pathobionts like *Clostridium difficile* [22].

A review of the literature of compounds that are clinically approved in humans or animals

identified several compounds that inhibit parasite growth in vitro and/or in murine models of

toxoplasmosis [21]. For example, repurposing of guanabenz, an FDA-approved drug that interferes

with translation, showed activity against acute and chronic toxoplasmosis in mice [23], although

this affect was dependent on the strain of mouse [24], and treatment did not eliminate cysts,

resulting in rebound after discontinuation [25]. A repurposing screen for inhibition of tachyzoite

growth in vitro using the Tocriscreen Total Library, which consists of 1280 biologically active

small molecules with confirmed molecular targets, identified multiple compounds that affect

dopaminergic and estrogen signaling, including tamoxifen that was shown to act by upregulating

xenophagy to restrict parasite growth and cause clearance [26]. Additionally, a number of new

investigational compounds have shown an ability to block tachyzoite growth in vitro and during

acute infection in animal models, including several that have activity against bradyzoite growth

or chronic infection [17].

The majority of efforts to identify new compounds with activity against *T. gondii* have

focused on in vitro assays using tachyzoite growth as a readout [17, 21]. Differentiation of

bradyzoites in vitro can be achieved by treatment with stress, such as high pH [27]; however, the

stages that develop under stress often continue to express tachyzoite markers [28, 29], thus

complicating screening efforts. Nonetheless, combining dual promoters to drive firefly luciferase

in the cytosol of bradyzoites and Nanoluc luciferase (nLuc) that was engineered to be secreted

into the cyst matrix, allowed evaluation of compounds for selective activity against chronic

stages [30]. Another promising development is the recent development of an in vitro system for

development of bradyzoites using a specialize KD3 muscle cell line where spontaneous

differentiation occurs [31]. Consistent with the lack of available treatments on tissue cysts in

chronically infected mice, treatment with pyrimethamine and/or sulfadiazine was not effective in

restricting the growth of mature bradyzoites formed in vitro in this system [31]. Hence, this system

could provide a useful platform for testing compounds for activity against bradyzoites, although

it requires the use of a specialized culture system and the treatment and recovery phase needed

for evaluation takes ~ 50 days.

As an alternative, we recently described a type II strain called Tg68 that has a high

propensity to differentiate into bradyzoite in vitro under condition of stress that include high pH

or cultivation in high glutamine, low glucose, that forces metabolism based on glutaminolysis [32].

Unlike other type II strains that undergo partial differentiation, Tg68 forms fully mature

bradyzoites without associated breakthrough of tachyzoites following stress induction in vitro [32].

Here, we engineered this strain to express Firefly luciferase (Fluc) under the control of a

constitutive promoter, and separately generated a lines expressing nLuc under the control of a

bradyzoite promoter. We used these reporter lines to develop a high throughput screen (HTS) and

used it to evaluate the Library of Pharmacological Active Compounds (LOPAC). Several

compounds with activity against both stages were identified, providing proof of concept for

further HTS projects designed to find new treatments for chronic toxoplasmosis.

Results and Discussion

To facilitate HTS using the Tg68 strain, we generated a clonal line of Tg68 constitutively

expressing firefly-luciferase (Fluc) under the p*TUB1* promoter (**Figure 1A**). The pTub1:Fluc

plasmid also contained a resistant DHFR cassette that was integrated into the genome after

electroporation followed by selection with pyrimethamine. Following passage in HFF cells, a

cloned line constitutively expressing Fluc was isolated and is referred to as Tg68-pTub1:Fluc.

Confluent HFF cells grown in 384-well plates were infected with Tg68-pTub1:Fluc tachyzoites

at 2×10^4 parasites per well, and the infection was allowed to proceed for 72 hr under 5% CO_2 at

37°C (**Figure 1B**). Comparison of Fluc activity between day 3 and day 0 (4 hr postinfection)

increased significantly, representing growth of the parasite (**Figure 1C**). In addition, we analyzed

Fluc expression from replicate 14 wells, across 5 plates and determined the Coefficient of

Variation (CV) for positive wells was 7±3% (**Table S1**). Addition of BRD7929, which targets

parasite phenylalanine tRNA synthetase [33], completely inhibited parasite growth at a

concentration of 10 µM (**Figure 1D**). Comparison of the luciferase signal from untreated and

treated wells from 5 plates demonstrated an average Z′ value of 0.77±0.11 (**Table S1**).

Additionally, the Tg68-pTub1:Fluc line demonstrated comparable sensitivity to BRD7929 and

atovaquone compared to the reference type II strain TgMe49-Fluc (**Figure 1D**). Taken together,

these findings indicate that Tg68-pTub1:Fluc provides a robust readout for HTS of compounds

against tachyzoite growth.

To expand the use of Tg68 for HTS of compounds with activity against bradyzoites, we
generated a clonal line expressing nLuc under control of the p*BAG1* promoter, which is strongly
upregulated upon differentiation [34]. The pBAG1:nLuc plasmid also contained a resistant DHFR
cassette that was integrated into the genome after electroporation followed by selection with
pyrimethamine. Following serial passage in HFF cells, a positive clone was isolated and referred
to as Tg68-pBAG1:nLuc. (**Figure 2A**). In previous studies, we have shown that the Tg68 strain
undergoes highly efficient differentiation to bradyzoites in alkaline media at low CO_2 in vitro and
that it retains a mature bradyzoite profile even at relatively high multiplicities of infection [32].
Tg68 also undergoes efficient differentiation when grown in glucose free medium supplemented
with 10 mM glutamine, a process that forces glutaminolysis for energy production via the
mitochondrion [32]. Confluent HFF cells in 384-well plates were infected with pBAG1:nLuc
parasites ($3x10^3$ parasites per well) for 2 hr, followed by washing and shifting to alkaline or
glutamine media. Cultures were maintained for 10 days under CO_2-free conditions (ambient air),
with media changes on day 3 and 6. On day 6, compounds were added and on day 10 luciferase
assays were performed (**Figure 2B**). Expression of nLuc was very low in cultures of tachyzoites
and in the initial culture conditions at day 0 (4 hr postinfection) in normal or differentiation
media (**Figure 2C**). Expression of nLuc increased dramatically by over 4 logs by day 3 and
continued to increase significantly at day 6 and at day 10 (**Figure 2C**). On day 10, analysis of
luciferase values from 14 wells showed an optimal Coefficient of Variation (CV) of 10±2% for
alkaline media and 7±2% for glutamine media, indicating suitability for high-throughput
screening (HTS) (**Table S1**). Treatment with BRD7929 at a concentration of 10 μM completely
inhibited growth of the parasites and comparison of nLuc activity from treated and untreated
wells revealed a Z´ value of 0.67±0.05 for alkaline media and 0.76±0.06 for glutamine media
(**Table S1**). We also compared the potency of BRD7929 and atovaquone under both conditions
that induced bradyzoite differentiation (**Figure 2D**). Both compounds showed a reduction in
EC_{50} values under glutamine differentiation when compared to alkaline conditions, although this
was much more dramatic for atovaquone that showed a > 30-fold shift (**Figure 2D**). The greater
potency of compounds in glutamine medium may reflect a decreased ability of the parasite to
generate energy stores from glutaminolysis vs. glycolysis as suggested previously by the modest
growth defects in knockouts of *T. gondii* hexokinase [35] and glucose transporter 1 [36].
Consequently, the use of glutamine medium for bradyzoite development may preferentially

reveal compounds that act on the mitochondrion, as is the case for atovaquone that inhibits the

bc1 complex [37].

The LOPAC library (1157 of 1280 total compounds) was plated in 384 well format and tested

for parasite growth inhibition in duplicate using a single concentration of 10 μM in the

tachyzoite and both alkaline and glutamine-induced bradyzoite assays. Growth inhibition was

averaged from the two replicates and plotted as a Venn diagram summarizing the outcome of

each of the three assays (**Figure 3A**). A total of 27 compounds showed selective inhibition of

tachyzoite growth without affecting growth in the other assays (**Figure 3A**). The largest number

of compounds was identified in the glutamine induced bradyzoite assay, perhaps reflecting the

metabolic liability of this growth condition (**Figure 3A**). In total, 21 compounds that inhibited

parasite growth in all three assays by 50% or more were identified as Primary Hits (**Figure 3A**).

Furthermore, we identified 9 compounds that inhibited bradyzoite growth in both assays by 50%

or more but not tachyzoite growth (**Figure 3A**). Finally, we identified 14 compounds that

specifically inhibited glutamine-induced bradyzoite growth by $\geq 80\%$, but were not effective in

growth inhibition in the other two assays (**Figure 3A**). Robust validation of the within plate and

between replicates was performed through collective calculation of the coefficient of variation

(CV) and Z' analysis of assay plates (**Table S2**). Of the 44 Primary Hit compounds defined above

(**Figure 3B**), 36 compounds were available in quantity at Calibr and were chosen for further

testing (**Table S3**). These 36 compounds were tested in a dilution series and 9 compounds were

found to have EC$_{50}$ values of ≤ 2 μM in either tachyzoite and/or bradyzoite assays, thus defining

a set of Top Hits (**Table S3, Figure 3B**).

We were able to source 8 of the 9 Top HIT compounds from commercial sources while one

of the top hit compounds Emetine dihydrochloride hydrate (TgEC50 ~70nM) was not available

from commercial sources for further study. The 8 available compounds were tested for

cytotoxicity against HepG2 and THP-1 cell lines, along with a control drug, atovaquone (**Figure**

3B). Most compounds exhibited a favorable selectivity index (SI), which is a measure of the

compound's EC$_{50}$ against the parasite compared to the CC$_{50}$ against host cells (**Table 1**).

However, a few compounds, namely Idarubicin hydrochloride and MS012, showed high toxicity,

while Auranofin and JFD00244 showed modest toxicity towards the cell lines. Auranofin is a

gold containing compound that has been approved for treatment of rheumatoid arthritis and it has

previously been reported to have activity against several parasites [38]. Consistent with this profile,

auranofin has previously been shown inhibit replication of the type I RH strain in vitro, reduce
infection in a chicken embryo model [39]and reduce the burden of cysts in chronically infected
mice [40]. However, the high level of growth inhibition for HepG2 and THP-1 cells treated with
auranofin in the present study would appear to limit the potential of this compound for further
clinical development. Additionally, Brefeldin A, which blocks ER to Golgi transport showed a
favorable SI in HepG2 cells but not in THP-1 cells (**Table 1**), suggesting that block of protein
export has very different consequences on host cell growth in different lineages. As a result,
these compounds were excluded from further assays due to their undesirable cytotoxic effects.

Table 1 Potency and selectivity of Top Hits

Compound	Tg68Fluc EC_{50} (Tachyzoite)	Me49Fluc EC_{50} (Tachyzoite)	Tg68nLuc EC_{50} (Alkaline)	Tg68nLuc EC50 (Glutamine)	HepG2 CC_{50}	SI	THP-1 CC_{50}	SI
Diphenyleneiodonium Sulfate	0.01	0.01	0.27	0.06	1.28	96	4.85	365
Sanguinarine Sulfate	0.03	0.01	4.00	1.25	9.44	238	4.34	130
Brefeldin A	0.06	0.05	0.40	0.16	60.00	987	0.07	1
Idarubicin hydrochloride	0.19	0.67	0.30	0.82	0.14	1	0.59	3
T0070907	0.75	0.87	0.64	0.60	18.12	24	21.98	29
MS012	2.46	2.68	0.85	2.23	2.03	1	1.19	0.5
Auranofin	2.75	1.12	0.68	0.38	9.81	4	2.83	1
JFD00244	3.89	3.39	2.88	4.48	16.07	4	2.95	1
Atovaquone	0.16	0.10	2.07	0.06	9.67	62	14.26	91
BRD7929	0.11	0.05	0.22	0.08	6.08	54	2.87	25

Values in μM, average of two or more biological replicates,

SI = selectivity index (Host cell CC_{50} / EC_{50} Tg68Fluc).

Although Tg68 undergoes efficient conversion to bradyzoites in vitro, with a transcriptional
profile that resemble in vivo bradyzoites [32], it may still lack some of the features of mature tissue
cysts. Hence, we tested select Top Hits, along with several reference compounds, against tissue
cysts that were harvested from chronically infected mice (**Figure 4A**). For these assays, we used
the ME49 EW strain, which produces high numbers of cysts in vivo [46]. As shown in Table 1, the
ME49 strain has a very similar sensitivity to the Top Hits and reference compounds, thus
validating the choice of strain. Compounds were tested using continuous exposure of ex vivo
bradyzoites to 3XEC_{90} as measured on tachyzoites. Alternatively, ex vivo bradyzoites were
treated at 3XEC_{90} for only 4 hr followed by washout to ascertain how irreversibly they might act

(**Figure 4A**). All three of the reference compounds BRD7929, atovaquone, and pyrimethamine showed potent inhibition of bradyzoites when used continuously (**Figure 4B**). Although atovaquone and BRD7929 are active on bradyzoites (**Table 1**), the activity of pyrimethamine in this continuous treatment assay is likely because it inhibits the outgrowth of tachyzoites. Consistent with this specificity, pyrimethamine was largely ineffective when used for only 4 hr, indicating it has minimal effects on bradyzoites present at the start of the assay (**Figure 4B**). Similar to pyrimethamine, testing of diphenyleneiodonium and T0070907 in the ex vivo bradyzoite assay revealed that they only work when used in continuous treatment, suggesting their activity is static rather than cidal (**Figure S1**). Diphenyleneiodonium is an inhibitor of NADPH oxidase that separately induces oxidative stress [41]. Previous studies have shown diphenyleneiodonium inhibits growth of *T. gondii* tachyzoites in ARPE-19 cells through the production of ROS [42]. In a separate study, it also showed activity against *P. falciparum* with an EC_{50} of 0.06 nM [43], and it has also been shown to have broad spectrum antibacterial activity [44]. T0070907 is an inhibitor of peroxisome proliferator activator receptor γ that induces G2/M arrest and thus has activity against cancer cells [45]. This nuclear hormone pathway is not conserved in *T. gondii*, and this compound has not been described to have anti-microbial activity previously, so the potential mechanism of action is uncertain.

Atovaquone showed partial inhibition in the washout ex vivo bradyzoite assay, and consistent with previous studies showing it is partially active in reducing cysts numbers during chronic infection in vivo [47]. BRD7929 was highly effective in preventing the outgrowth of bradyzoites even when removed after 4 hr (**Figure 4B**), consistent with previous findings [33]. Testing of the Top Hit Sanguinarine sulfate revealed potent activity in the ex vivo bradyzoite wash out assay, thus confirming it has activity against bradyzoites as well as tachyzoites (**Figure 4B**). Sanguinarine sulfate is a natural product produced by the opium poppy and several other plants and it consists of a benzoquinoline alkaloid [48]. Sanguinarine has anti-inflammatory, anti-tumor and antimicrobial activities and it is thought to act on numerous signaling pathways in human cells [49]. Sanguinarine has been described to be toxic to host cell, thought to be due to its action on Na+/K+ ATPase, although other mechanisms have also been described [48]. Although we did not measure appreciable cell toxicity in our assays, other studies have indicated sanguinarine induces apoptosis or blocks cell growth [49]. Additionally, the LD_{50} in mouse is 18 mg/kg by i.p. injection [49], likely limiting its use in efficacy testing for toxoplasmosis.

Conclusions

The newly discovered Tg68 strain is permissive for bradyzoite differentiation and forms the foundation for the screening protocol in this study. We have developed protocols for HTS of compound libraries that target for tachyzoites and bradyzoites and further validated hits in assays against ex vivo produced bradyzoites. Several of the hits identified in this screen have precedent for being antimicrobial; however, they pose challenges for selectively and lack of toxicity. Nonetheless, the methodology develop here offers promise for future HTS to identify potent and selective leads.

Experimental Section

Compounds and liquid handling. The LOPAC compound library was plated in a 384-well plate format for primary screening and determination of half-maximal effective concentration (EC_{50}) of primary hits by Calibr at Scripps Research (La Jolla, CA). The plates were stored at -80°C prior to use. All liquid handling steps (host and parasite cell seeding, media exchange, compound transfer to assay plates, and addition of luciferase reagents) were carried out in a semi-automated facility to ensure efficient and consistent execution of assays across all replicates (High-Throughput Screening Center, Washington University School of Medicine). Eight top hit compounds and two control compounds, atovaquone (Sigma#A7986) and pyrimethamine (Sigma#46706), were obtained from commercial sources, and follow-up assays were manually performed in-house.

Construction of transgenic parasites and parasite culture. Tg68 pTub1:Fluc,DHFR parasite lines were generated by utilizing a pre-existing pTub:Fluc plasmid [33], which provides constitutive firefly luciferase (Fluc) expression from the alpha tubulin promoter, and a pyrimethamine-resistant DHFR selectable marker. Tg68 tachyzoites were electroporated with 50 μg of the plasmid and subsequently selected using pyrimethamine (3 μM).

The plasmid pSAG1:EGFP-DHFR-BAG1:nLuc was constructed by assembling fragments encoding EGFP driven by the *SAG1* promoter, nLuc driven by the *BAG1* promoter, DHFR pyrimethamine resistance marker, and pNJ-26 vector using NEBuilder HiFi DNA Assembly Master Mix (NEB). Tg68 pBAG1:nLuc, DHFR tachyzoites (Tg68pSAG1:EGFP-DHFR-BAG1:nLuc) were electroporated with 50 μg of this plasmid and selected with pyrimethamine (3

μM) in order to establish stable parasite lines. Detailed primer information can be found in **Table**
S4.

Stable clones were isolated through limiting dilution, and the expression of the transgene was
confirmed by luciferase expression. Clonal transgenic Tg68 tachyzoites were maintained by
serial passage in T25 flasks with confluent HFFs in D10 medium (Dulbecco's Modified Eagle's
Medium, DMEM; Thermo Fisher, 10% fetal bovine serum (FBS), 2 mM glutamine (Sigma), and
10 μg/mL gentamicin (Thermo Fisher)) at 37°C and 5% CO_2.

In Vitro Assays for bradyzoite and tachyzoite growth inhibition.

Host cell culture and seeding

Human Foreskin Fibroblasts (HFF-1 SCRC-1041, ATCC) cells were cultured in T175 flasks
using D10 medium. Prior to seeding the cells to the assay plates, a cell suspension was prepared
by treating with trypsin (0.5 g/l porcine trypsin and 0.2 g/L EDTA in Hank's Balanced Salt
Solution with phenol red) for 5 min. EC_{50} assays (half maximal effective concentration) were
conducted in a 384-well plate format. Confluent HFF cells were seeded 4 days in advance using
a Multidrop Combi dispensing 80 uL/well, while stirring the cell suspension at 350 rpm. The
plates were stacked in a Cytomat rack on a level surface at room temperature until all plates were
completed, and then they were placed in a 37°C incubator with 5% CO_2. To minimize edge
effects, only the inner 240 wells of each plate were utilized.

Bradyzoite Primary Screening and EC_{50} Assay

The medium from the HFF cell plates was aspirated using a Biotek ELx405CW, leaving 20
μL/well, before infection with parasites. Freshly harvested Tg68pBAG1:nLuc,DHFR parasites
(3×10^3) in a 40 μL volume were then added to each well (total volume of 60 μL/well, containing
0.1% DMSO in D10) using a Fluotics BXi-50F while stirring the parasite suspension at 350 rpm.
The plates were incubated for 2 hr at 37°C in a 5% CO_2 incubator to allow parasite invasion.
Afterwards, the culture medium was removed and switched to either 80 μL/well of Alkaline
(RPMI 1640 (Sigma, #R6504) containing 1% FBS and 50 mM HEPES (Sigma), adjusted to pH
8.2), or Glutamine (glucose-free RPMI 1640 (Sigma, #R1383) containing 1% FBS and 50 mM
HEPES (Sigma), 10 mM glutamine, adjusted to pH 7.2) medium. The culture was then incubated
at 37°C in ambient CO_2 and maintained for 10 days, with media changes occurring on day 3 and
6 using a Biotek ELx405CW and Multidrop Combi inside a biohood to minimize contamination.
On day 6, before the compound was added, the plates were aspirated, and 40 μL/well of fresh

media was added, followed by the transfer of 40 µL/well of 2X compound using a Fluotics BXi-

50F. On day 10, the plates were equilibrated to room temperature for 30 min prior to the assay

readout. To read the plates, the medium was aspirated, 20 uL of Promega Nano-Glo reagent

(Promega N1150) was dispensed into each well using a Multidrop Combi, and the plates were

covered with a black lid. The plates were then incubated for 10 min at room temperature, and the

readout was performed using Envision.

Tachyzoite Primary Screening and EC_{50} Assay

Before being infected with parasites, the medium was removed from the HFF cell plates using a

Biotek ELx405CW, leaving 20 µL/well. Freshly harvested Tg68pTub1:Fluc parasites (2×10^4) in

a 20 µL volume were then added to each well while stirring the parasite suspension at 350 rpm,

and 40 µL of 2X compound solutions were transferred (80 µL/well total volume, containing

0.1% DMSO in D10) using a Fluotics BXi-50F. The plates were incubated for 72 hr at 37 °C in a

5% CO_2 incubator. Before the assay readout, the plates were equilibrated to room temperature for

30 min. To read the plates, the medium was aspirated to 20 µL using a Biotek ELx405CW, 20 µL

of Promega Bright-Glo reagent (Promega E2650) was dispensed into each well, and the plates

were covered with a black lid. The plates were then incubated for 10 min at room temperature

using a Multidrop Combi, and the readout was performed using Envision.

In Vitro Assays for Host cell cytotoxicity. Cytotoxicity against host cells host cells was tested

using human hepatocellular carcinoma cells (HepG2, ATCC-HB-8065) and human monocytic

tumor line (THP-1, ATCC-TIB-202), which were maintained according to ATCC

recommendations. Host cell lines were tested negative for mycoplasma using an e-Myco plus kit

(Intron Biotechnology). Compounds were diluted to 2X concentration, and a 10-dose serial

dilution series was prepared through stepwise, 3-fold dilutions in recommended media with 0.1%

DMSO. Host cells were seeded at a density of 10^4 cells/well (100 µL vol) to achieve sub-

confluent monolayers for a 72 hr growth assay. Prior to compound addition, THP-1 cells were

treated with 10 ng/mL phorbol 12-myristate 13 acetate for 24 hr to facilitate differentiation into

macrophages. HepG2 cells were treated with compounds 6 hr post-seeding (200 µL final volume,

0.05% DMSO) and incubated under 37 °C, 5% CO_2 culture conditions. At 72 hr post compound

addition, culture media were aspirated to 80 µL, and an equal volume of CellTiter-Glo

Luminescent Cell Viability Assay reagent (Promega G7571) was added. Luciferase activity was

measured using a BioTek Cytation 3 equipped with Gen5 software (v3.08). Each assay was

repeated with two technical replicates within two independent biological replicates. Statistical

analyses were performed using Prism 10 (GraphPad Software, Inc.). Dose-response inhibition

curves for host cell toxicity (CC_{50} values) were generated using the "Log(inhibitor) vs

normalized response variable slope" function. The reported values represent averages from three

biological replicates.

Ex-vivo Bradyzoite differentiation assay. CBA/CaJ mice (Strain # 000654) from Jackson

Laboratory were housed in an approved facility at Washington University School of Medicine,

and all animal studies followed ethical guidelines approved by the Institutional Animal Care and

Use Committee. CBA/CaJ mice were infected by oral gavage with 5-10 tissue cysts from the

brain homogenate of previously infected mice. The brains of CBA/CaJ mice infected with the

TgME49-EW strain [46] were collected at 1-2 mos post-infection, homogenized, and tissue cysts

were isolated using Percoll gradients, as described previously [50]. To release the bradyzoites from

the tissue cysts, purified tissue cysts were treated with an acid-pepsin solution (170 mM NaCl,

60 mM HCl), and a freshly prepared pepsin solution (0.1 mg/mL in 1xPBS) for 10 min at 37 °C.

The reaction was stopped by adding a neutralization buffer (94 mM Na_2CO_3). The liberated

bradyzoites were then evenly distributed into a duplicate set of 6-well plates (technical

replicates), each containing 5 mL of culture media. Each plate included a negative control (media

with 0.1% DMSO), the previously characterized PheRS inhibitor BRD7929 (0.5 µM)[33], and

pyrimethamine (2.0 µM). Compounds were tested at EC_{90} and $3XEC_{90}$ concentrations based on

their corresponding in vitro tachyzoite growth inhibition ($TgEC_{50}$) assays (Table S3). After a 4 hr

treatment, the compounds were removed from one set of plates by washing with 3 times in PBS

and replacement with compound-free D10 medium. The plates were then incubated at 37°C, 5%

CO_2 undisturbed for 12-14 days to form plaques. Plaque quantification was performed by fixing

the plates in 100% ethanol for 5 min at room temperature, followed by staining with a 0.1%

crystal violet solution for 10 min. After rinsing with water and air drying, plaque quantification

was conducted using a Nikon eclipse ts2 microscope equipped with a 4X objective. The number

of plaques from two biological replicates was normalized to DMSO control as a percentage.

Statistical analysis

Statistical comparisons were performed in Prism (GraphPad). Data were first analyzed for

normal distribution and according to the outcome, were then non-parametric tests applied.

Statistical test and *P* values are given in the figure legends.

Acknowledgments:

We thank Jennifer Barks for HFF cell culture and for providing administrative support for

research materials. We acknowledge Maxene Ilagan and Mike Prinsen from Washington

University's High Throughput Screening Center for their assistance with assay set-up and data

handling. Furthermore, we thank Kaycie Morwood and her team from Calibr – A Division of

Scripps Research for their assistance with compound management and plating. The ME49EW

was a kindly provided by Michael White, University of South Florida. Partial support provided

by an NIH grant (AI143857) and a pilot grant from the Center for Drug Discovery at Washington

University (PJ00027513).

Abbreviations used:

DMSO Dimethylsulfoxide

DHFR Dihydrofolate Reductase

EDTA Ethylenediaminetetraacetic acid

HFF Human Foreskin Fibroblasts

HepG2 human hepatocellular carcinoma cell

THP-1 human monocytic tumor cell

HTS High Throughput Screening

nLuc Nano Luciferase

Fluc Firefly Luciferase

PheRS Phenylalanine tRNA synthetase

Supplemental Materials

Table S1 HTS parameters for lines expressing luciferase.

Table S2 HTS parameters for the LOPAC screen.

Table S3 Activity of Primary Hits in growth inhibition assays.

Table S4 Primers used in the study.

Figure S1. Testing of additional TOP HIT compounds for activity against ex vivo bradyzoites.

References

1. Dubey JP. Toxoplasmosis of animals and humans. Boca Raton: CRC Press; 2010. 313 p.
2. Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis. Int J Parasitol. 2009;39(12):1385-94. Epub 2009/05/13. doi: S0020-7519(09)00184-2 [pii]
10.1016/j.ijpara.2009.04.003. PubMed PMID: 19433092.
3. Almeria S, Dubey JP. Foodborne transmission of Toxoplasma gondii infection in the last decade. An overview. Res Vet Sci. 2021;135:371-85. Epub 20201024. doi: 10.1016/j.rvsc.2020.10.019. PubMed PMID: 33148402.
4. Watts E, Zhao Y, Dhara A, Eller B, Patwardhan A, Sinai AP. Novel Approaches Reveal that Toxoplasma gondii Bradyzoites within Tissue Cysts Are Dynamic and Replicating Entities In Vivo. MBio. 2015;6(5):e01155-15. doi: 10.1128/mBio.01155-15. PubMed PMID: 26350965; PMCID: PMC4600105.
5. Elsheikha HM, Marra CM, Zhu XQ. Epidemiology, Pathophysiology, Diagnosis, and Management of Cerebral Toxoplasmosis. Clin Microbiol Rev. 2021;34(1). Epub 20201125. doi: 10.1128/CMR.00115-19. PubMed PMID: 33239310; PMCID: PMC7690944.
6. Torgerson PR, Mastroiacovo P. The global burden of congenital toxoplasmosis: a systematic review. Bull World Health Organ. 2013;91(7):501-8. Epub 2013/07/05. doi: 10.2471/BLT.12.111732
BLT.12.111732 [pii]. PubMed PMID: 23825877; PMCID: 3699792.
7. Jones LA, Alexander J, Roberts CW. Ocular toxoplasmosis: in the storm of the eye. Parasite Immunol. 2006;28:635-42.
8. Glasner PD, Silveira C, Kruszon-Moran D, Martins MC, Burnier M, Silveira S, Camargo ME, Nussenblatt RB, Kaslow RA, Belfort R. An unusually high prevalence of ocular toxoplasmosis in southern Brazil. Am J Opthalmol. 1992;114:136-44.
9. Holland GN. An epidemic of toxoplasmosis: lessons from Coimbatore, India. Arch Ophthalmol. 2010;128(1):126-8. doi: 10.1001/archophthalmol.2008.538. PubMed PMID: 20065229.
10. Palanisamy M, Madhavan B, Balasundaram MB, Andavar R, Venkatapathy N. Outbreak of ocular toxoplasmosis in Coimbatore, India. Indian J Ophthalmol. 2006;54(2):129-31. doi: 10.4103/0301-4738.25839. PubMed PMID: 16770035.
11. Pfaff AW, de-la-Torre A, Rochet E, Brunet J, Sabou M, Sauer A, Bourcier T, Gomez-Marin JE, Candolfi E. New clinical and experimental insights into Old World and neotropical ocular toxoplasmosis. Int J Parasitol. 2013. Epub 2013/11/10. doi: S0020-7519(13)00255-5 [pii]
10.1016/j.ijpara.2013.09.007. PubMed PMID: 24200675.
12. McPhillie M, Zhou Y, El Bissati K, Dubey J, Lorenzi H, Capper M, Lukens AK, Hickman M, Muench S, Verma SK, Weber CR, Wheeler K, Gordon J, Sanders J, Moulton H, Wang K, Kim TK, He Y, Santos T, Woods S, Lee P, Donkin D, Kim E, Fraczek L, Lykins J, Esaa F, Alibana-Clouser F, Dovgin S, Weiss L, Brasseur G, Wirth D, Kent M, Hood L, Meunieur B, Roberts CW, Hasnain SS, Antonyuk SV, Fishwick C, McLeod R. New paradigms for understanding and step changes in treating active and chronic, persistent apicomplexan infections. Sci Rep. 2016;6:29179. Epub 2016/07/15. doi: 10.1038/srep29179. PubMed PMID: 27412848; PMCID: PMC4944145.
13. Xiao J, Savonenko A, Yolken RH. Strain-specific pre-existing immunity: A key to understanding the role of chronic Toxoplasma infection in cognition and Alzheimer's

diseases? Neurosci Biobehav Rev. 2022;137:104660. Epub 20220409. doi:
10.1016/j.neubiorev.2022.104660. PubMed PMID: 35405182.

14. Wei HX, Wei SS, Lindsay DS, Peng HJ. A Systematic Review and Meta-Analysis of the
Efficacy of Anti-Toxoplasma gondii Medicines in Humans. PLoS One.
2015;10(9):e0138204. Epub 2015/09/24. doi: 10.1371/journal.pone.0138204. PubMed
PMID: 26394212; PMCID: PMC4578932.

15. McCabe RE. Antitoxoplasma chemotherapy. In: Joynson DHM, Wreghitt TG, editors.
Toxoplasmosis: a comprehensive clinical guide. Cambridge: Cambridge Univ. Press; 2001.
p. 319-59.

16. Ben-Harari RR, Goodwin E, Casoy J. Adverse Event Profile of Pyrimethamine-Based Therapy
in Toxoplasmosis: A Systematic Review. Drugs R D. 2017;17(4):523-44. doi:
10.1007/s40268-017-0206-8. PubMed PMID: 28879584; PMCID: PMC5694419.

17. Alday PH, Doggett JS. Drugs in development for toxoplasmosis: advances, challenges, and
current status. Drug Des Devel Ther. 2017;11:273-93. Epub 2017/02/10. doi:
10.2147/DDDT.S60973. PubMed PMID: 28182168; PMCID: PMC5279849.

18. Kieffer F, Wallon M. Congenital toxoplasmosis. Handb Clin Neurol. 2013;112:1099-101. Epub
2013/04/30. doi: 10.1016/B978-0-444-52910-7.00028-3. PubMed PMID: 23622316.

19. Berg-Candolfi M, Candolfi E. Depression of the N-demethylation of erythromycin,
azithromycin, clarithromycin and clindamycin in murine Toxoplasma infection. Int J
Parasitol. 1996;26(11):1321-3. Epub 1996/11/01. PubMed PMID: 9024879.

20. Derouin F, Almadany R, Chau F, Rouveix B, Pocidalo JJ. Synergistic activity of azithromycin
and pyrimethamine or sulfadiazine in acute experimental toxoplasmosis. Antimicrob
Agents Chemother. 1992;36(5):997-1001. Epub 1992/05/01. PubMed PMID: 1324642;
PMCID: PMC188824.

21. Neville AJ, Zach SJ, Wang X, Larson JJ, Judge AK, Davis LA, Vennerstrom JL, Davis PH.
Clinically Available Medicines Demonstrating Anti-Toxoplasma Activity. Antimicrob
Agents Chemother. 2015;59(12):7161-9. Epub 2015/09/24. doi: 10.1128/AAC.02009-15.
PubMed PMID: 26392504; PMCID: PMC4649158.

22. Slimings C, Riley TV. Antibiotics and hospital-acquired Clostridium difficile infection: update
of systematic review and meta-analysis. J Antimicrob Chemother. 2014;69(4):881-91.
Epub 2013/12/11. doi: 10.1093/jac/dkt477. PubMed PMID: 24324224.

23. Benmerzouga I, Checkley LA, Ferdig MT, Arrizabalaga G, Wek RC, Sullivan WJ, Jr.
Guanabenz repurposed as an antiparasitic with activity against acute and latent
toxoplasmosis. Antimicrob Agents Chemother. 2015;59(11):6939-45. Epub 2015/08/26.
doi: 10.1128/AAC.01683-15. PubMed PMID: 26303803; PMCID: PMC4604420.

24. Martynowicz J, Doggett JS, Sullivan WJ, Jr. Efficacy of Guanabenz Combination Therapy
against Chronic Toxoplasmosis across Multiple Mouse Strains. Antimicrob Agents
Chemother. 2020;64(9). Epub 20200820. doi: 10.1128/AAC.00539-20. PubMed PMID:
32540979; PMCID: PMC7449173.

25. Martynowicz J, Sullivan WJ, Jr. Rebound of cyst number following discontinuation of
guanabenz treatment for latent toxoplasmosis. Mol Biochem Parasitol. 2021;245:111411.
Epub 20210904. doi: 10.1016/j.molbiopara.2021.111411. PubMed PMID: 34492239;
PMCID: PMC8484070.

26. Dittmar AJ, Drozda AA, Blader IJ. Drug Repurposing Screening Identifies Novel Compounds
That Effectively Inhibit Toxoplasma gondii Growth. mSphere. 2016;1(2). Epub 20160302.
doi: 10.1128/mSphere.00042-15. PubMed PMID: 27303726; PMCID: PMC4894684.

27. Soête M, Fortier B, Camus D, Dubremetz JF. *Toxoplasma gondii*: kinetics of bradyzoite-
tachyzoite interconversion *in vitro*. Exp Parasitol. 1993;76:259-64.
28. Garfoot AL, Wilson GM, Coon JJ, Knoll LJ. Proteomic and transcriptomic analyses of early
and late-chronic Toxoplasma gondii infection shows novel and stage specific transcripts.
BMC Genomics. 2019;20(1):859. Epub 2019/11/16. doi: 10.1186/s12864-019-6213-0.
PubMed PMID: 31726967; PMCID: PMC6857164.
29. Bohne W, Heeseman J, Gross U. Coexistence of hetergeneous populations of *Toxoplasma*
gondii parasites within parasitophorous vacuoles of murine macrophages as revealed by a
bradyzoite-specific monoclonal antibody. Parasitol Res. 1993;79:485-7.
30. Smith D, Lunghi M, Olafsson EB, Hatton O, Di Cristina M, Carruthers VB. A High-
Throughput Amenable Dual Luciferase System for Measuring Toxoplasma gondii
Bradyzoite Viability after Drug Treatment. Anal Chem. 2023;95(2):668-76. Epub
20221222. doi: 10.1021/acs.analchem.2c02174. PubMed PMID: 36548400; PMCID:
PMC9850410.
31. Christiansen C, Maus D, Hoppenz E, Murillo-Leon M, Hoffmann T, Scholz J, Melerowicz F,
Steinfeldt T, Seeber F, Blume M. In vitro maturation of Toxoplasma gondii bradyzoites in
human myotubes and their metabolomic characterization. Nat Commun. 2022;13(1):1168.
Epub 20220304. doi: 10.1038/s41467-022-28730-w. PubMed PMID: 35246532; PMCID:
PMC8897399.
32. Xia J, Fu Y, Huang W, Sibley LD. Constitutive upregulation of transcription factors underlies
permissive bradyzoite differentiation in a natural isolate of Toxoplasma gondii. bioRxiv.
2024. Epub 20240228. doi: 10.1101/2024.02.28.582596. PubMed PMID: 38464000;
PMCID: PMC10925318.
33. Radke JB, Melillo B, Mittal P, Sharma M, Sharma A, Fu Y, Uddin T, Gonse A, Comer E,
Schreiber SL, Gupta AK, Chatterjee AK, Sibley LD. Bicyclic azetidines target acute and
chronic stages of Toxoplasma gondii by inhibiting parasite phenylalanyl t-RNA synthetase.
Nat Commun. 2022;13(1):459. Epub 2022/01/26. doi: 10.1038/s41467-022-28108-y.
PubMed PMID: 35075105; PMCID: PMC8786932 Washington University have applied
for a patent (PCT/US2019/051686 listing J.B.R., E.C., B.M. and L.D.S. as inventors) for
the use of bicyclic azetidines for antiparasitic therapies. All other authors declare no
competing interests.
34. Bohne W, Gross U, Ferguson DJP, Hessemann J. Cloning and characterization of a bradyzoite-
specifically expressed gene (*hsp30/bag1*) of *Toxoplasma gondii*, related to genes encoding
small heat-shock proteins of plants. Molecular Microbiology. 1995;16(6):1221-30.
35. Shukla A, Olszewski KL, Llinas M, Rommereim LM, Fox BA, Bzik DJ, Xia D, Wastling J,
Beiting D, Roos DS, Shanmugam D. Glycolysis is important for optimal asexual growth
and formation of mature tissue cysts by Toxoplasma gondii. Int J Parasitol.
2018;48(12):955-68. Epub 2018/09/04. doi: 10.1016/j.ijpara.2018.05.013. PubMed PMID:
30176233.
36. Blume M, Rodriguez-Contreras D, Landfear S, Fleige T, Soldati-Favre D, Lucius R, Gupta N.
Host-derived glucose and its transporter in the obligate intracellular pathogen Toxoplasma
gondii are dispensable by glutaminolysis. Proc Natl Acad Sci U S A. 2009;106(31):12998-
3003. Epub 2009/07/21. doi: 0903831106 [pii]
10.1073/pnas.0903831106. PubMed PMID: 19617561; PMCID: 2722337.

37. McFadden DC, Tomavo S, Berry EA, Boothroyd JC. Characterization of cytochrome b from
Toxoplasma gondii and Q$_o$ domain mutations as a mechanism of atovaquone-resistance.
Mol Biochem Parasitol. 2000;108(1):1-12.

38. Yildirim A, Ozbilgin A, Yereli K. Antiprotozoal activity of auranofin on Trypanosoma cruzi,
Leishmania tropica and Toxoplasma gondii: in vitro and ex vivo study. Trans R Soc Trop
Med Hyg. 2023;117(10):733-40. doi: 10.1093/trstmh/trad040. PubMed PMID: 37377375.

39. Andrade RM, Chaparro JD, Capparelli E, Reed SL. Auranofin is highly efficacious against
Toxoplasma gondii in vitro and in an in vivo experimental model of acute toxoplasmosis.
PLoS   Negl   Trop   Dis.   2014;8(7):e2973.   Epub   20140731.   doi:
10.1371/journal.pntd.0002973. PubMed PMID: 25079790; PMCID: PMC4117455.

40. Abou-El-Naga IF, Mogahed N. Repurposing auranofin for treatment of Experimental Cerebral
Toxoplasmosis. Acta Parasitol. 2021;66(3):827-36. Epub 20210208. doi: 10.1007/s11686-
021-00337-z. PubMed PMID: 33555553.

41. Riganti C, Gazzano E, Polimeni M, Costamagna C, Bosia A, Ghigo D. Diphenyleneiodonium
inhibits the cell redox metabolism and induces oxidative stress. J Biol Chem.
2004;279(46):47726-31. Epub 20040909. doi: 10.1074/jbc.M406314200. PubMed PMID:
15358777.

42. Sun PR, Gao FF, Choi HG, Zhou W, Yuk JM, Kwon J, Lee YH, Cha GH. Dipenyleneiodonium
Induces Growth Inhibition of Toxoplasma gondii through ROS Induction in ARPE-19
Cells.   Korean   J   Parasitol.   2019;57(2):83-92.   Epub   20190430.   doi:
10.3347/kjp.2019.57.2.83. PubMed PMID: 31104400; PMCID: PMC6526218.

43. Yuan J, Johnson RL, Huang R, Wichterman J, Jiang H, Hayton K, Fidock DA, Wellems TE,
Inglese J, Austin CP, Su XZ. Genetic mapping of targets mediating differential chemical
phenotypes in Plasmodium falciparum. Nat Chem Biol. 2009;5(10):765-71. Epub
20090906.   doi: 10.1038/nchembio.215.   PubMed   PMID: 19734910;   PMCID:
PMC2784992.

44. Pandey M, Singh AK, Thakare R, Talwar S, Karaulia P, Dasgupta A, Chopra S, Pandey AK.
Diphenyleneiodonium chloride (DPIC) displays broad-spectrum bactericidal activity. Sci
Rep. 2017;7(1):11521. Epub 20170914. doi: 10.1038/s41598-017-11575-5. PubMed
PMID: 28912539; PMCID: PMC5599662.

45. An Z, Muthusami S, Yu JR, Park WY. T0070907, a PPAR gamma inhibitor, induced G2/M
arrest enhances the effect of radiation in human cervical cancer cells through mitotic
catastrophe.   Reprod   Sci.   2014;21(11):1352-61.   Epub   20140318.   doi:
10.1177/1933719114525265. PubMed PMID: 24642720; PMCID: PMC4212328.

46. Vizcarra EA, Goerner AL, Ulu A, Hong DD, Bergersen KV, Talavera MA, Le Roch K, Wilson
EH, White MW. An ex vivo model of Toxoplasma recrudescence reveals developmental
plasticity of the bradyzoite stage. mBio. 2023;14(5):e0183623. Epub 2023/09/07. doi:
10.1128/mbio.01836-23. PubMed PMID: 37675999; PMCID: PMC10653814.

47. Scholer N, Krause K, Kayser O, Muller RH, Borner K, Hahn H, Liesenfeld O. Atovaquone
nanosuspensions show excellent therapeutic effect in a new murine model of reactivated
toxoplasmosis. Antimicrob Agents Chemother. 2001;45(6):1771-9. PubMed PMID:
11353624.

48. Singh N, Sharma B. Toxicological Effects of Berberine and Sanguinarine. Front Mol Biosci.
2018;5:21. Epub 20180319. doi: 10.3389/fmolb.2018.00021. PubMed PMID: 29616225;
PMCID: PMC5867333.

49. Huang LJ, Lan JX, Wang JH, Huang H, Lu K, Zhou ZN, Xin SY, Zhang ZY, Wang JY, Dai P,
Chen XM, Hou W. Bioactivity and mechanism of action of sanguinarine and its derivatives
in the past 10 years. Biomed Pharmacother. 2024;173:116406. Epub 20240308. doi:
10.1016/j.biopha.2024.116406. PubMed PMID: 38460366.
50. Wang Q, Sibley LD. Assays for Monitoring Toxoplasma gondii Infectivity in the Laboratory
Mouse. Methods Mol Biol. 2020;2071:99-116. Epub 2019/11/24. doi: 10.1007/978-1-
4939-9857-9_5. PubMed PMID: 31758448.
578

Figure Legends

Figure 1 Development of a High Throughput Screening assay for growth inhibition of Tg68

tachyzoites. (A) A tachyzoite-specific firefly-luciferase (Fluc) reporter strain of Tg68 was

generated using the pTUB1 promoter. (B) Confluent HFF cells in 384-well plates were infected

with Tg68 tachyzoites and luciferase activity was performed on day 3. (C) Replicate of 12 wells

from six independent plates was analyzed for luciferase activity. Fluc expression increased more

than 1000 fold and was found significantly higher at day 3 hr (72hr) compared to day 0 (4 hr

postinfection). Data accumulated from 12 wells, across 6 plates. Mann-Whitney test, $P < 0.02$.

(D) EC_{50} determination for Tg68-pTub1:Fluc and ME49-Fluc treated with serial dilutions of

BRD7929 or Atovaquone. All EC_{50} values are presented as the mean of three biological

replicates (n = 3)

Figure 2 Development of a High Throughput Screening assay for growth inhibition of Tg68
bradyzoites. (A) A bradyzoite-specific Nanoluc luciferase (nLuc) reporter strain of Tg68 was
generated using the p*BAG1* promoter. Parasites were grown in alkaline medium (pH 8.2), CO_2
free or in the absence of glucose supplemented with glutamine, which stimulates in vitro
development of bradyzoites. (B) Confluent HFFs in 384-well plates were infected with Tg68
tachyzoites for two hr, washed and then cultured either in D10 under normal conditions (Tz) or
switch to alkaline or glutamine conditions to induce bradyzoites. The cultures were then
maintained under CO_2-free conditions for 10 days with media changes on day 3 and 6, with a
compound treatment at day 6 and readout at day 10. (C) Luciferase signals from tachyzoites (Tz)
harvested at day 0 (4 hrs postinfection) postinfection or bradyzoites induced for different times
(day 0 (4 hr) to day 10) by culture in alkaline or glutamine media. Comparisons between
sequential time points using Mann Whitney test, **** $P < 0.0001$(D) Determination of EC_{50}
values for BRD7929 and atovaquone treatment of in vitro induced bradyzoites culture in alkaline
or glutamine media. All EC_{50} values are presented as the mean of four biological replicates (n =
4).

Figure 3 Summary of LOPAC screening for growth inhibition of *T. gondii*. (A) Venn diagram showing the number of compounds with ≥ 50% of growth inhibition at 10 μM in each of three growth assays. Tz = tachyzoite growth assay, Alk = alkaline induced bradyzoite growth assay, Gln = glutamine induced bradyzoite growth assay. Red circled numbers indicate the selection criteria for Primary Hits. (B). Summary of LOPAC screen and prioritization of Hits for follow up. Of 44 Primary Hits, 36 were available for dilution series to determine EC_{50} values. Of 9 Top Hits, 8 were available for further biological testing.

A.

B.

Figure 4. Testing of compounds for activity against ex vivo bradyzoites. (A) Schematic for

isolation of bradyzoites from chronically infect mice and testing in vitro during continuous

treatment or after 4 hr treatment and washout. (B) Testing of the Top Hit Sanguinarine and

reference compounds in continuous treatment vs. 4 hr treatment and washout. Values are

determined from plaque counts after 10-14 days of outgrowth and are normalized to the DMSO

control for each condition. Compounds were used at $3XEC_{90}$ based on sensitivity of Tg68Fluc

tachyzoites (**Table S3)**. Data from 2 biological replicates, bar graph represent the average of

percentage of number of plaques in the treatment group normalized to DMSO control.

Figure S1. Testing of additional TOP HIT compounds for activity against ex vivo bradyzoites.

Plaque number from 2 biological replicates for Diphenyleneiodonium, T0070907 and reference

compounds in continuous treatment vs. 4 hr treatment and washout presented in average

percentage with standard deviation bar.