Poly(allylamine)-tripolyphosphate polymeric nanoparticle as an NLRP3-dependent systemic adjuvant for the vaccine development

Gastón P. Rizzo[1], Rodrigo C. Sanches[2], Camila Chavero[1], Daiana S. Bianchi[1], Eugenia Apuzzo[3], Santiago E. Herrera[4], Maximiliano L. Agazzi[5], Irene A. Keitelman[6], Analía S. Trevani[6], Sergio C. Oliveira[2, 7], Omar Azzaroni[3], Paola L. Smaldini[1* †], Guillermo H. Docena[1 †].

[1]Instituto de Estudios Inmunológicos y Fisiopatológicos (IIFP), UNLP, CONICET, asociado a CIC PBA, Facultad de Ciencias Exactas, Departamento de Ciencias Biológicas, La Plata, Argentina (IIFP-UNLP-CONICET).

[2]Department of Biochemistry and Immunology, Institute of Biological Sciences, Federal University of Minas Gerais (ICB-UFMG).

[3]Instituto de Investigaciones Fisicoquímicas Teóricas y Aplicadas (INIFTA), (UNLP, CONICET), 1900 La Plata, Buenos Aires, Argentina.

[4]Instituto de Química de los Materiales, Ambiente y Energía (INQUIMAE), UBA, CONICET, Facultad de Ciencias Exactas y Naturales, Departamento de Química Inorgánica Analítica y Química Física. Buenos Aires, Argentina.

[5]Instituto para el Desarrollo Agroindustrial y de la Salud (IDAS), (UNRC, CONICET), Ruta Nacional 36 KM 601, 5800 Río Cuarto, Argentina.

[6]Laboratorio de Inmunidad Innata, Instituto de Medicina Experimental (IMEX), CONICET, Academia Nacional de Medicina, Buenos Aires, Argentina.

[7]Department of Immunology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil

24   †Shared senior authorship

26   ***Corresponding author:** Paola L. Smaldini; psmaldini@biol.unlp.edu.ar

ABSTRACT

Nanotechnology plays a crucial role in vaccine development and provides the opportunity to design functional nanoparticles (Np) of different compositions, sizes, charges and surface properties for biomedical applications. The present study aims to evaluate a complex coacervate-like Np composed of poly(allylamine hydrochloride) (PAH) and tripolyphosphate (Tpp) as a safe vehicle and adjuvant for systemic vaccines. We investigated the activation of different antigen-presenting cells (APCs) with Np-PAH and its adjuvanticity in Balbc/c and different KO mice that were intraperitoneally immunized with Np-OVA.

We found that Np-PAH increased the expression of CD86 and MHCII and promoted the production and secretion of interleukin-1β (IL-1β) and IL-18 through the inflammasome NLRP3 when macrophages and dendritic cells were co-incubated with LPS and Np-PAH. We evidenced an unconventional IL-1β release through the autophagosome pathway. The inhibition of autophagy with 3-methyladenine reduced the LPS/Np-PAH-induced IL-1β secretion. Additionally, our findings showed that the systemic administration of mice with Np-OVA triggered a significant induction of serum OVA-specific IgG and IgG2a, an increased secretion of IFN-γ by spleen cells, and high frequencies of LT $CD4^+IFN-\gamma^+$ and LT $CD8^+IFN-\gamma^+$. In conclusion, our findings show that PAH-based Np promoted the inflammasome activation of innate cells with Th1-dependent adjuvant properties, making them valuable for formulating of novel preventive or therapeutic vaccines for infectious and non-infectious diseases.

50   **Keywords:** Nanoparticle, Polymeric Poly(allylamine) nanoparticles, Adjuvant,

51   Inflammasome, NLRP3, Systemic vaccines.

INTRODUCTION

Vaccines have prevented millions of deaths in the last decades (mainly in the pediatric population), reduced medical care costs, and allowed us to coexist peacefully with pathogens. The use of adjuvants in vaccines promoted enhanced protection by the induction of robust and perdurable immunity. In this sense, nanotechnology has significantly impacted vaccine development, as nanoparticles are associated with inherent physical properties to protect the antigen cargo and increase immunogenicity and protection[1–3]. Very recently, nanovaccines showed to be safe and effective for immune intervention in COVID-19[4–6].

One of the critical elements in nanoparticle technology for vaccine formulation is polymers. Among these, amine-based polymers are interesting macromolecules since they can act as polycations depending on the pH. Due to this property, polyamine can serve as a building block for the construction of nanoparticles that are stabilized exclusively by soft interactions like electrostatics. An example is the formation of nano-sized complex coacervates by the interaction between polyamines and multivalent salts like citrate, phosphates and polyphosphates[7,8]. Poly(allylamine) hydrochloride (PAH) is one of the most frequently explored polyamines since its composition can be accurately determined and possesses only one type of amine: primary amines. PAH also finds application in oral drug formulations, as ligands for anion exchange chromatography to facilitate the subsequent processing of bio-pharmaceutical products, as reagents in diagnostics, and in formulations of bactericides for use as antibiotics and antiseptics[9–12]. A common complexing agent for developing polyamine-based nanoparticles and soft complexes is phosphate[13,14]. In particular, the tripolyphosphate (Tpp) anion has multiple industrial applications (e.g., food

additive), is readily available, carries enough charge to act as an ionic crosslinker (up to 5

negative charges per ion), and is an FDA- and GRAS-approved chemical. By this, Tpp is

one of the most used crosslinkers for synthesizing soft nanoparticles with biomedical

applications[15]. As we recently demonstrated, the mixture of PAH and Tpp in an aqueous

solution gives rise to the formation of a highly stable colloidal dispersion of complex

coacervates, characterized by size and surface charge that enable the loading of proteins

with different molecular weights and charges[16].

Organic and inorganic nanoparticles, including uric acid crystals, cholesterol crystals, silica

crystals, calcium pyrophosphate crystals (pseudogout), asbestos, and aluminum salts, are

capable of triggering innate immunity through the activation of the multiprotein complex

known as the inflammasome in antigen-presenting cells (APCs)[17,18]. Recently, polyamine-

based platforms, including polyethyleneimine, chitosan, and polylysine, have demonstrated

attractive properties as adjuvants[19–21].

The innate immune response is initiated by recognizing specific conserved molecular

patterns derived from microorganisms (MAMPs and PAMPs) and signals from host tissue

damage (DAMPs). A new designation for molecular patterns has been coined for non-

pathogenic nanoparticles as nanoparticle-associated molecular patterns or NAMPs[17,22].

PAMPs, DAMPs, and NAMPs interact with surface or intracellular pattern recognition

receptors, initiating pro-inflammatory programs that activate innate cells[23,24]. In particular,

NLRs detect NAMPs, and NLRP3 is the most extensively characterized cellular receptor[25].

It is primarily expressed in myeloid lineage cells, such as granulocytes, monocytes,

macrophages, and dendritic cells. Nanoparticles promote lysosomal membrane

permeabilization (LMP), and cathepsins are released into the cytosol[26]. Cathepsins are

99 cysteine proteases primarily found in late endosomes and lysosomes and play crucial roles

100 in various physiological processes, maintaining their proteolytic activity at physiologic pH.

101 These enzymes, mainly cathepsin B, can activate the NLRP3 inflammasome[27]. After the

102 inflammasome activation, the adaptor protein ASC (apoptosis-associated speck-like protein

103 containing a CARD) assembles into large helical fibers interacting with each other and

104 forming aggregates. The recruitment of pro-caspase-1 to ASC promotes the autocatalytic

105 activation of caspase-1, which leads to activating the pro-forms of IL-1β and IL-18 and the

106 pore-forming cytosolic protein gasdermin-D (GSDMD). Upon cleavage of the N-terminal

107 domain of GSDMD, these fragments oligomerize into a ring shape and insert into the

108 plasma membrane, forming a pore through which IL-1β and IL-18 are released. Moreover,

109 GSDMD permeabilizes the cell membrane, leading to ionic imbalance and subsequent

110 osmotic lysis, triggering a form of inflammatory programmed-cell death known as

111 pyroptosis, which can be observed by the release of the enzyme lactate dehydrogenase

112 (LDH) into the extracellular environment[28,29]. However, cytokines can be secreted through

113 an unconventional GSDMD–independent pathway, including intracellular vesicular

114 trafficking (exocytosis of lysosomes or autophagosomes) or microvesicle release without

115 inducing cell death[30,31]. For instance, multivesicular bodies (MVBs), loaded with

116 endosomes containing the cytokine, undergo lysis upon release into the extracellular

117 environment, releasing the soluble cytokine[32,33]. Moreover, there are non-canonical

118 pathways of inflammasome activation that proceed through the proteolytic action of

119 caspase-11[34,35].

120 The possibility that an immunogen encapsulated in PAH-based Np could trigger the

121 immune system by activating innate cells, endowing them with migratory capacities

towards the lymph node to promote the activation of naive B and T cells, makes them

potential candidates for vaccine formulation. The production of cytokines and chemokines,

along with the expression of CD80, CD86, and MHCII on the cell surface, transforms these

cells into APCs with an increased ability to activate T lymphocytes[36–38]. We previously

found that the positive PAH-based nanoparticles are non-toxic and are internalized by

phagocytic cells with intracellular localization in early and late endosomes[16]. In this study,

we found that dendritic cells and macrophages undergo activation upon nanoparticle

uptake, cell migration, and promotion of Th1-dependent adaptive immunity. The release of

IL-1β and IL-18 through the NLRP3 activation triggered the secretion of IFN-γ with the

induction of specific IgG antibodies.

MATERIALS AND METHODS

Preparation of PAH/Tpp nanoparticle

PAH/Tpp nanoparticles were prepared by following the procedure described in Apuzzo et al[16]. Briefly, 0.125 V_f ml of PAH (17.5 kDa, Sigma-Aldrich, St. Louis, MO, USA) 40mM (monomer concentration) at pH 8.5, 0.75 V_f ml of deionized water, and 0.125 V_f ml of Tpp (Sigma-Aldrich, St. Louis, MO, USA) 24mM at pH 8.5 were mixed under constant stirring at room temperature (RT). The addition of Tpp was made very quickly in order to prevent the system from collapsing into a macroscopic phase. After this procedure, a volume V_f of dispersed PAH/Tpp nanoparticles is obtained with a final concentration of PAH and Tpp of 5mM and 3mM, respectively. In this work, the Np concentration was expressed in terms of PAH monomer concentration (i.e., 0.5mM Np means a 1:10 dilution of the original dispersion). The protein-loaded nanoparticle dispersions were prepared by following the same procedure but adding an aqueous solution of chicken egg albumin (OVA) (EndoFit[(TM)] InvivoGen, San Diego, CA, USA) at the water addition stage. The volume of deionized water was re-calculated as $V_{water} = 0.75\,V_f - V_{OVA}$ and the OVA volume was fixed to obtain the desired OVA concentration in the final solution.

The percentage of encapsulated OVA was calculated by assessing the protein content of the supernatant using bicinchoninic acid (BCA) commercial kit (Thermo Fisher Scientific, Waltham, MA, USA).

Cell lines

HEK-hTLR4-reporter (InvivoGen®, San Diego, CA, USA), HT-29 and J774 cell lines were maintained in DMEM medium (Gibco® Thermo Fisher Scientific, Waltham, MA,

USA) containing 10% FBS (Internegocios, Mercedes, Bs As, Argentina), 100U/mL

penicillin and 100µg/mL streptomycin (Gibco® Thermo Fisher Scientific, Waltham, MA,

USA). The THP1-ASC-GFP cell line was maintained in RPMI medium (Gibco® Thermo

Fisher Scientific, Waltham, MA, USA) containing 10% FBS, 100U/mL penicillin and

100µg/mL streptomycin.

Bone marrow-derived macrophages (BMDMs) were cultured in DMEM medium, and bone

marrow-derived dendritic cells (BMDC) were cultured in RPMI medium containing 10%

FBS, 100U/mL penicillin and 100µg/mL streptomycin. Both cells were derived from wild-

type (WT) and knockout (KO) mice with 20% L929-conditioned medium (LCCM) in

DMEM and 20ng/mL recombinant granulocyte-macrophage colony-stimulating factor

(GM-CSF) (R&D Systems, Minneapolis, MN, USA) in RPMI for BMDM and BMDC,

respectively. Briefly, bone marrow cells were obtained from the femur and tibia of WT or

KO mice[39]. Cells were seeded on dish plate $5x10^5$ cell/mL (day 0) and maintained in a

conditioned medium and GM-CSF containing 10% FBS, 100U/mL penicillin and

100µg/mL streptomycin at 37°C, in a 5% CO_2 atmosphere for 7 days. On day 4, the

medium was fully replaced, and cells were stimulated with 0.5mM PAH/Tpp Nps. As a

positive control for inflammasome activation, cells were treated with 5mM Nigericin for

50min, after being primed with 1µg/mL LPS from *E. coli* (LPS E. coli) (Sigma-Aldrich, St.

Louis, MO, USA) for 3h at 37°C. Supernatants were collected and cells were lysed with

25µL/well M-PERR (Thermo Fisher Scientific, Waltham, MA, USA) supplemented with

1:100 protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO, USA). Both supernatants

and cell lysates were stored at −80°C until used.

Cell stimulation with Np and characterization

Flow cytometry for phenotype and cell activation characterization: BMDC were seeded in 24-well microplates at 1×10^6 cells/well and stimulated with ultrapure 1µg/mL LPS from *E. coli* for 3h at 37°C, followed with 0.5mM PAH/Tpp Nps overnight (ON). BMDC were washed with PBS and stained for 1h at 4°C with anti-CD11c (PE), anti-CD11b (PerCP-Cy5.5), anti-MHCII (APC) and anti-CD86 (FITC) (Thermo Fisher Scientific, Waltham, MA, USA). As a positive control for inflammasome induction, cells were primed with ultrapure 1µg/mL LPS from *E. coli* for 3h, followed by 5mM ATP (Sigma-Aldrich, St. Louis, MO, USA) for 3h at 37°C. For inhibition analysis, cells were pre-treated for 30min with 100µg/mL Z-VAD-FMK (Sigma-Aldrich, St. Louis, MO, USA), 1µg/mL cytochalasin D (Santa Cruz Biotechnology Inc, Santa Cruz, CA, USA) or 50µM CA-74ME (Santa Cruz Biotechnology Inc, Santa Cruz, CA, USA) and kept in culture during the experiment. Fluorescence acquisition was performed with a FACS Aria Fusion cytometer, and data was analyzed using FlowJo software.

Reagents for inhibitions: For inhibition analysis, cells were pre-treated for 30min with 100µg/mL Z-VAD-FMK, 1µg/mL cytochalasin D or 50µM CA-74ME, 5mM 3-methyl adenine (Cayman, Ann Arbor, MI, USA) and 1 µM VPS34-IN1 (Cayman, Ann Arbor, MI, USA) and 100nM Bafilomycin A1 (Cayman, Ann Arbor, MI, USA) and kept in culture during the experiment. When indicated, at 4 hours post-Np stimulation, supernatants were removed and replaced with either complete medium or Earle's Balanced Salt Solution (EBSS) for medium starvation.

Nanoparticle internalization analysis: BMDC and BMDM were grown on glass coverslips and stimulated ON with 0.5mM FITC-PAH/Tpp Np. Poly (fluorescein isothiocyanate

allylamine hydrochloride), Mw~56000 (Sigma-Aldrich, St. Louis, MO, USA) was used

instead of PAH in the PAH/Tpp formulation. Cells were fixed for 30min in 4%

paraformaldehyde (PFA) at RT and nuclei were stained with 1µg/mL propidium iodide (IP)

(Sigma-Aldrich, St. Louis, MO, USA). Cells were washed with PBS, blocked with 2%

BSA in phosphate buffer pH=7.4 for 30min at RT and incubated with mouse anti-human

LAMP1 (BD Biosciences, Franklin Lakes, NJ, USA) for 1h at RT, followed with anti-

mouse Alexa 647 (Abcam, Cambridge, UK) for 1h at RT. Cells were washed, coverslips

were mounted onto a slide containing 10µL of fluorescence mounting media (Dako,

Abcam, Cambridge, UK) and allowed to dry at 4°C ON. Fluorescence was analyzed with a

Leica SP5 Confocal Microscope.

Internalization inhibition assay: BMDC and J774 were stimulated with 0.5mM FITC-

PAH/Tpp Nps and phagocytosis was inhibited with 1µg/mL cytochalasin D. Cells were

characterized by flow cytometry with anti-CD11b (Percp-Cy5.5) and anti-F4/80 (APC)

(Thermo Fisher Scientific, Waltham, MA, USA) and confocal microscopy for intracellular

fluorescence. Nuclei were stained with 1µg/mL IP.

Intracellular immunostainings for the localization of IL-1β in autophagosomes: The J774

cell line was grown on glass coverslips and prime for 3h at 37°C with 1µg/mL LPS from E.

coli, followed with 0.5mM PAH/Tpp Np for 6h. After fixation with PFA 4% for 30 min,

cells were blocked with PBS-glycine (0.1 M) for 15 min, permeabilized with acetone

(−20°C) for 7min, and blocked with PBS-BSA 5% for 1h at RT. Then, macrophages were

incubated with the rabbit polyclonal anti-LC3B (Proteintech, Rosemont, IL, USA) and

mouse polyclonal anti-IL-1β (Cell Signaling Technology, Danvers, MA, USA) in blocking

buffer ON at 4°C and then incubated with the secondary anti-rabbit Alexa 594 (Abcam,

Cambridge, UK) and anti-mouse Alexa 488 (Invitrogen, Thermo Fisher Scientific,

Waltham, MA, USA) antibodies for 1h at RT. Nuclei were stained with methyl green

(Sigma-Aldrich, St. Louis, MO, USA) and mounted onto a slide containing fluorescence

mounting media (Dako, Abcam, cambrige, UK). Image acquisition was performed with an

SP5 Leica Confocal Microscope.

Epithelial cell stimulation: HT-29 cells were maintained in DMEM containing 10% FBS,

100U/mL penicillin and 100μg/mL streptomycin at 37°C with 5% CO_2, and $3x10^4$

cells/well were seeded in 96-well microplates. Cells were incubated ON to 70% confluence

and the medium was fully replaced. Then, cells were stimulated ON with 0.5mM FITC-

PAH/Tpp Np, or with 50ng/mL TNF-α (R&D Systems, Minneapolis, USA) for 30min at

37°C followed with FITC-PAH/Tpp Np. A mouse monoclonal anti-human EPCAM (Santa

Cruz Biotechnology, Santa Cruz, CA, USA) antibody, followed bywith an anti-mouse

Alexa 647 conjugated antibody (Abcam, Cambridge, UK), was used for epithelial cell

detection. Cell fluorescence was analyzed with an SP5 Leica Confocal Microscope.

Inflammasome activation assessment: THP1-ASC-GFP were maintained in RPMI medium

containing 10% FBS, 100U/mL penicillin and 100μg/mL streptomycin at 37°C in a 5%

CO_2 atmosphere and plated at $3.6×10^5$ cells/well in 96-well plates. Cells were stimulated

with 0.5mM PAH/Tpp Np or primed with ultrapure 1μg/mL LPS from *E. coli* followed

with 0.5mM PAH/Tpp Np ON. For a positive control, cells were primed with 1μg/mL LPS

from *E. coli* for 3h and transfected with 500ng/mL bacterial plasmid cDNA. pcDNA was

generated by cloning the PCR-generated full-length cDNA from a random non-related

gene. Transfection was carried out with Lipofectamine[TM] LTX Reagent with PLUS TM

Reagent (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA), following the

manufacturer´s instructions. Supernatants were harvested and stored at −80°C until used

and cells were stained with DAPI (nuclear staining) (Invitrogen, Thermo Fisher Scientific,

Waltham, MA, USA) and fixed with 2% PFA fixation for the microscopy analysis.

The frequency of ASC-GFP$^+$ cells and localization of fluorescent ASC specks were

assessed using a Nikon Eclipse Ti Fluorescence Microscope and analyzed with ImageJ

software. The total cells: ASC-GFP$^+$ cells ratio was calculated.

Quantification of secreted cytokines: Supernatants of stimulated BMDM, J774, BMDC,

THP1, THP1-ASC-GFP and HT-29 cells were assessed for mIL-1β, mIL-18, mIL-6, mIFN-

γ and mTNF-α using the Mouse DuoSet R&D Systems ELISA kit. The concentration of

hIL-1β and hIL-8 were measured using a human Invitrogen ELISA Kit (Thermo Fisher

Scientific, Waltham, MA, USA). All the protocols were used according to the

manufacturer's instructions. The concentration of released LDH (enzymatic assay) (LDH-L

Weiner Lab, Rosario, Santa Fe, Argentina) was quantified.

Immunoblotting analysis: Cell lysates and supernatants were analyzed by SDS-PAGE

followed by immunoblotting. Proteins suspended in loading buffer (SDS+β-

mercaptoethanol) were separated on a 15% SDS-PAGE gel, and transferred onto

nitrocellulose membranes (Amersham Biosciences, Uppsala, Sweden) using transfer buffer

(50mM Tris, 40mM glycine, 10% methanol). After blocking for 1h in TBS with 0.1%

Tween-20 containing 5% non-fat dry milk, membranes were incubated overnight at 4°C

with a mouse monoclonal antibody to the p20 subunit of caspase-1 (Adipogen, San Diego,

CA, USA) at a 1:1,000 dilution. A loading control blot was performed with an anti–β-actin

monoclonal antibody (Cell Signaling Technology, Danvers, MA, USA) at a 1:1,000

dilution. Following three washes in TBS with 0.1% Tween 20, membranes were incubated

for 1h at RT with the appropriate HRP-conjugated secondary antibody (Cell Signaling
Technology, Danvers, MA, USA) at a 1:1,000 dilution. Immunoreactive bands were
visualized using Luminol chemiluminescent HRP substrate (Millipore, Burlington, MA,
USA) and analyzed with ImageQuant TL Software (GE Healthcare, Buckinghamshire,
UK).

Analysis of the LPS content of PAH/Tpp nanoparticles

The HEK-hTLR4-reporter cell line was maintained in DMEM containing 10% FBS,
100U/mL penicillin and 100μg/mL streptomycin at 37°C with 5% CO_2. Thirty thousand
cells/well were seeded in 96-well microplates and incubated to 70% confluence. The
medium was fully replaced and cells were stimulated ON with different concentrations of
LPS, 0.5mM PAH, 0.3mM Tpp, or 0.5mM PAH/Tpp Np. The supernatants were harvested
and the secreted embryonic alkaline phosphatase (SEAP) was measured.

In vivo **assessment of adjuvant properties**

Inflammasome characterization: Male six to nine week-old wild-type and KO C57BL/6
Nlrp3$^{-/-}$, Casp1/11$^{-/-}$, Casp11$^{-/-}$, Gsdmd$^{-/-}$ and IL-1R$^{-/-}$ mice were purchased from the Federal
University of Minas Gerais and maintained under 12h cycles of light/dark. Femur and tibiae
were removed for BMDM preparation and for immunization assays. Furthermore, male five
to seven week-old wild-type Balb/c mice were purchased from the Laboratory of
Experimental Animals, School of Veterinary Sciences of the University of La Plata
(UNLP), Argentina.

Mice (C57BL/6 and Balb/c wild-type mice, and null C57BL/6 mice) were immunized once

a week for three consecutive weeks intraperitoneally (IP) with OVA-conjugated PAH/Tpp

Np (Np-OVA) or OVA adjuvanted with aluminum hydroxide (OVA+Alum). As a control,

mice were immunized only with 100μg OVA EndoFit$^{(TM)}$. One week after the last

immunization, mice were bled and sacrificed for the immune response analysis.

Humoral response assessment: OVA-specific antibodies were measured by indirect ELISA

in different biological samples. Briefly, MaxiSorp plates (NUNC, Roskilde, Denmark) were

coated with 1μg/mL OVA, 5mM PAH or 5mM Np in carbonate-bicarbonate buffer,

pH=9.6. Coated plates were blocked with 5% equine serum in saline for 1h at 37°C, and

subsequently incubated with sera (1/100) for 1h at 37°C. Bound immunoglobulins were

detected with anti-mouse IgG-HRP (Biorad, Hercules, CA, USA), anti- mouse IgG1-HRP,

anti-mouse IgG2a-HRP (BD Biosciences, Franklin Lakes, NJ, USA) or anti-mouse IgA-

HRP (Thermo Fisher Scientific, Waltham, MA, USA). The reaction was developed with

TMB (Sigma-Aldrich, St. Louis, MO, USA) and stopped with 2M H_2SO_4. OD was

measured at 450 nm with Varioskan Lux (Thermo Fisher Scientific).

Assessment of secreted cytokine from stimulated splenocytes: Spleen cell suspensions were

prepared in RPMI-1640 supplemented with 10% FBS, 10U/mL penicillin and 10mg/mL

streptomycin, seeded ($4x10^6$ cells/mL) in culture plates and cultured for 72h at 37°C under

controlled atmosphere. Cells were recalled with 10μg/mL OVA, and supernatants were

harvested for mIFN-γ, mIL-5 and mIL-13 quantification by ELISA, using commercially

available kits (R&D Systems, Minneapolis, MN, USA) and following the manufacturer´s

instructions.

Cytokines were also assessed by intracytoplasmic flow cytometry in spleen cells.

Splenocytes were incubated with brefeldin A (BD Biosciences, Franklin Lakes, NJ, USA)

for the last 4h of culture and stained with anti-CD4 (PerCP-Cy5.5) anti-CD8 (APC)

monoclonal Ab (Thermo Fisher Scientific, Waltham, MA, USA). Cells were fixed,

permeabilized, and stained with an anti-mouse IFN-γ (FITC) (Thermo Fisher Scientific,

Waltham, MA, USA). Sample acquisition was performed with a FACS Aria Fusion

cytometer, and data were analyzed using the FlowJo software.

Ethics: All experimental protocols were conducted in strict agreement with international

ethical standards for animal experimentation (Helsinki Declaration and its amendments,

Amsterdam Protocol of Welfare and Animal Protection and National Institutes of Health,

USA NIH, guidelines: Guide for the Care and Use of Laboratory Animals). Anesthetized

mice (isoflurane 5%) were killed by cervical dislocation by experienced research personnel,

who performed it humanely and effectively. All efforts were made to alleviate suffering

during the whole experiment. All protocols were approved by the Institutional Committee

for the Care and Use of Laboratory Animals of the School of Sciences (University of La

Plata) (Protocol Number: 006-37-21).

Statistical analysis

All experiments were repeated at least twice with consistent outcomes. Flow cytometry

data were analyzed using FlowJo X, images obtained by microscopy were analyzed using

ImageJ. Graphs and data analysis were performed using GraphPad Prism8. Statistical

analyses were carried out using one- or two-way ANOVA, followed by the Tukey test to

discern significant differences among measurements from different experimental groups or

by the t-Student test when applicable. Quantitative data are expressed as mean ± SEM, and

statistical significance was defined as $p < 0.05$.

RESULTS

Phagocytic cells internalized the nanoparticles and became activated

The positively charged PAH was mixed with the negatively charged crosslinker Tpp at a

ratio R=[Tpp]/[PAH]=0.6 (Figure 1A). The physicochemical properties of the self-

assembled spherical nanoparticles, with a diameter of 200 nm and a negative charge

potential of -23 mV, were measured using transmission electron microscopy (TEM) (Figure

1B) and dynamic light scattering (DLS) (Figure 1C)[16]. To study the interaction of PAH/Tpp

Nps with cells, we exposed the murine J774 macrophages to Np. We observed the

internalization and cellular distribution of fluorescent Nps by confocal microscopy after 4h

of incubation. Furthermore, we did not detect any internalization of the polymer PAH-FITC

(without Tpp) in macrophages (Figure S1). We observed that nanoparticles were well-

dispersed into cytoplasmic vesicles (fluorescent granules) and the cytosol (diffuse

fluorescence) (Figure 1D). Then, we analyzed the co-localization of the Np-FITC with

LAMP1 to define the Np localization in vesicles. As it can be observed in Figure 1D, Nps

were distributed between LAMP1$^+$- and LAMP1$^-$-vesicles, likely late and early endosomes

(yellow and green vesicles, respectively). The incubation of cells with the inhibitor of

phagocytosis cytochalasin D (Cyto D) confirmed that the internalization of Np was

dependent on phagocytosis. Additionally, we detected lower fluorescence by flow

cytometry in macrophages F4/80$^+$ CD11b$^+$ (Figure S1C) incubated with Cyto D than in

cells exposed to Np-FITC (Figure 1E). Next, bone marrow-derived dendritic cells were

exposed to Np and LPS and we analyzed the surface expression of Class II MHC and CD86

as cell activation markers (Figure 1F). We found that BMDC expressed significantly higher

levels of Class II MHC and CD86 when exposed to LPS or Np than cells exposed to

nanoparticles or medium. The LPS used as a positive control for dendritic cell activation

resulted in CD86 and Class II MHC expression similar to that in Np-exposed dendritic

cells. These findings indicate that the cells recognized and were activated by the Np.

A similar analysis of internalization was conducted using a human intestinal epithelial cells

line (HT-29). EPCAM was employed as a marker for epithelial cells, and various z-series

were acquired by confocal microscopy. Neither intracellular fluorescence nor hIL-8

secretion was observed when epithelial cells were incubated with Np. Pre-activated cells

with TNF-α (20ng/mL) rendered no Np up-take (Figure S2A) and production of hIL-8

(Figure S3B). As a first approach to investigate the Np-driven secretion of mIL-1β in a

phagocytic cell line, we performed a LPS (1µg/mL) prime stimulus follow by a dose-

response of Np. We found that 0.5mM Np promoted the highest level of secreted mIL-1β

(Figure S2C).

We next investigated the production of pro-inflammatory cytokines by phagocytic cells

exposed to 0.5mM of Np or controls (Figure 2). We found a significantly higher level of

IL-1β in murine J774 macrophages and BMDCs, and in human THP-1 monocytes exposed

to LPS+Np than cells exposed to LPS or Np (Figure 2A). We found that the different cells

exposed to Np did not significantly increase IL-1β concentration compared with the

medium; LPS+ATP was used as a positive control for IL-1β secretion[40]. These findings

collectively suggest that Np promoted IL-1β secretion and LPS was necessary.

Nevertheless, the quantification of secreted mIL-6 and hIL-8 showed no significant changes

when cells were incubated with LPS or LPS+Np, indicating that the cytokine secretion was

independent of the Np presence (Figure 2B). Similarly, the cytokine levels were basal when

the cells were exposed to Np (Figure 2B). Furthermore, when we analyzed mIL-18, we

observed a differential secretion in J774 macrophages and BMDC incubated with LPS+Np.

Cells exposed to medium, LPS or Nps showed basal levels, whereas LPS+ATP or LPS+Np

induced the highest concentration of mIL-18 in BMDC (Figure 2C). These findings

indicate that the secretion of IL-1β and IL-18 in APCs required LPS and Np.

Nanoparticles activated the intracellular inflammasome pathway

Inflammasomes, such as NLRP3, require the adaptor molecule ASC to activate caspase-1[41].

Following inflammasome activation, ASC assembles into a large protein complex, which

can be visualized as a "speck"[42]. Representative pictures are shown with white arrowheads

indicating GFP-specks (Figure 3A). The number of specks in LPS+Np-treated cells was

significantly higher (p ≤0.05) than in cells exposed to medium or LPS and equal to the

positive control with LPS+DNA (Figure 3B). This finding suggests that Np triggered the

assembly of the ASC complex. A similar analysis was performed using flow cytometry to

quantify ASC$^+$ cells[43], and it was observed that cells exposed to LPS+DNA or LPS+Np had

a significantly higher frequency of ASC$^+$ cells compared to resting cells (Figure 3C and D).

The ratio of GFP-ASC$^+$ cells:total cells triggered by Np was similar to the positive control

condition of LPS+DNA. Additionally, the quantification of hIL-1β in the supernatants

showed that LPS+DNA or LPS+Np induced higher levels of this cytokine than medium or

LPS (p<0.05) (Figure 3E). To investigate the inflammasome activation, we used different

pathway inhibitors. We found that cytochalasin D, the cathepsin B inhibitor CA-074Me and

the caspase-1 inhibitor Z-VAD-FMK significantly suppressed the mIL-1β secretion

induced by LPS+Np (p<0.001) (Figure 3F). The analysis of the mIL-18 secretion showed

similar results, with a higher secretion in cells exposed to LPS+Np than in cells incubated

with LPS or medium, and the inhibitors suppressed the cytokine secretion (Figure 3F). Our

results suggest that Np promoted inflammasome activation through cathepsin B and

caspase-1. These findings led us to perform an experiment using inflammasome-null mice

for NLRP3, caspase-1/11, caspase-11 and gasdermin D (GSDMD). BMDM obtained from

NLRP3$^{-/-}$, caspase1/11$^{-/-}$, caspase11$^{-/-}$ and GSDMD$^{-/-}$ mice were incubated with 1µg/mL

LPS and 0.5mM of Np. To analyze the inflammasome activation, we quantified secreted

mIL-1β and mTNF-α by ELISA in the supernatants. We observed no mIL-1β secretion in

cells stimulated with LPS or Np in wild-type and KO mice. However, exposure to LPS+Np

induced the production and release of this cytokine in wild-type, caspase11$^{-/-}$ and GSDMD^{-}

$^{/-}$ BMDM. In contrast, cells from NLRP3$^{-/-}$ and caspase1/11$^{-/-}$ mice showed decreased mIL-

1β secretion compared to wild-type animals (p<0.001) (Figure 4A). When the BMDM WT

and caspase11$^{-/-}$ were stimulated with the positive control LPS+Nigericin, we detected the

production of IL-1β. Meanwhile, the secretion of IL-1β in BMDM from NLRP3$^{-/-}$,

caspase1$^{-/-}$ and GSDMD$^{-/-}$ was significantly abrogated (Figure 4A). The quantification of

soluble mTNF-α showed that cells from wild-type and KO mice secreted similar high

amounts upon exposure to LPS, LPS+Np or LPS+Nigericin. No TNF- α was induced when

cells were incubated with Np (Figure 4B). LDH measured in the supernatants showed that

the stimuli did not significantly affect the cells. The consistent level of soluble LDH in cells

exposed to the medium, Np or LPS+Np indicated that cell viability was unaffected (Figure

S3A). Additionally, the lysate of LPS+Np-exposed cells from wild-type or KO mice

showed a minimal difference in the activation of caspase-1 (20 kDa) between cells derived

from wild-type and NLRP3$^{-/-}$ mice. Inactive caspase-1 was detected in resting and

LPS+Np-stimulated cells, while it was absent in samples from cells of caspase1/11$^{-/-}$

animals (Figure S3B).

As mentioned before, we found that the release of mIL-1β was independent of GSDMD

(Figure 4A). This prompted us to study the secretion of IL-1β and IL-18. Among the

multiple pathways described in myeloid cells, we investigated autophagy as responsible for

the secretion of IL-1β. As starvation has been reported to induce autophagy[44,45], we

performed experiments using the EBSS starvation medium following cell activation. We

observed a significant increase in the production of mIL-1β when macrophages were

stimulated with LPS+Np and supplemented with EBSS medium compared to the cells that

were stimulated with LPS+Np (Figure 4C). To confirm this finding, we used 3-

methyladenine (3-MA) (blocker of early stages of autophagy) and VPS34-IN1 (inhibitor of

the class III PI3K) as autophagy inhibitor. All treatments significantly decreased the mIL-

1β secretion in cells exposed to LPS+Np (Figure 4D). To confirm if autophagy is involved

in the secretion of mIL-1β, we measured a pro-inflammatory cytokine (mIL-6) that is

release through the canonical ER-Golgi and autophagy-independent pathway. We observed

unchanged high levels of mIL-6 in cells exposed to LPS+Np irrespective of the presence of

EBSS (Figure S3C). However, when the different inhibitory conditions were assessed, we

found that 3-MA led to a not significantly decreased cytokine level (Figure S3D).

Nevertheless, it is essential to note that this effect was negligible compared to that exerted

on mIL-1β secretion (Figure 4D). Additionally, we did not observe a significant increase in

the release of LDH in the cells, either with or without the autophagy inhibitors (Figure

S3E). Moreover, we observed the same results in BMDCs: the release of IL-1β was

abrogated after the treatment with the autophagy inhibitors (Figure S3F). Finally, we found

by confocal microscopy that mIL-1β co-localized with LC3B, a central protein in the

autophagy pathway and autophagosome biogenesis, when macrophages were stimulated

with LPS+Np (Figure 4E) compared to cells incubated with medium, Np or LPS (Figure

S3G).

Overall, these findings confirmed that the NLRP3-dependent inflammasome pathway was

involved in cell activation with IL-1β production, and that autophagy was involved in the

secretion of IL-1β induced by LPS+Np. However, considering the reduced level of mIL-1β

in cells from NLRP3[-/-] mice compared with wild-type animals and the active caspase-1

observed in lysates of NLRP3[-/-] cells (Figure S3B), we suggest that an NLRP3-independent

inflammasome pathway may also be involved in mIL-1β production. Besides, the

consistent level of soluble LDH in cells exposed to the medium, Np or LPS+Np indicated

that cell viability was unaffected upon activation.

The systemic administration of Np-OVA triggered a specific humoral and cellular immune

response

To characterize its adjuvanticity *in vivo*, we first measured the presence of LPS in PAH,

Tpp and Np with the human TLR4 HEK reporter cells and secreted SEAP, as readout. We

found that the SEAP activity were comparable to that in non-stimulated cells and lower

than 0.01 ng/mL of LPS (Figure S4). The percentage of OVA encapsulation by Np depends

on the final concentration, with a higher yield obtained at a concentration of 100µg/mL

(Figure S5).

Then, mice were intraperitoneally injected with Np-OVA in 3 weekly doses (Figure. 5A).

As a control, we administered a formulation containing the approved adjuvant for human

use, aluminum hydroxide (Alum)[46], with OVA. No weight loss was observed during the

whole experiment in mice (Figure 5B). We did not observe specific antibodies to the

nanoparticle or the polymer PAH in animals immunized with Np-OVA (Figure 5C).

Animals that received Np-OVA produced higher levels of serum OVA-specific IgG than

mice injected with OVA ($p<0.01$) and similar values compared with mice that received

OVA+Alum (Figure. 5D). The subclass analysis revealed that Np-OVA induced higher

titers of OVA-specific IgG2a than in mice injected with OVA+Alum (Figure 5E); in

addition, the ratio of IgG subclasses in Np-OVA and OVA+Alum injected mice showed a

higher production of OVA-specific IgG2a in Np-OVA immunized animals (Figure 5F). To

analyze the T cell-mediated immune response, spleen cells from mice immunized with Np-

OVA, OVA+Alum and OVA were stimulated with OVA and cytokines were measured in

the supernatants. We found that mice immunized with Np-OVA induced increased levels of

IFN-γ when cells were stimulated with OVA ($p<0.001$). Cells from mice that received

OVA or OVA+Alum did not secrete IFN-γ (Figure 5F). On the other hand, the Th2

cytokines mIL-5 and mIL-13 showed decreased concentration in the Np-OVA treated mice

as compared to mice administered with OVA+Alum (Figure 5F). To further evaluate the

cell source of IFN-γ, spleen cells were analyzed by intracellular cytokine staining following

in vitro OVA stimulation. We found that the frequency of $CD4^+IFN-\gamma^+$ and $CD8^+IFN-\gamma^+$

lymphocytes was significantly increased in mice immunized with Np-OVA (Figure 5G and

H). Overall, these findings support that Np-OVA promoted the induction of a Th1-

dependent immune response with specific IgG and IgG2a, and $CD4^+IFN-\gamma^+$ and $CD8^+IFN$-

γ^+ T cells following the immunization with OVA adjuvanted with Np.

To investigate the role of the inflammasome pathway in promoting the specific immune

response, C57BL/6 mice wild-type and NLRP3$^{-/-}$, caspase1/11$^{-/-}$ and IL-1R$^{-/-}$ mice were

intraperitoneally immunized with Np-OVA. As shown in Figure 6A, NLRP3$^{-/-}$ and

caspase1/11$^{-/-}$ mice exhibited lower levels of serum OVA-specific IgG than wild-type mice

($p<0.01$), whereas IL1-R null mice showed similar levels than wild-type mice (Figure 6A).

Finally, spleen cells from wild-type and KO mice were stimulated with OVA or medium,

and IFN-γ levels were assessed in the supernatants. We found that cells from wild-type

mice sensitized with Np-OVA showed higher levels of IFN-γ than cells from WT mice

sensitized with OVA, and NLRP3$^{-/-}$ and caspase1/11$^{-/-}$ mice sensitized with Np-OVA. Cells

from IL1-R$^{-/-}$ mice sensitized with Np-OVA showed a similar concentration of IFN-γ to

WT animals (Figure 6B). These findings showed that the NLRP3 pathway was critical to

promoting the OVA-specific Th1-dependent immune response.

DISCUSSION

Our findings showed that Np are exclusively internalized by phagocytic antigen-presenting cells, specifically dendritic cells and macrophages. Subsequently, they localize intracellularly within the endosome compartment, causing their disruption and leading to the translocation of their components into the cytosol. Thus, the internalization of Np triggers the cellular activation and inflammasome assembly, releasing the pro-inflammatory cytokines IL-1β and IL-18 to the extracellular medium. Remarkably, we did not observe the Np internalization in non-phagocytic mucosa epithelial cells. The permeabilization of the lysosomal membrane is one of the most frequently described mechanisms for activating the inflammasome by particles and Np[47,48]. We used several inhibitors to confirm the role of the different components of the pathway, and we corroborated the participation of cathepsin B as the early activator of the inflammasome[49,50]. Then, using BMDM KO for receptors and enzymes involved in this pathway, we confirmed that cellular activation depended on the NLRP3 receptor and caspase-1 but not on caspase-11. These findings demonstrate that the Np activates the NLRP3 inflammasome through the canonical pathway. Nevertheless, we found that the release of IL-1β was independent of gasdermin D pores, which is associated with the slow release of the cytokine and the absence of cellular death by pyroptosis[51]. The literature shows that the non-canonical autophagy pathway is involve in the secretion of inflammatory molecules such as IL-1β and HMGB1[52,53]. In macrophages and monocytes, LC3B is essential for the efficient secretion of IL-1β in response to lysosomal damage[45,54]. In this sense, we demonstrated that the inflammasome activation proceeded with autophagy induction for secreting IL-1β. All together these findings demonstrated that PAH-based Np activated phagocytic cells with increased expression of MHC II and CD86, along with the

secretion of IL-1β and IL-18, without inducing cell death, which make them a candidate

adjuvant to trigger an antigen-specific immune response.

It is known that aluminum salts, such as the extensively used in human vaccines aluminum

hydroxide (alum), can enhance the effectiveness of the immune response by activating

antigen-presenting cells through the depot effect and the aggregation of antigens to improve

the uptake and activation of APCs. Furthermore, it has been described that alum triggered

the NLRP3-dependent inflammasome pathway to further promote adaptive immunity and

memory[55]. Thus, we evaluated the *in vivo* adjuvant capacity of PAH/Tpp Np in mice that

were systemically administered with OVA encapsulated in Np. We observed the induction

of innate immunity via the inflammasome-dependent pathway to further promote a Th1-

driven adaptive humoral and cellular immunity. Our findings showed the induction of

serum IgG and IgG2a-specific antibodies, and the expansion of $CD4^+$- and $CD8^+$-IFN-γ-

producing T cells. Furthermore, we confirmed *in vivo* that the Th1-dependent isotype

synthesis and IFN-γ production were NLRP3- and caspase-1-dependent. It is reported that

the NLRP3 inflammasome pathway initiates a caspase-1–dependent IL-1β and IL-18

secretion promoting interferon-γ production and Th1 differentiation[56,57]. Similar to other

adjuvants, such as Alum and QS-21, PAH-based nanoparticles along with LPS as a first

inflammasome activation signal induced *in vitro* the NLRP3-ASC-caspase-dependent IL-1β

production. *In vivo*, however, the first signal may be triggered by a DAMP, such as uric

acid, generated at the immunization site[58,59]. Additionally, IL-18, constitutively expressed

by several cell types, can stimulate IL-1β transcription and may initiate a pro-inflammatory

cascade during vaccination[60]. The *in vivo* induction of OVA-specific $CD8^+$T cells indicates

that Np-OVA triggered the cross-presentation of OVA, which may proceed upon the

endosome is disrupted and the content is translocated into the cytosol. Further studies are

needed to confirm if macrophages and dendritic cells contribute to this mechanism, which

is critical when considering using PAH-based Np as an adjuvant in an anti-tumoral or viral

vaccine.

Muñoz-Wolf et al. reported that polymeric nanoparticles made of poly-lactic-co-glycolic

acid with a diameter of 50-60nm elicited a robust cellular immune response through the

caspase-11- and the pyroptosis effector gasdermin D-dependent non-canonical

inflammasome pathway. They described a division of labor for IL-1β and IL-18 to induce

the appropriate immunogenic context for Th1 and CD8[+] T cell induction, respectively[61]. In

our study, we characterized a polymeric nanoparticle of PAH that activated murine and

human macrophages and dendritic cells through the canonical caspase-1- and NLRP3-

dependent inflammasome pathway with the secretion of IL-1β and IL-18, without cell

death, and independent of caspase-11 and gasdermin D-dependent pyroptosis[62,63]. We

found that PAH-based Np induced autophagy for the secretion of the pro-inflammatory

cytokine IL-1β.

In conclusion, we characterized a novel polyallylamine-based self-assembled nanoparticle

as a systemic safe adjuvant that targeted the encapsulated OVA immunogen to phagocytic

cells, endowing these antigen-processing cells with T cell activation capacity that triggered

a Th1-dependent humoral and cellular immune response. These findings may pave the way

for designing and developing novel nanoparticle-based systemic vaccines for infectious and

non-infectious diseases.

CONFLICT OF INTEREST

None of the authors has a financial, commercial or personal relationship that could inappropriately influence or bias the content of the paper.

AUTHOR CONTRIBUTIONS

GPR Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing**.** *RCS* Data curation, Methodology. *CC* Data curation, Formal analysis, Investigation, Methodology**.** *DB* Data curation, Formal analysis, Investigation, Methodology**.** *EA* Data curation, Methodology**.** *SEH* Data curation, Methodology, Writing – original draft, Writing – review & editing**.** *MLA* Data curation, Methodology**.** IK Formal analysis, Methodology, Writing-review & editing. AT Formal analysis, Methodology, Writing-review & editing. *SCO* Supervision, Visualization, Writing – original draft, Writing – review & editing**.** *OA* Supervision, Visualization, Writing – original draft, Writing – review & editing**.** *PLS* Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing**.** *GHD* Conceptualization, Formal analysis, Funding acquisition, Investigation, Supervision, Visualization, Writing – original draft, Writing – review & editing.

FUNDINGS

This research was supported by grants from the Universidad Nacional de La Plata [UNLP,

grant No. 11/X695] and the Agencia Nacional de Promoción Científica y Tecnológica

[ANPCyT, grant No. PICT 2019-0368 and 2020-3166] to PLS and GHD, Fundação de

Amparo à Pesquisa do Estado de São Paulo (FAPESP) grant No 2023/02577-5 to SCO and

National Institutes of Health (NIH) grant No R01 AI116453

ACKNOWLEDGMENTS

SHE, MLA, AT, OA, PLS and GHD are researchers of CONICET; RCS and SCO are

researchers of *Federal University of Minas Gerais and University of São Paulo; GPR, DB,*

EA and IK are fellows of CONICET; CC is fellow of ANPCyT.

REFERENCES

1. Zhao, L. *et al.* Nanoparticle vaccines. *Vaccine* **32**, 327–337 (2014).

2. Irvine, D. J. & Read, B. J. Shaping humoral immunity to vaccines through antigen-displaying nanoparticles. *Curr. Opin. Immunol.* **65**, 1–6 (2020).

3. Huang, M., Zhang, M., Zhu, H., Du, X. & Wang, J. Mucosal vaccine delivery: A focus on the breakthrough of specific barriers. *Acta Pharm. Sin. B* **12**, 3456–3474 (2022).

4. Walls, A. C. *et al.* Elicitation of Potent Neutralizing Antibody Responses by Designed Protein Nanoparticle Vaccines for SARS-CoV-2. *Cell* **183**, 1367-1382.e17 (2020).

5. Arunachalam, P. S. *et al.* Adjuvanting a subunit COVID-19 vaccine to induce protective immunity. *Nature* **594**, 253–258 (2021).

6. Dalvie, N. C. *et al.* Engineered SARS-CoV-2 receptor binding domain improves manufacturability in yeast and immunogenicity in mice. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

7. Herrera, S. E. *et al.* Polyelectrolyte-multivalent molecule complexes: physicochemical properties and applications. *Soft Matter* (2023) doi:10.1039/d2sm01507b.

8. Herrera, S. E. *et al.* Polyelectrolyte–Ion Complexes as Stimuli-Responsive Systems for Controlled Drug Delivery. in *Supramolecular Nanotechnology* 729–764 (John Wiley & Sons, Ltd, 2023). doi:https://doi.org/10.1002/9783527834044.ch27.

9. Lawrence, P. G., Patil, P. S., Leipzig, N. D. & Lapitsky, Y. Ionically Cross-Linked

Polymer Networks for the Multiple-Month Release of Small Molecules. *ACS Appl.*

Mater. Interfaces **8**, 4323–4335 (2016).

10. Goldsmith, D. R., Scott, L. J., Cvetković, R. S. & Plosker, G. L. Sevelamer

hydrochloride: A review of its use for hyperphosphataemia in patients with end-stage

renal disease on haemodialysis. *Drugs* **68**, 85–104 (2008).

11. Aoyama, S., Matsumoto, Y., Mori, C. & Sota, K. Application of novel mixed mode

chromatography (MMC) resins having a hydrophobic modified polyallylamine

ligand for monoclonal antibody purification. *J. Chromatogr. B* **1191**, 123072 (2022).

12. Chang, Y. C. *et al.* Rapid detection of gut microbial metabolite trimethylamine n-

oxide for chronic kidney disease prevention. *Biosensors* **11**, (2021).

13. Herrera, S. E. *et al.* Layer-by-Layer Formation of Polyamine-Salt

Aggregate/Polyelectrolyte Multilayers. Loading and Controlled Release of Probe

Molecules from Self-Assembled Supramolecular Networks. *Macromol. Chem. Phys.*

220, 1–10 (2019).

14. Cuenca, V. E. *et al.* Polyphosphate Poly(amine) Nanoparticles: Self-Assembly,

Thermodynamics, and Stability Studies. *Langmuir* **35**, 14300–14309 (2019).

15. Bugnicourt, L. & Ladavière, C. Interests of chitosan nanoparticles ionically cross-

linked with tripolyphosphate for biomedical applications. *Prog. Polym. Sci.* **60**, 1–17

(2016).

16. E. Apuzzo, M. Agazzi, S. Herrera, A. Picco, G. Rizzo, C. Chavero, D. Bianchi, P.

Smaldini, L. Cortez, W. Marmisollé, G. Padula, A. Seoane, M. Alomar, M.

656   Denofrio, G. Docena, O. A. Poly (allylamine) -tripolyphosphate ionic assemblies as

657   nanocarriers : friend or foe? (2023) doi:10.1021/acsabm.3c00489.

658 17. Boraschi, D. *et al.* Nanoparticles and innate immunity: new perspectives on host

659   defence. *Semin. Immunol.* **34**, 33–51 (2017).

660 18. Jo, E. K., Kim, J. K., Shin, D. M. & Sasakawa, C. Molecular mechanisms regulating

661   NLRP3 inflammasome activation. *Cell. Mol. Immunol.* **13**, 148–159 (2016).

662 19. Lei, H. *et al.* Cationic nanocarriers as potent adjuvants for recombinant S-RBD

663   vaccine of SARS-CoV-2. *Signal Transduct. Target. Ther.* **5**, 4–6 (2020).

664 20. Zhao, L., Skwarczynski, M. & Toth, I. Polyelectrolyte-Based Platforms for the

665   Delivery of Peptides and Proteins. *ACS Biomater. Sci. Eng.* **5**, 4937–4950 (2019).

666 21. Malik, A., Gupta, M., Gupta, V., Gogoi, H. & Bhatnagar, R. Novel application of

667   trimethyl chitosan as an adjuvant in vaccine delivery. *Int. J. Nanomedicine* **13**, 7959–

668   7970 (2018).

669 22. Fadeel, B. Clear and present danger ? Engineered nanoparticles and the immune

670   system. 1–9 (2012) doi:10.4414/smw.2012.13609.

671 23. Kawai, T. & Akira, S. The role of pattern-recognition receptors in innate immunity :

672   update on Toll-like receptors. *Nat. Publ. Gr.* **11**, 373–384 (2010).

673 24. Iwasaki, A. & Medzhitov, R. Control of adaptive immunity by the innate immune

674   system. **16**, 343–353 (2015).

675 25. Martinon, F., Burns, K. & Tschopp, J. The Inflammasome: A molecular platform

676   triggering activation of inflammatory caspases and processing of proIL-β. *Mol. Cell*

10, 417–426 (2002).

26.   Serrano-Puebla, A. & Boya, P. Lysosomal membrane permeabilization in cell death:

new evidence and implications for health and disease. *Ann. N. Y. Acad. Sci.* **1371**,

30–44 (2016).

27.   Campden, R. I. & Zhang, Y. The role of lysosomal cysteine cathepsins in NLRP3

inflammasome activation. *Arch. Biochem. Biophys.* **670**, 32–42 (2019).

28.   Zhivaki, D. *et al.* Inflammasomes within Hyperactive Murine Dendritic Cells

Stimulate Long-Lived T Cell-Mediated Anti-tumor Immunity. *Cell Rep.* **33**, 108381

(2020).

29.   Evavold, C. L. *et al.* The Pore-Forming Protein Gasdermin D Regulates Interleukin-

1 Secretion from Living Macrophages. *Immunity* **48**, 35-44.e6 (2018).

30.   Monteleone, M. *et al.* Interleukin-1β Maturation Triggers Its Relocation to the

Plasma Membrane for Gasdermin-D-Dependent and -Independent Secretion. *Cell*

Rep. **24**, 1425–1433 (2018).

31.   Saeki, A. *et al.* Gasdermin D-independent release of interleukin-1β by living

macrophages in response to mycoplasmal lipoproteins and lipopeptides. *Immunology*

161, 114–122 (2020).

32.   Brubaker, S. W., Bonham, K. S., Zanoni, I. & Kagan, J. C. *Innate Immune Pattern*

Recognition : A Cell Biological Perspective. (2015). doi:10.1146/annurev-immunol-

032414-112240.

33.   Qu, Y., Franchi, L., Nunez, G. & Dubyak, G. R. Nonclassical IL-1β Secretion

Stimulated by P2X7 Receptors Is Dependent on In?ammasome Activation and

Correlated with Exosome Release in Murine Macrophages. (2015)

doi:10.4049/jimmunol.179.3.1913.

34. Abu Khweek, A. & Amer, A. O. Pyroptotic and non-pyroptotic effector functions of

caspase-11. *Immunol. Rev.* **297**, 39–52 (2020).

35. Kayagaki, N. *et al.* Non-canonical inflammasome activation targets caspase-11.

Nature **479**, 117–121 (2011).

36. Bourque, J. & Hawiger, D. *Immunomodulatory bonds of the partnership between*

dendritic cells and T cells. *Critical Reviews in Immunology* vol. 38 (2018).

37. Reddy, S. T., Swartz, M. A. & Hubbell, J. A. Targeting dendritic cells with

biomaterials: developing the next generation of vaccines. *Trends Immunol.* **27**, 573–

579 (2006).

38. Babensee, J. E. Interaction of dendritic cells with biomaterials. *Semin. Immunol.* **20**,

101–108 (2008).

39. Weischenfeldt, J. & Porse, B. Bone marrow-derived macrophages (BMM): Isolation

and applications. *Cold Spring Harb. Protoc.* **3**, 1–7 (2008).

40. Cullen, S. P., Kearney, C. J., Clancy, D. M. & Martin, S. J. Diverse Activators of the

NLRP3 Inflammasome Promote IL-1β Secretion by Triggering Necrosis. *Cell Rep.*

11, 1535–1548 (2015).

41. Chan, A. H. & Schroder, K. Inflammasome signaling and regulation of interleukin-1

family cytokines. *J. Exp. Med.* **217**, 1–10 (2020).

42. Hoss, F., Rodriguez-Alcazar, J. F. & Latz, E. Assembly and regulation of ASC

specks. *Cell. Mol. Life Sci.* **74**, 1211–1229 (2017).

43. Sester, D. P. *et al.* A Novel Flow Cytometric Method To Assess Inflammasome

Formation. *J. Immunol.* **194**, 455–462 (2015).

44. Iula, L. *et al.* Autophagy mediates interleukin-1β secretion in human neutrophils.

Front. Immunol. **9**, 1–14 (2018).

45. Dupont, N. *et al.* Autophagy-based unconventional secretory pathway for

extracellular delivery of IL-1β. *EMBO J.* **30**, 4701–4711 (2011).

46. Kwissa, M., Kasturi, S. P. & Pulendran, B. The science of adjuvants. *Expert Rev.*

Vaccines **6**, 673–684 (2007).

47. Hornung, V. *et al.* Silica crystals and aluminum salts activate the NALP3

inflammasome through phagosomal destabilization. *Nat. Immunol.* **9**, 847–856

(2008).

48. Ma, L. *et al.* Characterization of Cathepsin B in Mediating Silica Nanoparticle-

Induced Macrophage Pyroptosis via an NLRP3-Dependent Manner. 4537–4545

(2022).

49. Chevriaux, A. *et al.* Cathepsin B Is Required for NLRP3 Inflammasome Activation

in Macrophages, Through NLRP3 Interaction. *Front. Cell Dev. Biol.* **8**, 1–8 (2020).

50. Sahoo, M., Ceballos-Olvera, I., Del Barrio, L. & Re, F. Role of the inflammasome,

IL-1β, and IL-18 in bacterial infections. *ScientificWorldJournal.* **11**, 2037–2050

(2011).

51.   Baroja-Mazo, A. *et al.* Early endosome autoantigen 1 regulates IL-1β release upon

741   caspase-1 activation independently of gasdermin D membrane permeabilization. *Sci.*

742   *Rep.* **9**, 1–10 (2019).

743   52.   Ponpuak, M. *et al.* Secretory autophagy. *Curr. Opin. Cell Biol.* **35**, 106–116 (2015).

744   53.   Deretic, V. Autophagy in inflammation, infection, and immunometabolism.

745   *Immunity* **54**, 437–453 (2021).

746   54.   Kimura, T. *et al.* Dedicated SNARE s and specialized TRIM cargo receptors

747   mediate secretory autophagy . *EMBO J.* **36**, 42–60 (2017).

748   55.   Sweatt,S.K, Gower, B.A, Chieh, A.Y, Liu, Y, Li, L. Crucial role for the Nalp3

749   inflammasome in the immunostimulatory properties of aluminium adjuvants.

750   *Physiol. Behav.* **176**, 139–148 (2016).

751   56.   Daniel E Shumer, Natalie J Nokoff, N. P. S. & Erin C. Dowd, M.D.a, Michael J.

752   Frank, Ph.D.b, Anne Collins, Ph.D.c, James M. Goldd, and Deanna M. Barch, P. D. .

753   T helper 1 immunity requires complement-driven NLRP3 inflammasome activity in

754   CD4+ T cells. *Physiol. Behav.* **176**, 139–148 (2017).

755   57.   Wagatsuma, K. & Nakase, H. Contradictory effects of NLRP3 inflammasome

756   regulatory mechanisms in colitis. *Int. J. Mol. Sci.* **21**, 1–24 (2020).

757   58.   Marty-Roix, R. *et al.* Identification of QS-21 as an inflammasome-activating

758   molecular component of saponin adjuvants. *J. Biol. Chem.* **291**, 1123–1136 (2016).

759   59.   Kool, M. *et al.* Alum adjuvant boosts adaptive immunity by inducing uric acid and

760   activating inflammatory dendritic cells. *J. Exp. Med.* **205**, 869–882 (2008).

60. Content, R. Alum ' s Adjuvant Effect Are Mediated by NLRP3 1. **181**, 17–21

(2014).

61. Muñoz-Wolf, N. *et al.* Non-canonical inflammasome activation mediates the

adjuvanticity of nanoparticles. *Cell Reports Med.* **4**, (2023).

62. Kaplanski, G. Interleukin-18: Biological properties and role in disease pathogenesis.

Immunol. Rev. **281**, 138–153 (2018).

63. Lacaille-Dubois, M. A. Updated insights into the mechanism of action and clinical

profile of the immunoadjuvant QS-21: A review. *Phytomedicine* **60**, 152905 (2019).

FIGURES

Figure 1: Cell uptake of Np and activation. **(A)** Schematic Np composition; **(B)** TEM

image of the Np prepared at a Tpp/PAH=0.6 ratio; **(C)** Particle size measured by DLS. **(D)**

Np-FITC internalized by J774 and colocalization with the late endosome marker LAMP-1

by confocal microscopy after 4h stimulation. **(E)** Inhibition of phagocytosis of Np-FITC

with cytochalasin D in BMDM by confocal microscopy and flow cytometry. **(F)** BMDC

surface expression of CD86 and MHCII in BMDC (CD11b$^+$CD11c$^+$) by flow cytometry.

All experiments were performed in triplicate and data are expressed as mean±SEM. P-value

was determined by ordinary one-way ANOVA *P<0,05; **P<0,01.

Figure 2: Secretion of pro-inflammatory cytokine by macrophages, dendritic cells and

monocytes. J774, THP-1 and BMDC were prime with LPS (1μg/mL) for 3h; the medium

was fully replaced and stimulated with ATP (50nM) for 3h as a positive control or with Np

(0.5mM) overnight. Controls of LPS and Np were stimulated overnight. **(A)** IL-1β and **(B)**

IL-6 and IL-8 and **(C)** IL-18 secreted cytokine by phagocyte cells were measured in

supernatants by ELISA. All experiments were carried out in triplicate and date are

expressed as mean±SEM. P-value was determined by ordinary One-way ANOVA *P<0,05;

**P<0,01; ***P<0,001; ****p<0,0001.

Figure 3: Inflammasome activation and cytokine secretion in THP1-ACS-GFP cells.

Cells were primed with LPS for 3 hours. The medium was fully replaced, and the cells were

either stimulated with Np overnight or transfected with pcDNA as positive controls. **(A)**

ASC-GFP speck within the cells are shown with white arrow, performed by

epifluorescence microscopy; **(B)** Frequency of speck ASC-GFP[+]cells. Three independent

experiments were performed. Each bar represents the average number of total speck[+]

cells/total cells per field; 3 fields for each treatment were counted (2 replicates/treatment);

(C) Flow cytometry gate strategy to define ASC-GFP[+]cells; **(D)** Frequency of speck ASC-

GFP[+]cells by flow cytometry; **(E)** Quantification of secreted hIL-1β by ELISA. **(F)** BMDC

were stimulated overnight with LPS+Np and simultaneously inhibited with cytochalasin D

for phagocytosis, CA-74Me for cathepsin B and Z-VAD-FMK (Z-VAD) for caspase-1

during 30min, and quantification of secreted mIL-1β and mIL-18 by ELISA. All

experiments were carried out in triplicate and date are expressed as mean±SEM. P-value

was determined by ordinary one-way ANOVA *p<0,05; **p<0,01; ***p<0,001 or t-

Student #p<0,05; ##p<0,01.

Figure 4: Inflammasome pathway analysis in BMDM from C57BL/6 (WT), NLRP3$^{-/-}$,

Casp1/11$^{-/-}$, Casp11$^{-/-}$, GSDMD$^{-/-}$ mice. Cells were stimulated with LPS, Np, or

LPS+Np, and **(A)** mIL-1β, **(B)** mTNFα were measured in supernatants by ELISA. The

effect of autophagy stimulation by starvation on macrophage IL-1β secretion was

investigated in J774 cell lines. After being primed for 3 hours with LPS, the cells were

treated with Np. Six hours later, the cells were either left untreated in medium or had their

supernatants replaced with Earle's Balanced Salt Solution (EBSS) for an additional hour of

culture and quantification **(C)** mIL-1β in the supernatants by ELISA. J774 cells line were

stimulated for 6h with Np in the presence of inhibitors of the autophagy VPS34-IN1 and 3-

methiladenine (3-MA) and quantification **(D)** mIL-1β. **(E)** Cytokine mIL-1β colocalized

with LC3B by confocal microscopy after 6 hours of stimulation with LPS+Np (MG:

Methyl green). Data are expressed as mean±SEM. P-value was determined by ordinary

one-way ANOVA *$P < 0,05$; **$P < 0,01$; ***$P < 0,001$; ****$p < 0,0001$.

Figure 5: Intraperitoneally immunization of Balb/c mice with Np-OVA and specific

humoral and cellular immune responses. (A) Immunization scheme; **(B)** Body weight

monitoring; **(C)** quantification of serum specific IgG to PAH, Np and OVA and **(D)** OVA-

Specific IgG and **(E)** IgG2a titers in serum of immunization animals and **(F)** serum optical

density ratio of isotypes between Np-OVA and OVA+Alum; **(G)** Quantification of mIFN-

γ, mIL-5 and mIL-13 in the supernatants of spleen cells from immunized mice assessed by

ELISA. **(H)** Spleen cells from immunized mice were incubated with brefeldin A for the last

4h of culture and then were stained with anti-CD4 (PerCP-Cy5.5) or anti-CD8 (APC); flow

cytometry gate strategy and Mean Fluorescent Intensity (MFI) **(I)** quantification for

$CD4^+IFN-\gamma^+$ and $CD8^+IFN-\gamma^+$ T cells. The immunization experiment was carried out in

duplicate and all experiments were performed in triplicate; data are expressed as

mean±SEM. P-value was determined by ordinary one-way ANOVA *p<0,05; **p<0,01;

***p<0,001 or t-Student test #p<0,05.

Figure 6: Intraperitoneal immunization with Np-OVA in WT and KO C57BL/6 mice

and specific humoral and cellular response. C57BL/6 (WT), $NLRP3^{-/-}$, $Casp1/11^{-/-}$, IL-

$1R^{-/-}$ mice were immunized following the same scheme as in Figure 5A. **(A)** Measurement

of serum OVA-specific IgG in serum (dilution 1/100) by ELISA and **(B)** secreted mIFN-γ

by spleen cells from OVA or Np-OVA immunized mice by ELISA. The immunization

experiment was carried out in duplicate and all experiments were performed in triplicate;

data are expressed as mean±SEM. P-value was determined by ordinary one-way ANOVA

*p<0,05; **p<0,01; ***p<0,001 or t-Student test ##p<0,01.

SUPPLEMENTARY FIGURES

Figure S1: Internalization of (A) PAH-FITC (B) and Np-FITC in J774 macrophages by confocal microscopy. Representative images are shown. **(C)** Gate strategy to select F4/80 CD11b-positive macrophages cells.

Figure S2: **Interaction of Np-FITC with human epithelial cell lines and dose-**

dependent cytokine response of cells to different stimuli. (A) HT-29, different z-axis

positions were scanned by confocal microscopy and EPCAM were used as a marker of

epithelial cells. Representative images are shown. **(B)** Quantification of secreted hIL-8 in

HT-29 cells exposed to medium, TNF-α, TNF-α+Np and Np; **(C)** Quantification of

secreted mIL-1β in J774 macrophages exposed to medium, LPS, LPS+ATP, LPS+Np and

Np. All experiments were carried out in triplicate and date are expressed as mean±SEM. P-

value was determined by ordinary one-way ANOVA *p<0,05; **p<0,01; ***p<0,001;

****p<0,0001.

F

G

Figure S3: **Inflammasome pathway analysis in BMDM from C57BL/6 (WT),**

NLRP3$^{-/-}$, Casp1/11$^{-/-}$, Casp11$^{-/-}$, GSDMD$^{-/-}$ mice. Cells were stimulated with medium,

Np or LPS+Np, and **(A)** %LDH were quantified in the supernatants and **(B)** caspase-1

activation was analyzed by immunoblotting in cell lysates using an antibody specific for the

p20 subunit (20kDa) or the procaspase-1 (45kDa). J774 cell lines remained unstimulated or

were stimulated with LPS. Three hours later, they were treated with Np. At 6 hours post-Np

stimulation, cells were either left untreated (in medium) or their supernatants were removed

and replaced with EBSS for an additional hour of culture. Subsequently J774 macrophages

were stimulated for 6h with Np in the presence of inhibitors of the autophagy VPS34-IN1

and 3-methiladenine (3-MA) and quantification **(C-D)** mIL-6 and **(E)** %LDH in the

supernatants. **(F)** Quantification of mIL-1β and mIL-6 in the supernatants of BMDCs after

the treatment with autophagy inhibitors. **(G)** Medium, Np and LPS controls for the mIL-1β-

LC3B colocalization by confocal microscopy. P-value was determined by ordinary one-

way ANOVA *p<0,05; **p<0,01; ***p<0,001; ****p<0,0001.

Figure S4: **Analysis of the stimulation of the Hek-hTLR4 cell line with LPS**. Different

concentrations of LPS were included as positive controls; SEAP activity was measured in

the supernatants. This experiment was performed in triplicate and data are expressed as

mean±SEM. P-value was determined by ordinary one-way ANOVA *p<0,05; **p<0,01;

***p<0,001; Φ a result significantly different from LPS.

Figure S5: Analysis of the OVA encapsulation by Np. Free OVA was analyzed in the

supernatant of the Np-OVA formulation by protein quantification (BCA). This experiment

was performed in triplicate and data are expressed as mean±SEM. P-value was determined

by Student's test ### p<0,001.