Cryptic recombination and transposition drive structural variation to shape genomic

plasticity and life history traits in a host generalist fungal plant pathogen

Mark C Derbyshire[1,*], Toby E Newman[1], Yuphin Khentry[1], Pippa J Michael[1], Sarita Jane

Bennett[1], Ashmita Rijal Lamichhane[1], Carolyn Graham-Taylor[1] Subhash Chander[2],

Claudia Camplone[3], Simone Vicini[3], Laura Esquivel-Garcia[4], Lone Buchwaldt[5], Cathy

Coutu[5], Dwayne Hegedus[5], John Clarkson[6], Kurt Lindbeck[7], Lars G Kamphuis[1]

[1]Centre for Crop and Disease Management, Curtin University, Perth, Western Australia,

Australia

[2]Oilseeds Section, Department of Genetics and Plant Breeding, CCS Haryana Agricultural

University, Hisar-125004, India

[3]Department of Agricultural, Food and Environmental Sciences, University of Perugia,

Perugia, Italy.

[4]Plant Science, McGill University, Sainte-Anne-de-Bellevue, Quebec, Canada.

[5]Agriculture and Agri-Food Canada, Saskatoon, Saskatchewan, Canada.

[6]Warwick Crop Centre, School of Life Sciences, University of Warwick, Warwick, UK.

[7]New South Wales Department of Primary Industries, Wagga Wagga, New South Wales,

Australia

[*]Author for correspondence: mark.derbyshire@curtin.edu.au

Abstract

Background: An understanding of plant pathogen evolution is important for sustainable management of crop diseases. Plant pathogen populations must maintain adequate heritable phenotypic variability to survive. Polymorphisms >= 50 bp, known as structural variants (SVs), could contribute strongly to this variability by disrupting gene activities. SV acquisition is largely driven by mobile genetic elements called transposons, though a less appreciated source of SVs is erroneous meiotic double-strand break repair. The relative impacts of transposons and recombination on SV diversity and the overall contribution of SVs to phenotypic variability is elusive, especially in host generalists.

Results: We use 25 high quality genomes to create a graphical pan-genome of the globally distributed host-generalist crop pathogen *Sclerotinia sclerotiorum*. Outcrossing and recombination rates in this self-fertile species have been debated. Using bisulfite sequencing, and short read data from 190 strains, we show that *S. sclerotiorum* has many hallmarks of eukaryotic meiosis, including recombination hot and cold spots, centromeric and genic recombination suppression, and rapid linkage disequilibrium decay. Using a new statistic that captures average pairwise structural variation, we show that recombination and transposons make distinct contributions to SV diversity. Furthermore, despite only 5 % of genes being dispensable, SVs often had a stronger impact than other variants across 14 life history traits measured in 103 distinct strains.

Conclusion: Transposons and recombination make distinct contributions to SV diversity in *S. sclerotiorum*. Despite limited gene content diversity, SVs may strongly impact phenotypic variability. This sheds light on the genomic forces shaping adaptive flexibility in host generalists.

Keywords

Recombination, structural variant, genome graph, plant pathogen, fungus

Background

An understanding of the evolutionary processes underpinning plant pathogen adaptation is crucial for developing better disease management strategies, such as resistant cultivars, prediction of epidemics and monitoring of fungicide resistance [1–3]. Population genetic approaches can be used to understand the evolutionary characteristics of plant pathogens [4,5], although their application has been limited in the past to variants that can be confidently genotyped using short reads. However, with the now widespread use of long read sequencing, more plant pathogen pan-genomes of increasing quality are becoming available for evolutionary studies [6–12].

Aside from simple genotypic variants, such as single nucleotide polymorphisms (SNPs) and small insertions/deletions (InDels), complete genomes assembled using long reads can be used to identify structural variants (SVs), which are generally defined as polymorphisms of more than or equal to 50 bp [13]. These can be confidently genotyped using long reads assemblies and incorporated into a data structure known as a pan-genome graph [14,15]. Based on this underlying representation of genomic variation, SVs can be genotyped in a broader set of individuals using short reads [16,17]. This approach also improves the accuracy of non-SV calls by improving short read placement and reducing reference bias [18].

In many species, this and similar techniques have revealed that previously invisible SVs are strongly linked with phenotypic variability. For instance, the tomato graph pan-genome showed that SVs are a major component of 'missing heritability' [19], explaining much of the phenotypic variance not captured by simpler variants called against a single reference. Furthermore, in the important fungal wheat pathogen *Zymoseptoria tritici*, SVs have been shown to make a substantial contribution to important life history traits, such as tolerance of fungicides [20]

Two key processes underpinning evolutionary adaptation are mutation, including *de novo*

acquisition of SVs, and meiotic recombination. The traditional view is that mutation creates

new alleles and meiotic recombination shuffles alleles to create new haplotypes [21].

Shuffling of alleles into novel haplotypes allows beneficial alleles to spread without the

burden of linked deleterious alleles. Without meiotic exchange, populations are likely to

gradually accumulate deleterious mutations that cannot be lost without also losing

beneficial mutations, a process known as Muller's ratchet [22–24].

Though evolutionary theory often ascribes distinct roles to mutation and meiotic

recombination in creating and shuffling alleles, respectively, the two processes may not be

completely orthogonal. Meiosis itself may be powerfully mutagenic, as it requires the

induction of numerous double-strand DNA breaks. Through erroneous repair of these

breaks, meiosis has been linked with exceptionally high *de novo* mutation rates [12,20,21].

All types of mutations can occur through faulty repair of double-strand breaks, although

meiosis-induced double strand breaks may be particularly prone to creating new SVs [21].

In humans, for example, double-strand breaks induced by meiosis lead to a 400 to 1,000-

fold increase in the rate of SV acquisition, and many SVs induced in recombination

hotspots are pathogenic, highlighting the impact of meiotic mutagenesis on phenotype and

human disease [12, 25]. Recently, a machine learning approach showed that multiple

genomic features, including local recombination rate, were highly predictive of SVs

induced in haploid offspring of crosses of *Z. tritici* [20]. In the plant pathogen *Fusarium*

graminearum, local recombination rate was also shown to be associated with SVs across

four high quality genomes [12].

In addition to meiosis, transposition is a highly potent instigator of structural variation in

genomes. This occurs when active mobile elements called transposons duplicate or

relocate themselves in the genome [26]. In addition, the repetitive nature of transposons

can create SVs through pairing of distant genomic copies during DNA damage repair via

the homologous recombination pathway [26]. Though transposons can be destabilising to

genomes, occasionally they create beneficial mutations, which are an important source of

adaptive evolution [27].

In plant pathogens, transposition is widely appreciated as one of the main driving forces of

genomic plasticity. Though meiotic exchange has been linked with *de novo* acquisition of

SVs in plant pathogenic fungi, the link between meiotic exchange and genome stability has

not been widely explored in plant pathogen populations, and little is known about how

meiosis and transposition interact to shape SV diversity. Despite several long reads

pathogen pan-genomes, the overall contribution of SVs to variability in life history traits is

also poorly understood.

To date, much of the research on the evolution of plant pathogen genomes has also been

conducted on host specialists, which are under acute selective pressure to maintain

virulence on a single species. In contrast, the fungus *Sclerotinia sclerotiorum* infects

hundreds of plant species in at least 74 documented families [28]. Though its genome may

harbour some polymorphic regions [29,30], in contrast to many host specialists, its

predicted effectors are largely conserved [30] and several are likely compatible with

diverse hosts [31,32]. This suggests that, like many niche-generalists, *S. sclerotiorum* has

evolved an energetically-optimised and multifunctional genome, which facilitates its

colonisation of diverse hosts [33–35].

Sporulation in *S. sclerotiorum* occurs through obligate sexual reproduction. However, since

it is self-fertile (homothallic), sexual reproduction can create genotypically uniform progeny,

allowing certain genotypes to persist for long periods of time as clones [36]. The extent to

which *S. sclerotiorum* outcrosses to generate new diversity has been debated, with some

suggesting homothallism promotes universal outcrossing [29,37–44] and others

suggesting that outcrossing is extremely rare [45–47]. Consequently, the overall

contribution of meiotic exchange to genome stability and evolution in this species is

particularly poorly understood.

Here, we present a global graphical pan-genome of *S. sclerotiorum* and use 25 reference-

quality genomes and 190 short reads samples to investigate species-wide SV diversity. To

capture this diversity, we present a new statistic called 'SVπ', which describes the average

number of SVs between all pairs of individuals. Using population genetics techniques, we

establish *S. sclerotiorum* as an outcrossing species with many of the hallmarks of

eukaryotic meiotic recombination, such as rapid linkage disequilibrium decay, suppressed

recombination at centromeres, recombination hot and cold spots and enhanced

recombination outside of coding sequences. We find that both recombination rate and

transposable element content are independently positively correlated with total number of

SVs and SVπ though not positively correlated with one another.

Overall, unlike that of most host specialists studied to date, we show that gene content in

the *S. sclerotiorum* genome is largely stable, despite numerous small, unstable, repeat-

rich, gene-sparse regions. SVs often had a stronger effect than other variants on 14 life

history traits assessed across 103 strains, and we find that a 48 bp InDel is significantly

associated with tolerance of the fungicide azoxystrobin. Overall, our data suggest that

transposition and meiotic recombination make distinct contributions to SV diversity in *S.*

sclerotiorum, and that SVs may be an important driver of phenotypic plasticity, despite the

stability in gene content of the species. These insights shed new light on the genomic

processes underpinning the evolution of host generalism in plant pathogens.

Results and discussion

Development of a *Sclerotinia sclerotiorum* graphical pan-genome

To create a high-quality set of *S. sclerotiorum* genomes for SV analysis, we generated

Illumina-corrected Oxford Nanopore long reads assemblies of the genomes of 24 diverse

strains from Australia (10 strains), Europe (5 strains) and Canada (9 strains). Overall, 23 of

the strains had telomere-to-telomere assemblies for >= 10 of the 16 *S. sclerotiorum*

chromosomes, and seven had telomere-to-telomere assemblies for >= 14 chromosomes.

There were few gaps in assemblies on average, and 22 strains had gapless assemblies

for >= 10 chromosomes; three of these strains had gapless assemblies for all 16

chromosomes. All these assemblies are comparable to the reference *S. sclerotiorum*

genome [30], which has 14 telomere-to-telomere and 15 gapless chromosomes. BUSCO

scores ranged from 98.9 to 99.2, with a median of 99.05 (Supplementary Table 1),

confirming the completeness of these assemblies. These new assemblies are available in

NCBI under BioProject PRJNA1112094.

To explore structural variation in *S. sclerotiorum* we constructed a pan-genome graph from

the genomes of these 24 strains and the reference strain. In this graph, we identified

186,486 variants, including 154,892 SNPs, 5,877 multiple nucleotide polymorphisms

(MNPs), 20,061 InDels and 5,556 SVs. There were 9,876 complex variants with more than

one allele, including 2,892 (52 %) of the SVs.

To capture more genotypic diversity, we aligned Illumina short reads to the pan-genome

graph from an additional 190 strains, 181 of which were sequenced in this study (available

in NCBI under BioProject PRJNA1120954) (Supplementary Table 2). Overall, the

genotypes of 3,741 of the SVs from the pan-genome graph were captured in this broader

data set. This data set is the first graphical pan-genome of the important host generalist

pathogen *S. sclerotiorum*. It includes 215 strains, with 152 from Australia, 17 from North

America, 44 from Europe, and one each from South Africa and Morocco.

Figure 1. Genotypic clustering of *Sclerotinia sclerotiorum* strains from the global population sample. A A phylogenetic network with all strains in the dataset coloured according to geographical origin. The map inset shows where strains were collected with colours corresponding to those in the network. The sizes of circles on the map corresponds with the number of strains from each global region. **B** A phylogenetic network for the Australian strains. Circles are coloured according to geographical origin within Australia. Where circles are stacked on top of each other, isolates are a >= 98 %

genotypically identical group of clones. The map to the left shows where isolates were collected within Australia, with colours of circles corresponding to colours on the network. The sizes of circles represent the numbers of strains from each collection site. Haplotypes 1 (hap 1) and 2 (hap 2) are examples of frequently sampled and geographically widespread clones, with individuals from Western Australia and South Australia.

Sclerotinia sclerotiorum undergoes cryptic recombination whilst maintaining clonal lineages across large temporal and spatial distances

S. sclerotiorum produces ascospores through sexual reproduction. As it is homothallic, ascospores may be genotypically identical, which leads to an effectively clonal mode of propagation. Clonality is evident in the detection of temporally or spatially distant genotypically nearly identical strains. We identified 120 clonal lineages (>= 98 % identical) among the 215 strains (Supplementary Figure 1). Clonality was most prevalent among the Australian strains, whereas European and North American strains were mostly genotypically distinct (Figure 1). This was expected because most of the European and North American strains were previously shown to be distinct lineages using markers [48,49], whereas 99 of the Australian strains were collected from five sites (two of which were in the same locality) in Western Australia with no prior genotyping [50,51]. Confirming the long-term maintenance of clonal propagation, we found several clones from geographically distant regions, some of which were collected many years apart. For example, strains CU11.18 and F19064 were found in Western Australia in 2013 and South Australia in 2018, respectively. The most extreme example was the pair of clones S55 and MB57, which were collected in the USA in 1987 and Manitoba in 2010, respectively.

In our previous study, we found that the global *S. sclerotiorum* population forms two distinct sub-populations, between which there has been limited gene flow. SNP data from the 215 genomes confirmed this observation (Figure 1), showing that Australian/African

and European/North American strains formed mostly distinct sub-populations (referred to as AuAf and EuNA herein). Although we expanded the Australian collection, our study only contained the two African strains from our previous study [29], Sssaf from South Africa and Ss44 from Morocco, so the global relationship between Australian and African strains is still not fully resolved.

In the AuAf sub-population, we found evidence for three ancestral populations, and numerous admixed individuals. In the EuNA population, we identified three further ancestral populations with limited admixture (Figure 2 A). Two of the EuNA strains were admixed individuals containing alleles from either the AuAf ancestral populations or both the EuNA and AuAf ancestral populations. The widespread recent admixture among AuAf strains supports outcrossing between lineages from distinct ancestral populations.

To further explore outcrossing in the global *S. sclerotiorum* population, we investigated the rate of linkage disequilibrium decay in the 120 independent clonal lineages. We found that across the whole population, linkage disequilibrium decayed to half its maximum value at 428 bp (Figure 2 B). Three tests of the association between recombination and physical distance, neighbour similarity score [52], maximum X^2 [53], and pairwise homoplasy index [54], also supported statistically significant recombination between non-adjacent alleles (P = 0 for all tests and chromosomes, Supplementary Table 3).

The rate at which LD decays to half its maximum (LD_2) value is typically higher in predominantly outcrossing and lower in predominantly clonal species [55,56]. Species that rarely outcross often have LD_2 rates of more than 100 Kb, whereas highly outcrossing species have rates on the order of a few hundred bp [56]. Though we have no direct assessment of the rate of outcrossing in *S. sclerotiorum*, the very small LD_2 rate we observed suggests that it may be relatively frequent.

Like other outcrossing species, recombination was not uniform across the genome. In four non-structured subsamples comprising respectively 23, 13, 15 and 34 individuals (Figure 2

C), which we refer to as population-1, population-2, population-3 and population-4, we

identified 384 recombination hotspots (Supplementary Table 4). Like other outcrossing

species, we found that recombination rate was higher towards the ends of chromosomes

where chromatin is more likely to be relaxed (Figure 2 D). Furthermore, recombination rate

was negatively correlated with coding sequence density (Spearman's ρ = -0.15, P = 0,

Figure 2 E, Supplementary Figure 2) and recombination hotspots had a lower gene density

than other regions (P < 0.0001, Figure 2 F). Though recombination rate was weakly

negatively correlated with coding sequence density, the relationship between the two

variables was not monotonic. Instead, there was an optimal gene density at which

recombination rate peaked before declining rapidly (Figure 2 E, Supplementary Figure 2).

Our data suggest that recombination rate is generally higher outside of genic regions but

low in the most gene-sparse parts of the genome. This is consistent with observations in

other outcrossing species [57] where meiotic recombination within genes is selected

against as it can lead to polymorphisms due to erroneous double strand break (DSB)

repair, though meiosis is repressed in the most gene-sparse regions, which also tend to be

heterochromatic.

Across most chromosomes and all four population samples, there were clear

recombination coldspots that coincided with a single prominent drop in GC content and a

single prominent spike in cytosine methylation based on bisulfite sequencing data

generated in this study (Figure 2 G-H, Supplementary File 1, Supplementary Table 5).

Decreased GC content and increased cytosine methylation are both hallmarks of

eukaryotic centromeres [58,59], around which meiotic recombination is typically

suppressed [60]. The convergence of these three observations, and previous predictions

from optical mapping data [61], suggest that these sites are the centromeres of the *S.*

sclerotiorum chromosomes, and, as in other outcrossing species, meiotic recombination is

suppressed around them.

With the rapid decay of linkage disequilibrium, the presence of recombination hotspots,

and the conspicuous recombination-related features characteristic of eukaryotic meiosis,

we infer that *S. sclerotiorum* maintains genetic diversity across numerous populations

through sexual outcrossing. While clonal lineages may endure over extended periods via

self-fertilization, the ongoing process of sexual recombination among these lineages may

be important for creating genotypic diversity. Presently, meiotic exchange is cryptic, as

laboratory observations of sexual outcrossing are, to our knowledge, lacking.

Ecological theory suggests that loss of sexual reproduction initiates the gradual

accumulation of deleterious alleles inseparable from beneficial ones, a phenomenon

known as 'Muller's ratchet'. Consequently, strictly clonal populations are rare, with most

facing a trajectory toward extinction. Given the continuing pressure on *S. sclerotiorum* for

survival across numerous host species, coupled with its apparent lack of host preference,

it is not surprising that it exhibits many attributes indicative of sexual outcrossing. Drawing

from our findings and those of others [37,43,44], we suggest that homothallism in *S.*

sclerotiorum not only supports persistence of certain clonal lineages but also fosters

universal sexual compatibility.

Figure 2. Population structure and evidence of recombination. A Colours correspond

to ancestral populations making up individuals. Country of origin (above) is Au = Australia,

Mo = Morocco, SoA = South Africa, Ca = Canada, Fr = France, No = Norway, and UK =

UK. Below, states within Australia and Canada are indicated, where NSW = New South

Wales, SA = South Australia, WA = Western Australia, AB = Alberta, MB = Manitoba, and SK = Saskatchewan. **B** Linkage disequilibrium (y axis) decay with physical distance (x axis). Points are averages for unique distance measurements, and the red line is a general additive model fit. **C** The first two principal components of genotypic variance. Colours indicate geographical origin and point shapes the four population sub-samples used for recombination analysis. **D** Across chromosomes and population sub-samples, the distribution of Spearman's correlations between chromosome end distance and recombination rate. **E** Correlation between coding DNA sequence content (x axis) and recombination rate (y axis) of 50 Kb sliding windows. The line is a general additive model fit. **F** Boxplot showing percent gene content of 50 Kb windows containing and not containing recombination hotspots (*** = $P < 2e^{-16}$). Boxes and whiskers show interquartile range. **G** Circles show where windows containing putative centromeres lie on a plot of recombination rate (y axis) against log recombination rate (x axis). Putative centromeres are in regions of low recombination, before the inflection point. **H** The y axis is scaled (division by maximum) recombination rate, amount of methylation or GC content for sliding windows. The x axis shows position (Mb) across chromosome 6 (all chromosomes and population samples are in Supplementary File 1). All chromosomes had a dip in GC coincident with a spike in methylation, almost always coincident with a recombination cold spot.

The *Sclerotinia sclerotiorum* pan-genome graph suggests transposable elements create hotspots of structural diversity

To capture diversity of structural variation across the genome, we developed a statistic called 'SVπ'. Akin to nucleotide [62] and synteny diversity [63], this statistic captures the average number of SVs per Kb between all pairs of individuals. We found that SVπ was positively skewed when calculated for 50 Kb sliding windows across the genome (Figure 3

A). This suggests that the *S. sclerotiorum* genome is mostly stable, with a few regions of excessive structural variation. We defined SV hotspots as sliding windows with a SVπ value above the 95th percentile across the genome. Interestingly, more hotspots were detected on some chromosomes than others. For example, chromosome 12 contained six hotspots and had an average SVπ of 0.016, whereas, despite being larger, chromosome 6 contained only one hotspot and had an average SVπ of 0.009 (Figure 3 D, Supplementary Figure 3).

Transposable element and gene content were positively (Spearman's ρ = 0.28, P = 0) and negatively (Spearman's ρ = -0.33, P = 0) correlated with SVπ, respectively. The correlation between transposon/gene content and SVπ varied between chromosomes, with the highest correlations observed on chromosomes 10 and 12 (Spearman's ρ = 0.61 and 0.62, Spearman's ρ = -0.58 and -0.59, respectively (P = 0)) (Figure 3 B; Supplementary Table 8). The association between SVs and transposable elements was further supported by the observation that transposable elements were significantly closer than randomised loci to the nearest SV across all genomes (P = 0). Transposable elements in the long terminal repeat (LTR) family were significantly (P < 0.0001) closer to the nearest SV than those in eight out of 10 other families (Figure 3 C, Supplementary Table 6), suggesting they may strongly contribute to genome instability in *S. sclerotiorum*.

LTRs are a type of retrotransposon, which are transposons characterised by a copy and paste proliferation mechanism that involves transcription into RNA, reverse transcription into DNA and re-insertion into the genome [64]. Retrotransposons are unique to eukaryotes [65], and their replicative ability has made them often the dominant transposon class in eukaryote genomes [66]. Several studies have linked retrotransposons with virulence evolution in plant pathogens [67,68], including the host generalist species *Botrytis cinerea*, where they have been shown to encode small RNA effectors [69]. Our observations that retrotransposons are most strongly linked of all transposon classes to

SVs suggests that they are the most active mobile elements in *S. sclerotiorum*. Their ongoing contribution to structural variation may be important for genomic plasticity in this species.

Stable, gene-dense and repeat-poor, and unstable, gene-sparse and repeat-rich genomic regions are common across eukaryote genomes [70]. The accumulation of transposons and SVs in gene-sparse regions is likely a result of relaxed selective pressure and accumulation of largely selectively neutral alleles. These regions can be important for adaptive evolution because they harbour extensive diversity in gene content and gene sequences [71]. When the environment changes, previously selectively neutral mutations may confer an advantage, leading to ongoing maintenance of these regions, and the transposons within them, in populations [72]. Our data show that, like those of most eukaryotes, the *S. sclerotiorum* genome is also partitioned into stable and unstable regions, and unstable regions are likely most strongly shaped by LTR retrotransposons. Overall, transposon content in the 25 *S. sclerotiorum* genomes was relatively low at 5.51 to 6.91 % (Supplementary Table 7). Despite this, transposable elements are responsible for creating considerable diversity in SVs across the *S. sclerotiorum* genome.

Figure 3. Analysis of structural variation across the *Sclerotinia sclerotiorum* pan-

genome. A Distribution (y axis) of SVπ (x axis) for 50 Kb sliding windows. **B** For

chromosomes 10 and 12, correlation between SVπ (x axis) and proportion transposon (top

y axis) or coding DNA sequence (bottom y axis). Spearman's ρ and P value depicted top-

right. Blue lines show linear regression of y onto x and the shaded area 95 % confidence

interval. Red points are SVπ hotspot (> 95[th] percentile) windows. **C** The y axis shows

distance to nearest structural variant (SV) for transposon families. Transposon

classification is indicated at the top and family on the x axis. Boxes and whiskers show

interquartile range. LTR retrotransposons were generally closer than other transposons to

SVs (Kruskall-Wallis test shown in Supplementary Table 6). **D** The y axis is SVπ or percent

repeat for 50 Kb windows (scaled for visualisation). The x axis shows window start (Mb),

and plots show chromosomes 6 and 12, the latter having the highest average SVπ and the

most hotspots (shaded in pink). **E** Correlation between log recombination rate per Kb (y

axis) and SVπ (x axis) across 50 Kb sliding windows. Chromosomes are plotted in

different colours and data shown are for population-3. Spearman's ρ was 0.14-0.15 for all

populations (P = 0) but varied between chromosomes. **F** Distribution across chromosomes

(y axis) of Spearman's ρ for number of SVs and recombination rate in 50 Kb sliding

windows. Though correlation strength varied between chromosomes, correlations were

generally positive. **G** The y axis shows repeat content (top), SVπ (middle) and number of

SVs (bottom) for windows that did not (left) and did (right) contain recombination hotspots.

Boxes and whiskers show interquartile range; differences were significant according to a t-

test (*** = P < 2.2e^{-16}).

Recombination and transposable elements make distinct contributions to structural

variation

Several studies have shown that besides transposition, structural variation can be caused

by recombination. However, little is known about the overall impact of recombination on

structural variation in natural populations. In *S. sclerotiorum*, we found an overall

correlation between SVπ and recombination rate for all four population samples we used

for recombination rate estimation (Figure 3 E, Spearman's ρ = 0.14-0.15, P = 0). Though

this is suggestive of a link between SV diversity and recombination, it does not necessarily

imply that recombination creates SVs, as this relationship could also be caused by

increased haplotype diversity in regions with a high recombination rate. Therefore, to

determine whether genomic regions with a high recombination rate may be more prone to

development of SVs, we assessed the correlation between recombination rate and the

overall number of SVs called against the reference genome. Though the strength of

correlation between these parameters varied considerably between chromosomes and

populations, we found that, on average, there was a weak to moderate correlation

between total number of SVs and estimated recombination rate (Figure 3 F, mean

Spearman's ρ = 0.09). For 12, 8, 13, and 12 out of 16 chromosomes, for the four

respective populations, there was a significant positive correlation between recombination

rate and total number of SVs ($P < 0.05$, Supplementary Table 8). In contrast, only 1-3

chromosomes displayed a significant negative correlation between recombination rate and

number of SVs.

Despite the correlations between recombination rate and both SVπ and total SVs across

chromosomes and populations, there were far fewer instances of a positive correlation

between recombination rate and transposon content, and the overall average of all

Spearman's ρs was close to zero at -0.0018 (Supplementary Table 8). Furthermore,

recombination hotspots had a slightly but significantly lower average repeat content than

other parts of the genome (5.71 % vs 6.97 %, $P < 2.2e{-16}$), despite having elevated SVπ

(average of 0.012 vs 0.010, $P < 2.2e{-16}$) and more SVs (2.65 vs 2.40, $P < 2.2e{-16}$) (Figure

3 G). This suggests that meiotic recombination and transposition make orthogonal

contributions to structural variation. In agreement, we found that the number of SVs was

better described in a regression model by both average recombination rate and

transposon content than transposon content alone, though transposon content was the

dominant predictor in the model (likelihood ratio test P < 2.2e-16, transposon F = 74.86,

recombination rate F = 13.08).

Our analyses document an interesting link between estimated recombination rate and the

rate of structural variation in the *S. sclerotiorum* genome. This is not surprising given the

mutagenic properties of meiosis. Given the relatively low level of transposable element

content in the *S. sclerotiorum* genome, recombination through meiotic exchange could be

an additional important source of structural variation. Our regression model suggests that

recombination rate is far outweighed by transposon density as a predictor of genome

stability. However, since recombination rate was typically higher in regions of intermediate

gene density, recombination may have a greater chance of inducing SVs that impact gene

function.

***Sclerotinia sclerotiorum* has a closed pan-genome with relatively few non-syntenic**

blocks of genes

The gene-space within a pan-genome lies on a spectrum from high variability in certain

species to remarkable stability in others. Species harbouring a limited number of

dispensable genes are characterised by closed pan-genomes, while those with diverse

gene content are classified as having open pan-genomes [73]. To assess the openness of

the *S. sclerotiorum* pan-genome, we sampled from two to all 25 strains in our dataset and

plotted number of strains against number of novel genes. We found that the number of

additional genes brought by adding a new strain plateaued quickly at 5-10 strains,

indicating that most dispensable genes in the population are present in multiple strains

(Figure 4 A).

Across all strains, we identified 10,188 unique genes, of which 553 (5.43 %) were

dispensable. Few informative Gene Ontology terms were over-represented among these

genes, although we noted an over-representation of the term 'GO:0031177' (P = 0.012),

which is ascribed to genes containing a phosphopantetheine binding site. Since one of the

main functions of this site is in secondary metabolite biosynthesis by non-ribosomal peptide synthases (NRPSs) and polyketide synthases (PKSs), it is not surprising that we also found an over-representation of genes in secondary metabolite biosynthesis clusters among dispensable genes (odds ratio = 2.28, P = 1.44e-11). We found no significant over-representation of secreted proteins, regardless of size (odds ratio = 0.73, P = 0.18 for >= 300 amino acids; odds ratio = 1.38, P = 0.19 for <= 300 amino acids).

Based on a graphical representation of gene synteny, we identified 615 runs of one or more genes that were non-syntenic between strains (Figure 4 B). In keeping with graph terminology, we refer to these as 'gene bubbles'. The number of genes in gene bubbles ranged from 1 to 34, with most gene bubbles containing only a single gene (Figure 4 B). Consecutive runs of missing genes within bubbles ranged from 0 (i.e. the bubble was an inversion) to 13 (median = 1) (Figure 4 B). The largest three consecutive runs of missing genes within bubbles were identified on chromosome 12, which was the chromosome with the highest SVπ. Closer inspection of these runs identified a complex region partially duplicated in the strain R19, which was sampled in 2007 from buttercup in Warwickshire in the UK (Figure 4 C). Many of the genes in this region were likely transposon genes, as they were annotated with Pfam domains such as RNase H (PF00075), reverse transcriptase (PF00078), and endonuclease (PF14529) (Supplementary Table 9). However, there were also glycosyl hydrolases (PF00722), ubiquitins (PF00240) and a major facilitator superfamily transporter (PF07690). The fact that this region was the same in all strains apart from R19 could mean it is deleterious. Alternatively, it could be a relatively new polymorphism whose evolutionary fate has not yet been determined. So far, the polymorphism does not appear to be detrimental to infection on brassicas, since R19 is more aggressive than several other diverse isolates from the UK [74].

The closed *S. sclerotiorum* pan-genome contrasts the pan-genomes of host specialist fungal pathogens. For instance, in a population sample of 19 global isolates of *Z. tritici*,

approximately 40 % of genes were dispensable [8], and in 26 strains of the wheat

pathogen *Pyrenophora tritici-repentis* 43 % [75].

To our knowledge, little is known about what shapes pangenome openness in eukaryotes.

However, ecological theory suggests that selective pressure from the host is stronger on

host specialists than generalists [33]. To our knowledge, there are no *S. sclerotiorum*

strains unable to reproduce on a single host species or genotype. It is unlikely, therefore,

that a single virulence gene, such as an effector, would ever confer a strong host-driven

selective advantage in this species. Therefore, maintenance of a repertoire of dispensable

virulence proteins to ensure adaptability to a constantly changing host environment seems

unlikely. Instead, the closed pan-genome of *S. sclerotiorum* aligns with previous research

suggesting that it, and other host generalists, have evolved toward energetic optimisation

of core virulence genes that function on multiple host species [31,34].

Figure 4 Gene content variability in the *Sclerotinia sclerotiorum* pan-genome. A The

relationship between total number of unique genes (y axis) and number of genomes

sampled (x axis). **B** Number of gene bubbles (y axis) and number of genes they contained

(top) or number of consecutive missing genes they contained (bottom). **C** A region in the

1980 reference genome that had a partial duplication in the isolate R19 and no other

isolates. This region contained the largest three gene bubbles, indicated here with B1

(orange), B2 (yellow) and B3 (green). Start and end genes for each called bubble are

indicated in their respective colours and non-syntenic genes within bubbles are in light

grey. Neighbouring genes are in dark grey. The shaded area connects homologous

regions and the pink region is duplicated in R19.

Structural variation may have a strong impact on adaptive flexibility of life history

traits

Adaptive flexibility and fitness of a population are underpinned by genotypic variation that

impacts life history traits. As a global host generalist agricultural pest, *S. sclerotiorum* is

exposed to diverse environments and must be adaptable to a range of temperatures and

stressors, such as host metabolites. To assess global phenotypic diversity in *S.*

sclerotiorum, we measured 14 life history traits across 103 genotypically distinct strains,

including relative growth on the Brassicaceae defence compounds brassinin and

camalexin, the Fabaceae defence compound medicarpin, the reactive oxygen species

H_2O_2 (ROS), and the two fungicides azoxystrobin and tebuconazole; growth and relative

growth at 15, 20 and 25 °C; and fecundity-related traits including number, and average and

total weight of sclerotia.

We found significant differences between isolates from different geographical origins for

eight of these traits. Both European and Australian strains grew faster at 15 °C than

Canadian strains (Figure 5 A) (P = 0.014 and 0.007, respectively). At 20 °C, European

strains grew significantly faster than Australian but not Canadian strains (P = 0.003 and

0.52, respectively), though Canadian strains grew at a similar rate to Australian strains (P

= 0.40). At 25 °C, European strains grew faster than both Canadian and Australian strains

(P = 0.035 and 0.00072, respectively). Relative growth (growth divided by growth at 20 °C,

generally considered the middle of the optimum range [76,77]) at 15 °C was significantly

lower for both European and Canadian strains compared with Australian strains (P = 0.035

and 0.049, respectively), though relative growth at 25 °C was not significantly different

between strains from different global regions.

Differences in growth rate at different temperatures between these populations could be a

result of adaptation to prevailing temperatures during the growing season for major host

crops, a phenomenon that has been previously observed at a local level in Australia

[50,51,78]. However, it is difficult to completely align our observations with the likely

reproductive cycle of *S. sclerotiorum* in these three global cropping regions, as different

host crops are likely to be available at different times of year. For example, the major host

species *Brassica napus* usually flowers in spring in the UK, where temperatures are often

lower than when *B. napus* flowers in Western Australia in July. On the other hand, some

hosts, such as lettuce, may be also present later in the season in the UK. A weaker

adaptation to lower temperatures is possible for Canadian strains, which would likely infect

B. napus when it is flowering during the hotter summer months.

Among host antimicrobial metabolites, we found a significant increase in growth in

European strains compared with Canadian on camalexin (P = 0.015) and a significant

decrease on ROS (0.0015) compared with Australian strains. Growth on ROS was lower

for both European and Canadian compared with Australian strains, though the range in

growth of Canadian strains meant the difference between Australian and Canadian strains

was not significant (P = 0.12). Similarly, European strains had a significantly lower growth

rate on azoxystrobin compared with Australian strains (P = 0.035), and Canadian strains

were in the middle (P = 0.69 and 0.64 compared with Australian and European strains,

respectively).

Interestingly, Canadian strains produced a greater total sclerotia weight compared with

strains from Australia and Europe (P = 0.034 and 0.036, respectively). This seemed to be

due to a slight increase in the mean of both sclerotia number and weight. The size of

sclerotia has been previously linked with the rate of germination [79] and number of

apothecia per sclerotium [80], suggesting it is an important component of fecundity. Phenotypic variation in this trait may have important implications for pathogen proliferation and epidemic potential of different populations.

The traits we measured had complex genetic synergisms and antagonisms with one another, for instance brassinin and camalexin tolerance were positively correlated (Pearson's ρ = 0.43) and shared positive genetic covariance (0.93) (Figure 5 B, Supplementary Table 10). The same was true of camalexin and medicarpin tolerance (Pearson's ρ = 0.48, genetic covariance = 0.65). Other traits were negatively correlated and had negative genetic covariance, for instance growth at 20 °C and azoxystrobin tolerance (Pearson's ρ = -0.33, genetic covariance = -0.97), and total sclerotia weight and relative growth at 15 °C (Pearson's ρ = -0.39, genetic covariance = -0.63). This suggests that complex trade-offs and synergisms between life history traits may influence fitness [5]. Several studies have shown that SVs have a major role in creating phenotypic diversity, and graph pan-genomes in which SVs can be reliably called have shown that they can be a major component of missing heritability [19,81]. To test the relative impact of SVs on life history traits, we conducted a genome-wide association study (GWAS). For all traits, quantile-quantile plots suggested that the model we used adequately controlled P value inflation due to population structure (Supplementary Figure 4).

We found that the average absolute effect size of SVs was higher than that of non-SVs across 11 of the 14 traits, significantly higher for eight (P < 0.05), and lower for 2 of the 14 traits, tebuconazole and ROS tolerance. Notably absolute effect size was on average 0.015 and 0.020 points higher for relative growth on azoxystrobin and total sclerotia weight, respectively (P < 0.0001, Figure 5 C, Supplementary Table 11). On average, SVs had a lower average minor allele frequency than other variants (0.18 vs 0.21), which could lead to an increase in the variance of effect size estimates. Therefore, we took 500 random samples of non-SVs of equivalent size and minor allele frequency distribution to SVs and

assessed how many times their average absolute effect size was more than or equal to

that of the SVs. Based on this test, azoxystrobin tolerance, growth rate at 25 °C and

average sclerotia weight, were impacted more strongly by SVs than other variants in 100

% of random samples ($P = 0$), as well as having significant increases for the first test ($P <$

0.005) (Figure 5 C, Supplementary Table 11). Three other traits, medicarpin tolerance,

growth at 25 °C and total sclerotia weight were more strongly impacted compared with

more than 90 % of random samples ($P < 0.1$), as well as showing a significant increase

according to the first test ($P < 0.005$). According to the randomisation test, ROS and

tebuconazole tolerance were also significantly more weakly impacted on average by SVs

than other variants ($P = 1$ – all randomisations had a higher mean absolute effect size).

We infer from this analysis that SVs could have a larger impact on many of these traits

than other variants, though genetic architecture with respect to SVs likely varies

considerably between traits.

As an alternative to assessing the absolute effect size, we performed regressions on

different genomic relationship matrices, which either included or did not include all variants

in linkage disequilibrium with SVs. Using cross-validation, we found that for medicarpin

and ROS tolerance, growth at 15 °C and 20 °C, and sclerotia number, models that

included variants in LD with SVs had a better predictive ability than models that did not

(Pearson's $\rho = 0.25$ vs 0.23, 0.38 vs 0.37, 0.03 vs 0.02, 0.34 vs 0.29 and 0.14 vs -0.14,

respectively (Supplementary Table 12)). Given absolute effect size for ROS tolerance was

lower on average among SVs, it is possible that although individual SVs have a relatively

weak impact on this trait, as a collective they explain a relatively larger proportion of

additive heritability. For the other traits, the improvement in predictive ability was in

accordance with the increase in absolute effect size according to at least one of the

analyses mentioned previously.

Overall, we identified 15 variants with a significant (Benjamini-Hochberg adjusted P < 0.05)

impact on phenotype across six traits. 13 of these variants were intergenic SNPs or

InDels, one was a synonymous SNP and the other was a disruptive in-frame InDel. The

genes associated with these variants had diverse functions, which may be speculatively

associated with each of the traits (Supplementary Table 13). Though none of these

variants were SVs or in linkage disequilibrium with neighbouring SVs, one of them was a

48 bp InDel, close to the 50 bp cutoff we used for designating variants as 'structural'. This

variant, at position 107,298 on chromosome 12, was a deletion that significantly increased

relative growth on azoxystrobin ($P = 1.6e^{-5}$) (Figure 5 E). There were two genes either side

of this variant on opposite strands, one encoding a 78 amino acid protein with no known

domains and the other a 1,516 amino acid protein containing a centrosomin domain

(PF07989). The variant was, respectively, 1,177 and 2,571 bases away from the

transcription start sites of the shorter and longer genes. The amino acid sequences of both

proteins were well-conserved in fungi, with 80-100 % similarity to homologues from

species in the Helotiales, suggesting they are genuine genes (Supplementary Table 13).

With the current data, it is not possible to determine which, if either, of these genes'

functions is linked with growth rate on azoxystrobin. However, in fungi, centrosomins are

localised to spindle pole bodies, which are structures analogous to animal centrosomes,

the main sites of coordination of microtubule activity during mitosis [82]. It is conceivable

that the relative rate of cell division on a particular stressor could be impacted by mutations

affecting genes encoding the cell division machinery.

Our analyses are in accordance with several studies showing that SVs may have an

outsized impact on phenotype [19,83,84]. In our analyses, which used a GWAS approach,

we were restricted to common (minor allele frequency > 0.05), biallelic SVs. Therefore, we

likely favoured less deleterious SVs, since deleterious SVs with the strongest phenotypic

impacts are usually rare in populations [84,85]. Despite this, we observed an overall

stronger impact of SVs than other types of variants on many life history traits. This was

also despite the observation that SV diversity tended to cluster in polymorphic, repeat-rich

genome regions, which can often be sites of selective neutrality. Since the *S. sclerotiorum*

genome is relatively gene-dense, containing few repeat-rich regions, this could suggest

that the amount of selectively neutral structural variation it contains is relatively low.

Alternatively, the variation across many of the traits we assessed could be selectively

neutral or perhaps deleterious. In this case, the stronger link between SVs and phenotype

is indicative of SVs underlying a particularly large amount of adaptive potential.

Figure 5 Life history traits assessed across a subset of *Sclerotinia sclerotiorum*

strains. A Boxplots of measurements for life history traits (indicated to the right of the plot)

in four categories (to the left). The distribution is plot for strains from the three major

geographical regions, Australia, Canada and Europe. Points are the individual data points

and box and whisker plots show interquartile range. The letters a and b above plots

indicate significant differences between groups. **B** The top panel is a heatmap (rows are in

the same order as columns), showing Pearson's ρ between measurements for the 14 life

history traits. Colouring goes from green (negative correlation) to red (positive). The

dendrogram shows hierarchical clustering of the traits. **C** The distribution (y axis) across

the 14 traits of mean effect size of non-SVs subtracted from mean effect size of SVs (ΔE),

where effect sizes are absolute. **D** The y axis shows the density of measurements of

absolute effect size for 500 random samples with an identical minor allele frequency

distribution to that of SVs. The blue line shows the observed absolute effect size for SVs.

For these three traits, the P value of this test was 0. **E** A region surrounding a quantitative

trait locus (QTL) for relative rate of growth on azoxystrobin (top), with -log(P) on the y axis

and position in Mb on the x. The colours of points represent linkage disequilibrium of

variants at the different positions with the QTL (the purple point with associated P value).

The red line is a P value of 0.01. Below this region, the two genes neighbouring the 48 bp

InDel underlying the QTL are illustrated. These included a small gene with no known

domains and a larger centrosomin-encoding gene.

Conclusion

Collectively, our results portray *S. sclerotiorum* as both a clonal and sexually outcrossing

pathogen with limited diversity in gene content. Despite this limited diversity, *S.*

sclerotiorum isolates vary considerably in life history traits. SVs may make a particularly

strong contribution to this variability for some traits, and are likely generated through two

distinct mechanisms, meiotic recombination and transposition, the latter being the

dominant mechanism.

The limited genic diversity of *S. sclerotiorum* contrasts the highly variable open pan-

genomes of many host specialist species. Such stable gene content aligns with the

hypothesis that *S. sclerotiorum*, a niche generalist, is a 'jack of all trades', with a core,

multifunctional infective arsenal enabling fitness on hundreds of host species.

At this stage, the relative importance of meiosis and transposition in the generation of

adaptively advantageous SVs is unknown. Given the likely general evolution of *S.*

sclerotiorum towards a stable, repeat-poor genome, and the stronger contribution of SVs

to variability in life history traits, it is possible that meiosis, through its tendency to create

SVs, may have a significant role in adaptation beyond its well-recognised role in

recombination of alleles into new haplotypes.

Overall, our data shed considerable light on the evolutionary processes at play in an

important host generalist plant pathogen of agricultural significance.

Methods

Assessment of life history traits

A sclerotium of each isolate was cut in half, placed on a potato dextrose agar (PDA) plate,

and incubated at 20 °C for 4 to 5 days. Hyphae from the leading edge of the mycelium

were cut with a 3-mm cork-borer, placed onto fresh PDA plates and incubated at 20 °C for

2 days to source actively growing mycelium. Actively growing mycelia were subcultured

with a 3-mm cork-borer onto appropriate PDA plates for trait assessments.

To measure the effect of temperature on mycelium growth, mycelia were subcultured onto

PDA and grown at 15 °C, 20 °C and 25 °C for 1 day. To measure the effect of host

metabolites on mycelium growth, mycelia were subcultured onto PDA supplemented with

50 µM brassinin (Sigma-Aldrich), 20 µM camalexin (Sigma-Aldrich), 20 µg/mL medicarpin

(TargetMol), or 200 µg/mL hydrogen peroxide H_2O_2 (Westlab) and grown at 20 °C for 1

day. Brassinin, camalexin and medicarpin were all dissolved in DMSO prior to PDA

supplementation. To measure the effect of fungicides on mycelium growth, mycelia were

subcultured onto PDA supplemented with 0.2 µg/mL azoxystrobin and 50 µM

salicylhydroxamic acid (SHAM), or 0.16 µg/mL tebuconazole and grown at 20 °C for 1 day.

Azoxystrobin and tebuconazole were dissolved in ethanol and SHAM was dissolved in

water prior to PDA supplementation. All *S. sclerotiorum* strains were also grown on PDA

supplemented with the equivalent concentration of DMSO, ethanol, or ethanol and SHAM

as used for the aforementioned compounds. Though tebuconazole was dissolved in

ethanol, ethanol control plates were not available for the experiment, so growth on

tebuconazole was normalised to growth on DMSO. We did this because neither DMSO nor

ethanol strongly impacted growth, whereas growth rates were often variable between

experiments. For mycelium growth measurements on PDA, photographs were taken of

each inoculated PDA plate. The colony area was measured using Image J software.

Colony area relative to growth at 20 °C on PDA (plus appropriate solvent or SHAM) was

calculated for each isolate.

To measure the effect of temperature on sclerotia formation, mycelia were subcultured

onto PDA and grown at 20 °C for 1 month. Mature sclerotia were then air-dried for 3 days.

The number and weight of sclerotia per plate were recorded.

DNA extraction and sequencing

To extract high molecular weight DNA, sclerotia were cut with a sterile scalpel and placed

with the cut side touching the surface of the medium on potato dextrose agar (PDA) plates.

After three to four days at room temperature in darkness, strains were sub-cultured onto

fresh PDA plates from agar plugs using a sterile cork borer and forceps. After two further

days at room temperature in darkness, strains were sub-cultured again by placing four

plugs for each strain into 100 ml of potato dextrose broth (PDB) in 250 ml conical flasks.

These liquid cultures were incubated at room temperature with ambient light conditions in

the laboratory with shaking at 150 rotations per minute (RPM) for three days and used to

generate protoplasts.

Protoplasts were generated by removing fungal cultures that had grown around plugs and

placing two plugs each in 250 ml conical flasks with 40 ml enzymatic digestion solution

containing 0.8 M mannitol, 200 mM citric acid/tri-sodium citrate buffer and 1.5 % w/v lysing

enzymes from *Trichoderma harzianum* (L1412, Sigma, now discontinued). Digestions were

incubated for three hours at 30 °C with shaking at 80 RPM. All protoplasts from each

conical flask were then filtered through a 100 µm cell strainer (CLS431752, Merck) into

one 50 ml falcon tube and pelleted using a swinging bucket rotor centrifuge at 1,000-2,000

x g for 2-3 minutes at 4 °C. Protoplast pellets were re-suspended in 200 µl Tris-EDTA (pH

8.0).

The resuspended protoplasts were then used as input for the MagAttract high molecular

weight DNA extraction kit (67563, Qiagen), which was used with the manufacturer's

protocol for blood cells with the following modifications: 80 µl proteinase K was used

instead of 20 µl, 20 µl of RNase A was used instead of 4 µl, 600 µl of buffer AL was used

instead of 150 µl, 25 µl of MagAttract Suspension G was used instead of 15 µl, 600 µl of

buffer MB was used instead of 280 µl; before adding MagAttract Suspension G, samples

were also filtered through miracloth to remove debris. High molecular weight DNA was

then sequenced on an Oxford Nanopore MinION using an SQK-LSK109 library prep kit

multiplexed with the native barcoding expansion pack EXP-NBD104 on a R9.4.1 version

flowcell.

To extract DNA for Illumina sequencing, the same procedure was used for initial culturing

of *S. sclerotiorum* strains. Cultures from PDB were then snap frozen in liquid nitrogen and

freeze-dried overnight. Portions of approximately 1 g of freeze-dried samples were then

cut with a sterile scalpel and placed using forceps into 2 ml screw-capped Eppendorf tubes

with a single ball bearing. To each tube, 700 µl lysis buffer (50 mM Tris-HCL, 50 mM

EDTA, 3 % sodium dodecyl sulfate, 1 % 2-mercaptoethanol) was added, and samples

were ground in a MiniG model 1600 at 15000 RPM for 2 minutes. Samples were then

centrifuged at 17000 RPM for one minute, and ball bearings were removed. To each tube,

100 µl RNase A was added and tubes were then incubated for 1 hour at 65 °C. To each

tube, 700 µl chloroform:phenol (50:50) was added and tubes were vortexted. Tubes were

then centrifuged at maximum speed for 5 minutes before removal of the aqueous phase.

Then, 700 µl of chloroform:isamyl alcohol was added, the tubes vortexed and centrifuged

again at full speed for 5 minutes. The aqueous phase was again removed and DNA was

then precipitated using 6 M sodium acetate. Paired end Illumina sequencing was

conducted at Genomics WA on a NovaSeq 6000 flowcell at 2 x 150 cycles to yield 1.2 Gb

per sample. For genomic DNA methylation analysis, *S. sclerotiorum* 1980 (ATCC 18683)

was propagated on minimal salts – glucose (1% w/v) (MS–Glu) agar. The inoculum for all

experiments was prepared by grinding 2 g of sclerotia in 200 mL of MS–Glu in a Waring

blender for 4 min. The volume was increased to 500 mL in a 1 L baffled flask and the

culture incubated at 20 °C with shaking (60 r/min) for 3 days. 1 g of mycelia (wet mass)

was spread over a 5-cm-diameter area of *B. napus* leaf surface and incubated in a

humidified chamber. Leaves from 45-day-old plants were used. 3 biological replicates (3

different flasks of culture inoculated onto different leaves) were collected. The mycelial mat

was collected from the lesion using forceps at 48 hours post-inoculation, plant material

was removed, and the samples frozen immediately in liquid nitrogen. Samples were

ground in liquid nitrogen using a mortar and pestle, then genomic DNA was extracted from

a 100 mg sample using the DNeasy Plant Mini kit (Qiagen). Genome Quebec performed

whole genome bisulfite sequencing using the NEB Next kit, then sequenced 2x250 bp on

an illumina NovaSeq6000.

Trimming and demultiplexing reads

FAST5 files from the Oxford Nanopore were basecalled using Dorado version 0.3.2 and

de-multiplexed using Guppy version 6.5.7. Illumina whole genome sequencing reads were

trimmed using cutadapt version 2.8 [86] with appropriate adaptor sequences. Bisulfite

sequencing reads were assessed for quality and low quality bases and adapters were

trimmed using CLC genomics workbench 20.0.2.

Analysis of bisulfite sequencing data

Methylation analysis was performed using the CLC genomics workbench 20.0.2. Reads

were mapped to the *S. sclerotiorum* genome (GCF_000146945.2) using "Map Bisulfite

reads" (directional mapping and default mapping options). Methylated residues were

identified using "Call Methylation Levels" with default settings with the following

exceptions: exhaustive context-independent calls, minimum read depth of 10 reads. The

data presented in the results section is a count of methylated bases per 50 Kb sliding

window.

Genome assembly

Genomes were assembled from Nanopore reads using Flye version 2.8.1-b1676 [87] and

polished using Illumina reads, either from [29] or generated in this study, with one round of

Polypolish version 0.5.0 [88], followed by one round of Pilon version 1.24 [89]. Before

subsequent analyses, mitochondrial contigs were removed from assemblies using the

following procedure. Within Geneious Prime version 21.2.2, the Minimap2 version 2.24

[90] plug-in was used to align a published *S. sclerotiorum* mitochondrial genome (NCBI

accession KX351425) to each of the polished genomes. Contigs that aligned to this

accession with more than 95 % identity were separated from nuclear contigs, which we

focus on in this study.

Polished nuclear chromosomes for each genome were scaffolded to the *S. sclerotiorum*

reference genome [30] using the command 'scaffold', with the flags '-u -w -o', from RagTag

version 2.1.0 [91]. We then used the following process to finalise the scaffolded

assemblies. First, Nanopore reads for each assembly were self-corrected using Canu

version 2.2 [92]. Within Geneious Prime version 21.2.2, corrected reads were then aligned

to their respective assemblies using the Minimap2 version 2.24 plug-in and used to

manually add telomeres and subtelomeric sequences to the ends of chromosomes where

they could be recovered from reads. Gaps between scaffolded contigs were also removed

if there was extensive read support for joining the contigs.

Commands from Mummerplot version 3.1 [93] were then used to check for misassemblies.

First, 'nucmer' was used to align each assembly individually to the *S. sclerotiorum*

reference genome with the option '--mum'. The 'delta-filter' command was then used, with

the options '-1 -i 95 -l 10000 -u 100' to filter the output of nucmer. The filtered output was

then passed to the command 'show-coords' to produce coordinates of scaffold mappings

to the *S. sclerotiorum* reference genome. The command 'awk 'NR > 5 {strand="+"; if($2 <

$1){strand="-"};print $12"\t"$1"\t"$2"\t"$13"\t.\t"strand}'' was then used to convert these

coordinates into browser extensible data (BED) format. In Geneious, the BED file

containing alignments to 1980 and mappings of self-corrected reads were used to judge

whether chromosome segments had been artificially joined by the assembler. For

instance, if one chromosome was unusually large and contained two different segments

mapped to different 1980 chromosomes, it was split in two if (i) very few reads supported

the join and (ii) reads showed evidence of extensive soft clipping either side of the join.

Where chromosomes were split, genomes were scaffolded a second time using RagTag

and gaps between joined chromosome segments were removed if aligned reads

supported the join. The majority of chromosomes had zero gaps after the first round of

scaffolding and only two in each of three strains were broken and re-scaffolded based on

the latter procedure.

Genome annotation

For comparative purposes, all genomes, including the reference genome, were annotated

with the same procedure. First, repetitive sequences were annotated using EDTA version

2.2 [94] with the flag '--anno 1'. Then, Braker3 [95] was used to annotate genes with both

RNA sequencing and amino acid sequences as evidence with the additional flags '--

fungus', '--prot_seq=Fungi.fa', '--august_args="--species=botrytis_cinerea"'. The RNA

sequencing data used for annotation were derived from 32 samples from the sequence

read archive (SRA) detailed in Supplementary Table 14. Reads from these samples that

were derived from infected plant tissue were first filtered by alignment to their respective

host genomes (Supplementary Table 14) with Hisat2 version 2.1.0 [96] and keeping

unmapped reads with '--un-conc' for paired end reads or '--un' for single end reads.

Filtered reads, and reads not from plants, were then aligned to each of the *S. sclerotiorum*

genomes with Hisat2, converted to bam format with samtools version 1.10 [97] 'view' and

used as input for Braker3. Amino acid sequences from Braker3 annotations were

combined into a non-redundant set of genes for all isolates using cd-hit version 4.8.1 [98],

and non-redundant proteins were annotated with InterProScan version 5.54-87.0 [99].

Secondary metabolite clusters in this set of proteins were identified using antiSMASH

version 7.0 [100], and secreted proteins were identified with SignalP version 6.0 [101].

Pan-genome graph construction and variant calling

Using the 24 Nanopore assemblies and the *S. sclerotiorum* reference genome (GenBank

reference GCA_001857865.1), a pan-genome graph genome was constructed with cactus

2.5.2 [102]. Illumina reads from [29] and those generated in the current study were

mapped to the pan-genome GBZ formatted graph using the 'giraffe' command of vg

version 1.52.0 [18]. The resulting GAM formatted files, one for each set of Illumina reads,

were filtered using the vg command 'filter', with the flags '--min-primary 0.90 --frac-score --

substitutions --min-end-matches 1 --min-mapq 15 --defray-ends 999'. Filtered GAM files

were then passed to the vg command 'pack' to create pack formatted read support files for

each variant, with the flag '--min-mapq 5'. The 'call' command from vg was used to call

variants from the pan-genome graph using the Illumina reads with the flags '--ploidy 1 --

genotype-snarls' and create a variant call format (VCF) file. The VCF files for all samples

were combined into a single file by first converting them to gzip format with 'bgzip', then

merging them with the bcftools version 1.10.1 [103] command 'merge', with the option '--

all'. We then filtered this VCF with vcftools version 0.1.16 with the options '--minQ 30' and

'--minDP 5'. To do this, we had to first set all variants to 'PASS' because bcftools merge

adds a filter to the whole variant if only a single sample is filtered in one of the inputs. We

did this using a simple Awk script.

After calling variants present in the pan-genome graph, additional variants present in

Illumina reads but not in the 25 genomes that made up this graph were called using the

following procedure. First, filtered GAM files were converted to binary alignment map

(BAM) files using the vg command 'surject' and sorted using the samtools command 'sort'.

Then, the command 'mpileup' from bcftools was used with the flags '--max-depth 1000 --

output-type u' and the BAM files as input. The output of 'mpileup' was piped to the bcftools

command 'call', which was run with the flags '--output-type v --multiallelic-caller --ploidy 1'

to create a VCF file. We then filtered this VCF using vcftools with the options '--minQ 30', '-

-minGQ 30' and '--minDP 5'.

Finally, we used vcftools to remove variants called by vg from the VCF created using

bcftools with the options '--min-alleles 2', '--mac 1' and '--exclude-positions', and

concatenated the resulting VCF with the one produced using vg with the bcftools

command 'concat' with the option '--allow-overlaps'. We further filtered the final VCF with a

Python script (Supplementary File 2) to remove variants with a missing call rate of >= 0.2.

Population structure characterisation

To identify clones, a VCF containing variants called against the graph pan-genome was

used with plink version 1.9 [104] to generate an identical by state relationship matrix, with

the flags '--snps-only', '--biallelic-only', '--double-id', '--geno 0.2', '--mind 0.2' and '--make-

rel square 1-ibs'. Then, the matrix was used to construct a distance matrix and

dendrogram using hierarchical clustering. Clones were identified based on a relatedness

of 98 % identical by state with the R base function 'cutree'.

Population structure was analysed using ADMIXTURE version 1.3 [105]. As for the

identical by state relationship matrix, we considered only biallelic SNPs. These were first

filtered using plink with the flag '--indep-pairwise 50 10 0.1', and admixture was run for 1 to

10 ancestral populations with cross-validation. A scree plot was used to determine the

most appropriate number of populations to use based on cross-validation error. Principal

component analysis was also performed with plink using the flag '--pca 4'.

Assessment of structural variant diversity

To assess the diversity of SVs across the genome, we developed a novel statistic that we

refer to as SV_π, which is calculated as follows. First, we calculate SV_n, which is the sum of

the number of SVs between all pairs of individuals, excluding self-comparisons.

$$SV_n = \sum_{i=1}^{n-1} \sum_{j=i+1}^{n} SV_{ij}$$

865 Where SV_{ij} is the number of variants that are >= 50 bp in at least one individual for

866 individuals i and j in the set of n individuals in the sample. SV_n is then normalised in the

867 following way to obtain SV_π:

$$SV_\pi = \frac{SV_n}{\dfrac{k}{n^2}}$$

868 This divides SV_n by the number of possible pairs of individuals and the length of the

869 sequence under consideration, k. Since k varies between individuals depending on the SV

870 alleles they contain, it is calculated in the following way:

$$\frac{1}{n}\sum_{i=1}^{n} k_i$$

871 That is, k is the average value of all k_i sequence lengths, in Kb, in the set of n individuals.

872 The statistic is trivial for regions containing only biallelic SVs but more computationally

873 challenging for regions with multi-allelic variants.

The statistic SV_π is an estimate of the average number of SVs that are present per Kb between all pairs of individuals in the sample. It is an approximation of the genome stability in a region and may be better at identifying unstable genomic regions than considering simpler statistics such as proportion of rearranged sites or number of SVs relative to a single reference. The reason we developed this statistic was because we aimed to better capture the potential evolutionary rate of a region. For example, if considering the fraction of non-syntenic bases, a single large variant would create a high value, even if it is the only variant present. On the contrary, many diverse, small SVs would possibly cause a deflated estimate of the SV diversity of the region if their total length was a small proportion of the region's overall length. Though we do not present a detailed exposition of the method here, we present it as an intuitive and hopefully useful complementary technique for investigating structural diversity in pan-genomes. Our software for its calculation across sliding windows, svstats, is freely available on GitHub (https://github.com/markcharder/svstats). We used the program in this study to calculate SVpi in 50 Kb sliding windows across the genome with an increment of 1 Kb.

Analysis of linkage disequilibrium and recombination

To assess linkage disequilibrium decay with physical distance, linkage disequilibrium was first calculated for all pairs of variants between variants with the plink flags '--ld-window-r2 0', '–ld-window-kb 300' and '--r2 dprime'. R^2 was averaged for each physical distance and the distance at which average R^2 reached half its maximum value was recorded. The program phipack (obtained from https://www.maths.otago.ac.nz/~dbryant/software/PhiPack.tar.gz) was used to conduct three tests of the association between distance and linkage disequilibrium, the pairwise homoplasy index, maximum X^2, and nearest neighbour score tests.

To assess recombination rate, we selected four genotypically fairly uniform populations that had no obvious population structure. Recombination rate was calculated for these

populations using ldhat version 2.2 [57] and recombination hotspots were identified with

ldhot version 8.30 [106]. To run ldhat and ldhot, plink was first used to convert the pan-

genome VCF file to plink PED and MAP files with the flags '--recode', and '--biallelic-only'

and '--snps-only' to keep only biallelic SNPs. These were used as input for the command

'plink2ldhat' from our program 'svstats' (https://github.com/markcharder/svstats) to convert

to ldhat or ldhot format. A finite sites version of Watterson's theta was calculated using the

command 'watfsites' from svstats to provide a parameter for generating ldhat lookup

tables. The ldhat program 'complete' was then used, with flags '-rhomax 100' and '-n_pts

101', and the appropriate number of '-n' individuals, to create look-up tables for calculating

variable recombination rates across chromosomes. The ldhat interval program was then

used, with the appropriate look-up table, to calculate variable recombination rates with the

flags '-exact' '-its 10000000' and '-samp 3500'. Reversible jump Monte Carlo Markov

Chains were run starting with block penalties ranging from 5 to 50 (with an increment of 5)

and chains were assessed for convergence. Posterior distributions of rates and bounds

from the chains were estimated using the ldhat command 'stat', with the flag '-burn 35'.

Using the output of ldhat interval, ldhot was run using the appropriate look-up table with

the additional flag '--nsim 1000'.

Assessment of correlation between population-wide statistics and genomic features

The command 'makewindows' from Bedtools version 2.27.1 [107] was used to create

sliding windows of 50,000 bp, with an increment of 1,000 bp, across the *S. sclerotiorum*

genome. To calculate gene and repeat density, the bedtools command 'coverage' was

used with Braker3 and EDTA annotations, respectively, and the sliding windows. To

calculate GC content for windows, the command 'nuc' was used. Methylation rates from

bisulfite sequencing data were converted to BED format using a custom script in R, and

bedtools 'intersect' with the flag '-c' was used to calculate the number of methylated sites

per sliding window. A BED file was also created from the ldhat output, using a simple Awk

script, and used to calculate recombination rate for sites in sliding windows. The rate was

summed across sliding windows for comparison. Comparison between recombination rate

and other statistics of interest was conducted in R using Spearman's rank correlation.

The bedtools command 'closest' was used to determine the distance between transposon

annotations from EDTA and the nearest structural variant in all genomes. A Kruskall-Wallis

test in R was then used to determine whether any transposon classes were significantly

closer than other classes to the nearest SV.

Genome-wide association and trait correlation analyses

Before conducting GWAS and whole genome regression analyses, phenotype data were

normalised with the R package bestNormalize version 3.5. Two GWASs were run. One

(GWAS1) used variants that were filtered so that they were in approximate linkage

equilibrium using plink version 1.9 with the flag '--indep-pairwise 50kb 50 0.8', whereas the

other (GWAS2) did not. Both GWASs were conducted using GAPIT [108] with the BLINK

model. This model has been shown to adequately correct for population structure whilst

maintaining statistical power, and there were no non-genetic confounding factors between

populations as phenotypic data were collected in the same environment. We therefore

included no further population structure correction with, for example, principal components

or a kinship matrix. GWAS1 was used to identify significant marker trait associations as it

had fewer correlated markers than GWAS2 and therefore more statistical power. GWAS2

was used for the comparison of average absolute effects from structural and non-structural

variants.

To determine whether SVs had a larger impact on traits than other variants, we conducted

three tests. Firstly, we simply used standard t tests to compare the mean distributions of

absolute effect sizes of non-SVs and SVs. Since SVs had a lower minor allele frequency

on average than non-SVs, and this could affect variance of the test statistic, we developed

a randomisation test. This test sampled non-SVs 500 times, each time creating a random

set of non-SVs matching in number the total count of SVs. This random set was sampled

so that proportions of variants with all possible minor allele frequencies (rounded to three

significant digits) matched the minor allele frequency proportions in the SV set. The

average absolute effect size from GWAS2 was recorded for each of these 500 samples

and the number of times this effect size was larger than or equal to that of the mean

absolute effect size of the SVs was treated as the empirical P value.

In our third test, we partitioned variants into those that were in linkage disequilibrium with

structural variants and those that were not. We did this by first creating a file recording R^2

for all pairs of neighbouring variants within 2 Kb with the plink flags '--r2' and '--ld-window-

kb'. From this file, we created a list of variants that had an r2 of >= 0.5 with at least one

SV. This list, combined with the list of SVs themselves, was used to create two VCF files,

one containing SVs and variants in approximate linkage disequilibrium with them and the

other containing variants that were not SVs and were not in linkage disequilibrium with any

SVs. The two VCFs were filtered so that variants were not in strong linkage disequilibrium

with the plink command '--indep-pairwise 50kb 50 0.8'. Genomic relationship matrices

[109] were created for each of these sets of variants and for the whole set of variants used

in GWAS1 with the plink flag '--make-rel square'.

To assess genetic correlations between traits and determine whether adding SVs

improved predictive ability, we fit univariate and multivariate linear mixed models with the

R package sommer version 4.3.4 [110]. In these models, random effects for individuals

were estimated with assumed variance and covariance described by the genomic

relationship matrix proposed by Yang et al. (2011) [109]. To assess trait genetic

correlations, the genomic relationship matrix was estimated using all biallelic variants of

minor allele frequency >= 0.05. To assess improvement of prediction accuracy when

including SVs, we fit models with one random effect with variance structured by a

relationship matrix estimated with only non-SVs, and models with two random effects, one

structured by a non-SV and the other structured by an SV-only relationship matrix. For

each trait, we performed 'leave-one-out' cross validation. For each recording of each

phenotype for each strain, the recording was masked and the two models fit with this

recording missing. The predicted BLUP value based on the rest of the strains was

recorded for this version of the model and Pearson's correlation coefficient between model

predictions and phenotypes was recorded.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are available in NCBI,

under BioProjects PRJNA1112094 and PRJNA1120954, in the supplementary material, or

available upon reasonable request from the corresponding author.

Competing interests

The authors declare they have no competing interests.

Funding

MCD, YK, TN, PM, SJB, ARL, CG-T and LGK are funded by a co-investment between the

Grains Research and Development Corporation of Australia and Curtin University (project

code CUR00023). C-GT is also supported by a research training program (RTP)

scholarship from the Australian Government. SC acknowledges financial support from the

Institutional Development Plan (IDP) under National Agricultural Higher Educational

Project (NAHEP) of the Indian Council of Agricultural Research (ICAR) and World Bank.

Authors' contributions

MCD collected data, designed and executed experiments, oversaw research, conducted all main analyses and wrote the first manuscript draft. YK designed and executed experiments and edited the manuscript. TN designed and executed experiments and edited the manuscript. PM collected data, oversaw research, and edited the manuscript. SJB oversaw research and edited the manuscript. ARL collected data. CG-T collected data. SC designed and executed experiments and edited the manuscript. LB collected data and provided feedback on the manuscript. C Camplone executed experiments. SV executed experiments. DH designed and oversaw experiments and edited the manuscript. C Coutou designed and executed experiments and edited the manuscript. LGK initiated the project, oversaw research and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by resources provided by the Pawsey Supercomputing Research Centre's Setonix Supercomputer (https://doi.org/10.48569/18sb-8s43) and Nimbus research cloud (https://doi.org/10.48569/v0j3-qd51), with funding from the Australian Government and the Government of Western Australia.

References

1. Zhan J, Thrall PH, Papaïx J, Xie L, Burdon JJ. Playing on a Pathogen's Weakness: Using Evolution to Guide Sustainable Plant Disease Control Strategies. Annu Rev Phytopathol. 2015;53:19-43.

2. Burdon JJ, Thrall PH. Pathogen evolution across the agro-ecological interface: implications for disease management. Evol Appl. 2008;1:57-65.

3. Le May C, Montarry J, Morris CE, Frenkel O, Ravigné V. Editorial: Plant Pathogen Life-

History Traits and Adaptation to Environmental Constraints. Front Plant Sci.

2020;10:510672.

4. Atwell S, Corwin JA, Soltis N, Zhang W, Copeland D, Feusier J, et al. Resequencing

and association mapping of the generalist pathogen *Botrytis cinerea*. bioRxiv.

2018;489799.

5. Dutta A, Hartmann FE, Francisco CS, McDonald BA, Croll D. Mapping the adaptive

landscape of a major agricultural pathogen reveals evolutionary constraints across

heterogeneous environments. ISME J. 2021;15:1402-19.

6. Wyka S, Mondo S, Liu M, Nalam V, Broders K. A large accessory genome and high

recombination rates may influence global distribution and broad host range of the fungal

plant pathogen *Claviceps purpurea*. PLoS One. 2022;17.

7. Chen H, King R, Smith D, Bayon C, Ashfield T, Torriani S, et al. Combined pangenomics

and transcriptomics reveals core and redundant virulence processes in a rapidly evolving

fungal plant pathogen. BMC Biol. 2023;21:24.

8. Badet T, Oggenfuss U, Abraham L, McDonald BA, Croll D. A 19-isolate reference-quality

global pangenome for the fungal wheat pathogen *Zymoseptoria tritici*. BMC Biol.

2020;18:12.

9. Gourlie R, McDonald M, Hafez M, Ortega-Polo R, Low KE, Abbott DW, et al. The

pangenome of the wheat pathogen *Pyrenophora tritici-repentis* reveals novel transposons

associated with necrotrophic effectors ToxA and ToxB. BMC Biol. 2022;20:239.

10. Guo L, Dong Q, Wang B, Guo M, Ye K. Structural variants contribute to pangenome

evolution of a plant pathogenic fungus. bioRxiv. 2021;2021.04.14.439764.

11. Plissonneau C, Hartmann FE, Croll D. Pangenome analyses of the wheat pathogen

Zymoseptoria tritici reveal the structural basis of a highly plastic eukaryotic genome.

2018;16:5.

12. Dhakal U, Kim H-S, Toomajian C. The Landscape and Predicted Roles of Structural

Variants in *Fusarium graminearum* Genomes. G3 (Bethesda). 2024;14: jkae065.

13. Mahmoud M, Gobet N, Cruz-Dávalos DI, Mounier N, Dessimoz C, Sedlazeck FJ.

Structural variant calling: The long and the short of it. Genome Biol. 2019;20:1-14.

14. Hickey G, Monlong J, Ebler J, Novak AM, Eizenga JM, Gao Y, et al. Pangenome graph

construction from genome alignments with Minigraph-Cactus. Nat Biotechnol.

2023;42:663-73.

15. Eizenga JM, Novak AM, Sibbesen JA, Heumos S, Ghaffaari A, Hickey G, et al.

Pangenome Graphs. Annu Rev Genomics Hum Genet. 2020;21:139-62.

16. Hickey G, Heller D, Monlong J, Sibbesen JA, Sirén J, Eizenga J, et al. Genotyping

structural variants in pangenome graphs using the vg toolkit. Genome Biol. 2020;21:35.

17. Sirén J, Monlong J, Chang X, Novak AM, Eizenga JM, Markello C, et al. Pangenomics

enables genotyping of known structural variants in 5202 diverse genomes. Science.

2021;374: abg8871.

18. Garrison E, Sirén J, Novak AM, Hickey G, Eizenga JM, Dawson ET, et al. Variation

graph toolkit improves read mapping by representing genetic variation in the reference.

Nat Biotechnol. 2018;36:875-79.

19. Zhou Y, Zhang Z, Bao Z, Li H, Lyu Y, Zan Y, et al. Graph pangenome captures missing

heritability and empowers tomato breeding. Nature. 2022;606:527-34.

20. Badet T, Fouché S, Hartmann FE, Zala M, Croll D. Machine-learning predicts genomic

determinants of meiosis-driven structural variation in a eukaryotic pathogen. Nat Commun.

2021;12:3551.

21. Arbel-Eden A, Simchen G. Elevated Mutagenicity in Meiosis and Its Mechanism.

BioEssays. 2019;41:1800235.

22. Sakamoto T, Innan H. Muller's ratchet of the Y chromosome with gene conversion.

Genetics. 2022;220: iyab204.

23. Gordo I, Charlesworth B. The degeneration of asexual haploid populations and the

speed of Muller's ratchet. Genetics. 2000;154:1379-87.

24. Ruan Y, Wang H, Zhang L, Wen H, Wu C-I. Sex, fitness decline and recombination –

Muller's ratchet vs. Ohta's ratchet. bioRxiv. 2020;2020.08.06.240713.

25. Hinch R, Donnelly P, Hinch AG. Meiotic DNA breaks drive multifaceted mutagenesis in

the human germ line. Science (1979). 2023;382: eadh2531.

26. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten

things you should know about transposable elements. Genome Biology. 2018;19:1-12.

27. Oggenfuss U, Croll D. Recent transposable element bursts are associated with the

proximity to genes in a fungal plant pathogen. PLoS Pathog. 2023;19:e1011130.

28. Derbyshire MC, Newman TE, Khentry Y, Owolabi Taiwo A. The evolutionary and

molecular features of the broad-host-range plant pathogen *Sclerotinia sclerotiorum*. Mol

Plant Pathol. 2022;23:1075-90.

29. Derbyshire MC, Denton-Giles M, Hane JK, Chang S, Mousavi-Derazmahalleh M,

Raffaele S, et al. A whole genome scan of SNP data suggests a lack of abundant hard

selective sweeps in the genome of the broad host range plant pathogenic fungus

Sclerotinia sclerotiorum. PLoS One. 2019;14:e0214201.

30. Derbyshire M, Denton-Giles M, Hegedus D, Seifbarghi S, Rollins J, Kan J Van, et al.

The Complete Genome Sequence of the Phytopathogenic Fungus *Sclerotinia sclerotiorum*

Reveals Insights into the Genome Architecture of Broad Host Range Pathogens. Genome

Biol Evol. 2017;9:593.

31. Lyu X, Shen C, Fu Y, Xie J, Jiang D, Li G, et al. A Small Secreted Virulence-Related

Protein Is Essential for the Necrotrophic Interactions of *Sclerotinia sclerotiorum* with Its

Host Plants. PLoS Pathog. 2016;12:e1005435.

32. Newman TE, Kim H, Khentry Y, Sohn KH, Derbyshire MC, Kamphuis LG. The broad

host range pathogen *Sclerotinia sclerotiorum* produces multiple effector proteins that

induce host cell death intracellularly. Mol Plant Pathol. 2023;24:866-81.

33. Newman TE, Derbyshire MC. The Evolutionary and Molecular Features of Broad Host-

Range Necrotrophy in Plant Pathogenic Fungi. Front Plant Sci. 2020;11:591733.

34. Badet T, Peyraud R, Mbengue M, Navaud O, Derbyshire M, Oliver RP, et al. Codon

optimization underpins generalist parasitism in fungi. Elife. 2017;6: e22472.

35. Derbyshire MC, Raffaele S. Till death do us pair: Co-evolution of plant–necrotroph

interactions. Curr Opin Plant Biol. 2023;76:102457.

36. Bolton MD, Thomma BPHJ, Nelson BD. *Sclerotinia sclerotiorum* (Lib.) de Bary: biology

and molecular traits of a cosmopolitan pathogen. Mol Plant Pathol. 2006;7:1-16.

37. Attanayake RN, Tennekoon V, Johnson DA, Porter LD, del Río-Mendoza L, Jiang D, et

al. Inferring outcrossing in the homothallic fungus *Sclerotinia sclerotiorum* using linkage

disequilibrium decay. Heredity. 2014;113:353-63.

38. Atallah ZK, Larget B, Chen X, Johnson DA. High Genetic Diversity, Phenotypic

Uniformity, and Evidence of Outcrossing in *Sclerotinia sclerotiorum* in the Columbia Basin

of Washington State. Phytopathol. 2004;94:737-42.

39. Gomes EV, Breseguello L, Augusto M, Nasser LCB, Petrofeza S. Microsatellite

Markers Reveal Genetic Variation within *Sclerotinia sclerotiorum* Populations in Irrigated

Dry Bean Crops in Brazil. Journal of Phytopathology. 2011;159:94-9.

40. Sexton AC, Whitten AR, Howlett BJ. Population structure of *Sclerotinia sclerotiorum* in

an Australian canola field at flowering and stem-infection stages of the disease cycle.

Genome. 2006;49:1408-15.

41. Mert-Türk F, Ipek M, Mermer D, Nicholson P. Microsatellite and Morphological Markers

Reveal Genetic Variation within a Population of *Sclerotinia sclerotiorum* from Oilseed Rape

in the Çanakkale Province of Turkey. Journal of Phytopathology. 2007;155:182-7.

42. Hemmati R, Javan-Nikkhah M, Linde CC. Population genetic structure of *Sclerotinia*

sclerotiorum on canola in Iran. Eur J Plant Pathol. 2009;125:617-28.

43. Attanayake RN, Carter PA, Jiang D, Del Río-Mendoza L, Chen W. *Sclerotinia*

sclerotiorum Populations Infecting Canola from China and the United States Are

Genetically and Phenotypically Distinct. Phytopathol. 2013;103:750-61.

44. Attanayake RN, Porter L, Johnson DA, Chen W. Genetic and phenotypic diversity and

random association of DNA markers of isolates of the fungal plant pathogen *Sclerotinia*

sclerotiorum from soil on a fine geographic scale. Soil Biol Biochem. 2012;55:28-36.

45. Kohn LM. The clonal dynamic in wild and agricultural plant-pathogen populations.

Canad J Bot. 2011;73: 1231-40.

46. Anderson JB, Kohn LM. Clonality in soilborne, plant-pathogenic fungi. Annu Rev

Phytopathol. 1995;33:369-91.

47. Hambleton S, Walker C, Kohn LM. Clonal lineages of *Sclerotinia sclerotiorum*

previously known from other crops predominate in 1999-2000 samples from Ontario and

Quebec soybean. Canad J of Plant Pathol. 2002;24:309-15.

48. Buchwaldt L, Garg H, Puri KD, Durkin J, Adam J, Harrington M, et al. Sources of

genomic diversity in the self-fertile plant pathogen, *Sclerotinia sclerotiorum*, and

consequences for resistance breeding. PLoS One. 2022;17:e0262891.

49. Clarkson JP, Warmington RJ, Walley PG, Denton-Giles M, Barbetti MJ, Brodal G, et al.

Population Structure of *Sclerotinia subarctica* and *Sclerotinia sclerotiorum* in England,

Scotland and Norway. Front Microbiol. 2017;8:490.

50. Michael PJ, Lui KY, Thomson LL, Stefanova K, Bennett SJ. Carpogenic germinability of

diverse *Sclerotinia sclerotiorum* populations within the southwestern Australian grain belt.

Plant Dis. 2020;104:2891-7.

51. Michael PJ, Lui KY, Thomson LL, Lamichhane AR, Bennett SJ. Impact of

preconditioning temperature and duration period on carpogenic germination of diverse

Sclerotinia sclerotiorum populations in Southwestern Australia. Plant Dis. 2021;105:1798-805.

52. Jakobsen IB, Easteal S. A program for calculating and displaying compatibility matrices as an aid in determining reticulate evolution in molecular sequences. Comput Appl Biosci. 1996;12:291-5.

53. Smith JM. Analyzing the mosaic structure of genes. J Mol Evol. 1992;34:126-9.

54. Bruen TC, Philippe H, Bryant D. A simple and robust statistical test for detecting the presence of recombination. Genetics. 2006;172:2665-81.

55. Fayyaz A, Robinson G, Chang PL, Bekele D, Yimer S, Carrasquilla-Garcia N, et al. Hiding in plain sight: Genome-wide recombination and a dynamic accessory genome drive diversity in *Fusarium oxysporum* f.sp. *ciceris*. Proc Natl Acad Sci U S A. 2023;120:e2220570120.

56. Nieuwenhuis BPS, James TY. The frequency of sex in fungi. Philosophical Transactions of the Royal Society B: Biological Sciences. 2016;371:20150540.

57. McVean GAT, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. The Fine-Scale Structure of Recombination Rate Variation in the Human Genome. Science (1979). 2004;304:581-4.

58. Scelfo A, Fachinetti D. Keeping the Centromere under Control: A Promising Role for DNA Methylation. Cells. 2019;8: 912.

59. Talbert PB, Henikoff S. What makes a centromere? Exp Cell Res. 2020;389:111895.

60. Carneiro M, Ferrand N, Nachman MW. Recombination and Speciation: Loci Near Centromeres Are More Differentiated Than Loci Near Telomeres Between Subspecies of the European Rabbit (Oryctolagus cuniculus). Genetics. 2009;181:593:593-606.

61. Amselem J, Cuomo CA, van Kan JAL, Viaud M, Benito EP, Couloux A, et al. Genomic Analysis of the Necrotrophic Fungal Pathogens *Sclerotinia sclerotiorum* and *Botrytis cinerea*. PLoS Genet. 2011;7:e1002230.

62. Nei M, Li WH. Mathematical model for studying genetic variation in terms of restriction

endonucleases. Proc Natl Acad Sci U S A. 1979;76:5269-73.

63. Jiao WB, Schneeberger K. Chromosome-level assemblies of multiple Arabidopsis

genomes reveal hotspots of rearrangements with altered evolutionary dynamics. Nature

Communications. 2020;11:1-10.

64. Havecker ER, Gao X, Voytas DF. The diversity of LTR retrotransposons. Genome Biol.

2004;5:1-6.

65. Eickbush TH. Retrotransposons. Encyclopedia of Genetics. 2001;1699-701.

66. Boeke JD. The Unusual Phylogenetic Distribution of Retrotransposons: A Hypothesis.

Genome Res. 2003;13:1975-83.

67. Fouché S, Oggenfuss U, Chanclud E, Croll D. A devil's bargain with transposable

elements in plant pathogens. Trends in Genetics. 2022;38:222-30.

68. Amselem J, Vigouroux M, Oberhaensli S, Brown JKM, Bindschedler L V., Skamnioti P,

et al. Evolution of the EKA family of powdery mildew avirulence-effector genes from the

ORF 1 of a LINE retrotransposon. BMC Genomics. 2015;16:917.

69. Porquier A, Tisserant C, Salinas F, Glassl C, Wange L, Enard W, et al.

Retrotransposons as pathogenicity factors of the plant pathogenic fungus *Botrytis cinerea*.

Genome Biol. 2021;22:1-19.

70. Torres DE, Oggenfuss U, Croll D, Seidl MF. Genome evolution in fungal plant

pathogens: looking beyond the two-speed genome model. Fungal Biol Rev. 2020;34:136-

43.

71. Croll D, McDonald BA. The Accessory Genome as a Cradle for Adaptive Evolution in

Pathogens. PLoS Pathog. 2012;8:e1002608.

72. Charlesworth B. The Maintenance of Transposable Elements in Natural Populations.

Basic Life Sci. 1988;47:189-212.

73. Golicz AA, Bayer PE, Bhalla PL, Batley J, Edwards D. Pangenomics Comes of Age:

From Bacteria to Plant and Animal Applications. Trends in Genetics. 2020;36:132-45.

74. Taylor A, Coventry E, Jones JE, Clarkson JP. Resistance to a highly aggressive isolate

of *Sclerotinia sclerotiorum* in a *Brassica napus* diversity set. Plant Pathol. 2015;64:932-40.

75. Moolhuijzen PM, See PT, Shi G, Powell HR, Cockram J, Jørgensen LN, et al. A global

pangenome for the wheat fungal pathogen *Pyrenophora tritici-repentis* and prediction of

effector protein structural homology. Microb Genom. 2022;8:000872.

76. Clarkson JP, Fawcett L, Anthony SG, Young C. A Model for *Sclerotinia sclerotiorum*

Infection and Disease Development in Lettuce, Based on the Effects of Temperature,

Relative Humidity and Ascospore Density. PLoS One. 2014;9:e94049.

77. Shahoveisi F, del Río Mendoza LE. Effect of Wetness Duration and Incubation

Temperature on Development of Ascosporic Infections by *Sclerotinia sclerotiorum*. Plant

Dis. 2020;104:1817-23.

78. Uloth MB, You MP, Cawthray G, Barbetti MJ. Temperature adaptation in isolates of

Sclerotinia sclerotiorum affects their ability to infect *Brassica carinata*. Plant Pathol.

2015;64:1140-8.

79. Hao JJ, Subbarao K V., Duniway JM. Germination of *Sclerotinia minor* and *S.

sclerotiorum* Sclerotia Under Various Soil Moisture and Temperature Combinations.

Phytopathol. 2007;93:443-50.

80. Taylor A, Coventry E, Handy C, West JS, Young CS, Clarkson JP. Inoculum potential of

Sclerotinia sclerotiorum sclerotia depends on isolate and host plant. Plant Pathol.

2018;67:1286-95.

81. Edwards D, Batley J. Graph pangenomes find missing heritability. Nature Genetics

2022 54:7. 2022;54:919-20.

82. Jaspersen SL. Anatomy of the fungal microtubule organizing center, the spindle pole

body. Curr Opin Struct Biol. 2021;66:22.

83. Zong W, Wang J, Zhao R, Niu N, Su Y, Hu Z, et al. Associations of genome-wide

structural variations with phenotypic differences in cross-bred Eurasian pigs. J Anim Sci

Biotechnol. 2023;14:1-20.

84. Hämälä T, Wafula EK, Guiltinan MJ, Ralph PE, dePamphilis CW, Tiffin P. Genomic

structural variants constrain and facilitate adaptation in natural populations of *Theobroma*

cacao, the chocolate tree. Proc Natl Acad Sci U S A. 2021;118:e2102914118.

85. Chakraborty M, Emerson JJ, Macdonald SJ, Long AD. Structural variants exhibit

widespread allelic heterogeneity and shape variation in complex traits. Nature

Communications 2019;10:1.

86. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet J. 2011;17:10-2.

87. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using

repeat graphs. Nat Biotechnol. 2019;37:540-6.

88. Wick RR, Holt KE. Polypolish: Short-read polishing of long-read bacterial genome

assemblies. PLoS Comput Biol. 2022;18: e1009802.

89. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An

Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly

Improvement. PLoS One. 2014;9:e112963.

90. Li H. Minimap and miniasm: fast mapping and de novo assembly for noisy long

sequences. Bioinformatics. 2016;32:2103-10.

91. Alonge M, Lebeigle L, Kirsche M, Jenike K, Ou S, Aganezov S, et al. Automated

assembly scaffolding using RagTag elevates a new tomato system for high-throughput

genome editing. Genome Biol. 2022;23:1-19.

92. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable

and accurate long-read assembly via adaptive k-mer weighting and repeat separation.

Genome Res. 2017;27:722-36.

93. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile

and open software for comparing large genomes. Genome Biol. 2004;5:R12.

94. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al. Benchmarking

transposable element annotation methods for creation of a streamlined, comprehensive

pipeline. Genome Biol. 2019;20:1-18.

95. Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3:

Fully automated genome annotation using RNA-seq and protein evidence with GeneMark-

ETP, AUGUSTUS and TSEBRA. bioRxiv. 2023;2023.06.10.544449

96. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. Nat Methods. 2015;12:357-60.

97. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078-79.

98. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation

sequencing data. Bioinformatics. 2012;28:3150-52.

99. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, et al.

InterProScan: protein domains identifier. Nucleic Acids Res. 2005;33:W116-20.

100. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, et al. antiSMASH

7.0: new and improved predictions for detection, regulation, chemical structures and

visualisation. Nucleic Acids Res. 2023;51:W46-50.

101. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et

al. SignalP 6.0 predicts all five types of signal peptides using protein language models.

Nature Biotechnology. 2022;40:1023-5.

102. Armstrong J, Hickey G, Diekhans M, Fiddes IT, Novak AM, Deran A, et al.

Progressive Cactus is a multiple-genome aligner for the thousand-genome era. Nature.

2020;587:246-51.

103. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years

of SAMtools and BCFtools. Gigascience. 2021;10:1-4.

104. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK:

A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am J

Hum Genet. 2007;81:559-75.

105. Alexander DH, Lange K. Enhancements to the ADMIXTURE algorithm for individual

ancestry estimation. BMC Bioinformatics. 2011;12:1-6.

106. Auton A, McVean G. Recombination rate estimation in the presence of hotspots.

Genome Res. 2007;17:1219-27.

107. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26:841-2.

108. Wang J, Zhang Z. GAPIT Version 3: Boosting Power and Accuracy for Genomic

Association and Prediction. Genomics Proteomics Bioinformatics. 2021;19:629-40.

109. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: A Tool for Genome-wide Complex

Trait Analysis. Am J Hum Genet. 2011;88:76-82.

110. Covarrubias-Pazaran G. Genome-Assisted Prediction of Quantitative Traits Using the

R Package sommer. PLoS One. 2016;11:e0156744.

1301

Figures, tables and additional files

Figure legends

Figure 1. Genotypic clustering of *Sclerotinia sclerotiorum* strains from the global

population sample. A A phylogenetic network with all strains in the dataset coloured

according to geographical origin. The map inset shows where strains were collected with

colours corresponding to those in the network. The sizes of circles on the map

corresponds with the number of strains from each global region. **B** A phylogenetic network

for the Australian strains. Circles are coloured according to geographical origin within

Australia. Where circles are stacked on top of each other, isolates are a >= 98 %

genotypically identical group of clones. The map to the left shows where isolates were

collected within Australia, with colours of circles corresponding to colours on the network.

The sizes of circles represent the numbers of strains from each collection site. Haplotypes

1 (hap 1) and 2 (hap 2) are examples of frequently-sampled and geographically-

widespread clones, with individuals from Western Australia and South Australia.

Figure 2. Population structure and evidence of recombination. A Colours correspond

to ancestral populations making up individuals. Country of origin (above) is Au = Australia,

Mo = Morocco, SoA = South Africa, Ca = Canada, Fr = France, No = Norway, and UK =

UK. Below, states within Australia and Canada are indicated, where NSW = New South

Wales, SA = South Australia, WA = Western Australia, AB = Alberta, MB = Manitoba, and

SK = Saskatchewan. **B** Linkage disequilibrium (y axis) decay with physical distance (x

axis). Points are averages for unique distance measurements, and the red line is a general

additive model fit. **C** The first two principal components of genotypic variance. Colours

indicate geographical origin and point shapes the four population sub-samples used for

recombination analysis. **D** Across chromosomes and population sub-samples, the

distribution of Spearman's correlations between chromosome end distance and

recombination rate. **E** Correlation between coding DNA sequence content (x axis) and

recombination rate (y axis) of 50 Kb sliding windows. The line is a a general additive model

fit. **F** Boxplot showing percent gene content of 50 Kb windows containing and not

containing recombination hotspots (*** = $P < 2e^{-16}$). Boxes and whiskers show interquartile

range. **G** Circles show where windows containing putative centromeres lie on a plot of

recombination rate (y axis) against log recombination rate (x axis). Putative centromeres

are in regions of low recombination, before the inflection point. **H** The y axis is scaled

(division by maximum) recombination rate, amount of methylation or GC content for sliding

windows. The x axis shows position (Mb) across chromosome 6 (all chromosomes and

population samples are in Supplementary File 1). All chromosomes had a dip in GC coincident with a spike in methylation, almost always coincident with a recombination cold spot.

Figure 3. Analysis of structural variation across the *Sclerotinia sclerotiorum* pan-genome. A Distribution (y axis) of SVπ (x axis) for 50 Kb sliding windows. **B** For chromosomes 10 and 12, correlation between SVπ (x axis) and proportion transposon (top y axis) or coding DNA sequence (bottom y axis). Spearman's ρ and P value depicted top-right. Blue lines show linear regression of y onto x and the shaded area 95 % confidence interval. Red points are SVπ hotspot (> 95[th] percentile) windows. **C** The y axis shows distance to nearest structural variant (SV) for transposon families. Transposon classification is indicated at the top and family on the x axis. Boxes and whiskers show interquartile range. LTR retrotransposons were generally closer than other transposons to SVs (Kruskall-Wallis test show in Supplementary Table 6). **D** The y axis is SVπ or percent repeat for 50 Kb windows (scaled for visualisation). The x axis shows window start (Mb), and plots show chromosomes 6 and 12, the latter having the highest average SVπ and the most hotspots (shaded in pink). **E** Correlation between log recombination rate per Kb (y axis) and SVπ (x axis) across 50 Kb sliding windows. Chromosomes are plotted in different colours and data shown are for population-3. Spearman's ρ was 0.14-0.15 for all populations (P = 0) but varied between chromosomes. **F** Distribution across chromosomes (y axis) of Spearman's ρ for number of SVs and recombination rate in 50 Kb sliding windows. Though correlation strength varied between chromosomes, correlations were generally positive. **G** The y axis shows repeat content (top), SVπ (middle) and number of SVs (bottom) for windows that did not (left) and did (right) contain recombination hotspots. Boxes and whiskers show interquartile range; differences were significant according to a t-test (*** = $P < 2.2e^{-16}$).

Figure 4 Gene content variability in the *Sclerotinia sclerotiorum* pan-genome. A The relationship between total number of unique genes (y axis) and number of genomes sampled (x axis). **B** Number of gene bubbles (y axis) and number of genes they contained (top) or number of consecutive missing genes they contained (bottom). **C** A region in the 1980 reference genome that had a complex rearrangement in the isolate R19 and no other isolates. This region contained the largest three gene bubbles, indicated here with B1 (orange), B2 (yellow) and B3 (green). Start and end genes for each called bubble are indicated in their respective colours and non-syntenic genes within bubbles are in light grey. Neighbouring genes are in dark grey. The shaded area connects homologous regions and the pink region is duplicated in R19.

Figure 5 Life history traits assessed across a subset of *Sclerotinia sclerotiorum* strains. A Boxplots of measurements for life history traits (indicated to the right of the plot) in four categories (to the left). The distribution is plot for strains from the three major geographical regions, Australia, Canada and Europe. Points are the individual data points and box and whisker plots show interquartile range. The letters a and b above plots indicate significant differences between groups. **B** The top panel is a heatmap (rows are in the same order as columns), showing Pearson's ρ between measurements for the 14 life history traits. Colouring goes from green (negative correlation) to red (positive). The dendrogram shows hierarchical clustering of the traits. **C** The distribution (y axis) across the 14 traits of mean effect size of non-SVs subtracted from mean effect size of SVs (ΔE), where effect sizes are absolute. **D** The y axis shows the density of measurements of absolute effect size for 500 random samples with an identical minor allele frequency distribution to that of SVs. The blue line shows the observed absolute effect size for SVs. For these three traits, the P value of this test was 0. **E** A region surrounding a quantitative trait locus (QTL) for relative rate of growth on azoxystrobin (top), with -log(P) on the y axis and position in Mb on the x. The colours of points represents linkage disequilibrium of

1387   variants at the different positions with the QTL (the purple point with associated P value).

1388   The red line is a P value of 0.01. Below this region, the two genes neighbouring the 48 bp

1389   InDel underlying the QTL are illustrated. These included a small gene with no known

1390   domains and a larger centrosomin-encoding gene.

1391

Supplementary Material

Supplementary Figures

Supplementary Figure 1. A dendrogram showing the percentage of alleles identical

by state between strains in the collection. The green vertical line shows the cutoff used

to identify groups of individuals representing a single clone (blue).

Supplementary Figure 2. The relationship between recombination rate (y axis) and

coding sequence density (x axis) of 50 Kb sliding windows. The line is a general

additive model and the shading represents 95 % confidence intervals.

Supplementary Figure 3. SVπ and repeat content in 50 Kb windows across the

genome. The same as Figure 3 D but shown for all chromosomes.

Supplementary Figure 4. Q-Q plots for GWASs conducted for all traits. The y axis

shows observed P values and the x axis shows the expected P values given a normal

distribution. All plots show that most points are on (adequate correction) or below (over-

correction in some cases) the line, and P values are not inflated.

Supplementary Tables

Supplementary Table 1. A BUSCO scores for all strains used to construct the pan-

genome graph. **B.** Gaps and telomeres in each chromosome of each assembly. In the

TELOMERES column, L stands for 'left' and R stands for 'right', referring to the two

(arbitrary) ends of the chromosome in the assembly FASTA.

Supplementary Table 2. Strains, excluding the reference strain, 1980, used to create the

Sclerotinia sclerotiorum pan-genome and call structural variants. Strains with Nanopore

and Illumina data were used to create the pan-genome graph whereas strains with only

Illumina data (previous or current study) were used for mapping and variant calling against

the graph.

Supplementary Table 3. Results of phipack tests for recombination across the 120

independent *Sclerotinia sclerotiorum* lineages. These include the Neighbour Similarity

Score (NSS), the Maximum Chi^2 (MAX_CHI2), and the Pairwise Homoplasy Index (PHI)

tests. All tests were significant, with a P value of zero, indicating increasing levels of

recombination between alleles with distance.

Supplementary Table 4. Recombination hotspots identified relative to the *Sclerotinia*

sclerotiorum reference genome. Four non-structured population subsamples were used to

identify hotspots.

Supplementary Table 5. Cytosine methylation data from alignment of bisulfite sequencing

reads to the 1980 genome. The first column is the NCBI chromosome accession. The

columns for these tables are described in the CLC genomic workbench manual here:

https://resources.qiagenbioinformatics.com/manuals/clcgenomicsworkbench/current/index.

php?manual=Call_Methylation_Levels.html. Each spreadsheet represents one of the

samples, for example, 0 HPI R1 is 0 hours post-inoculation replicate 1.

Supplementary Table 6. Results of a Kruskal-Wallis test to determine whether LTR

transposons were significantly closer to structural variants across all genomes than other

transposons. Transposon classifications are taken from EDTA.

Supplementary Table 7. Transposable element content of the 24 Sclerotinia sclerotiorum

genomes based on EDTA annotations.

Supplementary Table 8. Spearman's correlation between estimated recombination rate

and SVπ, SV count and transposon content of 50,000 bp sliding windows. Rows coloured

in green are significant positive correlations, those in red are significant negative

correlations and those not coloured are not significant. Overall, the majority of

chromosomes and populations showed a correlation between recombination rate and both

SVπ and SV count but not transposon content.

Supplementary Table 9. Functional terms associated with genes in the largest gene

bubble. Results are from an InterProScan analysis. The Gene IDs are based on a cd-hit

grouping of Braker3 annotations across all genomes.

Supplementary Table 10. Genetic and actual correlations between life history traits.

Where genetic correlations are above 1, below -1 or 'NA', the model was likely poorly or

over-fit.

Supplementary Table 11. Tests for overall impact of SVs on phenotype. Grey cells are for

test statistics that were not significant. Green cells are for test statistics that indicate in

increase in SV impact on phenotype. Red cells are for test statistics that indicate a

decrease in SV impact on phenotype.

Supplementary Table 12. Linear mixed models testing improvement in predictive ability

(Pearson's ρ) from models with no SVs in the genomic relationship to matrix to models

with two terms, one for SVs and the other for non-SVs, or to models with only SVs.

Improvements in predictive ability were variable but some traits showed a relatively large

improvement.

Supplementary Table 13 A. Results of a GWAS for 14 life history traits. **B** BLASTp hits

for gene downstream of 48 bp InDel azoxystrobin QTL, which encodes a centrosomin. **C**

BLASTp hits for gene upstream of 48 bp InDel azoxystrobin QTL, which encodes a protein

with no known functional domains.

Supplementary table 14 A. RNA sequencing data used for Braker3 annotation of

genomes. **B** The host genomes used for filtering RNA sequencing reads used in Braker3

annotation.