Adult-neurogenesis allows for representational stability and flexibility in early olfactory system

Zhen Chen [1], Krishnan Padmanabhan [2]

1. Department of Brain and Cognitive Sciences, University of Rochester, Rochester, NY 14627

2. Department of Neuroscience, University of Rochester School of Medicine and Dentistry, Rochester, NY 14642.

* corresponding author: Krishnan Padmanabhan

Email: Krishnan_Padmanabhan@urmc.rochester.edu

Zhen Chen: ORCID iD: https://orcid.org/0000-0002-5590-3552

Krishnan Padmanabhan: ORCID iD: https://orcid.org/0000-0002-3255-8346

Abstract

In the olfactory system, adult-neurogenesis results in the continuous reorganization of synaptic connections and network architecture throughout the animal's life. This poses a critical challenge: How does the olfactory system maintain stable representations of odors amidst this ongoing circuit instability? Utilizing a detailed spiking network model of early olfactory circuits, we uncovered dual roles for adult-neurogenesis: one that both supports representational stability to faithfully encode odor information and also one that facilitates plasticity to allow for learning and adaptation. In the main olfactory bulb, adult-neurogenesis affects neural codes in individual mitral and tufted cells but preserves odor representations at the neuronal population level. By contrast, in the olfactory piriform cortex, both individual cell responses and overall population dynamics undergo progressive changes due to adult-neurogenesis. This leads to representational drift, a gradual alteration in stimulus-evoked activity patterns. Both processes are dynamic and depend on experience such that repeated exposure to specific odors reduces the drift due to adult-neurogenesis; thus, when the odor environment is stable over the course of adult-neurogenesis, it is spike-timing-dependent plasticity that leads representations to remain stable in piriform cortex; when those olfactory environments change, adult-neurogenesis allows cortical representations to track environmental change. Whereas perceptual stability and plasticity due to learning are often thought of as two distinct, often contradictory processes in neuronal coding, we find that adult-neurogenesis serves as a shared mechanism for both. In this regard, the quixotic presence of adult-neurogenesis in the mammalian olfactory bulb that has been the focus of considerable investigation in chemosensory neuroscience may be the mechanistic underpinning behind an array of complex computations.

Introduction

In the rodent brain, the main olfactory bulb (MOB) is one of two regions where neurogenesis persists throughout the animal's lifetime [1]. Adult-born cells in the sub-ventricular zone migrate to the MOB, where nearly 95% differentiate into inhibitory adult-born granule cells (abGCs) forming reciprocal dendro-dendritic connections [2,3] with the principal excitatory cells of the MOB, the mitral and tufted (M/T) cells. M/T cells in turn relay the chemosensory information to the downstream targets including the olfactory or piriform cortex (PCx) [4–6]. Although both the cellular composition and synaptic organization of this circuit undergo constant changes due to adult-neurogenesis [7], animals are nonetheless able to perform incredibly complex behavioral tasks related to the determination of odor identity and concentration, requiring that they maintain stable representations of odors. This raises two critical questions: Does the encoding of odors via patterns of neural activity throughout the early olfactory system undergo changes as a result of this ongoing plasticity, and if so, how? If changes in neural activity due to adult-neurogenesis percolate throughout early olfactory circuits, what computational principles allow the system to represent the same odor stably over time?

Previous studies have explored the plasticity at synapses between M/T cells and abGCs, particularly in relation to olfactory learning [8–10]. It is known that sensory experience influences the survival and synaptic turnover of abGCs with M/T cells [7,11], impacting both the normal activity of these cells [12] and animal's performance in odor discrimination tasks [13]. However, the implications of these local synaptic changes in the bulb on the downstream areas, including in the PCx, which is critical for olfactory perception and learning [14,15], remain largely unexplored.

The three-layer PCx is thought to be a region where the components of an odor are assembled into an olfactory percept or representation. Subsets of piriform cells (PCs) respond when multiple M/T cells are activated in complex temporal sequences [16–18]. A given population of M/T cells from a single glomerulus, a neuropil like structure where inputs from individual olfactory receptor neuron types converge [19,20], project randomly to a distributed array of PCs [4–6]. As odor information in PCx is coded for through the combinatorial activity of PCs [21,22] it is hypothesized that PCx is where stable representations of odors are maintained [23]. However, recent evidence is beginning to challenge this idea [24]. First, NMDA mediated long-term potentiation (LTP) in both afferent and associate fibers to PCx [25,26] is similar to that observed in the CA1 region of hippocampus. PCs thus undergo synaptic plasticity on scales observed in other regions where representations are not stable [27]. Second, recent studies examining the responses of PCs over months demonstrate that patterns of activity in both individual cells and populations of cells can drift over time [24]. This representational drift has led to the hypothesis that PCx, rather than being a primary sensory area, may be more like an association cortical region [28,29]. However, the mechanisms by which this drift in PCx occurs remain unknown.

One clue that adult-neurogenesis in MOB may be the mechanism underlying these diverse processes is that neurogenesis occurs on a similar time scale to the representational drift in PCx, and the turnovers of abGCs is highly experience dependent [9] which is also reflected in the PCx drift [24]. More frequent exposure to the same odor slows down the representational drift in PCx, while the drift rate increases again once this frequent exposure was halted [24].

We therefore hypothesized that the long-term plasticity conferred by adult-
neurogenesis in MOB percolates through the olfactory system serving as the mechanism
of representational drift in PCx. To test this hypothesis, we used a spiking neuronal
network model that replicated the circuit architecture within and between the MOB and
PCx, integrating both the ongoing network restructuring due to adult-neurogenesis and
the short-term plasticity of abGCs. We found that adult-neurogenesis differentially
modulated odor responses and representations in M/T cells in the MOB and PCs in PCx.
While the MOB maintained stable population representations despite changes in
individual M/T cell responses due to neurogenesis, the PCx shows variation at both
individual and population levels, leading to representational drift. Furthermore, when we
incorporated spike-timing-dependent plasticity (STDP) at the synapses on abGCs,
repeated odor exposure stabilized the PCx representations resulting in a reduction in drift
rate. Taken together, we identified how the rules of plasticity on short-term time scales
such as STDP and long-term time scales like adult-neurogenesis allow networks to both
preserve sensory representations in some circuits like the MOB, while allowing other
representations to change like those in PCx. Our work reveals the nuanced role played
by adult-neurogenesis in balancing stability and adaptability in the chemosensory system.
102
Results
Spiking network model of adult-neurogenesis in early olfactory system
To understand the functional role of adult-neurogenesis of GCs on the odor
representations in both MOB and PCx, we used a detailed spiking neural model that
recapitulates the circuit architecture and neural dynamics of the early olfactory system
(**Fig.1A1**, Methods). To this, we added adult-neurogenesis, which we modeled as a
process where a subset of granule cells are killed off and new granule cells are integrated
into the network as abGCs (**Fig.1A1**) [7,30]. We modeled this process by reshuffling the
synaptic weights of 10% of GCs in the network with the total number of GCs fixed. The
reshuffled synaptic weights include the weights between these abGCs and M/Ts, abGCs
and other GCs or short axon cells (SACs), and the top-down feedback projections from
PCs to abGCs (**Fig.1A2**). These new synaptic weights for abGCs were sampled from the
same distributions that generated the initial network model, such that although individual
weight values and connectivity were being modulated, the total distribution of synaptic
weights remain stable (**Fig.1B2**). We simulated a total of 11 days of neurogenesis from
Day-0 and Day-10 by which point almost all GCs in the model have gone through adult-
neurogenesis. Over the course, the synaptic weights between M/Ts and GCs, GCs/SACs
and GCs, PCs and GCs were changed to match the observed changes in experiments
(**Fig.1B1-1B3, FigS1**). Although the empirical rate of adult-neurogenesis has been found
to vary in rodent depending on the measurement approach and the behavioral paradigms
being used [31], we selected a rate of 10% for a reasonable computational time. Notably,
changing this rate in our model only affected the scale of the observations without
qualitatively changing the core results. These gradual changes in the network connectivity
allowed us to probe the long-term effect of adult-neurogenesis on odor processing and
the effect it had on modulating odor representations in MOB and PCx over time.
Odors drive coordinated activity in structures called glomeruli from which M/T cells
receive their information about the odors. To match the glomerular activation patterns
both in terms of the identify and onset timing in response to real odors [32–34], we generated

a panel of 100 model odors where each one activated 6%~20% of the total 50 glomeruli
in the model with different onset latencies (**Fig.1C1, FigS1**). Each model odor was
presented during a 250ms time window (4Hz sniff), corresponding to a single sniff that is
ethologically and behaviorally relevant time-scale for rodents [35–37]. Our generated panel
of model odors spanned a wide range of similarities, ranging from distinct odor pairs with
low pairwise correlations (see Methods) to highly similar odor pairs with high pairwise
correlations (**Fig.1C2**). The input to our network, the model odors at the glomerular level,
remained unchanged across simulated days consistent with previous observations of
stable odor responses of glomeruli [38]. Using this experimental paradigm, we could study
how the same odor inputs were differently encoded for over the course of adult-
neurogenesis by both the principal cells in the main olfactory bulb and the piriform cortex
[14,39].
143
Modulation of individual cell responses and preservation of population
representations in MOB
First, we examined how adult-neurogenesis affected the responses of both individual M/T
cells in the bulb and the population. We found that individual M/T cells changed their
responses to the same odor across days due to adult-neurogenesis, with some cells
decreasing the firing rate responses (**Fig.2A1** top) while other cells increased the
magnitude of their responses (**Fig.2A2** bottom, **Fig.S2**), similar to what has been
observed by others using calcium imaging of M/T cell activity [38,40]. These changes in gain
are consistent with a model of normalization, wherein the firing of an individual neuron is
modulated within a regime. Notably, individual M/T cell responses also exhibited some
extent of variability across repeated presentations of the same odor within a single
session (**Fig. S2**, error bars). This within-session variability occurs on a much shorter time
scale than adult-neurogenesis and therefore cannot be attributed to it. We return to this
point and its implications in the Discussion. However, when we examined the M/T
responses to the same odor across the population (**Fig.2B2**), we found that the overall
pattern of population responses was preserved across days despite the ongoing
neurogenesis. In this example, the odor activated 6 of the 50 glomeruli. Although
individual M/T responses changed (**Fig.2A1, 2A2** arrows), the groups of M/T cells driven
by the same glomerulus preserved the temporal structure of their firing as a whole. Each
individual M/T cell received certain amount of inhibition from a subset of GCs. With the
incorporation of abGCs that eliminated some connections, and randomly established new
ones, the synaptic weights of abGCs changed the inhibition at the individual M/T cell level,
but preserved the distribution of inhibition across the population of M/T cells associated
with each glomerulus. As a result, adult neurogenesis effectively preserved the ensemble
response to each odor for M/T cells associated with a single glomerulus but led to a
reshuffling of the identity of which M/T cells responded at what time epoch and with what
magnitude.
Next, we wanted to quantify the changes in M/T odor response across days due
to adult-neurogenesis. We took the odor-evoked firing rate over the 250ms window
across all M/T cells as a high-dimensional ensemble vector and computed their
correlation between Day-0 and each other day. We named this correlation as "*full-*
ensemble correlation". Regardless of using either single-trial (**Fig.2A3**) or trial-averaged
responses (**Fig.2A4**), the within-odor full-ensemble correlation decreased significantly

over time. This decrease is consistent with experimental observations [40]. The across-
odor correlations, computed from responses evoked by different odors, remained low
across all days (**Fig.2A4**). Therefore, these results indicated that the gradual changes in
the odor-evoked responses of individual M/T cells (**Fig.2A1**) accumulated over time.
However, as we observed above, the overall pattern of M/T population responses
was preserved (**Fig.2A2**), suggesting that the covariance, a measure of the shared
fluctuations across M/T population might remain stable. To test this, we projected M/T
population responses into a low-dimensional space (**Fig.2B1, 2B2,** see Methods), where
the time varying odor-evoked responses across the population constituted a trajectory in
the odor representation space of M/T cell population activity. For the two example odors,
the M/T trajectories of the same odor stayed quite close on the two example days
(**Fig.2B1**). If we simplified the single-trial trajectories by only plotting the maximal-distance
points to the origin (**Fig.2B2**), we found that the points of the same odor (color) stayed
clustered together in the space across all days (darkness of the color) despite the ongoing
circuit restricting due to adult-neurogenesis. The correlation of the low-dimensional
trajectories, which we named as "*reduced-space correlation*", remained high across all
days (**Fig.2B3**), and only dropped to ~0.7-0.8 on Day-10 (**Fig.2B4**). This is consistent
with a recent experimental study using calcium-imaging on M/T cells [41]. Additionally, while
the authors [41], observed relative stability of odor responses in some of the individual M/T
cells, they also observed that the full-ensemble showed a clear monotonic increase in
response differences over a long time intervals of months. In this regard, our results
showed that although adult-neurogenesis gradually varied the responses of individual
M/T cells, the low-dimensional representations of M/T cells in the reduced space
remained stable across days despite the changes in local synaptic organization due to
adult-neurogenesis.
202
Modulation of both individual cell responses and population representations in
PCx
M/T cells in MOB send random projections to PCx, such that each individual piriform cell
receives input from a random subset of M/T cells associated with different glomeruli [4,6].
PCx cells incorporate this feedforward information about odor identity and concentration
from the individual M/T cells in the MOB [14] and assembles those components into an
olfactory percept [21,23]. Although perceptual stability is a hallmark of sensory processing
and piriform cortex has historically been thought to be a place where perceptual stability
is established in the population code, recent evidence has started to challenge this idea
[24]. We hypothesized that the gradual changes in the individual M/T responses, passed
through the nonlinearities of synaptic integration, would affect the responses of PCs at
both individual and population level. We studied this by looking at the activity of individual
PCs and the ensembles of PCs in PCx.
Like M/T cells, the responses of individual PCs to the same odor varied across
days. Some PCs increased their responses while others decreased responses (**Fig.3A1**,
Fig.S3), consistent with recent chronic recordings in PCx (**Fig.S4**) [24]. However, unlike
M/T cells, the overall pattern of PCs responses varied as a result of adult-neurogenesis
(**Fig.3A2**). Some PCs responding strongly on Day-1 became inactive on Day-9, whereas
some PCs that did not respond on Day-1 became strongly activated on Day-9.
Consequently, the combinatorial patterns of activated PCs by the same odor were

reorganized by adult-neurogenesis on different days. Additionally, we also found that the
temporal structure of firing rate in individual PCs was highly variable. For example, some
PCs changed the onset and duration of their activation over the course of neurogenesis
(**Fig.3A2**). As a consequence, the within-odor full-ensemble correlation showed a
significant drop for PCs (**Fig.3A3, 3A4**). This is consistent with recent studies examining
the responses of PCs over time and demonstrating that patterns of PCs responses in
both individual cells and populations of cells can *drift* over time [24].
Changes in the firing rates reflect the changes in the variance of individual cells
(**Fig.S5**), but they do not say anything about the covariance, nor do they capture the
nonlinear interactions that shape the representations in PC populations when synaptic
information is integrated across M/T cell inputs and passes through the threshold
nonlinearity in individual PCs reflective of this integration. To study this, we calculated
the covariance across population of PCs was impacted by adult-neurogenesis. In
contrast to the M/T population where the representational trajectories of the same odor
followed one another closely in the space across different days, we found that the low-
dimensional trajectories of PCs showed a large variability across different days (**Fig.3B1**).
As a result, the maximal-distance points for the trajectories on each day were broadly
spread in the space across progressive days of neurogenesis (**Fig.3B2**). These results
are consistent with representational drift in the odor encoding space. Quantitatively, the
within-odor reduced-space correlation decreased substantially across days (**Fig.3B3,
3B4**). Our data suggests that as a consequence of both the random projections of M/T
cells to different PCs and the ways in which the cortical cells integrate those inputs, PCx
representations drift with ongoing adult neurogenesis in the bulb.
Next, we asked how adult-neurogenesis geometrically reshaped the odor manifold
and representational trajectories in the high-dimensional space? Recent computational
models that have been built on longstanding frameworks of population coding suggest
that neural representations reside in a high-dimensional manifold [42,43]. First, we reasoned
that the structure of the odor manifold would be defined as the aggregation of the odor-
evoked temporal responses within a sniff cycle to all the odors based on evidence
showing that different odors evoke complex and distinct temporal activities in both M/T
cells [44–46] and PCs [47–49]. For visualization, we plotted the odor manifold in the three-
dimensional PCA space (**Fig.S6**). We found the odor manifolds of M/T cells were highly
overlapping with each other, whereas for PCs they were much more separated. Across
individual odors, we quantified the geometrical reshaping of representational trajectories
by calculating the cosine similarity (see **Methods**) which measures the similarity between
two vectors. It has a value 1 if the two vectors have the same direction, and a value 0 if
they are orthogonal. We found that cosine similarity using the population firing rate of
M/T cells and PCs had a similar degree of decrease as the function of intervals (i.e.,
number of days between two representations) (**Fig.S7A1**). However, the cosine similarity
using representational trajectories remained stable for M/T cell but reduced for PCs
(**Fig.S7B1**).
If representations drift as a result of adult-neurogenesis, then behaviors too should
be impacted, drifting in a similar timescale. To unpack this connection, we used the K-
nearest neighbor algorithm as the decoder (see **Methods**). We found that the decoding
accuracy for discriminating against two different odors using M/T cell population

responses was high but accuracy in PCs dropped substantially capturing the
representational flexibility in PCx (**Fig.S7**).
270

Experience-dependent plasticity enhances representational stability in PCx
So far, we have shown that adult-neurogenesis, a process that changes the circuit
structure of the early olfactory system, is one mechanism by which representational drift
in PCx happens. Interestingly, it has been reported that repeated experience of an
odorant stabilizes odor representations and thus reduces the drift in PCx [24]. The
mechanism which drives drift also appears to stabilize that drift when the olfactory
environment is stable, suggesting that a core feature of the mechanism should be
plasticity. Since the synapses of abGCs have been shown to be highly plastic and
experience-dependent [8,9,50], we reasoned that experience-dependent plasticity of the
same abGCs that drive representational in PCx may also stabilize that drift in response
to the experience of encountering a stable olfactory environment.
To model experience-dependent changes at the level of synapses, we
implemented a spike-timing-dependent plasticity (STDP) rule [51–53] across the diversity of
the synapses related to an abGC (**Fig.4A1**). Depending on the spike timing interval of
pre- and post-synaptic neurons, the synaptic weight could be either facilitated or
suppressed (**Fig.4A2**, see Methods). Experience then was modeled as repeated
presentations of the same set of odors every day (50 trials for each odor per day) to our
network model, while adult-neurogenesis was occurring as previously described, with the
network presumably learning these stable representations in the pattern of synaptic
connectivity. In this example GC, the adult-neurogenesis that took place on day-4,
resulted in a new cell being added to the circuit, while the existing cell was removed, a
process that randomly reset a group of synaptic weights (say, S1 and S2, **Fig.4B**) across
two days of neurogenesis. Without STDP, the synaptic weights S1 and S2 would keep
constant or fluctuate by some stochastic process, regardless of what was happening in
the environment, including the effects of the repeated experience of an odor (black line,
Fig.4B). The implementation of STDP meant that the synaptic weight may change on a
trial-by-trial basis when an odor was repeatedly presented (blue line, **Fig.4B**). In the end,
the weight S1 may be enhanced while the weight S2 may be suppressed. Although odor
representations of M/T cells remained stable with and without STDP (**Fig.4C1**), we found
that odor representations in PCx were stabilized with STDP (**Fig.4C2**). The correlation
drops from day-0 were significantly reduced (0.8 without STDP and 0.4 with STDP by
day-10, **Fig.4C2** left), and the drift rate (° per day), which quantified the daily angle
changes of the population responses (see **Methods**), was reduced by 26% (7.9 without
STDP and 5.8 without STDP, **Fig.4C2** right). Adding STDP to the network and using
repeated presentations of an odor meant that the synapses that would otherwise
randomly change due to adult-neurogenesis converged asymptotically to some weight
(**Fig.S8**). Importantly, we identified that while the brain is constantly undergoing change,
especially in the MOB and the PCx due to ongoing adult-neurogenesis, the impact of
these changes on both population responses and the representation depends as much
on the plasticity rules as it does on the ongoing statistics of the environment.
311

Discussion

In our study, we modeled adult-neurogenesis as a dynamic reshuffling of granule cells
(GCs) within a detailed spiking neuronal network that replicated the circuit architecture of
the main olfactory bulb (MOB) and piriform cortex (PCx). This modeling revealed how
adult-neurogenesis differentially modulates odor responses and representations in mitral
and tufted (M/T) cells and PCs.
In the MOB, individual M/T cells exhibited variable odor responses with changing
firing rate magnitudes over time. This is consistent with earlier experimental studies using
calcium-imaging [38,40,41]. Variability in M/T cell responses arises on multiple time scales.
Experimental studies show pronounced trial-to-trial variability within single recording
sessions [37,54–58], which cannot be attributed to adult-neurogenesis. This fast variability
likely reflects ongoing network dynamics [59], behavioral state [38,60,61], attention [62], or top-
down modulation [48,63,64], none of which are included in our model. Our model therefore
captures only a limited amount of within-session variability and instead focuses on slower
changes that accumulate across days. In this regime, adult-neurogenesis provides a
mechanism for gradual changes in individual M/T responses. It would be interesting for
future work to integrate both fast and slow sources of variability within a unified modeling
framework.
Despite these fluctuations in individual cells, the overall pattern of M/T cells
population responses remained stable. This stability is attributed to consistent glomerular
input, as reflected in the low-dimensional M/T cell trajectories and their proximity in
reduced PCA space. This stability of M/T cells we observed in the reduced space is
consistent with a recent experimental study [41] using both linear (PCA) and nonlinear (t-
SNE) dimensionality reduction methods. Prior experimental studies report heterogeneous
effects of experience on mitral cell odor responses. Repeated odor exposure has been
associated with increased sparsening and reduced response amplitude in some
experiments [38], whereas others report substantial reorganization of ensemble
representations across days without consistent sparsening [40]. Our model best
recapitulates the findings of the latter, showing that adult-neurogenesis can reorganize
individual M/T responses while preserving stable low-dimensional population structure.
Differences in behavioral features across studies may reflect variations in behavioral
context or top-down modulation, factors that are not explicitly modeled here.
Conversely, in the PCx, both individual and population patterns of PCs responses
were in constant flux due to the ongoing adult-neurogenesis. This resulted in a geometric
reshaping of odor representations and a drift in the odor manifold defined by those
representations, consistent with experimental finding[24]. Our results reveal how the
process of adult-neurogenesis may support a number of computations in the early
olfactory circuits, including how a sensory representation may remain stable, and how it
might change due to plasticity.
We note that our simulations do not assume complete replacement of the granule
cell population over the 11-day period. Rather, each simulated "day" represents a discrete
epoch used to implement plasticity across multiple time scales. While some
developmentally born granule cells can persist for much of the animal's lifetime [65], some
fraction of the population undergoes turnover through adult neurogenesis [7,66,67]. In our
model, turnover is applied uniformly across granule cells, without preferential survival.
Additionally, different simulated time points may also be thought of as reflecting regimes
ranging from higher turnover to partial preservation of existing granule cells. While

behavioral factors such as novelty or running can modulate neurogenesis rates [68–70],
these effects are not included in our study and would be important for future studies.

Second, we found that one of the roles for STDP in the early olfactory system is
reducing representational drift in the PCx.   Repeated exposure to an odor significantly
reduced the representational drift in the PCx, consistent with experimental observations
[24]. Rather counterintuitively, this result suggests that a role of STDP and plasticity in
general is stabilizing representations despite ongoing changes in the circuit. This
implicates STDP at abGC-related synapses in a broader functional context. Previous
experimental studies have shown that abGCs integrate into the olfactory bulb in an
activity-dependent manner and preferentially stabilize circuits engaged by behaviorally
relevant odors [9,50,71]. Our model is consistent with this view, but approaches the problem
at the level of synaptic plasticity rather than cell survival. We show that STDP acting on
abGC synapses can reduce representational drift by selectively stabilizing dimensions of
population activity that are repeatedly activated due to experience. Here, representational
stability arises because of the stability of the sensory world; Representational
permanence, rather than being a feature of the circuit, is a feature of the sensory
landscape, a stabilizing force against the inherent variability introduced by continuous
adult neurogenesis. Adult-neurogenesis, which occurs throughout the animal's life, may
confer a mechanistically different computational framework as compared to vision or
audition, for which critical periods delineate the bounds of plasticity, and define the
periods over which sensory representations are changed or stabilized.

Third, our various dimensionality reduction analyses aim to illustrate the structure
of population activity in terms of the variance and covariance of neuronal responses;
which features of this activity are most relevant likely depend on what downstream brain
regions receive this input and what computations these regions ultimately perform. In this
framework, both low and high-dimensional representations would be functionally relevant
but may serve different roles. In the bulb, the low-dimensional population structure
remains stable despite ongoing reorganization in the full ensemble, consistent with its
role in providing a reliable sensory code. In contrast, PCx exhibits drift even in its low-
dimensional space, reflecting its more flexible and higher-dimensional representational
geometry. This higher dimensionality arises from divergent bulb inputs, dense recurrent
connectivity, and integration of contextual and feedback signals. Importantly, STDP
reduces drift in piriform cortex by stabilizing the dimensions reinforced by experience.
Residual drift persists because the cortical population retains many orthogonal
dimensions through which representations can continue to evolve.

Fourth, our model assumes broadly distributed projections from MOB to PCx and
random intra-piriform connectivity [22,28,72]. These divergent projections from MOB to PCx
is central to the combinatorial coding strategy of PCx, allowing individual pyramidal
neurons to represent complex mixtures and higher-order odor features [14,73,74]. On the
other hand, recent studies have revealed increasing structure in olfactory connectivity,
including molecularly defined projection neuron subtypes, long-range functional loops,
and experience-dependent motifs within piriform cortex [75–77]. These findings point to
additional constraints that may shape learning and perception. An important direction for
future work will be to incorporate such structured connectivity and cortical plasticity into
our model to examine how they interact with adult-neurogenesis in the bulb to influence
representational drift in PCx.

In addition to PCx, M/T cells also project to other cortical areas such as the anterior olfactory nucleus (AON). Although our model does not differentiate between mitral and tufted cells, it has been recently reported that mitral and tufted cells have distinct preferred cortical targets, with mitral cells preferentially projecting to PCx and tufted cells preferentially projecting to AON [75]. Tufted cells substantially outperform mitral cells in decoding both odor identity and intensity, suggesting that tufted cells and the AON where they project to are ideal for stably odor identity than the mitral cells and PCx [75]. As mitral cells are located in the deeper layers of the bulb, they interact with abGCs, whereas tufted cells in the superficial layer may preferentially interact with existing GCs that were born in the neonatal period [78,79]. Therefore, the activity of mitral cells may be more impacted by the turn-overs of abGCs, and the effect of adult-neurogenesis percolates onto PCx. By contrast, adult-neurogenesis has less impact on the tufted cells and the AON, potentially contributing to the stable encoding of odors in AON. This is one prediction of our model.

While our study focused on the effects of adult-neurogenesis in the olfactory bulb on odor representations, it does not negate the possibility of other forms of plasticity [80] contributing to the representational drift observed in the PCx [24]. Indeed, similar representational drifts have been noted in other brain areas like the posterior parietal cortex (PPC) [81], primary motor cortex [82], and hippocampus [83,84]. Of these regions, adult-neurogenesis only occurs in the hippocampus, where adult-born granule cells integrate into the existing circuitry of dentate gyrus [85]. Interestingly, similar to what we found in the early olfactory system (i.e., new-born cells in MOB and drift in PCx), the cells downstream of the dentate in the CA3 region [86] and CA1 region [27,83,84] exhibit drift over days and weeks.

Both PCx and hippocampus are evolutionarily old archi-cortical structures. This conservation of adult-neurogenesis in these brain regions may suggest a broader evolutionary strategy for balancing stability and flexibility. First, adult-neurogenesis in the olfactory bulb may support ongoing plasticity due to continuous peripheral turnover of olfactory receptor neurons [7,87]. In this context, abGCs may provide a mechanism for adapting bulb circuitry to changing input statistics, injury, or learning demands. Second, adult-neurogenesis may confer selective advantages for navigating dynamic environments. By ensuring flexibility in neural circuits, such as through adult-neurogenesis, these regions can rapidly update the representations of odors and spatial information, respectively, in response to changing environmental cues. This flexibility may enable the early organisms to effectively represent and adapt to the dynamic worlds they inhabited, enhancing their chances of survival and reproduction across evolutionary time scales [90–92]. As organisms evolved, evolutionarily newer brain regions such as the PPC, lacking significant adult-neurogenesis, developed alternative mechanisms to balance plasticity and stability [80]. Our work thus offers a new perspective on how the brain adapts to an ever-changing sensory environment.

446 **Acknowledgments**

447 This study was supported by funding from the National Institutes of Health (NIH) and the
448 National Science Foundation (NSF). KP was funded by NIH R01 MH113924,
449 DC021141, NS135763, and NSF CAREER 1749772. This manuscript has been
450 released as a pre-print.
451

452 **Author contributions**

453 K.P. conceived and supervised the project. Z.C. performed the experiments
454 and analyses. Z.C. and K.P. created the figures and wrote the manuscript.
455 Both authors approved the submitted version.
456
457 **Competing interests**

458 The authors declare no competing interests.

References

1. Lledo, P.-M., Alonso, M. & Grubb, M. S. Adult neurogenesis and functional plasticity in neuronal circuits. *Nat Rev Neurosci* **7**, 179–193 (2006).
2. Urban, N. N. & Sakmann, B. Reciprocal intraglomerular excitation and intra- and interglomerular lateral inhibition between mouse olfactory bulb mitral cells. *The Journal of Physiology* **542**, 355–367 (2002).
3. Schoppa, N. E. & Urban, N. N. Dendritic processing within olfactory bulb circuits. *Trends in Neurosciences* **26**, 501–506 (2003).
4. Sosulski, D. L., Bloom, M. L., Cutforth, T., Axel, R. & Datta, S. R. Distinct representations of olfactory information in different cortical centres. *Nature* **472**, 213–216 (2011).
5. Ghosh, S. *et al.* Sensory maps in the olfactory cortex defined by long-range viral tracing of single neurons. *Nature* **472**, 217–220 (2011).
6. Miyamichi, K. *et al.* Cortical representations of olfactory input by trans-synaptic tracing. *Nature* **472**, 191–196 (2011).
7. Lepousez, G., Valley, M. T. & Lledo, P.-M. The Impact of Adult Neurogenesis on Olfactory Bulb Circuits and Computations. *Annu. Rev. Physiol.* **75**, 339–363 (2013).
8. Lepousez, G. *et al.* Olfactory learning promotes input-specific synaptic plasticity in adult-born neurons. *Proceedings of the National Academy of Sciences* **111**, 13984–13989 (2014).
9. Livneh, Y. & Mizrahi, A. Experience-dependent plasticity of mature adult-born neurons. *Nat Neurosci* **15**, 26–28 (2012).
10. Nissant, A., Bardy, C., Katagiri, H., Murray, K. & Lledo, P.-M. Adult neurogenesis promotes synaptic plasticity in the olfactory bulb. *Nat Neurosci* **12**, 728–730 (2009).
11. Lledo, P.-M. & Valley, M. Adult Olfactory Bulb Neurogenesis. *Cold Spring Harb Perspect Biol* **8**, a018945 (2016).
12. Breton-Provencher, V., Lemasson, M., Peralta, M. R. & Saghatelyan, A. Interneurons produced in adulthood are required for the normal functioning of the olfactory bulb network and for the execution of selected olfactory behaviors. *Journal of Neuroscience* **29**, 15245–15257 (2009).
13. Kouremenou, I., Piper, M. & Zalucki, O. Adult Neurogenesis in the Olfactory System: Improving Performance for Difficult Discrimination Tasks? *BioEssays* **42**, 2000065 (2020).
14. Bekkers, J. M. & Suzuki, N. Neurons and circuits for odor processing in the piriform cortex. *Trends in Neurosciences* **36**, 429–438 (2013).
15. Choi, G. B. *et al.* Driving Opposing Behaviors with Ensembles of Piriform Neurons. *Cell* **146**, 1004–1015 (2011).
16. Smear, M., Resulaj, A., Zhang, J., Bozza, T. & Rinberg, D. Multiple perceptible signals from a single olfactory glomerulus. *Nat Neurosci* **16**, 1687–1691 (2013).
17. Chong, E. *et al.* Manipulating synthetic optogenetic odors reveals the coding logic of olfactory perception. *Science* **368**, eaba2357 (2020).
18. Gill, J. V. *et al.* Precise Holographic Manipulation of Olfactory Circuits Reveals Coding Features Determining Perceptual Detection. *Neuron* **108**, 382-393.e5 (2020).
19. Mombaerts, P. *et al.* Visualizing an Olfactory Sensory Map. *Cell* **87**, 675–686 (1996).
20. Buck, L. & Axel, R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* **65**, 175–187 (1991).
21. Davison, I. G. & Ehlers, M. D. Neural Circuit Mechanisms for Pattern Detection and Feature Combination in Olfactory Cortex. *Neuron* **70**, 82–94 (2011).
22. Stern, M., Bolding, K. A., Abbott, L. & Franks, K. M. A transformation from temporal to ensemble coding in a model of piriform cortex. *eLife* **7**, e34831 (2018).

23. Pashkovski, S. L. *et al.* Structure and flexibility in cortical representations of odour space. *Nature*
583, 253–258 (2020).
24. Schoonover, C. E., Ohashi, S. N., Axel, R. & Fink, A. J. P. Representational drift in primary olfactory
cortex. *Nature* **594**, 541–546 (2021).
25. Kanter, E. D. & Haberly, L. B. NMDA-dependent induction of long-term potentiation in afferent and
association fiber systems of piriform cortex in vitro. *Brain research* **525**, 175–179 (1990).
26. Jung, M. W. & Larson, J. Further characteristics of long-term potentiation in piriform cortex. *Synapse*
18, 298–306 (1994).
27. Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nature neuroscience* **16**, 264–266
(2013).
28. Haberly, L. B. & Bower, J. M. Analysis of association fiber system in piriform cortex with intracellular
recording and staining techniques. *Journal of Neurophysiology* **51**, 90–112 (1984).
29. Haberly, L. B. Parallel-distributed Processing in Olfactory Cortex: New Insights from Morphological
and Physiological Analysis of Neuronal Circuitry. *Chem Senses* **26**, 551–576 (2001).
30. Aimone, J. B. & Gage, F. H. Modeling new neuron function: a history of using computational
neuroscience to study adult neurogenesis. *European Journal of Neuroscience* **33**, 1160–1169 (2011).
31. Pignatelli, A. & Belluzzi, O. Neurogenesis in the Adult Olfactory Bulb. in *The Neurobiology of*
Olfaction (ed. Menini, A.) (CRC Press/Taylor & Francis, Boca Raton (FL), 2010).
32. Fantana, A. L., Soucy, E. R. & Meister, M. Rat Olfactory Bulb Mitral Cells Receive Sparse Glomerular
Inputs. *Neuron* **59**, 802–814 (2008).
33. Gschwend, O., Beroud, J., Vincis, R., Rodriguez, I. & Carleton, A. Dense encoding of natural odorants
by ensembles of sparsely activated neurons in the olfactory bulb. *Sci Rep* **6**, 36514 (2016).
34. Vincis, R., Gschwend, O., Bhaukaurally, K., Beroud, J. & Carleton, A. Dense representation of natural
odorants in the mouse olfactory bulb. *Nat Neurosci* **15**, 537–539 (2012).
35. Rinberg, D., Koulakov, A. & Gelperin, A. Speed-Accuracy Tradeoff in Olfaction. *Neuron* **51**, 351–358
(2006).
36. Uchida, N. & Mainen, Z. F. Speed and accuracy of olfactory discrimination in the rat. *Nat Neurosci* **6**,
1224–1229 (2003).
37. Wesson, D. W., Carey, R. M., Verhagen, J. V. & Wachowiak, M. Rapid Encoding and Perception of
Novel Odors in the Rat. *PLoS Biol* **6**, e82 (2008).
38. Kato, H. K., Chu, M. W., Isaacson, J. S. & Komiyama, T. Dynamic Sensory Representations in the
Olfactory Bulb: Modulation by Wakefulness and Experience. *Neuron* **76**, 962–975 (2012).
39. Uchida, N., Poo, C. & Haddad, R. Coding and Transformations in the Olfactory System. *Annu. Rev.*
Neurosci. **37**, 363–385 (2014).
40. Yamada, Y. *et al.* Context- and Output Layer-Dependent Long-Term Ensemble Plasticity in a Sensory
Circuit. *Neuron* **93**, 1198-1212.e5 (2017).
41. Shani-Narkiss, H., Beniaguev, D., Segev, I. & Mizrahi, A. Stability and flexibility of odor
representations in the mouse olfactory bulb. *Front. Neural Circuits* **17**, (2023).
42. Langdon, C., Genkin, M. & Engel, T. A. A unifying perspective on neural manifolds and circuits for
cognition. *Nat Rev Neurosci* 1–15 (2023) doi:10.1038/s41583-023-00693-x.
43. Kriegeskorte, N. & Wei, X.-X. Neural tuning and representational geometry. *Nat Rev Neurosci* **22**,
703–718 (2021).
44. Baker, K. L., Vasan, G., Gumaste, A., Pieribone, V. A. & Verhagen, J. V. Spatiotemporal dynamics of
odor responses in the lateral and dorsal olfactory bulb. *PLoS Biol* **17**, e3000409 (2019).
45. Gire, D. H. *et al.* Temporal Processing in the Olfactory System: Can We See a Smell? *Neuron* **78**, 416–
432 (2013).
46. Schaefer, A. T. & Margrie, T. W. Spatiotemporal representations in the olfactory system. *Trends in*
Neurosciences **30**, 92–100 (2007).

47. Rennaker, R. L., Chen, C.-F. F., Ruyle, A. M., Sloan, A. M. & Wilson, D. A. Spatial and Temporal
Distribution of Odorant-Evoked Activity in the Piriform Cortex. *Journal of Neuroscience* **27**, 1534–
1542 (2007).
48. Chen, Z. & Padmanabhan, K. Top-down feedback enables flexible coding strategies in the olfactory
cortex. *Cell Reports* **38**, 110545 (2022).
49. Haddad, R. *et al.* Olfactory cortical neurons read out a relative time code in the olfactory bulb. *Nat*
Neurosci **16**, 949–957 (2013).
50. Wu, A. *et al.* Context-dependent plasticity of adult-born neurons regulated by cortical feedback. *Sci.*
Adv. **6**, eabc8319 (2020).
51. Froemke, R. C. & Dan, Y. Spike-timing-dependent synaptic modification induced by natural spike
trains. *Nature* **416**, 433–438 (2002).
52. Bi, G. & Poo, M. Synaptic Modification by Correlated Activity: Hebb's Postulate Revisited. *Annual*
Review of Neuroscience **24**, 139–166 (2001).
53. Markram, H., Gerstner, W. & Sjöström, P. J. Spike-Timing-Dependent Plasticity: A Comprehensive
Overview. *Front. Synaptic Neurosci.* **4**, (2012).
54. Padmanabhan, K. & Urban, N. N. Intrinsic biophysical diversity decorrelates neuronal firing while
increasing information content. *Nat Neurosci* **13**, 1276–1282 (2010).
55. Angelo, K. & Margrie, T. W. Population diversity and function of hyperpolarization-activated current
in olfactory bulb mitral cells. *Sci Rep* **1**, 50 (2011).
56. Kapoor, V. & Urban, N. N. Glomerulus-Specific, Long-Latency Activity in the Olfactory Bulb Granule
Cell Network. *Journal of Neuroscience* **26**, 11709–11719 (2006).
57. Friedrich, R. W. & Laurent, G. Dynamic Optimization of Odor Representations by Slow Temporal
Patterning of Mitral Cell Activity. *Science* **291**, 889–894 (2001).
58. Smear, M., Shusterman, R., O'Connor, R., Bozza, T. & Rinberg, D. Perception of sniff phase in mouse
olfaction. *Nature* **479**, 397–400 (2011).
59. Laurent, G. Olfactory network dynamics and the coding of multidimensional signals. *Nature reviews*
neuroscience **3**, 884–895 (2002).
60. Schreck, M. R. *et al.* State-dependent olfactory processing in freely behaving mice. *Cell Reports* **38**,
(2022).
61. Chockanathan, U. *et al.* Changes in pairwise correlations during running reshape global network
state in the main olfactory bulb. *Journal of neurophysiology* **125**, 1612–1623 (2021).
62. D'Souza, R. D. & Vijayaraghavan, S. Paying attention to smell: cholinergic signaling in the olfactory
bulb. *Front. Synaptic Neurosci.* **6**, (2014).
63. Boyd, A. M., Sturgill, J. F., Poo, C. & Isaacson, J. S. Cortical Feedback Control of Olfactory Bulb
Circuits. *Neuron* **76**, 1161–1174 (2012).
64. Otazu, G. H., Chae, H., Davis, M. B. & Albeanu, D. F. Cortical Feedback Decorrelates Olfactory Bulb
Output in Awake Mice. *Neuron* **86**, 1461–1477 (2015).
65. Magavi, S. S. P., Mitchell, B. D., Szentirmai, O., Carter, B. S. & Macklis, J. D. Adult-Born and
Preexisting Olfactory Granule Neurons Undergo Distinct Experience-Dependent Modifications of
their Olfactory Responses In Vivo. *J. Neurosci.* **25**, 10729–10739 (2005).
66. Kaplan, M. S., McNelly, N. A. & Hinds, J. W. Population dynamics of adult-formed granule neurons of
the rat olfactory bulb. *Journal of Comparative Neurology* **239**, 117–125 (1985).
67. Petreanu, L. & Alvarez-Buylla, A. Maturation and Death of Adult-Born Olfactory Bulb Granule
Neurons: Role of Olfaction. *J. Neurosci.* **22**, 6106–6113 (2002).
68. Kamimura, S. *et al.* New granule cells in the olfactory bulb are associated with high respiratory input
in an enriched odor environment. *Neuroscience Research* **182**, 52–59 (2022).

69. van Praag, H., Christie, B. R., Sejnowski, T. J. & Gage, F. H. Running enhances neurogenesis, learning,
and long-term potentiation in mice. *Proceedings of the National Academy of Sciences* **96**, 13427–
13431 (1999).
70. Gheusi, G. & Lledo, P.-M. Adult Neurogenesis in the Olfactory System Shapes Odor Memory and
Perception. in *Progress in Brain Research* vol. 208 157–175 (Elsevier, 2014).
71. Alonso, M. *et al.* Olfactory Discrimination Learning Increases the Survival of Adult-Born Neurons in
the Olfactory Bulb. *Journal of Neuroscience* **26**, 10508–10513 (2006).
72. Franks, K. M. *et al.* Recurrent Circuitry Dynamically Shapes the Activation of Piriform Cortex. *Neuron*
72, 49–56 (2011).
73. Apicella, A., Yuan, Q., Scanziani, M. & Isaacson, J. S. Pyramidal Cells in Piriform Cortex Receive
Convergent Input from Distinct Olfactory Bulb Glomeruli. *Journal of Neuroscience* **30**, 14255–14260
(2010).
74. Blazing, R. M. & Franks, K. M. Odor coding in piriform cortex: mechanistic insights into distributed
coding. *Current Opinion in Neurobiology* **64**, 96–102 (2020).
75. Chae, H., Banerjee, A., Dussauze, M. & Albeanu, D. F. Long-range functional loops in the mouse
olfactory system and their roles in computing odor identity. *Neuron* S0896627322008108 (2022)
doi:10.1016/j.neuron.2022.09.005.
76. Zeppilli, S. *et al.* Molecular characterization of projection neuron subtypes in the mouse olfactory
bulb. *eLife* **10**, e65445 (2021).
77. Fink, A. J. P. *et al.* Experience-dependent reorganization of inhibitory neuron synaptic connectivity.
2025.01.16.633450 Preprint at https://doi.org/10.1101/2025.01.16.633450 (2025).
78. Lemasson, M., Saghatelyan, A., Olivo-Marin, J.-C. & Lledo, P.-M. Neonatal and adult neurogenesis
provide two distinct populations of newborn neurons to the mouse olfactory bulb. *Journal of
Neuroscience* **25**, 6816–6825 (2005).
79. Whitman, M. C. & Greer, C. A. Adult neurogenesis and the olfactory system. *Progress in
Neurobiology* **89**, 162–175 (2009).
80. Qin, S. *et al.* Coordinated drift of receptive fields in Hebbian/anti-Hebbian network models during
noisy representation learning. *Nat Neurosci* **26**, 339–349 (2023).
81. Driscoll, L. N., Pettit, N. L., Minderer, M., Chettih, S. N. & Harvey, C. D. Dynamic reorganization of
neuronal activity patterns in parietal cortex. *Cell* **170**, 986–999 (2017).
82. Rokni, U., Richardson, A. G., Bizzi, E. & Seung, H. S. Motor learning with unstable neural
representations. *Neuron* **54**, 653–666 (2007).
83. Lee, J. S., Briguglio, J. J., Cohen, J. D., Romani, S. & Lee, A. K. The statistical structure of the
hippocampal code for space as a function of time, context, and value. *Cell* **183**, 620–635 (2020).
84. Gonzalez, W. G., Zhang, H., Harutyunyan, A. & Lois, C. Persistence of neuronal representations
through time and damage in the hippocampus. *Science* **365**, 821–825 (2019).
85. Toda, T., Parylak, S. L., Linker, S. B. & Gage, F. H. The role of adult hippocampal neurogenesis in
brain health and disease. *Mol Psychiatry* **24**, 67–87 (2019).
86. Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory engrams in the
hippocampus. *Nature* **558**, 292–296 (2018).
87. Lazarini, F. & Lledo, P.-M. Is adult neurogenesis essential for olfaction? *Trends in Neurosciences* **34**,
20–30 (2011).
88. Gheusi, G. *et al.* Importance of newly generated neurons in the adult olfactory bulb for odor
discrimination. *Proceedings of the National Academy of Sciences* **97**, 1823–1828 (2000).
89. Brann, J. H. & Firestein, S. J. A lifetime of neurogenesis in the olfactory system. *Front. Neurosci.* **8**,
(2014).
90. Gonçalves, J. T., Schafer, S. T. & Gage, F. H. Adult Neurogenesis in the Hippocampus: From Stem
Cells to Behavior. *Cell* **167**, 897–914 (2016).

91. Jurkowski, M. P. *et al.* Beyond the Hippocampus and the SVZ: Adult Neurogenesis Throughout the
Brain. *Frontiers in Cellular Neuroscience* **14**, (2020).
92. Sailor, K. A., Schinder, A. F. & Lledo, P.-M. Adult neurogenesis beyond the niche: its potential for
driving brain plasticity. *Current Opinion in Neurobiology* **42**, 111–117 (2017).
93. Izhikevich, E. M. Simple model of spiking neurons. *IEEE Trans. Neural Netw.* **14**, 1569–1572 (2003).
94. Connors, B. W. & Gutnick, M. J. Intrinsic firing patterns of diverse neocortical neurons. *Trends in
Neurosciences* **13**, 99–104 (1990).
95. Padmanabhan, K. & Urban, N. N. Disrupting information coding via block of 4-AP-sensitive
potassium channels. *Journal of Neurophysiology* **112**, 1054–1066 (2014).
96. Burton, S. D. & Urban, N. N. Rapid Feedforward Inhibition and Asynchronous Excitation Regulate
Granule Cell Activity in the Mammalian Main Olfactory Bulb. *Journal of Neuroscience* **35**, 14103–
14122 (2015).
97. Egger, V., Svoboda, K. & Mainen, Z. F. Dendrodendritic Synaptic Signals in Olfactory Bulb Granule
Cells: Local Spine Boost and Global Low-Threshold Spike. *Journal of Neuroscience* **25**, 3521–3530
(2005).
98. Gibson, J. R., Beierlein, M. & Connors, B. W. Two networks of electrically coupled inhibitory neurons
in neocortex. *Nature* **402**, 75–79 (1999).
99. Suzuki, N. & Bekkers, J. M. Microcircuits Mediating Feedforward and Feedback Synaptic Inhibition in
the Piriform Cortex. *Journal of Neuroscience* **32**, 919–931 (2012).
100.   Dhawale, A. K., Hagiwara, A., Bhalla, U. S., Murthy, V. N. & Albeanu, D. F. Non-redundant odor
coding by sister mitral cells revealed by light addressable glomeruli in the mouse. *Nat Neurosci* **13**,
1404–1412 (2010).
101.   Koulakov, A. A. & Rinberg, D. Sparse Incomplete Representations: A Potential Role of Olfactory
Granule Cells. *Neuron* **72**, 124–136 (2011).
102.   Lledo, P.-M., Gheusi, G. & Vincent, J.-D. Information Processing in the Mammalian Olfactory
System. *Physiological Reviews* **85**, 281–317 (2005).
103.   Wachowiak, M., Denk, W. & Friedrich, R. W. Functional organization of sensory input to the
olfactory bulb glomerulus analyzed by two-photon calcium imaging. *PNAS* **101**, 9097–9102 (2004).
104.   Padmanabhan, K. *et al.* Diverse Representations of Olfactory Information in Centrifugal
Feedback Projections. *Journal of Neuroscience* **36**, 7535–7545 (2016).
105.   Sailor, K. A. *et al.* Persistent Structural Plasticity Optimizes Sensory Information Processing in the
Olfactory Bulb. *Neuron* **91**, 384–396 (2016).
106.   Wanner, A. A. & Friedrich, R. W. Whitening of odor representations by the wiring diagram of the
olfactory bulb. *Nat Neurosci* **23**, 433–442 (2020).
107.   Willhite, D. C. *et al.* Viral tracing identifies distributed columnar organization in the olfactory
bulb. *Proceedings of the National Academy of Sciences* **103**, 12592–12597 (2006).
108.   Egger, V. & Urban, N. N. Dynamic connectivity in the mitral cell–granule cell microcircuit.
Seminars in Cell & Developmental Biology **17**, 424–432 (2006).
109.   Schoppa, N. E. & Westbrook, G. L. AMPA autoreceptors drive correlated spiking in olfactory bulb
glomeruli. *Nat Neurosci* **5**, 1194–1202 (2002).
110.   Soucy, E. R., Albeanu, D. F., Fantana, A. L., Murthy, V. N. & Meister, M. Precision and diversity in
an odor map on the olfactory bulb. *Nat Neurosci* **12**, 210–220 (2009).
111.   Luna, V. M. & Schoppa, N. E. GABAergic Circuits Control Input-Spike Coupling in the Piriform
Cortex. *Journal of Neuroscience* **28**, 8851–8859 (2008).
112.   Nunez-Parra, A., Maurer, R. K., Krahe, K., Smith, R. S. & Araneda, R. C. Disruption of centrifugal
inhibition to olfactory bulb granule cells impairs olfactory discrimination. *Proc Natl Acad Sci USA*
110, 14777–14782 (2013).

113.   Padmanabhan, K. *et al.* Centrifugal Inputs to the Main Olfactory Bulb Revealed Through Whole
695   Brain Circuit-Mapping. *Front. Neuroanat.* **12**, 115 (2019).
696   114.   Large, A. M., Vogler, N. W., Mielo, S. & Oswald, A.-M. M. Balanced feedforward inhibition and
697   dominant recurrent inhibition in olfactory cortex. *Proc Natl Acad Sci USA* **113**, 2276–2281 (2016).
698

 Figures and Tables

Figure 1. Spiking network model of adult-neurogenesis. (A1). Schematic of early olfactory circuit. M/T: mitral/tufted cells, GC: granule cells, abGCs: adult-born GCs, PCs: piriform cortical cell, FFIN: feedforward inhibitory neuron, FBIN: feedback inhibitory neuron. Example plus signs indicate excitatory synapses and minus signs indicate inhibitory synapses. (A2). Schematic of synaptic reshuffling of GCs due to adult-neurogenesis. On each day, 10% of GCs have their weights reshuffled, including the synaptic weights from M/T cells, other GCs or short-axon cells (SAC), and PCs to GC (feedback). (B1). Partial weight matrix on two example days between GCs and M/T cells (top), other GCs or SACs (middle), and feedback from PCs (bottom). (B2). Histograms of all synaptic weights across days. (B3). Weight matrix dissimilarity between Day-0 and each other day. (C1). An example model odor generated by stimulating different combinations of glomerular identity and timing of activation. (C2). Pair-wise correlations between model odors show the relative similarities and differences across all model odors.

Figure 2. Adult-neurogenesis modulates individual M/T cell responses but preserves population representation. (A1). Trail-averaged firing rate of two example M/T cells responding to the same odor (within-odor) on three different days (mean ± SD, n = 10 trials). (A2). Firing rate patterns of odor-activated M/T cells to the same odor on three days. M/T cells driven by different glomeruli are separated by white dashed line. The two arrows correspond to the two example cells in (A1). (A3). Pairwise within-odor full-ensemble correlation of single-trial M/T cell responses between each day averaged across all odors (n = 100). Each small box separated by the dashed lines is a 10x10 matrix (n = 10 trials) corresponding to autocorrelation (same day on diagonal) and cross-correlation (different days off diagonal). (A4). Full-ensemble correlation of trial-averaged M/T cell responses for within-odor (black solid line with error bar: mean ± SD, n = 100 odors).) and across-odor (black dashed line with error bar: mean ± SD, n = 10 pairs) between Day-0 and each other day. Grey dashed line with no error bar: same as the black solid line in (B4). (B1). Low-dimensional trajectories of M/T responses to two example odors (color) on two different days (darkness of color). Thin curves: single-trial trajectories, thick curves: trial-averaged trajectories, points: maximal-distance points on trajectory to the origin. (B2). Only maximal-distance points are shown for different odors on different days. (B3). Similar to (A3) but for reduced-space correlation computed using the low-dimensional M/T trajectories (single-trial). (B4). Similar to (A4) but for reduced-space correlation computed using the low-dimensional M/T trajectories (trial-averaged). Grey dashed line with no error bar: same as the black solid line in (A4).

737
Figure 3. Adult-neurogenesis modulates both individual PCs responses and
population representations. (A1). Trail-averaged firing rate of two example PCs
responding to the same odor on three different days (mean ± SD, n = 10 trials). (A2).
Firing rate patterns of partial PCs responding to the same odor on three days. The two
arrows correspond to the two example cells in (A1). (A3). Pairwise within-odor full-
ensemble correlation of single-trial PCs responses between each day averaged across
all odors (n = 100). Each small box separated by the dashed lines is a 10x10 matrix (n =
10 trials) corresponding to autocorrelation (same day on diagonal) and cross-correlation
(different days off diagonal). (A4). Full-ensemble correlation of trial-averaged cell (PCs:
purple; M/T: black) responses for within-odor (solid line with error bar: mean ± SD, n =
100 odors).) and across-odor (dashed line with error bar: mean ± SD, n = 10 pairs)
between Day-0 and each other day. The curves for M/T cells are the same lines as in
Fig.2. (B1). Low-dimensional trajectories of PCs responses to two example odors (color)
on two different days (darkness of color). Thin curves: single-trial trajectories, thick curves:
trial-averaged trajectories, points: maximal-distance points on trajectory to the origin. (B2).
Only maximal-distance points are shown for different odors on different days. (B3). Similar
to (A3) but for reduced-space correlation computed using the single-trial PCA trajectories
of PCs. (B4). Similar to (A4) but for reduced-space correlation computed using the trial-
averaged PCA trajectories of PCs.

Figure 4. Experience-dependent plasticity enhances representational stability in PCx. (A1). Different types of plastic synapses (blue) related to an abGC. Circle: inhibitory; triangle: excitatory. (A2). Dependence of synaptic modification on pre/post inter-spike interval used for the spike-timing-dependent plasticity (STDP). S1 and S2 are two example synapses with different modification – S1: facilitated; S2: suppressed. (B). Weight history of the two example synapses in A2 when an odor is repeatedly presented. Adult-neurogenesis happens on day-4 which randomly resets the weights. The synaptic weights stay constant without plasticity (black) while change trial by trial with plasticity (blue). (C1). Left: reduced-space correlation of trial-averaged M/T trajectories between Day-0 and each day. Blue solid line with error bar: within-odor and with STDP (mean ± SD, n = 10 odors); black dashed line without error bar: within-odor and without STDP. Black dashed line with error bar: across-odors, mean ± SD, n = 10 pairs. Right: drift rate for M/T trajectories with and without STDP. (C2). Same as C1 but for PCs trajectories.

Methods

Organization and architecture of the model

The MOB consisted of 50 glomeruli (G) corresponding to the olfactory receptor neuron (ORN) inputs into the MOB [19]. Each glomerulus was connected to 25 mitral/tufted (M/T) cells for a total 1250 M/T cells. Within the MOB, a local population of 12,500 inhibitory granule cells (GCs) formed reciprocal and lateral inhibitory connections with M/T cells. Individual M/T cell "projections" formed random excitatory connections with 10,000 piriform cortical cells (PCs) in PCx. These PCs in turn "projected" back to the olfactory bulb, provided excitatory centrifugal feedback onto the inhibitory granule cells in the bulb. Within PCx, two types of inhibitory interneurons were included: a population of 1250 feedforward inhibitory neurons (FFIs) that received excitatory input from M/T cells and inhibited both PCs and other FFIs, and a population of 1250 local feedback inhibitory neurons (FBIs) that received input from a random subset of PCs and subsequently inhibited PCs and other FBIs.

Voltage dynamics of individual neurons

The voltage dynamics of individual cells in the network are modeled as spiking neurons [93] described by a two-dimensional (2D) system of ordinary differential equations of the form,

$$\frac{dv}{dt} = 0.04v^2 + 5v + 140 - u + I$$
$$\frac{du}{dt} = a(bv - u)$$

(1)

with the after-spiking resetting

$$\text{if } v \geq 30mV, \text{ then } v \leftarrow c, u \leftarrow u + d \qquad (2)$$

Here v represents the voltage (mV) of the neuron and u represents a dimensionless membrane recovery variable accounting for the activation or inactivation of ionic currents; t is time and has unit of ms; a, b, c and d are the parameters by tuning which various firing patterns can be generated; I represents synaptic currents or injected dc-currents to the neuron.

We choose to use this neuron model to simulate the voltage dynamics of individual neurons because: 1). It combines the biological plausibility of the Hodgkin–Huxley neuron model and the computational efficiency of leaky integrate-and-fire neuron model, allowing us to simulate tens of thousands of spiking neurons simultaneously in our network; 2). Different combinations of the parameter values a, b, c and d can reproduce a diversity of firing patterns of neurons of known types, so we can capture the biophysical diversity in the firing properties for different types of neurons in olfactory system, such as the mitral/tufted (M/T) cells and granule cells in the main olfactory bulb (MOB), and piriform cortical cells and other local inhibitory interneurons in piriform cortex (PCx). In order to achieve heterogeneity such that different cells within the same type exhibit different dynamics, we introduced randomness in the parameter assignment (see Table 1). The r_i is a random variable uniformly distributed on the interval $[0,1]$ and i denotes the neuron index. For example, the parameter a will be distributed on the interval $[0.02, 0.1]$ within which various firing patterns can emerge. We also used r_i^2 or r_i^4 to bias the distribution to different extent for different cell types. Overall, based on our choice of parameters in the Izhikevich model, the spiking patterns of PCs and M/T cells largely fall into the category of regular spiking, intrinsically bursting or chattering neurons [21,94,95]. Inhibitory

816 neurons including GCs and FFIs/FBIs in the network can generate spiking patterns as
817 fast spiking neurons and low-threshold spiking neurons [96–99].
818
819 **Table 1** Parameters of Izhikevich neuron model for different cell types

	M/T cells	GCs	PCs	FFIs/FBIs
a	$0.1 - 0.08r_i^4$	$0.1 - 0.08r_i^2$	$0.02 + 0.08r_i$	$0.1 - 0.08r_i^2$
b	0.2	0.2	0.2	0.2
c	-65	$-65 + 15r_i^2$	-65	$-65 + 15r_i^2$
d	$2 + 6r_i^4$	2	$8 - 6r_i$	2

820
821   Synaptic input I to each neuron depends on the neuron type. For a cell i in MOB,
822 I_i is a linear superposition of various sources

$$I_i = I_i^{mc-ex} + I_i^{gc-in} + I_i^{osn} + I_i^{feedback} + \xi_i \tag{3}$$

Here, I_i^{mc-ex} represents excitation from M/T cells and exists for both M/T cells and GCs.
For GCs, when a M/T cells fires, the excitatory post-synaptic current (EPSC) I_i^{mc-ex} into
different GCs are delayed by different latencies, resulting in different spiking latencies of
GCs, consistent with previous experimental findings in the olfactory bulb granule cell
network [56]. The I_i^{gc-in} represents inhibition from GCs and exists for both M/T cells and
GCs. I_i^{osn} represents glomerular input and only exists for M/T cells. When a glomerulus
is activated by a model odor, it provides correlated inputs I_i^{osn} to the M/T cells driven by
that glomerulus. When a glomerulus is activated, the input current I_i^{osn} that an associated
M/T cell receives is modelled as a step function with Gaussian noise added. Since each
glomerulus receives inputs from a set of receptor neurons expressing the same olfactory
gene type, the inputs to individual glomerulus from receptor neurons are substantially
correlated [100–103]. Therefore, we assumed that the glomerular inputs to the apical
dendrites received by the M/T cells associated with the same glomerulus were correlated.
No input correlation between M/T cells associated with different glomeruli is assumed.
$I_i^{feedback}$ represents excitatory centrifugal input from piriform cells and is non-zero only
for GCs when feedback is ON. We set it to zero for all GCs when feedback is OFF. The
ξ_i represents Gaussian white noise input with zero mean and standard deviation $\sigma = 1.75$
for M/T cells and $\sigma = 0.8$ for GCs.
Similarly, for a cell i in PCx, I_i is composed of

$$I_i = I_i^{mob} + I_i^{pc-ex} + I_i^{in} + \eta_i \tag{4}$$

where I_i^{mob} represents input from M/T cells in MOB and only exists for piriform cortical
cells (PCs) and feedforward inhibitory neurons (FFIs); I_i^{pc-ex} represents excitation from
PCs and exists for both PCs and feedback inhibitory neurons (FBIs); I_i^{in} represents
inhibition from local inhibitory neurons including FFIs and FBIs; η_i represents Gaussian
white noise input (zero mean and standard deviation $\sigma = 0.9$) and only exists for PCs.
Each action potential fired by a presynaptic neuron will evoke a jump in the corresponding
synaptic inputs of all its postsynaptic targets by an amount equal to the appropriate
synaptic strength. For example, action potentials of a M/T cell induce jumps in the
excitatory currents of their postsynaptic target neurons, including I_i^{mc-ex} in M/T cells and
GCs in MOB, and I_i^{mob} in FFIs and PCs in PCx. These synaptic inputs then decay to
zero with time constant $10ms$. The height of the jump is determined by the pairwise

synaptic strength between any two neurons and their values are given in the synaptic
weight matrix which will described in the next section.
857
Synaptic strength and model network architecture

The MOB consists of 50 glomeruli, each of which drives 25 M/T cells, thus a total 1250
M/T cells in MOB. Besides, a local population of 12,500 inhibitory GCs formed reciprocal
and lateral inhibitory connections with M/T cells. Thus, within the MOB, we have a weight
matrix $\mathbf{W}_{mob}$ of 13,750 by 13,750 with its entry W_{mob}^{ij} representing the synaptic strength
from presynaptic neuron j to postsynaptic neuron i. Depending on the cell type, this
matrix $\mathbf{W}_{mob}$ can be partitioned into four sub-matrices, i.e., from M/T cell to M/T cell, from
M/T cell to GC, from GC to M/T cell and from GC to GC. The specific value of each entry
in $\mathbf{W}_{mob}$ was assigned randomly according to two parameters we chose for each sub-
matrix. One is the connection density (the percentage of non-zero synaptic weights) and
the other is the average synaptic strength (mean of a uniform distribution from which
individual synaptic weights are sampled). Each sub-matrix has its own value of the
connection density and average synaptic strength. In particular, the connection density
and average synaptic strength between M/T cells driven by the same glomerulus are
higher than between M/T cells driven by different glomeruli.
Individual M/T cell "projections" form random excitatory connections with 10,000
PCs and 1250 FFIs in PCx, giving rise to a feedforward weight matrix $\mathbf{W}_{ff}$ of 11,250 by
1250. Within PCx, PCs form recurrent excitations with each other. The FFIs inhibit both
PCs and other FFIs, and another population of 1250 FBIs that receive input from a
random subset of PCs inhibit PCs and other FBIs. Therefore, we have a matrix $\mathbf{W}_{pcx}$ of
12,500 by 12,500 that identifies all synaptic weights between cells in PCx. PCs "project"
back to the MOB, providing excitatory centrifugal feedback to GCs, giving rise to a
feedback weight matrix $\mathbf{W}_{fb}$ of 12,500 by 10,000. Under the condition of centrifugal
feedback OFF, this $\mathbf{W}_{fb}$ is set to be a zero matrix. The connection density and average
synaptic strength for all sub-matrices can be found in Table 2. The parameters are all
chosen heuristically based on previous theoretical and experimental studies listed in
Table 2.
Feedback projections from piriform cortex to the bulb may be structured, as
retrograde rabies tracing demonstrates that piriform cells projecting to GC populations in
the bulb tend to be spatially clustered [104]. Furthermore, a number of studies suggest that
GC synapses are especially sensitive to plasticity [9,105], either through adult neurogenesis
or more traditional mechanisms of synaptic reorganization. To implement all of these
features, we structure the feedback projections to GCs such that the PCs receiving
feedforward inputs from the M/T cells of certain glomeruli project back to the GCs which
are reciprocally connected with M/T cells associated with other glomeruli. Reciprocal
connectivity between M/T cells and GCs is defined as: M/T-1 excites GC-1 and GC-1
inhibits M/T-1, as observed by many studies [106,107]. Across the M/T population, there are
291 ± 9 (mean$\pm$SD, $n = 1250$ M/T cells) GCs that are reciprocally connected with each
M/T cell. As a result, each PC projects to 7368 ± 64 GCs (mean$\pm$SD, $n = 10000$ PCs)
with weight magnitude larger than 0.01. All feedback synaptic weights are randomly
generated with small magnitude less than 0.05, and this structure gives rise to a dense
but weak connectivity matrix $\mathbf{W}_{fb}$. Due to the sparsity of the PC firings when feedback is

ON, this dense and weak top-down connectivity ensures robust influence of PCs on GC
activity and thus the contribution of PCx on odor processing in MOB.
902
903 **Table 2** Network parameters controlling the connectivity between cell types.

	Connection density	Average synaptic strength	References
MCtoMC (same glomerulus)	0.8	0.25	2,108–110
MCtoMC (different glomeruli)*	0	0	
MC2GC	0.3	0.25	3,56,96,107
GC2MC	0.02	-0.4	
GC2GC	0.05	-0.1	63,96
MC2PC	0.5	0.06	4,6,22,111
MC2FFI	0.2	0.2	
PC2GC	0.9	0.03	64,112,113
PC2PC	0.01	0.1	22,72,99,114
FFI2PC	0.1	-0.1	
FBI2PC	0.8	-0.1	
FFI2FFI	0.01	-1.0	
PC2FBI	0.02	0.3	
FBI2FBI	0.02	-0.5	

904 *: We assumed zero connection between M/T cells associated with different glomeruli.
905

Model odor definition

Model odors are defined by the combinatorial patterns of glomeruli which are activated
successively with different glomerular timing, a pattern recapitulating the spatiotemporal
structure of odor inputs (Rubin and Katz, 1999; Meister and Bonhoeffer, 2001).
Specifically, when a model odor is presented, 3~10 glomeruli will be activated ($6 - 20\%$
of all glomeruli) and all the M/T cells associated with those glomeruli will receive
correlated glomerular input I^{osn} which lasts for 90ms. A table of 100 model odors were
defined as the odor inputs to our network.
914

Modeling adult-neurogenesis as weight reshuffling

Adult-neurogenesis of the granule cells constantly removes old granule cells and replace
them with adult-born ones. As a consequence, all the synapses from and onto old granule

cells are removed and new synapses with adult-born ones are built. We modeled this
process by weight reshuffling in the network with the total number of granule cells fixed.
On each simulation day, 10% of the granule cells had their synaptic weights reshuffled.
For each granule cell, the values of synaptic weights changed randomly on each day.
The distributions from which new synaptic weights were sampled from were the same
distributions as building the network (see Table 3.2). We only reshuffled the individual
weight values without changing the whole weight distributions.
925

Principal component analysis (PCA)
Spiking activity of each mitral/tufted cell and each piriform cell was binned into a $5ms$
sliding time window and averaged across trials (each model odor was presented in 10
trials). To perform the PCA analysis, we concatenated the trial-averaged responses of all
mitral/tufted cells to all 100 model odors on all simulation days, resulting in a large matrix
of 1250 cells by 247 time bins $\times 100$ odors $\times 11$ days. Response covariance matrices
(1250 by 1250) were computed for this concatenated matrix (after subtracting the mean
responses averaged across time bins, odors, and days). This gave us a single set of
eigenvectors, thus the same eigenspace into which cell responses for all days can be
projected and compared. Each 1250-dimensional cell response vector was then
projected onto the first 3 principal eigenvectors for visualization and the first 50 principal
eigenvectors for computations. The same procedure was also done for piriform cells. In
our simulations, these PCs captured the majority of variance relevant for odor identity
(~60–70% for M/T cells and ~55–65% for piriform cortex).
940

Population vectors of firing rates and PCA trajectories
We constructed the population vectors using either the firing rates of all cells or the PCA
trajectories of the first 50 dimensions. When using the firing rates, for each odor on each
day, the single-trial responses all the cells was a matrix of 1250 cells by 247 time bins for
M/T cells or $10,000$ cells by 247 time bins for piriform cells. We then converted the matrix
into a long vector of lengths 1250×247 for M/T cells and $10,000 \times 247$ for piriform cells.
When using PCA trajectories, the same procedures were applied, with only the cell
number replaced by the reduced dimensionality.
949

Pairwise correlation between population vectors
To measure the similarity of population responses over time, we computed the Pearson's
pairwise correlations of the population vectors constructed either by firing rates or PCA
trajectories on two comparison days (for example, day-i and day-j). For within-odor
correlation, the two population vectors were the responses (either single-trial or trial-
averaged) to the same odor on day-i and day-j. For across-odor correlation, the two
population vectors were the responses to any given two different odors on day-i and day-
j, and we computed that for all different odor pairs on those two days. When $i = j$, the
within-odor correlation was computed by comparing responses of even and odd trials.
959

Cosine similarity between population vectors
To gain a geometric perspective of the drift over days, we computed the cosine similarity
of the population vectors by $\theta_{i,j} = \boldsymbol{u}_i \cdot \boldsymbol{u}_j / \|\boldsymbol{u}_i\| \times \|\boldsymbol{u}_j\|$, where $\theta_{i,j}$ is the cosine similarity
between day-i and day-j, and $\boldsymbol{u}_i$ ($\boldsymbol{u}_j$) is the trial-averaged population vector on day-i (day-

j). We also estimated the within-day variability for each odor on each day by computing
the cosine similarity between the even trial-averaged and odd trial-averaged responses.
For within-day cosine similarity when $i = j$, we subtracted the estimated within-day
variability from the computed cosine similarity.
968

Decoding analysis: K nearest neighbor algorithm
The K nearest neighbor approach was used to decode odor identity from the projected
ensemble responses of PCs to any given odor pair [54]. Consistent with the computation
of symmetrized Kullback–Leibler divergence D_{KL}, analysis was performed in the space of
the first 50 principal components. The original data was broken up into testing and
training sets. The training sets established the location of PC responses to known odors
(i.e., known PC responses) in the principal component space and the testing sets were
probed with respect to these known PC responses. The Euclidian distance of the
unknown odors to all PC responses was then calculated and the K nearest neighbors
were used to determine to which odor the unknown PC activity was responding to. This
process of generating testing and training sets was repeated 30 times, with each repeat
reflecting a different random population of testing and training to ensure that the decoding
accuracy was not a result of artifacts of selecting a single testing/training population. Free
parameters in the K nearest neighbor algorithm include the ratio of testing to training data,
and the number of nearest neighbors used in the calculation. For training/testing, we used
ratios of 50%, 70% and 90%. We also examined the algorithm's accuracy when 3, 5 and
7 nearest neighbors were used.
986

Spike-timing-dependent plasticity (STDP)
To model the effect of spike trains on synaptic weights, we used the suppression model
given in [51]. Each pre- and post-synaptic spike was assigned with an efficacy $\epsilon_i = 1 -$
$e^{-(t_i - t_{i-1})/\tau_s}$, where ϵ_i is the efficacy of the ith spike, t_i and t_{i-1} are the timing of the ith
and $(i-1)$th spike respectively, and τ_s is the suppression time constant. The effect of
each pair of pre- and post-synaptic spikes on synaptic modification was given by $\Delta w_{ij} =$
$\epsilon_i^{pre} \epsilon_j^{pos} F(\Delta t_{ij})$, where ϵ_i^{pre} and ϵ_j^{pos} are the efficacies of the ith presynaptic spike and
jth postsynaptic spike, respectively, and Δt_{ij} is the interval between the two spikes $t_j^{pos} -$
t_i^{pre}. The function F represents the effect of temporal window for STDP, defined as:

996
$$F(\Delta t) \begin{cases} A_+ e^{-|\Delta t|/\tau_+} & if \quad \Delta t > 0 \\ A_- e^{-|\Delta t|/\tau_-} & if \quad \Delta t < 0 \end{cases} \tag{5}$$

Where A is the scaling factor, τ is the time constant, $+$ means long-term potentiation (LTP)
and $-$ means long-term depression (LTD). We chose $\tau_s^{pre} = 34ms$, $\tau_s^{pos} = 75ms$. $A_{+/-}$
and $\tau_{+/-}$ are drawn from normal distributions for each synapse, where $A_+ \sim \mathcal{N}(1.03, 0.1)$,
$A_- \sim \mathcal{N}(-0.51, 0.01)$, $\tau_+ \sim \mathcal{N}(13.3, 1.7)$ (in ms), $\tau_- \sim \mathcal{N}(34.5, 1.6)$ (in ms), and $\mathcal{N}(\mu, \sigma)$
represents normal distribution with mean μ and standard deviation σ. In our model, STDP
acts on two sets of connections. It applies to the synapses *onto* abGCs from M/T cells,
GC/SAC cells, and PCx neurons. It also applies to the synapses *from* abGCs, including
those onto M/T cells and GC/SAC cells.
1005

Supplemental information

Adult-neurogenesis allows for representational stability and flexibility in early olfactory system

Zhen Chen, Krishnan Padmanabhan

1012

Fig.S1 Odor definition and synaptic weight changes

(A). Glomerular activation latency for all 100 odors we defined. Each column corresponds to one odor. (B).

Histogram of the number of activated glomeruli by all odors ($n = 100$ odors). (C). Histogram of correlation

coefficients between each pair of odors, measuring the similarity between odors. (D). Sparsely sampled

synaptic weights on day-0 (x axis) and day-10 (y axis). Each dot is a synapse. Left: M/T to GCs; middle:

GC/SAC to GCs; right: PCs to GCs.

1019

Fig.S2 Example M/T cell responses on day-0 and day-10

Three example M/T cells (column) in response to the five example odors (row). Black: response on day-0; red: response on day-10. Some cells increase their responses while some cells decrease their responses from day-0 to day-10 (mean $\pm$ SD, $n = 10$ trials).

1024

Fig.S3 Example Piriform cell responses on day-0 and day-10

Three example piriform cells (column) in response to the five example odors (row). Black: response on day-0; red: response on day-10 (mean $\pm$ SD, $n = 10$ trials).

1028

Fig.S4 Example piriform cell responses (trial-averaged) on all 10 days

(A). Another three example piriform cells in response to four odors from day-0 and day=10. Each curve is the response averaged across 10 trials (smoothed by a gaussian kernel with 20ms width). Note some piriform cells increase or decrease their responses, while some cells change their temporal profile (cell-3).

1033

Fig.S5 Cell response changes between day-0 and day-10

(A). Histogram showing the differences between the trial-averaged response on day-10 and the trial-averaged response on day-0 for each odor-activated M/T cell (n=100 odors). Positive values indicate that the responses on day-10 are larger than day-0. (B). Same as (A) but for piriform cells.

1038

Fig.S6 Odor manifold and geometric reshaping of odor representations in PCx.

1039

Odor manifolds of M/T cells (top) and PCs (bottom) on two example days (red and blue) are plotted by
clusters of transparent round dots in the PCA space. Each column contains the trial-averaged trajectories
on two days (red and blue) evoked by three example odors. M/T trajectories stay close while PCs
trajectories are reshaped in different ways: shifted (A), rotated (B), and warped (C).

1040
1041
1042
1043

1044

Fig.S7 Quantifying geometric reshaping and decoding analysis on M/T and PCs

1045

1046   (A1). Cosine similarity to quantify the geometric reshaping using population firing rate as a function of
1047   intervals (number of days separated). (A2). Decoding accuracy by K-nearest neighbors algorithm trained
1048   on M/T population firing rate of each sample day and tested on M/T population firing rate of each reference
1049   day. (A3). Same as (A2) but using PCs population firing rate. (B). Same as (A) but using PCA trajectories.

1050

Fig.S8 Synaptic weight history of STDP

(A). Traces of five example synapses from M/T cells to GCs. STDP was applied when adult-neurogenesis happened at the start of the day. STDP drove the synaptic weight (y axis) to certain values through trial-by-trial exposure to certain odors. For some synapses, adult-neurogenesis only happened on one day while for other it happened on multiple days. (B). With STDP applied, three example piriform cells (column) in response to the three example odors (row). Black: response on day-0; red: response on day-10 (mean ± SD, n = 10 trials). (C). Slope of the synaptic weight traces over the last five trials of odor exposure with STPD applied. The slopes were all close to zero indicating the convergence of synaptic weights.